

GFUN AND THE AGM

Bruno Salvy

January 1998

Let α and β be two positive real numbers, with $\beta < \alpha$. The *arithmetic-geometric* mean of α and β is classically defined as the common limit of the sequences a_k and b_k defined by

$$a_{k+1} = \frac{\alpha_k + \beta_k}{2}, b_{k+1} = \sqrt{\alpha_k \beta_k}, \text{ with } a_0 = \alpha \text{ and } b_0 = \beta.$$

That the sequences converge to the same limit can be inferred from

$$a_{k+1}^2 - b_{k+1}^2 = \left(\frac{\alpha_k - \beta_k}{2} \right)^2.$$

This common limit is known by Maple as [GaussAGM](#) (α, β) . It was discovered by Gauss that the arithmetic-geometric mean is related to hypergeometric functions by

> **GaussAGM(a,b)=a/hypergeom([1/2, 1/2],[1],1-b^2/a^2);**

$$\text{GaussAGM}(\alpha, \beta) = \frac{\alpha}{\text{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2}\right], [1], 1 - \frac{\beta^2}{\alpha^2}\right)}$$

> **eval(subs(a=3.,b=2.,"));**

$$2.474680436 = 2.474680437$$

This worksheet, largely inspired by [1], shows how [gfun](#) can be used to guess and then prove this result, as well as a generalization of it due to J. M. Borwein and P. B. Borwein.

The functional equation

Following [1], we start by introducing a generalization of the arithmetic-geometric mean obtained by considering the following iteration where $1 < N$ is an integer:

$$a_{k+1} = \frac{\alpha_k + (N-1)\beta_k}{N}, b_{k+1} = \left(\alpha_{k+1}^N - \left(\frac{\alpha_k - \beta_k}{N} \right)^N \right)^{\frac{1}{N}}.$$

where the second equality is motivated by

$$a_{k+1}^N - b_{k+1}^N = \left(\frac{\alpha_k - \beta_k}{N} \right)^N$$

from which follows that both sequences converge to a common limit, which is denoted by

$M_N(a, b)$. The arithmetic-geometric mean corresponds to the case $N = 2$.

The function $M_N(a, b)$ is easily seen to be homogeneous: $M_N(\lambda a, \lambda b) = \lambda M_N(a, b)$, for $0 < \lambda$.

Together with the obvious property that $M_N(a_0, b_0) = M_N(a_1, b_1)$, this implies that for x in $(0, 1)$,

$$M_N\left(1, (1-x)^{\frac{1}{N}}\right) = (1 + (N-1)x) M_N\left(1, \frac{1-x}{1 + (N-1)x}\right) .$$

Defining the function $A_N(x)$ by

$$A_N(x) = \frac{1}{M_N\left(1, (1-x)^{\frac{1}{N}}\right)}$$

the equation above translates into the following *functional equation* for $A_N(x)$:

> $funeq := (1 + (N-1)x) A_N(x)^N = A_N\left(1 - \frac{1-x}{1 + (N-1)x}\right)^N$

which plays a central rôle in this worksheet. It is not too difficult to show that $A_N(x)$ is analytic in the neighborhood of the origin and that the functional equation above has a unique analytic solution in this neighborhood.

The quadratic case

This is the case $N = 2$ and Gauss's theorem is equivalent to stating that

$$A_2(x) = \text{hypergeom}\left[\left[\frac{1}{2}, \frac{1}{2}\right], [1], x\right] .$$

We now use [gfun](#) to first guess and then prove this result. The first step is to use the functional equation to compute a series expansion of $A_2(x)$, then we use this series to guess a possible closed form which turns out to be analytic, then we show that this analytic function does satisfy the functional equation.

Series expansion

Starting from the functional equation,

> **funeq2:=subs(N=2,A[2]=A,op(1,funeq)-op(2,funeq));**

$$funeq2 := (1+x) A(x^2) - A\left(1 - \frac{(1-x)^2}{(1+x)^2}\right)$$

a series solution is easily obtained by a method of undetermined coefficients:

> **sol:=1:**

> **for i to 12 do**

sol:=sol+x^i*solve(op(1,series(eval(subs(A=unapply(sol+a*x^i,x),funeq2)),x,i+2)),a) od:sol;

$$1 + \frac{1}{4}x + \frac{9}{64}x^2 + \frac{25}{256}x^3 + \frac{1225}{16384}x^4 + \frac{3969}{65536}x^5 + \frac{53361}{1048576}x^6 + \frac{184041}{4194304}x^7$$

$$+ \frac{41409225}{1073741824}x^8 + \frac{147744025}{4294967296}x^9 + \frac{2133423721}{68719476736}x^{10} + \frac{7775536041}{274877906944}x^{11}$$

$$+ \frac{457028729521}{17592186044416}x^{12}$$

Guessing the solution

From this series, [gfun](#) guesses a differential equation which could be satisfied by $A(x)$:

> **deq:=op(1,gfun[seriestodiffeq](series(sol,x,13),y(x),[ogf]));**

$$deq := \{y(0) = 1, y(x) + (-4 + 8x) \left(\frac{\partial}{\partial x} y(x) \right) + (-4x + 4x^2) \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} y(x) \right) \right), D(y)(0) = \frac{1}{4}\}$$

It turns out that Maple's dsolve function is unable to solve this differential equation:

> **dsolve(deq,y(x));**

We then use [gfun\[diffeqtorec\]](#) which deduces from this differential equation the recurrence satisfied by the Taylor coefficients of its solutions:

> **gfun[diffeqtorec](deq,y(x),u(n));**

$$\{(1 + 4n + 4n^2)u(n) + (-8n - 4 - 4n^2)u(n+1), u(0) = 1, u(1) = \frac{1}{4}\}$$

From this first order linear recurrence, a solution is easily found:

> **rsolve(",u(n));**

$$\frac{\Gamma\left(n + \frac{1}{2}\right)^2}{\Gamma(n+1)^2 \pi}$$

hence the sum:

> **y(x)=sum("x^n,n=0..infinity);**

$$y(x) = \text{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2}\right], [1], x\right)$$

Proving the result of the guess

The proof consists in showing that the function $y(x)$, which is obviously analytic, satisfies the functional equation

> **subs(A=y,funeq2)=0;**

$$(1+x)y(x^2) - y\left(1 - \frac{(1-x)^2}{(1+x)^2}\right) = 0$$

Our approach consists in using closure properties of solutions of linear differential equations that are implemented in [gfun](#) to compute a linear differential equation satisfied by the left-hand side of this equation. The proof then reduces to showing that 0 is the only solution of this differential equation that is compatible with the initial conditions, which are 0 up to a large order by construction of y .

It turns out that this proof can be performed directly from the differential equation, and would apply even if no closed-form had been found.

Given a linear differential equation satisfied by a series $y(x)$, the function [gfun\[algebraicsubs\]](#) computes a linear differential equation satisfied by $y(f(x))$ for any algebraic function, given by a polynomial P such that $P(x, f(x)) = 0$. Thus a differential equation satisfied by

$$y\left(1 - \frac{(1-x)^2}{(1+x)^2}\right) \text{ is easily computed from that satisfied by } y(x) :$$

> **deq:=op(select(has,deq,x)):**

> **deq1:=gfun[algebraicsubs](deq,numer(y-(1-(1-x)^2/(1+x)^2)),y(x));**

deq1 :=

$$(-1+x)y(x) + (-x^3 - 3x^2 - x + 1)\left(\frac{\partial}{\partial x}y(x)\right) + (-x^4 - x^3 + x^2 + x)\left(\frac{\partial}{\partial x}\left(\frac{\partial}{\partial x}y(x)\right)\right)$$

Similarly, $y(x^2)$ satisfies

> **gfun[algebraicsubs](deq,y-x^2,y(x));**

$$y(x)x + (-1 + 3x^2)\left(\frac{\partial}{\partial x}y(x)\right) + (-x + x^3)\left(\frac{\partial}{\partial x}\left(\frac{\partial}{\partial x}y(x)\right)\right)$$

and its product by $-(1+x)$ satisfies

> **deq2:=gfun[`diffeq*diffeq`](,y(x)+1+x,y(x));**

$$(-1-x)y(x) + (-1-x)(-x^3 - 3x^2 - x + 1)\left(\frac{\partial}{\partial x}y(x)\right) + (-1-x)(-x^4 - x^3 + x^2 + x)\left(\frac{\partial}{\partial x}\left(\frac{\partial}{\partial x}y(x)\right)\right)$$

$$aeqz := (1-x)y(x) + (x+x^2+3x-1) \left(\frac{\partial}{\partial x} y(x) \right) + (-x-x^2+x^3+x^4) \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} y(x) \right) \right)$$

From there, we deduce a differential equation satisfied by the left-hand side of the functional equation when applied to the hypergeometric function we have guessed:

> **gfun[`diffeq+diffeq`](deq1,deq2,y(x));**

$$(1-x)y(x) + (x+x^3+3x^2-1) \left(\frac{\partial}{\partial x} y(x) \right) + (-x-x^2+x^3+x^4) \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} y(x) \right) \right)$$

Analytic solutions of this equation have a coefficient sequence which satisfies

> **gfun[diffeqtorec](",y(x),u(n));**

$$\{n^2 u(n) + (2+n^2+4n)u(n+1) + (-2n+1-n^2)u(n+2) + (-6n-9-n^2)u(n+3), \\ u(1) = 4_C_0, u(2) = _C_0, u(0) = 4_C_0\}$$

and thus the first three zeroes of the Taylor expansion of the left-hand side of the functional equation conclude the proof.

The cubic case

It has been discovered by J. M. Borwein and P. B. Borwein that a hypergeometric expression also exists when $N=3$. Again, the same steps as above lead to guessing and then proving the following result by [gfun](#).

Theorem . [Borwein & Borwein 90] *The function $A_3(x)$ corresponding to the AGM iteration of order 3 has the following closed form:*

$$A_3(x) = \text{hypergeom} \left(\left[\begin{matrix} 1 & 2 \\ 3 & 3 \end{matrix} \right], [1], x \right) .$$

Series expansion

We start from the functional equation

> **funeq3:=subs(N=3,A[3]=A,op(1,funeq)-op(2,funeq));**

$$funeq3 := (1+2x) A(x^3) - A \left(1 - \frac{(1-x)^3}{(1+2x)^3} \right)$$

and compute the first terms of the series expansion of the solution:

> **sol:=1;**

> **for i to 12 do**

sol:=sol+x^i*solve(op(1,series(eval(subs(A=unapply(sol+a*x^i,x),funeq3)),x,i+2)),a) od:sol;

$$1 + \frac{2}{9}x + \frac{10}{81}x^2 + \frac{560}{6561}x^3 + \frac{3850}{59049}x^4 + \frac{28028}{531441}x^5 + \frac{1905904}{43046721}x^6 + \frac{14780480}{387420489}x^7$$

$$\begin{aligned}
& + \frac{116858170}{3486784401} x^8 + \frac{75957810500}{2541865828329} x^9 + \frac{616777421260}{22876792454961} x^{10} + \frac{5056555387520}{205891132094649} x^{11} \\
& + \frac{376081306946800}{16677181699666569} x^{12}
\end{aligned}$$

Guessing the solution

Again, this is a lucky situation where a differential equation can be guessed:

> **deq:=op(1,gfun[seriestodiffeq](series(sol,x,13),y(x),[ogf]));**

deq:=

$$\left\{ D(y)(0) = \frac{2}{9}, y(0) = 1, 2 y(x) + (-9 + 18x) \left(\frac{\partial}{\partial x} y(x) \right) + (-9x + 9x^2) \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} y(x) \right) \right) \right\}$$

From there, we find a closed-form as before

> **gfun[diffeqtorec](deq,y(x),u(n));**

$$\left\{ u(1) = \frac{2}{9}, u(0) = 1, (2 + 9n + 9n^2) u(n) + (-18n - 9 - 9n^2) u(n+1) \right\}$$

> **rsolve(",u(n));**

$$\frac{1}{2} \frac{\Gamma\left(n + \frac{2}{3}\right) \Gamma\left(n + \frac{1}{3}\right) \sqrt{3}}{\Gamma(n+1)^2 \pi}$$

> **y(x)=sum("x^n,n=0..infinity);**

$$y(x) = \text{hypergeom}\left(\left[\frac{1}{3}, \frac{2}{3}\right], [1], x\right)$$

Proving the result of the guess

The same routine applies:

> **subs(A=y,funeq3)=0;**

$$(1 + 2x) y(x^3) - y\left(1 - \frac{(1-x)^3}{(1+2x)^3}\right) = 0$$

> **deq:=op(select(has,deq,x));**

> **deq1:=gfun[algebraicsubs](deq,numer(y-(1-(1-x)^3/(1+2*x)^3)),y(x));**

$$deq1 := (2 - 4x + 2x^2) y(x) + (-1 + 8x^5 + 12x^4 + 4x^3 + 4x^2) \left(\frac{\partial}{\partial x} y(x) \right)$$

$$+ (4x^6 + 4x^5 + x^4 - 4x^3 - 4x^2 - x) \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} y(x) \right) \right)$$

> **gfun[algebraicsubs](deq,y-x^3,y(x));**

$$2y(x)x^2 + (-1 + 4x^3) \left(\frac{\partial}{\partial x} y(x) \right) + (-x + x^4) \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} y(x) \right) \right)$$

> **deq2:=gfun[`diffeq*diffeq`](",y(x)+1+2*x,y(x));**

$$\begin{aligned} \text{deq2} := & (2 - 4x + 2x^2)y(x) + (-1 + 8x^5 + 12x^4 + 4x^3 + 4x^2) \left(\frac{\partial}{\partial x} y(x) \right) \\ & + (4x^6 + 4x^5 + x^4 - 4x^3 - 4x^2 - x) \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} y(x) \right) \right) \end{aligned}$$

> **gfun[`diffeq+diffeq`](deq1,deq2,y(x));**

$$\begin{aligned} & (2 - 4x + 2x^2)y(x) + (-1 + 8x^5 + 12x^4 + 4x^3 + 4x^2) \left(\frac{\partial}{\partial x} y(x) \right) \\ & + (4x^6 + 4x^5 + x^4 - 4x^3 - 4x^2 - x) \left(\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} y(x) \right) \right) \end{aligned}$$

> **gfun[diffeqtorec](",y(x),u(n));**

$$\begin{aligned} & \{(4n^2 + 4n)u(n) + (16n + 12 + 4n^2)u(n+1) + (n^2 + 7n + 12)u(n+2) \\ & + (-16 - 16n - 4n^2)u(n+3) + (-46 - 4n^2 - 28n)u(n+4) \\ & + (-10n - 25 - n^2)u(n+5), u(2) = 0, u(4) = -C_0, u(3) = \frac{1}{2}C_0, u(0) = \frac{9}{4}C_0, \\ & u(1) = \frac{9}{2}C_0\} \end{aligned}$$

and thus the first five zeroes of the Taylor expansion of the left-hand side of the functional equation conclude the proof.

Conclusion

These results are very good examples of the use of **gfun** : experiments first lead to conjecture a general form for the solution to a problem and then a completely different process leads to a proof. However, the apparent ease with which the problems treated here are solved using **gfun** hides the preliminary work which led to the form under which this approach could work. For example this approach does not seem to work for higher values of N , where similar results might exist.

Bibliography

[1] Arithmetic-Geometric Means Revisited. Jonathan M. Borwein, Petr Lisonek and John A. Macdonald. *MapleTech* , **4-1** , pp. 20-27 (1997).