

# On Additive Analogues of Certain Arithmetic Smarandache Functions

József Sándor

Department of Mathematics,

Babeş-Bolyai University, 3400 Cluj-Napoca, Romania

1. The Smarandache, Pseudo-Smarandache, resp. Smarandache-simple functions are defined as ([7], [6])

$$S(n) = \min\{m \in \mathbb{N} : n|m!\}, \quad (1)$$

$$Z(n) = \min\left\{m \in \mathbb{N} : n \mid \frac{m(m+1)}{2}\right\}, \quad (2)$$

$$S_p(n) = \min\{m \in \mathbb{N} : p^n|m!\} \text{ for fixed primes } p. \quad (3)$$

The duals of  $S$  and  $Z$  have been studied e.g. in [2], [5], [6]:

$$S_*(n) = \max\{m \in \mathbb{N} : m!|n\}, \quad (4)$$

$$Z_*(n) = \max\left\{m \in \mathbb{N} : \frac{m(m+1)}{2} | n\right\}. \quad (5)$$

We note here that the dual of the Smarandache simple function can be defined in a similar manner, namely by

$$S_{p^*}(n) = \max\{m \in \mathbb{N} : m!|p^n\} \quad (6)$$

This dual will be studied in a separate paper (in preparation).

**2.** The additive analogues of the functions  $S$  and  $S_*$  are real variable functions, and have been defined and studied in paper [3]. (See also our book [6], pp. 171-174). These functions have been recently further extended, by the use of Euler's gamma function, in place of the factorial (see [1]). We note that in what follows, we could define also the additive analogues functions by the use of Euler's gamma function. However, we shall apply the more transparent notation of a factorial of a positive integer.

The additive analogues of  $S$  and  $S_*$  from (1) and (4) have been introduced in [3] as follows:

$$S(x) = \min\{m \in \mathbb{N} : x \leq m!\}, \quad S : (1, \infty) \rightarrow \mathbb{R}, \quad (7)$$

resp.

$$S_*(x) = \max\{m \in \mathbb{N} : m! \leq x\}, \quad S_* : [1, \infty) \rightarrow \mathbb{R} \quad (8)$$

Besides of properties relating to continuity, differentiability, or Riemann integrability of these functions, we have proved the following results:

**Theorem 1.**

$$S_*(x) \sim \frac{\log x}{\log \log x} \quad (x \rightarrow \infty) \quad (9)$$

(the same for  $S(x)$ ).

**Theorem 2.** *The series*

$$\sum_{n=1}^{\infty} \frac{1}{n(S_*(n))^\alpha} \quad (10)$$

is convergent for  $\alpha > 1$  and divergent for  $\alpha \leq 1$  (the same for  $S_*(n)$  replaced by  $S(n)$ ).

**3.** The additive analogues of  $Z$  and  $Z_*$  from (2), resp. (4) will be defined as

$$Z(x) = \min \left\{ m \in \mathbb{N} : x \leq \frac{m(m+1)}{2} \right\}, \quad (11)$$

$$Z_*(x) = \max \left\{ m \in \mathbb{N} : \frac{m(m+1)}{2} \leq x \right\} \quad (12)$$

In (11) we will assume  $x \in (0, +\infty)$ , while in (12)  $x \in [1, +\infty)$ .

The two additive variants of  $S_p(n)$  of (3) will be defined as

$$P(x) = S_p(x) = \min\{m \in \mathbb{N} : p^x \leq m!\}; \quad (13)$$

(where in this case  $p > 1$  is an arbitrary fixed real number)

$$P_*(x) = S_{p^*}(x) = \max\{m \in \mathbb{N} : m! \leq p^x\} \quad (14)$$

From the definitions follow at once that

$$Z(x) = k \Leftrightarrow x \in \left( \frac{(k-1)k}{2}, \frac{k(k+1)}{2} \right] \text{ for } k \geq 1 \quad (15)$$

$$Z_*(x) = k \Leftrightarrow x \in \left[ \frac{k(k+1)}{2}, \frac{(k+1)(k+2)}{2} \right) \quad (16)$$

For  $x \geq 1$  it is immediate that

$$Z_*(x) + 1 \geq Z(x) \geq Z_*(x) \quad (17)$$

Therefore, it is sufficient to study the function  $Z_*(x)$ .

The following theorems are easy consequences of the given definitions:

**Theorem 3.**

$$Z_*(x) \sim \frac{1}{2}\sqrt{8x+1} \quad (x \rightarrow \infty) \quad (18)$$

**Theorem 4.**

$$\sum_{n=1}^{\infty} \frac{1}{(Z_*(n))^\alpha} \text{ is convergent for } \alpha > 2 \quad (19)$$

and divergent for  $\alpha \leq 2$ . The series  $\sum_{n=1}^{\infty} \frac{1}{n(Z_*(n))^\alpha}$  is convergent for all  $\alpha > 0$ .

**Proof.** By (16) one can write  $\frac{k(k+1)}{2} \leq x < \frac{(k+1)(k+2)}{2}$ , so  $k^2 + k - 2x \leq 0$  and  $k^2 + 3k + 2 - 2x > 0$ . Since the solutions of these quadratic equations are  $k_{1,2} = \frac{-1 \pm \sqrt{8x+1}}{2}$ , resp.  $k_{3,4} = \frac{-3 \pm \sqrt{8x+1}}{2}$ , and remarking that  $\frac{\sqrt{8x+1}-3}{2} \geq$

1  $\Leftrightarrow x \geq 3$ , we obtain that the solution of the above system of inequalities is:

$$\begin{cases} k \in \left[1, \frac{\sqrt{1+8x}-1}{2}\right] & \text{if } x \in [1, 3); \\ k \in \left(\frac{\sqrt{1+8x}-3}{2}, \frac{\sqrt{1+8x}-1}{2}\right] & \text{if } x \in [3, +\infty) \end{cases} \quad (20)$$

So, for  $x \geq 3$

$$\frac{\sqrt{1+8x}-3}{2} < Z_*(x) \leq \frac{\sqrt{1+8x}-1}{2} \quad (21)$$

implying relation (18).

Theorem 4 now follows by (18) and the known fact that the generalized harmonic series  $\sum_{n=1}^{\infty} \frac{1}{n^\theta}$  is convergent only for  $\theta > 1$ .

The things are slightly more complicated in the case of functions  $P$  and  $P_*$ . Here it is sufficient to consider  $P_*$ , too.

First remark that

$$P_*(x) = m \Leftrightarrow x \in \left[\frac{\log m!}{\log p}, \frac{\log(m+1)!}{\log p}\right). \quad (22)$$

The following asymptotic results have been proved in [3] (Lemma 2) (see also [6], p. 172)

$$\log m! \sim m \log m, \quad \frac{m \log \log m!}{\log m!} \sim 1, \quad \frac{\log \log m!}{\log \log(m+1)!} \sim 1 \quad (m \rightarrow \infty) \quad (23)$$

By (22) one can write

$$\frac{m \log \log m!}{\log m!} - \frac{m}{\log m!} \log \log p \leq \frac{m \log x}{\log m!} \leq \frac{m \log \log(m+1)!}{\log m!} - (\log \log p) \frac{m}{\log m!},$$

giving  $\frac{m \log x}{\log m!} \rightarrow 1$  ( $m \rightarrow \infty$ ), and by (23) one gets  $\log x \sim \log m$ . This means that:

**Theorem 5.**

$$\log P_*(x) \sim \log x \quad (x \rightarrow \infty) \quad (24)$$

The following theorem is a consequence of (24), and a convergence theorem established in [3]:

**Theorem 6.** *The series  $\sum_{n=1}^{\infty} \frac{1}{n} \left( \frac{\log \log n}{\log P_*(n)} \right)^{\alpha}$  is convergent for  $\alpha > 1$  and divergent for  $\alpha \leq 1$ .*

Indeed, by (24) it is sufficient to study the series  $\sum_{n \geq n_0}^{\infty} \frac{1}{n} \left( \frac{\log \log n}{\log n} \right)^{\alpha}$  (where  $n_0 \in \mathbb{N}$  is a fixed positive integer). This series has been proved to be convergent for  $\alpha > 1$  and divergent for  $\alpha \leq 1$  (see [6], p. 174).

## References

- [1] C. Adiga and Taekyun Kim, *On a generalization of Sándor's function*, Proc. Jangjeon Math. Soc. **5**(2002), no. 2, 121-124.
- [2] J. Sándor, *On certain generalizations of the Smarandache function*, Notes Number Th. Discr. Math. **5**(1999), no. 2, 41-51.
- [3] J. Sándor, *On an additive analogue of the function S*, Notes Number Th. Discr. Math. **7**(2001), no. 3.
- [4] J. Sándor, *On the pseudo-Smarandache function*, Smarandache Notions J. **12**(2001), no. 1-2-3, 59-62.
- [5] J. Sándor, *On a dual of the pseudo-Smarandache function*, Smarandache Notions J. **13**(2002), no. 1-2-3, 18-23.
- [6] J. Sándor, *Geometric theorems, diophantine equations and arithmetic functions*, American Research Press, Rehoboth, NM, 2002 (see pp. 141-149, 156-158, 161-166, 171-174) [MR 1 906 446 (Review); Zbl. 1004.11001; Zbl. Math. Di 2002e.03960]
- [7] F. Smarandache, *A function in the Number theory*, An. Univ. Timișoara, Ser. Șt. Mat. **28**(1980), no. 1, 79-88.