

# Polygon Dissections and Standard Young Tableaux

Richard P. Stanley<sup>1</sup>  
Department of Mathematics  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
rstan@math.mit.edu

## ABSTRACT

A simple bijection is given between dissections of a convex  $(n+2)$ -gon with  $d$  diagonals not intersecting in their interiors and standard Young tableaux of shape  $(d+1, d+1, 1^{n-1-d})$ .

---

<sup>1</sup>Partially supported by National Science Foundation grant #DMS-9206374.

For  $0 \leq d \leq n - 1$ , let  $f(n, d)$  be the number of ways to draw  $d$  diagonals in a convex  $(n + 2)$ -gon, such that no two diagonals intersect in their interior. For instance,  $f(n, n - 1)$  is just the Catalan number  $C_n = \frac{1}{n+1} \binom{2n}{n}$ . A result going back to Kirkman [3], Prouhet [4], and Cayley [1] (with Cayley giving the first complete proof) asserts that

$$f(n, d) = \frac{1}{n + d + 2} \binom{n + d + 2}{d + 1} \binom{n - 1}{d}. \quad (1)$$

K. O'Hara and A. Zelevinsky observed (unpublished) that the right-hand side of (1) is just the number of standard Young tableaux (as defined, e.g., in [5, p. 66]) of shape  $(d + 1, d + 1, 1^{n-1-d})$ , where  $1^{n-1-d}$  denotes a sequence of  $n - 1 - d$  1's. It is natural to ask for a bijection between the polygon dissections and the standard Young tableaux. If one is willing to accept the formula for the number of standard Young tableaux of a fixed shape (either in the original form due to MacMahon or the hook-length formula of Frame-Robinson-Thrall), then one obtains a simple proof of equation (1). In this note we give a simple bijection of the desired type.

First we recall that there is a well-known bijection [2] between dissections  $D$  of an  $(n + 2)$ -gon with  $d$  diagonals and integer sequences  $\psi(D) = (a_1, a_2, \dots, a_{n+d+1})$  such that (a) either  $a_i = -1$  or  $a_i \geq 1$ , (b) exactly  $n$  terms are equal to  $-1$ , (c)  $a_1 + a_2 + \dots + a_i \geq 0$  for all  $i$ , and (d)  $a_1 + a_2 + \dots + a_{n+d+1} = 0$ . This bijection may be defined recursively as follows. Fix an edge  $e$  of the dissected polygon  $D$ . When we remove  $e$  from  $D$ , we obtain a sequence of dissected polygons  $D_1, D_2, \dots, D_k$  (where  $k + 1$  is the number of sides of the region of  $D$  to which  $e$  belongs), arranged in clockwise order, with  $D_i$  and  $D_{i+1}$  intersecting at a single vertex. If  $D_i$  consists of a single edge, then define  $\psi(D_i) = -1$ , and set recursively  $\psi(D) = (k - 1, \psi(D_1)^*, \psi(D_2)^*, \dots, \psi(D_{k-1})^*, \psi(D_k))$ , where  $\psi(D_k)^*$  denotes  $\psi(D_k)$  with a  $-1$  appended at the end.

Given a sequence  $(a_1, a_2, \dots, a_{n+d+1})$  as above, define a standard Young tableau  $T$  of shape  $(d + 1, d + 1, 1^{n-1-d})$  as follows. We insert the elements  $1, 2, \dots, n + d + 1$  successively into  $T$ . Once an element is inserted, it remains in place. (There is no "bumping" as in the Robinson-Schensted correspondence.) Suppose that the positive  $a_i$ 's are given by  $b_1, b_2, \dots, b_{d+1}$ , in that order. The insertion is then defined by the following three rules:

- If  $a_i > 0$ , then insert  $i$  at the end of the first row. (We write our tableaux in “English” style, so the longest row is at the top.)
- If  $a_i = -1$  and the number of  $-1$ ’s preceding  $a_i$  is given by  $b_1 + b_2 + \cdots + b_j$  for some  $j \geq 0$ , then insert  $i$  at the end of the second row.
- If  $a_i = -1$  and the number of  $-1$ ’s preceding  $a_i$  is not of the form  $b_1 + b_2 + \cdots + b_j$ , then insert  $i$  at the bottom of the first column.

It is an easy exercise to check that the above procedure yields the desired bijection.

**Example.** Let the sequence corresponding to a dissection  $D$  (with  $n = 14$ ,  $d = 6$ ) be given by

$(4, 2, -\mathbf{1}, 1, -1, -1, 3, -1, -\mathbf{1}, 1, 1, -1, -\mathbf{1}, -\mathbf{1}, -1, -1, -\mathbf{1}, 2, -\mathbf{1}, -\mathbf{1}, -1)$ .

We have  $(b_1, \dots, b_7) = (4, 2, 1, 3, 1, 1, 2)$ . We have printed in boldface those  $-1$ ’s that are preceded by  $b_1 + \cdots + b_j$   $-1$ ’s for some  $j$ . The corresponding standard tableau  $\psi(D)$  is given by

1	2	4	7	10	11	18
3	9	13	14	17	19	20
5						
6						
8						
12						
15						
16						
21						

## References

- [1] A. CAYLEY, On the partitions of a polygon, *Proc. London Math. Soc.* (1) **22** (1890–1891), 237–262.

- [2] I. M. H. ETHERINGTON, Some problems of non-associative combinations (1), *Edinburgh Math. Notes* **32** (1940), 1–6.
- [3] T. P. KIRKMAN, On the  $k$ -partitions of the  $r$ -gon and  $r$ -ace, *Phil. Trans. Royal Soc. London* **147** (1857), 217–272.
- [4] E. PROUHET, Question 774, *Nouvelles Annales Math.* **5** (1866), 384.
- [5] B. E. SAGAN, “The Symmetric Group,” Wadsworth & Brooks/Cole, Pacific Grove, CA, 1991.