

On a sequence related to the Josephus problem

RALF STEPHAN*

In this short note, we show that an integer sequence defined on the minimum of differences between divisor complements of its partial products is connected with the Josephus problem ($q=3$).

We prove the following theorem and, finally, state the relatedness of two constants.

Theorem. *Let a_n and b_n be recursively defined as*

$$a_0 = 4, \quad a_n = \min (|d_j - p_n/d_j| > 1), \quad p_n = \prod_{k=0}^{n-1} a_k, \\ d_j | p_n, \quad 1 \leq j \leq \sigma(p_n). \\ b_1 = 1, \quad b_n = \lceil \frac{1}{2} \sum_{k=1}^{n-1} b_k \rceil.$$

(1) *Then $a_n = 2^{b_n}$, for $n > 2$.*

The first terms of a_n and b_n are [S][Z]

$$a_{n \geq 0} = \{4, 3, 4, 2, 4, 8, 16, 64, \dots\}, \quad b_{n \geq 1} = \{1, 1, 1, 2, 3, 4, 6, 9, 14, \dots\}.$$

We need two lemmata.

Lemma 1. *For $k > 0$,*

$$(2) \quad \sigma(3 \cdot 2^k) = 2k + 2.$$

Proof. This is true for $k = 1$, and the set of divisors of $3 \cdot 2^{k+1}$ is the set of divisors of $3 \cdot 2^k$ plus 2^{k+1} and $3 \cdot 2^{k+1}$ itself. \square

Lemma 2. *Let $\delta(m)$ denote the smallest absolute value of the differences between complementary divisors of $m > 1$:*

$$\delta(m) = \min \left(\left| d_j - \frac{m}{d_j} \right| \right), \quad d_j | m, \quad 1 \leq j \leq \sigma(m).$$

Then

$$(3) \quad \delta(3 \cdot 2^k) = 2^{\lceil k/2 \rceil}, \quad k > 0.$$

Proof. Let us sort the divisors of $3 \cdot 2^k$ by size and call these D_j :

$$D_{1 \leq j \leq 2k+2} = \{1, (2, 3), \dots, (2^i, \frac{3}{2}2^i), (2^{i+1}, \frac{3}{2}2^{i+1}), \dots, (2^k, \frac{3}{2}2^k), 3 \cdot 2^k\}.$$

Any smallest complementary divisor difference must be the one where the divisors are in the exact middle of the sorted list, which, using (2), is $k + 1$. And so, $\delta(3 \cdot 2^k) = D_{k+2} - D_{k+1}$.

*The author can be reached at <mailto:ralf@ark.in-berlin.de>.

Now, the proposition (3) is true for $k = 1$. For every increase of k by one, $\sigma(m)$ increases by two, and the index of the wanted pair of divisors increases by one, so $D_{k+2} - D_{k+1}$ goes through the values

$$\begin{aligned}\frac{3}{2}2^i - 2^i &= 2^{i-1} \\ 2^{i+1} - \frac{3}{2}2^i &= 2^{i-1} \\ \frac{3}{2}2^{i+1} - 2^{i+1} &= 2^i \\ 2^{i+2} - \frac{3}{2}2^{i+1} &= 2^i\end{aligned}$$

so it doubles every second step which is just the meaning of (3). \square

Fixing the induction base at $\delta(p_3 = 48) = 2^1 = 2^{b_3}$ to make sure that $D_{k+2} - D_{k+1} > 1$, the main proposition (1) is now obvious, since the powers of two in a_n behave the same way under multiplication as unity does in b_n under addition.

Because the asymptotics of b_n are known[C], with

$$b_n = \left\lceil c \cdot \left(\frac{3}{2}\right)^n - \frac{1}{2} \right\rceil, \quad c = 0.36050455619661495910154466\dots,$$

the investigation of $a_{n \geq 3} = 2^{b_n}$ is settled, except for the closed form for c . Reble already proved[R] that b_n is connected to the Josephus problem. Independently, our numerics show that

$$(4) \quad c = \frac{2}{9}K(3),$$

with $K(3)$ the universal constant in the same problem with $q = 3$, a constant already discussed ([OW][HH]), and whose closed form is still unknown.

REFERENCES

- [C] B. Cloitre, OEIS, 11/2002, A073941.
- [GKP] R. L. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*, 2nd ed., Addison-Wesley, 1994
- [HH] L. Halbeisen and N. Hungerbühler, *The Josephus problem*, <http://citeseer.nj.nec.com/235856.html>.
- [OEIS] N. Sloane, *Online Encyclopedia of Integer Sequences*, <http://www.research.att.com/~njas/sequences/Seis.html>
- [OW] A. M. Odlyzko and H. S. Wilf, *Functional iteration and the Josephus problem*, *Glasgow Math. J.* 33 (1991), 235–240. <http://citeseer.nj.nec.com/odlyzko91functional.html>
- [R] D. Reble, message to seqfan mailing list, ID <3EA7336C.BBAE31C1@nk.ca>, 04/2003.
- [S] R. Stephan, OEIS, 04/2003, A082125.
- [Z] R. Zumkeller, OEIS, 11/2002, A073941.