

ENUMERATION OF CONCAVE INTEGER PARTITIONS

JAN SNELLMAN AND MICHAEL PAULSEN

ABSTRACT. An integer partition $\lambda \vdash n$ corresponds, via its Ferrers diagram, to an artinian monomial ideal $I \subset \mathbb{C}[x, y]$ with $\dim_{\mathbb{C}} \mathbb{C}[x, y]/I = n$. If λ corresponds to an integrally closed ideal we call it *concave*. We study generating functions for the number of concave partitions, unrestricted or with at most r parts.

1. CONCAVE PARTITIONS

By an *integer partition* $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots)$ we mean a weakly decreasing sequence of non-negative integers, all but finitely many of which are zero. The non-zero elements are called the *parts* of the partition. When writing a partition, we often will only write the parts; thus $(2, 1, 1, 0, 0, 0, \dots)$ may be written as $(2, 1, 1)$.

We write $r = \langle \lambda \rangle$ for the number of parts of λ , and $n = |\lambda| = \sum_i \lambda_i$; equivalently, we write $\lambda \vdash n$ if $n = |\lambda|$. The set of all partitions is denoted by \mathcal{P} , and the set of partitions of n by $\mathcal{P}(n)$. We put $|\mathcal{P}(n)| = p(n)$. By subscripting any of the above with r we restrict to partitions with at most r parts.

We will use the fact that \mathcal{P} forms a monoid under component-wise addition.

For an integer partition $\lambda \vdash n$ we define its *Ferrers diagram* $F(\lambda) = \{(i, j) \in \mathbb{N}^2 \mid i < \lambda_{j+1}\}$. In figure 1 the black dots comprise the Ferrers diagram of the partition $\mu = (4, 4, 2, 2)$.

Then $F(\lambda)$ is a finite *order ideal* in the partially ordered set (\mathbb{N}^2, \leq) , where $(a, b) \leq (c, d)$ iff $a \leq c$ and $b \leq d$. In fact, integer partitions correspond precisely to finite order ideals in this poset.

The complement $I(\lambda) = \mathbb{N}^2 \setminus F(\lambda)$ is a monoid ideal in the additive monoid \mathbb{N}^2 . Recall that for a monoid ideal I the *integral closure* \bar{I} is

$$\{\mathbf{a} \mid \ell \mathbf{a} \in I \text{ for some } \ell \in \mathbb{Z}_+\} \quad (1)$$


and that I is *integrally closed* iff it is equal to its integral closure.

Definition 1. The integer partition λ is *concave* iff $I(\lambda)$ is integrally closed. We denote by $\bar{\lambda}$ the unique partition such that $I(\bar{\lambda}) = \bar{I}(\lambda)$.

Now let R be the complex monoid ring of \mathbb{N}^2 . We identify \mathbb{N}^2 with the set of commutative monomials in the variables x, y , so that $R \simeq \mathbb{C}[x, y]$. Then a monoid ideal $I \subset \mathbb{N}^2$ corresponds to the monomial ideal J in R generated by the monomials $\{x^i y^j \mid (i, j) \in I\}$. Furthermore, since the monoid ideals of the form $I(\lambda)$ are precisely those with finite complement to \mathbb{N}^2 , those

1991 *Mathematics Subject Classification.* 05A17; 13B22.

Key words and phrases. Integer partitions, monomial ideals, integral closure.

FIGURE 1. μ and $\bar{\mu}$

monoid ideals will correspond to monomial ideals $J \subset R$ such that R/J has a finite \mathbb{C} -vector space basis (consisting of images of those monomials not in J). By abuse of notation, such monomial ideals are called *artinian*, and the \mathbb{C} -vector space dimension of R/J is called the *colength* of J .

We get in this way a bijection between

- (1) integer partitions of n ,
- (2) order ideals in (\mathbb{N}^2, \leq) of cardinality n ,
- (3) monoid ideals in \mathbb{N}^2 whose complement has cardinality n , and
- (4) monomial ideals in R of colength n .

Recall that if \mathfrak{a} is an ideal in the commutative unitary ring S , then the *integral closure* $\bar{\mathfrak{a}}$ consists of all $u \in S$ that fulfill some equation of the form

$$s^n + b_1 s^{n-1} + \cdots + b_0, \quad b_i \in \mathfrak{a}^i \quad (2)$$

Then \mathfrak{a} is always contained in its integral closure, which is an ideal. The ideal \mathfrak{a} is said to be *integrally closed* if it coincides with its integral closure.

For the special case $S = R$, we have that the integral closure of a monomial ideal is again a monomial ideal, and that the latter monomial ideal corresponds to the integral closure of the monoid ideal corresponding to the former monomial ideal. Hence, we have a bijection between

- (1) concave integer partitions of n ,
- (2) integrally closed monoid ideals in \mathbb{N}^2 whose complements have cardinality n , and
- (3) integrally closed monomial ideals in R of colength n .

Fröberg and Barucci [3] studied the growth of the number of ideals of colength n in certain rings, among them local noetherian rings of dimension 1. Studying the growth of the number of monomial ideals of colength n in R is, by the above, the same as studying the partition function $p(n)$. In this article, we will instead study the growth of the number of integrally closed monomial ideals in R , that is, the number of concave partitions of n .

2. INEQUALITIES DEFINING CONCAVE PARTITIONS

It is in general a hard problem to compute the integral closure of an ideal in a commutative ring. However, for monomial ideals in a polynomial ring, the following theorem, which can be found in e.g. [6], makes the problem feasible.

Theorem 2. Let $I \subset \mathbb{N}^2$ be a monoid ideal, and regard \mathbb{N}^2 as a subset of \mathbb{Q}^2 in the natural way. Let $\text{conv}_{\mathbb{Q}}(I)$ denote the convex hull of I inside \mathbb{Q}^2 . Then the integral closure of I is given by

$$\text{conv}_{\mathbb{Q}}(I) \cap \mathbb{N}^2 \quad (3)$$

Example 3. The partition $\mu = (4, 4, 2, 2)$ corresponds to the monoid ideal $((0, 4), (2, 2), (4, 0))$, which has integral closure $((0, 4), (1, 3), (2, 2), (3, 1), (4, 0))$. It follows that $\bar{\mu} = (4, 3, 2, 1)$. In figure 1 we have drawn the lattice points belonging to $F(\mu)$ as dots, and the lattice points belonging to $I(\lambda)$ as crosses.

The above theorem gives the following characterization of concave partitions:

Lemma 4. Let $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots)$ be a partition. Then λ is concave iff for all positive integers $i < j < k$,

$$\lambda_j < 1 + \lambda_i \frac{k-j}{k-i} + \lambda_k \frac{j-i}{k-i} \quad (4)$$

or, equivalently, if

$$\lambda_i(j-k) + \lambda_j(k-i) + \lambda_k(i-j) < k-i \quad (5)$$

3. GENERATING FUNCTIONS FOR SUPER-CONCAVE PARTITIONS

We will enumerate concave partitions by considering another class of partitions which is more amenable to enumeration, yet is close to that of concave partitions.

Definition 5. Let $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots)$ be a partition. Then λ is *super-concave* iff for all positive integers $i < j < k$,

$$\lambda_i(j-k) + \lambda_j(k-i) + \lambda_k(j-i) \leq 0 \quad (6)$$

The reader should note that it is actually a *stronger* property to be super-concave than to be concave. Unlike the latter property, it is not necessarily preserved by conjugation: the partition (2) is super-concave, hence concave, but its conjugate (1, 1) is concave but not super-concave.

Theorem 6. Let $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots)$ be a partition, and let $\mu = (\mu_1, \mu_2, \mu_3, \dots)$ be its conjugate, so that $|\{j | \mu_j = i\}| = \lambda_i - \lambda_{i+1}$ for all i . Then the following are equivalent:

- (i) λ is super-concave,
- (ii) for all positive ℓ ,

$$-\lambda_\ell + 2\lambda_{\ell+1} - \lambda_{\ell+2} \leq 0 \quad (7)$$

- (iii) for all positive ℓ ,

$$\lambda_{\ell+1} - \lambda_\ell \geq \lambda_{\ell+2} - \lambda_{\ell+1} \quad (8)$$

- (iv) $|\{k | \mu_k = i\}| \geq |\{k | \mu_k = j\}|$ whenever $i \leq j$.

Proof. (i) \iff (ii): Let \mathbf{e}_i be the vector with 1 in the i 'th coordinate and zeros elsewhere, let $\mathbf{f}_j = -\mathbf{e}_j + 2\mathbf{e}_{j+1} - \mathbf{e}_{j+2}$, and let $\mathbf{t}_{i,j,k} = (j-k)\mathbf{e}_i + (k-i)\mathbf{e}_j + (j-i)\mathbf{e}_k$. Clearly, (6) is equivalent with $\mathbf{t}_{i,j,k} \cdot \lambda \leq 0$, and (7) is equivalent with $\mathbf{f}_j \cdot \lambda \leq 0$. We have that $\mathbf{f}_\ell = \mathbf{t}_{\ell,\ell+1,\ell+2}$. Conversely, we

claim that $\mathbf{t}_{i,j,k}$ is a positive linear combination of different \mathbf{f}_ℓ . From this claim, it follows that if λ fulfills (7) for all ℓ then λ is super-concave.

We can without loss of generality assume that $i = 1$. Then it is easy to verify that

$$\mathbf{t}_{1,j,k} = \sum_{\ell=1}^{j-2} \ell(k-j)\mathbf{f}_\ell + \sum_{\ell=j-1}^{k-2} \ell(j-1)(k-\ell-1)\mathbf{f}_\ell \quad (9)$$

(ii) \iff (iii) \iff (iv) : This is obvious. \square

The *difference operator* Δ is defined on partitions by

$$\Delta(\lambda_1, \lambda_2, \lambda_3, \dots) = (\lambda_1 - \lambda_2, \lambda_2 - \lambda_3, \lambda_3 - \lambda_4, \dots) \quad (10)$$

We get that the *second order difference operator* Δ^2 is given by

$$\begin{aligned} \Delta^2(\lambda_1, \lambda_2, \lambda_3, \dots) &= \Delta(\Delta(\lambda_1, \lambda_2, \lambda_3, \dots)) = \\ &= (\lambda_1 - 2\lambda_2 + \lambda_3, \lambda_2 - 2\lambda_3 + \lambda_4, \lambda_3 - 2\lambda_4 + \lambda_5, \dots) \end{aligned} \quad (11)$$

Corollary 7. *The super-concave partitions are precisely those with non-negative second differences.*

Definition 8. Let $p_{sc}(n)$ denote the number of super-concave partitions of n , and $p_{sc}(n, r)$ denote the number of super-concave partitions of n with at most r parts. Let similarly $p_c(n)$ and $p_c(n, r)$ denote the number of super-concave partitions of n , and the number of super-concave partitions of n with at most r parts, respectively. For a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ let $\mathbf{x}^\lambda = x_1^{\lambda_1} x_2^{\lambda_2} \dots$, and define

$$\begin{aligned} PS(\mathbf{x}) &= \sum_{\lambda \text{ super-concave}} \mathbf{x}^\lambda \\ PS_r(x_1, \dots, x_r) &= PS(x_1, x_2, \dots, x_r, 0, 0, 0, \dots) = \sum_{\substack{\lambda \text{ super-concave} \\ \lambda_{r+1}=0}} \mathbf{x}^\lambda \\ PC(\mathbf{x}) &= \sum_{\lambda \text{ concave}} \mathbf{x}^\lambda \\ PC_r(x_1, \dots, x_r) &= PC(x_1, x_2, \dots, x_r, 0, 0, 0, \dots) = \sum_{\substack{\lambda \text{ concave} \\ \lambda_{r+1}=0}} \mathbf{x}^\lambda \end{aligned} \quad (12)$$

Partitions with non-negative second differences have been studied by Andrews [2], who proved that there are as many such partitions of n as there are partitions of n into triangular numbers.

Canfield et al [4] have studied partitions with non-negative m 'th differences. Specialising their results to the case $m = 2$, we conclude:

Theorem 9. *Let n, r be denote positive integers.*

(i) *There is a bijection between partitions of n into triangular numbers and super-concave partitions.*

(ii) The multi-generating function for super-concave partitions is given by

$$PS(\mathbf{x}) = \frac{1}{\prod_{i=1}^{\infty} \left(1 - \prod_{j=1}^i x_j^{1+i-j}\right)}$$

$$= 1 + x_1 + x_1^2 + x_1^3 + x_1^4 + x_1^2 x_2 + x_1^5 + x_1^4 x_2 + x_1^3 x_2 + \dots \quad (13)$$

(iii) The multi-generating function for super-concave partitions with at most r parts is given by

$$PS_r(x_1, x_2, \dots, x_r) = \frac{1}{\prod_{i=1}^r \left(1 - \prod_{j=1}^i x_j^{1+i-j}\right)} \quad (14)$$

(iv) The generating function for super-concave partitions is

$$PS(t) = \sum_{n=0}^{\infty} p_{sc}(n)t^n = \prod_{i=1}^{\infty} \frac{1}{1 - t^{\frac{i(i+1)}{2}}} \quad (15)$$

and the one for super-concave partitions with at most r parts is

$$PS_r(t) = \sum_{n=0}^{\infty} p_{sc}(n, r)t^n = \prod_{i=1}^r \frac{1}{1 - t^{\frac{i(i+1)}{2}}} \quad (16)$$

(v) The proportion of super-concave partitions with at most r parts among all partitions with at most r parts is

$$\frac{r!}{\prod_{i=1}^r \frac{i(i+1)}{2}}. \quad (17)$$

(vi) As $n \rightarrow \infty$,

$$p_{sc}(n) \sim cn^{-3/2} \exp(3Cn^{1/3})$$

$$C = 2^{-1/3} [\zeta(3/2)\Gamma(3/2)]^{2/3}, \quad c = \frac{\sqrt{3}}{12} \left(\frac{C}{\pi}\right)^{3/2} \quad (18)$$

The sequence $(p_{sc}(n))_{n=0}^{\infty}$ is identical to sequence A007294 in OEIS [8]. We have submitted the sequences $(p_{sc}(n, r))_{n=0}^{\infty}$, for $r = 3, 4$, in OEIS [8], as A086159 and A086160. The sequence for $r = 2$ was already in the database, as A008620.

3.1. Other appearances of super-concave partitions in the literature.

The bijection between partitions into triangular numbers and partitions with non-negative second difference is mentioned in A007294 in OEIS [8], together with a reference to Andrews [2]. That sequence has been contributed by Mira Bernstein and Roland Bacher; we thank Philippe Flajolet for drawing our attention to it.

Gert Almkvist [1] gives an asymptotic analysis of $p_{sc}(n)$ which is finer than (18).

Another derivation of the generating functions above can found in a forthcoming paper ‘‘Partition Bijections, a Survey’’ [7] by Igor Pak. He observes that the set of super-concave partitions with at most r parts consists of the lattice points of the unimodular cone spanned by the vectors $v_0 = (1, \dots, 1)$ and $v_i = (i - 1, i - 2, \dots, 1, 0, 0, \dots)$ for $1 \leq i \leq r$.

Corteel and Savage [5] calculate rational generating functions for classes of partitions defined by linear homogeneous inequalities. This applies to super-concave partitions, but not directly to concave partitions, since the inequalities (5) defining them are inhomogeneous.

4. GENERATING FUNCTIONS FOR CONCAVE PARTITIONS

Theorem 10. *Let r be a positive integer. Then*

$$PC_r(x_1, \dots, x_r) = \frac{Q_r(x_1, \dots, x_r)}{\prod_{i=1}^r \left(1 - \prod_{j=1}^i x_j^{1+i-j}\right)} \quad (19)$$

where $Q_r(x_1, \dots, x_r)$ is a polynomial satisfying

- (i) $Q_r(x_1, \dots, x_r)$ has integer coefficients,
- (ii) $Q_r(1, \dots, 1) = 1$,
- (iii) all exponent vectors of the monomials that occur in Q_r are weakly decreasing, and
- (iv) $Q_r(x_1, \dots, x_r) = Q_{r+1}(x_1, \dots, x_r, 0)$.

Furthermore,

$$PC(\mathbf{x}) = \frac{Q(\mathbf{x})}{\prod_{i=1}^{\infty} \left(1 - \prod_{j=1}^i x_j^{1+i-j}\right)} \quad (20)$$

where $Q(\mathbf{x})$ is a formal power series with the property that for each ℓ , $Q(x_1, \dots, x_\ell, 0, 0, \dots) = Q_\ell(x_1, \dots, x_\ell)$; in other words,

$$Q = 1 + \sum_{i=1}^{\infty} (Q_i - Q_{i-1})$$

Proof. Let A be the matrix with r columns whose rows consists of all truncations of the vectors $\mathbf{t}_{i,j,k}$ introduced in the proof of Theorem 6, for $i < j < k$, $k < r + 2$. For example, if $r = 3$ and if we order the 3-subsets of $\{1, 2, 3, 4 <\}$ lexicographically we get that

$$A = \begin{pmatrix} -1 & 2 & -1 \\ -2 & 3 & 0 \\ -1 & 0 & 3 \\ 0 & -1 & 2 \end{pmatrix}$$

Then a super-concave partition with at most r parts corresponds to a solution to

$$A\mathbf{z} \leq \mathbf{0}, \quad \mathbf{z} \geq \mathbf{0} \quad (21)$$

whereas a concave partition with at most r parts corresponds to a solution to

$$A\mathbf{z} \leq \mathbf{b}, \quad \mathbf{z} \geq \mathbf{0} \quad (22)$$

where the entry of \mathbf{b} which corresponds to the row of A indexed by (i, j, k) is $i - k$. It follows from a theorem in Stanley's "green book" [9] that the multi-generating functions of these two solution sets have the same denominator, and that their numerator evaluates to the same value after substituting 1 for each formal variable.

All monomials in

$$\prod_{i=1}^r \left(1 - \prod_{j=1}^i x_j^{1+i-j} \right)$$

have weakly decreasing exponent vectors, hence this is also true for $PC_r(x_1, \dots, x_r)$.

The assertion about $PC(\mathbf{x})$ follows by passing to the limit. \square

Our calculations indicate that

$$\begin{aligned} Q_1(\mathbf{x}) &= 1 \\ Q_2(\mathbf{x}) &= 1 + x_1x_2 - x_1^2x_2 \\ Q_3(\mathbf{x}) &= Q_2(\mathbf{x}) + x_3(x_1^5x_2^3 - x_1^4x_2^3 - 2x_1^3x_2^2 + x_1^2x_2^2 + x_1x_2) \end{aligned} \quad (23)$$

Corollary 11. (i) *The generating function for concave partitions with at most r parts is given by*

$$PC_r(t) = \sum_{n=0}^{\infty} p_c(n, r)t^n = \frac{Q_r(t)}{\prod_{i=1}^r \left(1 - t^{\frac{i(i+1)}{2}} \right)} \quad (24)$$

where $Q_r(1) = 1$, and the numerator has degree strictly smaller than $r^3/6 + r^2/2 + r/3$.

(ii) *The proportion of concave partitions with at most r parts among all partitions with at most r parts is*

$$\frac{r!}{\prod_{i=1}^r \frac{i(i+1)}{2}}. \quad (25)$$

Proof. The only thing which does not follow immediately from substituting $x_i = t$ in the previous theorem is the assertion about the degree of the numerator. From Stanley's "grey book" [10, Theorem 4.6.25] we have that the rational function $PC_r(t, \dots, t)$ is of degree < 0 . The degree of the denominator is

$$\sum_{i=1}^r \frac{i(i+1)}{2} = \frac{r^3}{6} + \frac{r^2}{2} + \frac{r}{3}$$

so the result follows. \square

We can therefore say with absolute certainty that the first $Q_r(t)$ are as follows:

$$\begin{aligned} Q_1(t) &= 1 \\ Q_2(t) &= 1 + t^2 - t^3 \\ Q_3(t) &= 1 + t^2 + t^5 - 2t^6 - t^8 + t^9 \\ Q_4(t) &= 1 + t^2 + t^4 + t^5 - t^6 - t^7 + 2t^9 - 2t^{10} - t^{11} - 2t^{12} + \\ &\quad + 2t^{13} - t^{14} - t^{15} + t^{16} + t^{17} + t^{18} - t^{19} \end{aligned} \quad (26)$$

Hence, we believe that

$$PC(t) = \frac{1 + t^2 + O(t^3)}{\prod_{i=1}^{\infty} \left(1 - t^{\frac{i(i+1)}{2}} \right)} \quad (27)$$

We've calculated that

$$\begin{aligned}
 PC(t) = \sum_{n=0}^{\infty} p_c(n)t^n = & 1 + t + 2t^2 + 3t^3 + 4t^4 + 7t^5 + 9t^6 + 11t^7 + \\
 & + 17t^8 + 23t^9 + 28t^{10} + 39t^{11} + 48t^{12} + 59t^{13} + 79t^{14} + \\
 & + 100t^{15} + 121t^{16} + 152t^{17} + 185t^{18} + 225t^{19} + 280t^{20} + O(t^{21}) \quad (28)
 \end{aligned}$$

It seems likely that $\log p_c(n)$ grows as $n^{1/3}$ (i.e. approximately as fast as pseudo-convex partitions), but we can not prove this, since we have no estimates of the numerator in (27).

We have submitted $(p_c(n))_{n=0}^{\infty}$ to the OEIS [8]; it is A084913. The sequences $(p_c(n, r))_{n=0}^{\infty}$ are A086161, A086162, and A086163 for $r = 2, 3, 4$.

REFERENCES

- [1] Gert Almkvist. Asymptotics of various partitions. manuscript.
- [2] George E. Andrews. MacMahon's partition analysis. II. Fundamental theorems. *Ann. Comb.*, 4(3-4):327–338, 2000. Conference on Combinatorics and Physics (Los Alamos, NM, 1998).
- [3] Valentina Barucci and Ralf Fröberg. On the number of ideals of finite colength. In *Geometric and combinatorial aspects of commutative algebra (Messina, 1999)*, volume 217 of *Lecture Notes in Pure and Appl. Math.*, pages 11–19. Dekker, New York, 2001.
- [4] Rod Canfield, Sylvie Corteel, and Pawel Hitczenko. Random partitions with non-negative r -th differences. *Adv. in Appl. Math.*, 27(2-3):298–317, 2001. Special issue in honor of Dominique Foata's 65th birthday (Philadelphia, PA, 2000).
- [5] S. Corteel and C. D. Savage. Partitions and compositions defined by inequalities. math.CO/0309110.
- [6] David Eisenbud. *Commutative Algebra with a View Toward Algebraic Geometry*, volume 150 of *Graduate Texts in Mathematics*. Springer Verlag, 1995.
- [7] Igor Pak. Partition bijections, a survey. *Ramanujan Journal*, to appear.
- [8] Neil J. A. Sloane. The on-line encyclopedia of integer sequences. <http://www.research.att.com/~njas/sequences/index.html>.
- [9] Richard P. Stanley. *Combinatorics and Commutative Algebra*, volume 41 of *Progress in Mathematics*. Birkhäuser, 2 edition, 1996.
- [10] Richard P. Stanley. *Enumerative combinatorics. Vol. 1*. Cambridge University Press, Cambridge, 1997. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original.

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, SE-10691 STOCKHOLM, SWEDEN

E-mail address: Jan.Snellman@math.su.se