

Overlap-free words and spectra of matrices

Raphaël M. Jungers

*Division of Applied Mathematics, Université catholique de Louvain, 4 avenue
Georges Lemaitre, B-1348 Louvain-la-Neuve, Belgium, Email:
raphael.jungers@uclouvain.be*

Vladimir Yu. Protasov

*Department of Mechanics and Mathematics, Moscow State University, Vorobyovy
Gory, Moscow, 119992, Russia, Email: v-protassov@yandex.ru*

Vincent D. Blondel

*Division of Applied Mathematics, Université catholique de Louvain, 4 avenue
Georges Lemaitre, B-1348 Louvain-la-Neuve, Belgium, Email:
vincent.blondel@uclouvain.be*

Abstract

Overlap-free words are words over the binary alphabet $A = \{a, b\}$ that do not contain factors of the form $xvxx$, where $x \in A$ and $v \in A^*$. We analyze the asymptotic growth of the number u_n of overlap-free words of length n as $n \rightarrow \infty$. We obtain explicit formulas for the minimal and maximal rates of growth of u_n in terms of spectral characteristics (the lower spectral radius and the joint spectral radius) of certain sets of matrices of dimension 20×20 . Using these descriptions we provide new estimates of the rates of growth that are within 0.4% and 0.03% of their exact values. The best previously known bounds were within 11% and 3% respectively. We then prove that the value of u_n actually has the same rate of growth for “almost all” natural numbers n . This “average” growth is distinct from the maximal and minimal rates and can also be expressed in terms of a spectral quantity (the Lyapunov exponent). We use this expression to estimate it. In order to obtain our estimates, we introduce new algorithms to compute spectral characteristics of sets of matrices. These algorithms can be used in other contexts and are of independent interest.

Key words: Overlap-free words, Combinatorics on words, Joint spectral radius, Lyapunov exponent.

1 Introduction

Binary overlap-free words have been studied for more than a century. These are words over the binary alphabet $A = \{a, b\}$ that do not contain factors of the form $xvxvx$, where $x \in A$ and $v \in A^*$. For instance, the word $baabaa$ is overlap free, but the word $baabaab$ is not, since it can be written $xuxux$ with $x = b$ and $u = aa$. See [1] for a recent survey. Thue [22, 23] proved in 1906 that there are infinitely many overlap-free words. Indeed, the well-known Thue-Morse sequence¹ is overlap-free, and so the set of its factors provides an infinite number of different overlap-free words. The asymptotics of the number u_n of such words of a given length n was analyzed in a number of subsequent contributions². The number of factors of length n in the Thue-Morse sequence is proved in [6] to be larger than $3n$, thus providing a linear lower bound on u_n :

$$u_n \geq 3n.$$

The next improvement was obtained by Restivo and Salemi [20]. By using a certain decomposition result, they showed that the number of overlap-free words grows at most polynomially:

$$u_n \leq Cn^r,$$

where $r = \log(15) \approx 3.906$. This bound has been sharpened successively by Kfoury [11], Kobayashi [12], and finally by Lepisto [13] to the value $r = 1.37$. One could then suspect that the sequence u_n grows linearly. However, Kobayashi [12] proved that this is not the case. By enumerating the subset of overlap-free words of length n that can be infinitely extended to the right he showed that $u_n \geq Cn^{1.155}$ and so we have

$$C_1 n^{1.155} \leq u_n \leq C_2 n^{1.37}.$$

It is worth noting that the sequence u_n is 2-regular, as shown by Carpi [7]. On Figure 1(a) we show the values of the sequence u_n for $1 \leq n \leq 200$ and on Figure 1(b) we show the behavior of $\log u_n / \log n$ for larger values of n . One can see that the sequence u_n is not monotonic, but is globally increasing with n . Moreover the sequence does not appear to have a polynomial growth since the value $\log u_n / \log n$ does not seem to converge. In view of this, a natural question arises: is the sequence u_n asymptotically equivalent to n^r for some r ? Cassaigne proved in [8] that the answer is negative. He introduced the lower and the upper exponents of growth:

¹ The Thue-Morse sequence is the infinite word obtained as the limit of $\theta^n(a)$ for $n \rightarrow \infty$ with $\theta(a) = ab$, $\theta(b) = ba$; see [8].

² The number of overlap-free words of length n is referenced in the On-Line Encyclopedia of Integer Sequences under the code A007777; see [21]. The sequence starts 1, 2, 4, 6, 10, 14, 20, 24, 30, 36, 44, 48, 60, 60, 62, 72,...




Fig. 1. The values of u_n for $1 \leq n \leq 200$ (a) and $\log u_n / \log n$ for $1 \leq n \leq 10000$ (b).

$$\begin{aligned} \alpha &= \sup \left\{ r \mid \exists C > 0, u_n \geq Cn^r \right\}, \\ \beta &= \inf \left\{ r \mid \exists C > 0, u_n \leq Cn^r \right\}, \end{aligned} \quad (1)$$

and showed that $\alpha < \beta$. Cassaigne made a real breakthrough in the study of overlap-free words by characterizing in a constructive way the whole set of overlap-free words. By improving the decomposition theorem of Restivo and Salemi he showed that the numbers u_n can be computed as sums of variables that are obtained by certain linear recurrence relations. These relations are explicitly given in the next section and all numerical values can be found in Appendix A. As a result of this description, the number of overlap-free words of length n can be computed in logarithmic time. For the exponents of growth Cassaigne has also obtained the following bounds: $\alpha < 1.276$ and $\beta > 1.332$. Thus, combining this with the earlier results described above, one has the following inequalities:

$$1.155 < \alpha < 1.276 \quad \text{and} \quad 1.332 < \beta < 1.37. \quad (2)$$

In this paper we develop a linear algebraic approach to study the asymptotic behavior of the number of overlap-free words of length n . Using the results of Cassaigne we show in Theorem 2 that u_n is asymptotically equivalent to the norm of a long product of two particular matrices A_0 and A_1 of dimension 20×20 . This product corresponds to the binary expansion of the number $n-1$. Using this result we express the values of α and β by means of certain joint spectral characteristics of these matrices. We prove that $\alpha = \log_2 \check{\rho}(A_0, A_1)$ and $\beta = \log_2 \hat{\rho}(A_0, A_1)$, where $\check{\rho}$ and $\hat{\rho}$ denote, respectively, the lower spectral radius and the joint spectral radius of the matrices A_0, A_1 (we define these notions in the next section). In Section 3, we estimate these values and we obtain the following improved bounds for α and β :

$$1.2690 < \alpha < 1.2736 \quad \text{and} \quad 1.3322 < \beta < 1.3326. \quad (3)$$

Our estimates are, respectively, within 0.4% and 0.03% of the exact values. In addition, we show in Theorem 3 that the smallest and the largest rates of growth of u_n are effectively attained, and there exist positive constants C_1, C_2 such that $C_1 n^\alpha \leq u_n \leq C_2 n^\beta$ for all $n \in \mathbb{N}$.

Although the sequence u_n does not exhibit an asymptotic polynomial growth, we then show in Theorem 5 that for “almost all” values of n the rate of growth is actually equal to $\sigma = \log_2 \bar{\rho}(A_0, A_1)$, where $\bar{\rho}$ is the Lyapunov exponent of the matrices. For almost all values of n the number of overlap-free words does not grow as n^α , nor as n^β , but in an intermediary way, as n^σ . This means in particular that the value $\frac{\log u_n}{\log n}$ converges to σ as $n \rightarrow \infty$ along a subset of density 1. We obtain the following bounds for the limit σ , which provides an estimation within 0.8% of the exact value:

$$1.3005 < \sigma < 1.3098.$$

These bounds clearly show that $\alpha < \sigma < \beta$.

To compute the exponents α and σ we introduce new efficient algorithms for estimating the lower spectral radius $\check{\rho}$ and the Lyapunov exponent $\bar{\rho}$ of matrices. These algorithms are both of independent interest as they can be applied to arbitrary matrices.

Our linear algebraic approach not only allows us to improve the estimates of the asymptotics of the number of overlap-free words, but also clarifies some aspects of the nature of these words. For instance, we show that the “non purely overlap-free words” used in [8] to compute u_n are asymptotically negligible when considering the total number of overlap-free words.

The paper is organized as follows. In the next section we formulate and prove the main theorems (except for Theorem 2, whose proof is quite technical and is given in Appendix B). Then in Section 3 we present algorithms for estimating the joint spectral radius, the lower spectral radius, and the Lyapunov exponent of linear operators. Applying them to those special matrices we obtain the estimations for α, β and σ . In the appendices we write explicit forms of the matrices and initial vectors used to compute u_n , we give a proof of Theorem 2 and present the results of our numerical algorithms.

2 The asymptotics of the overlap-free words

In the sequel we use the following notation: \mathbb{R}^d is the d -dimensional space, inequalities $x \geq 0$ and $A \geq 0$ mean that all the entries of the vector x (respectively, of the matrix A) are nonnegative. We denote $\mathbb{R}_+^d = \{x \in \mathbb{R}^d, x \geq 0\}$, by $|x|$ we denote a norm of the vector $x \in \mathbb{R}^d$, and by $\|\cdot\|$ any matrix norm. In particular, $|x|_1 = \sum_{i=1}^d |x_i|$, $\|A\|_1 = \sup_{|x|_1=1} |Ax| = \max_{j=1,\dots,d} \sum_{i=1}^d |A_{ij}|$. We write $\mathbf{1}$ for the vector $(1, \dots, 1)^T \in \mathbb{R}^d$, $\rho(A)$ for the spectral radius of the matrix A , that is, the largest magnitude of its eigenvalues. If $A \geq 0$, then there is a vector $v \geq 0$ such that $Av = \rho(A)v$ (the so-called Perron-Frobenius eigenvector). For two functions f_1, f_2 from a set Y to \mathbb{R}_+ the relation $f_1(y) \asymp f_2(y)$ means that there are positive constants C_1, C_2 such that $C_1 f_1(y) \leq f_2(y) \leq C_2 f_1(y)$ for all $y \in Y$.

To compute the number u_n of overlap-free words of length n we use several results from [8] that we summarize in the following theorem:

Theorem 1 *Let $F_0, F_1 \in \mathbb{R}^{30 \times 30}$, and let $w, y_3, \dots, y_{15} \in \mathbb{R}_+^{30}$ be as given in Appendix A. For $n \geq 16$, let y_n be the solution of the following recurrence equations*

$$\begin{aligned} y_{2n} &= F_0 y_n \\ y_{2n+1} &= F_1 y_n. \end{aligned} \tag{4}$$

Then, for any $n \geq 16$, the number of overlap-free words of length n is equal to $w^T y_{n-1}$.

It follows from this result that the number u_n of overlap-free words of length $n \geq 16$ can be obtained by first computing the binary expansion $d_k \cdots d_1$ of $n - 1$, i.e., $n - 1 = \sum_{j=0}^{k-1} d_{j+1} 2^j$, and then defining

$$u_n = w^T F_{d_1} \cdots F_{d_{k-4}} y_m \tag{5}$$

where $m = d_{k-3} + d_{k-2} 2 + d_{k-1} 2^2 + d_k 2^3$. To arrive at the results summarized in Theorem 1, Cassaigne builds a system of recurrence equations allowing the computation of a vector U_n whose entries are the number of overlap-free words of certain types (there are 16 different types). These recurrence equations also involve the recursive computation of a vector V_n that counts other words of length n , the so-called ‘‘single overlaps’’. The single overlap words are not overlap-free, but have to be computed, as they generate overlap-free words of larger lengths. We now present the main result of this section which improves the above theorem in two directions. First we reduce the dimension of the matrices from 30 to 20, and second we prove that u_n is given asymptotically by the norm of a matrix product. The reduction of the dimension to 20 has a straightforward interpretation: when computing the asymptotic growth of

the number of overlap-free words, one can neglect the number of “single overlaps” V_n defined by Cassaigne. We call the remaining words *purely overlap-free words*, as they can be entirely decomposed in a sequence of overlap-free words via Cassaigne’s decomposition (see [8] for more details).

Theorem 2 *Let $A_0, A_1 \in \mathbb{R}_+^{20 \times 20}$ be the matrices defined in Appendix A (Equation A.3), let $\|\cdot\|$ be a matrix norm, and let $A(n) : \mathbb{N} \rightarrow \mathbb{R}_+^{20 \times 20}$ be defined as $A(n) = A_{d_1} \cdots A_{d_k}$ with $d_k \dots d_1$ the binary expansion of $n - 1$. Then,*

$$u_n \asymp \|A(n)\|. \quad (6)$$

Observe that the matrices F_0, F_1 in Theorem 1 are both nonnegative and hence possess a common invariant cone $K = \mathbb{R}_+^{30}$. We say that a cone K is invariant for a linear operator B if $BK \subset K$. All cones are assumed to be solid, convex, closed, and pointed. We start with the following simple result proved in [18].

Lemma 1 *For any cone $K \subset \mathbb{R}^d$, for any norm $|\cdot|$ in \mathbb{R}^d and any matrix norm $\|\cdot\|$ there is a homogeneous continuous function $\gamma : K \rightarrow \mathbb{R}_+$ positive on $\text{int}K$ such that for any $x \in \text{int}K$ and for any matrix B that leaves K invariant one has*

$$\gamma(x)\|B\| \cdot |x| \leq |Bx| \leq \frac{1}{\gamma(x)}\|B\| \cdot |x|.$$

Corollary 1 *Let two matrices A_0, A_1 possess an invariant cone $K \subset \mathbb{R}^d$. Then for any $x \in \text{int}K$ we have $|A_{d_1} \cdots A_{d_k}x| \asymp \|A_{d_1} \cdots A_{d_k}\|$ for all k and for all indices $d_1, \dots, d_k \in \{0, 1\}$.*

In view of Corollary 1 and of Eq. (5), Theorem 2 may seem obvious, at least if we consider the matrices F_i instead of A_i . One can however not directly apply Lemma 1 and Corollary 1 to the matrices A_0, A_1 or to the matrices F_0, F_1 because the vector corresponding to x is not in the interior of the positive orthant, which is an invariant cone of these matrices. To prove Theorem 2 we construct a wider invariant cone of A_0 and A_1 by using special properties of these matrices. We detail the construction of this cone in the proof given in Appendix B. Theorem 2 allows us to express the rates of growth of the sequence u_n in terms of norms of products of the matrices A_0, A_1 and then to use joint spectral characteristics of these matrices to estimate the rates of growth. More explicitly, Theorem 2 yields the following corollary:

Corollary 2 *Let $A_0, A_1 \in \mathbb{R}_+^{20 \times 20}$ be the matrices defined in Appendix A and let $A(n) : \mathbb{N} \rightarrow \mathbb{R}_+^{20 \times 20}$ be defined as $A(n) = A_{d_1} \cdots A_{d_k}$ with $d_k \dots d_1$ the binary expansion of $n - 1$. Then*

$$\frac{\log u_n}{\log n} - \log \|A(n)\|^{1/k} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (7)$$

Proof. Since $\frac{\log_2 n}{k} \rightarrow 1$ as $n \rightarrow \infty$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \left(\frac{\log_2 u_n}{\log_2 n} - \frac{\log_2 \|A_{d_1} \cdots A_{d_k}\|}{k} \right) &= \\ \lim_{n \rightarrow \infty} \frac{\log_2 u_n - \log_2 \|A_{d_1} \cdots A_{d_k}\|}{k} &= \lim_{n \rightarrow \infty} \frac{\log_2 (u_n \cdot \|A_{d_1} \cdots A_{d_k}\|^{-1})}{k}. \end{aligned}$$

By Theorem 2 the value $\log_2 (u_n \cdot \|A_{d_1} \cdots A_{d_k}\|^{-1})$ is bounded uniformly over $n \in \mathbb{N}$, hence it tends to zero, being divided by k . \square

We first analyze the smallest and the largest exponents of growth α and β defined in Eq. (1). For a given set of matrices $\Sigma = \{A_1, \dots, A_m\}$ we denote by $\check{\rho}$ and $\hat{\rho}$ its lower spectral radius and its joint spectral radius:

$$\begin{aligned} \check{\rho}(\Sigma) &= \lim_{k \rightarrow \infty} \min_{d_1, \dots, d_k \in \{1, \dots, m\}} \|A_{d_1} \cdots A_{d_k}\|^{1/k}, \\ \hat{\rho}(\Sigma) &= \lim_{k \rightarrow \infty} \max_{d_1, \dots, d_k \in \{1, \dots, m\}} \|A_{d_1} \cdots A_{d_k}\|^{1/k}. \end{aligned} \quad (8)$$

Both limits are well-defined and do not depend on the chosen norm. Moreover, for any product $A_{d_1} \cdots A_{d_k}$ we have

$$\check{\rho} \leq \rho(A_{d_1} \cdots A_{d_k})^{1/k} \leq \hat{\rho} \quad (9)$$

(see [16, 5] for surveys on these notions).

Theorem 3 For $k \geq 1$, let $\alpha_k = \min_{2^{k-1} < n \leq 2^k} \frac{\log u_n}{\log n}$ and $\beta_k = \max_{2^{k-1} < n \leq 2^k} \frac{\log u_n}{\log n}$.

Then

$$\alpha = \lim_{k \rightarrow \infty} \alpha_k = \log_2 \check{\rho}(A_0, A_1) \quad \text{and} \quad \beta = \lim_{k \rightarrow \infty} \beta_k = \log_2 \hat{\rho}(A_0, A_1), \quad (10)$$

where the matrices A_0, A_1 are defined in Appendix A. Moreover, there are positive constants C_1, C_2 such that

$$C_1 \leq \min_{2^{k-1} < n \leq 2^k} u_n n^{-\alpha} \quad \text{and} \quad C_1 \leq \max_{2^{k-1} < n \leq 2^k} u_n n^{-\beta} \leq C_2 \quad (11)$$

for all $k \in \mathbb{N}$.

The proof of this theorem is based on the following auxiliary result taken from [18]. For a given set of indices $\{i_1, \dots, i_p\} \subset \{1, \dots, d\}$, $p \geq 1$ we call the subspace $L_{i_1, \dots, i_p} = \{x \in \mathbb{R}^d, x_{i_1} = \dots = x_{i_p} = 0\}$ a *coordinate plane*.

Proposition 1 [18] Let A_0, A_1 be matrices with a common invariant cone. Then there is a positive constant c_1 such that

$$\max_{d_1, \dots, d_k} \|A_{d_1} \cdots A_{d_k}\| \geq c_1 \hat{\rho}^k \quad \text{and} \quad \min_{d_1, \dots, d_k} \|A_{d_1} \cdots A_{d_k}\| \geq c_1 \check{\rho}^k, \quad k \in \mathbb{N}.$$

If, moreover, these matrices have no common invariant subspace among the coordinate planes, then there is a positive constant c_2 such that

$$\max_{d_1, \dots, d_k} \|A_{d_1} \cdots A_{d_k}\| \leq c_2 \hat{\rho}^k, \quad k \in \mathbb{N}.$$

Proof of Theorem 3. The equalities in Eq. (10) follow immediately from Corollary 2 and the definitions given on Eq. (8). To prove the inequalities given at Eq. (11) we apply Proposition 1 for our matrices A_0, A_1 that have an invariant cone \mathbb{R}_+^{20} . Theorem 2 yields

$$u_n n^{-\alpha} \asymp \|A_{d_1} \cdots A_{d_k}\| 2^{-\alpha k} = \|A_{d_1} \cdots A_{d_k}\| \check{\rho}^{-k}.$$

Taking the minimum over $n = 2^{k-1} + 1, \dots, 2^k$ and invoking Proposition 1 we conclude that $\min_{2^{k-1} < n \leq 2^k} u_n n^{-\alpha} \geq C_1$. The same with the inequality $\max_{2^{k-1} < n \leq 2^k} u_n n^{-\beta} \geq C_1$. To prove the upper bound in Eq. (11) we note that the matrices A_0, A_1 have no common invariant subspaces among the coordinate planes (to see this observe, for instance, that $(A_0 + A_1)^5$ has no zero entry). \square

Corollary 3 *There are positive constants C_1, C_2 such that*

$$C_1 n^\alpha \leq u_n \leq C_2 n^\beta, \quad n \in \mathbb{N}.$$

In the next section we will see that $\alpha < \beta$. In particular, the sequence u_n does not have a constant rate of growth, and the value $\frac{\log u_n}{\log n}$ does not converge as $n \rightarrow \infty$. This was already noted by Cassaigne in [8]. Nevertheless, it appears that the value $\frac{\log u_n}{\log n}$ actually has a limit as $n \rightarrow \infty$, not along all the natural numbers $n \in \mathbb{N}$, but along a subsequence of \mathbb{N} of density 1. In other terms, the sequence converges with probability 1. The limit, which differs from both α and β can be expressed by the so-called Lyapunov exponent $\bar{\rho}$ of the matrices A_0, A_1 . To show this we apply the following result proved by Oseledets in 1968. For the sake of simplicity we formulate it for two matrices, although it can be easily generalized to any finite set of matrices.

Theorem 4 [14] *Let A_0, A_1 be arbitrary matrices and d_1, d_2, \dots be a sequence of independent random variables that take values 0 and 1 with equal probabilities 1/2. Then the value $\|A_{d_1} \cdots A_{d_k}\|^{1/k}$ converges to some number $\bar{\rho}$ with probability 1. This means that for any $\varepsilon > 0$ we have $P\left(\left|\|A_{d_1} \cdots A_{d_k}\|^{1/k} - \bar{\rho}\right| > \varepsilon\right) \rightarrow 0$ as $k \rightarrow \infty$.*

The limit $\bar{\rho}$ in Theorem 4 is called the *Lyapunov exponent* of the set $\{A_0, A_1\}$. This value is given by the following formula:

$$\bar{\rho}(A_0, A_1) = \lim_{k \rightarrow \infty} \left(\prod_{d_1, \dots, d_k} \|A_{d_1} \cdots A_{d_k}\|^{1/k} \right)^{1/2^k} \quad (12)$$

(for the proof see, for instance, [19]). To understand what this gives for the asymptotics of our sequence u_n we introduce some further notation. Let \mathcal{P} be some property of natural numbers. For a given $k \in \mathbb{N}$ we denote

$$P_k(\mathcal{P}) = 2^{-(k-1)} \text{Card}\{n \in \{2^{k-1} + 1, \dots, 2^k\}, \quad n \text{ satisfies } \mathcal{P}\}.$$

Thus, P_k is the probability that the integer n uniformly distributed on the set $\{2^{k-1} + 1, \dots, 2^k\}$ satisfies \mathcal{P} . Combining Proposition 2 and Theorem 4 we obtain

Theorem 5 *There is a number σ such that for any $\varepsilon > 0$ we have*

$$P_k\left(\left|\frac{\log u_n}{\log n} - \sigma\right| > \varepsilon\right) \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Moreover, $\sigma = \log_2 \bar{\rho}$, where $\bar{\rho}$ is the Lyapunov exponent of the matrices $\{A_0, A_1\}$ defined in Appendix A.

Thus, for almost all numbers $n \in \mathbb{N}$ the number of overlap-free words u_n has the same exponent of growth $\sigma = \log_2 \bar{\rho}$. If positive a and b are large enough and $a < b$, then for a number n taken randomly from the segment $[a, b]$ the value $\frac{\log u_n}{\log n}$ is close to σ . Let us recall that a subset $\mathcal{A} \subset \mathbb{N}$ is said to have density 1 if $\frac{1}{n} \text{Card}\{r \leq n, r \in \mathcal{A}\} \rightarrow 1$ as $n \rightarrow \infty$. We say that a sequence f_n converges to a number f along a set of density 1 if there is a set $\mathcal{A} \subset \mathbb{N}$ of density 1 such that $\lim_{n \rightarrow \infty, n \in \mathcal{A}} f_n = f$. Theorem 5 yields

Corollary 4 *The value $\frac{\log u_n}{\log n}$ converges to σ along a set of density 1.*

Proof. By Theorem 5 there exists $k_1 \in \mathbb{N}$ such that at least half of the natural numbers $n \leq 2^{k_1}$ satisfy the inequality $\left|\frac{\log u_n}{\log n} - \sigma\right| < \frac{1}{2}$. Denote the set of such numbers $n \leq 2^{k_1}$ by \mathcal{A}_1 . Further, there exists $k_2 > k_1$ such that at least $\frac{3}{4}$ of the numbers $n \in \{2^{k_1} + 1, \dots, 2^{k_2}\}$ satisfy the inequality $\left|\frac{\log u_n}{\log n} - \sigma\right| < \frac{1}{4}$. Denote the set of such numbers by \mathcal{A}_2 . The remainder is by induction: having a number k_{r-1} we take a number $k_r > k_{r-1}$ such that at least $\frac{2^r - 1}{2^r}$ of the numbers $n \in \{2^{k_{r-1}} + 1, \dots, 2^{k_r}\}$ satisfy the inequality $\left|\frac{\log u_n}{\log n} - \sigma\right| < \frac{1}{2^r}$ and denote the set of such numbers by \mathcal{A}_r . The set $\mathcal{A} = \cup_{r \in \mathbb{N}} \mathcal{A}_r$ has density 1 and $\frac{\log u_n}{\log n}$ tends to σ along \mathcal{A} as $n \rightarrow \infty$. \square

3 Estimations of the exponents

Theorems 2 and 5 reduce the problem of estimating the exponents of growth of u_n to computing joint spectral characteristics of the matrices A_0 and A_1 . In order to estimate the joint spectral radius we use a modified version of the “ellipsoidal norm algorithm” [3]. For the lower spectral radius and for the Lyapunov exponent we present new algorithms, which seem to be relatively efficient for nonnegative matrices. The results we obtain can be summarized in the following theorem:

Theorem 6

$$\begin{aligned} 1.2690 < \alpha < 1.2736 \\ 1.3322 < \beta < 1.3326 \\ 1.3005 < \sigma < 1.3098 \end{aligned} \tag{13}$$

In this section we also make (and give arguments for) the following conjecture:

Conjecture 1

$$\beta = \log_2 \sqrt{\rho(A_0 A_1)} = 1.3322 \dots$$

3.1 Estimation of β and the joint spectral radius

By Theorem 3 to estimate the exponent β one needs to estimate the joint spectral radius of the set $\{A_0, A_1\}$. A lower bound for $\hat{\rho}$ can be obtained by applying inequality (9). Taking $k = 2$ and $d_1 = 0, d_2 = 1$ we get

$$\hat{\rho} \geq \left[\rho(A_0 A_1) \right]^{1/2} = 2.5179, \tag{14}$$

and so $\beta > \log_2 2.5179 > 1.3322$ (this lower bound was already found in [8]). Upper bounds for the joint spectral radius of sets of matrices $\Sigma = \{A_1, \dots, A_m\}$ are usually derived from the following simple inequality

$$\hat{\rho} \leq \max_{d_1, \dots, d_k \in \{1, \dots, m\}} \|A_{d_1} \cdots A_{d_k}\|^{1/k}, \tag{15}$$

which holds for every $k \geq 1$ and converges to $\hat{\rho}$ as $k \rightarrow \infty$. This, at least theoretically, gives arbitrarily sharp estimations for $\hat{\rho}$. However, in our case, due to the size of the matrices A_0, A_1 , this method leads to computations that are too expensive even for relatively small values of k . Faster convergence

can be achieved by finding an appropriate norm. To do this we use the so-called ellipsoidal norm: $\|A\|_P = \max_x \sqrt{\frac{x^T A^T P A x}{x^T A x}}$, where P is a positive definite matrix. This is the matrix norm induced by the vector norm $|x|_P = (x^T P x)^{1/2}$. The crucial idea is that the optimal P , for which the right hand side in (15) for $k = 1$ is minimal, can be found by solving a simple semidefinite programming problem. This algorithm can be iterated using the relation $\rho(\Sigma^k) = \rho(\Sigma)^k$. In the sequel we denote $\Sigma^k = \{A_{d_1} \cdots A_{d_k}, 1 \leq d_i \leq m, i = 1, \dots, k\}$. Thus one can consider the set Σ^k as a new set of matrices, and approximate its joint spectral radius with the best possible ellipsoidal norm. In Appendix C we give an ellipsoidal norm such that each matrix in Σ^{14} has a norm smaller than 2.5186^{14} . This implies that $\hat{\rho} \leq 2.5186$, which gives $\beta < 1.3326$. Combining this with the inequality $\beta > 1.3322$ we complete the proof of the bounds for β in Theorem 6.

We have not been able to improve the lower bound of Eq. (14). However, the upper bound we obtain is very close to this lower bound, and the upper bounds obtained with an ellipsoidal norm for Σ^k get closer and closer to this value when k increases. Moreover, it has already been observed that for many sets of matrices for which the joint spectral radius is known exactly, and in particular matrices with nonnegative integer entries, there always is a product that achieves the joint spectral radius, i.e., a product $A \in \Sigma^t$ such that $\hat{\rho} = \rho(A)^{(1/t)}$ [9]. For these reasons, we conjecture that the exponent β is actually equal to the lower bound.

3.2 Estimation of α and the lower spectral radius

An upper bound for $\check{\rho}(A_0, A_1)$ can be obtained using Eq.(9) for $k = 1$ and $d_1 = 0$. We have

$$\alpha = \log_2(\check{\rho}) \leq \log_2(A_0) = 1.276\dots \quad (16)$$

This bound for α was first derived in [8]. This bound is however not optimal. Taking the product $A_1^{10}A_0$ (i.e., $k = 11$ in inequality (9)), we get a better estimate:

$$\alpha \leq \log_2 \left[(\rho(A_1^{10}A_0))^{1/11} \right] = 1.2735\dots \quad (17)$$

One can verify numerically that this product gives the best possible upper bound among all the matrix products of length $k \leq 14$.

We now estimate α from below. The problem of approximating the lower spectral radius is NP-hard [4] and to the best of our knowledge, no algorithm is known to compute $\check{\rho}$. Here we propose two new algorithms. We first consider nonnegative matrices. As we observed above, for any k we have $\check{\rho}(\Sigma^k) = \check{\rho}(\Sigma)^k$. Without loss of generality it can be assumed that the matrices of the set Σ do not have a common zero column. Otherwise, by suppressing this column and the corresponding row we obtain a set of matrices of smaller dimension with

the same lower spectral radius. The vector of ones is denoted by $\mathbf{1}$.

Theorem 7 *Let Σ be a set of nonnegative matrices that do not have any common zero column. If for some $r \in \mathbb{R}^+$, $s \leq t \in \mathbb{N}$, there exists $x \in \mathbb{R}^d$ satisfying the following system of linear inequalities*

$$\begin{aligned} B(Ax - rx) &\geq 0, \quad \forall B \in \Sigma^s, A \in \Sigma^t, \\ x &\geq 0, \quad (x, \mathbf{1}) = 1, \end{aligned} \tag{18}$$

then $\check{\rho}(\Sigma) \geq r^{1/t}$.

Proof. Let x be a solution of (18). Let us consider a product of matrices $A_k \dots A_1 \in \Sigma^{kt} : A_i \in \Sigma^t$. We show by induction in k that $A_k \dots A_1 x \geq r^{k-1} A_k x$: For $k = 2$, we have $A_2(A_1 x - rx) = CB(A_1 x - rx) \geq 0$, with $B \in \Sigma^s, C \in \Sigma^{t-s}$. For $k > 2$ we have $A_k \dots A_1 x = A_k A_{k-1} \dots A_1 x \geq r^{k-2} A_k A_{k-1} x \geq r^{k-1} A_k x$. In the last inequality the case for $k = 2$ was reused. Hence, $\|A_k \dots A_1\| = \mathbf{1}^T A_k \dots A_1 \mathbf{1} \geq r^{k-1} \mathbf{1}^T A_k x \geq r^k C$, where $C = (\min_k \mathbf{1}^T A_k x)/r > 0$. The last inequality holds because $A_k x = 0$, together with the first inequality in (18), imply that $-rBx = 0$ for all $B \in \Sigma^s$, which implies that all $B \in \Sigma^s$ have a common zero column. This is in contradiction with our assumption because the matrices in Σ^s share a common zero column if and only if the matrices in Σ do. \square

We were able to find a solution to the linear programming problem (18) with $r = 2.41$, $t = 16$, $s = 6$. Hence we get the following lower bound: $\alpha \geq \log r > 1.2690$. The corresponding vector x is given in Appendix D. This completes the proof of Theorem 6.

Theorem 7 handles nonnegative matrices, and we propose now a way to generalize this result to arbitrary real matrices. The idea is to lift the matrices to a larger vector space, so that all the matrices share an invariant cone. This kind of lifting is rather classical and is known under several names in the literature as for instance semidefinite lifting or symmetric algebras [2, 17, 15]. The idea is to consider the matrices $A_i \in \Sigma$ as linear operators acting on the cone of positive semidefinite matrices S as $S \rightarrow A_i^T S A_i$. It is not difficult to prove that the lower spectral radius of this new set of linear operators is equal to $\check{\rho}(\Sigma)^2$. We use the notation $A \succeq B$ to denote that the matrix $A - B$ is positive semidefinite. Recall that $A \succeq 0 \Leftrightarrow \forall y, y^T A y \geq 0$.

Theorem 8 *Let Σ be a set of matrices in $\mathbb{R}^{d \times d}$ and $s \leq t \in \mathbb{N}$. Suppose that there are $r > 0$ and a symmetric matrix $S \succeq 0$ such that*

$$\begin{aligned} B^*(A^* S A - rS)B &\succeq 0 \quad \forall A \in \Sigma^t, B \in \Sigma^s \\ S &\succ 0 \end{aligned} \tag{19}$$

then $\check{\rho}(\Sigma) \geq r^{1/2t}$.

Proof. The proof is formally similar to the previous one: Let S be a solution of (19). We denote M_k the product $A_1 \dots A_k$. It is easy to show by induction that $M_k^* S M_k \succeq r^{k-1} (A_k^* S A_k)$. This is obvious for $k = 2$ for similar reasons as in the previous theorem, and for $k > 2$, if, by induction,

$$\forall y, \quad y^* M_{k-1}^* S M_{k-1} y \geq r^{k-2} y^* A_{k-1}^* S A_{k-1} y,$$

then, with $y = A_k x$, for all x ,

$$x^* M_k^* S M_k x \geq r^{k-2} x^* A_k^* A_{k-1}^* S A_{k-1} A_k x \geq r^{k-1} x^* A_k^* S A_k x.$$

Thus,

$$\sup \left\{ \frac{x^* M_k^* S M_k x}{x^* S x} \right\} \geq r^{k-1} \sup \left\{ \frac{x^* A_k^* S A_k x}{x^* S x} \right\}.$$

Finally, $\|M_k\|_S \geq r^{k/2} C$, where C is a constant. \square

For a given $r > 0$ the existence of a solution S can be established by solving the semidefinite programming problem (19), and the optimal r can be found by bisection in logarithmic time.

3.3 Estimation of σ and the Lyapunov exponent

The exponent of the average growth σ is obviously between α and β , so $1.2690 < \sigma < 1.3326$. To get better bounds we need to estimate the Lyapunov exponent $\bar{\rho}$ of the matrices A_0, A_1 . The first upper bound can be given by the so-called 1-radius ρ_1 :

$$\rho_1 = \lim_{k \rightarrow \infty} \left(2^{-k} \sum_{d_1, \dots, d_k} \|A_{d_1} \cdots A_{d_k}\| \right)^{1/k}.$$

For matrices with a common invariant cone we have $\rho_1 = \frac{1}{2} \rho(A_0 + A_1)$ [18]. Therefore, in our case $\rho_1 = \frac{1}{2} \rho(A_0 + A_1) = 2.479\dots$. This exponent was first computed in [8], where it was shown that the value $\sum_{j=0}^{n-1} u_j$ is equivalent to n^η , where $\eta = 1 + \log_2 \rho_1 = 2.310\dots$. It follows immediately from the inequality between the arithmetic mean and the geometric mean that $\bar{\rho} \leq \rho_1$. Thus, $\sigma \leq \eta$. In fact, as we show below, σ is strictly smaller than η . We are not aware of any approximation algorithm for the Lyapunov exponent, except by application of Definition (12). It is easily seen that for any k the value $r_k = \left(\prod_{d_1, \dots, d_k} \|A_{d_1} \cdots A_{d_k}\| \right)^{\frac{1}{k2^k}}$ gives an upper bound for $\bar{\rho}$, that is $\bar{\rho} \leq r_k$ for any $k \in \mathbb{N}$. Since $r_k \rightarrow \bar{\rho}$ as $k \rightarrow \infty$, we see that this estimation can be arbitrarily sharp for large k . But for the dimension 20 this leads to extensive numerical computations. For example, for the norm $\|\cdot\|_1$ we have $r_{20} = 2.4865$, which is

even larger than ρ_1 . In order to obtain a better bound for $\bar{\rho}$ we state the following results. For any k and $x \in \mathbb{R}^d$ we denote $p_k(x) = \left(\prod_{d_1, \dots, d_k} |A_{d_1} \cdots A_{d_k} x| \right)^{\frac{1}{2^k}}$ and $m_k = \sup_{x \geq 0, |x|=1} p_k(x)$.

Proposition 2 *Let A_0, A_1 be nonnegative matrices in \mathbb{R}^d . Then for any norm $|\cdot|$ and for any $k \geq 1$ we have $\bar{\rho} \leq (m_k)^{1/k}$.*

Proof. For any $x \geq 0$ and for any $n, k \in \mathbb{N}$ we have $p_{k+n}(x) \leq m_k p_n(x)$. Therefore, $p_{tk}(x) \leq (m_k)^t$. Whence, $\lim_{t \rightarrow \infty} [p_{tk}(x)]^{1/tk} \leq (m_k^t)^{1/tk} = (m_k)^{1/k}$. On the other hand, by Corollary 1 for $x > 0$ we have $r_n \asymp [p_n(x)]^{1/n}$, and consequently $\lim_{t \rightarrow \infty} [p_{tk}(x)]^{1/tk} \rightarrow \bar{\rho}$ as $t \rightarrow \infty$. Thus, $\bar{\rho} \leq (m_k)^{1/k}$. \square

Proposition 3 *Let A_0, A_1 be nonnegative matrices in \mathbb{R}^d that do not have common invariant subspaces among the coordinate planes. If $\check{\rho} < \hat{\rho}$, then $\bar{\rho} < \rho_1$.*

Proof. Let v_* be the eigenvector of the matrix $\frac{1}{2}(A_0^* + A_1^*)$ corresponding to its largest eigenvalue ρ_1 . Since the matrices have no common invariant coordinate planes, it follows that $v_* > 0$. Consider the norm $|x| = (x, v_*)$ on \mathbb{R}_+^d . Take some $k \geq 1$ and $y \in \mathbb{R}_+^d$, $|y| = (y, v_*) = 1$, such that $p_k(y) = m_k$. We have

$$\begin{aligned} m_k = p_k(y) &\leq 2^{-k} \sum_{d_1, \dots, d_k} |A_{d_1} \cdots A_{d_k} y| = 2^{-k} \sum_{d_1, \dots, d_k} (A_{d_1} \cdots A_{d_k} y, v_*) \\ &= \left(y, 2^{-k} (A_0^* + A_1^*)^k v_* \right) = \rho_1^k (y, v_*) = \rho_1^k. \end{aligned}$$

Thus, $m_k \leq \rho_1^k$, and the equality is possible only if all 2^k values $|A_{d_1} \cdots A_{d_k} y|$ are equal. Since $\check{\rho} < \hat{\rho}$, there must be a k such that the inequality is strict. Thus, $m_k < \rho_1^k$ for some k , and by Proposition 2 we have $\bar{\rho} \leq (m_k)^{1/k} < \rho_1$. \square

We are now able to estimate $\bar{\rho}$ for the matrices A_0, A_1 . For the norm $|x| = (x, v_*)$ used in the proof of Proposition 3 the value $-\frac{1}{k} \log_2 m_k$ can be found as the solution of the following convex minimization problem with linear constraints:

$$\begin{aligned} \min & -\frac{1}{k 2^k \ln 2} \sum_{d_1, \dots, d_k \in \{0,1\}} \ln \left(x, A_{d_1}^* \cdots A_{d_k}^* v_* \right) \\ \text{s.t.} & x \geq 0, \quad (x, v_*) = 1. \end{aligned} \tag{20}$$

The optimal value of this optimization problem is equal to $-\frac{1}{k} \log_2 m_k$, which gives an upper bound for $\sigma = \log_2 \bar{\rho}$ (Proposition 2). Solving this problem for $k = 12$ we obtain $\sigma \leq 1.3098$. We finally provide a theorem that allows us to derive a lower bound on σ . The idea is identical to the one used in Theorem 7,

but transposed to the Lyapunov exponent.

Theorem 9 *Let Σ be a set of nonnegative matrices that do not have any common zero column. If for some $r_i \in \mathbb{R}_+$, $s \leq t \in \mathbb{N}$, there exists $x \in \mathbb{R}_+^d$ satisfying the following system of linear inequalities*

$$\begin{aligned} B(A_i x - r_i x) &\geq 0, & \forall B \in \Sigma^s, A_i \in \Sigma^t, \\ x &\geq 0, & (x, \mathbf{1}) = 1, \end{aligned} \tag{21}$$

then $\bar{\rho}(\Sigma) \geq \prod_i r_i^{1/(t2^t)}$.

The proof is similar to the proof of Theorem 7 and is left to the reader. Also, a similar theorem can be stated for general matrices (with negative entries), but involving linear matrix inequalities. Due to the number of different variables r_i , one cannot hope to find the optimal x with SDP and bisection techniques. However, by using the vector x computed for approximating the lower spectral radius (given in Appendix D), with the values $s = 8, t = 16$ for the parameters, one gets a good lower bound for σ : $\sigma \geq 1.3005$.

4 Conclusions

The goal of this paper is to precisely characterize the asymptotic rate of growth of the number of overlap-free words. Based on Cassaigne’s description of these words with products of matrices, we first prove that these matrices can be simplified, by decreasing the state space dimension from 30 to 20. This improvement is not only useful for numerical computations, but allows to characterize the overlap-free words that “count” for the asymptotics: we call these words *purely overlap free*, as they can be expressed iteratively as the image of shorter purely overlap free words.

We have then proved that the lower and upper exponents α and β defined by Cassaigne are effectively reached for an infinite number of lengths, and we have characterized them respectively as the logarithms of the *lower spectral radius* and the *joint spectral radius* of the simplified matrices that we constructed. This characterization, combined with new algorithms that we propose to approximate the lower spectral radius, allow us to compute them within 0.4%. The algorithms we propose can of course be used to reach any degree of accuracy for β (this seems also to be the case for α and σ , but no theoretical result is known for the approximation of the lower spectral radius). The computational results we report in this paper have all been obtained in a few minutes of computation time on a standard PC desktop and can therefore easily be improved.

Finally we have shown that for almost all values of n , the number of overlap-free words of length n do not grow as n^α , nor as n^β , but in an intermediary way as n^σ , and we have provided sharp bounds for this value of σ .

This work opens obvious questions: Can joint spectral characteristics be used to describe the rate of growth of other languages, such as for instance the more general repetition free languages ? The generalization does not seem to be straightforward for several reasons: first, the somewhat technical proofs of the links between u_n and the norm of a corresponding matrix product take into account the very structure of these particular matrices, and second, it is known that a bifurcation occurs for the growth of repetition-free words: for some members of this class of languages the growth is polynomial, as for overlap-free words, but for some others the growth is exponential [10], and one could wonder how the joint spectral characteristics developed in this paper could represent both kinds of growth.

Acknowledgment

We would like to thank Prof. Stephen Boyd (Stanford University), Yuri Nesterov, and François Glineur (Université catholique de Louvain) for their helpful suggestions on semi-definite programming techniques. This research was carried out during the visit of the second author to the Université catholique de Louvain (Louvain-la-Neuve, Belgium). That author is grateful to the university for its hospitality.

References

- [1] J. Berstel. Growth of repetition-free words—a review. *Theoretical Computer Science*, 340(2):280–290, 2005.
- [2] V. D. Blondel and Y. Nesterov. Computationally efficient approximations of the joint spectral radius. *SIAM Journal of Matrix Analysis*, 27(1):256–272, 2005.
- [3] V. D. Blondel, Y. Nesterov, and J. Theys. On the accuracy of the ellipsoid norm approximation of the joint spectral radius. *Linear Algebra and its Applications*, 394(1):91–107, 2005.
- [4] V. D. Blondel and J. N. Tsitsiklis. The lyapunov exponent and joint spectral radius of pairs of matrices are hard - when not impossible - to compute and to approximate. *Mathematics of Control, Signals, and Systems*, 10:31–40, 1997.
- [5] V. D. Blondel and J. N. Tsitsiklis. A survey of computational complexity results in systems and control. *Automatica*, 36(9):1249–1274, 2000.

- [6] S. Brlek. Enumeration of factors in the thue-morse word. *Discrete Applied Mathematics*, 24:83–96, 1989.
- [7] Arturo Carpi. Overlap-free words and finite automata. *Theoretical Computer Science*, 115(2):243–260, 1993.
- [8] J. Cassaigne. Counting overlap-free binary words. *STACS 93, Lecture Notes in Computer Science*, 665:216–225, 1993.
- [9] R. M. Jungers and V. D. Blondel. On the finiteness conjecture for rational matrices. To appear in *Linear Algebra and its Applications*, doi:10.1016/j.laa.2007.07.007, 2007.
- [10] Juhani Karhumäki and Jeffrey Shallit. Polynomial versus exponential growth in repetition-free binary words. *Journal of Combinatorial Theory Series A*, 105(2):335–347, 2004.
- [11] A. J. Kfoury. A linear time algorithm to decide whether a binary word contains an overlap. *Theoretical Informatics and Applications*, 22:135–145, 1988.
- [12] Y. Kobayashi. Enumeration of irreducible binary words. *Discrete Applied Mathematics*, 20:221–232, 1988.
- [13] A. Iepistö. A characterization of 2+-free words over a binary alphabet, Master Thesis, university of turku, finland, 1995.
- [14] V. I. Oseledets. A multiplicative ergodic theorem. Lyapunov characteristic numbers for dynamical systems. *Transactions of the Moscow Mathematical Society*, 19:197–231, 1968.
- [15] P. Parrilo and A. Jadbabaie. Approximation of the joint spectral radius of a set of matrices using sum of squares. 2007. To appear in Alberto Bemporad, Antonio Bicchi and Giorgio Buttazzo (Editors), *Hybrid Systems: Computation and Control* Springer Lecture Notes in Computer Science.
- [16] V. Y. Protasov. The joint spectral radius and invariant sets of linear operators. *Fundamentalnaya i prikladnaya matematika*, 2(1):205–231, 1996.
- [17] V. Y. Protasov. The generalized spectral radius. a geometric approach. *Izvestiya Matematika*, 61(5):995–1030, 1997.
- [18] V. Y. Protasov. On the asymptotics of the partition function. *Sbornik Matematika*, 191(3-4):381–414, 2000.
- [19] V. Y. Protasov. On the regularity of de rham curves. *Izvestiya Matematika*, 68(3):27–68, 2004.
- [20] A. Restivo and S. Salemi. Overlap-free words on two symbols. *Lecture Notes in Computer Science, Automata on Infinite Words*, 192:198–206, 1985.
- [21] N. J. A. Sloane. On-line encyclopedia of integer sequences. Url : <http://www.research.att.com/~njas/sequences>.

- [22] A. Thue. Über unendliche zeichenreihen. *Kra. Vidensk. Selsk. Skrifter. I. Mat. Nat. Kl.*, 7:1–22, 1906.
- [23] A. Thue. Über die gegenseitige lage gleicher teile gewisser zeichenreihen. *Kra. Vidensk. Selsk. Skrifter. I. Mat. Nat. Kl.*, 1:1–67, 1912.

A Numerical values

We introduce the following auxiliary matrices. For the sake of simplicity our notation do not follow exactly those of [8].

$$D_1 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 2 & 1 \\ 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, B_1 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, C_1 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 4 & 2 \\ 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$B_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, C_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, C_4 = \begin{pmatrix} 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Now, defining

$$F_0 = \left(\begin{array}{cc|cc} C_1 & 0_{10 \times 10} & C_2 & 0_{10 \times 5} \\ D_1 & B_1 & 0_{10 \times 5} & B_2 \\ \hline 0_{5 \times 10} & 0_{5 \times 10} & C_4 & 0_{5 \times 5} \\ 0_{5 \times 10} & 0_{5 \times 10} & 0_{5 \times 5} & 0_{5 \times 5} \end{array} \right), F_1 = \left(\begin{array}{cc|cc} D_1 & B_1 & 0_{10 \times 5} & B_2 \\ 0_{10 \times 10} & C_1 & 0_{10 \times 5} & C_2 \\ \hline 0_{5 \times 10} & 0_{5 \times 10} & 0_{5 \times 5} & 0_{5 \times 5} \\ 0_{5 \times 10} & 0_{5 \times 10} & 0_{5 \times 5} & C_4 \end{array} \right), \quad (\text{A.1})$$

$$w = (1, 2, 2, 2, 1, 2, 2, 1, 2, 1, 0_{1 \times 20})^T,$$

$$y_3 = (2, 0, 2, 0, 2, 0, 0, 0, 0, 2, 0, 2, 2, 0, 0, 0, 2, 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0)^T,$$

$$y_4 = (0, 2, 2, 0, 0, 0, 2, 2, 0, 0, 0, 0, 0, 0, 0, 2, 2, 0, 2, 2, 2, 0, 2, 0, 2, 0, 0, 0, 0, 0)^T,$$

$$y_5 = (2, 2, 0, 2, 2, 2, 0, 2, 0, 2, 0, 0, 0, 0, 0, 4, 2, 0, 2, 0, 2, 2, 0, 2, 0, 0, 2, 0, 0, 0)^T,$$

$$y_6 = (4, 2, 0, 2, 0, 2, 2, 0, 2, 0, 0, 2, 0, 0, 0, 4, 2, 2, 2, 4, 2, 0, 0, 2, 2, 0, 0, 0, 0, 0)^T,$$

$$y_7 = (4, 2, 2, 2, 4, 2, 0, 0, 2, 2, 0, 0, 0, 0, 0, 4, 4, 4, 2, 0, 2, 2, 0, 2, 0, 0, 0, 0, 0, 2)^T,$$

$$y_8 = (4, 4, 4, 2, 0, 2, 2, 0, 2, 0, 0, 0, 0, 0, 2, 6, 4, 4, 2, 4, 2, 0, 4, 2, 2, 0, 0, 0, 0, 0)^T,$$

$$y_9 = (6, 4, 4, 2, 4, 2, 0, 4, 2, 2, 0, 0, 0, 0, 0, 8, 4, 4, 2, 0, 4, 4, 4, 0, 0, 0, 0, 0, 0, 0)^T,$$

$$y_{10} = (8, 4, 4, 2, 0, 4, 4, 4, 0, 0, 0, 0, 0, 0, 0, 8, 4, 6, 4, 8, 2, 0, 4, 2, 4, 0, 0, 0, 0, 0)^T,$$

$$y_{11} = (8, 4, 6, 4, 8, 2, 0, 4, 2, 4, 0, 0, 0, 0, 0, 8, 6, 6, 2, 0, 2, 6, 4, 2, 0, 2, 0, 2, 2, 0)^T,$$

$$y_{12} = (8, 6, 6, 2, 0, 2, 6, 4, 2, 0, 2, 0, 2, 2, 0, 10, 6, 4, 4, 8, 2, 0, 4, 2, 4, 0, 0, 0, 0, 0)^T,$$

$$y_{13} = (10, 6, 4, 4, 8, 2, 0, 4, 2, 4, 0, 0, 0, 0, 0, 12, 6, 4, 4, 0, 6, 6, 4, 2, 0, 0, 0, 0, 0, 0)^T,$$

$$y_{14} = (12, 6, 4, 4, 0, 6, 6, 4, 2, 0, 0, 0, 0, 0, 0, 10, 6, 8, 6, 12, 4, 0, 0, 4, 4, 0, 0, 0, 0, 0)^T,$$

$$y_{15} = (10, 6, 8, 6, 12, 4, 0, 0, 4, 4, 0, 0, 0, 0, 0, 8, 10, 6, 6, 0, 4, 8, 4, 4, 0, 2, 2, 0, 0, 0)^T,$$

and introducing the recurrence relation

$$y_{2n} = F_0 y_n, \quad y_{2n+1} = F_1 y_n, \quad n \geq 8$$

one has the relation [8]:

$$u_{n+1} = w^T y_n. \tag{A.2}$$

We finally introduce two new matrices in $\mathbb{R}^{20 \times 20}$ that rule the asymptotics of u_n :

$$A_0 = \begin{pmatrix} C_1 & 0_{10 \times 10} \\ D_1 & B_1 \end{pmatrix}, \quad A_1 = \begin{pmatrix} D_1 & B_1 \\ 0_{10 \times 10} & C_1 \end{pmatrix}. \tag{A.3}$$

B Proof of Theorem 2

In this appendix we give a proof of Theorem 2. We first find a common invariant cone K for the matrices A_0, A_1 . We then show that the products $F(n) = F_{d_1} \cdots F_{d_k}$ are asymptotically equivalent to their corresponding product $A(n) = A_{d_1} \cdots A_{d_k}$. We then shed some light on the vectors z_n , the restriction of y_n to \mathbb{R}^{20} : their norms can be considered as norms of the products $A(n)$. We finally show that $\|A_{d_1} \cdots A_{d_k}\|$ is equivalent to $\|A_{d_1} \cdots A_{d_{k-4}}\|$. We end with the proof of Theorem 2 that puts all this together.

Let us first establish a special property of the matrices A_0, A_1 . Consider the following sets:

$$P = \left\{ x \in \mathbb{R}^{20}, x \geq 0, \quad x_i > 0 \text{ for all } i \notin \{5, 10, 17\} \right\},$$

$$Q = \left\{ x \in \mathbb{R}^{20}, x \geq 0, \quad x_i > 0 \text{ for all } i \notin \{7, 15, 20\} \right\}.$$

Let $S = P \cup Q$. For any $\varepsilon \geq 0$ let $p_\varepsilon \in \mathbb{R}^{20}$, $(p_\varepsilon)_i = -\varepsilon$ for $i \in \{5, 10, 17\}$ and $(p_\varepsilon)_i = 1$ otherwise, let also $q_\varepsilon \in \mathbb{R}^{20}$, $(q_\varepsilon)_i = -\varepsilon$ for $i \in \{7, 15, 20\}$ and $(q_\varepsilon)_i = 1$ otherwise. This is easy to verify by direct calculation that there is $\varepsilon > 0$ such that $A_0 p_\varepsilon, A_0 q_\varepsilon \in P$ and $A_1 q_\varepsilon, A_1 p_\varepsilon \in Q$ (this is the case for instance for $\varepsilon = 1/4$). Take any such ε and denote $K = \{v + tp_\varepsilon + sq_\varepsilon, v \in \mathbb{R}_+^{20}, t, s \geq 0\}$. Clearly, K is a convex closed pointed cone.

Lemma 2 *We have $A_i K \subset K$ and $A_i S \subset S$ for $i = 0, 1$. Moreover, $S \subset \text{int}K$.*

Proof. Let $x \in K : x = v + tp_\varepsilon + sq_\varepsilon, v \geq 0, t, s \geq 0$. Since $A_i v \geq 0, A_i p_\varepsilon \geq 0$ and $A_i q_\varepsilon \geq 0$ we see that $A_i x \geq 0$. Thus, $A_i K \subset K$.

Further, for an arbitrary $x \in S$ let $x_{\min} = \min_{x_i > 0} x_i$. Assume $x \in P$ (the proof for the case $x \in Q$ is literally the same). Since $x \geq x_{\min} p_0$ we have $A_0 x \geq x_{\min} A_0 p_0 \geq x_{\min} p_0 \in P$ and $A_1 x \geq x_{\min} A_1 p_0 \geq x_{\min} q_0 \in Q$. Therefore $A_i x \in S$.

Finally, for any $x \in S$ and $y \in \mathbb{R}^{20}$ such that $|y|_\infty < \min\{\frac{1}{2}\varepsilon x_{\min}, \frac{1}{2}\varepsilon\}$ we have $x + y \in K$. This proves that $x \in \text{int}K$. \square

Corollary 5 *For any $x \in S$ and for any sequence d_1, \dots, d_k we have $A_{d_1} \cdots A_{d_k} x \in \text{int}K$.*

Lemma 3 *Suppose $n \geq 1$ and consider the binary expansion $d_k \dots d_1$ of the number $n - 1$. We define $A(n) = A_{d_1} \cdots A_{d_k}$ and similarly for $F(n)$. Then, for any matrix norm one has:*

$$\|F(n)\| \asymp \|A(n)\|.$$

Proof. Since all matrix norms are equivalent, it suffices to consider any norm. Obviously $\|F_{d_1} \cdots F_{d_k}\|_1 \geq \|A_{d_1} \cdots A_{d_k}\|_1$ (because the matrices A_i are submatrices of F_i), hence it remains to prove the opposite inequality: there is a positive constant C such that $\|A_{d_1} \cdots A_{d_k}\|_1 \geq C \|F_{d_1} \cdots F_{d_k}\|_1$ for all $k \in \mathbb{N}$ and d_1, \dots, d_k . We consider the case where $d_k = 1$; the proof for the other case is similar. Let $m \leq k - 1$ be the biggest number such that $d_m = 0$. If the sequence has no zero, we fix $m = 0$. Let A_i, H_i , and R_i denote respectively

the upper left, the upper right and the lower right corners of the matrix F_i in bloc representation A.1. Then the product $F_{d_1} \dots F_{d_k}$ has the following form: the left upper block is $A_{d_1} \dots A_{d_k}$, the left lower block is zero, the right lower block is $R_{d_1} \dots R_{d_k}$, finally, the right upper block is

$$\sum_{p=1}^k \left(\prod_{j=1}^{p-1} A_{d_j} \right) \cdot H_{d_p} \cdot \left(\prod_{j=p+1}^k R_{d_j} \right). \quad (\text{B.1})$$

By convention, the product over an empty set is one. Since $R_0 R_1 = 0$, the right lower block is zero and block (B.1) becomes $\sum_{p=m}^k \left(\prod_{j=1}^{p-1} A_{d_j} \right) H_{d_p} R_1^{k-p}$, whose norm can be estimated from above as

$$H \sum_{p=m}^k \left\| \prod_{j=1}^{p-1} A_{d_j} \right\|_1 \cdot \left\| R_1^{k-p} \right\|_1, \quad (\text{B.2})$$

where $H = \max\{\|H_0\|_1, \|H_1\|_1\}$. It was shown in [8] that the sum of entries of the matrix R_i^l does not exceed $C2^l$ for any $i = 0, 1$ and $l \geq 1$, where $C > 0$ is a constant. Hence $\|R_1^{k-p}\|_1 \leq C2^{k-p}$. Thus,

$$\|F_{d_1} \dots F_{d_k}\|_1 \leq \|A_{d_1} \dots A_{d_k}\|_1 + HC \sum_{p=m}^k 2^{k-p} \left\| \prod_{j=1}^{p-1} A_{d_j} \right\|_1. \quad (\text{B.3})$$

On the other hand, for any $p \geq m$ we have $|A_{d_1} \dots A_{d_k} \mathbf{1}| = |A_{d_1} \dots A_{d_{p-1}} (A_{d_p} A_1^{k-p} \mathbf{1})|$, where $\mathbf{1}$ is the vector of ones. By Corollary 5 the vectors $A_{d_p} A_1^r \mathbf{1}$ belong to $\text{int}K$ for all $r \in \mathbb{N}$. Moreover, the vector $A_1^r \mathbf{1} / |A_1^r \mathbf{1}|$ converges to v_1 (the Perron-Frobenius eigenvector of A_1) as $r \rightarrow \infty$ and $v_1 \in S$, hence $v_1 \in \text{int}K$ (Lemma 2). Therefore there is a constant $C_1 > 0$ such that $\gamma(A_{d_p} A_1^r \mathbf{1}) \geq C_1$ for all $r \in \mathbb{N}$. Applying now Lemma 1 for $x = A_{d_p} A_1^{k-p} \mathbf{1}$, we get

$$\|A_{d_1} \dots A_{d_k}\| \geq C_2 |A_{d_1} \dots A_{d_{p-1}} (A_{d_p} A_1^{k-p} \mathbf{1})|$$

$$\geq C_2 C_1 \|A_{d_1} \dots A_{d_{p-1}}\| \cdot |A_{d_p} A_1^{k-p} \mathbf{1}| \geq C_3 \lambda^{k-p} \|A_{d_1} \dots A_{d_{p-1}}\|,$$

where $\lambda = \rho(A_1)$ (by the same reasoning we have $|A_{d_p} A_1^r \mathbf{1}| \geq \gamma(A_1^r \mathbf{1}) \|A_{d_p}\| \cdot |A_1^r \mathbf{1}| \geq C \lambda^r$, indeed, $A_1^r \mathbf{1} \in \text{int}K$, and thus $\|A_{d_p} A_1^r \mathbf{1}\| \asymp \|A_{d_p}\| \cdot |A_1^r \mathbf{1}|$). Thus, $\|A_{d_1} \dots A_{d_{p-1}}\| \leq C_3^{-1} \lambda^{p-k} \|A_{d_1} \dots A_{d_k}\|$ for all $p \leq k$. Substituting this in (B.3) and taking into account that $\frac{2}{\lambda} < 1$ (because $\lambda = \rho(A_1) > 2.42$) we take the sum of the geometrical progression and get $\|F_{d_1} \dots F_{d_k}\|_1 \leq C_4 \|A_{d_1} \dots A_{d_k}\|_1$, where C_4 is some constant. This concludes the proof. \square

Lemma 4 *For any n we have $u_{n+1} \leq 2u_n$.*

Proof. If a word of length $n + 1$ is overlap-free then so is its prefix of length n . On the other hand, at most two overlap-free words of length $n + 1$ have the

same prefix of length n . \square

Lemma 5 *Let the vectors $y_m \in \mathbb{R}^{30}$ be the solution of the recurrence equation 4, and $z_m \in \mathbb{R}^{20}$ be the vector with the first 20 entries of y_n . We have $z_m \in S$ for each $m = 64, \dots, 127$.*

Proof. The proof is by direct calculation. \square

Lemma 6 *Suppose $n \in \mathbb{N}$ and $d_k \dots d_1$ is the binary expansion of $n - 1$; then $\|A(n)\| \asymp \|A'(n)\|$, where $A(n) = A_{d_1} \cdots A_{d_k}$ and $A'(n) = A_{d_1} \cdots A_{d_{k-4}}$.*

Proof. The inequality $\|A_{d_1} \cdots A_{d_k}\| \leq C \|A_{d_1} \cdots A_{d_{k-4}}\|$ is obvious by submultiplicativity of the norm. For the other direction, we have:

$$\|A_{d_1} \cdots A_{d_k}\| \asymp |A_{d_1} \cdots A_{d_k} \mathbf{1}| \asymp |A_{d_1} \cdots A_{d_{k-4}} (A_{d_{k-3}} \cdots A_{d_k} \mathbf{1})|. \quad (\text{B.4})$$

Corollary 5 yields $A_{d_{k-3}} \cdots A_{d_k} \mathbf{1} \in \text{int}K$ for all $d_{k-3}, \dots, d_k \in \{0, 1\}$. Applying now Lemma 1 we get

$$\delta = \min_{d_{k-3}, \dots, d_k \in \{0, 1\}} \gamma(A_{d_{k-3}} \cdots A_{d_k} \mathbf{1}) > 0.$$

Therefore, for some $C_1 > 0$,

$$\begin{aligned} C_1 \|A_{d_1} \cdots A_{d_{k-4}}\| &\leq \delta \|A_{d_1} \cdots A_{d_{k-4}}\| \cdot |A_{d_{k-3}} A_{d_{k-1}} A_{d_k} \mathbf{1}| \\ &\leq |A_{d_1} \cdots A_{d_k} \mathbf{1}|. \end{aligned} \quad (\text{B.5})$$

Combining this with (B.4) we get $\|A_{d_1} \cdots A_{d_{k-4}}\| \leq C_2 \|A_{d_1} \cdots A_{d_k}\|$. \square
We are now able to prove Theorem 2:

Proof of Theorem 2. Let g be the vector of \mathbb{R}^{30} , whose first 20 entries are ones and the last 10 entries are zeros. Let also $m = d_{k-3} + d_{k-2}2 + 2^2 d_{k-1} + 2^3 d_k$. Since $w \leq 2g$, we have

$$\begin{aligned} u_n &\leq 2(y_{n-1}, g) = 2(F_{d_1} \cdots F_{d_{k-4}} y_m, g) \\ &\leq C_0 \|F_{d_1} \cdots F_{d_{k-4}}\| \asymp \|A_{d_1} \cdots A_{d_{k-4}}\|, \end{aligned} \quad (\text{B.6})$$

where C_0 does not depend on n (the first two relations are direct from fundamental assertions A.2, the third relation comes from the fact that y_m and g are bounded, and the last equivalence is by Lemma 3). Combining Lemma 6 and (B.6) gives $u_n \leq C_3 \|A_{d_1} \cdots A_{d_k}\|$.

Let us now prove the opposite inequality. Lemma 4, together with the fact that by construction, the first ten entries of y_n are equal to the entries 11, \dots , 20 of y_{n-1} , imply that $u_n \geq \frac{1}{3}(u_n + u_{n+1}) \geq \frac{1}{6}(y_{n-1}, g)$. Furthermore, for $n > 2^7$ we have $(y_{n-1}, g) = (F_{d_1} \cdots F_{d_{k-7}} y_l, g)$, where $l = \sum_{j=0}^6 d_{k-6+j} 2^j$. Thus,

$$u_n \geq \frac{1}{6}(F_{d_1} \cdots F_{d_{k-7}} y_l, g). \quad (\text{B.7})$$

On the other hand, defining $z_l \in \mathbb{R}^{20}$ as the vector with the first 20 entries of $y_l : (F_{d_1} \cdots F_{d_{k-7}} y_l, g) \geq (A_{d_1} \cdots A_{d_{k-7}} z_l, \mathbf{1}) = |A_{d_1} \cdots A_{d_{k-7}} z_l|_1$. By Lemma 5 we have $z_l \in \text{int}K$ for all $l \in \{64, \dots, 127\}$, and we can define $h = \min_{64 \leq l \leq 127} \gamma(z_l) > 0$ such that $|A_{d_1} \cdots A_{d_{k-7}} z_l|_1 \geq h C_4 \|A_{d_1} \cdots A_{d_{k-7}}\|$, where $C_4 = \min_{64 \leq l \leq 127} |z_l|_1$. Combining this with (B.7), we obtain

$$u_n \geq C_5 \|A_{d_1} \cdots A_{d_{k-7}}\|. \quad (\text{B.8})$$

Now, by submultiplicativity of the norm,

$$u_n \geq C_6 \|A_{d_1} \cdots A_{d_k}\|.$$

□

C The ellipsoidal norm

Define

$$P_1 = \begin{pmatrix} 31.3 & 7.5 & 2.3 & 3.3 & -0.4 & -0.3 & 0.3 & 0.4 & 3.7 & 0.3 \\ 7.5 & 57.7 & 10 & 6.3 & 18.4 & 35 & 16.3 & -5.8 & 13.8 & 5 \\ 2.3 & 10 & 59.9 & 11.3 & 0.4 & 29.2 & 4.2 & 10.1 & 8.2 & 0.8 \\ 3.3 & 6.3 & 11.3 & 48.5 & 4.6 & 13.5 & 10.8 & 2 & 6.9 & 1 \\ -0.4 & 18.4 & 0.4 & 4.6 & 36.4 & 23.5 & 22.6 & 4.4 & 8.9 & -1.2 \\ -0.3 & 35 & 29.2 & 13.5 & 23.5 & 105.9 & 38.4 & 9.5 & 33.7 & 6.1 \\ 0.3 & 16.3 & 4.2 & 10.8 & 22.6 & 38.4 & 59 & 2.7 & 17.4 & 9.2 \\ 0.4 & -5.8 & 10.1 & 2 & 4.4 & 9.5 & 2.7 & 38.6 & 14.8 & -1.7 \\ 3.7 & 13.8 & 8.2 & 6.9 & 8.9 & 33.7 & 17.4 & 14.8 & 57.5 & 8.6 \\ 0.3 & 5 & 0.8 & 1 & -1.2 & 6.1 & 9.2 & -1.7 & 8.6 & 42.3 \end{pmatrix},$$

$$P_2 = \begin{pmatrix} -10.4 & -1.7 & -18.1 & -0.4 & -5.8 & -5.1 & -4.9 & -0.8 & -2.7 & -0.9 \\ -11.1 & -22.4 & -8.2 & -14.7 & -9.9 & -30.3 & -16.7 & -11.3 & -16.9 & -6.6 \\ -2.2 & -16.4 & -15.8 & -5 & -8.5 & -7.2 & -5.4 & -18.5 & -3.5 & -3.4 \\ -0.2 & -13.6 & -5.2 & -9 & -10.7 & -14.6 & -9.2 & -1.6 & -11.3 & -1.1 \\ -4.6 & -17 & -13 & -9.1 & -0.6 & -11.2 & -23.9 & -7 & -12.1 & 0.3 \\ -5.9 & -26.4 & -27.4 & -17.4 & -31 & -37.6 & -28 & -4.4 & -27.3 & -7.4 \\ -1.4 & -19.3 & -11.6 & -10.8 & -22.3 & -17.9 & -11.7 & -11.3 & -12 & -9.8 \\ -6.3 & 2.1 & 1.7 & -3.4 & 3.2 & -7.6 & 0.2 & -5.2 & -3.1 & -1.4 \\ -7.4 & -15.9 & -4.7 & -6.7 & -12.2 & -17.3 & -11.6 & -5.3 & -6.8 & -1.6 \\ 1.3 & -5.7 & -3.6 & -3.2 & -0.4 & -6.1 & -9 & -1.4 & -6.9 & 0.4 \end{pmatrix},$$

$$P_4 = \begin{pmatrix} 29.1 & 8.3 & -1.6 & 4.8 & -1.3 & -4.4 & 0.6 & 1.7 & 7.5 & 1.1 \\ 8.3 & 47.3 & 13.6 & 2.8 & 11.7 & 19.8 & 17.4 & 0.6 & 10 & 3.7 \\ -1.6 & 13.6 & 46.6 & 10.4 & 6.5 & 24.9 & 11.8 & 6.5 & 12.5 & 1.4 \\ 4.8 & 2.8 & 10.4 & 47.6 & 5.1 & 8 & 7.6 & 5.1 & 3.7 & 1.8 \\ -1.3 & 11.7 & 6.5 & 5.1 & 32.8 & 19.5 & 19.4 & 7.6 & 6.7 & -0.2 \\ -4.4 & 19.8 & 24.9 & 8 & 19.5 & 64.8 & 16.2 & 11.4 & 13.8 & 6.8 \\ 0.6 & 17.4 & 11.8 & 7.6 & 19.4 & 16.2 & 56.7 & 7.6 & 12.2 & 6.5 \\ 1.7 & 0.6 & 6.5 & 5.1 & 7.6 & 11.4 & 7.6 & 38.7 & 11.2 & -1 \\ 7.5 & 10 & 12.5 & 3.7 & 6.7 & 13.8 & 12.2 & 11.2 & 55.6 & 4.2 \\ 1.1 & 3.7 & 1.4 & 1.8 & -0.2 & 6.8 & 6.5 & -1 & 4.2 & 43.8 \end{pmatrix},$$

$$P = \begin{pmatrix} P_1 & P_2 \\ P_2^T & P_4 \end{pmatrix}.$$

Then one has the relations:

$$A^t P A - (2.5186)^{28} P \prec 0, \quad \forall A \in \Sigma^{14}.$$

As explained in [3] this suffices to prove that $\rho(\Sigma) \leq 2.5186$.

D The vector x

Define

$$x = (153, 0, 60, 0, 50, 56, 99, 0, 58, 1, 157, 81, 0, 113, 0, 72, 0, 99, 0, 0)^T.$$

Then one has the relation

$$\begin{aligned} B(Ax - rx) &\geq 0, \quad \forall B \in \Sigma^6, A \in \Sigma^{16}, \\ x &\geq 0, \end{aligned} \tag{D.1}$$

with $r = 2.41^{16}$. this proves that $\check{\rho}(\Sigma) \geq 2.41$.