

Pattern Avoidance in “Flattened” Partitions

DAVID CALLAN

Department of Statistics
 University of Wisconsin-Madison
 1300 University Ave
 Madison, WI 53706-1532
callan@stat.wisc.edu

February 15, 2008

Abstract

To flatten a set partition (with apologies to *Mathematica*[®]) means to form a permutation by erasing the dividers between its blocks. Of course, the result depends on how the blocks are listed. For the usual listing—increasing entries in each block and blocks arranged in increasing order of their first entries—we count the partitions of $[n]$ whose flattening avoids a single 3-letter pattern. Five counting sequences arise: a null sequence, the powers of 2, the Fibonacci numbers, the Catalan numbers, and the binomial transform of the Catalan numbers.

1 Introduction

There is an extensive literature on pattern avoidance in permutations. Klazar [1, 2, 3] considered an analogous notion for set partitions and Sagan [4] introduced a second such notion based on restricted growth functions (see also [5, 6]). Here we consider set partitions avoiding a permutation in the following sense. Suppose partitions Π of $[n] = \{1, 2, \dots, n\}$ are written in some pre-specified standard form, say *standard increasing* form: increasing entries in each block and blocks arranged in increasing order of their first entries. Then define $\text{Flatten}(\Pi)$ to be the permutation of $[n]$ obtained by erasing the dividers between the blocks of Π . For example, $\Pi = 136-279-4-58$ is in standard increasing form and $\text{Flatten}(\Pi) = 136279458$. (The computer algebra system *Mathematica*[®] implements this operation with the command `Flatten`). For a permutation π on an initial segment of

the positive integers (a pattern permutation) we say the partition Π avoids π or Π is π -avoiding if the permutation $\text{Flatten}(\Pi)$ avoids π (in the classical sense). We write $\Pi \vdash [n]$ if Π is a partition of $[n]$. Set $\mathcal{U}(n; \pi) = \{\Pi \vdash [n] : \text{Flatten}(\Pi) \text{ avoids } \pi\}$.

In §2, we fix standard increasing as the form for writing partitions of $[n]$ and count $\mathcal{U}(n; \pi)$ for all 3-letter pattern permutations π .

2 Set partitions in standard increasing form

2.1 123-avoiding

This case is not very interesting; the counting sequence $(|\mathcal{U}(n; 123)|)_{n \geq 1}$ is $(1, 2, 1, 0, 0, 0, \dots)$.

2.2 132-avoiding

A partition Π of $[n]$ is in $\mathcal{U}(n; 132)$ if and only if $\text{Flatten}(\Pi)$ is the identity permutation. This is because the first entry of $\text{Flatten}(\Pi)$ is always 1 and will be the ‘1’ of a 132 pattern unless $\text{Flatten}(\Pi)$ is an increasing sequence, that is, the identity permutation. So any subset of the $n - 1$ spaces between $1, 2, \dots, n$ can serve as the dividers to form Π and $|\mathcal{U}(n; 132)| = 2^{n-1}$.

2.3 213-avoiding

First, we claim a partition Π of $[n]$ is in $\mathcal{U}(n; 213)$ if and only if (i) the first block of Π has the form $I \cup J$ with I a nonempty initial segment of $[n]$ and J a terminal segment of $[n]$ (possibly empty) disjoint from I , and (ii) the remaining blocks, when standardized, themselves form a 213-avoiding partition. (To standardize means to replace smallest entry by 1, second smallest by 2, and so on.)

Clearly, these two conditions are sufficient and condition (ii) is necessary. If condition (i) fails for $\Pi \in \mathcal{U}(n; 213)$, let a be the smallest element of $[n]$ not in the first block; a is necessarily the first element of the second block. Because the condition fails there exist b, c in $[n]$ with $c > b > a$, b in the first block and c in a later block. Hence c occurs after a and bac is a 213-pattern in $\text{Flatten}(\Pi)$, a contradiction. So condition (i) is necessary

also.

Now let $u(n) = |\mathcal{U}(n; 213)|$ and set $u(n, k) = |\{\Pi \in \mathcal{U}(n; 213) : \text{first block of } \Pi \text{ has length } k\}|$. Clearly, $u(n, n) = 1$ and for $1 \leq k \leq n$, the first block is determined by I and there are k choices for I , namely, $([i])_{i=1}^k$. Hence we have the system of equations

$$\begin{aligned} u(n, n) &= 1 && \text{for } n \geq 1 \\ u(n, k) &= ku(n-k) && \text{for } 1 \leq k < n \\ u(n) &= \sum_{k=1}^n u(n, k) && \text{for } n \geq 1 \end{aligned}$$

with solution involving the Fibonacci numbers ($F_{-1} := 1, F_0 = 0, F_1 = 1$)

$$\begin{aligned} u(n, j) &= jF_{2n-2j-1} && \text{for } 1 \leq j < n \\ u(n) &= F_{2n-1}. \end{aligned}$$

2.4 231-avoiding

This case gives rise to the Catalan numbers via Touchard's identity [7],

$$C_n = \sum_{k \geq 0} \binom{n-1}{2k} 2^{n-1-2k} C_k. \quad (1)$$

For a permutation p of $[n]$, a *descent terminator* is an entry smaller than its immediate predecessor and, by convention, the first entry is also considered a descent terminator. A *right-to-left (R-L) minimum* of p is an entry smaller than all the entries after it. Clearly, for a partition in standard increasing form and its associated permutation, $\{\text{descent terminators}\} \subseteq \{\text{block initiators}\} \subseteq \{\text{R-L minima}\}$. For $\Pi \vdash [n]$, let $M(\Pi)$ denote the set of R-L minima of $\text{Flatten}(\Pi)$ that are not descent terminators, and set $\mathcal{U}(n, k; 231) = \{\Pi \in \mathcal{U}(n; 231) : |M(\Pi)| = k\}$. We claim $|\mathcal{U}(n, k; 231)| = \binom{n-1}{k} 2^k C_{\frac{n-1-k}{2}}$ where C_n is the Catalan number and $C_n := 0$ when n is not an integer. Touchard's identity (1) then implies $|\mathcal{U}(n; 231)| = C_n$.

To establish the claim, it suffices to show

$$|\mathcal{U}(n, 0; 231)| = C_{\frac{n-1}{2}} \quad \text{for } n \geq 1, \text{ and} \quad (2)$$

$$|\mathcal{U}(n, k; 231)| = \binom{n-1}{k} 2^k |\mathcal{U}(n-k, 0; 231)| \quad \text{for } 1 \leq k < n. \quad (3)$$

To show (2), let $\Pi \in \mathcal{U}(n, 0; 231)$. Then the R-L minima and descent terminators of $p := \text{Flatten}(\Pi)$ coincide. The last entry of p is certainly a R-L minimum, hence a descent terminator, and so it must form a singleton block in Π . Each non-last block has length ≤ 2 because if (a, b, c, \dots) is a block of length ≥ 3 , then bcd is a 231-pattern where d is the first entry of the next block: certainly $b < c$ and we also have $d < b$ because if $b < d$, then $b < \text{all}$ entries that follow it. This would make b a R-L minimum that was not a descent terminator, a contradiction. On the other hand, each non-last block has length ≥ 2 because a non-last singleton block would imply that the first entry of the next block was a R-L minimum that was not a descent terminator. Hence all but the last block have length 2 and so n is odd, say $n = 2r + 1$, and Π is of the form $a_1 b_1 - a_2 b_2 - \dots - a_r b_r - a_{r+1}$.

Clearly, $a_1 = 1$. Also, $a_2 = 2$ because otherwise, since a_2 is a R-L minimum, 2 would occur to the left of a_2 and this would force $b_2 = 2$. But then a_2 would be a R-L minimum that was not a descent terminator. Next, we claim $a_{i+2} \leq 2i + 2$ for $1 \leq i \leq r - 1$. Suppose contrariwise that $a_{i+2} > 2i + 2$ for some i . Then none of $3, 4, \dots, 2i + 2$ can occur after a_{i+2} because a_{i+2} is a R-L minimum. This forces the first $i + 1$ blocks to consist of the first $2i + 2$ positive integers leaving b_{i+1} a R-L minimum, which is not possible. Hence the sequence $(c_i)_{i=1}^{r-1}$ with $c_i := a_{i+2} - 2$ satisfies

$$1 \leq c_1 < c_2 < \dots < c_{r-1}, \quad \text{and} \quad c_i \leq 2i \quad \text{for } 1 \leq i \leq r - 1. \quad (4)$$

We have exhibited a map from $\mathcal{U}(2r + 1, 0; 231)$ to sequences $(c_i)_{i=1}^{r-1}$ satisfying (4). This map is in fact a bijection and here is its inverse. Given such a sequence, for example with $r = 9$, $(c_i) = (1, 2, 4, 5, 7, 12, 13, 15)$, we can immediately recover the a_i 's and must determine the b_i 's (blank squares in Fig. 1).

$$\begin{array}{cccccccccc} a_1 & b_1 & a_2 & b_2 & \dots & & \dots & a_r & b_r & a_{r+1} \\ 1 & \square & 2 & \square & 3 & \square & 4 & \square & 6 & \square & 7 & \square & 9 & \square & 14 & \square & 15 & \square & 17 \end{array}$$

Fig. 1

Fill in the blank squares using $B = [2r + 1] \setminus (a_i)_{i=1}^{r+1}$ from right to left as follows. Place the smallest element of B that exceeds a_{r+1} in the b_r square and, in general, place the smallest not-yet-placed element of B that exceeds a_{i+1} in the b_i square. The example has $B = \{5, 8, 10, 11, 12, 13, 16, 18, 19\}$, yielding $(b_i)_{i=1}^r = (13, 12, 5, 11, 8, 10, 19, 16, 18)$.

There is a nice graphical way to visualize the result of this algorithmic procedure using Dyck paths. Recall that the Catalan number C_r counts sequences $(c_i)_{i=1}^{r-1}$ satisfying (4) [8, Ex. 6.19, item t]. Indeed, given a Dyck path of semilength r let c_i denote the number

of steps preceding the $(i + 1)$ st upstep for $1 \leq i \leq r - 1$. This is a bijection from Dyck r -paths to the sequences $(c_i)_{i=1}^{r-1}$ satisfying (4). So, sketch the Dyck path corresponding to the sequence $(c_i)_{i=1}^{r-1}$, prepend an upstep, and number all $2r + 1$ steps in order left to right, as in Fig.2 for our running example.




Fig. 2

Every upstep in a Dyck path has a matching downstep: the first one encountered directly east from the upstep or, more precisely, the terminal downstep of the shortest Dyck subpath starting at the upstep. The a_i 's are evident in the augmented Dyck path as the labels on the upsteps, and the b_i 's are also discernible: b_i is the label on the matching downstep for the *next* upstep after a_i , $1 \leq i \leq r$. It is now clear that the a_i 's are increasing and that $a_i < b_i > a_{i+1}$ for $1 \leq i \leq r$; hence $(a_i)_{i=1}^{r+1}$ is both the set of R-L minima and the set of descent terminators in $\text{Flatten}(\Pi)$ and so $\Pi \in \mathcal{U}(2r + 1, 0; 231)$. It is also easy to verify that Π is 231-avoiding. Indeed, since all entries following a_i are $> a_i$, the first two entries of a putative 231 pattern would have to be b 's, say $b_i < b_j$ with $i < j$, and a_j would be the last upstep preceding b_i (or else b_j would be $< b_i$). Hence, for all $k > j$, upstep a_k occurs after b_i and so $b_k > a_k > b_i$ for $k > j$. Since b_i is the '2' of the 231 pattern and we have just seen that all later entries are larger than b_i , no entry after b_j can serve as the '1' of the pattern. We conclude that the partition $a_1 b_1 - a_2 b_2 - \dots - a_r b_r - a_{r+1}$ is in $\mathcal{U}(n, 0; 231)$ as required.

To prove (3), consider $\Pi \in \mathcal{U}(n, k; 231)$. Let K denote the set of R-L minima that are not descent terminators in $\text{Flatten}(\Pi)$. Thus $|K| = k$ and $K \subseteq [2, n]$. Let L denote the set of elements in K that initiate a block in Π . Thus $L \subseteq K$. Let Π_0 denote the partition obtained from Π by deleting each element i of K from its block and, if i is also in L , concatenating this block with the currently preceding block. Then $\Pi_0 \in \mathcal{U}(n - k, 0; 231)$. For example, $\Pi = 1-24-37-568$ yields $K = \{2, 6, 8\}$, $L = \{2\}$, and $\Pi_0 = \text{standardize}(14-37-5) = 13-25-4$. An example where three consecutive blocks are concatenated to form Π_0 is $\Pi = 1-2-35-4$ with $K = \{2, 3\}$, $L = \{3\}$, and $\Pi_0 = \text{standardize}(15-4) = 13-2$. We claim the map $\mathcal{U}(n, k; 231) \rightarrow (K, L, \Pi_0)$ is a bijection to all triples (K, L, Π_0) with K a k -element subset of $[2, n]$, L an arbitrary subset

of K , and Π_0 a partition in $\mathcal{U}(n - k, 0; 231)$, and (3) then follows from (2). To establish the claim, suppose given such a triple (K, L, Π_0) , and build up Π as follows from Π_0 . For each $a \in K$ in turn from smallest to largest, locate the last block in the current partition whose first entry is $< a$; then, to get the next partition, after adding 1 to each entry $\geq a$ insert a into the located block at the appropriate position to ensure an increasing block. The end result will be a partition of $[n]$ in which the descent terminators are the block initiators and no element of K is a block initiator. Finally, for each element of K that is in the subset L , place a divider just before that element so that it initiates a block. This procedure yields Π and shows the map is invertible.

2.5 312-avoiding

We claim a partition Π of $[n]$ is in $\mathcal{U}(n; 312)$ if and only if (i) the first block of Π is all of $[n]$ or has the form $I \setminus \{a\}$ where I is an initial segment of $[n]$ of length ≥ 2 and $a \geq 2$ is in I , and (ii) the remaining blocks, when standardized, themselves form a 312-avoiding partition.

The conditions are sufficient because if they hold and a 312 pattern involved the first block, then only the ‘3’ could occur in the first block leaving the ‘1’ and ‘2’ to occur in later blocks. This however is impossible because at most one letter smaller than the ‘3’ is missing from the first block. So we merely need to show that condition (i) is necessary. Suppose then that condition (i) is not met. Let c denote the largest entry in the first block and a the smallest letter missing from the first block. Then by supposition there is a letter b missing from the first block with $a < b < c$. Since a must be the first entry of the second block, b occurs after a and cab is a 312 pattern in $\text{Flatten}(\Pi)$, a contradiction.

Now, if the first block has length $k < n$, there are exactly k choices for a , namely, $2, 3, \dots, k + 1$. This observation leads to the very same recurrence relation as in the 213-avoiding case, and another Fibonacci counting sequence: $|\mathcal{U}(n; 312)| = F_{2n-1}$.

2.6 321-avoiding

This case is counted by the binomial transform of the Catalan numbers: $|\mathcal{U}(n+1; 321)| = \sum_{k=0}^n \binom{n}{k} C_k$. Our proof is quite similar to that of the 231-avoiding case but with Touchard’s

identity replaced by the following one involving the Riordan numbers R_n ,

$$\sum_{k=0}^n \binom{n}{k} 2^k R_{n-k} = \sum_{k=0}^n \binom{n}{k} C_k, \quad (5)$$

where $R_n := \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} C_j$. The identity (5) is easily proved by reversing the order of summation after substituting for R_{n-k} .


The Riordan number R_n (A005043 in OEIS) is well known to count, among other things, Dyck n -paths with no short descents. (A ‘descent’ is a maximal sequence of contiguous downsteps and ‘short’ means of length 1.) Mimicking Section 2.4, define $\mathcal{U}(n, k; 321) = \{\Pi \in \mathcal{U}(n; 321) : |M(\Pi)| = k\}$. We claim $|\mathcal{U}(n, k; 321)| = \binom{n-1}{k} 2^k R_{n-1-k}$ for $0 \leq k \leq n-1$, and the identity (5) then implies $|\mathcal{U}(n; 321)| = \sum_{k=0}^{n-1} \binom{n-1}{k} C_k$.

To establish the claim, it suffices to show

$$|\mathcal{U}(n, 0; 321)| = R_{n-1} \quad (6)$$

$$|\mathcal{U}(n, k; 321)| = \binom{n-1}{k} 2^k |\mathcal{U}(n-k, 0; 321)| \quad \text{for } 1 \leq k \leq n-1 \quad (7)$$

To prove assertion (6) there is a bijection (essentially due to Krattenthaler [9]) from $\mathcal{U}(n, 0; 321)$ to Dyck $(n-1)$ -paths with no short descents, illustrated with $n=9$:



Bijection $\mathcal{U}(n, 0; 321) \longrightarrow$ to Dyck $(n-1)$ -paths with no short descents

The proof of assertion (7) uses the same bijection $\Pi \rightarrow (K, L, \Pi_0)$ as in the proof of (3) and is omitted.

It would be interesting to investigate permutation-avoidance for other canonical representations of a set partition where less familiar counting sequences seem to arise.

References

- [1] Martin Klazar, On abab-free and abba-free set partitions, *European J. Combin.* **17**, 1 (1996), 53–68.
- [2] Martin Klazar, Counting pattern-free set partitions I. A generalization of Stirling numbers of the second kind, *European J. Combin.* **21**, 3 (2000), 367378.
- [3] Martin Klazar, Counting pattern-free set partitions II. Noncrossing and other hypergraphs, *Electronic J. Combin.* **7** (2000), Research Paper 34, 25 pp. (electronic).
- [4] Bruce E. Sagan, [Pattern avoidance in set partitions](#), preprint, 2006.
- [5] Adam M. Goyt, [Avoidance of partitions of a three-element set](#), preprint, 2006.
- [6] Vit Jelinek and Toufik Mansour, [On pattern-avoiding partitions](#), preprint, 2007.
- [7] Louis W. Shapiro, A short proof of an identity of Touchard’s concerning Catalan numbers, *J. Combinatorial Theory A* **20**, No. 3, (1976) 375–376.
- [8] Richard P. Stanley, *Enumerative Combinatorics* Vol. 2, Cambridge University Press, 1999. Exercise 6.19 and related material on Catalan numbers are available online at <http://www-math.mit.edu/~rstan/ec/> .
- [9] Christian Krattenthaler, [Permutations with restricted patterns and Dyck paths](#), *Adv. Appl. Math.* **27** (2001), 510–530.