

Preprint, arXiv:0911.3060

FIBONACCI NUMBERS MODULO CUBES OF PRIMES

ZHI-WEI SUN

Department of Mathematics, Nanjing University
 Nanjing 210093, People's Republic of China
 zwsun@nju.edu.cn
<http://math.nju.edu.cn/~zwsun>

ABSTRACT. Let p be an odd prime. It is well known that $F_{p-(\frac{p}{5})} \equiv 0 \pmod{p}$ where $\{F_n\}_{n \geq 0}$ is the famous Fibonacci sequence and $(-)$ is the Jacobi symbol. In this paper we show that if $p \neq 5$ then we may determine $F_{p-(\frac{p}{5})} \pmod{p^3}$ in the following way:

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}}{(-16)^k} \equiv \left(\frac{p}{5}\right) \left(1 + \frac{F_{p-(\frac{p}{5})}}{2}\right) \pmod{p^3}.$$

We also use Lucas quotients to determine $\sum_{k=0}^{(p-1)/2} \binom{2k}{k} / m^k \pmod{p^2}$ for any integer $m \not\equiv 0 \pmod{p}$; in particular, we obtain

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}}{16^k} \equiv \left(\frac{3}{p}\right) \pmod{p^2}.$$

In addition, we raise two conjectures for further research.

1. INTRODUCTION

The well known Fibonacci sequence $\{F_n\}_{n \geq 0}$, defined by

$$F_0 = 0, \quad F_1 = 1, \quad \text{and } F_{n+1} = F_n + F_{n-1} \quad (n = 1, 2, 3, \dots),$$

plays an important role in many fields of mathematics. This sequence also has nice number-theoretic properties; for example, E. Lucas showed that $(F_m, F_n) = F_{(m,n)}$ for any $m, n \in \mathbb{N} = \{0, 1, \dots\}$, where (m, n) denotes the greatest common divisor of m and n . It is known that $F_{p-(\frac{p}{5})} \equiv 0 \pmod{p}$ for any prime $p \neq 2, 5$, where $(-)$ denotes the Jacobi symbol. In 1992

2010 *Mathematics Subject Classification.* Primary 11B39, 11B65; Secondary 05A10, 11A07.

Keywords. Fibonacci numbers, central binomial coefficients, congruences, Lucas sequences.

Supported by the National Natural Science Foundation (grant 11171140) of China.

Z. H. Sun and Z. W. Sun [SS] proved that if $p^2 \nmid F_{p-(\frac{p}{5})}$ then the Fermat equation $x^p + y^p = z^p$ has no integral solutions with $p \nmid xyz$. A prime $p > 5$ satisfying $F_{p-(\frac{p}{5})} \equiv 0 \pmod{p^2}$ is called a Wall-Sun-Sun prime (cf. [CDP] and [CP, p. 32]). Now it is known that there are no Wall-Sun-Sun primes below 9.7×10^{14} (cf. [DK]) though it is conjectured that there should be infinitely many (but rare) such primes. There are some congruences for the Fibonacci quotient $F_{p-(\frac{p}{5})}/p$ modulo p (cf. [W], [SS] and [ST]); for example, in 1982 H. C. Williams [W] proved that

$$\frac{F_{p-(\frac{p}{5})}}{p} \equiv \frac{2}{5} \sum_{k=1}^{\lfloor \frac{4}{5}p \rfloor} \frac{(-1)^k}{k} \pmod{p}.$$

Quite recently H. Pan and Z. W. Sun [PS] proved that for any prime $p \neq 2, 5$ and positive integer a we have

$$\sum_{k=0}^{p^a-1} (-1)^k \binom{2k}{k} \equiv \left(\frac{p^a}{5}\right) \left(1 - 2F_{p^a-(\frac{p^a}{5})}\right) \pmod{p^3},$$

which was a conjecture in [ST].

Now we give the first theorem of this paper.

Theorem 1.1. *Let p be an odd prime and let a be a positive integer. If $p \neq 5$, then*

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-16)^k} \equiv \left(\frac{p^a}{5}\right) \left(1 + \frac{F_{p^a-(\frac{p^a}{5})}}{2}\right) \pmod{p^3}. \quad (1.1)$$

If $p \neq 3$, then

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-32)^k} \equiv \left(\frac{2}{p^a}\right) \left(1 + \frac{2^{p^a-1}-1}{6} - \frac{(2^{p^a-1}-1)^2}{8}\right) \pmod{p^3}. \quad (1.2)$$

Let p be an odd prime and let $a \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$. For $k = 0, 1, \dots, (p^a-1)/2$, clearly

$$\frac{\binom{(p^a-1)/2}{k}}{\binom{-1/2}{k}} = \prod_{j=0}^{k-1} \left(1 - \frac{p^a}{2j+1}\right) \equiv 1 \pmod{p}$$

and hence

$$\binom{(p^a-1)/2}{k} \equiv \binom{-1/2}{k} = \frac{\binom{2k}{k}}{(-4)^k} \pmod{p}. \quad (1.3)$$

Thus it is easy to see that

$$\begin{aligned} \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{m^k} &\equiv \sum_{k=0}^{(p^a-1)/2} \binom{(p^a-1)/2}{k} \left(-\frac{4}{m}\right)^k \\ &= \left(1 - \frac{4}{m}\right)^{(p^a-1)/2} \equiv \left(\frac{m(m-4)}{p^a}\right) \pmod{p} \end{aligned}$$

for any integer $m \not\equiv 0 \pmod{p}$. In view of Lucas' theorem (cf. [St, p. 44]), for any $k = (p^a + 1)/2, \dots, p^a - 1$ we have

$$\binom{2k}{k} = \binom{p^a + (2k - p^a)}{0p^a + k} \equiv \binom{2k - p^a}{k} = 0 \pmod{p}.$$

Recently the author [Su10] managed to determine $\sum_{k=0}^{p^a-1} \binom{2k}{k}/m^k \pmod{p^2}$ in terms of Lucas sequences. (See also [SSZ], [GZ] and [Su11a] for related results on p -adic valuations.)

Let $A, B \in \mathbb{Z}$. The Lucas sequences $u_n = u_n(A, B)$ ($n \in \mathbb{N}$) and $v_n = v_n(A, B)$ ($n \in \mathbb{N}$) are defined by

$$u_0 = 0, \quad u_1 = 1, \quad \text{and} \quad u_{n+1} = Au_n - Bu_{n-1} \quad (n = 1, 2, 3, \dots)$$

and

$$v_0 = 2, \quad v_1 = A, \quad \text{and} \quad v_{n+1} = Av_n - Bv_{n-1} \quad (n = 1, 2, 3, \dots).$$

The sequence $\{v_n\}_{n \geq 0}$ is called the companion of $\{u_n\}_{n \geq 0}$. (Note that $F_n = u_n(1, -1)$ and those $L_n = v_n(1, -1)$ are called Lucas numbers.) It is known that for any prime p not dividing $2B$ we have

$$u_p \equiv \left(\frac{\Delta}{p}\right) \pmod{p} \quad \text{and} \quad u_{p-(\frac{\Delta}{p})} \equiv 0 \pmod{p}$$

where $\Delta = A^2 - 4B$ (see, e.g., [Su10, Lemma 2.3]); the integer $u_{p-(\frac{\Delta}{p})}/p$ is called a *Lucas quotient*. The reader may consult [Su06] for connections between Lucas quotients and quadratic fields.

Our second theorem is as follows.

Theorem 1.2. *Let p be an odd prime and let $a \in \mathbb{Z}^+$. Let m be any integer not divisible by p . Then*

$$\begin{aligned} \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{m^k} &\equiv \left(\frac{m(m-4)}{p^a}\right) \\ &+ \left(\frac{-m}{p}\right) \left(\frac{m(m-4)}{p^{a-1}}\right) \bar{m} u_{p-(\frac{4-m}{p})}(4, m) \pmod{p^2}, \end{aligned} \tag{1.4}$$

where

$$\bar{m} = \begin{cases} 1 & \text{if } m \equiv 4 \pmod{p}, \\ 2 & \text{if } (\frac{4-m}{p}) = 1, \\ 2/m & \text{if } (\frac{4-m}{p}) = -1. \end{cases} \quad (1.5)$$

We also have

$$\sum_{k=0}^{(p^a-1)/2} \frac{C_k}{m^k} \equiv \frac{4-m}{2} \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{m^k} + \frac{m}{2} - 2p \delta_{a,1} \left(\frac{-m}{p} \right) \pmod{p^2}, \quad (1.6)$$

where C_k denotes the Catalan number $\frac{1}{k+1} \binom{2k}{k} = \binom{2k}{k} - \binom{2k}{k+1}$.

Now we present two consequences of Theorem 1.2.

Corollary 1.1. *Let p be an odd prime and let $a \in \mathbb{Z}^+$. Then*

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{8^k} \equiv \left(\frac{2}{p^a} \right) \pmod{p^2} \quad (1.7)$$

and

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{16^k} \equiv \left(\frac{3}{p^a} \right) \pmod{p^2}. \quad (1.8)$$

Corollary 1.2. *Let $p > 3$ be a prime. Then*

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}}{(2k-1)^2 16^k} \equiv \left(\frac{-1}{p} \right) \frac{3(\frac{p}{3}) + 1}{4} \pmod{p^2}, \quad (1.9)$$

that is,

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}}{(2k-1)^2 16^k} \equiv \begin{cases} 1 \pmod{p^2} & \text{if } p \equiv 1 \pmod{12}, \\ -1/2 \pmod{p^2} & \text{if } p \equiv 5 \pmod{12}, \\ -1 \pmod{p^2} & \text{if } p \equiv 7 \pmod{12}, \\ 1/2 \pmod{p^2} & \text{if } p \equiv 11 \pmod{12}. \end{cases} \quad (1.10)$$

We will show Theorems 1.1 and 1.2 in Sections 2 and 3 respectively. Section 4 is devoted to the proofs of Corollaries 1.1-1.2.

To conclude this section we pose two conjectures.

Conjecture 1.1. *For any $n \in \mathbb{N}$ we have*

$$\frac{1}{(2n+1)^2 \binom{2n}{n}} \sum_{k=0}^n \frac{\binom{2k}{k}}{16^k} \equiv \begin{cases} 1 \pmod{9} & \text{if } 3 \mid n, \\ 4 \pmod{9} & \text{if } 3 \nmid n. \end{cases}$$

Also,

$$\frac{1}{3^{2a}} \sum_{k=0}^{(3^a-1)/2} \frac{\binom{2k}{k}}{16^k} \equiv (-1)^a 10 \pmod{27}$$

for every $a = 1, 2, 3, \dots$.

Let $p > 3$ be a prime. In 2007 A. Adamchuk [A] conjectured that if $p \equiv 1 \pmod{3}$ then

$$\sum_{k=1}^{\lfloor \frac{2}{3}p \rfloor} \binom{2k}{k} \equiv 0 \pmod{p^2}.$$

Motivated by this and Theorems 1.1 and 1.2, we raise the following conjecture based on the author's computation via the software **Mathematica**.

Conjecture 1.2. *Let p be an odd prime and let $a \in \mathbb{Z}^+$.*

(i) *If $p \equiv 1 \pmod{3}$ or $a > 1$, then*

$$\sum_{k=0}^{\lfloor \frac{5}{6}p^a \rfloor} \frac{\binom{2k}{k}}{16^k} \equiv \left(\frac{3}{p^a} \right) \pmod{p^2}.$$

(ii) *If $p^a \equiv 1, 2 \pmod{5}$, or $a > 1$ and $p \not\equiv 3 \pmod{5}$,*

$$\sum_{k=0}^{\lfloor \frac{4}{5}p^a \rfloor} (-1)^k \binom{2k}{k} \equiv \left(\frac{5}{p^a} \right) \pmod{p^2}.$$

If $p^a \equiv 1, 3 \pmod{5}$, or $a > 1$ and $p \not\equiv 2 \pmod{5}$, then

$$\sum_{k=0}^{\lfloor \frac{3}{5}p^a \rfloor} (-1)^k \binom{2k}{k} \equiv \left(\frac{5}{p^a} \right) \pmod{p^2}.$$

(iii) *If $p \equiv 1, 7 \pmod{10}$ or $a > 2$, then*

$$\sum_{k=0}^{\lfloor \frac{7}{10}p^a \rfloor} \frac{\binom{2k}{k}}{(-16)^k} \equiv \left(\frac{5}{p^a} \right) \pmod{p^2}.$$

If $p \equiv 1, 3 \pmod{10}$ or $a > 2$, then

$$\sum_{k=0}^{\lfloor \frac{9}{10}p^a \rfloor} \frac{\binom{2k}{k}}{(-16)^k} \equiv \left(\frac{5}{p^a} \right) \pmod{p^2}.$$

2. PROOF OF THEOREM 1.1

Lemma 2.1. *Let p be an odd prime and let $k \in \{0, \dots, (p^a - 1)/2\}$ with $a \in \mathbb{Z}^+$. Then*

$$\begin{aligned} & \binom{(p^a - 1)/2 + k}{2k} - \frac{\binom{2k}{k}}{(-16)^k} \\ & \equiv (-1)^{k-1} \left(\frac{-1}{p^a} \right) \binom{p^a - 1 - 2k}{(p^a - 1)/2 - k} \sum_{0 < j \leq k} \frac{p^{2a}}{(2j-1)^2} \pmod{p^3}. \end{aligned} \quad (2.1)$$

Proof. Clearly

$$\begin{aligned} \binom{(p^a - 1)/2 + k}{2k} &= \frac{\prod_{j=1}^k (p^{2a} - (2j-1)^2)}{4^k (2k)!} \\ &= \frac{\prod_{j=1}^k (-(2j-1)^2)}{4^k (2k)!} \prod_{j=1}^k \left(1 - \frac{p^{2a}}{(2j-1)^2} \right) \\ &\equiv \frac{\binom{2k}{k}}{(-16)^k} \left(1 - \sum_{j=1}^k \frac{p^{2a}}{(2j-1)^2} \right) \pmod{p^4} \end{aligned}$$

(which was observed by the author's brother Z. H. Sun [S11, Lemma 2.2] in the case $a = 1$), and in view of (1.3) we have

$$\begin{aligned} \frac{\binom{2k}{k}}{(-16)^k} &\equiv \binom{(p^a - 1)/2}{k} 4^{-k} = \binom{(p^a - 1)/2}{(p^a - 1)/2 - k} 4^{-k} \\ &\equiv \binom{p^a - 1 - 2k}{(p^a - 1)/2 - k} (-4)^{-((p^a - 1)/2 - k)} 4^{-k} \\ &\equiv \left(\frac{-1}{p^a} \right) (-1)^k \binom{p^a - 1 - 2k}{(p^a - 1)/2 - k} \pmod{p}. \end{aligned}$$

So (2.1) holds. \square

Lemma 2.2. *Let p be an odd prime and let $a \in \mathbb{Z}^+$. Then*

$$1 + \frac{2^{p^a-1} - 1}{6} + \frac{(2^{p^a-1} - 1)^2}{24} \equiv \left(\frac{2}{p^a} \right) \frac{2^{p^a+1}}{3 \times 2^{(p^a-1)/2}} \pmod{p^3}. \quad (2.2)$$

When $p \neq 5$, we have

$$\frac{L_{p^a} - 1}{5} - \left(\frac{p^a}{5} \right) F_{p^a} + 1 \equiv -\frac{1}{2} F_{p^a - (\frac{p^a}{5})}^2 \pmod{p^4}. \quad (2.3)$$

Proof. Note that

$$2^{(p^a-1)/2} = \left(2^{\frac{p-1}{2}}\right)^{\sum_{k=0}^{a-1} p^k} \equiv \left(\frac{2}{p}\right)^a = \left(\frac{2}{p^a}\right) \pmod{p}$$

and

$$\begin{aligned} \frac{2^{p^a-1} - 1}{p} &= \frac{2^{(p^a-1)/2} - \left(\frac{2}{p^a}\right)}{p} \left(2^{(p^a-1)/2} + \left(\frac{2}{p^a}\right)\right) \\ &\equiv 2 \left(\frac{2}{p^a}\right) \frac{2^{(p^a-1)/2} - \left(\frac{2}{p^a}\right)}{p} \pmod{p}. \end{aligned}$$

Thus

$$\begin{aligned} (2^{p^a-1} - 1)^2 &\equiv 4 \left(2^{(p^a-1)/2} - \left(\frac{2}{p^a}\right)\right)^2 \\ &= 4(2^{p^a-1} - 1) + 8 - 8 \left(\frac{2}{p^a}\right) 2^{(p^a-1)/2} \pmod{p^3}. \end{aligned}$$

It follows that

$$\begin{aligned} &\frac{2^{(p^a-1)/2}}{2} (2^{p^a-1} - 1) + \frac{1}{8} \left(\frac{2}{p^a}\right) (2^{p^a-1} - 1)^2 \\ &\equiv \left(\frac{2}{p^a}\right) (2^{p^a} + 1) - 3 \times 2^{(p^a-1)/2} \pmod{p^3}, \end{aligned}$$

which is equivalent to (2.2) times $3 \times 2^{(p^a-1)/2}$. So (2.2) is valid.

Now assume that $p \neq 5$. As $L_{2n} = 5F_n^2 + 2(-1)^n = L_n^2 - 2(-1)^n$ for all $n \in \mathbb{N}$, by [SS, Corollary 1] we have $L_{p-(\frac{5}{p})} \equiv 2(\frac{p}{5}) \pmod{p^2}$. Thus, in view of [Su10, Lemma 2.3],

$$L_{p^a-(\frac{p^a}{5})} \equiv (-1)^{((\frac{5}{p})-(\frac{5}{p^a}))/2} L_{p-(\frac{5}{p})} \equiv 2 \left(\frac{p^a}{5}\right) \pmod{p^2}.$$

Write $L_{p^a-(\frac{p^a}{5})} = 2(\frac{p^a}{5}) + p^2Q$ with $Q \in \mathbb{Z}$. Then

$$\begin{aligned} 5F_{p^a-(\frac{p^a}{5})}^2 &= L_{p^a-(\frac{p^a}{5})}^2 - 4(-1)^{p^a-(\frac{p^a}{5})} \\ &= -4 + \left(2 \left(\frac{p^a}{5}\right) + p^2Q\right)^2 \equiv 4p^2 \left(\frac{p^a}{5}\right) Q \pmod{p^4}. \end{aligned}$$

Note that

$$L_{p^a} = F_{p^a} + 2F_{p^a-1} = 2F_{p^a+1} - F_{p^a} = 2F_{p^a-(\frac{p^a}{5})} + \left(\frac{p^a}{5}\right) F_{p^a}$$

and

$$2L_{p^a} = 5F_{p^a-1} + L_{p^a-1} = 5F_{p^a+1} - L_{p^a+1} = 5F_{p^a-(\frac{p^a}{5})} + \left(\frac{p^a}{5}\right) L_{p^a-(\frac{p^a}{5})}.$$

Therefore

$$\begin{aligned} & \frac{L_{p^a} - 1}{5} - \left(\frac{p^a}{5}\right) F_{p^a} + 1 \\ &= 2F_{p^a-(\frac{p^a}{5})} - \frac{4}{5}(L_{p^a} - 1) \\ &= 2F_{p^a-(\frac{p^a}{5})} - \frac{4}{5} \left(\frac{5}{2}F_{p^a-(\frac{p^a}{5})} + \frac{1}{2} \left(\frac{p^a}{5}\right) L_{p^a-(\frac{p^a}{5})} - 1 \right) \\ &= \frac{4}{5} - \frac{2}{5} \left(\frac{p^a}{5}\right) \left(2 \left(\frac{p^a}{5}\right) + p^2 Q \right) \equiv -\frac{1}{2} F_{p^a-(\frac{p^a}{5})}^2 \pmod{p^4}. \end{aligned}$$

This proves (2.3). \square

The following Lemma was posed as [Su11c, Conjecture 1.1].

Lemma 2.3. *Let p be an odd prime and let $H_k^{(2)} = \sum_{0 < j \leq k} 1/j^2$ for $k = 0, 1, 2, \dots$. If $p \neq 5$ then*

$$\sum_{k=0}^{p-1} (-1)^k \binom{2k}{k} H_k^{(2)} \equiv \left(\frac{p}{5}\right) \frac{5}{2} q^2 \pmod{p}, \quad (2.4)$$

where q denotes the Fibonacci quotient $F_{p-(\frac{p}{5})}/p$. If $p > 3$, then

$$\sum_{k=0}^{p-1} (-2)^k \binom{2k}{k} H_k^{(2)} \equiv \frac{2}{3} q_p(2)^2 \pmod{p}, \quad (2.5)$$

where $q_p(2)$ denotes the Fermat quotient $(2^{p-1} - 1)/p$.

Proof. The desired congruences essentially follow from [MT, (36)]. Here we provide the details. Putting $t = -1, -1/2$ in [MT, (36)] we get

$$\sum_{k=0}^{p-1} (-1)^k \binom{2k}{k} H_k^{(2)} \equiv -2 \sum_{k=1}^{p-1} \frac{u_k(3, 1)}{k^2} \pmod{p} \quad (2.6)$$

and

$$\sum_{k=0}^{p-1} (-2)^k \binom{2k}{k} H_k^{(2)} \equiv -2 \sum_{k=1}^{p-1} \frac{u_k(5/2, 1)}{k^2} \pmod{p}. \quad (2.7)$$

Note that $u_k(3, 1) = F_{2k}$ and $u_k(5/2, 1) = 2(2^k - 2^{-k})/3$ for all $k = 0, 1, 2, \dots$

If $p > 3$, then

$$\sum_{k=1}^{p-1} \frac{2^k}{k^2} \equiv -q_p(2)^2 \pmod{p} \quad \text{and} \quad \sum_{k=1}^{p-1} \frac{1}{k^2 2^k} \equiv -\frac{q_p(2)^2}{2} \pmod{p}$$

by [Gr] and [S08, Theorem 4.1(iv)] respectively, hence (2.7) implies that

$$\sum_{k=0}^{p-1} (-2)^k \binom{2k}{k} H_k^{(2)} \equiv -\frac{4}{3} \sum_{k=1}^{p-1} \left(\frac{2^k}{k^2} - \frac{1}{k^2 2^k} \right) \equiv \frac{2}{3} q_p(2)^2 \pmod{p}.$$

Below we assume that $p \neq 5$. Recall that for any $n \in \mathbb{N}$ we have

$$F_n = \frac{\alpha^n - \beta^n}{\alpha - \beta} \quad \text{and} \quad L_n = \alpha^n + \beta^n,$$

where α and β are the two roots of the equation $x^2 - x - 1 = 0$. By [PS, (3.2) and (3.7)],

$$2\beta^{2p} \sum_{k=1}^{p-1} \frac{\alpha^{2k}}{k^2} \equiv -2 \sum_{k=1}^{p-1} \frac{\alpha^k}{k^2} - \left(\frac{L_p - 1}{p} \right)^2 \equiv \left(\frac{2\alpha^p - 1}{5} - 1 \right) \left(\frac{L_p - 1}{p} \right)^2.$$

Since $\alpha\beta = -1$ and $\alpha^{2p} = \alpha^p + 1$, we have

$$\sum_{k=1}^{p-1} \frac{\alpha^{2k}}{k^2} \equiv \frac{(\alpha^p + 1)(\alpha^p - 3)}{5} \left(\frac{L_p - 1}{p} \right)^2 = -\frac{\alpha^p + 2}{5} \left(\frac{L_p - 1}{p} \right)^2 \pmod{p}.$$

Hence

$$\begin{aligned} 5 \sum_{k=1}^{p-1} \frac{F_{2k}}{k^2} &= (\alpha - \beta)^2 \sum_{k=1}^{p-1} \frac{F_{2k}}{k^2} = (\alpha - \beta) \sum_{k=1}^{p-1} \frac{\alpha^{2k} - \beta^{2k}}{k^2} \\ &\equiv (\alpha - \beta) \frac{\beta^p - \alpha^p}{5} \left(\frac{L_p - 1}{p} \right)^2 \equiv -\frac{(\alpha - \beta)^{p+1}}{5} \left(\frac{L_p - 1}{p} \right)^2 \\ &= -5^{(p-1)/2} \left(\frac{L_p - 1}{p} \right)^2 \equiv -\left(\frac{5}{p} \right) \frac{25}{4} q^2 \pmod{p}. \end{aligned}$$

since $2(L_p - 1) \equiv 5F_{p-(\frac{p}{5})} \pmod{p^2}$ by [ST, p. 139]. Combining this with (2.6) we obtain

$$\sum_{k=0}^{p-1} (-1)^k \binom{2k}{k} H_k^{(2)} \equiv 2 \times \left(\frac{5}{p} \right) \frac{5}{4} q^2 = \left(\frac{p}{5} \right) \frac{5}{2} q^2 \pmod{p}.$$

The proof of Lemma 2.3 is now complete. \square

Proof of Theorem 1.1. Let us first recall the following two identities:

$$F_{n+1} = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n-k}{k} \quad \text{and} \quad \sum_{k=0}^n \frac{\binom{n+k}{2k}}{2^k} = \frac{2^{2n+1} + 1}{3 \times 2^n}.$$

Thus we have

$$F_{p^a} = \sum_{k=0}^{(p^a-1)/2} \binom{p^a-1-k}{p^a-1-2k} = \sum_{j=0}^{(p^a-1)/2} \binom{(p^a-1)/2+j}{2j}$$

and

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{(p^a-1)/2+k}{2k}}{2^k} = \frac{2^{p^a} + 1}{32^{(p^a-1)/2}}$$

Therefore, with the help of (2.1),

$$\begin{aligned} & \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-16)^k} - F_{p^a} \\ & \equiv \sum_{k=0}^{(p^a-1)/2} \binom{(p^a-1-2k)}{(p^a-1)/2-k} (-1)^{(p^a-1)/2-k} \sum_{j=1}^k \frac{p^{2a}}{(2j-1)^2} \\ & = \sum_{k=0}^{(p^a-1)/2} \binom{2k}{k} (-1)^k \sum_{j=1}^{(p^a-1)/2-k} \frac{p^{2a}}{(2j-1)^2} \pmod{p^3} \end{aligned}$$

and

$$\begin{aligned} & \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-32)^k} - \frac{2^{p^a} + 1}{3 \times 2^{(p^a-1)/2}} \\ & \equiv \sum_{k=0}^{(p^a-1)/2} \binom{2k}{k} \frac{(-1)^k}{2^{(p^a-1)/2-k}} \sum_{j=1}^{(p^a-1)/2-k} \frac{p^{2a}}{(2j-1)^2} \pmod{p^3}. \end{aligned}$$

For $k = 0, \dots, (p^a-1)/2$, clearly

$$\begin{aligned} & \sum_{j=1}^{(p^a-1)/2-k} \frac{p^{2a}}{(2j-1)^2} = \sum_{j=k+1}^{(p^a-1)/2} \frac{p^{2a}}{(2((p^a-1)/2-j+1)-1)^2} \\ & \equiv \sum_{j=1}^{(p^a-1)/2} \frac{p^{2a}}{4j^2} - \sum_{j=1}^k \frac{p^{2a}}{4j^2} \\ & \equiv \sum_{i=1}^{(p-1)/2} \frac{p^{2a}}{4(p^{a-1}i)^2} - \sum_{i=1}^{\lfloor k/p^{a-1} \rfloor} \frac{p^{2a}}{4(p^{a-1}i)^2} \pmod{p^3}. \end{aligned}$$

Since

$$2 \sum_{i=1}^{(p-1)/2} \frac{1}{i^2} = \sum_{i=1}^{(p-1)/2} \left(\frac{1}{i^2} + \frac{1}{(p-i)^2} \right) = \sum_{i=1}^{p-1} \frac{1}{i^2} \equiv 0 \pmod{p},$$

by the above we have

$$\frac{-4}{p^2} \left(\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-16)^k} - F_{p^a} \right) \equiv \sum_{k=0}^{p^a-1} \binom{2k}{k} (-1)^k H_{\lfloor k/p^{a-1} \rfloor}^{(2)} \pmod{p}$$

and

$$\begin{aligned} & \frac{-4}{p^2} \left(\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-32)^k} - \frac{2^{p^a} + 1}{3 \times 2^{(p^a-1)/2}} \right) \\ & \equiv \left(\frac{2}{p^a} \right) \sum_{k=0}^{p^a-1} \binom{2k}{k} (-2)^k H_{\lfloor k/p^{a-1} \rfloor}^{(2)} \pmod{p}. \end{aligned}$$

Let $u \in \{1, 2\}$. Then

$$\begin{aligned} & \sum_{k=0}^{p^a-1} \binom{2k}{k} (-u)^k H_{\lfloor k/p^{a-1} \rfloor}^{(2)} \\ & = \sum_{k=0}^{p-1} \sum_{r=0}^{p^{a-1}-1} \binom{2p^{a-1}k + 2r}{p^{a-1}k + r} (-u)^{p^{a-1}k+r} H_k^{(2)} \\ & \equiv \sum_{k=0}^{p-1} (-u)^k H_k^{(2)} \sum_{r=0}^{p^{a-1}-1} \binom{2p^{a-1}k + 2r}{p^{a-1}k + r} (-u)^r \pmod{p}. \end{aligned}$$

For $k \in \{0, \dots, p-1\}$ and $r \in \{0, \dots, p^{a-1}-1\}$, by the Chu-Vandermonde identity (cf. [GKP, p. 169]),

$$\binom{2p^{a-1}k + 2r}{p^{a-1}k + r} = \sum_{j=0}^{p^{a-1}k+r} \binom{2p^{a-1}k}{j} \binom{2r}{p^{a-1}k + r - j}.$$

If $p^{a-1} \nmid j$ then

$$\binom{2p^{a-1}k}{j} = \frac{2p^{a-1}k}{j} \binom{2p^{a-1}k - 1}{j-1} \equiv 0 \pmod{p}.$$

Thus

$$\begin{aligned}
\binom{2p^{a-1}k + 2r}{p^{a-1}k + r} &= \sum_{j=0}^{p^{a-1}k+r} \binom{2p^{a-1}k}{j} \binom{2r}{p^{a-1}k + r - j} \\
&\equiv \sum_{j=0}^k \binom{2p^{a-1}k}{p^{a-1}j} \binom{2r}{p^{a-1}(k-j) + r} = \binom{2p^{a-1}k}{p^{a-1}k} \binom{2r}{r} \\
&\equiv \binom{2k}{k} \binom{2r}{r} \pmod{p} \quad (\text{by Lucas' theorem}).
\end{aligned}$$

Therefore

$$\begin{aligned}
&\sum_{k=0}^{p-1} (-u)^k H_k^{(2)} \sum_{r=0}^{p^{a-1}-1} \binom{2p^{a-1}k + 2r}{p^{a-1}k + r} \\
&\equiv \sum_{k=0}^{p-1} (-u)^k H_k^{(2)} \binom{2k}{k} \sum_{r=0}^{p^{a-1}-1} \binom{2r}{r} (-u)^r \pmod{p}.
\end{aligned}$$

By a basic result mentioned in Section 1,

$$\sum_{r=0}^{p^{a-1}-1} \frac{\binom{2r}{r}}{m^r} \equiv \left(\frac{m(m-4)}{p^{a-1}} \right) \pmod{p}$$

for any integer $m \not\equiv 0 \pmod{p}$. Thus

$$\sum_{r=0}^{p^{a-1}-1} \binom{2r}{r} (-1)^r \equiv \left(\frac{5}{p^{a-1}} \right) \pmod{p}$$

and

$$\sum_{r=0}^{p^{a-1}-1} \binom{2r}{r} (-2)^r \equiv \left(\frac{(p-1)/2 \times ((p-1)/2 - 4)}{p^{a-1}} \right) = 1 \pmod{p}.$$

Therefore

$$\begin{aligned}
&\frac{-4}{p^2} \left(\sum_{k=0}^{(p^{a-1})/2} \frac{\binom{2k}{k}}{(-16)^k} - F_{p^a} \right) \\
&\equiv \sum_{k=0}^{p-1} \binom{2k}{k} (-1)^k H_k^{(2)} \sum_{r=0}^{p^{a-1}-1} \binom{2r}{r} (-1)^r \\
&\equiv \sum_{k=0}^{p-1} \binom{2k}{k} (-1)^k H_k^{(2)} \left(\frac{5}{p^{a-1}} \right) \pmod{p},
\end{aligned}$$

and similarly

$$\frac{-4}{p^2} \left(\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-32)^k} - \frac{2^{p^a} + 1}{32(p^a-1)/2} \right) \equiv \left(\frac{2}{p^a} \right) \sum_{k=0}^{p-1} \binom{2k}{k} (-2)^k H_k^{(2)} \pmod{p}.$$

Now assume that $p \neq 3$. By the last congruence and (2.5),

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-32)^k} - \frac{2^{p^a} + 1}{3 \times 2^{(p^a-1)/2}} \equiv - \left(\frac{2}{p^a} \right) \frac{p^2}{6} q_p(2)^2 \pmod{p^3}.$$

Since $p^a \equiv p \pmod{\varphi(p^2)}$, we have $2^{p^a} \equiv 2^p \pmod{p^2}$ and hence

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-32)^k} - \frac{2^{p^a} + 1}{3 \times 2^{(p^a-1)/2}} \equiv - \left(\frac{2}{p^a} \right) \frac{(2^{p^a-1} - 1)^2}{6} \pmod{p^3}.$$

Combining this with (2.2) we immediately obtain (1.2).

Below we suppose that $p \neq 5$. By (2.4) and the above,

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-16)^k} - F_{p^a} \equiv - \frac{5}{8} \left(\frac{p^a}{5} \right) F_{p-(\frac{p}{5})}^2 \pmod{p^3}.$$

In view of [Su10, Lemma 2.3],

$$\frac{F_{p-(\frac{p}{5})}}{p} \equiv (-1)^{((\frac{5}{p}) - (\frac{5}{p^a})) / 2} \left(\frac{5}{p^{a-1}} \right) \frac{F_{p-(\frac{p}{5})}}{p} = \frac{F_{p-(\frac{p}{5})}}{p} \pmod{p}$$

and thus

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-16)^k} - F_{p^a} \equiv - \frac{5}{8} \left(\frac{p^a}{5} \right) F_{p^a-(\frac{p^a}{5})}^2 \pmod{p^3}.$$

Combining this with (2.3) we get

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-16)^k} - F_{p^a} \equiv \frac{L_{p^a} - 1}{4} \left(\frac{p^a}{5} \right) - \frac{5}{4} F_{p^a} + \frac{5}{4} \left(\frac{p^a}{5} \right) \pmod{p^3}.$$

Therefore

$$\begin{aligned} \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{(-16)^k} &\equiv \left(\frac{p^a}{5} \right) \frac{L_{p^a}}{4} - \frac{F_{p^a}}{4} + \left(\frac{p^a}{5} \right) \\ &= \left(\frac{p^a}{5} \right) \left(1 + \frac{1}{4} \left(L_{p^a} - \left(\frac{p^a}{5} \right) F_{p^a} \right) \right) \\ &= \left(\frac{p^a}{5} \right) \left(1 + \frac{1}{2} F_{p^a-(\frac{p^a}{5})} \right) \pmod{p^3}. \end{aligned}$$

This proves (1.1).

So far we have completed the proof of Theorem 1.1. \square

3. PROOF OF THEOREM 1.2

We need some knowledge of Lucas sequences.

Let $A, B \in \mathbb{Z}$ and $\Delta = A^2 - 4B$. The equation $x^2 - Ax + B = 0$ has two roots

$$\alpha = \frac{A + \sqrt{\Delta}}{2} \quad \text{and} \quad \beta = \frac{A - \sqrt{\Delta}}{2}$$

which are algebraic integers. It is well known that for any $n \in \mathbb{N}$ we have

$$u_n(A, B) = \sum_{0 \leq k < n} \alpha^k \beta^{n-1-k} \quad \text{and} \quad v_n(A, B) = \alpha^n + \beta^n.$$

If p is a prime then

$$v_p(A, B) = \alpha^p + \beta^p \equiv (\alpha + \beta)^p = A^p \equiv A \pmod{p}.$$

Lemma 3.1. *Let $A, B \in \mathbb{Z}$ and $n \in \mathbb{N}$. Then*

$$u_{n+1}(A, B) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n-k}{k} A^{n-2k} (-B)^k. \quad (3.1)$$

Remark 3.1. (3.1) is a well-known formula due to Lagrange, see, e.g., H. Gould [G, (1.60)].

Lemma 3.2. *Let $A, B \in \mathbb{Z}$ and let p be an odd prime not dividing $B\Delta$ where $\Delta = A^2 - 4B$. Then*

$$u_p(A, B) \equiv \frac{A}{2} B^{((\frac{\Delta}{p})-1)/2} u_{p-(\frac{\Delta}{p})}(A, B) + \left(\frac{\Delta}{p}\right) \frac{B^{p-1} + 1}{2} \pmod{p^2}. \quad (3.2)$$

Proof. For convenience we let $u_n = u_n(A, B)$ and $v_n = v_n(A, B)$ for all $n \in \mathbb{N}$.

Let α and β be the two roots of the equation $x^2 - Ax + B = 0$. Then

$$v_n^2 - \Delta u_n^2 = (\alpha^n + \beta^n)^2 - (\alpha^n - \beta^n)^2 = 4(\alpha\beta)^n = 4B^n$$

for any $n \in \mathbb{N}$. As $p \mid u_{p-(\frac{\Delta}{p})}$ (see, e.g., [Su10, Lemma 2.3]), p^2 divides

$$\begin{aligned} & v_{p-(\frac{\Delta}{p})}^2 - 4B^{p-(\frac{\Delta}{p})} \\ &= \left(v_{p-(\frac{\Delta}{p})} - 2 \left(\frac{B}{p}\right) B^{(p-(\frac{\Delta}{p}))/2} \right) \left(v_{p-(\frac{\Delta}{p})} + 2 \left(\frac{B}{p}\right) B^{(p-(\frac{\Delta}{p}))/2} \right). \end{aligned}$$

On the other hand, by [Su10, Lemma 2.3] we have

$$v_{p-(\frac{\Delta}{p})} \equiv 2B^{(1-(\frac{\Delta}{p}))/2} \equiv 2\left(\frac{B}{p}\right)B^{(p-(\frac{\Delta}{p}))/2} \pmod{p}.$$

Therefore

$$v_{p-(\frac{\Delta}{p})} \equiv 2\left(\frac{B}{p}\right)B^{(p-(\frac{\Delta}{p}))/2} \pmod{p^2}.$$

By induction, for $\varepsilon = \pm 1$ we have

$$Au_n + \varepsilon v_n = 2B^{(1-\varepsilon)/2}u_{n+\varepsilon}$$

for all $n \in \mathbb{Z}^+$. Thus

$$\begin{aligned} 2B^{(1-(\frac{\Delta}{p}))/2}u_p &= Au_{p-(\frac{\Delta}{p})} + \left(\frac{\Delta}{p}\right)v_{p-(\frac{\Delta}{p})} \\ &\equiv Au_{p-(\frac{\Delta}{p})} + \left(\frac{\Delta}{p}\right)2\left(\frac{B}{p}\right)B^{(p-(\frac{\Delta}{p}))/2} \pmod{p^2} \end{aligned}$$

and hence

$$\begin{aligned} 2u_p - AB^{((\frac{\Delta}{p})-1)/2}u_{p-(\frac{\Delta}{p})} &\\ \equiv \left(\frac{\Delta}{p}\right)\left(2\left(\frac{B}{p}\right)\left(B^{(p-1)/2} - \left(\frac{B}{p}\right)\right) + 2\right) &\\ \equiv \left(\frac{\Delta}{p}\right)(B^{p-1} - 1 + 2) \pmod{p^2}. & \end{aligned}$$

So (3.2) is valid. \square

Lemma 3.3. *Let p be an odd prime and let $a \in \mathbb{Z}^+$. Let m be an integer not divisible by p . Then*

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k+1}}{m^k} \equiv \frac{m-2}{2} \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{m^k} - \frac{m}{2} + 2p\delta_{a,1}\left(\frac{-m}{p}\right) \pmod{p^2}. \quad (3.3)$$

Proof. Observe that

$$\begin{aligned} &\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k} + \binom{2k}{k+1}}{m^k} \\ &= \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k+1}{k+1}}{m^k} = \frac{\binom{p^a}{(p^a+1)/2}}{m^{(p^a-1)/2}} + \frac{1}{2} \sum_{k=0}^{(p^a-3)/2} \frac{\binom{2k+2}{k+1}}{m^k} \\ &= \frac{p^a/m^{(p^a-1)/2}}{(p^a+1)/2} \binom{p^a-1}{(p^a-1)/2} + \frac{m}{2} \sum_{k=1}^{(p^a-1)/2} \frac{\binom{2k}{k}}{m^k} \\ &\equiv 2p\delta_{a,1}\left(\frac{-m}{p}\right) + \frac{m}{2} \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{m^k} - \frac{m}{2} \pmod{p^2} \end{aligned}$$

and hence (3.3) follows. \square

Proof of Theorem 1.2. Set $n = (p^a - 1)/2$. By Lemma 3.3,

$$\sum_{k=0}^n \frac{C_k}{m^k} \equiv \left(1 - \frac{m-2}{2}\right) \sum_{k=0}^n \frac{\binom{2k}{k}}{m^k} + \frac{m}{2} - 2p\delta_{a,1}\left(\frac{-m}{p}\right) \pmod{p^2}.$$

This proves (1.6). It remains to show (1.4).

By Lemmas 3.1 and 2.1,

$$\begin{aligned} u_{p^a}(4, m) &= \sum_{k=0}^n \binom{2n-k}{k} 4^{2n-2k} (-m)^k \\ &= \sum_{k=0}^n \binom{2n-k}{2(n-k)} 16^{n-k} (-m)^k = \sum_{k=0}^n \binom{n+k}{2k} 16^k (-m)^{n-k} \\ &\equiv \sum_{k=0}^n \binom{2k}{k} (-1)^k (-m)^{n-k} = (-m)^n \sum_{k=0}^n \frac{\binom{2k}{k}}{m^k} \pmod{p^2}. \end{aligned}$$

Note that

$$(-m)^n = \left((-m)^{(p-1)/2}\right)^{\sum_{s=0}^{a-1} p^s} \equiv \left(\frac{-m}{p}\right)^{\sum_{s=0}^{a-1} p^s} = \left(\frac{-m}{p^a}\right) \pmod{p}$$

and hence

$$\begin{aligned} (-m)^n - \left(\frac{-m}{p^a}\right) &\equiv \left((-m)^n - \left(\frac{-m}{p^a}\right)\right) \frac{(-m)^n + \left(\frac{-m}{p^a}\right)}{2\left(\frac{-m}{p^a}\right)} \\ &\equiv \frac{(-m)^{p^a-1} - 1}{2} \left(\frac{-m}{p^a}\right) \pmod{p^2}. \end{aligned}$$

Thus

$$(-m)^n \equiv \left(\frac{-m}{p^a}\right) \left(1 + \frac{m^{p^a-1} - 1}{2}\right) \equiv \frac{\left(\frac{-m}{p^a}\right)}{1 - (m^{p^a-1} - 1)/2} \pmod{p^2}$$

and hence

$$\begin{aligned} \sum_{k=0}^n \frac{\binom{2k}{k}}{m^k} &\equiv u_{p^a}(4, m) \left(\frac{-m}{p^a}\right) \left(1 - \frac{m^{p^a-1} - 1}{2}\right) \\ &\equiv u_{p^a}(4, m) \left(\frac{-m}{p^a}\right) \left(1 - \frac{m^{p-1} - 1}{2}\right) \end{aligned}$$

since $m^{p^a-1} \equiv m^{p-1} \pmod{p^2}$ by Euler's theorem. By [Su10, Lemma 2.3],

$$\begin{aligned} u_{p^a}(4, m) &\equiv \left(\frac{4^2 - 4m}{p^{a-1}} \right) u_p(4, m) \pmod{p^2} \\ &\equiv \left(\frac{4^2 - 4m}{p^a} \right) u_1(4, m) = \left(\frac{4-m}{p^a} \right) \pmod{p} \end{aligned}$$

Therefore

$$\begin{aligned} \sum_{k=0}^n \frac{\binom{2k}{k}}{m^k} &\equiv u_{p^a}(4, m) \left(\frac{-m}{p^a} \right) - u_{p^a}(4, m) \left(\frac{-m}{p^a} \right) \frac{m^{p-1} - 1}{2} \\ &\equiv \left(\frac{4-m}{p^{a-1}} \right) \left(\frac{-m}{p^a} \right) u_p(4, m) - \left(\frac{-m(4-m)}{p^a} \right) \frac{m^{p-1} - 1}{2} \\ &\equiv \left(\frac{-m}{p} \right) \left(\frac{m(m-4)}{p^{a-1}} \right) u_p(4, m) - \left(\frac{m(m-4)}{p^a} \right) \frac{m^{p-1} - 1}{2} \pmod{p^2}. \end{aligned}$$

In view of Lemma 3.2,

$$u_p(4, m) - \left(\frac{4-m}{p} \right) \frac{m^{p-1} - 1}{2} \equiv \bar{m} u_{p-(\frac{4-m}{p})}(4, m) + \left(\frac{4-m}{p} \right) \pmod{p^2}.$$

So, by the above, $\sum_{k=0}^n \binom{2k}{k}/m^k$ is congruent to

$$\begin{aligned} &\left(\frac{m(m-4)}{p^{a-1}} \right) \left(\frac{-m}{p} \right) \left(\bar{m} u_{p-(\frac{4-m}{p})}(4, m) + \left(\frac{4-m}{p} \right) \right) \\ &= \left(\frac{m(m-4)}{p^a} \right) + \left(\frac{-m}{p} \right) \left(\frac{m(m-4)}{p^{a-1}} \right) \bar{m} u_{p-(\frac{4-m}{p})}(4, m) \end{aligned}$$

modulo p^2 . This proves (1.4). We are done. \square

4. PROOFS OF COROLLARIES 1.1-1.3 AND FINAL REMARKS

Proof of Corollary 1.1. Note that $n = p - (\frac{4-8}{p}) \equiv 0 \pmod{4}$. The equation $x^2 - 4x + 8 = 0$ has two roots $2 \pm 2i$ where $i = \sqrt{-1}$. Thus

$$u_n(4, 8) = \frac{(2+2i)^n - (2-2i)^n}{4i} = \frac{(i(2-2i))^n - (2-2i)^n}{4i} = 0$$

and hence by Theorem 1.2 we have

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{8^k} \equiv \left(\frac{8(8-4)}{p^a} \right) = \left(\frac{2}{p^a} \right) \pmod{p^2}.$$

Clearly $k = p - (\frac{4-16}{p}) = p - (\frac{p}{3})$ is divisible by 3 and the two roots of the equation $x^2 - 4x + 16 = 0$ are

$$2 + 2\sqrt{-3} = -4\omega^2 \text{ and } 2 - 2\sqrt{-3} = -4\omega,$$

where $\omega = (-1 + \sqrt{-3})/2$ is a primitive cubic root of unity. Thus

$$u_k(4, 16) = \frac{(-4\omega^2)^n - (-4\omega)^n}{4\sqrt{-3}} = 0$$

since $3 \mid n$. Applying (1.4) with $m = 16$ we get

$$\sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{16^k} \equiv \left(\frac{16(16-4)}{p^a} \right) = \left(\frac{3}{p^a} \right) \pmod{p^2}.$$

The proof of Corollary 1.1 is now complete. \square

Proof of Corollary 1.2. Set $n = (p-1)/2$. Then

$$\begin{aligned} & \sum_{k=1}^n \frac{\binom{2k}{k}}{16^k} \left(\frac{1}{2k-1} + \frac{1}{(2k-1)^2} \right) \\ &= \sum_{k=1}^n \frac{2\binom{2k-1}{k}}{16^k} \cdot \frac{2k}{(2k-1)^2} = \frac{1}{4} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{(2k+1)16^k} \equiv 0 \pmod{p^2} \end{aligned}$$

with the help of [Su11b, (1.4)].

Observe that

$$\begin{aligned} \sum_{k=1}^n \frac{\binom{2k}{k}}{(2k-1)16^k} &= \sum_{k=1}^n \frac{2\binom{2k-1}{k}}{(2k-1)16^k} = 2 \sum_{k=1}^n \frac{\binom{2k-2}{k-1}}{k16^k} \\ &= 2 \sum_{j=0}^{n-1} \frac{C_j}{16^{j+1}} = \frac{1}{8} \sum_{k=0}^n \frac{C_k}{16^k} - \frac{C_n}{8 \times 4^{2n}}. \end{aligned}$$

Also,

$$\frac{C_n}{4^{2n}} = \frac{\binom{p-1}{(p-1)/2}}{4^{p-1}(p+1)/2} \equiv (-1)^{(p-1)/2} 2(1-p) \pmod{p^2}$$

in view of Morley's congruence ([Mo])

$$\binom{p-1}{(p-1)/2} \equiv (-1)^{(p-1)/2} 4^{p-1} \pmod{p^3}.$$

By (1.6) and (1.8),

$$\sum_{k=0}^n \frac{C_k}{16^k} \equiv -6 \left(\frac{3}{p} \right) + 8 - 2p \left(\frac{-1}{p} \right) \pmod{p^2}.$$

Therefore

$$\begin{aligned} \sum_{k=1}^n \frac{\binom{2k}{k}}{(2k-1)16^k} &\equiv \frac{8 - 6\left(\frac{3}{p}\right) - 2p\left(\frac{-1}{p}\right)}{8} - \frac{2(1-p)\left(\frac{-1}{p}\right)}{8} \\ &= 1 - \frac{\left(\frac{-1}{p}\right) + 3\left(\frac{3}{p}\right)}{4} = 1 - \left(\frac{-1}{p}\right) \frac{3\left(\frac{p}{3}\right) + 1}{4} \pmod{p^2} \end{aligned}$$

and hence

$$\sum_{k=1}^n \frac{\binom{2k}{k}}{(2k-1)^2 16^k} \equiv - \sum_{k=1}^n \frac{\binom{2k}{k}}{(2k-1)16^k} \equiv -1 + \left(\frac{-1}{p}\right) \frac{3\left(\frac{p}{3}\right) + 1}{4} \pmod{p^2},$$

which yields (1.9) and its equivalent form (1.10). We are done. \square

Now let us make some final remarks.

For $m \in \mathbb{Z}$ and $n \in \mathbb{N}$, we have the identity

$$\sum_{k=0}^n \left(1 - \frac{m-4}{2}k \right) \frac{\binom{2k}{k}}{m^k} = (2n+1) \frac{\binom{2n}{n}}{m^n}, \quad (4.1)$$

which can be easily proved by induction. Using this identity one can easily deduce the following result.

Proposition 4.1. *Let p be an odd prime and let $a \in \mathbb{Z}^+$. Let $m \in \mathbb{Z}$ with $m \not\equiv 0 \pmod{p}$. Then*

$$\frac{m-4}{2} \sum_{k=0}^{(p^a-1)/2} \frac{k \binom{2k}{k}}{m^k} \equiv \sum_{k=0}^{(p^a-1)/2} \frac{\binom{2k}{k}}{m^k} - p^a \left(\frac{-m}{p^a} \right) \pmod{p^{a+1}} \quad (4.2)$$

and

$$\frac{m-4}{2} \sum_{k=0}^{p^a-1} \frac{k \binom{2k}{k}}{m^k} \equiv \sum_{k=0}^{p^a-1} \frac{\binom{2k}{k}}{m^k} - p^a \pmod{p^{a+1}}. \quad (4.3)$$

Let p be an odd prime and let $m \in \mathbb{Z}$ with $p \nmid m$. Since we have determined $\sum_{k=0}^{(p-1)/2} \binom{2k}{k}/m^k \pmod{p^2}$ and $\sum_{k=0}^{p-1} \binom{2k}{k}/m^k \pmod{p^2}$, with

the help of Proposition 4.1 we know the sums $\sum_{k=0}^{(p-1)/2} k \binom{2k}{k} / m^k$ and $\sum_{k=0}^{p-1} k \binom{2k}{k} / m^k$ modulo p^2 . For example, if $p > 3$ then

$$\sum_{k=0}^{p-1} \frac{k \binom{2k}{k}}{2^k} \equiv p - \left(\frac{-1}{p} \right) \pmod{p^2}, \quad (4.4)$$

$$\sum_{k=0}^{p-1} \frac{k \binom{2k}{k}}{3^k} \equiv 2p - 2 \left(\frac{p}{3} \right) \pmod{p^2}, \quad (4.5)$$

$$\sum_{k=0}^{(p-1)/2} \frac{k \binom{2k}{k}}{8^k} \equiv \left(\frac{2}{p} \right) \frac{1 - (-1)^{(p-1)/2} p}{2} \pmod{p^2}, \quad (4.6)$$

$$\sum_{k=0}^{(p-1)/2} \frac{k \binom{2k}{k}}{16^k} \equiv \frac{\left(\frac{3}{p} \right) - (-1)^{(p-1)/2} p}{6} \pmod{p^2}. \quad (4.7)$$

REFERENCES

- [A] A. Adamchuk, *Comments on OEIS A066796 in 2007*, On-Line Encyclopedia of Integer Sequences, <http://www.research.att.com/~njas/sequences/A066796>.
- [CDP] R. Crandall, K. Dilcher and C. Pomerance, *A search for Wieferich and Wilson primes*, Math. Comp. **66** (1997), 433–449.
- [CP] R. Crandall and C. Pomerance, *Prime Numbers: A Computational Perspective*, Second Edition, Springer, New York, 2005.
- [DK] F. G. Dorais and D. W. Klyve, *Near Wieferich primes up to 6.7×10^{15}* , preprint, 2010. <http://www-personal.umich.edu/~dorais/docs/wieferich.pdf>.
- [G] H. W. Gould, *Combinatorial Identities*, Morgantown Printing and Binding Co., 1972.
- [Gr] A. Granville, *The square of the Fermat quotient*, Integers **4** (2004), #A22, 3pp (electronic).
- [GKP] R. L. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*, 2nd ed., Addison-Wesley, New York, 1994.
- [GZ] V. J. W. Guo and J. Zeng, *Some congruences involving central q-binomial coefficients*, Adv. in Appl. Math. **45** (2010), 303–316.
- [MT] S. Mattarei and R. Tauraso, *Congruences for central binomial sums and finite polylogarithms*, preprint, arXiv:1012.1308. <http://arxiv.org/abs/1012.1308>.
- [Mo] F. Morley, *Note on the congruence $2^{4n} \equiv (-1)^n (2n)!/(n!)^2$, where $2n+1$ is a prime*, Ann. Math. **9** (1895), 168–170.
- [PS] H. Pan and Z. W. Sun, *Proof of three conjectures on congruences*, submitted, arXiv:1010.2489. <http://arxiv.org/abs/1010.2489>.
- [St] R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, Cambridge Univ. Press, Cambridge, 1999.
- [SSZ] N. Strauss, J. Shallit and D. Zagier, *Some strange 3-adic identities*, Amer. Math. Monthly **99** (1992), 66–69.
- [S08] Z. H. Sun, *Congruences involving Bernoulli and Euler numbers*, J. Number Theory **128** (2008), 280–312.
- [S11] Z. H. Sun, *Congruences concerning Legendre polynomials*, Proc. Amer. Math. Soc. **139** (2011), 1915–1929.

- [SS] Z. H. Sun and Z. W. Sun, *Fibonacci numbers and Fermat's last theorem*, Acta Arith. **60** (1992), 371–388.
- [Su06] Z. W. Sun, *Binomial coefficients and quadratic fields*, Proc. Amer. Math. Soc. **134** (2006), 2213–2222.
- [Su10] Z. W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients*, Sci. China Math. **53** (2010), 2473–2488. <http://arxiv.org/abs/0909.5648>.
- [Su11a] Z. W. Sun, *p -adic valuations of some sums of multinomial coefficients*, Acta Arith. **148** (2011), 63–76.
- [Su11b] Z. W. Sun, *On congruences related to central binomial coefficients*, J. Number Theory **131** (2011), 2219–2238.
- [Su11c] Z. W. Sun, *On harmonic numbers and Lucas sequences*, Publ. Math. Debrecen **79** (2011), in press. <http://arxiv.org/abs/1001.0348>.
- [ST] Z. W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math. **45** (2010), 125–148.
- [W] H. C. Williams, *A note on the Fibonacci quotient $F_{p-\varepsilon}/p$* , Canad. Math. Bull. **25** (1982), 366–370.