

Generating trees and pattern avoidance in alternating permutations

Joel Brewster Lewis
Massachusetts Institute of Technology
jblewis@math.mit.edu

May 24, 2010

Abstract

We extend earlier work of the same author to enumerate alternating permutations avoiding the permutation pattern 2143. We use a generating tree approach to construct a recursive bijection between the set $A_{2n}(2143)$ of alternating permutations of length $2n$ avoiding 2143 and standard Young tableaux of shape $\langle n, n, n \rangle$ and between the set $A_{2n+1}(2143)$ of alternating permutations of length $2n + 1$ avoiding 2143 and shifted standard Young tableaux of shape $\langle n + 2, n + 1, n \rangle$. We also give a number of conjectures and open questions on pattern avoidance in alternating permutations and generalizations thereof.

1 Introduction

In [5], the author enumerated the set $A_{2n}(1234)$ of alternating permutations with no four-term increasing subsequence by showing that it is in bijection with the set of standard Young tableaux of shape $\langle n, n, n \rangle$. One of the two proofs given of this result may be understood as a demonstration that these permutations and tableaux have isomorphic generating trees, that is, that they have the same recursive structure. (The other proof used a modification of the RSK correspondence first used by Ouchterlony [6] in the context of doubly-alternating permutations.) In this paper, we show that this approach can be extended to enumerate alternating permutations avoiding the pattern

2143. In particular, we show that $A_{2n}(2143)$ is in bijection with standard Young tableaux of shape $\langle n, n, n \rangle$ (and so also with $A_{2n}(1234)$) and that $A_{2n+1}(2143)$ is in bijection with *shifted* standard Young tableaux of shape $\langle n+2, n+1, n \rangle$. We end with a large number of open problems.

In Section 2, we provide some definitions and notation that will be used throughout the paper. In Section 3, we describe a generating tree for standard Young tableaux of certain shapes and show that it obeys a simple two-parameter labeling scheme. In Section 4.1, we briefly review (without proof) one of the main results of [5], namely that alternating permutations of even length with no four-term increasing subsequence share the same generating tree as the tableaux of Section 3. In Section 4.2, we show that alternating permutations of even length avoiding the pattern 2143 also share this generating tree. In Section 5, we show that similar methods can be used to enumerate alternating permutations of *odd* length avoiding the pattern 2143, and that the resulting bijection is with shifted Young tableaux of certain shapes. In Section 6 we give several conjectures and open problems relating to pattern avoidance in alternating permutations. In Section 7 we provide data that forms the basis for several of these conjectures.

2 Preliminaries

In this section, we provide important definitions and notation that will be used throughout the rest of the paper.

2.1 Permutations, partitions, tableaux

A *permutation* w of length n is a word containing each of the elements of $[n] := \{1, 2, \dots, n\}$ exactly once. The set of permutations of length n is denoted S_n . Given a word $w = w_1 \cdots w_n$ and a permutation $p = p_1 \cdots p_k \in S_k$, we say that w *contains the pattern* p if there exists a set of indices $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that the subsequence $w_{i_1} w_{i_2} \cdots w_{i_k}$ of w is order-isomorphic to p , i.e., such that $w_{i_\ell} < w_{i_m}$ if and only if $p_\ell < p_m$. Otherwise, w *avoids* p . Given a pattern p and a set S of permutations, we denote by $S(p)$ the set of elements of S that avoid p . For example, $S_n(123)$ is the set of permutations of length n avoiding the pattern 123, i.e., the set of permutations with no three-term increasing subsequence. We denote the size of a set S by $|S|$, so $|S_3(123)| = 5$.

A permutation $w = w_1 w_2 \cdots w_n$ is *alternating* if $w_1 < w_2 > w_3 < \dots$ (Note that in the terminology of [9], these “up-down” permutations are *reverse alternating* while alternating permutations are “down-up” permutations. Luckily, this convention doesn’t matter: any pattern result on either set can be translated into a result on the other via *complementation*, i.e., by considering w^c such that $w_i^c = n + 1 - w_i$. Then results for the pattern 123 and up-down permutations are replaced by results for 321 and down-up permutations, and so on.) We denote by A_n the set of alternating permutations of length n .

A *partition* is a weakly decreasing, finite sequence of nonnegative integers. We consider two partitions that differ only in the number of trailing zeroes to be the same. We write partitions in sequence notation, as $\langle \lambda_1, \lambda_2, \dots, \lambda_n \rangle$.

Given a partition $\lambda = \langle \lambda_1, \lambda_2, \dots, \lambda_n \rangle$, the *Young diagram* of shape λ is the left-justified array of $\lambda_1 + \dots + \lambda_n$ boxes with λ_1 in the first row, λ_2 in the second row, and so on. We identify each partition with its Young diagram and speak of them interchangeably. If λ is a Young diagram with m boxes, a *standard Young tableau* of shape λ is a filling of the boxes of λ with $[m]$ so that each element appears in exactly one box and entries increase along rows and columns. We identify boxes in Young diagrams and tableaux using matrix coordinates, so the box in the first row and second column is numbered $(1, 2)$. We denote by $\text{SYT}(\lambda)$ the set of standard Young tableaux of shape λ .

2.2 Generating trees

Given a sequence $\{\Sigma_n\}_{n \geq 1}$ of nonempty sets with $|\Sigma_1| = 1$, a *generating tree* for this sequence is a rooted, labeled tree such that the vertices at level n are the elements of Σ_n and the label of each vertex determines the multiset of labels of its children. In other words, a generating tree is one particular type of recursive structure in which heredity is determined by some local data. We are particularly interested in generating trees for which the labels are (much) simpler than the objects they are labeling. In this case, we may easily describe a generating tree by giving the label L_1 of the *root vertex* (the element of Σ_1) and the *succession rule* $L \mapsto S$ that gives the set S of labels of the children in terms of the label L of the parent.

Beginning with work of Chung, Graham, Hoggatt, and Kleiman [2], generating trees have been put to good use in the study of pattern-avoiding permutations, notably in the work of West (see, e.g., [11, 12]). The usual

approach has been to consider subtrees of the tree of all permutations given by the rule that $v \in S_{n+1}$ is a child of $u \in S_n$ exactly when erasing the entry $n + 1$ from v leaves the word u . Because we are interested here in alternating permutations and the permutation that results from erasing the largest entry of an alternating permutation typically is not alternating, this tree is unsatisfactory. However, the inverse tree, in which we arrive at the children of u by inserting arbitrary values in the last position (rather than inserting the largest value in an arbitrary position), is well-suited to the case of alternating permutations. This motivates the following definitions.

Given a permutation $u \in S_n$ and an element $i \in [n + 1]$, there is a unique permutation $v = v_1 v_2 \cdots v_n v_{n+1} \in S_{n+1}$ such that $v_{n+1} = i$ and the word $v_1 v_2 \cdots v_n$ is order-isomorphic to u . We denote this permutation by $u \leftarrow i$ and refer to it as the *extension* of u by i . In other words, the operation of extending u by i replaces each entry $c \geq i$ in u by $c + 1$ and then attaches i to the end of the result. For example, $3142 \leftarrow 3 = 41523$.

Given a pattern p and a permutation $w \in S_n(p)$, we say that $c \in [n + 1]$ is *active* or *an active value* for w (with respect to p) if $w \rightarrow c$ avoids p .¹

There is a natural generating tree structure on alternating permutations of even length analogous to the tree on all permutations mentioned above: given an alternating permutation u of length $2n$, its children are precisely the alternating permutations v of length $2n + 2$ such that the prefix of v of length $2n$ is order-isomorphic to u . (Of course, there is also a similar tree for alternating permutations of odd length.) Since pattern containment is transitive, the set $\bigcup_{n \geq 1} A_{2n}(p)$ of alternating permutations of even length avoiding the pattern p is the set of vertices of a subtree. It is these trees (for the patterns 1234 and 2143) that we will consider in Section 4.

3 A generating tree for tableaux

For $n \geq 1$, the collection of standard Young tableaux of shape $\langle n, n, n \rangle$ has a simple associated generating tree: given a tableau $S \in \text{SYT}(n, n, n)$, its children are precisely the tableaux $T \in \text{SYT}(n + 1, n + 1, n + 1)$ such that removing the last column of T leaves a tableau that is order-isomorphic to S . Notice that the shape of the tree below S is determined entirely by the entries

¹ This terminology is borrowed from the “usual” case, in which a position is said to be active (or is called an “active site”) if one can insert $n + 1$ into that position while preserving pattern avoidance.

of the last column of S , and in particular by the entries $S(1, n)$ and $S(2, n)$ (since $S(3, n) = 3n$ for all $S \in \text{SYT}(n, n, n)$). Thus, we wish to choose a labeling for our tree that captures exactly this information. Our choice (one of several reasonable options) is to assign to each $S \in \text{SYT}(n, n, n)$ the label $(a, b) = (3n + 1 - S(2, n), 3n + 1 - S(1, n))$. This provides a two-label generating tree for

$$\bigcup_{n \geq 1} \text{SYT}(n, n, n),$$

whose root (the unique standard Young tableau of shape $\langle 1, 1, 1 \rangle$) has label $(4 - 2, 4 - 1) = (2, 3)$. In Proposition 4.7 of [5], the author established (in a more general context, and using an object called “good tableaux” rather than the equivalent generating tree we’ve defined here) the following result.

Proposition 3.1. *The generating tree that we have just described for standard Young tableaux with all columns of length three obeys the rule*

$$(a, b) \mapsto \{(x, y) \mid 2 \leq x \leq a + 1 \text{ and } x + 1 \leq y \leq b + 2\}.$$

For example, the root tableau $\begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline \end{array}$ has label $(2, 3)$ and has five children,

$$\begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 5 \\ \hline 3 & 6 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 5 \\ \hline 4 & 6 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 5 \\ \hline 4 & 6 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline 5 & 6 \\ \hline \end{array} \text{ and } \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & 6 \\ \hline \end{array},$$

which have labels $(2, 3)$, $(2, 4)$, $(2, 5)$, $(3, 4)$ and $(3, 5)$, respectively. We restate the proof of this result in our current language.

Proof. Choose a standard Young tableau $S \in \text{SYT}(n, n, n)$ with label (a, b) and a child $T \in \text{SYT}(n + 1, n + 1, n + 1)$ of S with label (x, y) . We wish to show that $2 \leq x \leq a + 1$ and $x + 1 \leq y \leq b + 2$. Expressing these conditions in terms of the entries of S and T , we must show

$$S(2, n) + 2 \leq T(2, n + 1) \leq 3n + 2 \tag{1}$$

and

$$S(1, n) + 1 \leq T(1, n + 1) \leq T(2, n + 1) - 1. \tag{2}$$

We first demonstrate that Equation 2 is valid. It follows from the relationship between S and T that $S(1, n) \leq T(1, n)$, and since T is a standard

Young tableau we have $S(1, n)+1 \leq T(1, n)+1 \leq T(1, n+1) \leq T(2, n+1)-1$, as needed.

Now we demonstrate that Equation 1 is valid. Since T is a standard Young tableau, we have $T(2, n+1) < T(3, n+1) = 3n+3$. Thus $T(2, n+1) \leq 3n+2$, which is the right half of the desired inequality. For the left half, we compute that

$$\begin{aligned} T(2, n+1) &= \left| \{(i, j) \mid T(i, j) < T(2, n+1)\} \right| \\ &\geq \left| \{(i, j) \mid T(i, j) < T(2, n) \text{ and } j \leq n\} \cup \{(2, n), (1, n+1)\} \right| \\ &= \left| \{(i, j) \mid S(i, j) < S(2, n)\} \cup \{(2, n), (1, n+1)\} \right| \\ &= S(2, n) + 2, \end{aligned}$$

as desired.

We've now established that Equations 1 and 2 hold for all children T of S , and so that the labels of the children of S satisfy the claimed inequalities. Now we must show that S has exactly one child with each of these labels.

Given a parent tableau S and prescribed values $T(1, n+1)$ and $T(2, n+1)$ satisfying Equations 2 and 1, we construct the tableau T in the only way possible: define $T(3, n+1) = 3n+3$ and for $1 \leq i \leq 3$ and $1 \leq j \leq n$, set

$$T(i, j) = \begin{cases} S(i, j), & S(i, j) < T(1, n+1), \\ S(i, j) + 1, & T(1, n+1) \leq S(i, j) < T(2, n+1), \\ S(i, j) + 2, & T(2, n+1) \leq S(i, j). \end{cases}$$

By construction, the object T that results is of shape $\langle n+1, n+1, n+1 \rangle$ and contains each of the elements of $[3n+3]$ exactly once. It is not difficult to check that Equations 1 and 2 imply that T is increasing along rows and columns and consequently that $T \in \text{SYT}(n+1, n+1, n+1)$, and also that T is a child of S . Thus, S has at least one child with each label satisfying the given conditions. Finally, it's clear that this T is unique. \square

4 Generating trees for $A_{2n}(1234)$ and $A_{2n}(2143)$

Recall from Section 2.2 that for any pattern p , there is a natural generating tree for the set $\bigcup_n A_{2n}(p)$ of p -avoiding alternating permutations of even

length: given a p -avoiding alternating permutation u of length $2n$, its children are precisely the p -avoiding alternating permutations v of length $2n + 2$ such that the prefix of v of length $2n$ is order-isomorphic to u . In this section, we study this tree for the patterns $p = 1234$ and $p = 2143$.

4.1 1234

One of the main results of [5] was that even-length alternating permutations with no four-term increasing subsequence share the generating tree for Young tableaux described in Section 3. We now briefly describe this labeling.

Given a 1234-avoiding alternating permutation w , assign to it a label (a, b) where a is the smallest entry in w that is the largest entry in a two-term increasing subsequence of w , i.e.,

$$a = \min\{w_j \mid \exists i < j \text{ s.t. } w_i < w_j\},$$

and b is the number of active values for w with respect to 1234, i.e., b is the number of choices of $c \in [2n + 1]$ such that $w \leftarrow c$ is 1234-avoiding. (Equivalently, we could define b to be the smallest entry in w that the largest term in a three-term increasing subsequence, or $b = 2n + 1$ if w contains no three-term increasing subsequence.) This labeling has previously been used (e.g., by West [11] and Bousquet-Mélou [1]) in the study of $S_n(1234)$.

For example, we have for the permutation $w = 27583614 \in A_8(1234)$ that $\{w_j \mid \exists i < j \text{ s.t. } w_i < w_j\} = \{3, 4, 5, 6, 7, 8\}$ and that the set of active values for w is $\{1, 2, 3, 4\}$. Thus $a = 3$ and $b = 4$.

As a second example, we have for the permutation $w = 68372514 \in A_8(1234)$ that $\{w_j \mid \exists i < j \text{ s.t. } w_i < w_j\} = \{4, 5, 7, 8\}$ and that w contains no three-term increasing subsequence so all of $[9]$ is active for w . Thus $a = 4$ and $b = 9$.

The root of this tree is the permutation $12 \in A_2(1234)$, with label $(2, 3)$. Propositions 4.3, 4.4 and 4.6 of [5] collectively establish that if $w \in A_{2n}(1234)$ has label (a, b) then the collection of labels of children of w is exactly the set of (x, y) such that $2 \leq x \leq a + 1$ and $x + 1 \leq y \leq b + 2$, each pair occurring with multiplicity one.

We now turn our attention to the pattern 2143.

4.2 2143

In this section, we show that alternating permutations of even length avoiding 2143 have a generating tree isomorphic to those mentioned in preceding sections.

Given a permutation $w \in A_{2n}(2143)$, assign to it a label (a, b) where $a = w_{2n-1}$ is the next-to-last entry of w and b is the number of active values for w in $[2n + 1]$.

For example, the permutation $w = 68143527 \in A_8(2143)$ has active values $\{1, 2, 3, 8, 9\}$ (the permutations 792546381, 791546382, 791546283, 691435278 and 681435279 avoid 2143 while the permutations 691435287, 791435286, 791436285 and 791536284 contain it). Thus $a = 2$ and $b = 5$.

As a second example, the permutation $w = 35462718 \in A_8(2143)$ has active values $\{1, 2, 9\}$. Thus $a = 1$ and $b = 3$.

We have that the root $12 \in A_2(2143)$ has label $(1, 3)$. We will show that under this labeling the generating tree for $\bigcup_n A_{2n}(2143)$ obeys the rule

$$(a, b) \mapsto \{(x, y) \mid 1 \leq x \leq a + 1 \text{ and } x + 2 \leq y \leq b + 2\},$$

and we will use this result to establish an isomorphism between this generating tree and those discussed in the preceding sections. We break the proof of this result into several smaller pieces: Propositions 4.3 and 4.5 form the meat of the argument establishing the succession rule, while Propositions 4.2 and 4.4 are helpful technical lemmas. We begin with a simple observation that will be of use in the subsequent proofs.

Observation 4.1. *Given a permutation $w \in S_n(2143)$, we have that $c \in [n + 1]$ is not active for w if and only if there exists $i < j < k$ such that $w_j < w_i < c \leq w_k$.*

Proposition 4.2. *If $u \in A_{2n}(2143)$ and $u_{2n-1} = a$ then $\{1, 2, \dots, a + 1\}$ are active values for u .*

Proof. Fix $u \in A_{2n}(2143)$ with $u_{2n-1} = a$, choose $c \leq a + 1$ and let $v = u \leftarrow c \in S_{2n+1}$. We wish to show that v avoids 2143, so suppose otherwise. Then there exist $i < j < k < 2n + 1$ such that $v_i v_j v_k c$ is an instance of 2143 in v . We use this (suppositional) instance to construct an instance of 2143 in u ; this contradiction establishes that c is active for u . In particular, we show that $u_i u_j u_{2n-2} u_{2n-1}$ is an instance of 2143 in u by showing that $v_i v_j v_{2n-2} v_{2n-1}$ is an instance of 2143 in v . In order to do this, it suffices to

show that $j < 2n - 2$ (so that $v_i v_j v_{2n-2} v_{2n-1}$ is a subsequence of v) and that $v_i < v_{2n-1}$ (so that this subsequence is order-isomorphic to 2143).

Since $v_i v_j v_k c$ is an instance of 2143 and $c \leq a + 1$, we have that $v_j < v_i < c \leq a + 1$ and thus $v_i \leq a = u_{2n-1} \leq v_{2n-1}$. There are at least three entries to the right of v_i in v but only two to the right of v_{2n-1} , so $v_i \neq v_{2n-1}$ and actually $v_i < v_{2n-1}$, one of the two conditions we need. It follows that $v_j < v_i < v_{2n-1} < v_{2n-2}$ and similarly $v_j < v_{2n}$, so v_j is smaller than all of $v_{2n-2}, v_{2n-1}, v_{2n}$ and v_{2n+1} . These entries form a suffix of v , so v_j must occur at an earlier position in v . That is, we have $j < 2n - 2$, the second necessary condition. Thus $v_i v_j v_{2n-2} v_{2n-1}$ is an instance of 2143 in v and so $u_i u_j u_{2n-2} u_{2n-1}$ is an instance of 2143 in u . Since u avoids 2143, this is a contradiction, so actually v avoids 2143 and c is active for u , as desired. \square

Proposition 4.3. *If $u \in A_{2n}(2143)$ has label (a, b) and w is a child of u with label (x, y) then $1 \leq x \leq a + 1$ and $x + 2 \leq y \leq b + 2$.*

Proof. Suppose that permutations $u \in A_{2n}$ and $w \in A_{2n+2}$ have the property that the first $2n$ entries of w are order-isomorphic to u , and set $a = u_{2n-1}$. If $w_{2n+1} > a + 1$ then also $w_{2n} > a + 1$ and $w_{2n+2} > a + 1$, while $w_{2n-1} = a$. Thus $w^{-1}(a + 1) \notin \{2n - 1, 2n, 2n + 1, 2n + 2\}$ and so defining $i = w^{-1}(a + 1)$ we have $i < 2n - 1$. Then $w_i w_{2n-1} w_{2n} w_{2n+1}$ is an instance of 2143 in w . Taking the contrapositive, if w avoids 2143 then $w_{2n+1} \leq a + 1$. Now fix $u \in A_{2n}(2143)$ with label (a, b) and a child $w \in A_{2n+2}(2143)$ of u with label (x, y) ; the preceding argument shows that $x \leq a + 1$. Clearly $x \geq 1$, so we have proved the first half of our assertion. We now proceed to bound y , the number of active values of w .

Define a one-to-one function $f: [2n] \rightarrow [2n + 2]$ by

$$f(z) = \begin{cases} z, & z < w_{2n+1} \\ z + 1, & w_{2n+1} \leq z < w_{2n+2} \\ z + 2, & w_{2n+2} \leq z, \end{cases}$$

so $f(u_\ell) = w_\ell$ for all $\ell \in [2n]$. We show that f is a map from nonactive values of u to nonactive values of w . To this end, choose any $c \in [2n]$ that is not active for u , and choose $i < j < k$ such that $u_j < u_i < c \leq u_k$. We have $(w_i, w_j, w_k, f(c)) = (f(u_i), f(u_j), f(u_k), f(c))$. One can easily see that f preserves order, so $w_j < w_i < f(c) \leq w_k$ and thus $f(c)$ is not active for w . Thus for each of the $2n - b$ choices of a nonactive value c for u

we have a corresponding nonactive value $f(c)$ for w and so w has at most $(2n+2) - (2n-b) = b+2$ active values, i.e., $y \leq b+2$.

Finally, we have by Proposition 4.2 that $\{1, 2, \dots, x+1\}$ are active values for w . We also have that $2n+3$ is active for w and that $2n+3 \notin \{1, 2, \dots, x+1\}$, so there are at least $x+2$ active values for w . Thus $x+2 \leq y$, which completes the proof of our claim. \square

Proposition 4.4. *Suppose $u \in A_{2n}(2143)$ has label (a, b) and active values $s_1 < s_2 < \dots < s_b$. If $x \leq a+1$ and $v = u \leftarrow x$ then $v \in A_{2n+1}(2143)$ and v has active values $1, 2, \dots, x, s_x+1, s_{x+1}+1, \dots, s_b+1$.*

For example, $u = 68143527 \in A_8(2143)$ has label $(2, 5)$ and active values $1, 2, 3, 8$ and 9 . For $x = 3 \leq 2+1$ we have $v = 791546283$ with active values $1, 2, 3, 4 (= 3+1), 9 (= 8+1)$ and $10 (= 9+1)$.

Proof. Choose $u \in A_{2n}(1234)$ with label (a, b) and choose $x \leq a+1$. Let $s_1 < \dots < s_b$ be the active values for u and let $v = u \leftarrow x$. Since $v_{2n+1} = x \leq a+1 \leq v_{2n-1}+1$ and $v_{2n-1} < v_{2n}$, we have $v_{2n+1} \leq v_{2n}$ and thus actually $v_{2n+1} < v_{2n}$. Thus v is alternating. Also, by Proposition 4.2, x is an active value for u and so v avoids 2143. This proves the first part of our claim, and we now must show that the set of active values for v is $\{1, 2, \dots, x, s_x+1, s_{x+1}+1, \dots, s_b+1\}$. We first show that each of these values is in fact an active value for v .

Case 1. Fix $m \geq x$ so that $s_m \geq x$ is an active value for u . We wish to show that s_m+1 is an active value for v . Let $w = w_1 \cdots w_{2n+2}$ be the result of extending v by s_m+1 , and suppose for sake of contradiction that w contains an instance $w_i w_j w_k w_\ell$ of 2143. Since v avoids 2143, we must have $\ell = 2n+2$, so $w_\ell = w_{2n+2} = s_m+1$. As $w_{2n+2} = s_m+1 > x = v_{2n+1}$, extending v by s_m+1 does not change the value of the $(2n+1)$ th entry and thus $w_{2n+1} = v_{2n+1} = x$, whence $w_{2n+2} > w_{2n+1}$. Since $w_k > w_{2n+2}$, we have $k \neq 2n+1$ and so w_{2n+1} is not part of our instance of 2143. Let v' be the permutation order-isomorphic to $w_1 w_2 \cdots w_{2n} w_{2n+2}$. Thus $v'_i v'_j v'_k v'_{2n+1}$ is an instance of 2143 in v' . However, we also have that $v' = u \leftarrow s_m$ and that s_m is active for u . This is a contradiction, so w cannot contain an instance of 2143, and we conclude that s_m+1 is an active value for v by definition.

Case 2. Fix $c \leq x$ so that $c = s_c$ is an active value for u . We wish to show that c is an active value for v . Let $w = w_1 \cdots w_{2n+1} w_{2n+2}$ be the

result of extending v by c , and suppose for sake of contradiction that w contains an instance $w_i w_j w_k w_\ell$ of 2143. If $\{k, \ell\} \neq \{2n+1, 2n+2\}$ then we can conclude by an argument nearly identical to the previous case, so assume that $k = 2n+1$ and $\ell = 2n+2$. Since $w_i < w_k = w_{2n+1}$ and $w_j < w_k = w_{2n+1}$ but $w_{2n} > w_{2n+1}$, $2n \notin \{i, j\}$ and so $w_i w_j w_{2n} w_{2n+2}$ is another instance of 2143 in w . But now we have an instance not including w_{2n+1} and so we may proceed as in the previous case. We conclude that c is an active value for v .

Finally, we show that these values are the only active values for v . In particular, we must show that for every $c > x$ that is not active for u , $c+1$ is not active for v . Fix such a c . Since c is not active for u , there exist $i < j < k$ such that $u_j < u_i < c \leq u_k$. We have $v_\ell \leq u_\ell + 1$ for all $\ell \in [2n]$, so $v_j < v_i < c+1$. Since $u_k > c-1 \geq x$, we have $v_k = u_k + 1$. Thus $v_j < v_j < c+1 \leq u_k + 1 = v_k$ and so $c+1$ is not an active value for v . We conclude that the active values for v are exactly $1, 2, \dots, x, s_x + 1, s_{x+1} + 1, \dots, s_b + 1$, as claimed. \square

Proposition 4.5. *If $u \in A_{2n}(2143)$ has label (a, b) and $1 \leq x \leq a+1$, $x+2 \leq y \leq b+2$, then there is a unique child $w \in A_{2n+2}(2143)$ of u with label (x, y) .*

Proof. Choose a permutation $u \in A_{2n}(2143)$ with label (a, b) and choose (x, y) such that $1 \leq x \leq a+1$ and $x+2 \leq y \leq b+2$. We will construct a child of u with label (x, y) .

Let $s_1 < \dots < s_b$ be the active values for u . Define $v = v_1 v_2 \dots v_{2n} v_{2n+1}$ and $w = w_1 w_2 \dots w_{2n} w_{2n+1} w_{2n+2}$ by $v = u \leftarrow x$ and $w = v \leftarrow (s_{b+2+x-y} + 1)$. We claim that w is the desired permutation. We must show that w belongs to $A_{2n+2}(2143)$ and that its label really is (x, y) .

It follows from Proposition 4.4 that $v \in A_{2n+1}(2143)$. Because $y \leq b+2$, we have $b+2+x-y \geq x$ and so $s_{b+2+x-y} + 1 > s_x = x = v_{2n+1}$. Thus w is alternating. We also have from Proposition 4.4 that $s_{b+2+x-y} + 1$ is an active value for v , so $w \in A_{2n+2}(2143)$. We have left to show that w has label (x, y) .

Since $w_{2n+2} > v_{2n+1}$, extending v by w_{2n+2} leaves the value of the entry in the $(2n+1)$ th position unchanged and so $w_{2n+1} = v_{2n+1} = x$. It remains to show that w has exactly y active values. We claim that the active values for w are precisely $1 < 2 < \dots < x+1 < s_{b+2+x-y} + 2 < s_{b+3+x-y} + 2 < \dots < s_b + 2$. By Proposition 4.2, we already know that $1, 2, \dots, x+1$ are active for w .

For any fixed i such that $b + 2 + x - y \leq i \leq b$, let $z = w \leftarrow (s_i + 2)$. We wish to show that z avoids 2143. We proceed by the same argument as in Case 1 of the proof of Proposition 4.4: since $s_i + 2 > w_{2n+2}$, we have $z_{2n+2} = w_{2n+2} < z_{2n+3}$. It follows that z_{2n+2} and z_{2n+3} cannot be part of the same 2143 pattern in z . Thus z contains an instance of 2143 if and only if $z' = z_1 \cdots z_{2n+1} z_{2n+3}$ does. However, z' is order-isomorphic to $v \leftarrow (s_i + 1)$. Since $s_i + 1$ is an active value for v , z' avoids 2143 and so z avoids 2143. Thus $s_i + 2$ is an active value for w . There are $b - (b + 2 + x - y) + 1 = y - x - 1$ such active values. We must show that there are no other active values for w larger than $x + 1$.

For any fixed $c > s_{b+2+x-y} + 2$ that is not of the form $s_i + 2$, we wish to show that $w \leftarrow c$ contains 2143. We proceed by the same arguments that follow Case 2 of the proof of Proposition 4.4: if $v_i v_j v_k$ are entries of v that can be used to form a 2143 pattern when v is extended by $c - 1$ then $w_i w_j w_k$ can be used to form a 2143 pattern when w is extended by c .

Finally, for any fixed c such that $x + 1 < c < s_{b+2+x-y} + 2$, let $z = w \leftarrow c$. We wish to show that z contains 2143. We have that $z_{2n+1} = x$, $z_{2n+2} = s_{b+2+x-y} + 2 > c$, and $z_{2n+3} = c > x + 1$, and we know that there exists $i < 2n + 1$ such that $z_i = x + 1$. Then $z_i z_{2n+1} z_{2n+2} z_{2n+3} = (x + 1)x(s_{b+2+x-y} + 2)c$ is an instance of 2143 in z , so c is not active for w .

The preceding four paragraphs account for all elements $[2n + 3]$. We've shown that exactly $(x + 1) + (y - x - 1) = y$ of these values are active for w . Putting everything together, we have that $w \in A_{2n+2}(2143)$ is a child of u with y active values and $w_{2n+1} = x$, i.e., w has label (x, y) . In fact, it follows from our proof that every child of u has a distinct label: we've exhausted the possible pairs of values for w_{2n+1} , w_{2n+2} such that $w \in A_{2n+2}(2143)$ is a child of u . \square

Theorem 4.6. *For all $n \geq 1$ we have*

$$|A_{2n}(2143)| = |\text{SYT}(n, n, n)| = |A_{2n}(1234)| = \frac{2 \cdot (3n)!}{n!(n+1)!(n+2)!}.$$

Proof. Proposition 3.1 shows that the generating tree for $\bigcup_{n \geq 1} \text{SYT}(n, n, n)$ has root $(2, 3)$ and rule

$$(a, b) \mapsto \{(x, y) \mid 2 \leq x \leq a + 1 \text{ and } x + 1 \leq y \leq b + 2\}$$

while Propositions 4.3 and 4.5 together show that the generating tree for $\bigcup_{n \geq 1} A_{2n}(2143)$ has root $(1, 3)$ and rule

$$(a, b) \mapsto \{(x, y) \mid 1 \leq x \leq a + 1 \text{ and } x + 2 \leq y \leq b + 2\}.$$

These two trees are isomorphic: replacing each label (a, b) in the first tree with $(a - 1, b)$ results in the second tree. Thus, there is a recursive bijection between $A_{2n}(2143)$ and $\text{SYT}(n, n, n)$, and we have the first claimed equality. The work in [5] summarized in Section 4.1 gives the second equality. Finally, applying the hook-length formula (see, e.g., [10, Chapter 7.21]) gives the third equality. \square

5 Generating tree for $A_{2n+1}(2143)$

Turning our attention to alternating permutations of odd length, we find that the generating trees of $A_{2n+1}(1234)$ and $A_{2n+1}(2143)$ are not isomorphic. Indeed, the two sequences enumerate differently: all sixteen alternating permutations of length five avoid 1234, but only twelve of them avoid 2143. Although this initially seems disappointing, it turns out that we can still use the methods of the preceding section to enumerate 2143-avoiding alternating permutations of odd length.

As in [5], it is convenient to consider the set A'_{2n+1} of *down-up* alternating permutations of odd length rather than up-down alternating permutations; note that results in either case may be translated into results in the other via reverse-complementation. Arguments very similar to those of Section 4.2 show that if we associate to the permutation $w \in A'_{2n+1}(2143)$ the label (a, b) , where $a = w_{2n}$ and b is the number of active values for w then the generating tree for $\bigcup_{n \geq 0} A'_{2n+1}(2143)$ is as follows:

Root: $(0, 2)$

Rule: $(a, b) \mapsto \{(x, y) \mid 1 \leq x \leq a + 1 \text{ and } x + 2 \leq y \leq b + 2\}$

Here the root permutation $1 \in A'_1(2143)$ has label $(0, 2)$. Its two children $213, 312 \in A'_3(2143)$ have labels $(1, 3)$ and $(1, 4)$, respectively, and so on.

We seek to enumerate $A'_{2n+1}(2143)$ by applying our approach for even-length permutations, i.e., by finding a family of objects with isomorphic generating tree that we already know how to enumerate. In the case at hand, these objects turn out to be *shifted standard Young tableaux* (henceforward SHSYT) of shape $\langle n + 2, n + 1, n \rangle$. (See [4] or [3, Chapter 10] for definitions, etc.)

Given a SHSYT T of shape $\langle n + 2, n + 1, n \rangle$, assign to it the label (a, b) , where $a = 3n + 4 - T(n + 2, 2)$ and $b = 3n + 4 - T(n + 2, 1)$. (Note that for $n \geq 1$, we have $T(n + 2, 3) = 3n + 3$ for every SHSYT of this shape, so this

label captures all the information we need to reconstruct the last column of T .) Then the root of the tree is the unique SHSYT $\begin{array}{|c|c|} \hline 1 & 2 \\ \hline & 3 \\ \hline \end{array}$ of shape $\langle 2, 1 \rangle$, which has label $(1, 2)$. Its children are the two SHSYT

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline & 3 & 5 \\ \hline & & 6 \\ \hline \end{array} \quad \text{and} \quad \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline & 4 & 5 \\ \hline & & 6 \\ \hline \end{array}$$

of shape $\langle 3, 2, 1 \rangle$, which have labels $(2, 3)$ and $(2, 4)$, respectively. In subsequent layers of the tree, the succession rule is identical to the rule for rectangular tableaux; indeed, the subtree below an SHSYT of shape $\langle n+2, n+1, n \rangle$ depends only on its last column (or equivalently, on its associated label), and so might as well be the subtree of an SYT of shape $\langle (n+1)^3 \rangle$ with the same label.

It follows immediately that $|A'_{2n+1}(2143)|$ (and so also $|A_{2n+1}(2143)|$) is the number of SHSYT of shape $\langle n+2, n+1, n \rangle$. As in the case of standard Young tableaux, there is a simple hook-length formula for SHSYT (see e.g. [4] or [3, pp. 187-190]). In our case, it gives the following result.

Proposition 5.1. *For $n \geq 0$ we have*

$$|A_{2n+1}(2143)| = \frac{2(3n+3)!}{n!(n+1)!(n+2)!(2n+1)(2n+2)(2n+3)}.$$

6 Open problems

In this section we pose a number of open enumerative problems related to pattern avoidance in alternating permutations.

6.1 Other equivalences for patterns of length 4

If permutations p and q satisfy $|A_{2n}(p)| = |A_{2n}(q)|$ for all $n \geq 1$, we say that p and q are *equivalent* for even-length alternating permutations. Note that if $p = p_1 \cdots p_k$ and $q = (k+1-p_k)(k+1-p_{k-1}) \cdots (k+1-p_1)$ (i.e., p and q are reverse-complements) then p and q are equivalent for even-length alternating permutations: for every n , the operation of reverse-complementation is a bijection between $A_{2n}(p)$ and $A_{2n}(q)$. Pairs of patterns that are equivalent for this reason are said to be *trivially equivalent*. Similarly, if $|A_{2n+1}(p)| =$

$|A_{2n+1}(q)|$ for all $n \geq 0$, we say that p and q are equivalent for odd-length alternating permutations, and if p is the reverse of q then we say they are trivially equivalent.

Numerical data (see Section 7) suggest the following conjecture.

Conjecture 6.1. *We have $|A_{2n}(p)| = |A_{2n}(1234)| (= |A_{2n}(2143)|)$ for all $n \geq 1$ and every $p \in \{1243, 2134, 1432, 3214, 2341, 4123, 3421, 4312\}$.*

Observe that these eight patterns come in four pairs of trivially equivalent patterns. The results of West [11] and computer investigations of short permutations suggest that some of these equivalences may be susceptible to a generating-tree attack. In particular, the generating trees for alternating permutations avoiding 1243 or 2134 may be isomorphic to the generating tree discussed in Sections 3 and 4.

For alternating permutations of even length, the only other possible equivalences not ruled out by numerical data are captured by the following conjecture.

Conjecture 6.2. *We have $|A_{2n}(3142)| = |A_{2n}(3241)| = |A_{2n}(4132)|$ and $|A_{2n}(2413)| = |A_{2n}(1423)| = |A_{2n}(2314)|$ for all $n \geq 1$.*

In both cases, the second of the two equalities is a trivial equivalence.

For odd-length alternating permutations, computational data suggest the following conjectures.

Conjecture 6.3. *We have $|A_{2n+1}(p)| = |A_{2n+1}(1234)|$ for all $n \geq 0$ and every $p \in \{2134, 4312, 3214, 4123\}$.*

We also have the trivial equivalence $|A_{2n+1}(1234)| = |A_{2n+1}(4321)|$. The equivalence between 4321 and 4312 may be amenable to generating tree methods.

Conjecture 6.4. *We have $|A_{2n+1}(p)| = |A_{2n+1}(2143)|$ for all $n \geq 0$ and every $p \in \{1243, 3421, 1432, 2341\}$.*

We also have the trivial equivalence $|A_{2n+1}(2143)| = |A_{2n+1}(3412)|$. The equivalence between 3412 and 3421 may be amenable to generating tree methods.

The only other possible equivalence for odd-length alternating permutations not ruled out by data is captured by the following conjecture.

Conjecture 6.5. *The permutations 2314, 4132, 2413, 3142, 1423 and 3241 are equivalent for odd-length alternating permutations.*

Other than the cases covered by Conjectures 6.1, 6.3 and 6.4, none of the sequences $\{|A_{2n}(p)|\}_n$ or $\{|A_{2n+1}(p)|\}_n$ for $p \in S_4$ are recognizable to the present author (and in particular they do not appear in the OEIS [7]). Numerical data (see Section 7) rule out simple product formulas similar to those of Theorem 4.6 and Proposition 5.1. However, in some cases it may be possible to give a generating tree and so perhaps to adapt the method of [1] to find generating functions or even closed formulas.

6.2 Other problems

The conjectures of the preceding section concern two equivalence relations on patterns of length four: equivalence for even-length alternating permutations and equivalence for odd-length alternating permutations. It happens that these relations are both refinements of the usual (Wilf-)equivalence relation for all permutations (see, for example, [8]). (Note that this is true on account of numerical data, regardless of the truth of any of the preceding conjectures.) This suggests the following conjecture.

Conjecture 6.6. *If permutations p and q are equivalent for alternating permutations of either parity then p and q are Wilf-equivalent for all permutations.*

More broadly, we can ask the following question.

Question 6.7. *Are there any large families of patterns that can be shown to be equivalent for alternating permutations (of either parity)?*

The work in [5] suggests two possible generalizations of alternating permutations for purposes of pattern avoidance. One of these (denoted $\mathcal{L}_{n,k}$ in [5]; it consists of the reading words of standard Young tableaux of the “thickened staircase” shape $\langle n+k-1, n+k-2, \dots, k+1, k \rangle / \langle n-1, n-2, \dots, 1 \rangle$) does not seem to give rise to nontrivial equivalences. However, an alternative (and perhaps more natural) generalization is the set $\text{Des}_{n,k}$ of permutations of length n with descent set $\{k, 2k, \dots\}$.

Question 6.8. *Is $\text{Des}_{n,k}$ a “good” context to study pattern avoidance? In particular, are there any pairs or families of patterns that can be shown to*

be Wilf-equivalent for these permutations? Is Conjecture 6.6 valid in this context?

In analogy with the case of alternating permutations, it is natural to consider separate cases depending on the congruence class of n modulo k . Finally, we note that it may also be fruitful to consider permutations whose descent set is contained in (rather than “is equal to”) $\{k, 2k, \dots\}$, a case on which our work here does not touch.

7 Appendix

The tables of data that follow form the basis for several of the conjectures in the preceding section. They were generated by brute-force computer enumeration.

Tables 1 and 2 give the number of alternating permutations avoiding patterns of length four, grouped by conjectural equivalence.

Patterns	1	3	5	7	9	11
(1234, 4321), (2134, 4312), (3214, 4123)	1	2	16	168	2112	30030
(2143, 3412), (1243, 3421), (1432, 2341)	1	2	12	110	1274	17136
(2314, 4132), (2413, 3142), (1423, 3241)	1	2	12	106	1138	13734
(1324, 4231)	1	2	12	110	1285	17653
(1342, 2431)	1	2	12	108	1202	15234
(3124, 4213)	1	2	16	168	2072	28298

Table 1: Values of $A_n(p)$ for odd n and $p \in S_4$. Parentheses indicate trivial equivalences.

Patterns	2	4	6	8	10	12
1234, (1243, 2134), (1432, 3214), 2143, (2341, 4123), (3421, 4312)	1	5	42	462	6006	87516
3142, (3241, 4132)	1	5	42	444	5337	69657
(1423, 2314), 2413	1	4	28	260	2844	34564
3412	1	4	29	290	3532	49100
1324	1	4	29	292	3620	51866
(1342, 3124)	1	5	42	453	5651	77498
(2431, 4213)	1	5	42	454	5680	78129
4231	1	5	42	462	6070	90686
4321	1	5	61	744	10329	157586

Table 2: Values of $A_n(p)$ for even n and $p \in S_4$. Parentheses indicate trivial equivalences.

Tables 3 and 4 give the number of alternating permutations avoiding certain patterns of length five. Only those patterns that might potentially have nontrivial equivalences are included, and they are grouped by these potential equivalences.

Patterns	1	3	5	7	9	11
(12534, 43521), (21534, 43512)	1	2	16	243	5291	144430
(12453, 35421), (21453, 35412)	1	2	16	243	5307	146013
(12354, 45321), (12543, 34521), (15432, 23451), (21354, 45312), (21543, 34512), (32154, 45123)	1	2	16	243	5330	148575
(12435, 53421), (21435, 53412)	1	2	16	243	5330	148764
(12345, 54321), (21345, 54312), (32145, 54123), (43215, 51234)	1	2	16	272	6531	194062

Table 3: Selected values of $A_n(p)$ for odd n and $p \in S_5$. Parentheses indicate trivial equivalences. All possible nontrivial equivalences are among the permutations in this table.

Patterns	2	4	6	8	10	12
(12534, 23145), (21534, 23154)	1	5	56	997	23653	679810
(34512, 45123), 45312	1	5	56	1004	24310	724379
(12435, 13245), (13254, 21435)	1	5	56	1004	24336	727807
(12453, 31245), (21453, 31254)	1	5	61	1194	30802	953088
12345, 21354, (12354, 21345), (12543, 32145), (15432, 43215), (21543, 32154), (23451, 51234), (34521, 54123), (45321, 54312)	1	5	61	1194	30945	970717

Table 4: Selected values of $A_n(p)$ for even n and $p \in S_5$. Parentheses indicate trivial equivalences. All possible nontrivial equivalences are among the permutations in this table.

Table 5 gives the number of permutations of length $3n$ with descent set $\{3, 6, \dots\}$ that avoid certain patterns of length four and five. Only those patterns that might potentially have nontrivial equivalences are included.

Patterns	3	6	9	12
2413, (1423, 2314)	1	9	153	3465
(1243, 2134), (2341, 4123)	1	9	153	3579
3142, (3241, 4132)	1	19	642	27453
2143, 4231, (1432, 3214), (3421, 4312)	1	19	642	29777
(12354, 21345), (23451, 51234)	1	19	887	66816
(15243, 32415), (35241, 52413)	1	19	1077	102051
(12543, 32154), (34521, 54123)	1	19	1134	114621
21354, 52341	1	19	1134	115515
(15432, 43215), (21543, 32154), (25431, 53214), (31542, 42153), (32541, 52143), (35421, 54213), (41532, 43152), (42531, 53142), (43251, 51432), (43521, 54132), (45321, 54312), (52431, 53241), (53421, 54231)	1	19	1513	211425

Table 5: Selected values of $\text{Des}_{n,3}(p)$ for n divisible by 3 and $p \in S_4$ or S_5 . Parentheses indicate trivial equivalences. All possible nontrivial equivalences are among the permutations in this table.

References

- [1] M. Bousquet-Mélou. Four classes of pattern-avoiding permutations under one roof: generating trees with two labels. *Electronic J. Combinatorics*, 9:R19, 2003.
- [2] F. R. K. Chung, R. L. Graham, V. E. Hoggatt, Jr., and M. Kleiman. The number of Baxter permutations. *J. Combinatorial Theory, Series A*, 24:382–394, 1978.
- [3] P. N. Hoffman and J. F. Humphreys. *Projective representations of the symmetric groups*. Oxford University Press, 1992.

- [4] C. Krattenthaler. Bijective proofs of the hook formulas for the number of standard Young tableaux, ordinary and shifted. *Electronic J. Combinatorics*, 2:R13, 1995.
- [5] J. B. Lewis. Pattern avoidance and RSK-like algorithms for alternating permutations and Young tableaux. [arXiv:0909.4966v2](https://arxiv.org/abs/0909.4966v2), 2010.
- [6] E. Ouchterlony. Pattern avoiding doubly alternating permutations. *Proc. FPSAC 2006*. [arXiv:0908.0255v1](https://arxiv.org/abs/0908.0255v1).
- [7] N. J. A. Sloane. The on-line encyclopedia of integer sequences. Available online at <http://www.research.att.com/njas/sequences/>, 2009.
- [8] Z. Stankova and J. West. A new class of Wilf-equivalent permutations. *J. Algebraic Combinatorics*, 15:271–290, 2002.
- [9] R. P. Stanley. *Enumerative Combinatorics, Volume 1*. Cambridge University Press, 1997.
- [10] R. P. Stanley. *Enumerative Combinatorics, Volume 2*. Cambridge University Press, 2001.
- [11] J. West. Generating trees and the Catalan and Schröder numbers. *Discrete Mathematics*, 146:247–262, 1995.
- [12] J. West. Generating trees and forbidden subsequences. *Discrete Mathematics*, 157:363–372, 1996.