

On Guessing Whether A Sequence Has A Certain Property

Samuel Alexander

alexander@math.ohio-state.edu

Abstract: A concept of “guessability” is defined for sets of sequences of naturals. Eventually, these sets are thoroughly characterized. To do this, a nonstandard logic is developed, a logic containing symbols for the ellipsis as well as for functions without fixed arity. New proofs are given for some seemingly-unrelated known results.

1 Motivation

Suppose we are playing a game. I am reading a fixed sequence, one entry at a time. You are trying to guess whether 0 is in the sequence. You can revise your guess with each new revealed entry, and you win if your guesses converge to the correct answer. You have an obvious strategy: always guess no, until 0 appears (if ever), then guess yes forever. The set of sequences containing 0 is guessable.

Suppose, instead, you are trying to guess whether my sequence contains *infinitely many* zeroes. We will see there is no strategy, not even if your computation power is unlimited. The set of sequences with infinitely many zeroes is unguessable.

A sequence $f : \mathbb{N} \rightarrow \mathbb{N}$ is *onto* if $\forall m \exists n f(n) = m$. This definition uses nested quantifiers: quantifiers appear in the scope of other quantifiers. Is it possible to give an alternate definition without nested quantifiers? The answer is “no”, but how to prove it? We’ll give a proof of a very strong negative answer, strong in the sense that nested quantifiers cannot be eliminated even in an extremely powerful language. Of course, the technique generalizes to a wide class of things, not just the onto sequences.

2 Basics

Let $\mathbb{N}^{\mathbb{N}}$ be the set of sequences $f : \mathbb{N} \rightarrow \mathbb{N}$, and let $\mathbb{N}^{<\mathbb{N}}$ be the set of finite sequences.

Definition 1. A function $G : \mathbb{N}^{<\mathbb{N}} \rightarrow \{0, 1\}$ *guesses* (and is a *guesser* for) a set $S \subseteq \mathbb{N}^{\mathbb{N}}$ if for every $f : \mathbb{N} \rightarrow \mathbb{N}$, there exists some $m > 0$ such that for all $n > m$,

$$G(f(0), \dots, f(n)) = \begin{cases} 1, & \text{if } f \in S; \\ 0, & \text{if } f \notin S. \end{cases}$$

A set $S \subseteq \mathbb{N}^{\mathbb{N}}$ is *guessable* if it has a guesser.

The next test is very useful for showing nonguessability, though its converse is not true.

Theorem 1. Let $S \subseteq \mathbb{N}^{\mathbb{N}}$. Suppose that, for every finite sequence $g \in \mathbb{N}^{<\mathbb{N}}$, there are sequences $g_1, g_2 \in \mathbb{N}^{\mathbb{N}}$ extending g with $g_1 \in S$ and $g_2 \notin S$. Then S is nonguessable.

Proof. Suppose S has a guesser G . I'll define a sequence $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $G(f(0), \dots, f(n))$ fails to converge, which violates the definition of guesser.

Clearly $S \neq \emptyset$, so let $s_1 : \mathbb{N} \rightarrow \mathbb{N}$ be some sequence in S . By definition of guesser, we can find some x_1 such that $G(s_1(0), \dots, s_1(x_1)) = 1$. Let $f(0) = s_1(0), \dots, f(x_1) = s_1(x_1)$.

Inductively, suppose I've defined $x_1 < \dots < x_k$ and $f(0), \dots, f(x_k)$ such that $G(f(0), \dots, f(x_i)) \equiv i \pmod{2}$ for $i = 1, \dots, k$. By the theorem's hypothesis, we can find some $s_{k+1} : \mathbb{N} \rightarrow \mathbb{N}$, extending the finite sequence $(f(0), \dots, f(x_k))$, such that s_{k+1} is in S iff $k+1 \equiv 1 \pmod{2}$. By definition of guesser, find $x_{k+1} > x_k$ such that $G(s_{k+1}(0), \dots, s_{k+1}(x_{k+1})) \equiv k+1 \pmod{2}$. Let $f(x_k+1) = s_{k+1}(x_k+1), \dots, f(x_{k+1}) = s_{k+1}(x_{k+1})$.

This defines sequences $f : \mathbb{N} \rightarrow \mathbb{N}$ and $x_1 < x_2 < \dots$ with the property that $G(f(0), \dots, f(x_i)) \equiv i \pmod{2}$ for every $i > 0$. This contradicts that $G(f(0), \dots, f(n))$ is supposed to converge. ■

Using this test, we can immediately confirm, for example, the set of sequences containing infinitely many zeroes is nonguessable, as is the set of onto sequences.

Remark 1. Theorem 1 is constructive up to certain choices. Starting with a set S satisfying the hypotheses of Theorem 1 and naively trying to guess it,

and being systematic in the choices from the proof, can lead to the creation of a concrete sequence which thwarts the naive guessing attempt. In an informal sense, it should be especially difficult for someone not in the know to guess whether the resulting sequence lies in S . And the more sophisticated the futile guessing attempt, the more difficult the resulting sequence becomes. For some explicit examples, see sequences A082691, A182659, and A182660 in Sloane’s OEIS [4].

Tsaban and Zdomskyy also briefly mention a somewhat similar notion of guessable sets in their paper [5].

3 A Logic for Ellipses

Because guessers are functions which do not have “arity” in the usual sense, instead being defined on the whole space $\mathbb{N}^{<\mathbb{N}}$ of finite sequences, and since we care so much about expressions like $G(f(0), \dots, f(n))$, we will extend logic to mesh better with these sorts of expressions. I assume familiarity with basic first-order logic, which Enderton [2] has written about extensively, as has Bilaniuk [1].

Definition 2. A *language with ellipses* is a standard language of first-order logic, with a constant symbol $\mathbf{0}$, together with a set of *function symbols of arity* $\mathbb{N}^{<\mathbb{N}}$ and a special logical symbol \dots_x for every variable x .

To avoid confusion, we’ll use \dots_x for the syntactical symbol and \dots for meta-ellipses. For example, $G(s(\mathbf{0}), \dots, s(\mathbf{2}))$ is a meta-abbreviation for

$$G(s(\mathbf{0}), s(\mathbf{1}), s(\mathbf{2})),$$

different than $G(s(\mathbf{0}), \dots_x, s(\mathbf{2}))$ which has no counterpart in classical logic.

Definition 3. If \mathcal{L} is a language with ellipses, then the *terms* of \mathcal{L} (and their free variables) are defined inductively:

1. x is a term, for any variable x , and $FV(x) = \{x\}$.
2. c is a term, for any constant symbol c , and $FV(c) = \emptyset$.
3. If f is a function symbol of arity n or arity $\mathbb{N}^{<\mathbb{N}}$, and t_1, \dots, t_n are terms, then $f(t_1, \dots, t_n)$ is a term with free vars $FV(t_1) \cup \dots \cup FV(t_n)$.

4. If G is an $\mathbb{N}^{<\mathbb{N}}$ -ary function symbol, and u, v are terms, and x is a variable, then $G(u(\mathbf{0}), \dots_x, u(v))$ is a term with free variables

$$(FV(u) \setminus \{x\}) \cup FV(v).$$

The *well-formed formulas* of \mathcal{L} are defined as usual from these terms. Term substitution is defined by the usual induction with two new cases:

- If $y \neq x$ then

$$G(u(\mathbf{0}), \dots_x, u(v))(y|t) = G(u(y|t)(\mathbf{0}), \dots_x, u(y|t)(v(y|t))).$$

- $G(u(\mathbf{0}), \dots_x, u(v))(x|t) = G(u(\mathbf{0}), \dots_x, u(v(x|t)))$.

A *model* for a language with ellipses \mathcal{L} is a model \mathcal{M} for the classical part of \mathcal{L} , together with a function $G^{\mathcal{M}} : \mathcal{M}^{<\mathbb{N}} \rightarrow \mathcal{M}$ for each $\mathbb{N}^{<\mathbb{N}}$ -ary function symbol G in \mathcal{L} . However, defining how an arbitrary model evaluates terms is difficult. We'll only be interested in one very specific family of models, where there is no trouble evaluating terms.

Definition 4. The following models lie at the heart of all later results.

- \mathcal{L}_{\max} is the language with ellipses which contains a constant symbol \mathbf{n} for every $n \in \mathbb{N}$, an n -ary function symbol \tilde{w} for every function $w : \mathbb{N}^n \rightarrow \mathbb{N}$ ($n > 0$), an n -ary predicate symbol \tilde{p} for every subset $p \subseteq \mathbb{N}^n$ ($n > 0$), an $\mathbb{N}^{<\mathbb{N}}$ -ary function symbol \tilde{G} for every function $G : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N}$, and one additional unary function symbol \mathbf{f} .
- For every function $f : \mathbb{N} \rightarrow \mathbb{N}$, \mathcal{M}_f is the model, on the language \mathcal{L}_{\max} , with universe \mathbb{N} , which interprets \mathbf{n} as n for every n , and interprets \tilde{w} as w for every $w : \mathbb{N}^n \rightarrow \mathbb{N}$, and which interprets \tilde{p} as p for every $p \subseteq \mathbb{N}^n$, and which interprets \tilde{G} as G for every $G : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N}$, and which interprets \mathbf{f} as f .

If $n \in \mathbb{N}$ then \bar{n} denotes the numeral \mathbf{n} of n .

Definition 5. Let $f : \mathbb{N} \rightarrow \mathbb{N}$. The semantics of \mathcal{M}_f are defined as follows. Let s be any assignment from the variables to \mathbb{N} .

- (\mathcal{M}_f, s) interprets terms t into naturals $t^{\mathcal{M}_f, s}$, or t^s if there is no ambiguity, according to the usual inductive definition, with one new case:

- If u, v are terms and x is a variable and G is an $\mathbb{N}^{<\mathbb{N}}$ -ary function symbol, then

$$G(u(\mathbf{0}), \dots_x, u(v))^s = G^{\mathcal{M}_f}(u(x|\mathbf{0})^s, \dots, u(x|\overline{v^s})^s).$$

- For example, the interpretation of $\tilde{G}(\mathbf{f}(x)(\mathbf{0}), \dots_x, \mathbf{f}(x)(\mathbf{99}))$ is

$$G(f(0), \dots, f(99)),$$

while the interpretation of $\tilde{G}(\mathbf{f}(x)(\mathbf{0}), \dots_x, \mathbf{f}(x)(\mathbf{f}(y)))$ is

$$G(f(0), \dots, f(f(y^s))).$$

- From here, the remaining semantics of \mathcal{M}_f are defined as usual.

In classical logic, every term with no free variables has the property that its interpretation in any model depends only on finitely many values of the interpretations of the function symbols in that model. For example, the interpretation of $5 + (2 \cdot 3)$ depends only on one value of \cdot and one value of $+$. Similar properties are true of our \mathcal{M}_f models.

Lemma 2. Suppose u is a term with no free variables, and c is a constant symbol. For any $f : \mathbb{N} \rightarrow \mathbb{N}$, $\mathcal{M}_f \models u = c$ iff there is some k such that whenever $g : \mathbb{N} \rightarrow \mathbb{N}$ extends $(f(0), \dots, f(k))$, $\mathcal{M}_g \models u = c$ and to check whether $\mathcal{M}_g \models u = c$ using the inductive definition of semantics for \mathcal{M}_g , it is not necessary to query $g(i)$ for any $i > k$.

Proof. (\Rightarrow): Induction on complexity of u .

- u cannot be a variable, since it has no free variables. If u is a constant symbol, the lemma is trivial.
- Maybe u is $h(u_1, \dots, u_n)$ for some n -ary (or $\mathbb{N}^{<\mathbb{N}}$ -ary) function symbol h other than \mathbf{f} , and some terms u_1, \dots, u_n with no free variables. If $\mathcal{M}_f \models h(u_1, \dots, u_n) = c$, then there are $a_1, \dots, a_n \in \mathbb{N}$ such that $h^{\mathcal{M}_f}(a_1, \dots, a_n) = c^{\mathcal{M}_f}$ and $\mathcal{M}_f \models u_i = \bar{a}_i$ for $i = 1, \dots, n$. Since \bar{a}_i is a constant symbol, by induction find k_1, \dots, k_n such that for any $i = 1, \dots, n$ and any $g : \mathbb{N} \rightarrow \mathbb{N}$ with $g(0) = f(0), \dots, g(k_i) = f(k_i)$, $\mathcal{M}_g \models u_i = \bar{a}_i$, and checking this by definition of semantics does not require querying $g(j)$ for any $j > a_i$. Then $k = \max\{k_1, \dots, k_n\}$ works (using the fact $h^{\mathcal{M}_g}$ does not depend on g since h is not \mathbf{f}).

- Maybe u is $\mathbf{f}(v)$ where v is a term with no free variables. If $\mathcal{M}_f \models \mathbf{f}(v) = c$ then there's $a \in \mathbb{N}$ such that $f(a) = c^{\mathcal{M}_f}$ and $\mathcal{M}_f \models v = \bar{a}$. Since \bar{a} is a constant symbol, by induction find k_0 such that whenever $g(0) = f(0), \dots, g(k_0) = f(k_0)$, then $\mathcal{M}_g \models v = \bar{a}$, and checking $\mathcal{M}_g \models v = \bar{a}$ does not require querying $g(i)$ for any $i > k_0$. Let $k = \max\{k_0, a\}$. Suppose $g(0) = f(0), \dots, g(k) = f(k)$. Then $\mathbf{f}(v)^{\mathcal{M}_g} = \mathbf{f}^{\mathcal{M}_g}(v^{\mathcal{M}_g}) = g(a) = f(a) = c^{\mathcal{M}_f}$. So $\mathcal{M}_g \models \mathbf{f}(v) = c$, and to check so, we only had to query $g(a)$ in addition to any queries we had to make to check $\mathcal{M}_g \models v = \bar{a}$, so we did not have to query $g(i)$ for any $i > k$.
- Maybe u is $G(v(\mathbf{0}), \dots_x, v(w))$ where v, w are terms, x is a variable, $FV(w) = \emptyset$, $FV(v) \subseteq \{x\}$, and G is an $\mathbb{N}^{<\mathbb{N}}$ -ary function symbol. If $\mathcal{M}_f \models G(v(\mathbf{0}), \dots_x, v(w)) = c$ then

$$G^{\mathcal{M}_f} \left(v(x|\mathbf{0})^{\mathcal{M}_f}, \dots, v \left(x \mid \overline{w^{\mathcal{M}_f}} \right)^{\mathcal{M}_f} \right) = c^{\mathcal{M}_f}.$$

Since $\mathcal{M}_f \models w = \overline{w^{\mathcal{M}_f}}$, find some number k_{-1} such that whenever g extends $(f(0), \dots, f(k_{-1}))$, $\mathcal{M}_g \models w = \overline{w^{\mathcal{M}_f}}$ and checking so does not require queries beyond $g(k_{-1})$. Since $\mathcal{M}_f \models v(x|\mathbf{i}) = \overline{v(x|\mathbf{i})}$ for $i = 0, \dots, w^{\mathcal{M}_f}$, find $k_0, \dots, k_{w^{\mathcal{M}_f}}$ such that for each $i = 0, \dots, w^{\mathcal{M}_f}$, if $g(0) = f(0), \dots, g(k_i) = f(k_i)$ then $\mathcal{M}_g \models v(x|\bar{i}) = \overline{v(x|\bar{i})}$ can be confirmed without querying g beyond $g(k_i)$.

Let $k = \max\{k_{-1}, k_0, \dots, k_{w^{\mathcal{M}_f}}\}$. Suppose $g(0) = f(0), \dots, g(k) = f(k)$. Then $\mathcal{M}_g \models w = \overline{w^{\mathcal{M}_f}}$, so $w^{\mathcal{M}_g} = w^{\mathcal{M}_f}$. Similarly $v(x|\mathbf{i})^{\mathcal{M}_g} = \overline{v(x|\mathbf{i})^{\mathcal{M}_f}}$ for $i = 0, \dots, w^{\mathcal{M}_f}$. And $G^{\mathcal{M}_g} = G^{\mathcal{M}_f}$. It follows that

$$\mathcal{M}_g \models G(v(\mathbf{0}), \dots_x, v(w)) = c,$$

and checking so does not require any queries to $g(j)$ for any $j > k$.

(\Leftarrow). Suppose there's some k so that whenever g extends $(f(0), \dots, f(k))$ then $\mathcal{M}_g \models u = c$ (nevermind the extra information about queries). In particular, f itself extends $(f(0), \dots, f(k))$, so $\mathcal{M}_f \models u = c$. \blacksquare

Corollary 3. Let ϕ be a quantifier-free sentence. For any $f : \mathbb{N} \rightarrow \mathbb{N}$, $\mathcal{M}_f \models \phi$ iff there's some k such that for every $g : \mathbb{N} \rightarrow \mathbb{N}$ extending $(f(0), \dots, f(k))$, $\mathcal{M}_g \models \phi$, and in checking $\mathcal{M}_g \models \phi$ by the inductive definition of semantics, we never need to query $g(i)$ for any $i > k$.

Proof. • Maybe ϕ is $u = v$ for terms u, v with no free variables. Assume $\mathcal{M}_f \models u = v$. Then $\mathcal{M}_f \models u = \overline{u}^{\mathcal{M}_f}$ and $\mathcal{M}_f \models v = \overline{v}^{\mathcal{M}_f}$. By Lemma 2, find k big enough that whenever $g(0) = f(0), \dots, g(k) = f(k)$, then $\mathcal{M}_g \models u = \overline{u}^{\mathcal{M}_g}$ and $\mathcal{M}_g \models v = \overline{v}^{\mathcal{M}_g}$ and both facts can be confirmed without querying g beyond $g(k)$. For any such g , $\mathcal{M}_g \models u = v$, verifiable with no additional g -queries. The converse is trivial.

- Maybe ϕ is $\tilde{p}(u_1, \dots, u_n)$ for an n -ary predicate symbol \tilde{p} and terms u_1, \dots, u_n with no free variables. Then ϕ is equivalent (in every \mathcal{M} .) to $\tilde{g}(u_1, \dots, u_n) = \mathbf{1}$ where g is the characteristic function of p , so we're done by the previous case.
- Maybe ϕ is $\phi_1 \wedge \phi_2$. Assume $\mathcal{M}_f \models \phi$. Inductively, find k_1 and k_2 such that if g extends $(f(0), \dots, f(k_i))$ then $\mathcal{M}_g \models \phi_i$ is verifiable with no g -queries beyond $g(k_i)$. Then any g extending $(f(0), \dots, f(\max\{k_1, k_2\}))$ has $\mathcal{M}_g \models \phi$, verifiable without querying beyond $g(\max\{k_1, k_2\})$. The converse is trivial.
- The cases of other propositional connectives are similar. ■

Lemma 4. (*The Weak Substitution Lemma*) For a formula ϕ , an assignment s , and a constant symbol c , and for any $f : \mathbb{N} \rightarrow \mathbb{N}$, $\mathcal{M}_f \models \phi[s(x|\overline{c^s})]$ iff $\mathcal{M}_f \models \phi(x|c)[s]$.

Proof. In the Appendix. ■

A full Substitution Lemma is also true, but it requires a nonclassical definition of *substitutable*, which goes too far away from the matters this paper is actually concerned with.

4 Guessability and Quantifiers

Definition 6. Let $S \subseteq \mathbb{N}^{\mathbb{N}}$ be a set of sequences. Let ϕ be a sentence in \mathcal{L}_{\max} . We say that ϕ *defines* S if, for every $f : \mathbb{N} \rightarrow \mathbb{N}$, $\mathcal{M}_f \models \phi$ iff $f \in S$.

Lemma 5. Suppose $S \subseteq \mathbb{N}^{\mathbb{N}}$ is guessable. Then S is defined by some sentence $\exists x \forall y \phi$ and also by some sentence $\forall x \exists y \psi$, where ϕ and ψ are quantifier-free.

Proof. S is guessed by a guesser G . For any $f : \mathbb{N} \rightarrow \mathbb{N}$, $f \in S$ if and only if $G(f(0), \dots, f(n)) = 1$ for all n sufficiently large. Therefore S is defined by

$$\exists x \forall y ((y > x) \rightarrow \tilde{G}(\mathbf{f}(z)(\mathbf{0}), \dots_z, \mathbf{f}(z)(y)) = \mathbf{1}),$$

where “ $y > x$ ” is shorthand for $\tilde{>}(y, x)$. Similarly, S is also defined by

$$\forall x \exists y ((y > x) \wedge \tilde{G}(\mathbf{f}(z)(\mathbf{0}), \dots_z, \mathbf{f}(z)(y)) = \mathbf{1}).$$

■

We will prove the converse of Lemma 5 shortly. To that end, a piece of technical machinery is needed.

Definition 7. A set $S \subseteq \mathbb{N}^{\mathbb{N}}$ is *overguessable* if there is a function $\mu : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N} \cup \{\infty\}$ such that:

1. For any $f \in S$, $\mu(f(0), \dots, f(n))$ is eventually bounded by a finite number.
2. For any $f \notin S$, $\mu(f(0), \dots, f(n)) \rightarrow \infty$ as $n \rightarrow \infty$.

Lemma 6. Suppose $S \subseteq \mathbb{N}^{\mathbb{N}}$ is defined by the sentence $\exists x \forall y \phi$ where ϕ is quantifier-free. Then S is overguessable.

Proof. Given a tuple (n_0, \dots, n_k) , define $\mu(n_0, \dots, n_k)$ as follows. Let $h : \mathbb{N} \rightarrow \mathbb{N}$ be defined by $h(i) = n_i$ if $i \leq k$, $h(i) = 0$ otherwise. Given a pair $(a, b) \in \mathbb{N}^2$, consider the sentence $\phi(x, y | \bar{a}, \bar{b})$. Attempt to check whether $\mathcal{M}_h \models \phi(x, y | \bar{a}, \bar{b})$, using the inductive definition of the semantics of \mathcal{M}_h . If, in so doing, you must query $h(i)$ for some $i > k$, say that *the attempt failed*. Otherwise, say *the attempt succeeded*. If the attempt failed, *or* if $\mathcal{M}_h \models \phi(x, y | \bar{a}, \bar{b})$, then say that (a, b) is *nice*.

Call a number a *very nice* if (a, b) is nice for every b . If there is any very nice number, then let $\mu(n_0, \dots, n_k)$ be the smallest very nice number. Otherwise let $\mu(n_0, \dots, n_k) = \infty$.

I claim this μ works.

First, suppose $f \in S$. Since S is defined by $\exists x \forall y \phi$, $\mathcal{M}_f \models \exists x \forall y \phi$. By the Weak Substitution Lemma, for some a , $\mathcal{M}_f \models \phi(x, y | \bar{a}, \bar{b})$ for every b . When we attempt to check whether $\mathcal{M}_h \models \phi(x, y | \bar{a}, \bar{b})$ in the definition of $\mu(f(0), \dots, f(k))$, if the attempt succeeds, then $\mathcal{M}_h \models \phi(x, y | \bar{a}, \bar{b})$ because $\mathcal{M}_f \models \phi(x, y | \bar{a}, \bar{b})$ and we never had to look at the part of h which disagrees

with f . So (a, b) is nice for every b , so a is very nice, so $\mu(f(0), \dots, f(k))$ is bounded by a .

Next, suppose $f \notin S$. Let $a \in \mathbb{N}$, I claim $\mu(f(0), \dots, f(n)) \neq a$ for all n sufficiently large. Since $f \notin S$, $\mathcal{M}_f \not\models \exists x \forall y \phi$. By the Weak Substitution Lemma, there is some b such that $\mathcal{M}_f \not\models \phi(x, y | \bar{a}, \bar{b})$. Since ϕ is a quantifier-free sentence, we invoke Corollary 3 on $\neg\phi(x, y | \bar{a}, \bar{b})$ and find k such that $\mathcal{M}_g \not\models \phi(x, y | \bar{a}, \bar{b})$ whenever g extends $(f(0), \dots, f(k))$, and, to check whether $\mathcal{M}_g \models \phi(x, y | \bar{a}, \bar{b})$, we do not need to query $h(i)$ for $i > k$. Then, in the definition of $\mu(f(0), \dots, f(k))$, for pair (a, b) , the attempt succeeds and $\mathcal{M}_h \not\models \phi(x, y | \bar{a}, \bar{b})$, so (a, b) is not nice, so a is not very nice, so $\mu(f(0), \dots, f(k)) \neq a$, in fact, $\mu(f(0), \dots, f(j)) \neq a$ for all $j \geq k$. By arbitrariness of a , $\mu(f(0), \dots, f(n)) \rightarrow \infty$. ■

Proposition 7. The converse of Lemma 6 also holds. $S \subseteq \mathbb{N}^{\mathbb{N}}$ is defined by some sentence $\exists x \forall y \phi$ with ϕ quantifier-free, iff S is overguessable.

Proof. (\Rightarrow) is Lemma 6. Conversely, suppose S is overguessed by $\mu : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N} \cup \{\infty\}$. Define $\mu' : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}$ by saying $\mu'(n) = \mu(n) + 1$ if $\mu(n) \neq \infty$, $\mu'(n) = 0$ if $\mu(n) = \infty$. For any $f : \mathbb{N} \rightarrow \mathbb{N}$, $f \in S$ iff $\mu(f(0), \dots, f(n))$ is eventually bounded by some finite number, which is true iff $\mu'(f(0), \dots, f(n))$ is eventually bounded by some finite number and eventually nonzero. This latter equivalence can be expressed by

$$\exists m_1 \exists m_2 \forall m_3 ((m_3 > m_2) \rightarrow (0 < \mu'(f(0), \dots, f(m_3)) < m_1)).$$

Let $d : \mathbb{N} \rightarrow \mathbb{N}^2$ be any onto map from \mathbb{N} to \mathbb{N}^2 . Write $d(n) = (d_1(n), d_2(n))$, thus defining two functions $d_1, d_2 : \mathbb{N} \rightarrow \mathbb{N}$. Then the above formula is equivalent to

$$\exists m \forall m_3 ((m_3 > d_2(m)) \rightarrow (0 < \mu'(f(0), \dots, f(m_3)) < d_1(m))).$$

This can be formalized in \mathcal{L}_{\max} , providing a sentence $\exists x \forall y \phi$ which defines S , with ϕ quantifier-free. ■

Example 8. Every countable subset of $\mathbb{N}^{\mathbb{N}}$ is overguessable.

Proof. Let $S \subseteq \mathbb{N}^{\mathbb{N}}$ be countable. Define $g : \mathbb{N}^2 \rightarrow \mathbb{N}$ by saying $g(m, n) = h_m(n)$ where h_m is the m th element of S . Then S is defined by

$$\exists x \forall y \tilde{g}(x, y) = \mathbf{f}(y).$$

By Lemma 6, S is overguessable. ■

Theorem 9. A set $S \subseteq \mathbb{N}^{\mathbb{N}}$ is guessable if and only if it is defined by some sentence $\forall x \exists y \phi$ and also by some sentence $\exists x \forall y \psi$, where ϕ and ψ are quantifier-free.

Proof. (\Rightarrow) By Theorem 5.

(\Leftarrow) Suppose S is defined by sentences $\forall x \exists y \phi$ and $\exists x \forall y \psi$, where ϕ and ψ are quantifier-free.

By Lemma 6, find $\mu : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N} \cup \{\infty\}$ which overguesses S . And since S^c is defined by $\exists x \forall y \neg \phi$, use Lemma 6 again to find $\nu : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N} \cup \{\infty\}$ which overguesses S^c .

Define $G : \mathbb{N}^{<\mathbb{N}} \rightarrow \{0, 1\}$ by saying $G(n_0, \dots, n_k) = 1$ if $\mu(n_0, \dots, n_k) \leq \nu(n_0, \dots, n_k)$ and 0 otherwise. If $f \in S$ then $\mu(f(0), \dots, f(k))$ is eventually bounded by a finite number and $\nu(f(0), \dots, f(k)) \rightarrow \infty$, so $G(f(0), \dots, f(k))$ converges to 1. The other case is similar. ■

Remark 2. Guessable and overguessable sets of sequences are analogous to computable and computably enumerable sets of naturals, respectively. One shows that Δ_1 sets (in a much weaker logical setting than \mathcal{L}_{\max}) of naturals are computable by showing that they and their complements are c.e. by using the characterization of c.e. sets as sets which are Σ_1 -definable (in the weaker setting). By comparison, I've shown that Δ_2 sets of sequences (in a very strong logical setting) are guessable by showing that they and their complements are overguessable by using the characterization of overguessable sets as Σ_2 -definable (in the stronger setting). These analogous phenomena in computability theory have been written about by Rogers [3], Enderton [2], Bilaniuk [1], and many other authors.

Corollary 10. Suppose $S \subseteq \mathbb{N}^{\mathbb{N}}$ is definable by a sentence $\forall x \exists y \phi$ where ϕ is quantifier-free. If S is countable then S is guessable.

Proof. Suppose S is countable. In the proof of Example 8, we showed S is definable by a sentence $\exists x \forall y \psi$ where ψ is quantifier-free. By Theorem 9, S is guessable. ■

Example 11. There are uncountably many permutations of \mathbb{N} .

Proof. A function $f : \mathbb{N} \rightarrow \mathbb{N}$ is a permutation iff

$$\forall m_1 \forall m_2 \exists n ((f(m_1) = f(m_2) \rightarrow m_1 = m_2) \wedge f(n) = m_2).$$

By appropriately coding $\langle m_1, m_2 \rangle$, the set S of permutations is defined by a sentence $\forall x \exists y \phi$ where ϕ is quantifier-free.

Permutations are not guessable. If G were a permutation-guesser, it would diverge on the following sequence. Let $f(0) = 0$, $f(1) = 1$, and so on until $G(f(0), \dots, f(k_1)) = 1$ (this must happen since G would converge to 1 if we kept going forever). Then skip a number, $f(k_1 + 1) = k_1 + 2$, $f(k_1 + 2) = k_1 + 3$, and keep going until $G(f(0), \dots, f(k_2)) = 0$. Then fill in the gap, $f(k_2 + 1) = k_1 + 1$, and resume where we left off, $f(k_2 + 2) = k_2 + 2$, and so on until $G(f(0), \dots, f(k_3)) = 1$. This process shows permutations are unguessable.

By Corollary 10, S is uncountable. ■

Example 12. (*Cantor*) There are uncountably many real numbers.

Proof. Consider the set A of irrationals in the interval $(0, 1)$ which have infinitely many 5s in their decimal expansions. There is an obvious bijection between A and the set S of sequences $f : \mathbb{N} \rightarrow \{0, 9\}$ such that $f(n) = 5$ infinitely often. This set S is defined by $\forall x \exists y ((y > x) \wedge \mathbf{f}(y) = \mathbf{5})$. By Corollary 10, if S is countable then it is guessable. But it is not: if G were a guesser for S , then we could define a sequence on whose initial segments G does not converge. Namely, let $f(0) = \dots = f(x_k) = 0$ where x_k is big enough that $G(f(0), \dots, f(x_k)) = 0$, and then let $f(x_k + 1) = \dots = f(x_{k+1}) = 5$, where $x_{k+1} > x_k$ is big enough that $G(f(0), \dots, f(x_{k+1})) = 1$. And so on, alternating, forever. This shows S is not guessable, so S is not countable, so A is uncountable, so \mathbb{R} is uncountable. ■

Lemma 13. If $S \subseteq \mathbb{N}^{\mathbb{N}}$ is definable by a sentence ϕ without nested quantifiers (that is, no quantifier appearing in the scope of another), then S is guessable.

Proof. If so, then ϕ is a propositional combination of quantifier-free sentences and sentences of the form $\forall x \phi_0$ and $\exists x \phi_1$ where ϕ_0, ϕ_1 are quantifier-free. The sets defined by these component sentences are guessable by Theorem 9. Clearly guessable sets are closed under union, intersection, and complement, so S itself is guessable. ■

Example 14. The definition of onto functions cannot be simplified to get rid of nested quantifiers, not even with the full power of \mathcal{L}_{\max} .

Proof. By Lemma 13 and the fact the set of onto functions is not guessable (by Theorem 1). ■

Appendix: The Weak Substitution Lemma

We prove Theorem 4, which said: For a formula ϕ , an assignment s , and a constant symbol c , and for any $f : \mathbb{N} \rightarrow \mathbb{N}$, $\mathcal{M}_f \models \phi[s(x|\overline{c^s})]$ iff $\mathcal{M}_f \models \phi(x|c)[s]$.

Proof. By the inductive argument used to prove the full Substitution Lemma in classical logic, most of which we omit. But there are tricky new cases for our new terms.

Claim: For any terms u, v , constant symbol c , variables $x \neq y$, and assignment s ,

$$G(u(\mathbf{0}), \dots_x, u(v))(y|c)^s = G(u(\mathbf{0}), \dots_x, u(v))^{s(y|c)}.$$

The details are (using the induction hypothesis repeatedly):

$$\begin{aligned} G(u(\mathbf{0}), \dots_x, u(v))(y|c)^s &= G(u(y|c)(\mathbf{0}), \dots_x, u(y|c)(v(y|c)))^s \\ &= G^{\mathcal{M}_f} \left(u(y|c)(x|\mathbf{0})^s, \dots, u(y|c) \left(x \mid \overline{v(y|c)^s} \right)^s \right) \\ &= G^{\mathcal{M}_f} \left(u(x|\mathbf{0})^{s(y|c)}, \dots, u \left(x \mid \overline{v^{s(y|c)}} \right)^{s(y|c)} \right) \\ &= G(u(\mathbf{0}), \dots_x, u(v))^{s(y|c)}. \end{aligned}$$

Claim: For any terms u, v , constant symbol c , and variable x and assignment s ,

$$G(u(\mathbf{0}), \dots_x, u(v))(x|c)^s = G(u(\mathbf{0}), \dots_x, u(v))^{s(x|c)}.$$

Using the induction hypothesis repeatedly:

$$\begin{aligned} G(u(\mathbf{0}), \dots_x, u(v))(x|c)^s &= G(u(\mathbf{0}), \dots_x, u(v(x|c)))^s \\ &= G^{\mathcal{M}_f} \left(u(x|\mathbf{0})^s, \dots, u \left(x \mid \overline{v(x|c)^s} \right)^s \right) \\ &= G^{\mathcal{M}_f} \left(u(x|\mathbf{0})^s, \dots, u \left(x \mid \overline{v^{s(x|c)}} \right)^s \right) \\ &= G^{\mathcal{M}_f} \left(u(x|\mathbf{0})^{s(x|c)}, \dots, u \left(x \mid \overline{v^{s(x|c)}} \right)^{s(x|c)} \right) \\ &= G(u(\mathbf{0}), \dots_x, u(v))^{s(x|c)}. \end{aligned}$$

The 2nd-to-last equation is justified because the terms whose “exponents” are changed do not depend on x . ■

A full Substitution Lemma also holds in \mathcal{M}_f , but it requires a non-classical definition of substitutability and we don’t need it for our results anyway.

References

- [1] S. Bilaniuk, *A Problem Course in Mathematical Logic*. Self-published, 1991. <http://euclid.trentu.ca/math/sb/pcml/pcml-16.pdf>
- [2] H. Enderton, *A Mathematical Introduction to Logic*. Academic Press, London, 2nd Edition, 2001.
- [3] H. Rogers, *Theory of Recursive Functions and Effective Computability*. The MIT Press, Cambridge, 1987.
- [4] N. J. A. Sloane, Online Encyclopedia of Integer Sequences (OEIS), <http://www.oeis.org/>.
- [5] B. Tsaban & L. Zdomskyy, Combinatorial images of sets of reals and semifilter trichotomy, *J. of Symbolic Logic* **73** (2008), 1278-1288.