

AN ADIC DYNAMICAL SYSTEM RELATED TO THE DELANNOY NUMBERS

KARL PETERSEN

In memory of Dan Rudolph—friend, colleague, and inspiration.

ABSTRACT. We introduce an adic (Bratteli-Vershik) dynamical system based on a diagram whose path counts from the root are the Delannoy numbers. We identify the ergodic invariant measures, prove total ergodicity for each of them, and initiate the study of the dimension group and other dynamical properties.

1. INTRODUCTION

Adic, or Bratteli-Vershik, dynamical systems present the cutting and stacking constructions of ergodic theory in a form that is convenient for studying important relations such as orbit equivalence; they also draw connections with the theory of C^* algebras and group representations and raise interesting questions in combinatorics and number theory. In recent years some progress has been made on the dynamics of several natural non-stationary and non-simple adic systems, such as the Pascal, Euler, reverse Euler, and Stirling systems. See [4] for a discussion of how such systems arise from walks and reinforced walks on graphs and [1–3, 8, 9, 11, 12, 15, 16] and the references that these papers contain for background on these and other adic systems. In this paper, by investigating the dynamical properties of a particular example, which we call the *Delannoy adic*, we initiate study of a class of adic systems which, though non-stationary, still are highly regular in that each vertex has the same pattern of exiting edges. We begin with a brief discussion of how we came across this particular diagram.

In a music theory seminar at IRCAM in Paris, Norman Carey and David Clampitt discussed a diagram attributed to Nicomachus, and later studied by Nicole Oresme, which was supposed to have wonderful properties, including forming a basis for the generation of various kinds of musical scales [7]. The diagram, with entries $2^n 3^k$ at the points (n, k) of the integer lattice in the plane, is shown in Figure 1.

Naturally one may ask whether this might turn into a Bratteli diagram in such a way that the entries $2^a 3^b$ would be the dimensions of vertices, that is, the numbers of paths from the

Date: January 26, 2013.

1991 *Mathematics Subject Classification.* Primary: 37A05, 37A25, 05A10, 05A15; Secondary: 37A50, 37A55.

Key words and phrases. Adic transformation, invariant measure, ergodicity, Delannoy numbers.




FIGURE 1. The first part of the Nicomachus diagram.

root (lower left corner) to the vertices. Adding diagonals to the boxes makes this happen, as seen in Figure 2.




FIGURE 2. The Nicomachus diagram with added diagonals. Numbers next to edges indicate multiplicities greater than 1.

As we will see below, this does not produce an especially interesting adic dynamical system, since the only ergodic invariant measures are supported on the two boundary odometers. Reducing the edge multiplicities to 1 produces a figure whose path counts are the *Delannoy numbers* [5] $D(n, k)$, for $n, k \geq 0$; see Figure 3.




FIGURE 3. The Delannoy graph.

Notice, however, that this is not a legitimate Bratteli diagram, since it is not true that there are connections only from one level (the set of vertices a fixed graph distance from $(0, 0)$) to the next; see Figure 4.

This can be remedied by inserting “extra” vertices at the centers of the squares. While this complicates the graph, the space of infinite paths beginning at the root is the same as before, and so we may regard this as a Bratteli diagram and adic dynamical system that realizes the Delannoy numbers as the dimensions of vertices; see Figure 5.

Here are some known key formulas involving the Delannoy numbers (see, for example, [17, 18]).

$$\begin{aligned}
 (1.1) \quad & D(n, 0) = D(0, n) = 1 \text{ for all } n \geq 0; \\
 & D(n, k) = 0 \text{ if either } n \text{ or } k < 0; \\
 & D(n, k) = D(n, k - 1) + D(n - 1, k - 1) + D(n - 1, k) \text{ for all } n, k.
 \end{aligned}$$

$$(1.2) \quad \sum_{n, k \geq 0} D(n, k) x^n y^k = \frac{1}{1 - (x + y + xy)}.$$




FIGURE 4. Levels in the Delannoy graph.




FIGURE 5. The Delannoy graph made into a Bratteli diagram.

Assuming $n \geq k$,

$$\begin{aligned}
 D(n, k) &= \sum_{d=0}^k \binom{k}{d} \binom{n+k-d}{k} = \sum_{d=0}^k 2^d \binom{n}{d} \binom{k}{d} \\
 (1.3) \quad &= \sum_{d=0}^k \binom{k}{d} \binom{n+d}{k} = \sum_{d=0}^k \binom{k}{k-d} \binom{n+d}{k} \\
 &= \sum_{d=0}^k \binom{n+k-d}{k-d} \binom{n}{d} = \sum_{d=0}^k \binom{n+d}{d} \binom{n}{k-d}.
 \end{aligned}$$

$$(1.4) \quad D(n, n) \sim (3 + 2\sqrt{2})^n (.57\sqrt{n} - .067n^{-3/2} + .006n^{-5/2} + \dots) \text{ as } n \rightarrow \infty.$$

Now we define the two dynamical systems which concern us. Each is based on a countably infinite directed graph with an ordering on the edges that enter each vertex. The *Nicomachus graph* (shown in Figure 2) has vertices $(n, k) \in \mathbb{Z}_+^2$ and an edge from (n, k) to $(n+1, k)$, to $(n, k+1)$, and to $(n+1, k+1)$ for all $n, k \geq 1$. From each vertex $(n, 0), n \geq 0$, there are two edges to $(n+1, 0)$ one edge to $(n+1, 1)$, and, if $n \geq 1$, one edge to $(n, 1)$. From each vertex $(0, k), k \geq 0$, there are three edges to $(0, k+1)$, one edge to $(1, k+1)$, and, if $k \geq 1$, one edge to $(1, k)$. Each doublet or triplet of edges entering a vertex $(n, 0)$ or $(0, k)$ is ordered arbitrarily. The three edges entering a vertex $(n+1, k+1)$ with $n, k \geq 0$ are ordered so that the one from $(n+1, k)$ is the first and the one from $(n, k+1)$ is the last.

The *Delannoy graph* (shown in Figure 3) has vertices $(n, k) \in \mathbb{Z}_+^2$ and an edge from (n, k) to $(n+1, k)$, to $(n, k+1)$, and to $(n+1, k+1)$ for all $n, k \geq 0$. Thus it is obtained from the Nicomachus graph by replacing the multiple edges along the x and y axes by single edges. Again the three edges entering a vertex $(n+1, k+1)$ with $n, k \geq 0$ are ordered so that the one from $(n+1, k)$ is the first and the one from $(n, k+1)$ is the last.

In each case the phase space X of our dynamical system will consist of infinite paths x that begin at the origin. The paths have labelings, either corresponding uniquely to their sequences of edges $e_i = e_i(x), i \geq 1$ (with labels from an alphabet $A = \{h, d, v\}$, for horizontal, diagonal, or vertical, respectively, and $A = \{h, d, v, h_1, h_2, v_1, v_2, v_3\}$ when there can be more than one edge between a pair of vertices), or to the sequences $(n_i, k_i) = (n_i(x), k_i(x)), i \geq 0$, of vertices through which they pass. The space X is a compact totally disconnected metric space when we define the distance from x to y to be less than or equal to $1/2^N$ if $e_i(x) = e_i(y)$ for all $i \leq N$. The topology of X is generated by the family of (clopen) *cylinder sets* defined by finite sequences (a_i) from A ,

$$(1.5) \quad [a_1 a_2 \dots a_I] = \{x \in X : e_i(x) = a_i \text{ for all } i \leq I\}.$$

Two paths $x, y \in X$ are *tail equivalent* if they coincide from some point on: there is a smallest N such that $e_i(x) = e_i(y)$ for all $i > N$. In such a case we define $x < y$ if $e_N(x) < e_N(y)$. Define X_{\max} to be the set of maximal paths, i.e. those that include only

maximal edges, and define X_{\min} to be the set of minimal paths. For each of the two graphs, $X_{\max} \cup X_{\min}$ is countable. The *adic transformation* $T : X \setminus X_{\max} \rightarrow X \setminus X_{\min}$ is defined by letting Tx equal the smallest $y \in X$ such that $y > x$. Then T is a homeomorphism. We define T and T^{-1} to be the identity on each of X_{\max} and X_{\min} . As extended, T is not continuous on X . The orbits of T coincide with the equivalence classes of the tail relation.

Acknowledgment. The author thanks Jean-Paul Allouche, James Damon, Sarah Bailey Frick, and François Ledrappier for helpful discussions; the University of Paris 6, the Wrocław University of Technology, and the Institut Henri Poincaré for hospitality while this research was underway; and the referees for suggestions that improved the exposition.

2. INVARIANT MEASURES

In the following, by “measure” we will mean “Borel probability measure”. For an invariant measure of an adic system, all cylinder sets determined by initial path segments from the root R to a fixed vertex v have the same measure. For vertices u and v in the diagram of the system, denote by $\dim(u, v)$ the number of paths in the diagram from u to v . From the Ergodic Theorem it follows that if μ is an ergodic invariant measure and C is a cylinder set determined by an initial path segment ending at a vertex v , then

$$(2.1) \quad \mu(C) = \lim_{n \rightarrow \infty} \frac{\dim(v, x_n)}{\dim(R, x_n)}$$

for μ -a.e. path $x = (x_n)$ (indexed here by its vertices). In the case of the Delannoy graph it is enough to specify the measure of each cylinder set determined by an initial path segment that ends at a “regular” vertex (n_0, k_0) in the integer lattice, since the measure of any cylinder ending at one of the “extra” vertices $(n_0 + 1/2, k_0 + 1/2)$ equals that of any cylinder ending at $(n_0 + 1, k_0 + 1)$. We will abbreviate by saying such a cylinder set “ends at (n_0, k_0) ”.

Thus to identify the ergodic measures, it is important to have information about path counts between vertices in the diagram that defines the system. The asymptotics of Delannoy numbers were determined by Pemantle and Wilson [13] by analytic methods:

$$(2.2) \quad D(n, k) \sim \left(\frac{\sqrt{n^2 + k^2} - k}{n} \right)^{-n} \left(\frac{\sqrt{n^2 + k^2} - n}{k} \right)^{-k} \sqrt{\frac{nk/(2\pi)}{(n+k - \sqrt{n^2 + k^2})^2 \sqrt{n^2 + k^2}}},$$

uniformly if n/k and k/n are bounded.

Theorem 2.1. *The only ergodic (invariant probability) measures for the Nicomachus adic dynamical system (described by Figure 2) are the two unique measures supported on the two boundary odometers.*

Proof. Let (n_0, k_0) be an interior regular vertex, i.e. $n_0, k_0 > 0$. We claim that

$$(2.3) \quad \frac{\dim((n_0, k_0), (n_0 + n, k_0 + k))}{\dim((0, 0), (n_0 + n, k_0 + k))} = \frac{D(n, k)}{2^{n_0+n} 3^{k_0+k}} \rightarrow 0 \quad \text{as } n+k \rightarrow \infty.$$

It follows that any ergodic invariant measure must assign measure 0 to any cylinder set determined by an initial path segment that ends at an interior vertex.

Let $0 < \epsilon < 1$. We consider first (n, k) with $\theta = k/n \in [\epsilon, 1]$. According to the Pemantle-Wilson estimate, for large n, k , uniformly in this region,

$$(2.4) \quad D(n, k) \sim (\sqrt{1+\theta^2} + \theta)^n \left(\frac{\sqrt{1+\theta^2} + 1}{\theta} \right)^{\theta n} \frac{\sqrt{\theta/(2\pi n)}}{\sqrt{(1+\theta - \sqrt{1+\theta^2})^2 \sqrt{1+\theta^2}}}.$$

Let

$$(2.5) \quad A(\theta) = (\sqrt{1+\theta^2} + \theta) \left(\frac{\sqrt{1+\theta^2} + 1}{\theta} \right)^\theta, \quad G(\theta) = \log A(\theta) - \log 2 - \theta \log 3,$$

and $B(\theta) = \frac{\sqrt{\theta}}{\sqrt{(1+\theta - \sqrt{1+\theta^2})^2 \sqrt{1+\theta^2}}}.$

We note that $B(\theta)$ is bounded for $\theta \in [\epsilon, 1]$ and $G(\theta)$ has an absolute maximum at $\theta = 3/4$, so that

$$(2.6) \quad \frac{A(\theta)}{2 \cdot 3^\theta} \leq \frac{A(3/4)}{2 \cdot 3^{3/4}} = 1$$

for all $\theta \in [\epsilon, 1]$. Hence

$$(2.7) \quad \frac{D(n, k)}{2^n 3^k} \leq c \left(\frac{A(\theta)}{2 \cdot 3^\theta} \right)^n \frac{1}{\sqrt{n}} \rightarrow 0$$

uniformly as $n \rightarrow \infty$ in this region.

Second, we consider (n, k) in the region where $k \leq \epsilon n$. Let

$$(2.8) \quad H(\epsilon) = -\epsilon \log \epsilon - (1 - \epsilon) \log(1 - \epsilon) \quad \text{and} \quad \lambda = e^{H(\epsilon)}.$$

Note that $0 \leq H(\epsilon) \leq \log 2$ for all $\epsilon \in [0, 1]$ and $\lambda \rightarrow 1$ as $\epsilon \rightarrow 0^+$. By Stirling's Formula,

$$(2.9) \quad \binom{n}{\epsilon n} \sim \lambda^n \quad \text{and}$$

$$\binom{\epsilon n}{d} \leq \binom{\epsilon n}{\epsilon n/2} \sim \frac{2^{\epsilon n}}{\sqrt{\pi \epsilon n/2}} \quad \text{for } d \leq k \leq \epsilon n.$$

Thus, since in this region $D(n, k)$ and $C(n, k)$ are increasing in k ,

$$(2.10) \quad D(n, k) = \sum_{d=0}^k 2^d \binom{n}{d} \binom{k}{d} \leq D(n, \epsilon n) \leq \sum_{d=0}^{\epsilon n} 2^d \lambda^n \binom{\epsilon n}{d}$$

$$\leq \epsilon n 2^{\epsilon n} \lambda^n \frac{2^{\epsilon n}}{\sqrt{\pi \epsilon n/2}} \sim c \sqrt{\epsilon n} \lambda^n 2^{2\epsilon n} = c \sqrt{\epsilon n} (2^{2\epsilon} \lambda)^n.$$

Choosing ϵ small enough that

$$(2.11) \quad 2^{2\epsilon} \lambda < 2,$$

we have that $D(n, k)/2^n \rightarrow 0$ uniformly in this region as $n \rightarrow \infty$.

In the region $k \geq n$, similar but easier estimates apply, since $D(n, k)$ is symmetric in n and k , and $2^n 3^k$ is now much larger than before. \square

Theorem 2.2. *The non-atomic ergodic (invariant probability) measures for the Delannoy adic dynamical system (described by Figure 3 or Figure 5) are a one-parameter family $\{\mu_\alpha : \alpha \in [0, 1]\}$ given by choosing nonnegative α, β, γ with $\alpha + \beta + \gamma = 1$ and $\beta\gamma = \alpha$ and then putting weight β on each horizontal edge, weight γ on each vertical edge, and weight α on each diagonal edge. (The measure of any cylinder set is then determined by multiplying the weights on the edges that define it.)*

Proof. Each μ_α as above is adic-invariant (by the conditions). It describes an i.i.d. random walker who starts at $(0, 0)$ and at each time $t = 0, 1, 2, \dots$ can choose independently whether to take a diagonal, horizontal, or vertical step. The probability of each finite initial path segment is the probability of the cylinder set that it defines. We will give two proofs that each μ_α is ergodic, first by means of the Hewitt-Savage 0,1 Law (see [6]), and then by a collision argument as in [2], using known conditions for recurrence of random walks.

(1) Identify the set X of infinite paths from the root in the Delannoy graph with $\{h, d, v\}^{\mathbb{N}}$ by using the edge labels. Then μ_α is a Bernoulli measure for the stationary process in which the symbols h, d, v arrive independently with probabilities β, α, γ , respectively. If $A \subset X$ is a Borel set that is invariant under the adic transformation T , then it is also invariant under permutations of finitely many coordinates. Therefore the σ -algebra of adic-invariant Borel sets is contained in the σ -algebra of symmetric sets (those which are invariant under permutations of finitely many coordinates). Since the latter algebra is trivial by the Hewitt-Savage Theorem, A has measure 0 or 1, and thus the measure μ_α is ergodic.

(2) The ‘‘collision argument’’ presented in [2] establishes ergodicity of μ_α once we show that for $\mu_\alpha \times \mu_\alpha$ -a.e. pair of infinite paths $(x, y) \in X \times X$, with ‘‘regular’’ vertices (n_i, k_i) and (r_i, s_i) respectively, there are infinitely many i for which $(n_i, k_i) = (r_i, s_i)$.

As mentioned above, paths $x \in X$ are trajectories of a random walk in \mathbb{Z}^2 with independent increments $(1, 1), (0, 1), (1, 0)$ having probabilities α, β, γ , respectively. We want to show recurrence (i.e., infinitely many returns to the origin of the symmetric (mean 0) process $Z = X - X$). This process has independent increments as shown in the first array below, each with probability given in the second array below:

$$(2.12) \quad \begin{pmatrix} (0, 0) & (0, 1) & (1, 0) \\ (0, -1) & (0, 0) & (1, -1) \\ (-1, 0) & (-1, 1) & (0, 0) \end{pmatrix} \quad \begin{pmatrix} \alpha^2 & \alpha\beta & \alpha\gamma \\ \beta\alpha & \beta^2 & \beta\gamma \\ \gamma\alpha & \gamma\beta & \gamma^2 \end{pmatrix}$$

(the rows and columns are each indexed by the possible increments listed above). Since the walk in \mathbb{Z}^2 has mean 0 and finite second moment, it is recurrent [19, p. 83].

(3) Now we prove that every adic-invariant ergodic measure for the Delannoy system is Bernoulli for the shift, i.e. is one of the measures μ_α described above, by applying the isotropy

arguments of [11]. Let μ be an adic-invariant ergodic measure. Since μ is non-atomic and adic-invariant, $\mu[t] > 0$ for each $t \in A = \{d, h, v\}$. Let $C = [c_0 \dots c_n]$ be a cylinder set determined by a string $c_0 \dots c_n$ on the alphabet A . For a finite or infinite string w on A and $t \in A$, let $\sigma_t w = tw$. Denote $C^t = [c_0 \dots c_n t]$, and note that $\sigma_t C$ and C^t end at the same vertex. For a vertex $y = (n, k)$, let $y^h = (n-1, k)$, $y^v = (n, k-1)$, and $y^d = (n-1, k-1)$.

First we show that the given ergodic measure μ is shift invariant. Let $x \in X$ be an infinite path labeled by a string $(\omega_i) \in A^{\mathbb{Z}^+}$. Denote by $x_i = (n_i, k_i)$ the terminal vertex of the path $\omega_0 \dots \omega_i$. The set of paths from the terminal vertex of C to x_i breaks into three disjoint sets, depending on whether the last entry of the string labeling the path is d, h , or v . Thus

$$(2.13) \quad \dim(C, x_i) = \dim(C, x_i^d) + \dim(C, x_i^h) + \dim(C, x_i^v).$$

By isotropy of the graph, for each $t \in A$,

$$(2.14) \quad \dim(C, x_i^t) = \dim(\sigma_t C, x_i);$$

and

$$(2.15) \quad \sigma^{-1}C = \sigma_d C \cup \sigma_h C \cup \sigma_v C \quad (\text{disjoint union}).$$

Thus dividing the equation

$$(2.16) \quad \dim(C, x_i) = \dim(\sigma_d C, x_i) + \dim(\sigma_h C, x_i) + \dim(\sigma_v C, x_i)$$

through by $\dim(x_i)$ and letting $i \rightarrow \infty$, if x is generic for μ , we conclude that $\mu(C) = \mu(\sigma^{-1}C)$.

Now we will show that $\mu(C^t)/\mu(C)$ is independent of the cylinder set C for each $t \in A$, thereby concluding the proof. First, we show that

$$(2.17) \quad \frac{\mu(C^t)}{\mu(C)} = \frac{\mu(C^{st})}{\mu(C^s)} \quad \text{for all } s, t \in A.$$

Note that by isotropy (it doesn't matter where the extra step is inserted, we still arrive at the same terminal vertex),

$$(2.18) \quad \dim(C^t, x_i) = \dim(C^{st}, (\sigma_s x)_{i+1}), \quad \dim(C, x_i) = \dim(C^s, (\sigma_s x)_{i+1}).$$

Dividing, we have

$$(2.19) \quad \frac{\dim(C^t, x_i)}{\dim(C, x_i)} = \frac{\dim(C^{st}, (\sigma_s x)_{i+1})}{\dim(C^s, (\sigma_s x)_{i+1})}.$$

Divide top and bottom on the left by $\dim(x_i)$, on the right by $\dim((\sigma_s x)_{i+1})$, and let $i \rightarrow \infty$. As in [11], if E is the set of μ -measure 1 of points x that give the correct values for the limits on the left-hand side, then $\mu[s] > 0$ implies that $\mu(\sigma_s E) > 0$, and hence $\mu(\sigma_s E \cap E) > 0$. Thus using any $x \in \sigma_s E \cap E$ yields Formula (2.17).

We can then compute that

$$(2.20) \quad \frac{\mu(C^t)}{\mu(C)} = \frac{\mu[c_0 \dots c_{n-1}]^{c_n t}}{\mu[c_0 \dots c_{n-1}]^{c_n}} = \frac{\mu[c_0 \dots c_{n-1}]^t}{\mu[c_0 \dots c_{n-1}]} = \dots = \frac{\mu[c_0]^t}{\mu[c_0]}.$$

By adic invariance of μ , for $r, s, t \in A$,

$$(2.21) \quad \mu[st] = \mu[ts] \quad \text{and} \quad \mu[rst] = \mu[srt],$$

since the paths defining the cylinders end at the same vertex in each case. Then

$$(2.22) \quad \frac{\mu[s]^t}{\mu[s]} = \frac{\mu[s]^{rt}}{\mu[s]^r} = \frac{\mu[srt]}{\mu[sr]} = \frac{\mu[rst]}{\mu[rs]} = \frac{\mu[r]^{st}}{\mu[r]^s} = \frac{\mu[r]^t}{\mu[r]},$$

so that $\mu[c_0]^t/\mu[c_0]$ is independent of c_0 . \square

Remark 2.3. Suppose that μ_α is a measure on the Delannoy system as described in the statement of Theorem 2.2. Suppose that C is a cylinder set determined by a path segment from $(0, 0)$ to a regular vertex (n_0, k_0) . Because μ_α is Bernoulli, hence invariant and ergodic for the shift on the space of infinite one-sided sequences on the alphabet of three symbols $\{d, h, v\}$ (for diagonal, horizontal, or vertical steps taken by the random walker), for μ_α -a.e. infinite path $x = (n_i, k_i)$ starting at $(0, 0)$ in the Delannoy graph, we have

$$(2.23) \quad k_i/n_i \rightarrow \rho = \frac{\alpha + \gamma}{\alpha + \beta}.$$

From the ergodicity of μ_α , which we have proved above, it follows that for a.e. path (n_i, k_i) ,

$$(2.24) \quad \frac{\dim((n_0, k_0), (n_i, k_i))}{\dim((0, 0), (n_i, k_i))} \rightarrow (\sqrt{1 + \rho^2} + \rho)^{-n_0} \left(\frac{\sqrt{1 + \rho^2} + 1}{\rho} \right)^{-k_0} \quad \text{as } i \rightarrow \infty.$$

It is not clear that the Pemantle-Wilson asymptotics (2.2) yield this conclusion for all paths with $k_i/n_i \rightarrow \rho$. Similarly, it is not clear that the asymptotics would allow one to calculate the measure of any cylinder extended by a single edge and conclude that every ergodic measure is one of the measures μ_α .

3. TOTAL ERGODICITY

In this section we will show that with respect to each of its ergodic measures the Delannoy adic system is *totally ergodic*, i.e. has no proper roots of unity among its L^2 eigenvalues. The proof depends on congruence properties of the Delannoy numbers with respect to primes. If λ is an eigenvalue of T , then approximating an eigenfunction by linear combinations of characteristic functions of cylinders, many of whose points return after $\dim(R, x_n)$ iterates, shows that

$$(3.1) \quad \lambda^{D(n_i, k_i)} \rightarrow 1 \quad \text{for a.e. path } x = ((n_i, k_i)).$$

If $\lambda = e^{2\pi im/n}$ with m, n relatively prime, if p is a prime that divides n , and if a.e. path $x = ((n_i, k_i))$ includes infinitely many vertices for which $D(n_i, k_i)$ is not divisible by p , then it is impossible for the convergence in (3.1) to hold—see [14]. We will find “blocking sets”: configurations of vertices with $D(n, k)$ not divisible by p that must be hit infinitely many times by each infinite path in the diagram.

A basic ingredient in the following argument is the well-known formula of Lucas [10]: If p is a prime and $n = n_0 + n_1p + n_2p^2 + \dots$, $k = k_0 + k_1p + k_2p^2 + \dots$, are the base p expansions of n and k (each $n_i, k_i \in \{0, 1, \dots, p-1\}$) then

$$(3.2) \quad \binom{n}{k} \equiv_p \binom{n_0}{k_0} \binom{n_1}{k_1} \cdots$$

Lemma 3.1. *For p prime, $r \geq 0$, and $j = 0, 1, \dots, p^r - 1$,*

$$(3.3) \quad C(p^r - 1, j) = \binom{p^r - 1}{j} \equiv_p (-1)^j.$$

Proof. $C(p^r - 1, j) \equiv_p 1$ at $j = 0$ and $j = p^r - 1$, and $C(p^r - 1, j) + C(p^r - 1, j + 1) \equiv_p 0$ for $j = 0, 1, \dots, p^r - 2$ since, by Lucas' formula, $C(p^r, j) \equiv_p 0$ for $1 < j < p^r$.

Alternatively,

$$(3.4) \quad (i + 1)C(p - 1, i + 1) = (p - (i + 1))C(p - 1, i) \equiv_p -(i + 1)C(p - 1, i) \text{ for } i < p - 1,$$

so

$$(3.5) \quad C(p - 1, i + 1) \equiv_p -C(p - 1, i) \text{ if } i < p - 1.$$

Thus, again by Lucas' formula, increasing any entry i of the expansion of j base p multiplies $C(p^r - 1, j) \pmod p$ by -1 . And if the entries of the expansion of j base p are $(p - 1, p - 1, \dots, 0, j_s)$ with $j_s < p - 1$, then the entries of $j + 1$ base p are $(0, 0, \dots, 0, j_s + 1)$, and $C(p - 1, p - 1) = C(p - 1, 0)$, so again the sign changes when we pass from j to $j + 1$. \square

Lemma 3.2. *For p prime, $r \geq 0$, $j = 0, 1, \dots, p^r - 1$, and $i = 0, 1, \dots$,*

$$(3.6) \quad \binom{j + ip^r}{p^r - 1} \equiv_p \binom{j}{p^r - 1}.$$

Proof. Let $k = p^r - 1 = (p - 1) + (p - 1)p = \dots + (p - 1)p^{r-1}$ and $j = j_0 + j_1p + \dots + j_{r-1}p^{r-1} + j_r p^r + \dots$, with all $j_s \in \{0, 1, \dots, p-1\}$. (So the base p entries of k are $k_m = p - 1$ if $m = 0, 1, \dots, r - 1$, $k_m = 0$ for $m > r - 1$.)

Then for $i = 0, 1, \dots$, we have

$$(3.7) \quad j + ip^r = j_0 + j_1p + \dots + j_{r-1}p^{r-1} + \dots$$

When we pass from j to $j + ip^r$, in the base p expansion only entries j_m for $m \geq r$ can change, say to j'_m . But for $m \geq r$, $C(j'_m, 0) = C(j_m, 0) = 1$. Thus $C(j_m, k_m) = C(j'_m, k_m)$ for all m , and hence, by Lucas' formula, $C(j + ip^r, k) \equiv_p C(j, k)$. \square

Lemma 3.3. *For p prime, $r \geq 0$, and $j = 1, \dots, p^r - 1$,*

$$(3.8) \quad \binom{p^r - 1 + j}{p^r - 1} \equiv_p 0.$$

Proof. Let p be prime, $k = p^r - 1$, $j \in \{0, 1, \dots, p^r - 1\}$. Let k_m, j_m be the entries in the base p expansions of k and j , respectively, as in the proof of Lemma 3.2, so that some (first) $j_m > 0$.

Then $p^r + j = j_0 + j_1p + \dots + j_{r-1}p^{r-1} + p^r$, and when we form $p^r + j - 1$, we decrease the first nonzero j_m (which is in $\{1, \dots, p - 1\}$), so that the base p expansion of $p^r + j - 1$ has an entry $j'_i = j_i - 1 < p - 1$, whence $C(j'_i, p - 1) = 0$. \square

Theorem 3.4. *For p prime, $r \geq 0$, and $n = 0, 1, 2, \dots$,*

$$(3.9) \quad D(n, p^r - 1) \equiv_p (-1)^{(n \bmod p^r)}.$$

Proof. Let $k = p^r - 1$ and $m = (n \bmod p^r) \in \{0, 1, \dots, p^r - 1\}$. Using Lemma 3.2,

$$(3.10) \quad D(n, k) = \sum_{d=0}^k C(k, d)C(n + d, k) \equiv_p \sum_{d=0}^k C(k, d)C(m + d, k) = \sum_{d=0}^k C(k, d)C(m + k - d, k).$$

By Lemma 3.3, the only nonzero term mod p is at $m - d = 0$. Thus, if $n = \sum m_i p^i$ with each $m_i \in \{0, 1, \dots, p - i\}$, the sum reduces to

$$(3.11) \quad C(k, m) \equiv_p C(p - 1, m_0) \cdots C(p - 1, m_{r-1}) \neq 0.$$

In fact, by Lemma 3.1, $C(k, m) \equiv_p \pm 1$. \square

Theorem 3.5. *With respect to each of its ergodic (invariant probability) measures, the Delannoy adic dynamical system is totally ergodic (i.e., has among its eigenvalues no roots of unity besides 1).*


Proof. As in the proof of total ergodicity of the Pascal adic (see, for example, [1, 14, 15]), it is enough to show that for each prime p , every infinite path in the diagram must pass through infinitely many vertices whose dimensions are not divisible by p . The vertices $\{(n, p^r - 1) : n, r \geq 0\} \cup \{(p^r - 1, k) : k, r \geq 0\}$, form such a “blocking set”—see Figure 6. \square

4. DIMENSION GROUPS

We want to compute the dimension group of the Delannoy Bratteli diagram shown in Figure 5. The adjacency matrices A_i between levels i and $i + 1$, for $i \geq 0$, determine the sequence of ordered abelian group homomorphisms

$$(4.1) \quad \mathbb{Z} \xrightarrow{A_1} \mathbb{Z}^3 \xrightarrow{A_2} \mathbb{Z}^5 \xrightarrow{A_3} \mathbb{Z}^7 \xrightarrow{A_4} \dots$$

Writing as usual transposes of the vectors and matrices involved for ease of typing,



those equivalence classes which contain a nonnegative vector, and a distinguished order unit, the equivalence class of $1 \in \mathbb{Z}$. Equivalence classes are added by adding representatives that have equal dimensions.

We can also give a slightly more concrete representation of this dimension group as follows. We correspond to each $v \in \mathbb{Z}^{2k+1}$ the following pair of polynomials with coefficients from \mathbb{Z} :

$$(4.3) \quad r(v) = \sum_{i=0}^{k-1} v_{2i} x^i, \quad s(v) = \sum_{j=0}^k v_{2j+1} x^j.$$

Then $A_{k+1}v$ corresponds to the pair of polynomials

$$(4.4) \quad (r', s') = (s, xr(v) + (1+x)s(v)) = (r, s) \begin{pmatrix} 0 & x \\ 1 & 1+x \end{pmatrix}.$$

The connecting maps A_{k+1} can be realized by a single matrix

$$(4.5) \quad B = \begin{pmatrix} 0 & x \\ 1 & 1+x \end{pmatrix}, \quad \text{with inverse } B^{-1} = -\frac{1}{x} \begin{pmatrix} 1+x & -x \\ -1 & 0 \end{pmatrix}.$$

Thus the discrete union of $\{\mathbb{Z}^{2k+1} : k = 0, 1, 2, \dots\}$ maps onto the set of pairs of polynomials (r, s) with integer coefficients and $\deg(r) < \deg(s)$. Two pairs (r, s) and (u, v) are equivalent if there is $n \in \mathbb{Z}$ such that $(u, v) = B^n(r, s)$. Each nonzero equivalence class contains a pair (r, s) of smallest degree, which satisfies $r(0) \neq s(0)$. It can be found by starting with any member of the class and applying powers of B^{-1} repeatedly. Equivalence classes can be added by adding members of equal degrees. Unfortunately the map that takes $v \in \cup\{\mathbb{Z}^{2k+1} : k = 0, 1, 2, \dots\}$ to the pair of polynomials $(r(v), s(v))$ is many-to-one: for example, $(-1, 1, 0)$ and $(-1, 1, 0, 0, 0)$ have the same image. (The map could be made one-to-one by sending $v \in \mathbb{Z}^{2k+1}$ to the triple $(r(v), s(v), k)$.) So the set of equivalence classes of pairs of integer-coefficient polynomials is a quotient of the dimension group of the Delannoy adic. The distinguished order unit is the class of the image of $1 \in \mathbb{Z}$, namely the class of the pair of polynomials $r(x) = 1, s(x) = x + 1$. This class consists of the ‘‘Delannoy polynomials’’ $P_n(x)$ defined by $P_0(x) = 1, P_1(x) = x + 1, P_{n+1}(x) = (x + 1)P_n(x) + xP_{n-1}(x)$, whose coefficients form the array of Delannoy numbers.

5. REMARKS

(1) Recall that $A = \{h, d, v\}$ denotes the set of edge labels for paths in the Delannoy graph. Denote by $\sigma : A^{\mathbb{Z}} \rightarrow A^{\mathbb{Z}}$ the usual shift map $(\sigma\omega)_i = (\omega)_{i+1}, i \in \mathbb{Z}$. For the Delannoy system there is a countable set of paths $X' \subset X$ such that the map $\phi : X \setminus X' \rightarrow A^{\mathbb{N}}$ defined by $(\phi(x))_i = e_1(T^i x)$ is one-to-one (and Borel measurable) and satisfies $\phi \circ T = \sigma \circ \phi$. Thus for each of its invariant measures μ , the Delannoy system is isomorphic to a genuine topological dynamical system (a compact metric space together with a homeomorphism) with a shift-invariant Borel probability measure, namely the subshift Σ which is the closure of the image of ϕ together with σ and the image of μ under ϕ . The proof is the same as in [12].

(2) The just-mentioned subshift (Σ, σ) is topologically weakly mixing. The proof is the same as in [12].

(3) With each of its ergodic invariant measures, the Delannoy adic dynamical system is loosely Bernoulli. The proof is the same as in [8] (see also [9]).

(4) It may be possible to determine for (Σ, σ) limit laws for return times to (or hitting times of) cylinder sets and the complexity of the associated formal language, as in [12]. We do not know, for non-atomic ergodic measures, about weak mixing, multiplicity of the spectrum, or joinings of any μ_α with any μ_β .

(5) The Delannoy adic is another example, after the Pascal, of a system that is *isotropic* in the sense that there is the same pattern of edges leaving each vertex. It seems that section (3) of the proof of Theorem 2.2 (X. Méla’s isotropy argument) will extend to all such systems to identify their ergodic measures. These systems will be the subject of future work.

REFERENCES

- [1] Terrence M. Adams and Karl E. Petersen, *Binomial-coefficient multiples of irrationals*, *Monatsh. Math.* **125** (1998), no. 4, 269–278, DOI 10.1007/BF01305342. MR1621682 (99d:28026)
- [2] Sarah Bailey, Michael Keane, Karl Petersen, and Ibrahim A. Salama, *Ergodicity of the adic transformation on the Euler graph*, *Math. Proc. Cambridge Philos. Soc.* **141** (2006), no. 2, 231–238, DOI 10.1017/S0305004106009431. MR2265871 (2007m:37010)
- [3] S. Frick and K. Petersen, *Random permutations and unique fully supported ergodicity for the Euler adic transformation*, *Ann. Inst. Henri Poincaré Probab. Stat.* **44** (2008), no. 5, 876–885 (English, with English and French summaries). MR2453848 (2009h:37004)
- [4] ———, *Reinforced random walks and adic transformations*. To appear in *J. Theor. Prob.*
- [5] Cyril Banderier and Sylviane Schwer, *Why Delannoy numbers?*, *J. Statist. Plann. Inference* **135** (2005), no. 1, 40–54, DOI 10.1016/j.jspi.2005.02.004. MR2202337
- [6] P. Billingsley, *Probability and Measure*, John Wiley and Sons, Inc., 1995.
- [7] Norman Carey and David Clampitt, *Regions: A Theory of Tonal Spaces in Early Medieval Treatises*, *Journal of Music Theory* **40** (Spring, 1996), no. 1, 113–147.
- [8] Élise Janvresse and Thierry de la Rue, *The Pascal adic transformation is loosely Bernoulli*, *Ann. Inst. H. Poincaré Probab. Statist.* **40** (2004), no. 2, 133–139, DOI 10.1016/S0246-0203(03)00062-1 (English, with English and French summaries). MR2044811 (2004k:37006)
- [9] Sarah Bailey Frick, *Limited scope adic transformations*, *Discrete Contin. Dyn. Syst. Ser. S* **2** (2009), no. 2, 269–285, DOI 10.3934/dcdss.2009.2.269. MR2505638 (2010h:37001)
- [10] Edouard Lucas, *Théorie des Fonctions Numériques Simplement Périodiques*, *Amer. J. Math.* **1** (1878), no. 4, 289–321, DOI 10.2307/2369373 (French). MR1505176
- [11] Xavier Méla, *A class of nonstationary adic transformations*, *Ann. Inst. H. Poincaré Probab. Statist.* **42** (2006), no. 1, 103–123, DOI 10.1016/j.anihpb.2005.02.002 (English, with English and French summaries). MR2196974 (2006j:37003)
- [12] Xavier Méla and Karl Petersen, *Dynamical properties of the Pascal adic transformation*, *Ergodic Theory Dynam. Systems* **25** (2005), no. 1, 227–256, DOI 10.1017/S0143385704000173. MR2122921 (2005k:37012)
- [13] Robin Pemantle and Mark C. Wilson, *Asymptotics of multivariate sequences. I. Smooth points of the singular variety*, *J. Combin. Theory Ser. A* **97** (2002), no. 1, 129–161, DOI 10.1006/jcta.2001.3201. MR1879131 (2003a:05015)

- [14] Karl Petersen, *Information compression and retention in dynamical processes*, Dynamics and randomness (Santiago, 2000), Nonlinear Phenom. Complex Systems, vol. 7, Kluwer Acad. Publ., Dordrecht, 2002, pp. 147–217. MR1975578 (2005d:37020)
- [15] Karl Petersen and Klaus Schmidt, *Symmetric Gibbs measures*, Trans. Amer. Math. Soc. **349** (1997), no. 7, 2775–2811, DOI 10.1090/S0002-9947-97-01934-X. MR1422906 (99a:28016)
- [16] Karl Petersen and Alexander Varchenko, *The Euler adic dynamical system and path counts in the Euler graph*, Tokyo J. Math. **33** (2010), no. 2, 327–340.
- [17] N. Sloane, *The On-Line Encyclopedia of Integer Sequences*, <http://oeis.org>, 2010.
- [18] N. Sloane and S. Plouffe, *The Encyclopedia of Integer Sequences*, Academic Press Inc., San Diego, CA, 1995. With a separately available computer disk. MR1327059 (96a:11001)
- [19] Frank Spitzer, *Principles of Random Walk*, The University Series in Higher Mathematics, D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto-London, 1964. MR0171290 (30 #1521)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL, CHAPEL HILL,
NC 27599-3250, USA

E-mail address: `petersen@math.unc.edu`