

# UNIQUE PATH PARTITIONS: CHARACTERIZATION AND CONGRUENCES

CHRISTINE BESSENRODT, JØRN B. OLSSON, AND JAMES A. SELLERS

ABSTRACT. We give a complete classification of the unique path partitions and study congruence properties of the function which enumerates such partitions.

## 1. INTRODUCTION

The famous Murnaghan-Nakayama formula gives a combinatorial rule for computing the value of the irreducible character of the symmetric groups  $S_n$  labelled by the partition  $\lambda$  on the conjugacy class labelled by a partition  $\mu$  (see [2]). This value is the weighted sum over the  $\mu$ -paths in  $\lambda$ , as defined below, where the weight is a sign corresponding to the sum of the leg lengths of the rim hooks removed along the path.

If  $\mu = (a_1, a_2, \dots, a_k)$ , with  $a_1 \geq a_2 \geq \dots \geq a_k > 0$ , and  $\lambda$  are partitions of  $n$ , then a  $\mu$ -path in  $\lambda$  is a sequence of partitions,  $\lambda = \lambda_0, \lambda_1, \dots, \lambda_k = (0)$ , where for  $i = 1, \dots, k$  the partition  $\lambda_i$  is obtained by removing an  $a_i$ -hook from  $\lambda_{i-1}$ . As in [3], we call  $\mu$  a *unique path partition for  $\lambda$*  (or *up-partition for  $\lambda$*  for short) if the number of  $\mu$ -paths in  $\lambda$  is at most 1. We call  $\mu$  a *up-partition* if it is a *up-partition* for all partitions of  $n$ .

Thus, a *up-partition*  $\mu$  labels a conjugacy class where all non-zero irreducible character values are 1 or  $-1$ , i.e., they are *sign partitions* as defined in [3]. By [6, 7.17.4], the sign partitions  $\mu$  are exactly those for which the expansion of the corresponding power sum symmetric function into Schur functions is multiplicity-free.

Note that not every sign partition is a *up-partition* as cancellation may occur. For example, the partition  $(3, 2, 1)$  is a sign partition, but not a *up-partition*, since there are two  $(3, 2, 1)$ -paths in the partition  $(3, 2, 1)$ .

In this paper, we accomplish three goals. First, we provide an explicit classification of the unique path partitions in terms of partitions we call *strongly decreasing*. We then discuss numerous connections between *up-partitions* and certain types of binary partitions. Such connections are truly beneficial; they led us to the development of a generating function for, and a recurrence satisfied by  $u(n)$ , the number of *up-partitions* of the positive integer  $n$ . Thanks to this link between

---

*Date:* September 3, 2012.

*1991 Mathematics Subject Classification.* 05A17, 11P83, 20C30.

*Key words and phrases.* binary partitions, unique path partitions, rim hooks, symmetric group, character values, congruences.

$up$ -partitions and restricted binary partitions, we were encouraged to consider the arithmetic properties of  $u(n)$ . (Such a motivation is natural based on the literature that already exists on congruence properties satisfied by binary partitions. Indeed, Churchhouse [1] initiated the study of congruence properties satisfied by the unrestricted binary partition function in the late 1960's. This work was further extended by Rødseth and Sellers [4].) We close this paper by proving a number of congruence relations satisfied by  $u(n)$  modulo powers of 2.

## 2. THE CLASSIFICATION OF $up$ -PARTITIONS

We now collect the facts necessary for classifying the  $up$ -partitions in an elegant fashion. As usual, we gather equal parts together and write  $i^m$  for  $m$  parts equal to  $i$  in a partition.

**Lemma 2.1.** (1) *If  $\mu = (a_1, a_2, \dots, a_k)$  is a  $up$ -partition with  $a_k = 2$ , then  $\mu' = (a_1, a_2, \dots, a_{k-1}, 1^2)$  is also a  $up$ -partition.*

(2) *If  $\mu = (a_1, a_2, \dots, a_k)$  is a  $up$ -partition with  $k \geq 2$ , then  $\mu_2 = (a_2, \dots, a_k)$  is also a  $up$ -partition.*

*Proof.* (1) follows immediately from the definition.

(2) If a partition  $\lambda_2$  of  $n - a_1$  has two or more  $\mu_2$ -paths then any partition of  $n$  obtained by adding an  $a_1$ -hook to  $\lambda_2$  has two or more  $\mu$ -paths.  $\square$

**Lemma 2.2.** *Let  $\mu = (a_1, a_2, \dots, a_k)$  be a partition of  $n$  and  $a > n$ . Then  $\mu$  is a  $up$ -partition if and only if  $\mu' = (a, a_1, \dots, a_k)$  is a  $up$ -partition.*

*Proof.* By Lemma 2.1(2) we only need to show that if  $\mu$  is a  $up$ -partition then also  $\mu'$  is a  $up$ -partition. Let  $\lambda'$  be a partition of  $a + n$ . Since  $a > n$ ,  $\lambda'$  cannot contain two or more  $a$ -hooks. If  $\lambda'$  contains an  $a$ -hook, we let  $\lambda$  be the partition obtained by removing it. Since by assumption  $\mu$  is a  $up$ -partition for  $\lambda$ , we get that  $\mu'$  is a  $up$ -partition for  $\lambda'$ .  $\square$

We call an extension of a partition of  $n$  by a part  $a > n$  as in Lemma 2.2 *strongly decreasing*, or for short, an *sd-extension*. A partition  $\mu$  obtained from a partition  $\rho$  by several *sd*-extensions is then called an *sd-extension of  $\rho$* ; if  $\rho = (0)$ ,  $\mu$  is called an *sd-partition*. As stated in [3], a partition  $\mu = (a_1, a_2, \dots, a_k)$  is an *sd-partition* if and only if  $a_i > a_{i+1} + \dots + a_k$  for all  $i = 1, \dots, k - 1$ .

We have the following classification result for  $up$ -partitions:

**Theorem 2.3.** *A partition  $\mu$  is a  $up$ -partition if and only if one of the following holds:*

- (i)  *$\mu$  is an *sd-partition*.*
- (ii)  *$\mu$  is an *sd-extension of  $(1^2)$ .**

*Proof.* In the proof we use the well-known connection between first column hook lengths and hook removal as described in [2, Section 2.7].

As (0) and  $(1^2)$  are  $up$ -partitions, Lemma 2.2 shows that their *sd*-extensions are  $up$ -partitions. Suppose that  $n$  is minimal such that there exists a partition

$\mu = (a_1, a_2, \dots, a_k)$  of  $n$ , which is a *up*-partition but not an *sd*-extension of  $(0)$  or  $(1^2)$ . Obviously  $k \geq 2$ .

Assume  $a_2 = 1$ , i.e.,  $\mu = (n - k + 1, 1^{k-1})$ . If  $k > 3$ , then  $\mu$  is not a *up*-partition since  $(1^{k-1})$  is not. For  $k = 3$ , only  $(2, 1^2)$  and  $(1^3)$  are not *sd*-extensions of  $(1^2)$ , but these are not *up*-partitions. For  $k = 2$ ,  $\mu$  is an *sd*-partition or  $(1^2)$ .

Thus we may now assume that  $a_2 > 1$ . We put  $\mu_i = (a_i, \dots, a_k)$  and  $n_i = |\mu_i|$  for  $i = 2, \dots, k$ . Also  $n_{k+1} := 0$ .

Now suppose that  $a_1 = a_2$ . If  $k = 2$  then  $\mu$  is not a *up*-partition for  $\lambda = (a_1, a_1)$ . If  $k > 2$  then  $\mu$  is not a *up*-partition for  $\lambda = (n - a_1, 1^{a_1})$ .

Thus we may now assume  $a_1 > a_2 > 1$ . By Lemma 2.1,  $\mu_2 = (a_2, \dots, a_k)$  is a *up*-partition, and thus, by minimality, it is an *sd*-extension of  $(0)$  or  $(1^2)$ . Then  $\mu$  cannot be an *sd*-extension of  $\mu_2$ , and hence  $a_1 \leq n_2$ .

Now  $a_1 > a_2 > n_3$  and hence  $d := a_1 - n_3 - 1 > 0$ . Note that  $n_2 = n_3 + a_2 > n_3 + 1$ , and thus  $\lambda = (n_2, n_3 + 1, 1^d)$  is a partition of  $n_2 + n_3 + 1 + d = a_1 + n_2 = n$ . The set of first column hook lengths for  $\lambda$  is  $\{a_1 + a_2, a_1, d, d - 1, \dots, 1\}$ , as is easily calculated. As  $d \leq n_2 - n_3 - 1 = a_2 - 1$ ,  $\lambda$  has two  $a_1$ -hooks. After removing the  $a_1$ -hook in the second row we get the partition  $\lambda' = (n_2)$ . After removing the  $a_1$ -hook in the first row we get  $\{a_1, a_2, d, d - 1, \dots, 1\}$  as a set of a first column hook lengths for a partition  $\lambda''$ . Now  $\lambda''$  has an  $a_2$ -hook in the second row. Removing it we obtain the partition  $(n_3)$ . This shows that  $\mu$  is not a *up*-partition for  $\lambda$ , giving a contradiction.  $\square$

### 3. ON *up*-PARTITIONS AND RESTRICTED BINARY PARTITIONS

For each  $n \in \mathbb{N}$ , we denote the number of *up*-partitions of  $n$  by  $u(n)$ . For  $t \in \mathbb{N}$ , we define an *sd<sub>t</sub>*-partition to be an *sd*-extension of the partition  $(t)$ . The following lemma is obvious.

**Lemma 3.1.** *Let  $\mu$  be a partition of  $t$ . There is a bijection between *sd*-extensions of  $\mu$  and *sd<sub>t</sub>*-partitions obtained by replacing all the parts of  $\mu$  by one part  $t$ .  $\square$*

We denote the number of *sd*-partitions of  $n$  by  $s(n)$  and the number of *sd<sub>t</sub>*-partitions of  $n$  by  $s_t(n)$  so that  $s(n) = \sum_{t \geq 1} s_t(n)$ . Combining Theorem 2.3 with Lemma 3.1 we get the following:

**Corollary 3.2.** *For each  $n \geq 1$ ,*

$$u(n) = s(n) + s_2(n). \quad \square$$

Next, we focus our attention on  $s(n)$ .

**Proposition 3.3.** *For each  $n \geq 2$ ,*

$$s(n) = 2s_1(n) + s_2(n).$$

*Proof.* Let  $\lambda = (a_1, a_2, \dots, a_k)$  be an *sd<sub>t</sub>*-partition, i.e.,  $a_k = t$ . If we map  $\lambda$  onto  $(a_1, a_2, \dots, a_k - 1, 1)$  we get a bijection between the set of all *sd<sub>t</sub>*-partitions of  $n$  with  $t \geq 3$  and the set of all *sd<sub>1</sub>*-partitions of  $n$ . Thus  $s_1(n) = \sum_{t \geq 3} s_t(n)$ . The result follows, since  $s(n) = \sum_{t \geq 1} s_t(n)$ .  $\square$

Combining Corollary 3.2 and Proposition 3.3, we have the following:

**Theorem 3.4.** *For each  $n \geq 2$ ,  $u(n)$  is even. In fact,*

$$\frac{u(n)}{2} = s_1(n) + s_2(n). \quad \square$$

Thanks to their definition, it is clear that  $sd$ -partitions are closely related to non-squashing partitions and binary partitions as described in [5]. A partition  $\lambda = (a_1, a_2, \dots, a_k)$  is called *non-squashing* if  $a_i \geq a_{i+1} + \dots + a_k$  for  $1 \leq i \leq k-1$  and *binary* if all parts  $a_i$  are powers of 2. The difference between  $sd$ - and non-squashing partitions is whether or not the inequalities between  $a_i$  and  $a_{i+1} + \dots + a_k$  are strict. A binary partition is called *restricted* (for short, an *rb-partition*) if it satisfies the following condition: Whenever  $2^i$  is a part and  $i \geq 1$  then  $2^{i-1}$  is also a part. For  $t \in \mathbb{N}$ , an *rb<sub>t</sub>-partition* is an *rb-partition* where the largest part occurs with multiplicity  $t$ .

With this in mind, we can naturally connect the  $sd_t$ -partitions and the  $rb_t$ -partitions.

**Theorem 3.5.** *Let  $n, t \in \mathbb{N}$ . There is a bijection between the set of  $sd_t$ -partitions of  $n$  and the set of  $rb_t$ -partitions of  $n$ .*

*Proof.* Clearly, an  $sd$ -partition  $\lambda = (a_1, a_2, \dots, a_k)$  of  $n$  is uniquely determined by the positive integers  $d_i \in \mathbb{N}$ ,  $i = 1, \dots, k$ , defined by  $d_i = a_i - (a_{i+1} + \dots + a_k)$  for  $i = 1, \dots, k-1$ , and  $d_k = a_k$ . An easy calculation shows that with this notation  $n = d_1 + d_2 2 + \dots + d_k 2^{k-1}$ . Thus if we map  $\lambda$  onto the binary partition where  $2^j$  occurs with multiplicity  $d_{j+1}$ ,  $j = 0, 1, \dots, k-1$ , we get the desired bijection.  $\square$

**Remark 3.6.** Theorem 3.5 shows that  $s(n)$  equals the number of  $rb$ -partitions of  $n$ . Let  $S(q) := \sum_{n \geq 1} s(n)q^n$  be the generating function for  $s(n)$ . It is easy to write down the generating function for the number of  $rb$ -partitions which implies that

$$S(q) = \sum_{i \geq 1} q^{2^i - 1} \prod_{j=0}^{i-1} \frac{1}{1 - q^{2^j}}.$$

From its definition, one also gets the identity

$$S(q)(1 - q) = q(1 + S(q^2)).$$

Moreover, the generating function  $S_t(q)$  for the number of  $rb_t$ -partitions is given by

$$S_t(q) = \sum_{i \geq 1} q^{2^i - 1 + (t-1)2^{i-1}} \prod_{j=0}^{i-2} \frac{1}{1 - q^{2^j}},$$

and it satisfies the identity

$$(S_t(q) - q^t)(1 - q) = qS_t(q^2).$$

Hence, by Theorem 3.4, the generating function  $U(q)$  for the number of  $up$ -partitions is then

$$U(q) = 2(S_1(q) + S_2(q)).$$

We now exploit this connection between  $rb$ -partitions and  $sd$ -partitions to prove a number of facts about  $s(n)$  and related functions. The following results may alternatively also be proved by using the identities for the generating functions  $S(q)$  and  $S_t(q)$  stated above.

**Proposition 3.7.** *For each  $r \in \mathbb{N}$  we have*

$$\begin{aligned} s(2r) &= s(2r-1) \\ s(2r+1) &= s(2r) + s(r). \end{aligned}$$

*Proof.* An  $rb$ -partition must contain a part 1. Removing such a part from an  $rb$ -partition  $\lambda$  of  $2r$  gives an  $rb$ -partition  $\lambda'$  of  $2r-1$ . (A binary partition of an odd number must contain 1 as a part, so that  $\lambda'$  is still  $rb$ .) This map is then in fact a bijection between  $rb$ -partitions of  $2r$  and those of  $2r-1$ .

Removing a part 1 from an  $rb$ -partition  $\lambda$  of  $2r+1$  gives a binary partition  $\lambda'$  of  $2r$ . If  $\lambda'$  has a part equal to 1, it is an  $rb$ -partition and we put  $\lambda'' = \lambda'$ . Otherwise all parts of  $\lambda'$  are even and we may divide them all by 2 to get an  $rb$ -partition  $\lambda''$  of  $r$ . The process of going from  $\lambda$  to  $\lambda''$  may obviously be reversed. Thus  $s(2r+1) = s(2r) + s(r)$ .  $\square$

With Proposition 3.7 in mind, we define  $s^*(r) := s(2r) (= s(2r-1))$  for  $r \in \mathbb{N}$ .

**Proposition 3.8.** *We have  $s^*(1) = 1$  and*

$$s^*(r) = s^*(r-1) + s^*\left(\left\lfloor \frac{r}{2} \right\rfloor\right), \text{ for } r \geq 2.$$

*Proof.* Clearly  $s^*(1) = s(1) = 1$ . We prove the proposition by showing that for  $r' \in \mathbb{N}$  we have

$$s^*(2r') = s^*(2r'-1) + s^*(r') \text{ and } s^*(2r'+1) = s^*(2r') + s^*(r').$$

The equations are by definition of  $s^*$  equivalent to

$$s(4r') = s(4r'-2) + s(2r') \text{ and } s(4r'+2) = s(4r') + s(2r').$$

But these are easily deduced from Proposition 3.7.  $\square$

**Remark 3.9.** Proposition 3.8 proves that the sequence  $s^*(n)$  is listed in [7] as A033485 and thus that the sequence  $s(n)$  is listed as A040039. In particular, the comment by John McKay which appears in A40039 in [7] is confirmed.

We proceed to consider the numbers  $s_t(r)$  of  $rb_t$ -partitions.

**Proposition 3.10.** *Let  $t \in \mathbb{N}$ . We have  $s_t(1) = s_t(2) = \dots = s_t(t-1) = 0$ ,  $s_t(t) = 1$ ,  $s_t(t+1) = \dots = s_t(2t) = 0$ , and  $s_t(2t+1) = 1$ . Also,  $s_t(2r) = s_t(2r-1)$  whenever  $t \neq 2r, 2r-1$ , i.e., whenever  $r \neq \lfloor \frac{t+1}{2} \rfloor$ .*

*Proof.* The statements about  $s_t(j)$  for  $j \leq 2t+1$  are trivial. The final statement is proved in analogy with Proposition 3.7. Using the notation of that proof we have the following: If we assume that  $\lambda$  is  $rb_t$  then also  $\lambda'$  is  $rb_t$  with the exception of the case where  $\lambda = (1^t)$ . Also, if  $\lambda'$  is  $rb_t$  then  $\lambda$  is  $rb_t$  with the exception of the case where  $\lambda' = (1^t)$ . Thus we have  $s_t(2r) = s_t(2r-1)$  except when  $t \in \{2r, 2r-1\}$ .  $\square$

**Corollary 3.11.** *We have  $u(1) = 1, u(2) = 2$ , and for  $r \geq 2, u(2r) = u(2r-1)$ .  $\square$*

We now define

$$s_t^*(r) := \begin{cases} s_t(2r) & \text{if } r \text{ is odd} \\ s_t(2r-1) & \text{if } r \text{ is even} \end{cases}.$$

Proposition 3.10 shows that for all  $r \neq \lfloor \frac{t+1}{2} \rfloor$  we have  $s_t^*(r) = s_t(2r-1) = s_t(2r)$ . Also  $s_t^*(r) = 0$  for  $1 \leq r \leq t, r \neq \lfloor \frac{t+1}{2} \rfloor$  and  $s_t^*(t+1) = 1$ .

In analogy with Proposition 3.8 we have:

**Proposition 3.12.** *For all  $r \geq t+2, s_t^*(r) = s_t^*(r-1) + s_t^*(\lfloor \frac{r}{2} \rfloor)$ .*

*Proof.* The assumption on  $r$  and  $t$  implies that the partitions  $(1^r)$  and  $(1^{r-1})$  are not  $rb_t$ . Therefore the bijections in the proof of Proposition 3.7 work for  $rb_t$ -partitions of  $r$  as well. Thus we have the recursions

$$\begin{aligned} s_t(2r) &= s_t(2r-1) \\ s_t(2r+1) &= s_t(2r) + s_t(r). \end{aligned}$$

In the case  $t = 1, r = 3$  we have  $s_1^*(3) = s_1(6) = 1$  and  $s_1^*(2) + s_1^*(1) = s_1(3) + s_1(2) = 1 + 0 = 1$ . We assume  $r \geq 4$  and write  $r = 4s + i, s \in \mathbb{N}, i \in \{0, 1, 2, 3\}$ . We show

$$\begin{aligned} s_t^*(4s) &= s_t^*(4s-1) + s_t^*(2s) \\ s_t^*(4s+1) &= s_t^*(4s) + s_t^*(2s) \\ s_t^*(4s+2) &= s_t^*(4s+1) + s_t^*(2s+1) \\ s_t^*(4s+3) &= s_t^*(4s+2) + s_t^*(2s+1) \end{aligned}$$

By definition of  $s_t^*$  the equations are equivalent to

$$\begin{aligned} s_t(8s-1) &= s_t(8s-2) + s_t(4s-1) \\ s_t(8s+2) &= s_t(8s-1) + s_t(4s-1) \\ s_t(8s+3) &= s_t(8s+2) + s_t(4s+2) \\ s_t(8s+6) &= s_t(8s+4) + s_t(4s+2). \end{aligned}$$

These follow from the recursions for  $s_t$ .  $\square$

Lastly, we define  $w(n) = \frac{u(2n)}{2}$ . Theorem 3.4 and Proposition 3.12 yield the following:

**Proposition 3.13.** *For each  $n \geq 1, w(n) = s_1^*(n) + s_2^*(n)$ . Moreover, for  $n \geq 3, w(n) = w(n-1) + w(\lfloor \frac{n}{2} \rfloor)$ , and  $w(1) = w(2) = 1$ .  $\square$*

**Remark 3.14.** Proposition 3.13 shows that the sequence of numbers  $w(n)$  is listed in [7] as A075535. The simple recurrence relation is used in the next section to prove congruence results for the numbers  $w(n)$  and thus for the numbers  $u(n)$  of unique path partitions.

**Remark 3.15.** We may consider also  $w_2(n) := s_3^*(n) + s_4^*(n)$ . Then we have  $w_2(1) = 0, w_2(2) = 1, w_2(3) = 0, w_2(4) = 1$ , and for  $n \geq 5$

$$w_2(n) = w_2(n-1) + w_2(\lfloor \frac{n}{2} \rfloor).$$

Similar recurrence relations are more generally valid for

$$w_r(n) := s_{2r-1}^*(n) + s_{2r}^*(n)$$

which starts by  $w_r(1) = \dots = w_r(r-1) = 0, w_r(r) = 1, w_r(r+1) = \dots = w_r(2r-1) = 0, w_r(2r) = 1$ . This is an infinite family of sequences which may satisfy congruence relations similar to those satisfied by  $w_1(n) = w(n)$ .

In the next section we discuss congruences for  $w(n)$  and in part also for the  $w_i(n)$ 's.

#### 4. CONGRUENCES FOR THE NUMBER OF $up$ -PARTITIONS

In this section we investigate arithmetical properties of  $u(n)$ , the number of  $up$ -partitions of  $n$ . Since  $w(n) = \frac{u(2n)}{2}$ , any result on the  $w$ -sequences may be translated into a result on the  $u$ -sequence. In particular, as studying congruences of the  $u$ -sequence modulo  $2m$  is equivalent to studying the  $w$ -sequence modulo  $m$ , we will concentrate on the latter sequence.

At the start, we consider a more general situation that also covers the more general sequences defined in Remark 3.15; however, in the remaining part of this section we restrict our attention to the numbers  $w(n)$ .

**Proposition 4.1.** *Let  $(a(n))_{n \in \mathbb{N}}$  be a sequence with  $a(c), a(2c)$  odd for some  $c \in \mathbb{N}$ ,  $a(m)$  even when  $c < m < 2c$ , and  $a(n) = a(n-1) + a(\lfloor \frac{n}{2} \rfloor)$  for  $n \geq 2c$ . Then for  $n \geq c$ ,  $a(n)$  is odd exactly when  $n$  is of the form  $2^d c$ .*

*Proof.* Certainly the assertion is true for  $n = c$  and  $n = 2c$ . Assume the result holds up to some number  $n = 2^r c$ ,  $r \geq 1$ . Then

$$a(n+1) = a(n) + a(\lfloor \frac{n}{2} \rfloor) = a(2^r c) + a(2^{r-1} c) \equiv 0 \pmod{2}.$$

For any  $k$  with  $2 \leq k \leq 2^r c - 1$ , we then get by induction on  $k$  that

$$a(n+k) = a(n+k-1) + a(\lfloor \frac{n+k}{2} \rfloor) \equiv 0 \pmod{2}$$

since  $2^{r-1} c < \lfloor \frac{n+k}{2} \rfloor < 2^r c$ . For  $k = 2^r c$  we then obtain

$$a(2^{r+1} c) = a(2^{r+1} c - 1) + a(2^r c) \equiv 1 \pmod{2}.$$

Hence the assertion is proved.  $\square$

**Corollary 4.2.** *Let  $(a(n))_{n \in \mathbb{N}}$  be as in Proposition 4.1. Let  $m$  be an odd number such that  $2^b c + 1 < m \leq 2^{b+1} c - 1$  for some  $b$ . Then  $a(m) \equiv a(m-2) \pmod{4}$ . In particular,  $a(m) \equiv a(2^b c + 1) \pmod{4}$ .*

*Proof.* Since  $m$  is odd, we have

$$a(m) = a(m-1) + a(\lfloor \frac{m}{2} \rfloor) = a(m-2) + 2 a(\lfloor \frac{m}{2} \rfloor).$$

As  $m-1$  is not of the form  $2^d c$ ,  $\lfloor \frac{m}{2} \rfloor > 2^{b-1} c$  is not either. Hence,  $a(\lfloor \frac{m}{2} \rfloor)$  is even, and then the claim follows.  $\square$

Since  $w(1) = w(2) = 1$ , the following is immediate, and it gives corresponding congruences modulo 4 and 8 for  $u(n)$ :

**Corollary 4.3.** *For  $n \geq 1$ ,  $w(n)$  is even exactly when  $n$  is not a 2-power.*

*For any odd number  $m$  such that  $2^b + 1 \leq m \leq 2^{b+1} - 1$ ,*

$$w(m) \equiv w(2^b + 1) \pmod{4}.$$

□

Note that the first part of Corollary 4.3 implies infinitely many Ramanujan-like congruences modulo 4 satisfied by  $u(n)$ . To further understand the congruences of  $u(n) \pmod{8}$ , we first focus on the 2-powers. Set  $v(k) = w(2^k)$  for  $k \in \mathbb{N}_0$ .

**Proposition 4.4.** *For each  $k \geq 2$ ,*

$$v(k) \equiv 2v(k-1) + v(k-2) \pmod{4}.$$

*Proof.* Using Corollary 4.3, we have the following congruences mod 4:

$$\begin{aligned} v(k) &= w(2^k) = w(2^{k-1}) + w(2^k - 1) \equiv w(2^{k-1}) + w(2^{k-1} + 1) \\ &\equiv 2w(2^{k-1}) + w(2^{k-2}) = 2v(k-1) + v(k-2). \end{aligned}$$

□

**Proposition 4.5.** *For each  $k \geq 1$ ,*

$$v(k) = w(2^k) \equiv \begin{cases} k & \pmod{8} \text{ if } k \text{ is odd} \\ k+1 & \pmod{8} \text{ if } k \text{ is even} \end{cases}.$$

*Equivalently,*

$$v(k) \equiv 2 \left\lfloor \frac{k}{2} \right\rfloor + 1 \pmod{8}.$$

*Proof.* From the recursion formula we have

$$\begin{aligned} w(2^k) &= w(2^{k-1}) + w(2^k - 1) = w(2^{k-1}) + w(2^{k-1} - 1) + w(2^k - 2) \\ &= w(2^{k-1}) + 2w(2^{k-1} - 1) + w(2^k - 3) \\ &\quad \vdots \\ &= w(2^{k-1}) + 2w(2^{k-1} - 1) + \dots + 2w(2^{k-2} + 1) + w(2^{k-1} + 1) \\ &= 2w(2^{k-1}) + 2w(2^{k-1} - 1) + \dots + 2w(2^{k-2} + 1) + w(2^{k-2}) \end{aligned}$$

and we now investigate sums of the form  $\sum_{i=2^d+1}^{2^{d+1}} w(i)$ , for  $d \geq 1$ . We want to show

by induction that they are always congruent to 5 mod 8; for  $d = 1$ ,  $w(3) + w(4) = 2 + 3 = 5$ , so the claim holds. Now we have for any  $d \geq 2$  (using induction and the corollary):

$$\begin{aligned} \sum_{i=2^d+1}^{2^{d+1}} w(i) &= \sum_{i=2^{d-1}+1}^{2^d} w(2i) + \sum_{i=2^{d-1}+1}^{2^d} w(2i-1) \\ &= \sum_{i=2^{d-1}+1}^{2^d} w(i) + 2 \sum_{i=2^{d-1}+1}^{2^d} w(2i-1) \\ &\equiv 5 + 2^d w(2^d + 1) \pmod{8} \\ &\equiv 5 \pmod{8}. \end{aligned}$$



We can now continue to compute  $w(2^k) \pmod 8$  for  $k \geq 2$ :

$$\begin{aligned} w(2^k) &= 2 \sum_{i=2^{k-2}+1}^{2^{k-1}} w(i) + w(2^{k-2}) \\ &\equiv 2 + w(2^{k-2}) \pmod 8. \end{aligned}$$

Starting with  $w(2^0) = 1 = w(2^1)$ , the assertion now follows easily.  $\square$

We now obtain full information on the congruences modulo 8 for the  $u$ -sequence via the following result on the  $w$ -sequence modulo 4.

**Theorem 4.6.** *Let  $n \in \mathbb{N}$ ,  $n$  not a 2-power. Write  $n = \sum_{i=0}^k 2^{n_i}$  with  $n_0 < n_1 < \dots < n_k$ . Then we have*

$$w(n) \equiv \begin{cases} 0 \pmod 4 & \text{if } n_0 \equiv 3 \pmod 4 \\ & \text{or } n_0 \equiv 0 \pmod 4 \text{ and } n_k \text{ is even} \\ & \text{or } n_0 \equiv 2 \pmod 4 \text{ and } n_k \text{ is odd} \\ 2 \pmod 4 & \text{if } n_0 \equiv 1 \pmod 4 \\ & \text{or } n_0 \equiv 0 \pmod 4 \text{ and } n_k \text{ is odd} \\ & \text{or } n_0 \equiv 2 \pmod 4 \text{ and } n_k \text{ is even} \end{cases}$$

*Proof.* Assume that  $n_0 \geq 1$ ; then  $m = n - 1$  is an odd number such that  $2^{n_k} + 1 \leq m = n - 1 \leq 2^{n_k+1} - 1$ ; hence, using Corollary 4.3,  $w(n - 1) \equiv w(2^{n_k} + 1) = w(2^{n_k}) + w(2^{n_k-1}) \pmod 4$ . Then

$$w(n) = w(n - 1) + w\left(\sum_{i=0}^k 2^{n_i-1}\right) \equiv w(2^{n_k}) + w(2^{n_k-1}) + w\left(\sum_{i=0}^k 2^{n_i-1}\right) \pmod 4.$$

If  $n_0 > 1$ , we can repeat the argument to obtain (using Corollary 4.3 again)

$$\begin{aligned} w(n) &= w(n - 1) + w\left(\sum_{i=0}^k 2^{n_i-1}\right) \\ &\equiv v(n_k) + 2v(n_k - 1) + v(n_k - 2) + w\left(\sum_{i=0}^k 2^{n_i-2}\right) \pmod 4 \\ &\equiv 2v(n_k) + w\left(\sum_{i=0}^k 2^{n_i-2}\right) \pmod 4 \quad (\text{using Proposition 4.4}) \\ &\equiv 2 + w\left(\sum_{i=0}^k 2^{n_i-2}\right) \pmod 4. \end{aligned}$$

We now use this reduction to discuss the different cases for  $n_0$ .

If  $n_0 = 4j - 1$  for some  $j \in \mathbb{N}$ , then we can use the 2-step reduction above  $2j - 1$  times, then the 1-step reduction, and we obtain (using Corollary 4.3 again)

$$\begin{aligned}
w(n) &\equiv 2 + w\left(2 + \sum_{i=1}^k 2^{n_i - n_0 + 1}\right) \pmod{4} \\
&\equiv 2 + w(2^{n_k - n_0 + 1}) + w(2^{n_k - n_0}) + w\left(1 + \sum_{i=1}^k 2^{n_i - n_0}\right) \\
&\equiv 2 + w(2^{n_k - n_0 + 1}) + w(2^{n_k - n_0}) + w\left(1 + 2^{n_k - n_0}\right) \\
&\equiv 2 + w(2^{n_k - n_0 + 1}) + 2 w(2^{n_k - n_0}) + w(2^{n_k - n_0 - 1}) \\
&\equiv 2 + 2 v(n_k - n_0 + 1) \equiv 0 \pmod{4} \quad (\text{using Proposition 4.4}).
\end{aligned}$$

In the case  $n_0 = 4j + 1$  for some  $j \in \mathbb{N}$ , we are just doing one less 2-step reduction, hence in this case it follows that  $w(n) \equiv 2 \pmod{4}$ .

When  $n_0 = 4j$  for some  $j \in \mathbb{N}$ , we do again  $2j - 1$  2-step reductions and obtain

$$\begin{aligned}
w(n) &\equiv 2 + w\left(2^2 + \sum_{i=1}^k 2^{n_i - n_0 + 2}\right) \pmod{4} \\
&\equiv 2 + w\left(3 + \sum_{i=1}^k 2^{n_i - n_0 + 2}\right) + w\left(2 + \sum_{i=1}^k 2^{n_i - n_0 + 1}\right) \\
&\equiv 2 + w(2^{n_k - n_0 + 2} + 1) + 2 \\
&\equiv w(2^{n_k - n_0 + 2}) + w(2^{n_k - n_0 + 1}) \\
&\equiv v(n_k + 2) + v(n_k + 1) \pmod{4}.
\end{aligned}$$

With the previous result on the  $v$ -sequence, the assertion then follows.

When  $n_0 = 0$ , we are in the case of an odd  $n$ , where then (by Corollary 4.3)

$$w(n) = w(1 + 2^{n_k}) = w(2^{n_k}) + w(2^{n_k - 1})$$

and the result is the same as above for  $n_0 = 4j$ .

When  $n_0 = 4j - 2$  for some  $j \in \mathbb{N}$ , the result is complementary to the one above, by a shift of 2, as stated in the assertion.  $\square$

**Remark 4.7.** In Section 3 we have seen that the generating function  $W(q)$  of  $w(n)$  is the even part of  $S_1(q) + S_2(q)$ . The functional equations given in Remark 3.6 then yield

$$W(q) = q + \frac{1+q}{1-q} W(q^2).$$

Iterating this equation and considering congruences modulo 2 and modulo 4 then provides a different route to the congruence results obtained above.

We close by noting that there may also be very special behavior of the  $w$ -sequence modulo 8. (Indeed, the data strongly suggest this.) Obviously, this would then imply congruences modulo 16 for the numbers  $u(n)$ .

**Acknowledgments.** The first two authors would like to thank the Danish Research Council (FNU) for the support of their collaboration. Thanks go also to a referee for remarks on alternative proofs.

## REFERENCES

- [1] R. F. CHURCHHOUSE, Congruence properties of the binary partition function, *Proc. Camb. Phil. Soc.* **66** (1969), 371-376
- [2] G. JAMES, A. KERBER, *The representation theory of the symmetric group*. Encyclopedia of Mathematics and its Applications, 16, Addison-Wesley, Reading, Mass., 1981
- [3] J.B. OLSSON, Sign conjugacy classes in symmetric groups, *J. Algebra* **322** (2009), 2793-2800
- [4] Ø. RØDSETH, J. A. SELLERS, Binary partitions revisited, *J. Comb. Thy. Ser. A* **98** (2002), 33-45
- [5] N.J.A. SLOANE, J. A. SELLERS, On non-squashing partitions, *Discr. Math.* **294** (2005), 259-274.
- [6] R. P. STANLEY, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, New York/Cambridge, 1999
- [7] The On-Line Encyclopedia of Integer Sequences, published electronically at <http://oeis.org>, 2011

INSTITUT FÜR ALGEBRA, ZAHLENTHEORIE UND DISKRETE MATHEMATIK, LEIBNIZ UNIVERSITÄT  
HANNOVER, WELFENGARTEN 1, D-30167 HANNOVER, GERMANY

*E-mail address:* `bessen@math.uni-hannover.de`

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN  
5, DK-2100 COPENHAGEN Ø, DENMARK

*E-mail address:* `olsson@math.ku.dk`

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, 104 MCALLISTER BUILDING, UNI-  
VERSITY PARK, PA 16802, USA

*E-mail address:* `sellersj@math.psu.edu`