

K-THEORY OF FURSTENBERG TRANSFORMATION GROUP C^* -ALGEBRAS

KAMRAN REIHANI

ABSTRACT. The paper studies the K -theoretic invariants of the crossed product C^* -algebras associated with an important family of homeomorphisms of the tori \mathbb{T}^n called *Furstenberg transformations*. Using the Pimsner-Voiculescu theorem, we prove that given n , the K -groups of those crossed products, whose corresponding $n \times n$ integer matrices are unipotent of maximal degree, always have the same rank a_n . We show using the theory developed here, together with two computing programs - included in an appendix - that a claim made in the literature about the torsion subgroups of these K -groups is false. Using the representation theory of the simple Lie algebra $\mathfrak{sl}(2, \mathbb{C})$, we show that, remarkably, a_n has a combinatorial significance. For example, every a_{2n+1} is just the number of ways that 0 can be represented as a sum of integers between $-n$ and n (with no repetitions). By adapting an argument of van Lint (in which he answered a question of Erdős), a simple, explicit formula for the asymptotic behavior of the sequence $\{a_n\}$ is given. Finally, we describe the order structure of the K_0 -groups of an important class of Furstenberg crossed products, obtaining their complete Elliott invariant using classification results of H. Lin and N. C. Phillips.

CONTENTS

1. Introduction	2
2. K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$	5
3. Anzai transformation group C^* -algebras $\mathcal{A}_{n,\theta}$	7
4. A Poincaré type of duality	10
5. The rank a_n of the K -groups of $\mathcal{A}_{n,\theta}$	11
6. Combinatorial properties of the sequence $\{a_n\}$	12
6.1. Connections with representation theory of $\mathfrak{sl}(2, \mathbb{C})$	12
6.2. Generating functions for the sequence $\{a_n\}$	14
7. The positive cone of $K_0(\mathcal{F}_{\theta,f})$	18
8. Concluding remarks	21
Appendices	22
Appendix A. The Smith normal form	22
Appendix B. Nilpotent and unipotent linear mappings	23
Appendix C. Endomorphisms and derivations of exterior algebras	24
Appendix D. Representation theory of $\mathfrak{sl}(2, \mathbb{C})$	25
Appendix E. Some computer codes and a counterexample	26
Appendix F. Applications to $C^*(\mathcal{D}_n)$	27
References	29

2000 *Mathematics Subject Classification*. Primary: 19K14, 19K99, 46L35, 46L80, Secondary: 05A15, 05A16, 05A17, 15A36, 17B10, 17B20, 37B05, 54H20.

1. INTRODUCTION

Furstenberg transformations were introduced in [9] as the first examples of homeomorphisms of the tori, which under some necessary and sufficient conditions are minimal and uniquely ergodic. In some sense, they generalize the irrational rotations on the circle. They also appear in certain applications of ergodic theory to number theory (e.g. in Diophantine approximation [8]), and sometimes are called *skew product transformations* or *compound skew translations* of the tori. The terminology “Furstenberg transformation group C^* -algebra” is what we would like to use in this paper to call the crossed products associated with Furstenberg transformations, and we will denote them by $\mathcal{F}_{\theta, \mathbf{f}}$. There have been several contributions to the computations of K -theoretic invariants for some examples of these C^* -algebras in the literature (see [16, 18, 25, 30, 39] to name a few). However, a more general study of such invariants for these C^* -algebras has not been available to the best of our knowledge.

Remark 1.1. In independent (unpublished) work [16], R. Ji studied the K -groups of the C^* -algebras $\mathcal{F}_{\theta, \mathbf{f}}$ (denoted by $A_{F_{\mathbf{f}, \theta}}$ in there) associated with the descending affine Furstenberg transformations (denoted by $F_{\mathbf{f}, \theta}$ in there) on the tori. He comments that “explicitly computing the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_K \mathbb{Z} [A_{F_{\mathbf{f}, \theta}}$ for $\theta = 0]$ is still not an easy matter”. Moreover, he gives no information about the ranks of the K -groups or order structure of K_0 in general, which are studied in the present paper. As we shall see in Remark 1.7 below, the claim that he makes about the form of the torsion subgroup of $K_*(\mathcal{F}_{\theta, \mathbf{f}})$ is unfortunately not correct.

From the C^* -algebraic point view, when a Furstenberg transformation is minimal and uniquely ergodic, the associated transformation group C^* -algebra is simple and has a unique tracial state with a dense tracial range of the K_0 -group in the real line. Because of this, these C^* -algebras fit well into the classification program of G. Elliott by finding their K -theoretic invariants. In fact, in the class of transformation group C^* -algebras of uniquely ergodic minimal homeomorphisms on infinite compact metric spaces, K -theory is a complete invariant. More precisely, suppose that X is an infinite compact metric space with finite covering dimension and $h : X \rightarrow X$ is a uniquely ergodic minimal homeomorphism, and put $A := \mathcal{C}(X) \rtimes_h \mathbb{Z}$. Let τ be the trace induced by the unique invariant probability measure. Then τ is the unique tracial state on A . Let $\tau_* : K_0(A) \rightarrow \mathbb{R}$ be the induced homomorphism on $K_0(A)$ and assume that $\tau_* K_0(A)$ is dense in \mathbb{R} . Then the 4-tuple

$$(K_0(A), K_0(A)_+, [1_A], K_1(A))$$

is a complete algebraic invariant (called the *Elliott invariant* of A) [21, Corollary 4.8]. In this case, A has stable rank one, real rank zero and tracial topological rank zero in the sense of H. Lin [19]. The order on $K_0(A)$ is also determined by the unique trace τ , in the sense that an element $x \in K_0(A)$ is positive if and only if either $x = 0$ or $\tau_*(x) > 0$ [22, 29]. This implies, in particular, that the torsion subgroup of $K_0(A)$ contributes nothing interesting to the order information. In other words, the order on $K_0(A)$ is determined by the order on the free part. We will study the order structure of $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ in Section 7.

In order to compute the K -groups of a crossed product of the form $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ in general, we make use of the algebraic properties of $K_*(\mathcal{C}(\mathbb{T}^n))$ in Section 2. More precisely, $K_*(\mathcal{C}(\mathbb{T}^n))$ is an exterior algebra over \mathbb{Z}^n with a certain natural basis, and the induced automorphism α_* on $K_*(\mathcal{C}(\mathbb{T}^n))$ is in fact a ring automorphism, which makes computations much easier. In fact, it is shown in Theorem 2.1 that the problem of finding the K -groups of the transformation group C^* -algebra of a homeomorphism of the n -torus is completely computable in the sense that one only needs to calculate the kernels and cokernels of a finite number of integer matrices. These K -groups are finitely generated with the same rank (see Corollary 2.2). In the special case of Anzai transformation group C^* -algebras $\mathcal{A}_{n, \theta}$ associated with Anzai transformations on the n -torus, we denote this common rank by a_n , which we will study in detail in this paper. It is proved in Theorem 5.1 that a_n is the common rank of the K -groups of a larger class of transformation group C^* -algebras, including the C^* -algebras associated with Furstenberg transformations on \mathbb{T}^n . We describe a_n as the constant term in a certain Laurent polynomial (Theorem 6.7). Then we study

the combinatorial properties of the sequence $\{a_n\}$, which leads to a simple asymptotic formula.

To present the results and proofs of this paper we need some definitions about transformations on the tori and the corresponding C^* -crossed products. Throughout this paper, \mathbb{T}^n denotes the n -dimensional torus with coordinates $(\zeta_1, \zeta_2, \dots, \zeta_n)$.

Definition 1.2. An **affine transformation** on \mathbb{T}^n is given by

$$\beta(\zeta_1, \zeta_2, \dots, \zeta_n) = (e^{2\pi i t_1} \zeta_1^{b_{11}} \dots \zeta_n^{b_{1n}}, e^{2\pi i t_2} \zeta_1^{b_{21}} \dots \zeta_n^{b_{2n}}, \dots, e^{2\pi i t_n} \zeta_1^{b_{n1}} \dots \zeta_n^{b_{nn}}),$$

where $\mathbf{t} := (t_1, t_2, \dots, t_n) \in (\mathbb{R}/\mathbb{Z})^n$ and $\mathbf{B} := [b_{ij}]_{n \times n} \in \text{GL}(n, \mathbb{Z})$. We identify the pair (\mathbf{t}, \mathbf{B}) with β .

Note that any automorphism of \mathbb{T}^n followed by a rotation can be expressed in such a fashion. The set of affine transformations on \mathbb{T}^n form a group $\text{Aff}(\mathbb{T}^n)$, which can be identified with the semidirect product $(\mathbb{R}/\mathbb{Z})^n \rtimes \text{GL}(n, \mathbb{Z})$. More precisely, for two affine transformations $\beta = (\mathbf{t}, \mathbf{B})$ and $\beta' = (\mathbf{t}', \mathbf{B}')$ on \mathbb{T}^n , we have

$$\beta \circ \beta' = (\mathbf{t} + \mathbf{B}\mathbf{t}', \mathbf{B}\mathbf{B}') \text{ and } \beta^{-1} = (-\mathbf{B}^{-1}\mathbf{t}, \mathbf{B}^{-1}).$$

(In the expression $\mathbf{B}\mathbf{t}$, \mathbf{t} is a column vector, but for convenience we write it as a row vector.)

We remind the reader of an important fact before giving the next definition. Consider homotopy classes of continuous functions from \mathbb{T}^n to \mathbb{T} . It is well known that in each class there is a unique ‘‘linear’’ function $(\zeta_1, \dots, \zeta_n) \mapsto \zeta_1^{b_1} \dots \zeta_n^{b_n}$ for some $b_1, \dots, b_n \in \mathbb{Z}$. More precisely, every continuous function $f : \mathbb{T}^n \rightarrow \mathbb{T}$ can be written as

$$f(\zeta_1, \dots, \zeta_n) = e^{2\pi i g(\zeta_1, \dots, \zeta_n)} \zeta_1^{b_1} \dots \zeta_n^{b_n},$$

for some continuous function $g : \mathbb{T}^n \rightarrow \mathbb{R}$ and unique integer exponents b_1, \dots, b_n . In particular, the cohomotopy group $\pi^1(\mathbb{T}^n)$ is isomorphic to \mathbb{Z}^n . Following [8, p. 35], we denote the exponent b_i , which is uniquely determined by the homotopy class of f , as $b_i = A_i[f]$.

Definition 1.3. We define the following transformations in accordance with [16].

(a) A **Furstenberg transformation** $\varphi_{\theta, \mathbf{f}}$ on \mathbb{T}^n is given by

$$\varphi_{\theta, \mathbf{f}}^{-1}(\zeta_1, \zeta_2, \dots, \zeta_n) = (e^{2\pi i \theta} \zeta_1, f_1(\zeta_1)\zeta_2, f_2(\zeta_1, \zeta_2)\zeta_3, \dots, f_{n-1}(\zeta_1, \dots, \zeta_{n-1})\zeta_n),$$

where θ is a real number, each $f_i : \mathbb{T}^i \rightarrow \mathbb{T}$ is a continuous function with $A_i[f_i] \neq 0$ for $i = 1, \dots, n-1$, and $\mathbf{f} = (f_1, \dots, f_{n-1})$.

(b) An **affine Furstenberg transformation** α on \mathbb{T}^n is given by

$$\alpha^{-1}(\zeta_1, \zeta_2, \dots, \zeta_n) = (e^{2\pi i \theta} \zeta_1, \zeta_1^{b_{12}} \zeta_2, \zeta_1^{b_{13}} \zeta_2^{b_{23}} \zeta_3, \dots, \zeta_1^{b_{1n}} \zeta_2^{b_{2n}} \dots \zeta_{n-1}^{b_{n-1,n}} \zeta_n),$$

where θ is a real number and the exponents b_{ij} are integers and $b_{i,i+1} \neq 0$ for $i = 1, \dots, n-1$.

(c) An **ascending Furstenberg transformation** α on \mathbb{T}^n is given by

$$\alpha^{-1}(\zeta_1, \zeta_2, \dots, \zeta_n) = (e^{2\pi i \theta} \zeta_1, \zeta_1^{k_1} \zeta_2, \zeta_2^{k_2} \zeta_3, \dots, \zeta_{n-1}^{k_{n-1}} \zeta_n),$$

where θ is a real number and the exponents k_i are nonzero integers and $k_i \mid k_{i+1}$ for $i = 1, \dots, n-2$.

(d) In (c), if $k_i = 1$ for $i = 1, \dots, n-1$, the transformation is called an **Anzai transformation** $\sigma_{n, \theta}$ on \mathbb{T}^n . Thus it is given by

$$\sigma_{n, \theta}^{-1}(\zeta_1, \zeta_2, \dots, \zeta_n) = (e^{2\pi i \theta} \zeta_1, \zeta_1 \zeta_2, \dots, \zeta_{n-1} \zeta_n),$$

where θ is a real number. We usually drop the indices n and θ and write only σ for more convenience.

Note that one can easily verify that $\varphi_{\theta, \mathbf{f}}$ is a homeomorphism. Also in the above definition, we have converted ‘‘descending’’, which is used in [16, Definition 2.16], to ‘‘ascending’’ since the order of coordinates there is opposite to ours.

For certain Furstenberg transformations on \mathbb{T}^n we have the following theorem.

Theorem 1.4. ([9], 2.3) *If θ is irrational, then $\varphi_{\theta, \mathbf{f}}$ defines a minimal dynamical system on \mathbb{T}^n . If in addition, each f_i satisfies a uniform Lipschitz condition in ζ_i for $i = 1, \dots, n-1$, then $\varphi_{\theta, \mathbf{f}}$ is a uniquely ergodic transformation and the unique invariant measure is the normalized Lebesgue measure on \mathbb{T}^n . In particular, every affine Furstenberg transformation defines a minimal and uniquely ergodic dynamical system if θ is irrational.*

As a conclusion, we have the following result for the Furstenberg transformation group C^* -algebra $\mathcal{F}_{\theta, \mathbf{f}} := \mathcal{C}(\mathbb{T}^n) \rtimes_{\varphi_{\theta, \mathbf{f}}} \mathbb{Z}$ as introduced in [16].

Corollary 1.5. *$\mathcal{F}_{\theta, \mathbf{f}} = \mathcal{C}(\mathbb{T}^n) \rtimes_{\varphi_{\theta, \mathbf{f}}} \mathbb{Z}$ is a simple C^* -algebra for irrational θ . If in addition, each f_i satisfies a uniform Lipschitz condition in ζ_i for $i = 1, \dots, n-1$, then $\mathcal{F}_{\theta, \mathbf{f}}$ has a unique tracial state.*

Proof. For the first part, the minimality of the action as stated in the preceding theorem implies the simplicity of $\mathcal{F}_{\theta, \mathbf{f}}$ [4, 32]. For the second part, one can easily check that since θ is irrational, the action of \mathbb{Z} on \mathbb{T}^n generated by $\varphi_{\theta, \mathbf{f}}$ is free. So there are no periodic points in \mathbb{T}^n . This and the unique ergodicity of $\varphi_{\theta, \mathbf{f}}$ yield the result [38, Corollary 3.3.10, p. 91]. \square

Remark 1.6. Using the preceding corollary and much like the proof of Theorem 2.1 in [33], one can prove that for irrational θ , $\mathcal{F}_{\theta, \mathbf{f}}$ is in fact the unique C^* -algebra generated by unitaries U, V_1, \dots, V_n satisfying the commutator relations

$$(\mathbf{CR})_{\mathbf{f}} \quad [U, V_1] = e^{2\pi i \theta}, [U, V_2] = f_1(V_1), \dots, [U, V_n] = f_{n-1}(V_1, \dots, V_{n-1}),$$

where $[a, b] := aba^{-1}b^{-1}$ and all other pairs of operators from U, V_1, \dots, V_n commute.

Remark 1.7. In [16, Proposition 2.17], R. Ji claims to have proved

(*) *If $\varphi_{\theta, \mathbf{f}}$ is an ascending Furstenberg transformation on \mathbb{T}^n with the ascending sequence $\{k_1, k_2, \dots, k_{n-1}\}$, then the torsion subgroup of $K_*(\mathcal{F}_{\theta, \mathbf{f}})$ is isomorphic to $\mathbb{Z}_{k_1} \oplus \mathbb{Z}_{k_2}^{(m_2)} \oplus \dots \oplus \mathbb{Z}_{k_{n-1}}^{(m_{n-1})}$, where the group $\mathbb{Z}_{k_i}^{(m_i)}$ is the direct product of m_i copies of the cyclic group $\mathbb{Z}_{k_i} = \mathbb{Z}/k_i\mathbb{Z}$.*

From this claim one would immediately deduce that the K -groups of the C^* -algebra $\mathcal{A}_{n, \theta} := \mathcal{C}(\mathbb{T}^n) \rtimes_{\sigma} \mathbb{Z}$ generated by an Anzai transformation σ on \mathbb{T}^n should be torsion-free. However, we will show that this is not true in general. This type of example first appears for $n = 6$, which seems already beyond hand calculation. (We admit that hand calculation would be the most convincing method to use; however, it is not practicable.) As the first counterexample we obtained by computer, we will see in Example E.1 that $K_1(\mathcal{A}_{6, \theta}) \cong \mathbb{Z}^{13} \oplus \mathbb{Z}_2$ (also, see Example 3.3). In fact, the error in the proof of (*) is in [16, p. 29, 1.2]; there it is “clearly” assumed that using a matrix S in $\mathrm{GL}(2^n, \mathbb{Z})$, one can delete all entries denoted by \star 's in $K_* - \mathbf{l}$, where K_* is the $2^n \times 2^n$ integer matrix corresponding to $\mathcal{F}_{\theta, \mathbf{f}}$ that acts on $K_*(\mathcal{C}(\mathbb{T}^n)) = \Lambda^* \mathbb{Z}^n$ with respect to a certain ordered basis. This error arose originally from the general form of the matrix K_* in [16, p. 27], which is not correct. R. Ji went on to use the torsion subgroup in (*) as an invariant to classify the C^* -algebras generated by ascending transformations and matrix algebras over them [16, Theorem 3.6]. We do not know whether those classifications holds.

Question. Do there exist two different ascending Furstenberg transformations with the same parameter θ and with isomorphic transformation group C^* -algebras?

It is worth mentioning that explicit calculations of K -groups in terms of the parameters involved are possible in low dimensions (of the tori), and answer the question raised above negatively. However, such calculations in terms of the given integer parameters (exponents) of the ascending Furstenberg transformation become quickly cumbersome and impossible in higher dimensions. We have used the computer codes in Appendix E for several numerical values of the parameters in higher dimensions, and we have not found any examples leading to the negative answer to this question yet. The torsion subgroups of

the K -groups are usually larger than what R. Ji claimed in [16, Proposition 2.17]; it seems likely that the torsion subgroups depend on the parameters involved in such a way that Ji's classification result is still true. We are currently investigating this problem.

This paper is organized as follows. In Section 2, we review the general approach of exterior algebras for finding K -groups of transformation group C^* -algebras of homeomorphisms of the tori. In Section 3, we apply this method to the important case of Anzai transformations and give the K -groups of their transformation group C^* -algebras based on the tori of dimension up to 12 in Table 1 using the computer programs given in Appendix E. In Section 4, we establish a Poincaré type of duality for the cokernels of integer matrices that leads to some interesting facts about the K -groups when the dimension of the underlying torus is odd. In Section 5, we focus on the rank a_n of the K -groups of Anzai transformations group C^* -algebras based on the n -torus, and we show that a_n is, in fact, the rank of the K -groups of a large class of transformation group C^* -algebras including those associated with Furstenberg transformations on the n -torus. In Section 6, we first uncover an interesting connection between studying a_n and the irreducible representations of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$. This leads to a formula for a_n in terms of certain partitions of integers. Then we use this formula to show several interesting combinatorial properties of the sequence $\{a_n\}$. In Section 7, we study the order structure of the K_0 -group of a class of simple Furstenberg transformation group C^* -algebras to make their Elliott invariants more accessible. The appendices at the end are provided mainly for self-containment of the paper, but we sometimes refer to them for the proof of some lemmas or propositions that are somewhat far from the main concepts and goals of this paper by nature. Appendix F contains some applications of the results of this paper to our earlier work [33].

2. K -GROUPS OF $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$

In this section, we describe a general method to compute the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$, where α is an arbitrary homeomorphism of \mathbb{T}^n . (By abuse of notation, the automorphism α of $\mathcal{C}(\mathbb{T}^n)$ is defined by $\alpha(f) = f \circ \alpha^{-1}$ for $f \in \mathcal{C}(\mathbb{T}^n)$.) To do this, we will pay special attention to the algebraic structure of $K^*(\mathbb{T}^n)$ and how the induced automorphisms on it can be realized. Note that it is sufficient to consider the special case of “linear” homeomorphisms since, as stated before Definition 1.3, every continuous function $f : \mathbb{T}^n \rightarrow \mathbb{T}$ is homotopic to a unique “linear” function $(\zeta_1, \dots, \zeta_n) \mapsto \zeta_1^{b_1} \dots \zeta_n^{b_n}$ for some integer exponents b_1, \dots, b_n . Moreover, the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ depend (up to isomorphism) only on the homotopy class of α [3, Corollary 10.5.2].

It is well known that $K^*(\mathbb{T}^n)$ is a \mathbb{Z}_2 -graded ring, and by the Künneth formula (see [2, Corollary 2.7.15] or [36, Theorem 4.1]), it is an exterior algebra (over \mathbb{Z}) on n generators, where the elements of even degree are in $K^0(\mathbb{T}^n)$ and those of odd degree are in $K^1(\mathbb{T}^n)$. The generators of this exterior algebra correspond to the generators of the dual group \mathbb{Z}^n of \mathbb{T}^n [37, p. 185]. Indeed, in this case the Chern character

$$\text{ch} : K^*(\mathbb{T}^n) \longrightarrow \check{H}^*(\mathbb{T}^n, \mathbb{Q})$$

is integral and gives the Chern isomorphisms

$$\text{ch}_0 : K^0(\mathbb{T}^n) \longrightarrow \check{H}^{\text{even}}(\mathbb{T}^n, \mathbb{Z}),$$

$$\text{ch}_1 : K^1(\mathbb{T}^n) \longrightarrow \check{H}^{\text{odd}}(\mathbb{T}^n, \mathbb{Z}),$$

where $\check{H}^*(\mathbb{T}^n, \mathbb{Z}) \cong \Lambda_{\mathbb{Z}}^*(e_1, \dots, e_n)$ is the (Čech) cohomology ring of \mathbb{T}^n under the cup product, and $\check{H}^k(\mathbb{T}^n, \mathbb{Z}) \cong \Lambda_{\mathbb{Z}}^k(e_1, \dots, e_n)$. On the other hand, $K^*(\mathbb{T}^n) \cong K_*(\mathcal{C}(\mathbb{T}^n))$. So by introducing $e_i := [z_i]_1$, i.e. the class in $K_1(\mathcal{C}(\mathbb{T}^n))$ of the coordinate function $z_i : \mathbb{T}^n \rightarrow \mathbb{T}$ given by

$$z_i(\zeta_1, \dots, \zeta_n) = \zeta_i$$

as a unitary element of $\mathcal{C}(\mathbb{T}^n)$ for $i = 1, \dots, n$, we have the isomorphisms $K_*(\mathcal{C}(\mathbb{T}^n)) \cong \Lambda_{\mathbb{Z}}^*(e_1, \dots, e_n) \cong \Lambda^* \mathbb{Z}^n$, which respect the canonical embedding of \mathbb{Z}^n . Moreover, these isomorphisms are unique since only

the identity automorphism of the ring $\Lambda^*\mathbb{Z}^n$ fixes each element of \mathbb{Z}^n .

Now, we use the Pimsner-Voiculescu six term exact sequence [31] as the main tool for computing the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$. Let $\alpha_*(= K_*(\alpha))$ be the ring automorphism of $K_*(\mathcal{C}(\mathbb{T}^n))$ induced by α and let α_i be the restriction of α_* on $K_i(\mathcal{C}(\mathbb{T}^n))$ for $i = 0, 1$, and set $A := \mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$. Then we have the following exact sequence.

$$(1) \quad \begin{array}{ccccc} K_0(\mathcal{C}(\mathbb{T}^n)) & \xrightarrow{\alpha_0 - \text{id}} & K_0(\mathcal{C}(\mathbb{T}^n)) & \xrightarrow{j_0} & K_0(A) \\ \exp \uparrow & & & & \downarrow \partial \\ K_1(A) & \xleftarrow{j_1} & K_1(\mathcal{C}(\mathbb{T}^n)) & \xleftarrow{\alpha_1 - \text{id}} & K_1(\mathcal{C}(\mathbb{T}^n)) \end{array}$$

Here, $j : \mathcal{C}(\mathbb{T}^n) \rightarrow A$ is the canonical embedding of $\mathcal{C}(\mathbb{T}^n)$ in A , $j_0 := K_0(j)$ and $j_1 := K_1(j)$. Also, from now on id denotes the identity function on each underlying set. As a result, we have the following short exact sequences

$$(2) \quad 0 \longrightarrow \text{coker}(\alpha_0 - \text{id}) \longrightarrow K_0(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \longrightarrow \ker(\alpha_1 - \text{id}) \longrightarrow 0,$$

$$(3) \quad 0 \longrightarrow \text{coker}(\alpha_1 - \text{id}) \longrightarrow K_1(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \longrightarrow \ker(\alpha_0 - \text{id}) \longrightarrow 0.$$

Since all the groups involved are abelian and finitely generated, and $\ker(\alpha_i - \text{id})$ is torsion-free ($i = 0, 1$), these short exact sequences split (since projective \mathbb{Z} -modules are precisely free abelian groups), and we have

$$(4) \quad \boxed{K_0(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong \text{coker}(\alpha_0 - \text{id}) \oplus \ker(\alpha_1 - \text{id})},$$

$$(5) \quad \boxed{K_1(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong \text{coker}(\alpha_1 - \text{id}) \oplus \ker(\alpha_0 - \text{id})}.$$

So it suffices to determine the kernel and cokernel of $(\alpha_0 - \text{id})$ and $(\alpha_1 - \text{id})$ acting as endomorphisms on the finitely generated abelian groups $\Lambda_{\mathbb{Z}}^{\text{even}}(e_1, \dots, e_n) \cong \mathbb{Z}^{2^{n-1}}$ and $\Lambda_{\mathbb{Z}}^{\text{odd}}(e_1, \dots, e_n) \cong \mathbb{Z}^{2^{n-1}}$, respectively. Note that from the isomorphisms (4) and (5), the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ are finitely generated abelian groups. Now, since α_* is a ring homomorphism, it suffices to know the action of α_* on e_1, \dots, e_n . In fact, for a general basis element $e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_r}$ of $K_*(\mathcal{C}(\mathbb{T}^n)) \cong \Lambda_{\mathbb{Z}}^*(e_1, \dots, e_n)$ we have

$$\alpha_*(e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_r}) = \alpha_*(e_{i_1}) \wedge \alpha_*(e_{i_2}) \wedge \dots \wedge \alpha_*(e_{i_r}).$$

Thus if we consider $\{e_1, \dots, e_n\}$ as the canonical basis of \mathbb{Z}^n and take $\hat{\alpha} = \alpha_*|_{\mathbb{Z}^n}$, we have $\alpha_* = \wedge^* \hat{\alpha} = \bigoplus_{r=1}^n \wedge^r \hat{\alpha}$, $\alpha_0 = \wedge^{\text{even}} \hat{\alpha} = \bigoplus_{r \geq 0} \wedge^{2r} \hat{\alpha}$ and $\alpha_1 = \wedge^{\text{odd}} \hat{\alpha} = \bigoplus_{r \geq 0} \wedge^{2r+1} \hat{\alpha}$, where $\wedge^i \hat{\alpha}$ is the i -th exterior power of $\hat{\alpha}$, which acts on $\Lambda^i \mathbb{Z}^n$ for $i = 0, 1, \dots, n$. Now, let $\alpha^{-1} = (f_1, \dots, f_n)$ and $a_{ji} := A_j[f_i]$, or in other words, assume that f_i is homotopic to $z_1^{a_{1i}} \dots z_n^{a_{ni}} : (\zeta_1, \dots, \zeta_n) \mapsto \zeta_1^{a_{1i}} \dots \zeta_n^{a_{ni}}$ for $i = 1, \dots, n$. So we can write

$$\alpha_*(e_i) = \alpha_*[z_i]_1 = [\alpha(z_i)]_1 = [z_i \circ \alpha^{-1}]_1 = [f_i]_1 = [z_1^{a_{1i}} \dots z_n^{a_{ni}}]_1 = \sum_{j=1}^n a_{ji} [z_j]_1 = \sum_{j=1}^n a_{ji} e_j.$$

Therefore $\hat{\alpha}$ acts on \mathbb{Z}^n via the corresponding integer matrix $\mathbf{A} := [a_{ij}]_{n \times n} \in \text{GL}(n, \mathbb{Z})$, α_* acts on $\Lambda^* \mathbb{Z}^n$ via $\wedge^* \mathbf{A}$, and we have the following isomorphisms

$$K_0(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong \text{coker}(\alpha_0 - \text{id}) \oplus \ker(\alpha_1 - \text{id}) = \text{coker}(\bigoplus_{r \geq 0} \wedge^{2r} \hat{\alpha} - \text{id}) \oplus \ker(\bigoplus_{r \geq 0} \wedge^{2r+1} \hat{\alpha} - \text{id}),$$

so we can write

$$\boxed{K_0(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong \bigoplus_{r \geq 0} [\text{coker}(\wedge^{2r} \hat{\alpha} - \text{id}) \oplus \ker(\wedge^{2r+1} \hat{\alpha} - \text{id})]},$$

and similarly

$$K_1(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong \bigoplus_{r \geq 0} [\text{coker}(\wedge^{2r+1} \hat{\alpha} - \text{id}) \oplus \text{ker}(\wedge^{2r} \hat{\alpha} - \text{id})].$$

We summarize the arguments discussed above in the following theorem.

Theorem 2.1. *Let α be a homeomorphism of \mathbb{T}^n and $\hat{\alpha} \in \text{Aut}(\mathbb{Z}^n)$ be the restriction of α_* to \mathbb{Z}^n (as above). Then $\alpha_* = \wedge^* \hat{\alpha} = \bigoplus_{r=1}^n \wedge^r \hat{\alpha}$ on $K^*(\mathbb{T}^n) = \Lambda^* \mathbb{Z}^n$ and*

$$K_0(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong \bigoplus_{r \geq 0} [\text{coker}(\wedge^{2r} \hat{\alpha} - \text{id}) \oplus \text{ker}(\wedge^{2r+1} \hat{\alpha} - \text{id})],$$

$$K_1(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong \bigoplus_{r \geq 0} [\text{coker}(\wedge^{2r+1} \hat{\alpha} - \text{id}) \oplus \text{ker}(\wedge^{2r} \hat{\alpha} - \text{id})].$$

Therefore in order to compute the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$, we must find the kernel and cokernel of $\wedge^r \hat{\alpha} - \text{id}$ as an endomorphism of $\Lambda^r \mathbb{Z}^n$ for $r = 0, 1, \dots, n$. Note that the matrix of $\wedge^r \hat{\alpha} - \text{id}$ with respect to the canonical basis $\{e_{i_1} \wedge \dots \wedge e_{i_r} \mid 1 \leq i_1 < \dots < i_r \leq n\}$ with lexicographic order is $A_{n,r} := \wedge^r A - I_{\binom{n}{r}}$, which is an integer matrix of order $\binom{n}{r}$ (I_k is the identity matrix of order k - we often omit k whenever it is clear). So by computing the kernel and cokernel of $A_{n,r}$ for $r = 0, 1, \dots, n$ with appropriate tools (such as the Smith normal form), one can determine the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$. The author has written some Maple codes to handle such computations (see Appendix E).

Corollary 2.2. *The K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ are finitely generated abelian groups with the same rank. Moreover, this common rank equals*

$$\text{rank ker}(\wedge^* \hat{\alpha} - \text{id}) = \sum_{r=0}^n \text{rank ker}(\wedge^r \hat{\alpha} - \text{id}).$$

Proof. Use the previous proposition and note that for any $\varphi \in \text{End}(\mathbb{Z}^n)$ one has $\text{rank ker } \varphi = \text{rank coker } \varphi$ by the Smith normal form theorem (see Theorem A.2). \square

Corollary 2.3. *If α, β are homeomorphisms of \mathbb{T}^n , whose corresponding integer matrices $A, B \in \text{GL}(n, \mathbb{Z})$ are similar over \mathbb{Z} , then*

$$K_j(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong K_j(\mathcal{C}(\mathbb{T}^n) \rtimes_{\beta} \mathbb{Z}), \quad (j = 1, 2).$$

Proof. The assumption obviously implies that the automorphisms $\hat{\alpha}$ and $\hat{\beta}$ are conjugate in $\text{Aut}(\mathbb{Z}^n)$. This together with an easy application of the identity $\wedge^r(\hat{\phi} \circ \hat{\psi}) = (\wedge^r \hat{\phi}) \circ (\wedge^r \hat{\psi})$ (see Appendix C) imply that $\wedge^r \hat{\alpha}$ and $\wedge^r \hat{\beta}$ (and therefore $\wedge^r \hat{\alpha} - \text{id}$ and $\wedge^r \hat{\beta} - \text{id}$) are conjugate in $\text{Aut}(\Lambda^r \mathbb{Z}^n)$ for $r = 0, 1, \dots, n$. The result follows now from Theorem 2.1. \square

3. ANZAI TRANSFORMATION GROUP C*-ALGEBRAS $\mathcal{A}_{n,\theta}$

The simplest case of a Furstenberg transformation on an n -torus is an Anzai transformation σ , which was defined in part (d) of Definition 1.3. To study the K -groups of Anzai transformation group C^* -algebras $\mathcal{A}_{n,\theta} = \mathcal{C}(\mathbb{T}^n) \rtimes_{\sigma} \mathbb{Z}$ using methods of the previous section, we will first need the “linearized” form of the corresponding affine homeomorphism σ^{-1} , which is as follows

$$(\zeta_1, \zeta_2, \dots, \zeta_n) \mapsto (\zeta_1, \zeta_1 \zeta_2, \dots, \zeta_{n-1} \zeta_n).$$

So $\hat{\sigma}(e_i) = e_{i-1} + e_i$ for $i = 1, \dots, n$ ($e_0 := 0$). The matrix with respect to the canonical basis $\{e_1, \dots, e_n\}$ of \mathbb{Z}^n that corresponds to $\hat{\sigma}$ is the full Jordan block

$$S_n := \begin{pmatrix} 1 & 1 & 0 & \cdots & 0 \\ 0 & 1 & 1 & & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & & 0 & 1 & 1 \\ 0 & \cdots & 0 & 0 & 1 \end{pmatrix}_{n \times n}$$

The following examples illustrates the methods described in the previous section for computing the K -groups of Anzai transformation group C^* -algebras $\mathcal{A}_{n,\theta}$.

Example 3.1. We compute the K -groups of $\mathcal{A}_{3,\theta}$, which were computed in [39] by another method (the C^* -algebra was denoted by $A_\theta^{5,5}$ in there). In fact, the Chern character and noncommutative geometry were used in [39] to compute the kernel and cokernel of $\sigma_i - \text{id}$ for $i = 0, 1$. However, we compute the kernel and cokernel of $S_{3,r} := \wedge^r S_3 - I_{\binom{3}{r}}$ for $r = 0, 1, 2, 3$, where

$$S_3 = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$r = 0$) $S_{3,0} = \wedge^0 S_3 - I_1 = [0]$. So $\ker S_{3,0} = \mathbb{Z}$ and $\text{coker } S_{3,0} = \mathbb{Z}/\langle 0 \rangle \cong \mathbb{Z}$.

$r = 1$)

$$S_{3,1} = \wedge^1 S_3 - I_3 = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

So

$$\begin{aligned} \ker S_{3,1} &= \{(x, y, z) \in \mathbb{Z}^3 \mid y = z = 0\} = (\mathbb{Z}, 0, 0) \cong \mathbb{Z}, \\ \text{coker } S_{3,1} &= \mathbb{Z}^3 / S_{3,1} \mathbb{Z}^3 = \mathbb{Z}^3 / \langle e_1, e_2 \rangle \cong \mathbb{Z}. \end{aligned}$$

$r = 2$)

$$S_{3,2} = \wedge^2 S_3 - I_3 = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

So

$$\begin{aligned} \ker S_{3,2} &= \{(x, y, z) \in \mathbb{Z}^3 \mid y + z = z = 0\} = (\mathbb{Z}, 0, 0) \cong \mathbb{Z}, \\ \text{coker } S_{3,2} &= \mathbb{Z}^3 / S_{3,2} \mathbb{Z}^3 = \mathbb{Z}^3 / \langle e_1, e_2 \rangle \cong \mathbb{Z}. \end{aligned}$$

$r = 3$) $S_{3,3} = \wedge^3 S_3 - I_1 = [0]$. So $\ker S_{3,3} = \mathbb{Z}$ and $\text{coker } S_{3,3} = \mathbb{Z}/\langle 0 \rangle \cong \mathbb{Z}$.

Now, using Theorem 2.1 we have

$$K_0(A_\theta^{5,5}) = K_0(\mathcal{A}_{3,\theta}) \cong (\text{coker } S_{3,0} \oplus \text{coker } S_{3,2}) \oplus (\ker S_{3,1} \oplus \ker S_{3,3}) \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} = \mathbb{Z}^4,$$

$$K_1(A_\theta^{5,5}) = K_1(\mathcal{A}_{3,\theta}) \cong (\text{coker } S_{3,1} \oplus \text{coker } S_{3,3}) \oplus (\ker S_{3,0} \oplus \ker S_{3,2}) \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} = \mathbb{Z}^4.$$

Notation 3.2. We let $a_n := \text{rank } K_0(\mathcal{A}_{n,\theta}) = \text{rank } K_1(\mathcal{A}_{n,\theta})$ and $a_{n,r} := \text{rank } \ker(\wedge^r S_n - \text{I})$ for $r = 0, 1, \dots, n$. From Corollary 2.2 we have

$$a_n = \text{rank } \ker(\wedge^* S_n - \text{I}) = \sum_{r=0}^n a_{n,r}.$$

Example 3.3. Using the methods described in Section 2, we have obtained the K -groups of $\mathcal{A}_{n,\theta}$ by computer for $1 \leq n \leq 12$. The cases $n = 1, 2, 3$ have been calculated in the literature already: $\mathcal{A}_{1,\theta} = A_\theta$ in [34]; $\mathcal{A}_{2,\theta} = A_\theta^4$ in [25]; and $\mathcal{A}_{3,\theta} = A_\theta^{5,5}$ in [39]. However, there are no explicit computations for the higher dimensional cases starting with $\mathcal{A}_{4,\theta} = A_\theta^{6,10}$ as in [24] since hand calculations of kernels and cokernels of the maps become quickly impossible. Using the Maple codes given in Appendix E, we can find the kernel and cokernel of

$$S_{n,r} := \wedge^r S_n - \text{I}_{\binom{n}{r}}$$

for $r = 0, 1, \dots, n$ and $n = 1, \dots, 12$ by means of the Smith normal form theorem (see Appendix A), and therefore we can compute the K -groups. The results are illustrated in Table 1, where $\mathbb{Z}_k^{(m)}$ denotes the direct sum of m copies of the cyclic group $\mathbb{Z}_k = \mathbb{Z}/k\mathbb{Z}$.

TABLE 1. The K -groups of $\mathcal{A}_{n,\theta}$ for $1 \leq n \leq 12$

n	$K_0(\mathcal{A}_{n,\theta})$	$K_1(\mathcal{A}_{n,\theta})$	a_n
1	\mathbb{Z}^2	\mathbb{Z}^2	2
2	\mathbb{Z}^3	\mathbb{Z}^3	3
3	\mathbb{Z}^4	\mathbb{Z}^4	4
4	\mathbb{Z}^6	\mathbb{Z}^6	6
5	\mathbb{Z}^8	\mathbb{Z}^8	8
6	\mathbb{Z}^{13}	$\mathbb{Z}^{13} \oplus \mathbb{Z}_2$	13
7	\mathbb{Z}^{20}	\mathbb{Z}^{20}	20
8	$\mathbb{Z}^{32} \oplus \mathbb{Z}_8^{(2)}$	$\mathbb{Z}^{32} \oplus \mathbb{Z}_{18}^{(2)}$	32
9	$\mathbb{Z}^{52} \oplus \mathbb{Z}_3^{(2)} \oplus \mathbb{Z}_9^{(2)}$	$\mathbb{Z}^{52} \oplus \mathbb{Z}_3^{(2)} \oplus \mathbb{Z}_9^{(2)}$	52
10	$\mathbb{Z}^{90} \oplus \mathbb{Z}_{55}^{(4)}$	$\mathbb{Z}^{90} \oplus \mathbb{Z}_{11}^{(2)} \oplus \mathbb{Z}_{99} \oplus \mathbb{Z}_{198} \oplus \mathbb{Z}_{2574}$	90
11	$\mathbb{Z}^{152} \oplus \mathbb{Z}_{11}^{(12)} \oplus \mathbb{Z}_{143}^{(4)} \oplus \mathbb{Z}_{286}^{(2)}$	$\mathbb{Z}^{152} \oplus \mathbb{Z}_{11}^{(12)} \oplus \mathbb{Z}_{143}^{(4)} \oplus \mathbb{Z}_{286}^{(2)}$	152
12	$\mathbb{Z}^{268} \oplus \mathbb{Z}_{13}^{(14)} \oplus \mathbb{Z}_{26}^{(4)} \oplus \mathbb{Z}_{1716}^{(4)} \oplus \mathbb{Z}_{3432}^{(2)} \oplus \mathbb{Z}_{58344}^{(2)}$	$\mathbb{Z}^{268} \oplus \mathbb{Z}_{13}^{(4)} \oplus \mathbb{Z}_{26}^{(4)} \oplus \mathbb{Z}_{286}^{(6)} \oplus \mathbb{Z}_{4862}^{(2)} \oplus \mathbb{Z}_{68068}^{(2)}$	268

Due to computational limitations, we do not have any results yet for $n > 12$, except for the sequence of ranks $\{a_n\}$, which we will study in detail in Sections 5 and 6. We will show the importance of

this sequence in Section 5. Briefly, a_n is the common rank of the K -groups of a certain family of C^* -algebras including Furstenberg transformation group C^* -algebras $\mathcal{F}_{\theta, \mathbf{f}}$ based on \mathbb{T}^n . Also, we will prove that $\{a_n\}$ is a strictly increasing sequence (see Proposition 6.5). On the other hand, it seems that the K -groups of $\mathcal{A}_{n, \theta}$ have torsion in general. The first example is $K_1(\mathcal{A}_{6, \theta})$; this is in fact because $\text{coker } \mathbb{S}_{6,3} = \text{coker}(\wedge^3 \mathbb{S}_6 - \text{l}_{20}) \cong \mathbb{Z}^3 \oplus \mathbb{Z}_2$ (see Example E.1 and Remark 1.7). Also, it is seen that the K_0 - and K_1 -groups are isomorphic for odd values of n in Table 1. In fact, this is true for more general cases (see Theorem 4.3).

4. A POINCARÉ TYPE OF DUALITY

As stated in Theorem 2.1, the K -groups of a transformation group C^* -algebras of the form $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ are completely determined by the corresponding homomorphism $\hat{\alpha} \in \text{Aut}(\mathbb{Z}^n)$ and its exterior powers. From a computational point of view, we only need the cokernels of the maps involved, since we know that for any endomorphism ϱ on \mathbb{Z}^m , $\ker \varrho \cong \text{coker} \varrho / \text{tor}(\text{coker} \varrho)$, where $\text{tor}(G)$ denotes the torsion subgroup of the finitely generated abelian group G (see Appendix A). When $\det \hat{\alpha} = 1$, we don't even need to compute all the cokernels. This is due to the following proposition, which establishes a Poincaré type of duality between cokernels of certain integer matrices. We refer to Definition A.1 for the notion of equivalence of endomorphisms of \mathbb{Z}^n .

Proposition 4.1 (Poincaré duality). *Let $\hat{\alpha} \in \text{SL}(n, \mathbb{Z})$ (i.e. $\det \hat{\alpha} = 1$). Then $\wedge^r \hat{\alpha} - \text{id}$ and $\wedge^{n-r} \hat{\alpha} - \text{id}$ are equivalent as endomorphisms of $\wedge^r \mathbb{Z}^n = \wedge^{n-r} \mathbb{Z}^n = \mathbb{Z}^{\binom{n}{r}}$. Equivalently, $\text{coker}(\wedge^r \hat{\alpha} - \text{id})$ is isomorphic to $\text{coker}(\wedge^{n-r} \hat{\alpha} - \text{id})$ for $r = 0, 1, \dots, n$.*

Proof. We prove the equivalence of the endomorphisms for their corresponding integer matrices with respect to a certain basis. Let $\mathcal{E} = \{e_1, \dots, e_n\}$ be a basis for \mathbb{Z}^n and set $S = \{1, 2, \dots, n\}$. For $I = \{i_1, \dots, i_r\} \subset S$ with $1 \leq i_1 < \dots < i_r \leq n$, put $e_I = e_{i_1} \wedge \dots \wedge e_{i_r} \in \wedge^r \mathbb{Z}^n$. Then $\mathcal{E}_r := \{e_I \mid I \subset S, |I| = r\}$ is a basis for $\wedge^r \mathbb{Z}^n$. Let $\omega := e_1 \wedge \dots \wedge e_n$, which generates $\wedge^n \mathbb{Z}^n$. We have $\wedge^0 \hat{\alpha} - \text{id} = 0$, and $\wedge^n \hat{\alpha}(\omega) = \hat{\alpha}(e_1) \wedge \dots \wedge \hat{\alpha}(e_n) = (\det \hat{\alpha})(e_1 \wedge \dots \wedge e_n) = \omega$, so $\wedge^n \hat{\alpha} - \text{id} = 0$. Now, fix an $r \in \{1, \dots, n-1\}$. For an arbitrary subset $I \subset S$ with $|I| = r$, take $J = \mathcal{E} \setminus I = \{j_1, \dots, j_{n-r}\}$, so $|J| = n-r$. Then $e_I \wedge e_J = (\text{sgn } \mu) \omega$, in which $\mu \in S_n$ is the permutation that converts $(1, 2, \dots, n)$ to $(i_1, \dots, i_r, j_1, \dots, j_{n-r})$. It is easily seen that $\mu = \mu_1 \dots \mu_r$, where μ_k is the permutation that takes i_k from its position in $(1, 2, \dots, n)$ to its new position in $(i_1, \dots, i_r, j_1, \dots, j_{n-r})$. One can see that μ_k is the combination of $i_k - (r - k + 1)$ number of transpositions ($k = 1, \dots, r$). Thus

$$\text{sgn } \mu = \prod_{k=1}^r (-1)^{i_k - (r - k + 1)} = (-1)^{\ell(I) - \frac{r(r+1)}{2}},$$

where $\ell(I) := \sum_{k=1}^r i_k$. Now, take $m = \binom{n}{r} = \binom{n}{n-r}$ and let $\mathcal{E}_r = \{e_{I_1}, \dots, e_{I_m}\}$ be a basis for $\wedge^r \mathbb{Z}^n$. Write $\mathcal{E}_{n-r} = \{e_{J_1}, \dots, e_{J_m}\}$ as the basis for $\wedge^{n-r} \mathbb{Z}^n$ such that $J_k = \mathcal{E} \setminus I_k$ for $k = 1, \dots, m$. From the above argument one can write

$$e_{I_i} \wedge e_{J_j} = (-1)^{\ell(I_i) - \frac{r(r+1)}{2}} \delta_{ij} \omega,$$

since if $i \neq j$ then $I_i \cap J_j \neq \emptyset$ and $e_{I_i} \wedge e_{J_j} = 0$. Let $\mathbf{A} = [a_{ij}]_{m \times m}$ and $\mathbf{B} = [b_{ij}]_{m \times m}$ be the corresponding integer matrices of $\wedge^r \hat{\alpha}$ and $\wedge^{n-r} \hat{\alpha}$ with respect to \mathcal{E}_r and \mathcal{E}_{n-r} , respectively. So $\wedge^r \hat{\alpha}(e_{I_i}) = \sum_{p=1}^m a_{pi} e_{I_p}$ and $\wedge^{n-r} \hat{\alpha}(e_{J_j}) = \sum_{q=1}^m b_{qj} e_{J_q}$. What we want to show is that $\mathbf{A} - \mathbf{I}$ is equivalent to $\mathbf{B} - \mathbf{I}$. We have

$$\wedge^n \hat{\alpha}(e_{I_i} \wedge e_{J_j}) = (-1)^{\ell(I_i) - \frac{r(r+1)}{2}} \delta_{ij} \omega = \wedge^r \hat{\alpha}(e_{I_i}) \wedge \wedge^{n-r} \hat{\alpha}(e_{J_j}) = \sum_{p,q=1}^m a_{pi} b_{qj} (-1)^{\ell(I_p) - \frac{r(r+1)}{2}} \delta_{pq} \omega.$$

Therefore one obtains

$$(6) \quad \sum_{k=1}^m (-1)^{\ell(I_k) - \ell(I_i)} a_{ki} b_{kj} = \delta_{ij}.$$

Now, if we set $c_{ij} := (-1)^{\ell(I_j) - \ell(I_i)} a_{ji}$ and $C := [c_{ij}]_{m \times m}$, then $c_{ij} - \delta_{ij} = (-1)^{\ell(I_j) - \ell(I_i)} (a_{ji} - \delta_{ji})$. Therefore $C - I$ is obtained from $A - I$ by changing rows (and columns) and occasionally multiplying some rows (and columns) by -1 . This means that $C - I$ is equivalent to $A - I$. On the other hand, the equation (6) means that $CB = I$. So $C - I = C(B - I)(-I)$ and $B - I$ is also equivalent to $C - I$. Consequently, $A - I$ is equivalent to $B - I$. \square

Corollary 4.2. *If $\det \hat{\alpha} = 1$, then $\text{rank ker}(\wedge^r \hat{\alpha} - \text{id}) = \text{rank ker}(\wedge^{n-r} \hat{\alpha} - \text{id})$. In particular, using Notation 3.2, we have $a_{n,r} = a_{n,n-r}$ for $r = 0, 1, \dots, n$.*

We are now ready to apply our Poincaré duality to a the following K -theoretic result.

Theorem 4.3. *Let $A := \mathcal{C}(\mathbb{T}^{2m-1}) \rtimes_{\alpha} \mathbb{Z}$ be such that the corresponding homomorphism $\hat{\alpha}$ satisfies $\det \hat{\alpha} = 1$. Then $K_0(A) \cong K_1(A)$ as abelian groups, and the (common) rank of the K -groups of A is an even number. In particular, for every Furstenberg transformation group C^* -algebra $\mathcal{F}_{\theta, \mathbf{f}}$ based on an odd-dimensional torus (e.g. $\mathcal{A}_{2m-1, \theta}$), one has $K_0(\mathcal{F}_{\theta, \mathbf{f}}) \cong K_1(\mathcal{F}_{\theta, \mathbf{f}})$.*

Proof. Combining Theorem 2.1 and Proposition 4.1, one obtains

$$K_0(A) \cong K_1(A) \cong \bigoplus_{k=0}^{m-1} [\text{coker}(\wedge^k \hat{\alpha} - \text{id}) \oplus \text{ker}(\wedge^k \hat{\alpha} - \text{id})].$$

As a result, the rank of the K -groups of A is an even number since the ranks of the cokernel and kernel of an endomorphism coincide. Note that for $\mathcal{F}_{\theta, \mathbf{f}}$ the corresponding integer matrix of $\hat{\alpha}$ is an upper triangular matrix with 1's on the diagonal. Thus $\det \hat{\alpha} = 1$. \square

5. THE RANK a_n OF THE K -GROUPS OF $\mathcal{A}_{n, \theta}$

In this section, we study some general properties of a_n , the (common) rank of the K -groups of Anzai transformation group C^* -algebras based on \mathbb{T}^n . We specify a family of C^* -algebras, whose ranks of K -groups are given by the same sequence $\{a_n\}$. As an application, we characterize the rank of the K -groups of Furstenberg transformation group C^* -algebras $\mathcal{F}_{\theta, \mathbf{f}}$. In Appendix F, this study will have some applications to the classification of simple infinite dimensional quotients of the Heisenberg-type group C^* -algebras $C^*(\mathfrak{D}_n)$, which were studied in an earlier work [33]. We remind the reader of some linear algebraic properties of nilpotent and unipotent matrices in Appendix B.

We compare the ranks of the K -groups of a class of C^* -algebras of the form $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ in the following theorem, which shows that the rank a_n of the K -groups of $\mathcal{A}_{n, \theta}$ is somehow generic.

Theorem 5.1. *Let $A = \mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$, in which α is a homeomorphism of \mathbb{T}^n , whose corresponding integer matrix $A \in \text{GL}(n, \mathbb{Z})$ is unipotent of maximal degree (i.e. $\deg(A) = n$). Then*

$$\text{rank } K_0(A) = \text{rank } K_1(A) = a_n = \text{rank } K_0(\mathcal{A}_{n, \theta}) = \text{rank } K_1(\mathcal{A}_{n, \theta}).$$

In particular, the rank of the K -groups of any Furstenberg transformation group C^ -algebra $\mathcal{F}_{\theta, \mathbf{f}} = \mathcal{C}(\mathbb{T}^n) \rtimes_{\varphi_{\theta, \mathbf{f}}} \mathbb{Z}$ is equal to the rank of the K -groups of $\mathcal{A}_{n, \theta}$, namely, to a_n .*

Proof. Let $\hat{\alpha}$ denote the restriction of α_* to \mathbb{Z}^n and A be the corresponding matrix of $\hat{\alpha}$ acting on \mathbb{Z}^n . Also, let S_n be the corresponding matrix for $\mathcal{A}_{n, \theta}$ as denoted in Section 3. Since A is unipotent of maximal degree by assumption, and S_n is unipotent of maximal degree too, the matrices A and S_n are similar over \mathbb{C} (see Corollary B.4). In fact, the Jordan normal form of $A - I$ is precisely $S_n - I$. On the other hand, we know by Corollary 2.2 that the rank of the K -groups of A is equal to $\text{rank ker}(\wedge^* A - I)$. Note that by the Smith normal form theorem (see Theorem A.2), $\text{rank ker}(\wedge^* A - I) = \dim_{\mathbb{C}} \text{ker}(\wedge^* A - I)$. The similarity of A and S_n implies the similarity of $\wedge^* A - I$ and $\wedge^* S_n - I$ as matrices acting on $\wedge^* \mathbb{C}^n$. So $\dim_{\mathbb{C}} \text{ker}(\wedge^* A - I) = \dim_{\mathbb{C}} \text{ker}(\wedge^* S_n - I) = a_n$, which yields the result.

For the second part, note that the corresponding integer matrix of a Furstenberg transformation $\varphi_{\theta, \mathbf{f}}$ on \mathbb{T}^n is of the form

$$(\heartsuit) \quad \begin{pmatrix} 1 & b_{12} & b_{13} & \cdots & b_{1n} \\ 0 & 1 & b_{23} & & \vdots \\ 0 & \ddots & \ddots & \ddots & b_{n-2,n} \\ \vdots & & 0 & 1 & b_{n-1,n} \\ 0 & \cdots & 0 & 0 & 1 \end{pmatrix}_{n \times n}$$

which is unipotent of maximal degree since $b_{i,i+1} \neq 0$ for $i = 1, \dots, n-1$ (see Definition 1.3 and Example B.5). Now, the proof of the first part yields the result. \square

Remark 5.2. In the preceding theorem, the basis for \mathbb{Z}^n for the matrices involved is $\{e_1, \dots, e_n\}$, where $e_i := [z_i]_1$ as introduced at the beginning of Section 2. It is interesting to know that if $\hat{\alpha}$ is an arbitrary unipotent automorphism of \mathbb{Z}^n , then there is a basis for \mathbb{Z}^n with respect to which the integer matrix \mathbf{A} of $\hat{\alpha}$ is of the form (\heartsuit) above (but not necessarily with $b_{i,i+1} \neq 0$ for $i = 1, \dots, n-1$, unless $\hat{\alpha}$ is of maximal degree) [12, Theorems 16 and 18]. The unipotency of $\hat{\alpha}$ has also important effects on the dynamics of the generated flow on \mathbb{T}^n . For example, if α is an affine transformation on \mathbb{T}^n and $\hat{\alpha}$ is unipotent, then the dynamical system (\mathbb{T}^n, α) has quasi-discrete spectrum [12, Theorem 19]. More generally, let $\alpha = (\mathbf{t}, \mathbf{A})$ be an affine transformation on \mathbb{T}^n and take $Z_p(\mathbf{A}) = \ker(\mathbf{A}^p - \text{id}) \subset \mathbb{Z}^n$ for $p \in \mathbb{N}$ and consider the following conditions

- (1) $Z_1(\mathbf{A}) = Z_p(\mathbf{A}), \forall p \in \mathbb{N}$,
- (2) \mathbf{t} is rationally independent over $Z_1(\mathbf{A})$, i.e. if $\mathbf{k} = (k_1, \dots, k_n) \in Z_1(\mathbf{A})$ is such that $\langle \mathbf{t}, \mathbf{k} \rangle := \sum_{j=1}^n t_j k_j$ is a rational number, then $\mathbf{k} = \mathbf{0}$.
- (3) $Z_1(\mathbf{A}) \neq \{0\}$,
- (4) \mathbf{A} is unipotent.

Then (\mathbb{T}^n, α) is ergodic with respect to Haar measure if and only if α satisfies the conditions (1) and (2) [12]. Moreover, if α satisfies the conditions (1) through (4), then the dynamical system (\mathbb{T}^n, α) is minimal, uniquely ergodic with respect to Haar measure, and has quasi-discrete spectrum. Conversely, any minimal transformation on \mathbb{T}^n with topologically quasi-discrete spectrum is conjugate to an affine transformation which must satisfy the conditions (1) through (4) [13]. The C^* -algebras corresponding to such actions are therefore simple and have a unique tracial state.

6. COMBINATORIAL PROPERTIES OF THE SEQUENCE $\{a_n\}$

As mentioned before, one of our main goals is to describe a_n as the rank of the K -groups of $\mathcal{A}_{n,\theta}$. Since $a_n = \sum_{r=0}^n a_{n,r}$, it makes sense to first study $a_{n,r}$. So we begin by finding some combinatorial properties of $a_{n,r}$, which is the rank of $\ker(\wedge^r \hat{\sigma} - \text{id})$ for $r = 0, 1, \dots, n$, where $\hat{\sigma}$ is the automorphism of \mathbb{Z}^n corresponding to the Anzai transformation σ on \mathbb{T}^n , and is represented by the integer matrix \mathbf{S}_n as in the beginning of Section 3. In fact, we will show that $a_{n,r}$ equals the number of partitions of $[r(n+1)/2]$ to r distinct positive integers not greater than n . To do this, we will use properties of the irreducible representations of the simple Lie algebra $\mathfrak{sl}(2, \mathbb{C})$.

6.1. Connections with representation theory of $\mathfrak{sl}(2, \mathbb{C})$. The automorphism $\hat{\sigma}$ is realized through its action on the basis $\{e_1, \dots, e_n\}$ of \mathbb{Z}^n , where $e_i := [z_i]$ for $i = 1, \dots, n$ as in Section 2, and we have $\hat{\sigma}(e_i) = e_i + e_{i-1}$ with $e_0 := 0$. Therefore introducing a new endomorphism of \mathbb{Z}^n by $\hat{\varphi} := \hat{\sigma} - \text{id}$, we will get

$$\hat{\varphi}(e_i) = e_{i-1}.$$

This is precisely a relation that may be recognized as part of the data of the canonical representation π_n of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ on a complex vector space V with basis $\{e_1, \dots, e_n\}$. More precisely, we have

$$\hat{\varphi} = \pi_n(f),$$

where the canonical representation π_n and the (third) basis element $f \in \mathfrak{sl}(2, \mathbb{C})$ are defined in Appendix D. The endomorphism $\hat{\varphi}$ induces a derivation on $\Lambda^r V$, which is defined by

$$\mathcal{D}^r \hat{\varphi}(x_1 \wedge \dots \wedge x_r) = \sum_{i=1}^r x_1 \wedge \dots \wedge \hat{\varphi}(x_i) \wedge \dots \wedge x_r$$

for $r = 2, \dots, n$ and $x_i \in V$, and by setting $\mathcal{D}^0 \hat{\varphi} := 0$, and $\mathcal{D}^1 \hat{\varphi} := \hat{\varphi}$ (see Appendix C). Then the following result states that $a_{n,r}$ equals the nullity of the linear mapping $\mathcal{D}^r \hat{\varphi}$.

Proposition 6.1. *Let σ be an Anzai transformation on \mathbb{T}^n and σ_* be the corresponding induced homomorphism on $K_*(\mathcal{C}(\mathbb{T}^n)) = \Lambda^* \mathbb{Z}^n$. Let $\hat{\sigma}$ be the restriction of σ_* to \mathbb{Z}^n and consider the linear mapping $\hat{\sigma} \otimes 1$ on $V := \mathbb{Z}^n \otimes \mathbb{C}$. Take $\hat{\varphi} = \hat{\sigma} \otimes 1 - \text{id}$ and $\mathcal{D}^r \hat{\varphi}$ as above. Then*

$$a_{n,r} = \text{rank ker}(\wedge^r \hat{\sigma} - \text{id}) = \dim \ker \mathcal{D}^r \hat{\varphi}.$$

Proof. Since $\hat{\varphi}$ is a nilpotent mapping, we can use Corollary C.4 to conclude that $\wedge^r(\hat{\sigma} \otimes 1) - \text{id} \sim \mathcal{D}^r \hat{\varphi}$. Therefore

$$\text{rank ker}(\wedge^r \hat{\sigma} - \text{id}) = \dim \ker(\wedge^r(\hat{\sigma} \otimes 1) - \text{id}) = \dim \ker \mathcal{D}^r \hat{\varphi}.$$

□

Notation 6.2. Let n, k, r be positive integers. Then $P(n, r, k)$ denotes the number of partitions of k to r distinct positive integers not greater than n . In other words

$$P(n, r, k) = \text{card}\{(i_1, \dots, i_r) \mid i_1 + \dots + i_r = k, 1 \leq i_1 < \dots < i_r \leq n\}.$$

By convention, we set $P(n, 0, 0) = 1$ and $P(n, r, 0) = P(n, 0, k) = 0$ for $r, k \geq 1$.

We are ready now to state the main result of this section.

Theorem 6.3. *With the above notation, $a_{n,r} = P(n, r, \lceil r(n+1)/2 \rceil)$, where $\lceil x \rceil$ denotes the greatest integer not greater than x . In particular,*

$$a_n = \sum_{r=0}^n P(n, r, \lceil \frac{r(n+1)}{2} \rceil).$$

Proof. Let $\pi_n : \mathfrak{sl}(2, \mathbb{C}) \rightarrow \mathfrak{gl}(V)$ be the canonical representation of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ on the n -dimensional complex vector space V , and extend π_n to $\pi_n^r : \mathfrak{sl}(2, \mathbb{C}) \rightarrow \mathfrak{gl}(\Lambda^r V)$ with $\pi_n^1 = \pi_n$. More precisely, for every $X \in \mathfrak{sl}(2, \mathbb{C})$ define

$$\pi_n^r(X)(v_1 \wedge \dots \wedge v_r) = (\pi_n(X)v_1) \wedge v_2 \wedge \dots \wedge v_r + \dots + v_1 \wedge \dots \wedge v_{r-1} \wedge (\pi_n(X)v_r).$$

This means that we have $\mathcal{D}^r \hat{\varphi} = \pi_n^r(f)$. In particular, $a_{n,r}$ is the nullity of $\pi_n^r(f)$ by the previous proposition. Following Weyl's theorem (see Theorem D.1), since the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ is semisimple the representation π_n^r has to be completely reducible. This means we should have a decomposition $\Lambda^r V = \bigoplus_{p=1}^N W_p$, where W_p 's are some π_n^r -invariant irreducible subspaces of $\Lambda^r V$. Moreover, the number N of such subspaces is equal to $\dim E_0 + \dim E_1$, where

$$E_j = \{v \in \Lambda^r V \mid \pi_n^r(h)v = jv\}, \quad (j = 0, 1)$$

and h is the first basis element of $\mathfrak{sl}(2, \mathbb{C})$ as in Appendix D (see Theorem D.3). On the other hand, the number N is equal to the nullity of $\pi_n^r(f)$. In fact, since $\pi_n^r|_{W_p}$ is an irreducible representation of $\mathfrak{sl}(2, \mathbb{C})$ on W_p , it is equivalent to the canonical representation of $\mathfrak{sl}(2, \mathbb{C})$ on W_p by Theorem D.3. But the image of f in the canonical representation has a 1-dimensional kernel due to the part (c) of Proposition D.2. So the nullity of $\pi_n^r(f)$ counts the number of W_p 's. Therefore

$$a_{n,r} = \dim \ker \pi_n^r(f) = \dim E_0 + \dim E_1.$$

To compute the last two terms, note that using Proposition D.2 we have $\pi_n(h)e_i = (2i - n - 1)e_i$, which leads to

$$\pi_n^r(h)(e_{i_1} \wedge \dots \wedge e_{i_r}) = (2(i_1 + \dots + i_r) - r(n+1))e_{i_1} \wedge \dots \wedge e_{i_r}.$$

So for even $r(n+1)$ we have $E_1 = \{0\}$ and $\dim E_0 = P(n, r, r(n+1)/2)$, and for odd $r(n+1)$ we have $E_0 = \{0\}$ and $\dim E_1 = P(n, r, r(n+1)/2 - 1)$. To summarize, we have established the following equalities

$$a_{n,r} = \dim \ker \mathcal{D}^r \hat{\varphi} = \dim \ker \pi_n^r(f) = N = \dim E_0 + \dim E_1 = P(n, r, [r(n+1)/2]).$$

The desired formula for a_n is immediate now by writing $a_n = \sum_{r=0}^n a_{n,r}$. \square

Using the previous theorem, we can prove that $\{a_n\}$ is a strictly increasing sequence. We need a lemma first.

Lemma 6.4. $P(n+1, r, k+s) \geq P(n, r, k)$ for $s = 0, 1, \dots, r$.

Proof. For $s = 0$, the proof is clear. Now, let $1 \leq s \leq r$ and suppose that (j_1, \dots, j_r) is a partition of k such that $1 \leq j_1 < \dots < j_r \leq n$. Define $i_q := j_q$ for $1 \leq q \leq r-s$ and $i_q := j_q + 1$ for $r-s+1 \leq q \leq r$. Then (i_1, \dots, i_r) is a partition of $k+s$ and $1 \leq i_1 < \dots < i_r \leq n+1$. Thus $P(n+1, r, k+s) \geq P(n, r, k)$. \square

Proposition 6.5. $\{a_n\}$ is a strictly increasing sequence.

Proof. First, note that $a_{n,0} = a_{n,n} = P(n, 0, 0) = P(n, n, n(n+1)/2) = 1$, and from the previous theorem we have $a_n = \sum_{r=0}^n P(n, r, [r(n+1)/2])$. Fix $m \in \mathbb{N}$, and get

$$\begin{aligned} a_{2m+1} &= 1 + \sum_{r=0}^m P(2m+1, 2r, 2rm+2r) + \sum_{r=0}^{m-1} P(2m+1, 2r+1, 2rm+2r+m+1), \\ a_{2m} &= \sum_{r=0}^m P(2m, 2r, 2rm+r) + \sum_{r=0}^{m-1} P(2m, 2r+1, 2rm+m+r) \\ &= 1 + \sum_{r=0}^{m-1} P(2m, 2r, 2rm+r) + \sum_{r=0}^{m-1} P(2m, 2r+1, 2rm+m+r), \\ a_{2m-1} &= \sum_{r=0}^{m-1} P(2m-1, 2r, 2rm) + \sum_{r=0}^{m-1} P(2m-1, 2r+1, 2rm+m). \end{aligned}$$

Applying the previous lemma to the terms of the sums expressed above implies that

$$a_{2m+1} > a_{2m} > a_{2m-1}.$$

\square

6.2. Generating functions for the sequence $\{a_n\}$. In this part, we express the rank of the K -groups of $\mathcal{A}_{n,\theta}$ as explicitly as possible. In fact, we present them as the constant terms in the polynomial expansions of certain functions. First of all, we need the following basic lemma.

Lemma 6.6. Let $P(n, r, k)$ denote the number of partitions of k to r distinct positive integers not greater than n . Then $P(n, r, k)$ is the coefficient of $u^r t^k$ in the polynomial expansion of $F_n(u, t) := \prod_{i=1}^n (1 + ut^i)$. In other words,

$$\sum_{r,k \geq 0} P(n, r, k) u^r t^k = \prod_{i=1}^n (1 + ut^i).$$

Proof.

$$\prod_{i=1}^n (1 + ut^i) = 1 + \sum_{r=1}^n \sum_{\substack{(i_1, \dots, i_r) \\ 1 \leq i_1 < \dots < i_r \leq n}} (ut^{i_1}) \dots (ut^{i_r}) = 1 + \sum_{r=1}^n \sum_{k \geq 1} P(n, r, k) u^r t^k = \sum_{r,k \geq 0} P(n, r, k) u^r t^k.$$

\square

Now, we have the following result for the rank a_n of the K -groups of $\mathcal{A}_{n,\theta}$.

Theorem 6.7. *Let $a_n = \text{rank } K_0(\mathcal{A}_{n,\theta}) = \text{rank } K_1(\mathcal{A}_{n,\theta})$. Then for a nonnegative integer m we have*

(i) a_{2m+1} is the constant term in the Laurent polynomial expansion of

$$\prod_{j=-m}^m (1 + z^j),$$

(ii) a_{2m} is the constant term in the Laurent polynomial expansion of

$$(1 + z) \prod_{j=-m+1}^m (1 + z^{2j-1}).$$

Proof. We know that $a_n = \sum_{r=0}^n a_{n,r}$ and $a_{n,r} = P(n, r, [r(n+1)/2])$ by Theorem 6.3. We have $a_{2m+1} = \sum_{r=0}^{2m+1} P(2m+1, r, r(m+1))$. Now, take $y = ut^{m+1}$ and use the preceding lemma to get

$$F_{2m+1}(u, t) = F_{2m+1}(yt^{-m-1}, t) = \prod_{i=1}^{2m+1} (1 + yt^{i-m-1}) = \sum_{r,k \geq 0} P(2m+1, r, k) y^r t^{k-r(m+1)}.$$

In particular, we get the following identity for $y = 1$

$$\prod_{i=1}^{2m+1} (1 + t^{i-m-1}) = \sum_{r,k \geq 0} P(2m+1, r, k) t^{k-r(m+1)},$$

or equivalently, by setting $z = t$ and $j = i - m - 1$ we have

$$\prod_{j=-m}^m (1 + z^j) = \sum_{r,k \geq 0} P(2m+1, r, k) z^{k-r(m+1)}.$$

In particular, the constant term in the Laurent polynomial expansion of $\prod_{j=-m}^m (1 + z^j)$ is obtained when we take the sum of those terms for which $k = r(m+1)$ holds, namely

$$\sum_{r=0}^{2m+1} P(2m+1, r, r(m+1)),$$

which is precisely the expression for a_{2m+1} .

For part (ii), write

$$\begin{aligned} a_{2m} &= \sum_{r=0}^{2m} P(2m, r, [r(m + \frac{1}{2})]) = \sum_{r=0}^m P(2m, 2r, r(2m+1)) + \sum_{r=0}^{m-1} P(2m, 2r+1, 2rm+m+r) \\ &=: A_m + B_m. \end{aligned}$$

Let us determine A_m first. Note that using the preceding lemma we have

$$\frac{1}{2} \left\{ \prod_{i=1}^{2m} (1 + ut^i) + \prod_{i=1}^{2m} (1 - ut^i) \right\} = \sum_{r,k \geq 0} P(2m, r, k) \left\{ \frac{1 + (-1)^r}{2} \right\} u^r t^k = \sum_{r,k \geq 0} P(2m, 2r, k) u^{2r} t^k.$$

If we define $y := u^2 t^{2m+1}$, we have the following identity

$$\frac{1}{2} \left\{ \prod_{i=1}^{2m} (1 + y^{\frac{1}{2}} t^{i-(m+\frac{1}{2})}) + \prod_{i=1}^{2m} (1 - y^{\frac{1}{2}} t^{i-(m+\frac{1}{2})}) \right\} = \sum_{r,k \geq 0} P(2m, 2r, k) y^r t^{k-r(2m+1)},$$

which for $y = 1$ yields

$$\frac{1}{2} \left\{ \prod_{i=1}^{2m} (1 + t^{i-(m+\frac{1}{2})}) + \prod_{i=1}^{2m} (1 - t^{i-(m+\frac{1}{2})}) \right\} = \sum_{r,k \geq 0} P(2m, 2r, k) t^{k-r(2m+1)}.$$

Hence A_m is the constant term in the polynomial expansion of

$$\frac{1}{2} \left\{ \prod_{i=1}^{2m} (1 + t^{i-(m+\frac{1}{2})}) + \prod_{i=1}^{2m} (1 - t^{i-(m+\frac{1}{2})}) \right\}.$$

Similarly, for B_m we have

$$\frac{1}{2} \left\{ \prod_{i=1}^{2m} (1 + ut^i) - \prod_{i=1}^{2m} (1 - ut^i) \right\} = \sum_{r,k \geq 0} P(2m, r, k) \left\{ \frac{1 - (-1)^r}{2} \right\} u^r t^k = \sum_{r,k \geq 0} P(2m, 2r+1, k) u^{2r+1} t^k.$$

If we define $y^2 := u^2 t^{2m+1}$, we have the following identities

$$\begin{aligned} \frac{1}{2} \left\{ \prod_{i=1}^{2m} (1 + y^{\frac{1}{2}} t^{i-(m+\frac{1}{2})}) - \prod_{i=1}^{2m} (1 - y^{\frac{1}{2}} t^{i-(m+\frac{1}{2})}) \right\} \\ = \sum_{r,k \geq 0} P(2m, 2r+1, k) y^{2r+1} t^{k-(2rm+r+m)-\frac{1}{2}} \\ = t^{-\frac{1}{2}} \sum_{r,k \geq 0} P(2m, 2r+1, k) y^{2r+1} t^{k-(2rm+r+m)}, \end{aligned}$$

which for $y = 1$ yields

$$\frac{t^{\frac{1}{2}}}{2} \left\{ \prod_{i=1}^{2m} (1 + t^{i-(m+\frac{1}{2})}) - \prod_{i=1}^{2m} (1 - t^{i-(m+\frac{1}{2})}) \right\} = \sum_{r,k \geq 0} P(2m, 2r+1, k) t^{k-(2rm+r+m)}.$$

Hence B_m is the constant term in the polynomial expansion of

$$\frac{t^{\frac{1}{2}}}{2} \left\{ \prod_{i=1}^{2m} (1 + t^{i-(m+\frac{1}{2})}) - \prod_{i=1}^{2m} (1 - t^{i-(m+\frac{1}{2})}) \right\}.$$

Therefore $a_{2m} = A_m + B_m$ is the constant term in the polynomial expansion of

$$\frac{1}{2} \left\{ \prod_{i=1}^{2m} (1 + t^{i-(m+\frac{1}{2})}) + \prod_{i=1}^{2m} (1 - t^{i-(m+\frac{1}{2})}) + \frac{t^{\frac{1}{2}}}{2} \prod_{i=1}^{2m} (1 + t^{i-(m+\frac{1}{2})}) - \frac{t^{\frac{1}{2}}}{2} \prod_{i=1}^{2m} (1 - t^{i-(m+\frac{1}{2})}) \right\},$$

or equivalently, the constant term in the polynomial expansion of

$$\frac{1}{2} \left\{ (1+z) \prod_{i=1}^{2m} (1 + z^{2i-(2m+1)}) + (1-z) \prod_{i=1}^{2m} (1 - z^{2i-(2m+1)}) \right\},$$

which equals the constant term in the Laurent polynomial expansion of

$$(1+z) \prod_{j=-m+1}^m (1 + z^{2j-1}).$$

□

Thanks to this theorem, one can compute a_n for large values of n using a computer algebra program. Many more terms are also available online at OEIS (The Online Encyclopedia of Integer Sequences at www.oeis.org). Moreover, as the following corollaries suggest, such recognitions as constant terms of certain Laurent polynomials opens the door to finding even more interesting combinatorial properties of the sequence $\{a_n\}$, which have been of interest to Erdős, J. H. van Lint and R. C. Entringer to name a few (*cf.* [6, 23, 5]).

Corollary 6.8. *Let n be a nonnegative integer.*

(i) The integer a_{2n+1} is the number of solutions of the equation

$$\sum_{k=-n}^{k=n} k \epsilon_k = 0,$$

where $\epsilon_k = 0$ or 1 for $-n \leq k \leq n$. In other words, a_{2n+1} is the number of ways that a sum of integers between $-n$ and n (with no repetitions) equals to 0.

(ii) The integer a_{2n} is the number of solutions of the equation

$$\sum_{k=-n+1}^{k=n} (2k-1) \epsilon_k = 0 \text{ or } 1,$$

where $\epsilon_k = 0$ or 1 for $-n+1 \leq k \leq n$. In other words, a_{2n} is the number of ways that a sum of half-integers between $-n+1/2$ and $n-1/2$ (with no repetitions) equals to 0 or $1/2$.

Proof. Using Theorem 6.7, the number a_{2n+1} is the constant term in the Laurent polynomial expansion of $\prod_{k=-n}^n (1+z^k)$, which is a finite sum of the form $\sum A(n,m)z^m$. Obviously, the integer coefficient $A(n,m)$ is the number of all possible combinations from the terms $z^{-n}, \dots, z^0, \dots, z^n$, whose product makes a z^m . In other words, by putting $\epsilon_k = 1$ when z^k contributes to such a product making a z^m , and $\epsilon_k = 0$ otherwise, we conclude that

$$A(n,m) = \#\{(\epsilon_{-n}, \dots, \epsilon_0, \dots, \epsilon_n) \in \{0,1\}^{2n+1} : \sum_{k=-n}^{k=n} k \epsilon_k = m\}.$$

In particular, the constant term of the Laurent polynomial expansion is $A(n,0)$, and we have $a_{2n+1} = A(n,0)$. This proves part (i). For part (ii), we use the same idea for the Laurent polynomial expansion of

$$(1+z) \prod_{k=-m+1}^m (1+z^{2k-1}) = \prod_{k=-m+1}^m (1+z^{2k-1}) + z \prod_{k=-m+1}^m (1+z^{2k-1})$$

as suggested by part (ii) of Theorem 6.7. □

J. H. van Lint in [23] answered a question of Erdős by determining the asymptotic behavior of

$$A(n,0) = \#\{(\epsilon_{-n}, \dots, \epsilon_0, \dots, \epsilon_n) \in \{0,1\}^{2n+1} : \sum_{k=-n}^{k=n} k \epsilon_k = 0\}.$$

The idea in his proof is as follows. Since $A(n,0)$ is the constant term of the Laurent polynomial expansion of $\prod_{k=-n}^n (1+z^k)$, we can compute it as the Cauchy integral

$$\frac{1}{2\pi i} \oint_C \frac{\prod_{k=-n}^n (1+z^k)}{z} dz,$$

where C denotes the unit circle. By parameterizing C by $z = e^{2ix}$ for $x \in [0, \pi]$, applying the elementary identity $(1 + e^{2ikx})(1 + e^{-2ikx}) = 4 \cos^2 kx$, and a simple calculation we arrive at

$$A(n,0) = \frac{2^{2n+2}}{\pi} \int_0^{\frac{\pi}{2}} \prod_{k=1}^n \cos^2 kx dx.$$

We can then proceed by estimating the integrand near and far from 0 using some elementary inequalities, which lead to the asymptotic formula $A(n,0) \sim (3/\pi)^{\frac{1}{2}} 2^{2n+1} n^{-\frac{3}{2}}$ [23]. This will immediately give the asymptotic behavior of the sequence $\{a_{2n+1}\}$ by the previous corollary. One can adapt the arguments

used by J. H. van Lint to obtain a similar asymptotic behavior for the sequence $\{a_{2n}\}$ by estimating the corresponding integral

$$\frac{2^{2n+2}}{\pi} \int_0^{\frac{\pi}{2}} \cos^2 x \prod_{k=1}^n \cos^2(2k-1)x dx,$$

which leads to the asymptotic formula $a_{2n} \sim (3/\pi)^{\frac{1}{2}} 2^{2n} n^{-\frac{3}{2}}$. This gives rise to the following result.

Corollary 6.9. $a_n \sim \sqrt{\frac{24}{\pi}} 2^n n^{-\frac{3}{2}}$ when $n \rightarrow \infty$. In particular, $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 2$.

7. THE POSITIVE CONE OF $K_0(\mathcal{F}_{\theta, \mathbf{f}})$

In this section, we generalize a result of Kodaka on the order structure of the group K_0 of the crossed product by a Furstenberg transformation on the 2-torus [18, Theorem 5.2]. However, our approach is different, and follows the general guidelines of [30, Lemma 3.1]. We remind the reader that for a C^* -algebra A the positive cone of $K_0(A)$ is the set $K_0(A)_+ = \{[q] \in K_0(A) : q \in \mathcal{P}_\infty(A)\}$, where $\mathcal{P}_\infty(A)$ is the set of all projections in matrix algebras over A . Also, any positive trace τ on a C^* -algebra A induces a group homomorphism $\tau_* : K_0(A) \rightarrow \mathbb{R}$. As was indicated in the Introduction, when the Furstenberg transformation $\varphi_{\theta, \mathbf{f}}$ is minimal and uniquely ergodic, using the results of H. Lin and N. C. Phillips in [21] the transformation group C^* -algebra $\mathcal{F}_{\theta, \mathbf{f}}$ is classifiable by its Elliott invariant, and the order of $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ is determined by the unique tracial state τ on $\mathcal{F}_{\theta, \mathbf{f}}$ [22, 29]. The fact that $\tau_* K_0(\mathcal{F}_{\theta, \mathbf{f}}) = \mathbb{Z} + \mathbb{Z}\theta$ was first proved in the unpublished thesis of R. Ji [16]. However, we will study the effect of the trace on the order structure of K_0 using R. Exel's machinery of rotation numbers [7].

For a C^* -algebra A , we denote by $U_p(A)$ the set of unitary elements of $M_p(A)$. The following lemma is well known, but it is convenient to state and prove it for self-containment of the paper.

Lemma 7.1. *Let A and B be unital C^* -algebras and let $A \otimes B$ denote their minimal tensor product. Suppose that $u \in U_p(A)$ and $v \in U_q(B)$, and let $\phi : \mathcal{C}(\mathbb{T}^2) \rightarrow M_{pq}(A \otimes B)$ be the unique homomorphism mapping the coordinate unitaries $z_1, z_2 \in U(\mathcal{C}(\mathbb{T}^2))$ to the commuting unitaries $u \otimes 1_q, 1_p \otimes v \in U_{pq}(A \otimes B)$, respectively. Let $b(u, v) \in K_0(A \otimes B)$ denote the Bott element of u, v defined by $K_0(\phi)(b)$, where $\beta = [z_1] \wedge [z_2]$ is the Bott element in $K_0(\mathcal{C}(\mathbb{T}^2))$ so that $K_0(\mathcal{C}(\mathbb{T}^2)) = \mathbb{Z}[1] + \mathbb{Z}\beta$. Then $\tau_*(b(u, v)) = 0$ for any tracial state τ on $A \otimes B$.*

Proof. Since $\tau \circ \phi$ is a trace on \mathbb{T}^2 , there exists a Borel probability measure μ on \mathbb{T}^2 such that

$$(\tau \circ \phi)(f) = \int_{\mathbb{T}^2} f(x) d\mu(x), \quad f \in \mathcal{C}(\mathbb{T}^2).$$

Write $\beta = [p] - [q]$, where p, q are appropriate projections in some matrix algebra over $\mathcal{C}(\mathbb{T}^2)$, so we have

$$\tau_*(b(u, v)) = \tau_*(K_0(\phi)(\beta)) = (\tau \circ \phi)_*(\beta) = \int_{\mathbb{T}^2} \text{Tr}(p(x)) - \text{Tr}(q(x)) d\mu(x).$$

It is well known that for the Bott element b we have $\text{Tr}(p(x)) - \text{Tr}(q(x)) = 0$, namely, the projections $p(x)$ and $q(x)$ have the same rank for all $x \in \mathbb{T}^2$, and this common rank does not depend on x since \mathbb{T}^2 is connected (in fact, they are rank one projections). This can be proved either by a calculation of the traces of the projections $p(x)$ and $q(x)$ explicitly (cf. [1, p. 7]), or by using the naturality in the Künneth formula for \mathbb{T}^2 , which shows that the image under any point evaluation of β is zero. Briefly speaking, the map $x \mapsto \text{Tr}(p(x)) - \text{Tr}(q(x))$ belongs to $\mathcal{C}(\mathbb{T}^2, \mathbb{Z})$, so it has to assume a constant integer, which we call $\dim \beta$. In particular, $\dim \beta$ is invariant under the change of coordinate $(\zeta_1, \zeta_2) \mapsto (\zeta_1, \zeta_2^{-1})$, whereas the naturality of the Künneth homomorphism $\alpha_{1,1} : K_1(\mathcal{C}(\mathbb{T})) \otimes K_1(\mathcal{C}(\mathbb{T})) \rightarrow K_0(\mathcal{C}(\mathbb{T}^2))$, which maps $[z] \otimes [z]$ to β , implies that the Bott element β will transform into $-\beta$ under this change of coordinates since $[z] \otimes [z^{-1}] = [z] \otimes (-[z]) = -([z] \otimes [z])$. This means $\dim \beta = 0$. \square

We denote by u the unitary in $\mathcal{F}_{\theta, \mathbf{f}}$ implementing the action generated by the transformation $\varphi_{\theta, \mathbf{f}}$ on \mathbb{T}^n with irrational parameter θ , and by z_1 the unitary in $\mathcal{C}(\mathbb{T}^n)$ defined by $z_1(\zeta_1, \dots, \zeta_n) = \zeta_1$ as in Section 2. Then we have $uz_1u^{-1} = z_1 \circ \varphi_{\theta, \mathbf{f}}^{-1} = e^{2\pi i \theta} z_1$ so that $C^*(u, z_1) \cong A_\theta$, the irrational rotation algebra. Let $p_\theta \in C^*(u, z_1)$ be a Rieffel projection of trace θ as in [34]. It is obvious that $\tau_*([1]) = 1$. On the other hand, since the restriction of τ on the C^* -subalgebra $A_\theta \subseteq \mathcal{F}_{\theta, \mathbf{f}}$ has to be the unique tracial state on A_θ , we have $\tau_*([p_\theta]) = \theta$. The main result of this section will show that all the essential information about the order structure of $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ is encoded in the embedding of A_θ in $\mathcal{F}_{\theta, \mathbf{f}}$.

Theorem 7.2. *Let $\varphi_{\theta, \mathbf{f}}$ be a minimal uniquely ergodic Furstenberg transformation on \mathbb{T}^n with $\theta \in (0, 1)$ (e.g. when $\theta \in (0, 1) \setminus \mathbb{Q}$ and each f_i satisfies a uniform Lipschitz condition in ζ_i for $i = 1, \dots, n-1$). Let a_n and $\mathcal{T}_{\mathbf{f}}^0$ denote, respectively, the rank and the torsion subgroup of $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ so that $K_0(\mathcal{F}_{\theta, \mathbf{f}}) \cong \mathbb{Z}^{a_n} \oplus \mathcal{T}_{\mathbf{f}}^0$. Then the isomorphism of $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ with this group can be chosen in such a way that*

(i) *the unique tracial state τ on $\mathcal{F}_{\theta, \mathbf{f}}$ induces the map*

$$\tau_*(a[1] + b[p_\theta], \mathbf{c}, \mathbf{t}) = a + b\theta$$

on $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ for all $(a[1] + b[p_\theta], \mathbf{c}, \mathbf{t}) \in (\mathbb{Z}[1] + \mathbb{Z}[p_\theta]) \oplus \mathbb{Z}^{a_n-2} \oplus \mathcal{T}_{\mathbf{f}}^0 \cong \mathbb{Z}^{a_n} \oplus \mathcal{T}_{\mathbf{f}}^0$,

(ii) *the positive cone $K_0(\mathcal{F}_{\theta, \mathbf{f}})_+$ can be identified with*

$$\{(a[1] + b[p_\theta], \mathbf{c}, \mathbf{t}) \in (\mathbb{Z}[1] + \mathbb{Z}[p_\theta]) \oplus \mathbb{Z}^{a_n-2} \oplus \mathcal{T}_{\mathbf{f}}^0 : a + b\theta > 0\} \cup \{0\}.$$

Proof. The idea of the proof is to show that there exists a generating set for the finitely generated abelian group $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ including $[1]$ and $[p_\theta]$ such that the induced homomorphism τ_* vanishes at all generators, except for $[1]$ and $[p_\theta]$ for which we have $\tau_*([1]) = 1$ and $\tau_*([p_\theta]) = \theta$. Using Theorem 2.1 and setting $\alpha = \varphi_{\theta, \mathbf{f}}$ and $\alpha_j = K_j(\alpha)$ for $j = 1, 2$ we have

$$K_0(\mathcal{F}_{\theta, \mathbf{f}}) \cong \text{coker}(\alpha_0 - \text{id}) \oplus \ker(\alpha_1 - \text{id}) = \bigoplus_{r \geq 0} [\text{coker}(\wedge^{2r} \hat{\alpha} - \text{id}) \oplus \ker(\wedge^{2r+1} \hat{\alpha} - \text{id})],$$

where $\hat{\alpha}$ is the restriction of α_1 to the subgroup $\mathbb{Z}[z_1] + \dots + \mathbb{Z}[z_n]$ of $K_1(\mathcal{C}(\mathbb{T}^n))$ as in Section 2, and $z_j(\zeta, \dots, \zeta_n) = \zeta_j$ of $\mathcal{C}(\mathbb{T}^n)$ for $j = 1, \dots, n$. Note that by Definition 1.3, $\mathbf{f} = (f_1, \dots, f_{n-1})$ consists of continuous functions $f_{j-1} : \mathbb{T}^{j-1} \rightarrow \mathbb{T}$ for $j = 2, \dots, n$. First, we “linearize” each f_{j-1} by finding the unique “linear” function

$$(\zeta_1, \dots, \zeta_{j-1}) \mapsto \zeta_1^{b_{1j}} \dots \zeta_{j-1}^{b_{j-1,j}}, \quad (b_{j-1,j} \neq 0)$$

in the homotopy class of f_{j-1} . This allows us to calculate $\hat{\alpha}([z_j])$ by writing

$$\hat{\alpha}([z_1]) = [z_1 \circ \varphi_{\theta, \mathbf{f}}^{-1}] = [e^{2\pi i \theta} z_1] = [z_1],$$

$$\hat{\alpha}([z_j]) = [z_j \circ \varphi_{\theta, \mathbf{f}}^{-1}] = [f_{j-1}(z_1, \dots, z_{j-1})z_j] = [z_1^{b_{1j}} \dots z_{j-1}^{b_{j-1,j}} z_j] = b_{1j}[z_1] + \dots + b_{j-1,j}[z_{j-1}] + [z_j],$$

for $j = 2, \dots, n$. In other words, the integer matrix of $\hat{\alpha}$ with respect to the basis $\{[z_1], \dots, [z_n]\}$ of \mathbb{Z}^n is precisely in the form (♥) as in the proof of Theorem 5.1. Now, we can realize $\alpha_0 = \wedge^{\text{even}} \hat{\alpha}$ and $\alpha_1 = \wedge^{\text{odd}} \hat{\alpha}$ to calculate the K -groups of $\mathcal{F}_{\theta, \mathbf{f}}$ as in Section 2.

It is important to note that, by referring to the exact sequences (1), (2) and (3) in Section 1, the isomorphic image of $\text{coker}(\alpha_0 - \text{id})$ in $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ is precisely the image $\text{im } j_0$ of $K_0(\mathcal{C}(\mathbb{T}^n))$, and an isomorphic image of $\ker(\alpha_1 - \text{id})$ in $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ is obtained by finding the image of a splitting (injective) homomorphism $s : \ker(\alpha_1 - \text{id}) \rightarrow K_0(\mathcal{F}_{\theta, \mathbf{f}})$ for the exact sequence (2) so that $\partial \circ s = \text{id}$ on $\ker(\alpha_1 - \text{id})$. Any such splitting homomorphism is obtained as follows: fix a basis $\{\gamma_1, \dots, \gamma_q\}$ for the free finitely generated group $\ker(\alpha_1 - \text{id})$ of rank q , and find elements $\nu_1^{(0)}, \dots, \nu_q^{(0)} \in K_0(\mathcal{F}_{\theta, \mathbf{f}})$ such that $\partial \nu_j^{(0)} = \gamma_j$ for $j = 1, \dots, q$. Then define $s(\sum_j m_j \gamma_j) = \sum_j m_j \nu_j^{(0)}$ for $m_j \in \mathbb{Z}$. Clearly, $K_0(\mathcal{F}_{\theta, \mathbf{f}}) = \text{im } j_0 \oplus \text{im } s$.

Now, since $\wedge^0 \hat{\alpha} = \text{id}$ on $\Lambda^0 \mathbb{Z}^n = \mathbb{Z}$, and $\wedge^1 \hat{\alpha} = \hat{\alpha}$ on $\Lambda^1 \mathbb{Z}^n = \mathbb{Z}^n$, we can write the isomorphism

$$K_0(\mathcal{F}_{\theta, \mathbf{f}}) \cong \mathbb{Z} \oplus \ker(\hat{\alpha} - \text{id}) \oplus \bigoplus_{r \geq 1} [\ker(\wedge^{2r+1} \hat{\alpha} - \text{id}) \oplus \text{coker}(\wedge^{2r} \hat{\alpha} - \text{id})].$$

In fact, a single generator for the isomorphic image of \mathbb{Z} in $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ is $[1]$, and since $b_{j-1, j} \neq 0$ for $j = 2, \dots, n$ we have $\ker(\hat{\alpha} - \text{id}) = \mathbb{Z}e_1 = \mathbb{Z}[z_1]$. It is easy to see that $\partial([p_\theta]) = [z_1]$ (see the proposition in the appendix of [31]). Therefore there exists a basis $\{\gamma_1, \dots, \gamma_q\}$ for $\ker(\alpha_1 - \text{id}) = \bigoplus_{r \geq 0} \ker(\wedge^{2r+1} \hat{\alpha} - \text{id})$ with $\gamma_1 = [z_1]$ and a splitting homomorphism $s : \ker(\alpha_1 - \text{id}) \rightarrow K_0(\mathcal{F}_{\theta, \mathbf{f}})$ with $s([z_1]) = [p_\theta]$. Hence a single generator for the image of $\ker(\hat{\alpha} - \text{id})$ in $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ is $[p_\theta]$.

It remains to study the effect of τ_* on the isomorphic image of $\bigoplus_{r \geq 1} \text{coker}(\wedge^{2r} \hat{\alpha} - \text{id})$, which contains the torsion subgroup $\mathcal{T}_{\mathbf{f}}^0$, and the image of $\bigoplus_{r \geq 1} \ker(\wedge^{2r+1} \hat{\alpha} - \text{id})$ in $K_0(\mathcal{F}_{\theta, \mathbf{f}})$. For more convenience, set $e_j := [z_j]$ for $j = 1, \dots, n$ as in Section 2. First, we study the isomorphic image of $\bigoplus_{r \geq 1} \text{coker}(\wedge^{2r} \hat{\alpha} - \text{id})$. We show that τ_* vanishes on this whole subgroup by showing, equivalently, that τ_* vanishes on the image of the subgroup $\bigoplus_{r \geq 1} \Lambda_{\mathbb{Z}}^{2r}(e_1, \dots, e_n) \subset K_0(\mathcal{C}(\mathbb{T}^n))$ in $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ under the map j_0 . Let $\eta = e_{i_1} \wedge \dots \wedge e_{i_{2r}} \in K_0(\mathcal{C}(\mathbb{T}^n))$ for some $r \geq 1$ and $1 \leq i_1 < \dots < i_{2r} \leq n$. We want to show that $\tau_*(j_0(\eta)) = 0$, where $j_0 := K_0(j)$ and $j : \mathcal{C}(\mathbb{T}^n) \rightarrow \mathcal{F}_{\theta, \mathbf{f}}$ is the natural embedding in the structure of the crossed product $\mathcal{F}_{\theta, \mathbf{f}} = \mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$. By Künneth formula we have $\eta = b(u, z)$, where u is a unitary in some matrix algebra over $\mathcal{C}(\mathbb{T}^{n-1})$ with $[u] = e_{i_1} \wedge \dots \wedge e_{i_{2r-1}} \in K_1(\mathcal{C}(\mathbb{T}^{n-1}))$ and z is the canonical unitary in $\mathcal{C}(\mathbb{T})$ with $[z] = e_{i_{2r}} \in K_1(\mathcal{C}(\mathbb{T}))$. By Lemma 7.1, we have

$$\tau_*(j_0(\eta)) = \tau_*(K_0(j)(\eta)) = (\tau \circ j)_*(\eta) = (\tau \circ j)_*(b(u, z)) = 0.$$

Now, we study the isomorphic image of $\bigoplus_{r \geq 1} \ker(\wedge^{2r+1} \hat{\alpha} - \text{id})$ in $K_0(\mathcal{F}_{\theta, \mathbf{f}})$. We will show that τ_* assumes only integer values on this whole subgroup. In other words, if $\{\gamma_1, \dots, \gamma_q\}$ is a basis for the $\ker(\alpha_0 - \text{id})$ as above such that $\gamma_1 = [z_1]$ is a basis for $\ker(\hat{\alpha} - \text{id})$ and $\{\gamma_2, \dots, \gamma_q\}$ is a basis for $\bigoplus_{r \geq 1} \ker(\wedge^{2r+1} \hat{\alpha} - \text{id})$, then $\tau_*(\nu_1^{(0)}) = \theta$, and $\tau_*(\nu_j^{(0)}) = k_j$ for some $k_j \in \mathbb{Z}$ for $j = 2, \dots, q$, where $\nu_j^{(0)}$'s are chosen in $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ so that $\partial(\nu_j^{(0)}) = \gamma_j$ for $j = 1, \dots, q$ and $\nu_1^{(0)} = [p_\theta]$ as above. To demonstrate this, we prove that the determinant of any unitary representing an element in the subgroup $\bigoplus_{r \geq 1} \Lambda_{\mathbb{Z}}^{2r+1}(e_1, \dots, e_n)$ is the constant function 1. Then the rotation number homomorphism $\rho_{\alpha}^{\mu} : \ker(\alpha_1 - \text{id}) \rightarrow \mathbb{T}$ defined by R. Exel is the constant 1 on the subgroup $\bigoplus_{r \geq 1} \ker(\wedge^{2r+1} \hat{\alpha} - \text{id})$ [7, Theorem VI.11], hence the trace will be integer-valued on this subgroup because $\exp(2\pi i \tau_*(\eta)) = \rho_{\alpha}^{\mu} \circ \partial(\eta)$ for all $\eta \in K_0(\mathcal{F}_{\theta, \mathbf{f}})$ [7, Theorem V.12]. To calculate the determinant on $\bigoplus_{r \geq 1} \Lambda_{\mathbb{Z}}^{2r+1}(e_1, \dots, e_n)$, let $\gamma = e_{i_1} \wedge \dots \wedge e_{i_{2r+1}} \in K_1(\mathcal{C}(\mathbb{T}^n))$, set $\eta = e_{i_1} \wedge \dots \wedge e_{i_{2r}} \in K_0(\mathcal{C}(\mathbb{T}^{n-1}))$, and write $\eta = [p] - [q]$ as above. Then $e_{i_{2r+1}} = [z]$ for the canonical unitary z of $\mathcal{C}(\mathbb{T})$, and using the Künneth formula we have

$$\gamma = \eta \otimes [z] = ([p] - [q]) \otimes [z] = [p] \otimes [z] + [q] \otimes [z^{-1}] = [((1-p) \otimes 1 + p \otimes z)((1-q) \otimes 1 + q \otimes z^{-1})]$$

So, $\gamma = [\omega_1 \omega_2]$, where $\omega_1 := (1-p) \otimes 1 + p \otimes z$ and $\omega_2 := (1-q) \otimes 1 + q \otimes z^{-1}$ are unitaries in some matrix algebra of the same size over $\mathcal{C}(\mathbb{T}^n)$. Since for all $x \in \mathbb{T}^{n-1}$ the projections $p(x)$ and $q(x)$ have the same rank ρ as in the proof of the previous lemma, we have the following equivalence of projections in some matrix algebra $M_l(\mathbb{C})$

$$p(x) \sim \mathbf{1}_{\rho} \oplus \mathbf{0}_{l-\rho} \sim q(x),$$

where $\mathbf{1}_m, \mathbf{0}_m$ denote the identity and the zero matrix of order m , respectively, and \oplus is the direct sum of matrices. This implies the following unitary equivalence of projections in $M_{2l}(\mathbb{C})$

$$p(x) \oplus \mathbf{0}_l \sim_u \mathbf{1}_{\rho} \oplus \mathbf{0}_{2l-\rho} \sim_u q(x) \oplus \mathbf{0}_l.$$

In particular, we conclude the following unitary equivalence of unitary matrices for all $(x, \zeta) \in \mathbb{T}^{n-1} \times \mathbb{T}$

$$\omega_1(x, \zeta) \oplus \mathbf{1}_l \sim_u \zeta \mathbf{1}_{\rho} \oplus \mathbf{1}_{2l-\rho}, \quad \omega_2(x, \zeta) \oplus \mathbf{1}_l \sim_u \zeta^{-1} \mathbf{1}_{\rho} \oplus \mathbf{1}_{2l-\rho}.$$

Therefore $\text{Det} \omega_1(x, \zeta) = \zeta^{\rho}$ and $\text{Det} \omega_2(x, \zeta) = \zeta^{-\rho}$, hence $\text{Det}(\omega_1 \omega_2)(x, \zeta) = 1$, for all $(x, \zeta) \in \mathbb{T}^{n-1} \times \mathbb{T}$. This implies that $\text{Det}_*(\gamma) = 1 \in [\mathbb{T}^n, \mathbb{T}]$, where $[\mathbb{T}^n, \mathbb{T}]$ denotes the set of homotopy classes of continuous functions from \mathbb{T}^n to \mathbb{T} (see Definition VI.8 and Proposition VI.9 of [7]).

Finally, by setting $\nu_1 := \nu_1^{(0)} = [p_\theta]$ and $\nu_k := \nu_j^{(0)} - k_j[1]$ for $j = 2, \dots, q$ so that $\tau_*(\nu_1) = \theta$ and $\tau_*(\nu_j) = 0$ for $j = 2, \dots, q$, we can form a generating set with the desired property for $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ by taking the union of $\{\nu_1, \dots, \nu_q\}$ and a generating set including $[1]$ for the isomorphic image of $\text{coker}(\alpha_0 - \text{id})$. This proves

part (i).

For part (ii), we use part (i) together with the fact that the order on $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ is determined by the effect of the unique tracial state τ because \mathbb{T}^n is a finite dimensional infinite compact metric space and $\varphi_{\theta, \mathbf{f}}$ is a minimal homeomorphism of \mathbb{T}^n (see Theorem 5.1(1) of [22] or Theorem 4.5(1) of [29]). \square

Corollary 7.3. *Let $\varphi_{\theta, \mathbf{f}}$ be a minimal uniquely ergodic Furstenberg transformation on \mathbb{T}^n as above. Then linearizing the functions $f_i : \mathbb{T}^i \rightarrow \mathbb{T}$ in $\mathbf{f} = (f_1, \dots, f_{i-1})$ does not change the isomorphism class of the transformation group C*-algebra $\mathcal{F}_{\theta, \mathbf{f}}$.*

Proof. Since $\varphi_{\theta, \mathbf{f}}$ is minimal, θ must be irrational. So the range of the unique tracial state (by unique ergodicity) on $K_0(\mathcal{F}_{\theta, \mathbf{f}})$ is dense in \mathbb{R} as it is $\mathbb{Z} + \mathbb{Z}\theta$ by the above argument. Benefiting from the results of [21], such C*-algebras are completely classifiable by their Elliott invariants, which remain unchanged (up to isomorphism) after the linearization process: linearizing does not change the isomorphism classes of the K-groups, and the previous theorem guarantees that the order structure of the group K_0 is precisely the regular order inherited from \mathbb{R} on $\mathbb{Z} + \mathbb{Z}\theta$ before and after linearization. \square

8. CONCLUDING REMARKS

I) The method used in Section 1 for computing K-groups of the transformation group C*-algebras of homeomorphisms of the tori may be extended to more general settings. Let G be a compact connected Lie group with torsion-free fundamental group $\pi_1(G)$. (It is well known that the fundamental group of such spaces are finitely generated and abelian, so being torsion-free means $\pi_1(G) \cong \mathbb{Z}^l$, for some l .) Some important examples are any finite Cartesian products of the groups S^3 , $SO(2)$, $Sp(n)$, $U(n)$ and $SU(n)$. Then $K^*(G)$ is torsion-free and can be given the structure of a \mathbb{Z}_2 -graded Hopf algebra over the integers [14]. Moreover, regarded as a Hopf algebra, $K^*(G)$ is the exterior algebra on the module of the primitive elements, which are of degree 1. The module of the primitive elements of $K^*(G)$ may also be described as follows. Let $U(n)$ denote the group of unitary matrices of order n and let $U := \cup_{n=1}^{\infty} U(n)$ be the stable unitary group. Any unitary representation $\rho : G \rightarrow U(n)$, by composition with the inclusion $U(n) \subset U$, defines a homotopy class $\beta(\rho)$ in $[G, U] = K^1(G)$. The module of the primitive elements in $K^1(G)$ is exactly the module generated by all classes $\beta(\rho)$ of this type. If in addition, G is semisimple and simply connected of rank l , there are l basic irreducible representations ρ_1, \dots, ρ_l , whose maximum weights $\lambda_1, \dots, \lambda_l$ form a basis for the character group $\hat{\mathbf{T}}$ of the maximal torus \mathbf{T} of G and the classes $\beta(\rho_1), \dots, \beta(\rho_l)$ form a basis for the module of the primitive elements in $K^1(G)$ and $K^*(G) = \Lambda^*(\beta(\rho_1), \dots, \beta(\rho_l))$. In any case, to compute $K_*(\mathcal{C}(G) \rtimes_{\alpha} \mathbb{Z})$ it is sufficient to determine the homotopy classes of $\alpha \circ \rho$ for the irreducible representations ρ of G in terms of $\beta(\rho)$'s.

II) There is a relation between the K-theory of transformation group C*-algebras of the homeomorphisms of the tori and the topological K-theory of compact nilmanifolds. In fact, let $\alpha = (\mathbf{t}, \mathbf{A})$ be an affine transformation on \mathbb{T}^n satisfying the conditions (1) through (4) in Remark 5.2. Then it has been shown in [12] that α is conjugate (in the group of affine transformations on \mathbb{T}^n) to the transformation $\alpha' = (\mathbf{t}', \mathbf{A}')$, where \mathbf{A}' has an upper triangular matrix, whose bottom right $k \times k$ corner is the identity matrix I_k and $\mathbf{t}' = (0, \dots, 0, t'_1, \dots, t'_k)$. The transformation α' is called a *standard form* for α [28]. Assume that α is given in standard form. Then J. Packer associates an *induced flow* $(\mathbb{R}, N/\Gamma)$ to the flow $(\mathbb{Z}, \mathbb{T}^n)$ generated by α , where N is a simply connected nilpotent Lie group of dimension $n + 1$, the discrete group Γ is a cocompact subgroup of N , and the action of \mathbb{R} is given by translation on the left by $\exp sX$ for $s \in \mathbb{R}$ and some $X \in \mathfrak{n}$, the Lie algebra of N . One of the most important facts is that the C*-algebra $\mathcal{C}(N/\Gamma) \rtimes_{\beta} \mathbb{R}$ corresponding to the induced flow is strongly Morita equivalent to $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ [28, Proposition 3.1]. Consequently, one has

$$(7) \quad K_i(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong K_i(\mathcal{C}(N/\Gamma) \rtimes_{\beta} \mathbb{R}) \cong K^{1-i}(N/\Gamma); \quad i = 0, 1.$$

The second isomorphism here, is the Connes' Thom isomorphism. So the K -theory of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ is converted to the topological K -theory of the compact nilmanifold N/Γ . Following the proof of Proposition 3.1 in [28], one can conclude that for the special case of Anzai transformations we can take $N = \mathfrak{F}_{n-1}$ (the generic filiform Lie group of dimension $n + 1$) and $\Gamma = \mathfrak{D}_{n-1}$, which were defined in [33]. On the other hand, following [35, Theorem 3.6], one has the isomorphism

$$(8) \quad K_i(C^*(\Gamma)) \cong K^{i+n+1}(N/\Gamma); \quad i = 0, 1.$$

Combining (7) and (8) one gets

$$(9) \quad K_i(\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}) \cong K^{i+1}(N/\Gamma) \cong K_{i+n}(C^*(\Gamma)); \quad i = 0, 1.$$

Using the above isomorphisms, one can relate the algebraic invariants of the involved C^* -algebras and topological information of the corresponding nilmanifold. For example, since N/Γ is a classifying space for Γ , one has the following isomorphisms

$$(10) \quad H_{\text{dR}}^*(N/\Gamma) \cong \check{H}^*(N/\Gamma, \mathbb{R}) \cong H^*(\Gamma, \mathbb{R}) \cong H^*(N, \mathbb{R}) \cong H^*(\mathfrak{n}, \mathbb{R}),$$

where $H_{\text{dR}}^*(N/\Gamma)$ denotes the de Rham cohomology of the manifold N/Γ , $\check{H}^*(N/\Gamma, \mathbb{R})$ denotes the Čech cohomology of N/Γ with coefficients in \mathbb{R} , $H^*(\Gamma, \mathbb{R})$ denotes the group cohomology of Γ with coefficients in the trivial Γ -module \mathbb{R} , $H^*(N, \mathbb{R})$ denotes the Moore cohomology group of N (as a locally compact group) with coefficients in the trivial Polish N -module \mathbb{R} , and $H^*(\mathfrak{n}, \mathbb{R})$ denotes the cohomology of the Lie algebra \mathfrak{n} with coefficients in the trivial \mathfrak{n} -module \mathbb{R} . Now, using the Chern isomorphisms $\text{ch}_0 : K^0(N/\Gamma) \otimes \mathbb{Q} \rightarrow \check{H}^{\text{even}}(N/\Gamma, \mathbb{Q})$ and $\text{ch}_1 : K^1(N/\Gamma) \otimes \mathbb{Q} \rightarrow \check{H}^{\text{odd}}(N/\Gamma, \mathbb{Q})$, one concludes that the even and odd cohomology groups stated in (10) are all isomorphic to \mathbb{R}^k , where k is the (common) rank of the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ as in Corollary 2.2. As an example, if $N = \mathfrak{F}_{n-1}$, $\Gamma = \mathfrak{D}_{n-1}$, and $\mathfrak{n} = \mathfrak{f}_{n-1}$, then the even and odd cohomology groups stated in (10) are all isomorphic to \mathbb{R}^{a_n} , where a_n is the rank of the K -groups of $\mathcal{A}_{n,\theta}$ that was studied in detail in Sections 5 and 6. Conversely, one may use the topological tools for N/Γ to get some information about $\mathcal{C}(\mathbb{T}^n) \rtimes_{\alpha} \mathbb{Z}$ and $C^*(\Gamma)$. For example, we know that N/Γ as a compact nilmanifold can be constructed as a principal \mathbb{T} -bundle over a lower dimensional compact nilmanifold [10]. Then we can compute the topological K -groups of N/Γ using the six term Gysin exact sequence [17, IV.1.13, p. 187]. As an example, one can see that $\mathfrak{F}_{n-1}/\mathfrak{D}_{n-1}$ is a principal \mathbb{T} -bundle over $\mathfrak{F}_{n-2}/\mathfrak{D}_{n-2}$, and the corresponding Gysin exact sequence is in fact the topological version of the Pimsner-Voiculescu exact sequence for the crossed product $\mathcal{A}_{n,\theta} \cong \mathcal{A}_{n-1,\theta} \rtimes_{\alpha} \mathbb{Z}$ as in Theorem 2.1(d) in [33].

Acknowledgment. I would like to thank Graham Denham, George Elliott, Herve Oyono-Oyono, and Tim Steger for very helpful discussions. I would like to thank N. C. Phillips for his thoughtful suggestions and for bringing the possibility of generalizing Lemma 3.1 of [30] to my attention. I am grateful to Alan T. Paterson for reading the paper and suggesting several helpful comments. I would also like to thank Paul Milnes and the Department of Mathematics of the University of Western Ontario, where some parts of this work were done.

APPENDIX A. THE SMITH NORMAL FORM

The Smith normal form is a very important tool for studying integer matrices. We refer to [27] for this interesting topic and its applications.

Definition A.1. Let $\hat{\alpha}, \hat{\beta} \in \text{End}(\mathbb{Z}^m)$. We say that $\hat{\alpha}$ is equivalent to $\hat{\beta}$ over \mathbb{Z} (and write $\hat{\alpha} \text{ equiv } \hat{\beta}$) if there exist $\hat{u}, \hat{v} \in \text{Aut}(\mathbb{Z}^m)$ such that $\hat{u} \circ \hat{\alpha} \circ \hat{v} = \hat{\beta}$. Similarly, if \mathbf{A} and \mathbf{B} are integer $m \times m$ matrices, \mathbf{A} is equivalent to \mathbf{B} if there exist $\mathbf{U}, \mathbf{V} \in \text{GL}(m, \mathbb{Z})$ such that $\mathbf{UAV} = \mathbf{B}$.

Recall that $\hat{\alpha} \text{ equiv } \hat{\beta}$, if and only if $\text{coker } \hat{\alpha} \cong \text{coker } \hat{\beta}$, if and only if $\hat{\alpha}$ and $\hat{\beta}$ have the same Smith normal form. Also, $\mathbf{A} \text{ equiv } \mathbf{B}$ if and only if \mathbf{B} is obtainable from \mathbf{A} by a finite number of elementary operations. An elementary operation on an integer matrix is one of the following types: interchanging two rows (or two columns), adding an integer multiple of one row (or column) to another, and multiplying a row (or

column) by -1 .

Now, we recall a fundamental theorem for integer matrices (cf. [27, p. 26]).

Theorem A.2 (Smith Normal Form). *Let A be an $m \times m$ integer matrix. Then A is equivalent to a diagonal matrix $\text{diag}(d_1, \dots, d_r, 0, \dots, 0)$, where r is the rank of A , and the integers d_1, \dots, d_r satisfy $d_i | d_{i+1}$ for $i = 1, \dots, r - 1$.*

APPENDIX B. NILPOTENT AND UNIPOTENT LINEAR MAPPINGS

Definition B.1. Let V be a (complex) vector space. A mapping $\hat{\epsilon} \in \text{End}_{\mathbb{C}}V$ is called nilpotent (respectively, unipotent) if $\hat{\epsilon}^k = 0$ (respectively, $(\hat{\epsilon} - \text{id})^k = 0$) for some positive integer k . The minimum value of k with this property is called the *degree* of $\hat{\epsilon}$, denoted $\text{deg}(\hat{\epsilon})$.

As an example, every upper (respectively, lower) triangular matrix with zeros on the diagonal is nilpotent. Also, the matrix S_n defined in Section 2 is a unipotent matrix of degree n . Note that all eigenvalues of a nilpotent (respectively, unipotent) matrix are zeros (respectively, ones). In particular, every unipotent matrix is invertible, and every unipotent endomorphism is an automorphism.

Corollary B.2. *Let V be a finite dimensional complex vector space and $\hat{\epsilon}$ be a nilpotent (respectively, unipotent) endomorphism of V . Then $\text{deg}(\hat{\epsilon})$ is equal to the maximum order of its Jordan blocks.*

Proof. It suffices to prove the statement for the nilpotent case. Since all the eigenvalues of $\hat{\epsilon}$ are zero, each Jordan block is a zero matrix of order one or is of the form

$$\begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & & 0 & 0 & 1 \\ 0 & \cdots & 0 & 0 & 0 \end{pmatrix}$$

which is a nilpotent matrix and its degree is the same as its order, which is greater than 1. The rest of proof is clear. \square

Definition B.3. Let V be a finite dimensional complex vector space and $\hat{\epsilon} \in \text{End}_{\mathbb{C}}V$ be nilpotent (respectively, unipotent). We say that $\hat{\epsilon}$ is of *maximal degree* if $\text{deg}(\hat{\epsilon}) = \dim V$.

Corollary B.4. *Let V be an n -dimensional complex vector space and $\hat{\epsilon} \in \text{End}_{\mathbb{C}}V$ be nilpotent (respectively, unipotent). Then $\text{deg}(\hat{\epsilon}) \leq n$. If $\text{deg}(\hat{\epsilon}) = n$, then the Jordan normal form of $\hat{\epsilon}$ is the full Jordan block of order n with 0's on the diagonal. In particular, all nilpotent (respectively, unipotent) matrices of maximal degree acting on V are similar.*

Proof. Use the preceding corollary and the Jordan normal form theorem. \square

Example B.5. Let $B = [b_{ij}]_{n \times n}$ be any upper triangular matrix, whose diagonal elements are all zeros (respectively, ones), and whose entries $b_{i,i+1}$ for $i = 1, \dots, n - 1$ are all nonzero. Then B is nilpotent (respectively, unipotent) of maximal degree. In fact, let n be the order of B and let $b := \prod_{i=1}^{n-1} b_{i,i+1}$, which is a nonzero number. Then one can easily see that $B^n = 0$ (respectively, $(B - I)^n = 0$) and B^{n-1} (respectively, $(B - I)^{n-1}$) is a matrix with b appearing on the upper-right corner and zeros elsewhere. So $\text{deg}(B) = n$, i.e. B is of maximal degree.

Lemma B.6. *Let V be a complex vector space and $\hat{\epsilon} \in \text{End}_{\mathbb{C}}V$ be nilpotent of degree k . Then $\exp(\hat{\epsilon})$ is unipotent of degree k . Moreover, $\exp(\hat{\epsilon}) - \text{id}$ is similar to $\hat{\epsilon}$.*

Proof. For the first part, we know that $\exp(\hat{\epsilon}) - \text{id} = \hat{\epsilon} + \hat{\epsilon}^2/2! + \dots + \hat{\epsilon}^{k-1}/(k-1)! = \hat{\epsilon}\omega$, where $\omega := \text{id} + \hat{\epsilon}/2! + \dots + \hat{\epsilon}^{k-2}/(k-1)!$ commutes with $\hat{\epsilon}$ and is invertible since it is unipotent. So, $(\exp(\hat{\epsilon}) - \text{id})^r = (\hat{\epsilon}\omega)^r = \hat{\epsilon}^r \omega^r$ for all positive integers r . Thus $\exp(\hat{\epsilon}) - \text{id}$ is unipotent with the same degree of $\hat{\epsilon}$. For the second part, using the Jordan normal form of $\hat{\epsilon}$, it is sufficient to prove the statement for the special case when $\hat{\epsilon}$ is a Jordan block with zeros on the diagonal. Since in this case $\hat{\epsilon}$ is of maximal degree, by the first part, $\exp(\hat{\epsilon}) - \text{id}$ is also of maximal degree. Therefore they are similar by Corollary B.4. \square

APPENDIX C. ENDOMORPHISMS AND DERIVATIONS OF EXTERIOR ALGEBRAS

We refer to the Chapter 5 of [11] for general properties of exterior algebras and mappings between them.

Let V be a (complex) vector space and $\hat{\phi} : V \rightarrow V$ be a linear mapping. Then $\hat{\phi}$ can be extended in a unique way to a homomorphism $\wedge^* \hat{\phi} : \wedge^* V \rightarrow \wedge^* V$ such that $\wedge^* \hat{\phi}(1) = 1$, yielding

$$\wedge^* \hat{\phi}(x_1 \wedge \dots \wedge x_p) = \hat{\phi}(x_1) \wedge \dots \wedge \hat{\phi}(x_p), \quad (x_i \in V).$$

Also, $\hat{\phi}$ can be extended in a unique way to a derivation $\mathcal{D}^* \hat{\phi} : \wedge^* V \rightarrow \wedge^* V$, yielding

$$\mathcal{D}^* \hat{\phi}(x_1 \wedge \dots \wedge x_p) = \sum_{i=1}^p x_1 \wedge \dots \wedge \hat{\phi}(x_i) \wedge \dots \wedge x_p \quad (p \geq 2, x_i \in V).$$

We define $\wedge^r \hat{\phi} := \wedge^* \hat{\phi}|_{\wedge^r V}$ and $\mathcal{D}^r \hat{\phi} := \mathcal{D}^* \hat{\phi}|_{\wedge^r V}$ as induced linear mappings on the r -th exterior power of V for $r \geq 0$. Then we have

$$\wedge^* \hat{\phi} = \bigoplus_{r \geq 0} \wedge^r \hat{\phi}, \quad \mathcal{D}^* \hat{\phi} = \bigoplus_{r \geq 0} \mathcal{D}^r \hat{\phi}.$$

One can easily show that $\wedge^*(\hat{\phi} \circ \hat{\psi}) = (\wedge^* \hat{\phi}) \circ (\wedge^* \hat{\psi})$ and $\mathcal{D}^*([\hat{\phi}, \hat{\psi}]) = [\mathcal{D}^* \hat{\phi}, \mathcal{D}^* \hat{\psi}]$ (cf. equations (5.20) and (5.25) in [11]).

Lemma C.1. *With the above notation, if $\hat{\phi} : V \rightarrow V$ is nilpotent, then $\wedge^r \hat{\phi}$ and $\mathcal{D}^r \hat{\phi}$ are also nilpotent for $r \geq 1$. If V is finite dimensional, then $\mathcal{D}^* \hat{\phi}$ is nilpotent.*

Proof. Assume that $\hat{\phi}^t = 0$ for some $t \in \mathbb{N}$. We have $(\wedge^r \hat{\phi})^t(x_1 \wedge \dots \wedge x_r) = \hat{\phi}^t(x_1) \wedge \dots \wedge \hat{\phi}^t(x_r) = 0$. So, $(\wedge^r \hat{\phi})^t = 0$, which means that $\wedge^r \hat{\phi}$ is nilpotent. For $\mathcal{D}^r \hat{\phi}$, we know that $\mathcal{D}^r \hat{\phi}(x_1 \wedge \dots \wedge x_r) = \sum_{i=1}^r x_1 \wedge \dots \wedge \hat{\phi}(x_i) \wedge \dots \wedge x_r$ and one can easily deduce that

$$\mathcal{D}^r \hat{\phi}^p(x_1 \wedge \dots \wedge x_r) = \sum_{\substack{i_1 + \dots + i_r = p \\ (i_j \geq 0)}} \frac{p!}{(i_1)! \dots (i_r)!} \hat{\phi}^{i_1} x_1 \wedge \dots \wedge \hat{\phi}^{i_r} x_r.$$

Now since $i_1 + \dots + i_r = p$, there exists an i_j with $i_j \geq p/r$. So, if $p \geq rt$ then $\hat{\phi}^{i_j} = 0$ and $\mathcal{D}^r \hat{\phi}^p = 0$. Thus $\mathcal{D}^r \hat{\phi}$ is nilpotent.

For the next part, let $m := \dim V$. Since $\mathcal{D}^* \hat{\phi} = \bigoplus_{r \geq 0}^m \mathcal{D}^r \hat{\phi}$ and $\hat{\phi}_0 = 0$, from the first part we have $(\mathcal{D}^* \hat{\phi})^{mt} = 0$, hence $\mathcal{D}^* \hat{\phi}$ is nilpotent, too. \square

Lemma C.2. *Let $\hat{\phi} : V \rightarrow V$ be a nilpotent linear mapping. Then $\exp(\mathcal{D}^* \hat{\phi}) = \wedge^* \exp(\hat{\phi})$ on $\wedge^* V$.*

Proof. We have

$$\begin{aligned} \exp(\mathcal{D}^* \hat{\phi})(x_1 \wedge \dots \wedge x_r) &= \sum_{p \geq 0} \frac{1}{p!} \mathcal{D}^p \hat{\phi}(x_1 \wedge \dots \wedge x_r) \\ &= \sum_{p \geq 0} \frac{1}{p!} \left(\sum_{\substack{i_1 + \dots + i_r = p \\ (i_j \geq 0)}} \frac{p!}{(i_1)! \dots (i_r)!} \hat{\phi}^{i_1} x_1 \wedge \dots \wedge \hat{\phi}^{i_r} x_r \right) \end{aligned}$$

$$\begin{aligned}
&= \sum_{i_j \geq 0} \frac{1}{(i_1)! \dots (i_r)!} \hat{\phi}^{i_1} x_1 \wedge \dots \wedge \hat{\phi}^{i_r} x_r \\
&= \left(\sum_{i_1 \geq 0} \frac{1}{(i_1)!} \hat{\phi}^{i_1} x_1 \right) \wedge \dots \wedge \left(\sum_{i_r \geq 0} \frac{1}{(i_r)!} \hat{\phi}^{i_r} x_r \right) \\
&= (\exp(\hat{\phi})x_1) \wedge \dots \wedge (\exp(\hat{\phi})x_r) \\
&= \wedge^* \exp(\hat{\phi})(x_1 \wedge \dots \wedge x_r),
\end{aligned}$$

which yields the result. Note that all sums in the above equalities are finite according to the previous lemma. \square

Remark C.3. The nilpotency of $\hat{\phi}$ is not necessary in the preceding lemma. In fact, one may use the definition of $\exp : \mathfrak{gl}(\Lambda^*V) \rightarrow \mathrm{GL}(\Lambda^*V)$. More precisely, define $s : \mathbb{R} \rightarrow \mathrm{GL}(\Lambda^*V)$ by $s(t) = \wedge^* \exp(t\hat{\phi})$. Then one may check that s is the 1-parameter subgroup generated by $\mathcal{D}^* \hat{\phi}$ (i.e. $\dot{s}(0) = \mathcal{D}^* \hat{\phi}$), and we have $s(1) = \wedge^* \exp(\hat{\phi})$.

Corollary C.4. *Let $\hat{\phi} : V \rightarrow V$ be a nilpotent linear mapping, and set $\hat{e} := \hat{\phi} + \mathrm{id}$. Then $\wedge^r \hat{e} - \mathrm{id}$ is similar to $\mathcal{D}^r \hat{\phi}$ for $r \geq 0$.*

Proof. We know from Lemma B.6 that $\exp(\hat{\phi}) - \mathrm{id}$ is similar to $\hat{\phi}$, hence $\exp(\hat{\phi})$ is similar to $\hat{\phi} + \mathrm{id} = \hat{e}$. So

$$\wedge^r \hat{e} - \mathrm{id} \sim \wedge^r \exp(\hat{\phi}) - \mathrm{id} = \exp(\mathcal{D}^r \hat{\phi}) - \mathrm{id} \sim \mathcal{D}^r \hat{\phi}.$$

\square

APPENDIX D. REPRESENTATION THEORY OF $\mathfrak{sl}(2, \mathbb{C})$

We refer to Section II.7 of [15] for studying the irreducible representations of the Lie algebra $\mathfrak{sl}(2, \mathbb{C})$.

Let $\mathfrak{sl}(2, \mathbb{C})$ denote the special linear Lie algebra over \mathbb{C}^2 defined by $\mathfrak{sl}(2, \mathbb{C}) := \{a \in M_2(\mathbb{C}) \mid \mathrm{Tr}(a) = 0\}$. It is well known that $\mathfrak{sl}(2, \mathbb{C})$ is a 3-dimensional simple complex Lie algebra. One can check that

$$\mathfrak{B} := \left\{ h := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, e := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, f := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right\}$$

is a basis for this Lie algebra.

The following theorem is the foundation of representation theory of semisimple Lie algebras including $\mathfrak{sl}(2, \mathbb{C})$. It is stated and proved in the Subsection II.6.3 of [15].

Theorem D.1 (Weyl). *Every finite dimensional representation of a semisimple Lie algebra is completely reducible, namely, it can be decomposed into a direct sum of irreducible representations.*

Proposition D.2. *Let V be an n -dimensional complex vector space with a basis $\{e_1, \dots, e_n\}$. Then the following equalities (for $i = 1, \dots, n$)*

- (a) $\pi_n(h)e_i = (2i - n - 1)e_i$;
- (b) $\pi_n(e)e_i = i(n - i)e_{i+1}$; ($e_{n+1} := 0$)
- (c) $\pi_n(f)e_i = e_{i-1}$; ($e_0 := 0$).

define a representation $\pi_n : \mathfrak{sl}(2, \mathbb{C}) \rightarrow \mathfrak{gl}(V)$.

We call π_n defined in the previous proposition the *canonical representation of $\mathfrak{sl}(2, \mathbb{C})$ on V associated with the basis $\{e_1, \dots, e_n\}$* . We recall the following theorem from Section II.7 of [15].

Theorem D.3. *Let π_n be the representation described above. Then*

- (i) π_n is an irreducible representation of $\mathfrak{sl}(2, \mathbb{C})$.
- (ii) Any n -dimensional irreducible representation of $\mathfrak{sl}(2, \mathbb{C})$ is equivalent to π_n .
- (iii) Let V be a finite dimensional $\mathfrak{sl}(2, \mathbb{C})$ -module and define

$$V_\alpha = \{v \in V \mid h.v = \alpha v\}$$

for $\alpha \in \mathbb{C}$. Then V decomposes into a direct sum of irreducible submodules (Weyl), and in any such decomposition, the number of summands is precisely $\dim V_0 + \dim V_1$.

APPENDIX E. SOME COMPUTER CODES AND A COUNTEREXAMPLE

To compute the K -groups of $\mathcal{C}(\mathbb{T}^n) \rtimes_\alpha \mathbb{Z}$, one should first compute the kernels and cokernels of the following integer matrices

$$(11) \quad A_r := \wedge^r A - I_{\binom{n}{r}}$$

for $r = 0, 1, \dots, n$, where A is the integer matrix corresponding to α acting on \mathbb{Z}^n . We have written two Maple codes to obtain this goal.

The first one is an auxiliary procedure called `exterior(r,A)`, which computes the r -th exterior power of a given $n \times n$ integer matrix A for $r = 1, \dots, n$ as follows (note that $\wedge^0 A := I_1$)

```
> exterior:=proc(r,A)
> local n,N,Q,E,i,j;
> n:=linalg[rowdim](A);
> N:=binomial(n,r);
> Q:=combinat[choose](n,r);
> E:=array(1..N,1..N);
> for i from 1 to N do
> for j from 1 to N do
> E[i,j]:=linalg[det](linalg[submatrix](A,Q[i],Q[j]));
> od;
> od;
> RETURN(evalm(E));
> end;
```

The second code, which calls the first one, is called `po(A)`. It lists the factorized characteristic polynomials of the Smith normal forms of A_r for $r = 0, 1, \dots, n$ given a matrix A

```
> po:=proc(A)
> local r,n,x,p;
> n:=linalg[rowdim](A);
> print(x);
> for r from 1 to n do
> p:=factor(linalg[charpoly](linalg[ismith](exterior(r,A)-1),x));
> print(p);
> od;
> end;
```

These factorized polynomials encode the diagonal entries of the Smith normal forms of the matrices A_r for $r = 0, 1, \dots, n$. In particular, one can easily find the kernels and cokernels of these matrices. To see this, observe that the kernel of an $m \times m$ integer matrix is isomorphic to a torsion-free finitely generated abelian group, whose rank is the number of zeros on the diagonal of its Smith normal form, and the cokernel is isomorphic to $\bigoplus_{j=1}^m \mathbb{Z}_{k_j}$, where k_1, \dots, k_m are the diagonal entries of the Smith normal form.

The following example computes the K -groups of $\mathcal{A}_{6,\theta}$, which was mentioned in Remark 1.7 as a counterexample for [16, Proposition 2.17].

Example E.1. $K_0(\mathcal{A}_{6,\theta}) \cong \mathbb{Z}^{13}$ and $K_1(\mathcal{A}_{6,\theta}) \cong \mathbb{Z}^{13} \oplus \mathbb{Z}_2$.

Proof. Using the procedure $\text{po}(\mathbf{A})$ for the matrix $\mathbf{A} := \mathbf{S}_6$ as defined at the beginning of Section 3 we get $\text{po}(\mathbf{A})$;

$$\begin{array}{c} x \\ x(x-1)^5 \\ x^3(x-1)^{12} \\ x^3(x-2)(x-1)^{16} \\ x^3(x-1)^{12} \\ x(x-1)^5 \\ x \end{array}$$

Consequently, we have the following table representing the kernels and cokernels of $\mathbf{S}_{6,r} := \wedge^r \mathbf{S}_6 - 1_{\binom{6}{r}}$ for $r = 0, 1, \dots, 6$.

r	$\ker \mathbf{S}_{6,r}$	$\text{coker } \mathbf{S}_{6,r}$
0	\mathbb{Z}	\mathbb{Z}
1	\mathbb{Z}	\mathbb{Z}
2	\mathbb{Z}^3	\mathbb{Z}^3
3	\mathbb{Z}^3	$\mathbb{Z}^3 \oplus \mathbb{Z}_2$
4	\mathbb{Z}^3	\mathbb{Z}^3
5	\mathbb{Z}	\mathbb{Z}
6	\mathbb{Z}	\mathbb{Z}

Now, using Theorem 2.1 we have

$$\begin{aligned} K_0(\mathcal{A}_{6,\theta}) &\cong (\text{coker } \mathbf{S}_{6,0} \oplus \text{coker } \mathbf{S}_{6,2} \oplus \text{coker } \mathbf{S}_{6,4} \oplus \text{coker } \mathbf{S}_{6,6}) \oplus (\ker \mathbf{S}_{6,1} \oplus \ker \mathbf{S}_{6,3} \oplus \ker \mathbf{S}_{6,5}) \\ &\cong (\mathbb{Z} \oplus \mathbb{Z}^3 \oplus \mathbb{Z}^3 \oplus \mathbb{Z}) \oplus (\mathbb{Z} \oplus \mathbb{Z}^3 \oplus \mathbb{Z}) = \mathbb{Z}^{13}, \\ K_1(\mathcal{A}_{6,\theta}) &\cong (\text{coker } \mathbf{S}_{6,1} \oplus \text{coker } \mathbf{S}_{6,3} \oplus \text{coker } \mathbf{S}_{6,4}) \oplus (\ker \mathbf{S}_{6,0} \oplus \ker \mathbf{S}_{6,2} \oplus \ker \mathbf{S}_{6,4} \oplus \ker \mathbf{S}_{6,6}) \\ &\cong (\mathbb{Z} \oplus \mathbb{Z}^3 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}) \oplus (\mathbb{Z} \oplus \mathbb{Z}^3 \oplus \mathbb{Z}^3 \oplus \mathbb{Z}) = \mathbb{Z}^{13} \oplus \mathbb{Z}_2. \end{aligned}$$

□

APPENDIX F. APPLICATIONS TO $C^*(\mathfrak{D}_n)$

This appendix is devoted to some applications of the results of this paper to the C^* -algebras studied in [33], in which the authors promised to completely classify the simple infinite dimensional quotients of the group C^* -algebra of their interest $C^*(\mathfrak{D}_n)$ by their K -theoretic invariants in a later work. In this context, the discrete group \mathfrak{D}_n is a higher dimensional analogue of the discrete Heisenberg group H_3 , and is defined by

$$\mathfrak{D}_n = \langle x, y_0, y_1, \dots, y_n \mid xy_0 = y_0x, y_i y_j = y_j y_i \text{ for } 0 \leq i, j \leq n, [x, y_j] = y_{j-1} \text{ for } 1 \leq j \leq n \rangle,$$

where $[x, y] := xyx^{-1}y^{-1}$. The group \mathfrak{D}_n can be represented as a semidirect product $\mathbb{Z}^{n+1} \rtimes_{\eta} \mathbb{Z}$, where the group homomorphism $\eta : \mathbb{Z} \rightarrow \text{GL}(n+1, \mathbb{Z})$ is such that $\eta(k)$ is the matrix, whose (i, j) -entry is given by $\binom{k}{j-i}$ as defined in [33]. This realization allows us to study K -theory of $C^*(\mathfrak{D}_n)$.

Proposition F.1. $K_i(C^*(\mathfrak{D}_n)) \cong K_i(\mathcal{A}_{n+1,\theta})$ for $i = 0, 1$. In particular,

$$\text{rank } K_0(C^*(\mathfrak{D}_n)) = \text{rank } K_1(C^*(\mathfrak{D}_n)) = a_{n+1}.$$

Proof. Since $\mathfrak{D}_n \cong \mathbb{Z}^{n+1} \rtimes_{\tilde{\eta}} \mathbb{Z}$, so $C^*(\mathfrak{D}_n) \cong C^*(\mathbb{Z}^{n+1}) \rtimes_{\tilde{\eta}} \mathbb{Z} \cong \mathcal{C}(\mathbb{T}^{n+1}) \rtimes_{\tilde{\eta}} \mathbb{Z}$ and the integer matrix corresponding to $\tilde{\eta}$ is the $(n+1) \times (n+1)$ matrix M_n introduced in [33], which is precisely the matrix S_{n+1} defined in Section 5 to describe the linear structure of Anzai transformations on \mathbb{T}^{n+1} . The rest of the proof follows from the Theorem 2.1. \square

As some higher-dimensional analogues of the irrational rotation algebras A_θ , all simple infinite-dimensional quotients of $C^*(\mathfrak{D}_n)$ have been classified in Theorem 3.2 of [33]. These consist of the C^* -algebras $\mathcal{A}_{n,\theta}$ for some irrational parameter θ , and a few more classes of C^* algebras denoted by $A_i^{(n)}$, which are of the form $\mathcal{C}(\mathbf{Y}_i \times \mathbb{T}^{n-i}) \rtimes_{\phi_i} \mathbb{Z}$ for some suitable finite sets \mathbf{Y}_i and minimal homeomorphisms ϕ_i for $i = 1, 2, \dots, n-1$. Then it is proved in Theorem 4.8 of [33] that $A_i^{(n)} \cong M_{C_i}(B_i^{(n)})$, where $B_i^{(n)}$ is a certain transformation group C^* -algebra of some affine Furstenberg transformation on \mathbb{T}^{n-i} . We conclude the following results.

Corollary F.2. *Let A be a simple infinite dimensional quotient of $C^*(\mathfrak{D}_n)$. Then $\text{rank } K_0(A) = \text{rank } K_1(A) = a_{n-i}$ for some $i \in \{0, 1, \dots, n-1\}$ that is uniquely determined by the isomorphism $A \cong \mathcal{C}(\mathbf{Y}_i \times \mathbb{T}^{n-i}) \rtimes_{\phi_i} \mathbb{Z}$ as in Theorem 3.2 of [33].*

Proof. It is proved that A is isomorphic to a matrix algebra over a Furstenberg transformation group C^* -algebra $B_i^{(n)}$ on \mathbb{T}^{n-i} for some suitable $i \in \{0, 1, \dots, n\}$ [33, Theorem 4.8]. So $K_j(A) \cong K_j(B_i^{(n)})$ for $j = 0, 1$. The rest of the proof is clear from the preceding theorem. \square

We saw in Proposition 6.5 that $\{a_n\}$ is a strictly increasing sequence. Therefore the preceding corollary is a first step towards the classification of the simple infinite dimensional quotients of $C^*(\mathfrak{D}_n)$ by means of K -theory. But as is seen, the rank of the K -groups alone can not distinguish the algebras at the same ‘‘level’’ (i.e. those algebras that are included in the same class, but with different values of the parameters). The other powerful K -theoretic invariant that helps us do this is the trace invariant, i.e. the range of the unique tracial state acting on the K_0 -group.

Proposition F.3. *Suppose $A \cong \mathcal{C}(\mathbf{Y}_i \times \mathbb{T}^{n-i}) \rtimes_{\phi_i} \mathbb{Z}$ is a simple infinite dimensional quotient of $C^*(\mathfrak{D}_n)$ as in Theorem 3.2 in [33]. Then A has a unique tracial state $\tilde{\tau}$ and $\tilde{\tau}_* K_0(A) = \frac{1}{C_i}(\mathbb{Z} + \mathbb{Z}\vartheta_i)$, where $C_i = |\mathbf{Y}_i|$ and $e^{2\pi i\vartheta_i} = \zeta_i = (-1)^{C_i+1} \eta_i^{C_i}$ as in Lemma 4.6 and Theorem 4.8 in [33].*

Proof. Following Theorem 4.8 in [33], A is isomorphic to $M_{C_i}(B_i^{(n)}) = M_{C_i}(\mathbb{C}) \otimes B_i^{(n)}$, where $B_i^{(n)}$ is a simple Furstenberg transformation group C^* -algebra with the irrational parameter $\zeta_i = (-1)^{C_i+1} \eta_i^{C_i}$. By Corollary 1.5, $B_i^{(n)}$ has a unique tracial state τ . Moreover, $\tau_* K_0(B_i^{(n)}) = \mathbb{Z} + \mathbb{Z}\vartheta_i$, where $e^{2\pi i\vartheta_i} = \zeta_i$ (see [16, Theorem 2.23] or Theorem 7.2). Thus A has the unique tracial state $\tilde{\tau} = (\frac{1}{C_i} \text{Tr}) \otimes \tau$, in which Tr is the usual trace on $M_{C_i}(\mathbb{C})$, and so $\tilde{\tau}_* K_0(A) = \frac{1}{C_i}(\mathbb{Z} + \mathbb{Z}\vartheta_i)$ [16, Lemma 3.5]. \square

Finally, we can characterize all simple infinite-dimensional quotients of $C^*(\mathfrak{D}_n)$.

Proposition F.4. $\mathcal{A}_{n,\theta} \cong \mathcal{A}_{n',\theta'}$ if and only if $n = n'$ and there exists an integer k such that $\theta = k \pm \theta'$. More generally, let $A_i^{(n)} \cong \mathcal{C}(\mathbf{Y}_i \times \mathbb{T}^{n-i}) \rtimes_{\phi_i} \mathbb{Z}$ be a simple infinite dimensional quotient of $C^*(\mathfrak{D}_n)$ with the structure constants $\lambda, \mu_1, \dots, \mu_i$ as on p. 165-166 of [33], and let $A_{i'}^{(n')} \cong \mathcal{C}(\mathbf{Y}'_{i'} \times \mathbb{T}^{n'-i'}) \rtimes_{\phi'_{i'}} \mathbb{Z}$ be a simple infinite dimensional quotient of $C^*(\mathfrak{D}_{n'})$ with the structure constants $\lambda', \mu'_1, \dots, \mu'_{i'}$. Suppose that $C_i = |\mathbf{Y}_i|$ and $C'_{i'} = |\mathbf{Y}'_{i'}|$. Then $A_i^{(n)} \cong A_{i'}^{(n')}$ if and only if $n - i = n' - i'$, $C_i = C'_{i'}$, and

$$\lambda^{(C_i)} \mu_1^{(C_i)} \mu_2^{(C_i-1)} \dots \mu_i^{C_i} = \lambda'^{(C'_{i'})} \mu'_1^{(C'_{i'})} \mu'_2^{(C'_{i'}-1)} \dots \mu'_{i'}^{C'_{i'}}$$

or

$$\lambda^{(C_i)} \mu_1^{(C_i)} \mu_2^{(C_i-1)} \dots \mu_i^{C_i} = (\lambda'^{(C'_{i'})} \mu'_1^{(C'_{i'})} \mu'_2^{(C'_{i'}-1)} \dots \mu'_{i'}^{C'_{i'}})^{-1}.$$

Proof. Use the previous proposition and the fact that $\{a_n\}$ is a strictly increasing sequence (see Proposition 6.5). Note that

$$\zeta_i = (-1)^{C_i+1} \eta_i^{C_i} = \lambda^{\binom{C_i}{i+1}} \mu_1^{\binom{C_i}{i}} \mu_2^{\binom{C_i}{i-1}} \dots \mu_i^{C_i}$$

by the last equation on p. 171 of [33]. \square

Remark F.5. Note that $C_i = |\mathbf{Y}_i|$ is completely determined by the structural constants $\lambda, \mu_1, \dots, \mu_{i-1}$ (which are roots of unity). More precisely, by calculations on p. 165-166 of [33], we have

$$C_i = \min\{r \in \mathbb{N} \mid \lambda^r = \lambda^{\binom{r}{2}} \mu_1^r = \dots = \lambda^{\binom{r}{i}} \mu_1^{\binom{r}{i-1}} \mu_2^{\binom{r}{i-2}} \dots \mu_{i-1}^r = 1\}.$$

For an explicit example in lower dimensions, see [26, Lemma 5.4].

REFERENCES

- [1] J. Anderson, W. Paschke, *The rotation algebra*, Houston J. Math. **15**(1) (1989), 1–26.
- [2] M. Atiyah, *K-theory*, W. A. Benjamin, New York, 1967.
- [3] B. Blackadar, *K-theory for operator algebras*, Second edition, Mathematical Sciences Research Institute Publications 5, Cambridge University Press, Cambridge, 1998.
- [4] E.G. Effros and F. Hahn, *Locally compact transformation groups and C^* -algebras*, Mem. Amer. Math. Soc. No. 75, Providence, RI (1967).
- [5] R. C. Entringer, *Representation of m as $\sum_{k=-n}^n \epsilon_k k$* , Canad. Math. Bull. **11** (1968), 289–293.
- [6] P. Erdős, *Extremal problems in number theory*, 1965 Proc. Sympos. Pure Math., Vol. VIII, pp. 181–189, Amer. Math. Soc., Providence, R.I.
- [7] R. Exel, *Rotation numbers for automorphisms of C^* -algebras*, Pacific J. Math. **127**(7) (1981), 31–89.
- [8] H. Furstenberg, *Recurrence in ergodic theory and combinatorial number theory*, Princeton University Press, Princeton, NJ, 1981.
- [9] H. Furstenberg, *Strict ergodicity and transformation of the torus*, Amer. J. Math. **83** (1961), 573–601.
- [10] V. V. Gorbatsevich, A. L. Onishchik and E. B. Vinberg, *Foundations of Lie theory and Lie transformation groups*, Encyclopaedia of Mathematical Sciences 20, Springer-Verlag, Berlin Heidelberg, 1997.
- [11] W. Greub, *Multilinear algebra*, Second edition, Springer-Verlag, New York-Heidelberg, 1978.
- [12] F. J. Hahn, *On affine transformations of compact abelian groups*, Amer. J. Math. **85** (1963), 428–446.
- [13] F. J. Hahn and W. Parry, *Minimal systems with quasi-discrete spectrum*, J. London, Math. Soc. **40** (1965), 300–323.
- [14] L. Hodgkin, *On the K-theory of Lie groups*, Topology, **6** (1967), 1–36.
- [15] J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Second printing, revised, Graduate Texts in Mathematics 9, Springer-Verlag, New York-Berlin, 1978.
- [16] R. Ji, *On the crossed product C^* -algebras associated with Furstenberg transformations on tori*, Ph.D. dissertation, State University of New York at Stony Brook, August 1986.
- [17] M. Karoubi, *K-theory*, Springer-Verlag, Berlin Heidelberg, 1978.
- [18] K. Kodaka, *The positive cones of K_0 -groups of crossed products associated with Furstenberg transformations on the 2-torus*, Proc. Edinburgh Math. Soc. (2) **43** (1), (2000), 167–175.
- [19] H. Lin, *classification of simple C^* -algebras of tracial topological rank zero*, Duke Math. J. **125**(1) (2004), 91–119.
- [20] Q. Lin, N. C. Phillips, *Direct limit decomposition for C^* -algebras of minimal diffeomorphisms*, Operator algebras and applications, 107–133, Adv. Stud. Pure Math., 38, Math. Soc. Japan, Tokyo, 2004.
- [21] H. Lin, N. C. Phillips, *Crossed products by minimal homeomorphisms* J. Reine Angew. Math. **641** (2010), 95–122.
- [22] Q. Lin, N. C. Phillips, *Ordered K-theory for C^* -algebras of minimal homeomorphisms*, Operator algebras and operator theory (Shanghai, 1997), 289–314, Contemp. Math., **228**, Amer. Math. Soc., Providence, RI, 1998.
- [23] J. H. van Lint, *Representations of 0 as $\sum_{k=-N}^N \epsilon_k k$* , Proc. Amer. Math. Soc. **18** (1967), 182–184.
- [24] P. Milnes, *Groups and C^* -algebras from a 4-dimensional Anzai flow*, Math. Nachr. **235** (2002), 129–141.
- [25] P. Milnes and S. Walters, *Simple quotients of the group C^* -algebra of a discrete 4-dimensional nilpotent group*, Houston J. Math. **19** (1993), 615–636.
- [26] P. Milnes and S. Walters, *Simple infinite dimensional quotients of $C^*(G)$ for discrete 5-dimensional nilpotent groups* G, Illinois J. Math. **41**(2) (1997), 315–340.
- [27] M. Newman, *Integral matrices*, Pure and Applied Mathematics 45, Academic Press, New York and London, 1972.
- [28] J. A. Packer, *K-theoretic invariants for C^* -algebras associated to transformations and induced flows*, J. Funct. Anal. **67** (1986), 25–59.
- [29] N. C. Phillips, *Cancellation and stable rank for direct limits of recursive subhomogeneous algebras*, Trans. Amer. Math. Soc. **359** (2007), 4625–4652.
- [30] N. C. Phillips, *Examples of different minimal diffeomorphisms giving the same C^* -algebras*, Israel J. Math. **160** (2007), 189–217.

- [31] M. Pimsner and D. Voiculescu, *Exact sequences for K -groups and Ext -groups of certain crossed product C^* -algebras*, J. Operator Theory, **4** (1980), 93–118.
- [32] S. C. Power, *Simplicity of C^* -algebras associated to minimal dynamical systems*, J. London Math. Soc. **18** (1978), 534–538.
- [33] K. Reihani and P. Milnes, *Analysis on discrete cocompact subgroups of the generic filiform Lie groups*, Acta. Math. Hung., **112** (1-2) (2006), 157–179.
- [34] M. Rieffel, *C^* -algebras associated with irrational rotations*, Pacific J. Math. **93**(2) (1981), 415–429.
- [35] J. Rosenberg, *Group C^* -algebras and topological invariants*, in Operator algebras and group representations, vol. I, Pitman, London, 1984, pp. 95–115.
- [36] C. Schochet, *Topological methods for C^* -algebras II: Geometric resolutions and the Künneth formula*, Pacific J. Math. **98** (2) (1982), 443–458.
- [37] J. L. Taylor, *Banach algebras and topology*, Algebras in analysis, (J. H. Williamson, ed.), Academic Press, London, 1975, pp. 118–186.
- [38] J. Tomiyama, *Invitation to C^* -algebras and topological dynamics*, Advanced Series in Dynamical Systems, vol. 3, World Scientific, Singapore, 1987.
- [39] S. Walters, *K -groups and classification of simple quotients of group C^* -algebras of certain discrete 5-dimensional nilpotent groups*, Pacific J. Math. **202**(2) (2002), 491–509.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KS 66045-7594
E-mail address: reihani@math.ku.edu