

Large primes in generalized Pascal triangles

Gábor FARKAS

Eötvös Loránd University
email: farkasg@compalg.inf.elte.hu

Gábor KALLÓS

Széchenyi István University, Győr
email: kallos@sze.hu

Gyöngyvér KISS

Eötvös Loránd University
email: kissgyongyver@gmail.com

Abstract. In this paper, after presenting the results of the generalization of Pascal triangle (using powers of base numbers), we examine some properties of the 112-based triangle, most of all regarding to prime numbers. Additionally, an effective implementation of ECPP method is presented which enables Magma computer algebra system to prove the primality of numbers with more than 1000 decimal digits.

1 Generalized Pascal triangles using the powers of base numbers

As it is a well-known fact, the classic Pascal triangle has served as a model for various generalizations. Among the broad variety of ideas of generalizations we can find e.g.: the generalized binomial coefficients of s^{th} order (leading to generalized Pascal triangles of s^{th} order), the multinomial coefficients (leading to Pascal pyramids and hyperpyramids), special arithmetical sequences (leading to resulting triangles which we might call as Lucas, Fibonacci, Gaussian, Catalan, ... triangle) (details in [3]).

One of the present authors has devised, and then worked out in detail and published such a type of generalization, which is based on the idea of using

Computing Classification System 1998: G.4

Mathematics Subject Classification 2010: 05A10, 11Y11

Key words and phrases: generalized Pascal triangles, elliptic curve primality proving

“the powers of the base number”. Referring to our former results (presented in detail in [7] and [8]; here we don’t repeat/echo the theorems and propositions) we show here the first few rows of the 112-based triangle (Figure 1), which will gain outstanding importance below in this paper.

						1						
						1	1	2				
					1	2	5	4	4			
			1	3	9	13	18	12	8			
		1	4	14	28	49	56	56	32	16		
	1	5	20	50	105	161	210	200	160	80	32	
1	6	27	80	195	366	581	732	780	640	432	192	64
						...						

Figure 1: The 112-based triangle

Let us use the notation $E_{k,n}^{a_0 a_1 \dots a_{m-1}}$ for the k^{th} element in the n^{th} row of $a_0 a_1 \dots a_{m-1}$ -based triangle ($0 \leq a_0, a_1, \dots, a_{m-1} \leq 9$ are integers). Then we have the definition rule, as follows:

$$E_{k,n}^{a_0 a_1 \dots a_{m-1}} = a_{m-1} E_{k-m+1,n-1}^{a_0 a_1 \dots a_{m-1}} + a_{m-2} E_{k-m+2,n-1}^{a_0 a_1 \dots a_{m-1}} + \dots + a_1 E_{k-1,n-1}^{a_0 a_1 \dots a_{m-1}} + a_0 E_{k,n-1}^{a_0 a_1 \dots a_{m-1}}.$$

The indices in the rows and columns run from 0, elements with non-existing indices are considered to be zero. Applying this general form to the 112-based triangle (now: $m = 3$), we get the specific rule

$$E_{k,n}^{112} = 2E_{k-2,n-1}^{112} + E_{k-1,n-1}^{112} + E_{k,n-1}^{112}.$$

The historical overview of this special field is presented in [8]. In the last few years there were published several new results which are related to our topic (e.g. [2]). Moreover, besides that, up to about 2005, all generalized triangle sequences of the type $ax + by$ were added to the database On-line Encyclopedia of Integer sequences [11], since that time there have been several new applications, too, based on sequences appearing in our triangles. However, e.g. the sequences based on the general abc -based triangles are still not widely known.

Recalling the basic properties of generalized triangles—most of all in connection with powering the base number $a_0 a_1 \dots a_{m-1}$ and with the polynomial

$(a_0x_0 + a_1x_1 + \dots + a_{m-1}x_{m-1})^n$ —we can state that we have the “right” to call these types of triangles as generalized Pascal triangles (details in [8], summary in [6]).

2 Divisibility of elements and prime numbers

The classic divisibility investigations in Pascal triangle (for binomial coefficients) are very popular and even spectacular, if the traditional “strict” mathematical approach is moved toward coloring and fractals (details in [3]). For generalized binomial coefficients (with our notation: in triangles with bases $11 \dots 1$) we have similar results, too, with a remark that in these cases general proofs are harder, and there are many conjectures, too.

We recall here the beautiful result of Richard C. Bollinger, who proved for generalized Pascal triangles of p^{th} order that for large n , “almost every” element in the n^{th} row is divisible by p (see [3], p. 24). For example, for the 111-based triangle this means divisibility by 3. (We mention that the p^{th} order Pascal triangle is a triangle with base $11 \dots 1$, where we have p pieces of 1.)

Now we turn our attention specially to the 112-based triangle, and in the following we are interested mostly in prime numbers. It is obvious that the right part of the triangle contains only even numbers. Moreover, if we move to the right, the powers of 2 are usually (not always) growing as divisors. Analyzing connections with the multinomial theorem we can conclude that the left part of the triangle contains mostly (with possible exception of the first two places) composite numbers, too. Of course, this can be not true for the 0^{th} and 1^{st} numbers, which are the same as in the classic Pascal triangle. Moreover, using induction we can see that the center element in every row is always odd.

We can pose obviously two (not hard) questions in connection with prime numbers:

1. Can we find every prime number as an element in our triangle?
2. Can we find every prime number as an element in our triangle in non-trivial places?

The answer to question 1 is “yes”, as we already saw above (the 1^{st} elements in every row, however, this is a trivial match). To question 2, we fix first that primes are worth looking for only in the middle position.

With a computer investigation (using e.g. the Maple program) we can find 6 small primes up to the 100^{th} row (Figure 2).

Extending the examination up to the 1900^{th} row, we get only one more

Position (row, column)	Prime
2, 2	5
3, 3	13
8, 8	7393
15, 15	65753693
21, 21	175669746209
24, 24	9232029156001

Figure 2: Small primes in the 112-based triangle

positive answer, in position (156, 156), a 90-digit prime (candidate). So, the answer to our second question (considering only this triangle) is “no”.

Our possibilities are extended rapidly, if we look up not only pure prime numbers, but even decompositions. So now we modify our question 2 as “can we find every prime number as a factor of any element in our triangle?” (Examining only non-trivial places, so, positions 0 and 1 are in every row excluded.)

We see immediately that every one-digit prime occurs as a factor at least once up to the 4th row. Here 2 and 5 are triangle elements themselves; 3 is a factor of 9, 7 is a factor of 14.

Continuing with an easy computer examination for two-digit primes we find all but 4 up to the 12th row. For the rest of the numbers we get the following first occurrences (in number–row form): 79–14, 71–15, 59–17 and, surprisingly 41–27.

Now, we turn our attention to 3-digit primes. Here we need a much larger triangle-part. Let’s choose, say, a 100-row triangle in an easy-factorized form. With a small Maple program on a normal table-PC, we can generate the necessary data in a few minutes. (Easy factorization is very important here, otherwise, with full factorization the generation could take an extremely long time...) The output of the program in txt form will be approximately 1.15 MBytes.

From the 143 3-digit primes we find 105 up to 40th row. For the remaining 38 numbers, 18 numbers are situated in rows 41 – 50, 11 additional primes in rows 51 – 60, and 2 (823 and 827) in rows 61 – 70. The still missing “hardest” 3-digit primes finally give the following first occurrences (in number – row form): 479 – 74, 499 – 74, 677 – 76, 719 – 77, 859 – 72, 937 – 98 and 947 – 73. To the contrary, the “easiest” 3-digit primes are 103, 191 and 409 in the 7th row.

With this we give up the claim “to find all of the primes as divisors”.

Our next investigation focuses on very large prime factors (more accurately: prime candidates).

Computer investigations suggest that the largest prime factors in a given row occur very likely in the center position or very close to that place. Of course, this is not an absolute rule, but since our goal is “only” to find very large prime (candidate) factors, we can limit the investigation to the center element. (This has a significant importance to achieving: go as “deep” relatively quickly in the triangle as possible.)

Moreover, the center element carries special properties compared with other elements. Recalling Richard C. Bollinger’s result above, we can set up a similar interesting conjecture:

For large n , the center element in the n^{th} row “almost surely” will be divisible by 5 and 7 (but surely not by 2 and usually not by 3).

So, with a relatively simple Maple program we set out to the easy-factorization of the center element up to the 1900th row. On a normal table-PC, the execution time is approximately 11 hours, with an output file in txt form roughly 110 KBytes.

Analyzing the output we can deduce that prime divisors here follow the Knuth-observation [9], too: we usually find few small factors some of which are repetitive; composite (not decomposable with the ‘easy’ option) large factors are common, pure large prime factors are however rare or extremely rare.

Position (row, column)	Digits of the prime candidate
1726, 1726	1002
1793, 1793	1028
1794, 1794	1030

Figure 3: Large “pure” prime factors (candidates)—112-based triangle, center position

Considering only the primes (prime candidates) with digits more than 1000 we get 3 matches.

Here the second and third matches are especially interesting, since they can be considered as a special kind of “twin-primes” (candidates) in the triangle. In general, our chance to find “pure” large prime factors in consecutive rows is very little...

Here the factorization of element with position 1793, 1793 is as follows:

1793, 1793;“(5) *“(7)² *“(673) *“(65119) *“(1485703) *“(15578887875328
 926423851777567602680378792003694589981499750631818308971422277975
 902867850432471811687112334064063828539296067422531997963055491323
 406425659317001574425151788919713654021679547897110675223861482309
 644220358490739245691930715715021145166205571510978302005857149111
 239471032734380710285002174983967604232152940389858538629493812650
 108566716591594874813194189360195173091031608755605756723631900973
 625032697091409833078265261680211635427069757196618031458397872466
 034789488450265204214587550269112317436588892430166513888148357222
 480962630168478230243146450158020142586939406221546644931686618139
 068737541801842683626194613956159330873776421795220707554672321055
 658602305273678940456712151943459348907356567358277310497505925970
 210070347980231047308886323693790450859256057748541430119354204022
 527748661261790305800487349106563678280226712828838174678186252307
 070941149885645163684441661612796581751766644659424590726902531393
 104098376100305217952214533052008783687240950373043230661705142861
 901235736247002277563333)

In [6] we proved the primality of the largest factor of 1726, 1726 which has 1002 decimal digits. That time we used a freeware software developed by F. Morain. In the remaining part of this paper our goal is to present our selfmade program which is appropriate to prove the prime property of such large numbers. Let us denote the 1028 digits long factor of 1793, 1793 by n_1 and the 1030 digits long factor of 1794, 1794 by n_2 . We investigated n_1 and n_2 with our program, and have found that both of them are really primes. Moreover, the process of the proof and schematic structure of the evidence will be presented, too.

3 Atkin’s primality test

We described the theoretical foundations of the elliptic curve primality proving in [6]. Unfortunately, most computer algebra systems include just probability primality test, so we can not use them to reach our purposes. Although the Magma system (described below) is able to carry out primality proving with ECPP (Elliptic Curve Primality Proving), we did not get any result even after two days running for n_2 . Thus we have developed an own primality proving program presented in the next section.

According to the notation of [6] let us denote an elliptic curve over $\mathbb{Z}/n\mathbb{Z}$ by E_n . The first step in the basic ECPP algorithm is choosing randomly an

E_n elliptic curve, the second one is counting $|E_n|$, the order of E_n . The latter action is very time-consuming, so we had to find an improved version of ECPP. Finally we have implemented an algorithm suggested by A. O. L. Atkin. A specification of this method can be found in [1]. Lenstra and Lenstra published a heuristic running time analysis of Atkin's elliptic curve primality proving algorithm in [10]. They conjectured that with fast arithmetic methods the running time of ECPP can be reduced to $O(\ln^{4+\epsilon}(n))$.

Atkin brilliant idea was founding an appropriate m order in advance and then constructing E_n for this m avoiding the order-counting. Moreover, we get simultaneously two elliptic curves increasing the chance of the successful running of the test. m order has to be chosen from the algebraic integer of an imaginary quadratic field $\mathbb{Q}(\sqrt{D})$. An appropriate D , so-called *fundamental discriminant*, has some properties: $D \equiv 0 \pmod{4}$, or $D \equiv 1 \pmod{4}$, for every $k(> 1)$ D/k^2 is not a fundamental discriminant, $D \leq -7$ and $(D|n) = 1$, where $(D|n)$ is the Jacobi symbol.

The function NEXTD() gives a value D which meets the above mentioned requirements. A given D value is suitable if there exist such $x, y \in \mathbb{Z}$ for which

$$4n = (2x + yD)^2 - y^2D. \quad (1)$$

In that case we get two possible orders: $m = |\nu \pm 1|^2$, where

$$\nu = x + y \frac{D + \sqrt{D}}{2}.$$

If (1) is valid, then we can compute an x_0 root of the *Hilbert polynomial* $(\text{mod } n)$. The function HILBERT(n, D) returns with a root of the appropriate Hilbert polynomial. Then we get two elliptic curves with order $m = |\nu \pm 1|^2$. The rest of the algorithm works as we described in [6].

PROOF(E_n, m, f)

```

1  P ← RANDOMPOINT( $E_n$ )
2  if  $f \cdot P$  is not defined
3    then return COMPOSITE
4  if  $f \cdot P = O$ 
5    then goto 1
6  if  $mP \neq O$ 
7    then return NO
8  return YES
```

Here symbol \mathbf{O} means the “point infinitely far” e.g. the unit of the Abelian group. The function $\text{PROOF}()$ has three input values: E_n , m , f , where E_n is an elliptic curve with order m , $m = f \cdot s$, the factorization of f is known and s is probably prime. The output value COMPOSITE means that n is surely composite. If the output is NO , then n is composite or we have to choose the other elliptic curve. In case YES the next recursion step follows. In the following we present the pseudocode of the Atkin’s test.

$\text{ATKIN-PRIMALITY-TEST}(n)$

```

1   $D \leftarrow \text{NEXTD}()$ 
2   $\omega \leftarrow (D + \sqrt{D})/2$ 
3  if  $\exists x, y \in \mathbb{Z} : 4n = (2x + yD)^2 - y^2D$ 
4    then  $v \leftarrow x + y\omega$ 
5    else goto 1
6   $m \leftarrow |v + 1|^2$ 
7  if  $m = f \cdot s$ , where  $s$  “probably prime” and  $s > (\sqrt[4]{n} + 1)^2$ 
8    then goto 12
9   $m \leftarrow |v - 1|^2$ 
10 if  $m = f \cdot s$  can not be produced so that  $s$  is “probably prime”
    and  $s > (\sqrt[4]{n} + 1)^2$ 
11 then goto 1
12  $x_0 \leftarrow \text{HILBERT}(n, D)$ 
13  $c \leftarrow$  arbitrary integer for which  $(c/n) = -1$ 
14  $k \leftarrow$  arbitrary integer for which  $k \equiv x_0/(1728 - x_0) \pmod{n}$ 
15  $E_n \leftarrow \{(x, y) \mid y^2 = x^3 + 3kx + 2k\}$ 
16 if  $\text{PROOF}(E_n, m, f) = \text{COMPOSITE}$ 
17 then return  $\text{COMPOSITE}$ 
18 else if  $\text{PROOF}(E_n, m, f) = \text{YES}$ 
19 then goto 23
20  $E_n \leftarrow \{(x, y) \mid y^2 = x^3 + 3kc^2x + 2kc^3\}$ 
21 if  $\text{PROOF}(E_n, m, f) = \text{COMPOSITE}$  or  $\text{PROOF}(E_n, m, f) = \text{NO}$ 
22 then return  $\text{COMPOSITE}$ 
23 if  $s$  surely prime
24 then return  $\text{PRIME}$ 
25 else  $\text{ATKIN-PRIMALITY-TEST}(s)$ 

```


4 Magma Computer Algebra System

Magma [5] is a large software system specialized in high-performance computations in number theory, group theory, geometry, combinatorics and other branches of algebra. It was launched at the First Magma Conference on Computational Algebra held at Queen Mary and Westfield College, London, August 1993. It contains a large body of intrinsic functions (implemented in C language), but also allows the user to implement functions on top of this, making use of the Pascal-like user language and the programming environment that is provided.

4.1 Primality tests in Magma

Magma has several built-in functions for primality testing purposes.

`IsProbablyPrime(n: parameter) : RngIntElt ↦ BoolElt`

The function returns TRUE if and only if `n` is a probable prime. This function uses the Miller-Rabin test; setting the optional integer parameter `Bases` to some value `B`, the Miller-Rabin test will use `B` bases while testing compositeness. The default value is 20. This function will never declare a prime number composite, but with very small probability (much smaller than 2^{-B} , and by default less than 10^{-6}) it may fail to find a witness for compositeness, and declare a composite number probably prime.

`IsPrime(n: parameter) : RngIntElt ↦ BoolElt`

This function proves primality using ECPP which is of course more time-consuming. It is possible though to set the optional Boolean parameter `Proof` to FALSE; in which case the function uses the probabilistic Miller-Rabin test, with the default number of bases.

`PrimalityCertificate(n: parameter) : RngIntElt ↦ List`

This function proves primality and provides a certificate for it using ECPP. If the number `n` is proven to be composite or the test fails, a runtime error occurs.

`IsPrimeCertificate(c: parameter) : List ↦ BoolElt`

To verify primality from a given certificate `c` this function is used. This returns the result of the verification by default, a more detailed outcome can be obtained by setting the optional Boolean parameter `ShowCertificate` to TRUE.

The numbers n_1 and n_2 were tested with Magma's own ECPP, using the intrinsic `IsPrime` function, and with our ECPP implementation written in Magma language. We refer to Magma's ECPP algorithm as Magma-ECPP and to our implementation as modified-ECPP. Both tests were running in Magma 2.16 on a machine with 7425 MB RAM and four 2400 MHz Dual-Core AMD Opteron (TM) Processors.

The Magma-ECPP provided a primality proof for n_1 in 32763.52 seconds, but seemed to stuck after the third iteration during the test of n_2 ; the modified-ECPP provided proof for n_1 in 5666.96 seconds and for n_2 in 5153.37 seconds. As the modified-ECPP is not finished yet, the running time can still be improved.

4.2 The implementation of ECPP algorithm

The ECPP algorithm consists of iteration steps, where the i^{th} iteration step outputs an s_i which will be the input of the next iteration step. In one iteration step an attempt is made to factor order m_i of the group of points on a curve E_i . Curve E_i is defined using the input s_{i-1} and a discriminant of an imaginary quadratic field, read in from a list.

If the attempt is successful, factor s_i is the output; if not, we need to backtrack. A different discriminant in an iteration step results in a different s_i . The possible iteration chains that occur this way, can be represented as paths in a directed graph $G(\mathbf{n})$. The nodes of $G(\mathbf{n})$ are the s_i 's, the root represents \mathbf{n} , the edges are the iteration steps. An edge leads from s_i to s_{i+1} if there is an iteration that produces s_{i+1} with input s_i . Consider a path successful if the corresponding iteration-chain starts with input \mathbf{n} and ends with input s_l , where s_l is a small prime, which can be verified by easy inspection, or trial division. In the rest of the paper we refer to the s_i 's also as nodes.

Magma-ECPP uses a small fixed set of discriminants during the process. Each iteration goes through this set until it finds a discriminant which produces a new node. Using a small set of discriminants makes the algorithm faster, but increases the probability of producing no new node. If no discriminant produces new node in the set it backtracks to the previous node and retries that with the same set of discriminants but possibly stronger factorization methods to factor the m_i 's. If backtracking does not produce a new node, it will try to factor again with more effort; these hard factorizations may

consume a large amount of time, and the process appears to get stuck in a seemingly endless loop. This happened during the test of our number n_2 with Magma-ECPP.

4.2.1 Modifications

During the iteration steps certain limits are used; for example, the bound B on the primes found in factoring the m_i -s. Imposing a small B decreases the difference between the size of the s_i -s and thus may extend the path down to the small primes. On the other hand, setting a large B significantly increases the running time needed for factoring. Of course, choosing a more sophisticated factoring method smoothes the differences in running time, but the size of B still remains an important factor. Other important limits are the bound D on the discriminants and the limit S on the prime factors of the discriminants. Decreasing them leads to speed improvement but to a smaller set of discriminants, too.

The modified-ECPP uses a huge file which contains a list of fully factored discriminants up to 10^9 . During the selection of discriminants useful for the current input we extract a modular square root of its prime divisors and build up the square root of the discriminant by multiplication. After using one prime, the square root is stored, and thus it will be computed only once in an iteration step. The speed that we gain this way makes it possible to increase limits D , S in the iterations, which are adjusted to the size of the current input.

The steps can be extended to result in a *series* of s_i -s at a time instead of just a single one: if the iteration step does not stop at the first good discriminant but will collect several good ones. This way, we can select the input of the next step from a set of new nodes.

The numbers have individual properties, which makes a difference from the point of usability. The modified-ECPP predicts the minimal value of D which is still enough to produce at least one new node for each s_i produced by earlier steps and, building upon this prediction, sets up a priority between them. It selects the one with the highest priority as input for the next iteration step. If the step does not provide output the limit D will be increased in order to use a new set of discriminants next time when the node is selected. The priority is reevaluated after each step because either there are new nodes or in case of no output D is increased. This way the possibility of getting stuck is lower (details can be found in [4]).

Table 1: The first rows of the proof of n_1

i	s_i	a_i	b_i	x_i	y_i	f_i
1	165490139	148629518369919	154064198784106	1248188509129	156779851067219	1047222
2	173304931274467	83727826741233492116 021	26748895837956005585 121	23717486180315890605 060	11528948633455951893 545	503211105
3	87208965968598967476 679	27707400957247299977 4111947197	38465375268196111041 0774819991	28142518963869506523 604739355	20626644997054427792 7805954752	6877885
4	59981323890093733168 4100565579	44360713177559177572 547682547718474	11117040242765996352 2375118328811693	19775111407066701253 6508097973496758	18428910703962901519 3733563259937266	408110
5	24478978092786153542 2029989809998131	18957568139328887813 3604105862968021099	10802087315828632193 95669478393483623141	69127839011266503431 9346422505776608550	35593578508679194394 5359772187301402486	4853
6	11879648068429120313 59056929012033432579	42083366473007987927 62804820140778821039 92386685393	28055577648671991951 75203213427185880693 28257790262	18066828358351637326 30229471640748615572 09385839814	25827423550789508626 77731097240769328473 18909337972	486045540848
7	57740499705035302965 17991973651703959752 02732115661	11380443233447151682 41036572010957871951 7294768226496	12206202132034258692 19783734272205524829 2902663774895	96784583503642168856 07646244567179724818 972920682916	59710222084606095711 75786070086466618975 533228021487	24
8	13857719929208472711 77278058794700830584 4118454871693	41694582424591111764 43156453894775152544 45255867194500535484 548	63775518208646414603 03125686908495336379 05480470784855197824 527	34124747439640525997 72929688220778601290 76542652423518714847 638	13433669109612418315 5961778582791580192 42180614563041511515 824	53096907911
9	73580207893761171469 79557764345765335962 72772075969232836325 203	43072369720090946264 13350749758127925795 34831485143675218313 234009372	37871678106813894501 76380954770420227217 88363264525523890880 507401845	70859975965243426606 69499533970811791751 13479474574680812743 326336082	87503943548730935045 05661695045997191182 08741123870099436602 667558083	1271492
10	93556645695254179434 47331860903525804841 40605445293133424460 870453037	18992439425877599779 81729792883189473382 67763448710029974361 4358086171211009277	12661626283918399853 21153195255459648921 78508965806686649574 2905390780807339518	25459301193842798167 48588095939934589673 54516398347837548526 9028646284881593862	20234087314600234645 56960079052485667673 33324851231828927752 7747596369915016860	6762000366

4.3 The proof

On input n , a probable prime, the primality test results in a list, which provides sufficient data to prove the correctness of the sequence of the steps along the successful path. If we consider the length of the proof list as $\#L$, the i^{th} list element, as the proof runs in reverse order, starting from the smallest s_i , corresponds to the $\#L - i^{\text{th}}$ step in the sequence and consists of s_i , a_i , b_i , P_i , f_i , where $s_i f_i = m_i$ and s_i is a probable prime, the factorization of f_i is known, $y^2 = x^3 + a_i x + b_i$ is an elliptic curve of order m_i over $\mathbb{Z}/s_{i+1}\mathbb{Z}$, and P_i is a point on this curve that satisfies the condition $m_i P_i = 0$, $f_i P_i \neq 0$. P_i is given by its two coordinates x_i and y_i . The correctness proof guarantees recursively that all s_i are genuine primes, and eventually that the input n is prime.

Since the size of the above mentioned list is too large (approximately 809 KB in txt form), the exact details can not be presented in this paper. Instead of this, we give here only a small part of this file (see in Table 1). The full text can be downloaded from page: <http://compalg.inf.elte.hu/tanszek/farkasg/proof-tri.txt>

Acknowledgements

Prepared in the framework of application TÁMOP 4.1.1/A-10/1/KONV-2010-0005 with the support of Universitas-Győr Foundation (pages: 158–164); The Project is supported by the European Union and co-financed by the European Social Fund (grant agreement no. TÁMOP 4.2.1/B-09/1/KMR-2010-0003) (pages: 165–171).

References

- [1] A. O. L. Atkin, F. Morain, Elliptic curves and primality proving, *Math. Comp.* **61**, 203 (1993) 29–68. [⇒164](#)
- [2] H. Belbachir, S. Bouroubi, A. Khelladi, Connection between ordinary multinomials, Fibonacci numbers, Bell polynomials and discrete uniform distribution, *Ann. Math. Inform.* **35** (2008) 21–30. [⇒159](#)
- [3] B. A. Bondarenko, *Generalized Pascal Triangles and Pyramids, Their Fractals, Graphs and Applications*, The Fibonacci Association, Santa Clara, CA, USA, 1993. [⇒158, 160](#)

-
- [4] W. Bosma, A. Járari, Gy. Kiss, *Better paths for elliptic curve primality proofs*, <http://www.math.ru.nl/~bosma/pubs/reportfinal.pdf>, 2009. ⇒ 168
- [5] W. Bosma, J. Cannon, C. Playoust, The Magma algebra system. I. The user language, *J. Symbolic Comput.*, **24**, 3-4 (1997) 235–265. ⇒ 166
- [6] G. Farkas, G. Kallós, Prime numbers in generalized Pascal triangles, *Acta Tech. Jaur.* **1**, 1 (2008) 109-118. ⇒ 160, 163, 164
- [7] G. Kallós, *A Pascal háromszög általánosításai* (in Hungarian), Master thesis, Eötvös Loránd University, Budapest, 1993. ⇒ 159
- [8] G. Kallós, A generalization of Pascal’s triangle using powers of base numbers, *Ann. Math. Blaise Pascal* **13**, 1 (2006) 1–15. ⇒ 159, 160
- [9] D. E. Knuth, *The Art of Computer Programming, Vol. 2.*, Addison-Wesley, Reading, MA, USA, 1981. ⇒ 162
- [10] A. K. Lenstra, H. W. Lenstra, Algorithms in number theory, in: *Handbook of Theoretical Computer Science, Vol. A, Algorithms and Complexity*, Ed. J. van Leeuwen, Elsevier Science Publisher, B.V., Amsterdam; MIT Press, Cambridge, MA, 1990, pp. 673–715. ⇒ 164
- [11] *The On-line Encyclopedia of Integer Sequences*, Published electronically at <http://oeis.org>, 2011. ⇒ 159

Received: June 11, 2011 • Revised: September 14, 2011