

# Skyscraper Numbers

Tanya Khovanova  
MIT

Joel Brewster Lewis  
UMN

February 7, 2014

## Abstract

We introduce numbers depending on three parameters which we call *skyscraper numbers*. We discuss properties of these numbers and their relationship with Stirling numbers of the first kind, and we also introduce a skyscraper sequence.

## 1 Introduction

In skyscraper puzzles you have to put an integer from 1 to  $n$  in each cell of a square grid. Integers represent heights of buildings. Every row and column needs to be filled with buildings of different heights and the numbers outside the grid indicate how many buildings you can see from this direction. For example, in the sequence 213645 you can see three buildings from the left (2,3,6) and two from the right (5,6).

In mathematical terminology, we are asked to build a Latin square such that each row is a permutation of length  $n$  with a given number of left-to-right and right-to-left-maxima. The 7-by-7 skyscraper puzzle in Figure 1 is from the Eighth World Puzzle Championship.

Latin squares are notoriously complicated and difficult to enumerate. Thus, in this paper we discuss the combinatorics of a single row instead of asking about the entire puzzle. We give formal definitions and provide examples in Section 2. In Section 3 we give formulas for skyscraper numbers and in Section 4 we define the skyscraper sequence. In Section 5 we prove some properties of skyscraper numbers and the sequence.




Figure 1: A skyscraper puzzle

## 2 Definitions and Examples

What can you say about a row of a skyscraper puzzle if you ignore all information except the numbers given at the two ends of the row? In particular, how many permutations of the buildings are there that satisfy the given restrictions? Of course, the outside numbers have to be between 1 and  $n$ , and we leave as an exercise the proof that their sum must be between 3 and  $n+1$ . Now we consider some of the simplest cases.

Suppose the two given numbers are  $n$  and 1. In this case, the row is completely defined. There is only one possibility: the buildings should be arranged in the increasing order from the side where we see all of them.

Suppose the grid size is  $n$  and the outside numbers are  $a$  and  $b$ . Let us denote the total number of permutations by  $f_n(a, b)$ . We will assume that  $a$  is on the left and  $b$  is on the right. We call these number  $f_n(a, b)$  the *skyscraper numbers* and they are the object of our study.

In the previous example, we showed that  $f_n(n, 1) = 1$ . And of course we have  $f_n(a, b) = f_n(b, a)$ .

Let us discuss a couple of other examples.

First, we want to discuss the case when the sum of the border numbers is

the smallest, namely 3. In this case,  $f_n(1, 2)$  is  $(n - 2)!$ . Indeed, we need to put the tallest building on the left and the second tallest on the right. After that we can permute the leftover buildings any way we want.

Second, we want to discuss the case when the sum of the border numbers is the largest:  $n + 1$ . In this case  $f_n(a, n + 1 - a)$  is  $\binom{n-1}{a-1}$ . Indeed, the position of the tallest building is uniquely defined — it has to take the  $a$ -th spot from the left. After that we can pick a set of  $a - 1$  buildings that go to the left from the tallest building and the row is uniquely defined by this set.

## 3 Skyscraper Numbers

### 3.1 One-sided skyscraper numbers

Before going further let us see what happens if only one of the maxima is given. Let us denote by  $c(n, a)$  the number of permutations of  $n$  buildings so that you can see  $a$  buildings from the left. If we put the shortest building on the left then the leftover buildings need to be arranged so that you can see  $a - 1$  of them. If the shortest building is not on the left, then it can be in any of the  $n - 1$  places and we still need to rearrange the leftover buildings so that we can see  $a$  of them. We just proved that the function  $c(n, a)$  satisfies the recurrence

$$c(n, a) = c(n - 1, a - 1) + (n - 1)c(n - 1, a) = \sum_{k=1}^{n-1} \frac{(n - 1)!}{(n - k - 1)!} c(n - k, a - 1).$$

Actually  $c(n, a)$  is a well-known function. The numbers  $c(n, a)$  are called *unsigned Stirling numbers of the first kind* (see sequence A132393 in the Online Encyclopedia of Integer Sequences [1]); not only do they count permutations with a given number of left-to-right (or right-to-left) maxima, but they also count permutations with a given number of cycles, and they appear as the coefficients in the product  $(x + 1)(x + 2)(x + 3) \dots (x + n)$ , among other places; see for example [2, Chapter 1].

### 3.2 The tallest building is on the left

We are now equipped to calculate  $f_n(1, b)$ . The tallest building must be on the left, and the rest could be arranged so that, in addition to the tallest building,  $b - 1$  more buildings are seen from the right. That is,  $f_n(1, b) = c(n - 1, b - 1)$ .

	$b=2$	$b=3$	$b=4$	$b=5$	$b=6$	$b=7$
$n=2$	1					
$n=3$	1	1				
$n=4$	2	3	1			
$n=5$	6	11	6	1		
$n=6$	24	50	35	10	1	
$n=7$	120	274	225	85	15	1

Table 1: The Stirling numbers  $c(n - 1, b - 1) = f_n(1, b)$

The non-zero values of  $f_n(1, b)$  are collected in Table 1.

### 3.3 Skyscraper numbers

Now we have everything we need to consider the general case. In any permutation of length  $n$ , the left-to-right maxima consist of  $n$  and all left-to-right maxima that lie to its left; similarly, the right-to-left maxima consist of  $n$  and all the right-to-left maxima to its right. We can take any permutation counted by  $f_n(a, b)$  and split it into two parts: if the value  $n$  is in position  $k + 1$  for some  $0 \leq k \leq n - 1$ , the first  $k$  values form a permutation with  $a - 1$  left-to-right maxima and the last  $n - k - 1$  values form a permutation with  $b - 1$  right-to-left maxima, and there are no other restrictions. Thus,

$$f_n(a, b) = \sum_{k=0}^{n-1} \binom{n-1}{k} c(k, a-1) \cdot c(n-k-1, b-1).$$

Table 2 shows  $f_7(a, b)$ , of which we already calculated the first row:

## 4 The Skyscraper Sequence

We see that the first two rows of the puzzle above (see Figure 1) have the same pair of numbers, namely 2 and 3, outside. If we ignore all other constraints there are 675 ways to fill in each of the first two rows. Number 675 is the largest number in the Table 2. We can say that these two rows of the puzzle are the most difficult to fill in: the pair of numbers 2 and 3 is the least restrictive. Given  $n$ , we call such a pair the *maximizing pair*.



building consider a group of buildings that are immediately after it to the right and that are located before the next left-to-right maximum. Choose  $a - 1$  left-to-right maxima out of them and move them together with their groups to the right and put them in the reverse order.  $\square$

We want to add a different proof of this lemma through Stirling numbers.

*Proof.* We want to show that  $f_n(a, b) = \sum_{k=0}^{n-1} \binom{n-1}{k} \cdot c(k, a-1) \cdot c(n-k-1, b-1)$  is equal to  $\binom{a+b-2}{a-1} \cdot c(n, a+b-2)$ . We can do this using the fact that  $c(n, a+b-2)$  counts also permutations of length  $n-1$  with  $a+b-2$  cycles. Let us count permutations of length  $n-1$  with  $a+b-2$  cycles, of which  $a-1$  are colored red and  $b-1$  are colored blue, in two different ways: on one hand, we may first choose a permutation of length  $n-1$  with  $a+b-2$  cycles, then choose  $a-1$  of these cycles to be red, in  $\binom{a+b-2}{a-1} \cdot c(n-1, a+b-2)$  ways. On the other hand, for any  $k$  between 0 and  $n-1$  we may first choose  $k$  elements to be in red cycles, then make a permutation on these  $k$  entries with exactly  $a-1$  cycles, and make a permutation with exactly  $b-1$  cycles on the remaining  $n-k-1$  entries. Thus we have in total  $\sum_{k=0}^{n-1} \binom{n-1}{k} \cdot c(k, a-1) \cdot c(n-k-1, b-1)$  ways in this case. So the two things are equal, as claimed.  $\square$

Now we are ready to prove the promised theorem about maximizing pairs for the skyscraper sequence.

**Theorem 2.** *The numbers in the maximizing pairs do not differ by more than 1.*

*Proof.* The maximum number in a table corresponding to the given  $n$  is the maximum number in its line parallel to antidiagonal. As each diagonal is proportionate to a row in the Pascal's triangle, the maximum can be found in its middle.  $\square$

Now let us sum all numbers in each row of Table 2. We get the following sums: 720, 1764, 1624, 735, 175, 21, 1. These numbers bring us back to the sequence A132393 in the OEIS [1]. These numbers are a row in the triangle of unsigned Stirling numbers of the first kind. This brings us to the next lemma:

**Lemma 3.**  $\sum_b f_n(a, b) = f_{n+1}(a+1, 1)$ .

*Proof.* Consider a row of length  $n + 1$  with  $a + 1$  left-to-right maxima and 1 right-to-left maximum. In this configuration the tallest building is on the right. If we remove the tallest building, then we get a row of length  $n$  with  $a$  left-to-right maxima and an unknown number of right-to-left maxima.  $\square$

Skyscraper numbers are fun to play with. We are sure that they are hiding many more secrets.

## References

- [1] N. J. A. Sloane, Online Encyclopedia of Integer Sequences (OEIS). <http://www.research.att.com/~njas/sequences/>
- [2] Richard P. Stanley, *Enumerative Combinatorics, Volume 1*. Cambridge University Press, 2012.

---

2010 *Mathematics Subject Classification*: Primary 11B83; Secondary 11Y55.  
*Keywords*: sequences, skyscraper puzzle, Stirling numbers.

---

(Mentions A132393, A218531.)

---