

Refining enumeration schemes to count according to permutation statistics

Andrew M. Baxter

January 3, 2014

Abstract

We consider the question of computing the distribution of a permutation statistics over restricted permutations via enumeration schemes. The restricted permutations are those avoiding sets of vincular patterns (which include both classical and consecutive patterns), and the statistics are described in the number of copies of certain vincular patterns such as the descent statistic and major index. An enumeration scheme is a polynomial-time algorithm (specifically, a system of recurrence relations) to compute the number of permutations avoiding a given set of vincular patterns. Enumeration schemes' most notable feature is that they may be discovered and proven via only finite computation. We prove that when a finite enumeration scheme exists to compute the number of permutations avoiding a given set of vincular patterns, the scheme can also compute the distribution of certain permutation statistics with very little extra computation.

1 Introduction

Enumeration schemes are special recurrences to compute the number of permutations avoiding a set of vincular patterns. In this paper, we discuss how to refine enumeration schemes to compute the distributions of certain permutation statistics over a set of pattern-avoiding permutations. This extends previous work in [6, 9] which considers the distribution of only the inversion number.

Let $[n]$ be shorthand for the set $\{1, \dots, n\}$. For a word $w \in [n]^k$, we write $w = w_1 w_2 \cdots w_k$ and define the *reduction* $\text{red}(w)$ to be the word obtained by replacing the i^{th} smallest letter(s) of w with i . For example $\text{red}(839183) = 324132$. If $\text{red}(w) = \text{red}(w')$, we say that w and w' are *order-isomorphic* and write $w \sim w'$. We will commonly use the notation $|w|$ to denote the length of w .

Vincular patterns resemble classical patterns, with the constraint that some of the letters in a copy must be consecutive. Formally, a *vincular pattern of length k* is a pair (σ, X) where σ is a permutation in \mathfrak{S}_k and $X \subseteq \{0, 1, 2, \dots, k\}$ is a set of “adjacencies.” A permutation $\pi \in \mathfrak{S}_n$ *contains the vincular pattern* (σ, X) if there is a k -tuple $1 \leq i_1 < i_2 < \cdots < i_k \leq n$ such that the following three criteria are satisfied:

- $\text{red}(\pi_{i_1}\pi_{i_2}\cdots\pi_{i_k}) = \sigma$.
- $i_{x+1} = i_x + 1$ for each $x \in X \setminus \{0, k\}$.
- $i_1 = 1$ if $0 \in X$ and $i_k = n$ if $k \in X$.

In the present work we restrict our attention to patterns (σ, X) where $\sigma \in \mathfrak{S}_k$ and $X \subseteq [k-1]$, rendering the third containment criterion irrelevant.¹ The subsequence $\pi_{i_1}\pi_{i_2}\cdots\pi_{i_k}$ is called a *copy* of (σ, X) . In the permutation $\pi = 162534$, the subsequence 1253 is a copy of $(1243, \{3\})$, but the subsequence 1254 is not a copy since the 5 and 4 are not adjacent in π . The “classical pattern” σ is precisely the vincular pattern (σ, \emptyset) since no adjacencies are required, while the “consecutive pattern” σ is the vincular pattern $(\sigma, \{1, 2, \dots, k-1\})$ since all internal adjacencies are required.

In practice we write (σ, X) as a permutation with a dash between σ_j and σ_{j+1} if $j \notin X$. For example, $(1243, \{3\})$ is written 1-2-43. We occasionally refer to “the vincular pattern σ ” or even “the pattern σ ” without explicitly referring to X .

If the permutation π does not contain a copy of the pattern (σ, X) , then π is said to *avoid* σ . We will notation $\mathfrak{S}_n(\sigma)$ or $\mathfrak{S}_n((\sigma, X))$ to denote the set of permutations avoiding the (σ, X) , and $\mathfrak{S}_n(B)$ denotes those permutations avoiding every vincular pattern $(\sigma, X) \in B$.

Observe that a vincular pattern (σ, X) of length k exhibits similar symmetries to those of permutations, except for taking inverses. The reverse is given by $(\sigma, X)^r = (\sigma^r, k-X)$ where $k-X = \{k-x : x \in X\}$. For example, $(1-3-42)^r = 24-3-1$. The complement is $(\sigma, X)^c = (\sigma^c, X)$. For example, $(1-3-42)^c = 4-2-13$. It follows that that π avoids (σ, X) if and only if π^r avoids $(\sigma, X)^r$. Similarly, π avoids (σ, X) if and only if π^c avoids $(\sigma, X)^c$.

See Steingrímsson’s survey for a fuller history in [24]. From their earliest days, vincular patterns been linked to many of the common combinatorial structures such as set partitions and lattice paths in [14] as well as permutation statistics in [2].

Enumeration schemes were introduced by Zeilberger in [27] as an automated method to compute $|\mathfrak{S}_n(B)|$ for many different B . Vatter improved schemes in [25] with the introduction of gap vectors, and Zeilberger provided an alternate implementation in [28]. The greatest feature of schemes is that they may be discovered by a computer: the user need only input the set B (along with bounds to the computer search) and the computer will return an enumeration scheme (if one exists within the bounds of the search) which computes $|\mathfrak{S}_n(B)|$ in polynomial time. Pudwell extended these methods to consider pattern avoidance in permutations of a multiset in [22, 20], as well as barred-pattern avoidance in [21]. The author and Pudwell extended schemes to sets of vincular patterns in [9].

A *permutation statistic* is any function $f : \bigcup_{n \geq 0} \mathfrak{S}_n \rightarrow \mathbb{Z}$. The most-studied permutation statistic is the *inversion number* $\text{inv}(\pi) = |\{(i, j) : i < j \text{ and } \pi_i > \pi_j\}|$. In terms of vincular patterns, $\text{inv}(\pi)$ is the number

¹We enact this restriction partly for simplicity. It is likely that the prefix-focused arguments in [9] and below extend to patterns (σ, X) with $0 \in X$ with few modifications, but it is unlikely such an approach could work for patterns with $k \in X$.

of copies of 2-1 = (21, \emptyset). Similarly, the *descent number* $\text{des}(\pi) = |\{i : \pi_i > \pi_{i+1}\}|$ is the number of copies of 21 = (21, {1}). In this work we will primarily consider permutation statistics which count the number of copies of a given vincular pattern, sometimes called *pattern functions*. It is shown in [2] that many well-known permutation statistics can be framed as linear combinations of pattern functions. For a permutation statistic f and set $S \subseteq \bigcup_{n \geq 0} \mathfrak{S}_n$, the *distribution of f over S* is given by:

$$F(S, f, q) := \sum_{\pi \in S} q^{f(\pi)} \quad (1)$$

We also consider the simultaneous distribution of multiple statistics $\mathbf{f} = \langle f_1, \dots, f_m \rangle$ over the same set S with the indeterminates $\mathbf{q} = \langle q_1 \dots, q_m \rangle$. The distribution of \mathbf{f} over S is given by:

$$F(S, \langle f_1, \dots, f_m \rangle, \langle q_1, \dots, q_m \rangle) := \sum_{\pi \in S} q_1^{f_1(\pi)} q_2^{f_2(\pi)} \dots q_m^{f_m(\pi)}. \quad (2)$$

Distributions of statistics over sets of pattern-avoiding permutations have received increased attention of late, focusing primarily on sets of permutations avoiding classical patterns of length 3. Barucci et al. use generating trees to study the inversion number over $\mathfrak{S}_n(B)$ for a few examples of sets B in [3]. Barnabei et al. study copies of consecutive patterns over $\mathfrak{S}_n(1\text{-}2\text{-}3)$ and $\mathfrak{S}_n(3\text{-}1\text{-}2)$ in [4, 5]. Dokos et al. refine Wilf-equivalence in [15] by studying the inversion number and major index over $\mathfrak{S}_n(\tau)$ for classical patterns $\tau \in \mathfrak{S}_3$. Bona and Homberger study the total number of classical patterns $\sigma \in \mathfrak{S}_3$ over $\mathfrak{S}_n(\tau)$ for another classical pattern $\tau \in \mathfrak{S}_3$ in [11, 12, 16]. Most recently, Burstein and Elizalde in [13] study the total number of vincular patterns of length 3 over $\mathfrak{S}_n(\tau)$ for classical patterns $\tau \in \mathfrak{S}_3$.

Suppose that E is a finite enumeration scheme which gives a recurrence to compute $|\mathfrak{S}_n(B)|$ for a given set of patterns B . The work in [6, 9] demonstrates how to use E to compute $F(\mathfrak{S}_n(B), \text{inv}, q)$. The present work demonstrates how to use E to compute the distribution $F(\mathfrak{S}_n(B), \mathbf{f}, \mathbf{q})$ where each statistic f_i counts the number of copies of a vincular pattern of the form $(\sigma, [|\sigma| - 1])$ or $(\sigma, [|\sigma| - 2])$ or counts the number of right-to-left minima or right-to-left maxima. The results are implemented in the Maple package **Statter**, available for download from the author’s homepage.

The paper is organized as follows. Section 2 outlines the basics of enumeration schemes and their structure. Section 3 defines the notion of an “enumeration-scheme-compatible,” or “ES-compatible,” statistic. Subsection 3.2 presents three classes of ES-compatible statistics. Section 4 presents a technical result proving that any given finite enumeration scheme can be expanded to fit the additional requirements which ES-compatible statistics can impose. Section 5 presents three specific examples of how enumeration schemes can be applied to explore statistics over sets $\mathfrak{S}_n(B)$.

2 Overview of Enumeration Schemes

Enumeration schemes are succinct encodings for a family of recurrence relations enumerating a family of sets. The enumerated sets are actually subsets of $\mathfrak{S}_n(B)$ determined by prefixes.

For pattern $p \in \mathfrak{S}_k$, let $\mathfrak{S}_n(B)[p]$ be the set of permutations $\pi \in \mathfrak{S}_n(B)$ such that $\text{red}(\pi_1\pi_2 \dots \pi_k) = p$. We call p the *prefix pattern*. To refine further, let $w \in [n]^k$ and define $\mathfrak{S}_n(B)[p; w]$ to be those permutations in $\pi \in \mathfrak{S}_n(B)[p]$ such that $\pi_1\pi_2 \dots \pi_k = w$. For example,

$$\mathfrak{S}_5(1\text{-}2\text{-}3)[21; 53] = \{53142, 53214, 53241, 53412, 53421\}.$$

Since we are interested in enumeration, it will be handy to have the notation $s_n(B)[p] = |\mathfrak{S}_n(B)[p]|$ and $s_n(B)[p; w] = |\mathfrak{S}_n(B)[p; w]|$.

By looking at the prefix of a permutation, one can identify likely “trouble spots” where forbidden patterns may appear. For example, suppose we wish to avoid the (classical) pattern 1-2-3. Then the presence of the pattern 12 in the prefix indicates the potential for the whole permutation to contain a 1-2-3 pattern.

Enumeration schemes take a divide-and-conquer approach to enumeration. We define the *child* of a permutation $p \in \mathfrak{S}_k$ to be any permutation $p' \in \mathfrak{S}_{k+1}$ such that $\text{red}(p'_1p'_2 \dots p'_k) = p$. Any $\mathfrak{S}_n(B)[p]$ for $p \in \mathfrak{S}_k$ may be partitioned into the family of sets $\mathfrak{S}_n(B)[p']$ for each of its children $p' \in \mathfrak{S}_{k+1}(B)[p]$. The sets indexed by these children are then considered as described below, and their sizes are totaled to obtain $s_n(B)[p]$. In the end we have computed $|\mathfrak{S}_n(B)|$, since $\mathfrak{S}_n(B) = \mathfrak{S}_n(B)[\epsilon] = \mathfrak{S}_n(B)[1]$, where ϵ is the empty (i.e., length 0) permutation.

For $p \in \mathfrak{S}_k$ a set $\mathfrak{S}_n(B)[p]$ fits into one of three cases:

- (1) If $n = k$, then $\mathfrak{S}_n(B)[p]$ is either $\{p\}$ or \emptyset , depending on whether p avoids B .
- (2) For each $w \in [n]^k$ such that $\text{red}(w) = p$, one of the following happens:
 - (2a) $\mathfrak{S}_n(B)[p; w]$ is empty, and so $s_n(B)[p; w] = 0$.
 - (2b) $\mathfrak{S}_n(B)[p; w]$ is in bijection with some other $\mathfrak{S}_{\hat{n}}(B)[\hat{p}; \hat{w}]$ for $\hat{n} < n$, and so $s_n(B)[p; w] = s_{\hat{n}}(B)[\hat{p}; \hat{w}]$.
- (3) $\mathfrak{S}_n(B)[p]$ must be partitioned further, so $s_n(B)[p] = \sum_{p' \in \mathfrak{S}_{k+1}(B)[p]} s_n(B)[p']$.

Case (1) provides the base cases for our recurrence. If case (2) applies, then we will use it preferentially over case (3). If case (2) does not apply, we must divide $\mathfrak{S}_n(B)[p]$ as in case (3). Determining whether case (2) applies makes use of *gap vector criteria* to test (2a) and *reversible deletions* to form the bijection in (2b). These concepts are outlined in the following subsections.

2.1 Gap Vectors

The differences between the values of letters in the prefix may be great that a forbidden pattern *must* appear. To make this more precise, we follow our example above and compute $s_n(1\text{-}2\text{-}3)[12]$. Observe that $\mathfrak{S}_n(1\text{-}2\text{-}3)[12; w_1w_2]$ is empty if $w_1 < w_2 < n$, since otherwise if $\pi \in \mathfrak{S}_n(1\text{-}2\text{-}3)[12; w_1w_2]$ then

$\pi_i = n$ for some $i \geq 3$ and so $w_1 w_2 n$ forms a 1-2-3 pattern. Since the possibility for any $\pi_i > w_2$ for $i \geq 3$ prohibits the formation of a 1-2-3-avoiding permutation, we must restrict the space above w_2 .

To formalize this, consider $\mathfrak{S}_n(B)[p; w]$ and let c_i be the i^{th} smallest letter in w . Let $c_0 = 0$ and $c_{k+1} = n + 1$, and form the $(k + 1)$ -vector $\vec{g}(n, w)$ so that the i^{th} component is $g_i = c_i - c_{i-1} - 1$. Note that g_i counts the number of letters for any $\pi \in \mathfrak{S}_n(B)[p; w]$ which lie strictly between c_{i-1} and c_i , i.e., the number of letters π_j following the prefix ($j > k$) and $c_{i-1} \leq \pi_j \leq c_i$. We call $\vec{g}(n, w)$ the *spacing vector* for w .

In the example above, if $\vec{g}(n, w) \geq \langle 0, 0, 1 \rangle$ in the product order of \mathbb{N}^3 (i.e., component-wise), then $\mathfrak{S}_n(1-2-3)[12; w] = \emptyset$. We call $\langle 0, 0, 1 \rangle$ a gap vector for the prefix 12. More generally we may make the following definition:

Definition 1. *Given a set of forbidden patterns B and prefix p , then \vec{v} is a gap vector for prefix p with respect to B if, for all n , $\mathfrak{S}_n(B)[p; w] = \emptyset$ for any w such that $\vec{g}(n, w) \geq \vec{v}$. In this case we say that w satisfies the gap vector criterion for \vec{v} .*

Hence $\vec{v} = \langle 0, 0, 1 \rangle$ is a gap vector for $p = 12$ with respect to $B = \{1-2-3\}$, and any prefix set $w = w_1 w_2$ with $w_1 < w_2 < n$ satisfies the gap vector condition for v .

Observe that gap vectors for a given prefix $p \in \mathfrak{S}_k$ form an upper order ideal in \mathbb{N}^{k+1} , since if \vec{v} is a gap vector so is any $\vec{u} \geq \vec{v}$. Hence it suffices to determine only the minimal elements (which form a basis). For details on the discovery of gap vectors and automating the process, see [25, 28, 9].

Note that if the prefix p contains a pattern in B , then $\mathfrak{S}_n(B)[p; w] = \emptyset$ for any appropriate w , and so $\vec{0} = \langle 0, 0, \dots, 0 \rangle$ is a gap vector.

2.2 Reversible Deletability

When w fails the gap vector criterion for all gap vectors \vec{v} , we must rely on bijections with previously-computed $\mathfrak{S}_n(B)[\hat{p}; \hat{w}]$. To continue our example above, consider $\mathfrak{S}_n(1-2-3)[12; w_1 n]$. Here $w_1 n$ fails all gap vector criteria, because $\langle 0, 0, 1 \rangle$ forms the basis for the ideal of gap vectors and $\vec{g}(n, w_1 n) = \langle w_1 - 1, n - w_1 - 1, 0 \rangle \not\geq \langle 0, 0, 1 \rangle$. However, any $\pi \in \mathfrak{S}_n(1-2-3)[12; w_1 n]$ has $\pi_2 = n$, so we may use the map $d_2 : \pi_1 \pi_2 \dots \pi_n \mapsto \text{red}(\pi_1 \pi_3 \dots \pi_n)$ to form a bijection $\mathfrak{S}_n(1-2-3)[12; w_1 n] \rightarrow \mathfrak{S}_{n-1}(1-2-3)[1; w_1]$. The deletion of a letter always preserves pattern-avoidance properties when considering classical patterns, but inverting the map by inserting a letter has the potential for creating a forbidden pattern. Here, however, inserting an n at the second index cannot possibly create a 1-2-3, so we may safely reverse the deletion.

More generally define the deletion $d_r(\pi) := \text{red}(\pi_1 \dots \pi_{r-1} \pi_{r+1} \dots \pi_n)$, that is, the permutation obtained by omitting the r^{th} letter of π and reducing. Furthermore for a set R , define $d_R(\pi)$ to be the permutation obtained by deleting π_r for each $r \in R$ and then reducing. For a word w with no repeated letters, define $d_r(w)$ be the word obtained by deleting the r^{th} letter and then subtracting 1 from each remaining letter larger than w_r . Similarly, to construct $d_R(w)$ delete w_r for each $r \in R$ and

subtract $|\{r \in R : w_r < w_i\}|$ from each remaining w_i . For example $d_3(6348) = 537$ and $d_{\{1,3\}}(6348) = 36$. It can be seen that this definition is equivalent to the one given above when $w \in \mathfrak{S}_k$, and it allows for more succinct notation in the upcoming definition. In the unrestricted case, $d_R : \mathfrak{S}_n(\emptyset)[p; w] \rightarrow \mathfrak{S}_{n-|R|}(\emptyset)[d_R(p); d_R(w)]$ is a bijection for any set $R \subseteq [|p|]$. Sometimes we are lucky and the restriction to $\mathfrak{S}_n(B)[p; w]$ is a bijection with $\mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)]$, leading to the following definition:

Definition 2. *The set of indices R is reversibly deletable for p with respect to B if the map*

$$d_R : \mathfrak{S}_n(B)[p; w] \rightarrow \mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)]$$

is a bijection for all words w failing the gap vector criterion for every gap vector of p with respect to B (i.e., d_R is a bijection for all w such that $\mathfrak{S}_n(B)[p; w] \neq \emptyset$).

Note that the empty set $R = \emptyset$ is reversibly deletable for any p and B , but is uninteresting. Additionally, if $\vec{0}$ is a gap vector then any set $R \subseteq [|p|]$ is vacuously reversibly deletable since $\mathfrak{S}_n(B)[p; w] = \emptyset$ for any prefix w .

Proving that a set is reversibly deletable for prefix p with respect to B can be carried out by a finite list of verifications, and thus can be done via computer. This is proven in [25] for the case that B contains only classical patterns, and in [9] in the case that B contains vincular patterns. The process of automated discovery itself is not relevant to the present work and will be omitted.

2.3 Formal Definition of Enumeration Schemes

We will now formally define an enumeration scheme.

Definition 3. *Let B be a set of vincular patterns. An enumeration scheme for B is a set E of triples (p, G, R) , where p is a permutation (i.e., , the prefix pattern), G is a basis of gap vectors for p with respect to B , and R is a reversibly deletable set for p with respect to B . Furthermore, E must satisfy the following criteria:*

1. $(\epsilon, \emptyset, \emptyset) \in E$.
2. For each $(p, G, R) \in E$,
 - (a) If $R = \emptyset$ and $\vec{0} \notin G$, then there exists a triple $(p', G', R') \in E$ for each child p' of p .
 - (b) If $R \neq \emptyset$, then there exists a triple $(\hat{p}, \hat{G}, \hat{R}) \in E$ for $\hat{p} = d_R(p)$.

To compute $s_n(B)[p; w]$ for a fixed n , p , and w , the enumeration scheme E is “read” by finding the appropriate triple $(p, G, R) \in E$ and concluding:

1. If w satisfies the gap vector criteria for some $\vec{v} \in G$, then $s_n(B)[p; w] = 0$.
2. If w fails the gap criteria for all $\vec{v} \in G$ and $R \neq \emptyset$, then $s_n(B)[p; w] = s_{n-|R|}(B)[d_R(p); d_R(w)]$.

3. If w fails the gap criteria for all $\vec{v} \in G$ and $R = \emptyset$, then $s_n(B)[p] = \sum_{p' \in \mathfrak{S}_{k+1}(B)[p]} s_n(B)[p']$.

When combined with the initial conditions that $s_n(B)[p; w] = 1$ whenever p has length n and avoids B , the scheme provides a system of recurrences to compute $s_n(B)[p; w]$ and ultimately $|\mathfrak{S}_n(B)|$.

To illustrate, consider the enumeration scheme for $\mathfrak{S}_n(1-2-3)$:

$$\{(\epsilon, \emptyset, \emptyset), (1, \emptyset, \emptyset), (12, \{(0, 0, 1)\}, \{2\}), (21, \emptyset, \{1\})\} \quad (3)$$

Since $R_\epsilon = \emptyset$, the first condition above requires the presence of $(1, G_1, R_1)$. Starting with the pattern 1 yields no additional information, so $R_1 = \emptyset$ and thus explaining the presence of $(12, G_{12}, R_{12})$ and $(21, G_{21}, R_{21})$. As discussed above, $\{(0, 0, 1)\}$ forms a basis for the gap vectors for 12, and whenever w fails this gap vector criteria the second letter is reversibly deletable. For the fourth entry in the scheme, suppose that $\pi \in \mathfrak{S}_n(\emptyset)[21]$ contains a 1-2-3 pattern involving the first letter, say $\pi_1 < \pi_i < \pi_j$ for $i < j$. Then since $\pi_2 < \pi_1$, we see that $\pi_2 < \pi_i < \pi_j$ is another 1-2-3 pattern. Therefore π_1 cannot be the deciding factor for whether π contains 1-2-3. Hence the index 1 is reversibly deletable, so $R_{21} = \{1\}$.

Enumeration schemes exhibit a tree-like structure. The empty prefix ϵ serves as the root, and the children of each prefix are drawn as children in a rooted tree. When a prefix has nontrivial gap vector criteria, we list those basis vectors below it. When prefix p has a non-empty reversibly deletable set R , we draw an arrow from p to $d_R(p)$ labeled with “ d_R ”. See Figure 1 for an example.




Figure 1: Tree representation of the enumeration scheme for $\mathfrak{S}_n(1-2-3)$

If $|E|$ is finite, we say that B admits a finite enumeration scheme. A finite enumeration scheme gives us a polynomial-time algorithm to compute $s_n(B)[p; w]$. We construct the system of recurrences based on the partitions and bijections above, along with base cases as given by the gap vector criteria and the trivial cases when $\mathfrak{S}_n(B)[p] = \{p\}$ or \emptyset . For example, the above enumeration scheme in (3) translates into the following system of recurrences:

$$\begin{aligned}
|\mathfrak{S}_n(1-2-3)| &= s_n(1-2-3)[\epsilon] \\
&= s_n(1-2-3)[1] \text{ when } n > 0 \\
&= \sum_{i=1}^n s_n(1-2-3)[1; i] \\
s_n(1-2-3)[1; i] &= \sum_{j=1}^{i-1} s_n(1-2-3)[21; ij] + \sum_{j=i+1}^n s_n(1-2-3)[12; ij] \quad (4) \\
s_n(1-2-3)[12; ij] &= \begin{cases} 0 & \text{if } n-j \geq 1 \\ s_n(1-2-3)[1; i] & \text{otherwise} \end{cases} \\
s_n(1-2-3)[21; ij] &= s_n(1-2-3)[1; j]
\end{aligned}$$

The recurrences in (4) simplify to create the following recurrence:

$$s_n(1-2-3)[1; i] = \sum_{j=1}^i s_{n-1}(1-2-3)[1; j] \quad (5)$$

Among other things, one can then evaluate this recurrence by hand to identify the closed form $|\mathfrak{S}_n(1-2-3)| = \frac{1}{n+1} \binom{2n}{n}$, the Catalan numbers.

The length of the longest prefix p appearing in finite scheme E is called the *depth* of E . Not every set B admits a finite enumeration scheme, the simplest example being the classical pattern 2-3-1. Let E be the scheme for $\mathfrak{S}_n(2-3-1)$, and let $J_t = t(t-1)\cdots 21$ be the decreasing permutation of length t . It can be shown that for any t there are no gap vectors for J_t and no non-empty reversibly deletable sets. Hence E must contain the triple $(J_t, \emptyset, \emptyset)$ for each $t \geq 1$ and hence E is infinite.

It should be noted that the enumeration scheme for $\mathfrak{S}_n(1-3-2)$ is finite (of depth 2) and $|\mathfrak{S}_n(2-3-1)| = |\mathfrak{S}_n(1-3-2)|$ by symmetry. More generally, it can be seen that if B admits an enumeration scheme E_B of depth K then its set of complements $B^c = \{\sigma^c : \sigma \in B\}$ also admits an enumeration scheme E_{B^c} of depth K . The analogous statements regarding $B^r = \{\sigma^r : \sigma \in B\}$ do not hold and so B may not have a finite scheme while B^r does, as exhibited by $B = \{2-3-1\}$.

3 Enumeration-scheme-compatible statistics

3.1 Definitions and interaction with enumeration schemes

The author proves in [6] that if B admits a finite enumeration scheme, then the distribution of the statistic $\text{inv}(\pi)$ over $\mathfrak{S}_n(B)$ can be computed via the same enumeration scheme. This is the consequence of comparing the inversion number of a permutation and its image under the deletion map $d_R : \mathfrak{S}_n(B)[p; w] \rightarrow \mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)]$. In particular, for $\pi \in \mathfrak{S}_n$

the change in the inversion number after deleting the r^{th} letter is given by

$$\delta_r^{\text{inv}}(\pi) := \text{inv}(\pi) - \text{inv}(d_r(\pi)) = (\pi_r - 1) + \sum_{i < r} \text{sgn}(\pi_i - \pi_r), \quad (6)$$

where $\text{sgn}(x)$ is the signum function:

$$\text{sgn}(x) := \begin{cases} -1, & x < 0 \\ 0, & x = 0 \\ 1, & x > 0. \end{cases}$$

For the more general case, let $R = \{r_1, \dots, r_t\}$ where $r_j < r_{j+1}$. Then the deletion d_R has the following effect on inversion number:

$$\delta_R^{\text{inv}}(\pi) := \text{inv}(\pi) - \text{inv}(d_R(\pi)) = \sum_{r \in R} \delta_r(\pi) - \text{inv}(\pi_{r_1} \cdots \pi_{r_t}) \quad (7)$$

Observe that $\delta_R(\pi)$ can be written purely in terms of the letters π_1, \dots, π_{r_t} . Therefore if E is an enumeration scheme for pattern set B , and $(p, G, R) \in E$, then δ_R is constant over the set $\mathfrak{S}_n(B)[p; w]$ for any fixed w . Thus we define a new function $\Delta_R^{\text{inv}}(w, n)$ on prefix words w , which takes on the value $\delta_R^{\text{inv}}(\pi)$ given by a $\pi \in \mathfrak{S}_n(B)[p; w]$. Therefore we may recursively compute the distribution of inv over $\mathfrak{S}_n(B)[p; w]$ via

$$F(\mathfrak{S}_n(B)[p; w], \text{inv}, q) = q^{\Delta_R^{\text{inv}}(w, n)} F(\mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)], \text{inv}, q) \quad (8)$$

The above results motivate the following definitions:

Definition 4. Let $f : \bigcup_{n \geq 0} \mathfrak{S}_n \rightarrow \mathbb{Z}$ be a permutation statistic. For a permutation π let the R -deletion difference, denoted $\delta_R^f(\pi)$, be $f(\pi) - f(d_R(\pi))$.

For nonnegative integer m , a permutation statistic f is enumeration-scheme-compatible (or “ES-compatible”) with margin m if for any positive integer t and any $R \subseteq [t]$, the R -deletion differences $\delta_R^f(\pi) = \delta_R^f(\pi')$ whenever π and π' are two permutations of length $n \geq t + m$ such that $\pi_1 \cdots \pi_{t+m} = \pi'_1 \cdots \pi'_{t+m}$. We denote this constant value $\Delta_R^f(w, n)$ where $w = \pi_1 \cdots \pi_{t+m}$.

In other words, permutation statistic f is ES-compatible with margin m if $\delta_R^f(\pi)$ may be determined from only the length of π and its first $\max R + m$ letters. Note that if f is ES-compatible with margin m , then f is also ES-compatible with margin m' for any $m' \geq m$.

The results from [6] cited above may be rephrased as follows:

Theorem 5. The inversion number is ES-compatible with margin 0.

It follows from the definition of ES-compatible that enumeration schemes are amenable to computing the distribution for any ES-compatible statistic:

Theorem 6. Let f be a ES-compatible permutation statistic with margin m . If R is reversibly deletable for prefix $p \in \mathfrak{S}_k$ with respect to B and $\max R + m \leq k$, then

$$F(\mathfrak{S}_n(B)[p; w], f, q) = q^{\Delta_R^f(w, n)} F(\mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)], f, q), \quad (9)$$

where $\Delta_R^f(w, n)$ has the value $\delta_R^f(\pi)$ for any $\pi \in \mathfrak{S}_n(B)[p; w]$.

Proof. Since $\max R + m \leq |p|$ and f is ES-compatible with margin m , we see from the definition of ES-compatible that $\delta_R^f(\pi)$ is constant for all permutations $\pi \in \mathfrak{S}_n(B)[p; w]$. Therefore $\Delta_R^f(w, n)$ is well-defined. Furthermore, $f(\pi) = \Delta_R^f(w, n) + f(d_R(\pi))$ for any $\pi \in \mathfrak{S}_n(B)[p; w]$.

From the definition of reversibly-deletable, $d_R : \mathfrak{S}_n(B)[p; w] \rightarrow \mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)]$ is a bijection. Shifting focus to the weight enumerators, we see that

$$\begin{aligned}
F(\mathfrak{S}_n(B)[p; w], f, q) &:= \sum_{\pi \in \mathfrak{S}_n(B)[p; w]} q^{f(\pi)} \\
&= \sum_{\pi \in \mathfrak{S}_n(B)[p; w]} q^{\Delta_R^f(w, n)} q^{f(d_R(\pi))} \\
&= q^{\Delta_R^f(w, n)} \sum_{\pi \in d_R(\mathfrak{S}_n(B)[p; w])} q^{f(\pi)} \tag{10} \\
&= q^{\Delta_R^f(w, n)} \sum_{\pi \in \mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)]} q^{f(\pi)} \\
&= q^{\Delta_R^f(w, n)} F(\mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)], f, q)
\end{aligned}$$

Thus we have proven equation (9). \square

By a similar proof we get the following multivariate generalization of Theorem 6:

Theorem 7. *Let f_1, \dots, f_s be ES-compatible permutation statistics, each with margin at most m , and let $\mathbf{f} = \langle f_1, \dots, f_s \rangle$ and $\mathbf{q} = \langle q_1, \dots, q_s \rangle$. If R is reversibly deletable for prefix $p \in \mathfrak{S}_k$ with respect to B and $\max R + m \leq k$, then*

$$F(\mathfrak{S}_n(B)[p; w], \mathbf{f}, \mathbf{q}) = \left(q_1^{\Delta_R^{f_1}(w, n)} \dots q_s^{\Delta_R^{f_s}(w, n)} \right) F(\mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)], \mathbf{f}, \mathbf{q}), \tag{11}$$

where each $\Delta_R^{f_i}(w, n)$ has the value $\delta_R^{f_i}(\pi)$ for any $\pi \in \mathfrak{S}_n(B)[p; w]$.

To more clearly tie Theorems 6 and 7 to enumeration schemes, we introduce the following terminology:

Definition 8. *For nonnegative integer c , an enumeration scheme E has clearance c if for each $(p, G, R) \in E$, either $R = \emptyset$, $\vec{0} \in G$, or $|p| - \max R \geq c$.*

For example, the scheme for 1-2-3-avoiding permutations given in (3) has clearance 0 because of the triple $(12, \{(0, 0, 1)\}, \{2\})$. The clearance of an enumeration scheme describes the largest margin that the scheme could accomodate, as detailed in the following corollaries.

Corollary 9. *If f is a ES-compatible permutation statistic of margin m and E is an enumeration scheme for pattern set B with clearance at least m , then $F(\mathfrak{S}_n(B), f, q)$ may be computed in polynomial time (via enumeration scheme E).*

Corollary 10. *Let f_1, \dots, f_s be ES-compatible permutation statistics, each with margin at most m , and let $\mathbf{f} = \langle f_1, \dots, f_s \rangle$ and $\mathbf{q} = \langle q_1, \dots, q_s \rangle$. Let E be a finite enumeration scheme for pattern set B with clearance at least m . Then $F(\mathfrak{S}_n(B), \mathbf{f}, \mathbf{q})$ may be computed in polynomial time (via enumeration scheme E).*

Clearly corollaries 9 and 10 are impractical if there is no scheme E satisfying the conditions stated. It will be shown in Theorem 15 of Section 4 that a finite scheme of any clearance is sufficient for a polynomial time computation since a scheme can be “deepened” to create a scheme for the same pattern set with any desired clearance.

3.2 Examples of ES-compatible statistics

We now take some time to prove some well-known permutation statistics are indeed ES-compatible.

3.2.1 Copies of consecutive patterns

We first consider statistics based on the number of copies of a given consecutive pattern. Several well-studied statistics can be phrased in terms of the number of copies of certain consecutive patterns. The descent number, $\text{des}(\pi)$, is the number of copies of the consecutive pattern 21. The number of double-descents, i.e., indices i so that $\pi_i > \pi_{i+1} > \pi_{i+2}$, is the number of copies of the consecutive pattern 321. In Subsection 5.2 we discuss the distribution for the number of peaks, i.e., indices i so that $\pi_{i-1} < \pi_i$ and $\pi_i > \pi_{i+1}$, which is the total of the number of copies of 132 and the number of copies of 231.

Theorem 11. *Let $\sigma \in \mathfrak{S}_t$ and $f(\pi)$ be the number of copies of the (consecutive) pattern $(\sigma, [t-1])$ in π . Then f is a ES-compatible statistic with margin $t-1$.*

Note that $f(\pi)$ is defined for any word π , not just permutations. Furthermore, if $\pi \sim \pi'$, then $f(\pi) = f(\pi')$.

Proof. Fix pattern $\sigma \in \mathfrak{S}_t$ and let $k \geq 1$. We will prove that $f(\pi) - f(\pi_1 \cdots \pi_k) = f(d_R(\pi)) - f(d_R(\pi_1 \cdots \pi_k))$ for any $R \subseteq [k-t+1]$. From this it is clear that $\delta_R^f(\pi) = f(\pi_1 \cdots \pi_k) - f(d_R(\pi_1 \cdots \pi_k))$ for any $k \geq \max R + t - 1$, and so f is ES-compatible with margin $t-1$.

Let $f_i(\pi)$ be the number of copies of $(\sigma, [t-1])$ starting at π_i . Since $(\sigma, [t-1])$ is consecutive, we see that

$$f_i(\pi) = \chi[\pi_i \cdots \pi_{i+t-1} \sim \sigma],$$

where $\chi[P]$ is the characteristic function for statement P , i.e., $\chi[P]$ equals 1 if P is true and 0 otherwise. Therefore $f(\pi) = \sum_{i=1}^{n-t+1} f_i(\pi)$ for $\pi \in \mathfrak{S}_n$,

and splitting this sum implies that:

$$\begin{aligned}
f(\pi) &= \sum_{i=1}^{n-t+1} f_i(\pi) \\
&= \sum_{i=1}^{k-t+1} f_i(\pi) + \sum_{i=k-t+2}^{n-t+1} f_i(\pi) \\
&= \sum_{i=1}^{k-t+1} f_i(\pi_1 \cdots \pi_k) + \sum_{i=k-t+2}^{n-t+1} f_i(\pi_{k-t+2} \cdots \pi_n) \\
&= f(\pi_1 \cdots \pi_k) + f(\pi_{k-t+2} \cdots \pi_n)
\end{aligned}$$

Thus $f(\pi) = f(\pi_1 \cdots \pi_k) + f(\pi_{k-t+2} \cdots \pi_n)$. Similarly, let $\pi' = d_R(\pi)$ for $R \subseteq [k-t+1]$, and by the appropriate sum-splitting we see that $f(\pi') = f(\pi'_1 \cdots \pi'_{k-|R|}) + f(\pi'_{k-t+2-|R|} \cdots \pi'_{n-|R|})$. By the definition of d_R , $\pi_{k-t+2} \cdots \pi_n \sim \pi'_{k-t+2-|R|} \cdots \pi'_{n-|R|}$. Hence it follows that

$$\begin{aligned}
f(\pi) - f(\pi_1 \cdots \pi_k) &= f(\pi_{k-t+2} \cdots \pi_n) \\
&= f(\pi'_{k-t+2-|R|} \cdots \pi'_{n-|R|}) \\
&= f(\pi') - f(\pi'_1 \cdots \pi'_{k-|R|})
\end{aligned}$$

Thus we have confirmed $f(\pi) - f(\pi_1 \cdots \pi_k) = f(d_R(\pi)) - f(d_R(\pi_1 \cdots \pi_k))$ and our result follows. \square

Remark. *In terms relevant to enumeration schemes, for any permutation $\pi \in \mathfrak{S}_n(B)[p; w]$ for a prefix of length k ,*

$$\Delta_R^f(w, n) = f(w) - f(d_R(w)) = f(p) - f(d_R(p)), \quad (12)$$

where f counts the number of copies of a consecutive pattern.

3.2.2 Copies of vincular patterns

We next consider the number of copies of a vincular pattern of the form $\sigma_1 \cdots \sigma_{t-1} \sigma_t$. The proof will proceed similarly to that of Theorem 11. Note that the inversion number $\text{inv}(\pi)$ is the number of copies of 2-1, and so is a special case of this result. Subsection 5.3 extends the results in this section to apply the major index statistic.

Theorem 12. *Let $\sigma \in \mathfrak{S}_t$ and let $g(\pi)$ be the number of copies of the pattern $(\sigma, [t-2])$ in π . Then g is a ES-compatible statistic with margin $t-2$.*

Proof. Fix $\sigma \in \mathfrak{S}_t$ and let $k \geq 1$. We will prove that $\delta_R^g(\pi) := g(\pi) - g(d_R(\pi))$ is determined by the length of π together with the prefix $\pi_1 \cdots \pi_k$ for any $R \subseteq [k-t+2]$. From this it follows that g is ES-compatible with margin $t-2$.

For a permutation $\pi \in \mathfrak{S}_n$, define $g_i(\pi)$ to be the number of copies of $(\sigma, [t-2])$ starting at π_i . Clearly $g(\pi) = \sum_{i=1}^n g_i(\pi)$, and if π and π' are order-isomorphic words then $g_i(\pi) = g_i(\pi')$. In particular we see that

$g_i(\pi)$ can be computed in the following way based, on which π_j can be the last letter of a copy of $(\sigma, [t-2])$ starting at π_i :

$$\begin{aligned} g_i(\pi) &:= |\{j : j > i+t-2, \text{ and } \pi_i \cdots \pi_{i+t-2} \pi_j \sim \sigma\}| \\ &= |\{j : j \geq 1 \text{ and } \pi_i \cdots \pi_{i+t-2} \pi_j \sim \sigma\}| - |\{j : j < i \text{ and } \pi_i \cdots \pi_{i+t-2} \pi_j \sim \sigma\}| \end{aligned} \quad (13)$$

We now will show that $g_i(\pi)$ can be determined entirely from $|\pi|$ and $\pi_1 \cdots \pi_{i+t-2}$ by showing each of the addends in (13) requires such limited information.

We consider the first term, $|\{j : j \geq 1 \text{ and } \pi_i \cdots \pi_{i+t-2} \pi_j \sim \sigma\}|$. Given the fixed $\sigma \in \mathfrak{S}_t$, we can define $h_i(w, n)$ for words $w \in [n]^k$ as follows:

$$h_i(w, n) := \begin{cases} \min(w_i, \dots, w_{i+t-2}) - 1 & \text{if } \sigma_t = 1 \\ n - \max(w_i, \dots, w_{i+t-2}) & \text{if } \sigma_t = t \\ w_b - w_a - 1 & \text{if } 1 < \sigma_t < t, \sigma_a = \sigma_t - 1, \sigma_b = \sigma_t + 1. \end{cases} \quad (14)$$

If $\sigma_t = 1$, then $h_i(\pi, n)$ yields the number of letters π_j which are less than each of $\pi_i, \dots, \pi_{i+t-2}$, in which case $\pi_i \cdots \pi_{i+t-2} \pi_j \sim \sigma$. Similarly $\sigma_t = t$, then $h_i(\pi, n)$ yields the number of letters π_j which are greater than each of $\pi_i, \dots, \pi_{i+t-2}$. Last, if a and b are defined by $\sigma_a = \sigma_t - 1$ and $\sigma_b = \sigma_t + 1$, then $h_i(\pi, n)$ yields the number of letters π_j so that $\pi_{i+a-1} < \pi_j < \pi_{i+b-1}$ since every number between the values π_{i+a-1} and π_{i+b-1} appears somewhere in π . In each case, we see that

$$h_i(\pi, n) = |\{j : j \geq 1 \text{ and } \pi_i \cdots \pi_{i+t-2} \pi_j \sim \sigma\}|.$$

Observe that $h_i(\pi, n) = h_i(\pi_1 \cdots \pi_s, n)$ for any $s \geq i+t-2$, and so the first term of the sum in (13) is determined solely by $|\pi|$ and $\pi_i \cdots \pi_{i+t-2}$.

Similarly, the term $|\{j : j < i \text{ and } \pi_i \cdots \pi_{i+t-2} \pi_j \sim \sigma\}|$ can be determined entirely by $\pi_1 \cdots \pi_{i+t-2}$. Define $h'_i(w)$ for word w to be

$$h'_i(w) := \sum_{j=1}^{i-1} \chi[w_i \cdots w_{i+t-2} w_j \sim \sigma],$$

and so it is clear that $h'_i(\pi) = |\{j : j < i \text{ and } \pi_i \cdots \pi_{i+t-2} \pi_j \sim \sigma\}|$. Note that $h'_i(\pi) = h'_i(\pi_1 \cdots \pi_s)$ for any $s \geq i+t-2$.

Combining the above observations, we can decompose the sum $g(\pi) = \sum_{i=1}^n g_i(\pi)$ as follows:

$$\begin{aligned} g(\pi) &= \sum_{i=1}^n g_i(\pi) \\ &= \sum_{i=1}^{k-t+2} g_i(\pi) + \sum_{i=k-t+3}^n g_i(\pi) \\ &= \sum_{i=1}^{k-t+2} (h_i(\pi_1 \cdots \pi_k) - h'_i(\pi_1 \cdots \pi_k)) + \sum_{i=k-t+3}^n g_i(\pi) \end{aligned} \quad (15)$$

Fix $R \subseteq [k-t+2]$ and let $\pi' = d_R(\pi)$. The above observations also apply to π' to imply:

$$g(\pi') = \sum_{i=1}^{k-t-|R|+2} (h_i(\pi'_1 \cdots \pi'_{k-|R|}, n-|R|) - h'_i(\pi'_1 \cdots \pi'_{k-|R|})) + \sum_{i=k-t-|R|+3}^{n-|R|} g_i(\pi') \quad (16)$$

We now focus on the last terms of the equations (15) and (16). By the definition of d_R , $\pi_{k-t+3} \cdots \pi_n \sim \pi'_{k-t+3-|R|} \cdots \pi'_{n-|R|}$. Therefore $g_i(\pi) = g_{i-|R|}(\pi')$ for $i \geq k-t+3$, and so $\sum_{i=k-t+3}^n g_i(\pi) = \sum_{i=k-t-|R|+3}^{n-|R|} g_i(\pi')$.

Thus subtracting equation (16) from (15) we see that:

$$\begin{aligned} \delta_R^g(\pi) &= g(\pi) - g(\pi') \\ &= \sum_{i=1}^{k-t+2} (h_i(\pi_1 \cdots \pi_k, n) - h'_i(\pi_1 \cdots \pi_k)) - \sum_{i=1}^{k-t-|R|+2} (h_i(\pi'_1 \cdots \pi'_{k-|R|}, n-|R|) - h'_i(\pi'_1 \cdots \pi'_{k-|R|})) \end{aligned} \quad (17)$$

Thus we have proven that g is ES-compatible with margin $t-2$. \square

Remark. *In terms relevant to enumeration schemes, for any permutation $\pi \in \mathfrak{S}_n(B)[p; w]$ for a prefix of length k ,*

$$\Delta_R^g(w, n) = \sum_{i=1}^{k-t+2} (h_i(w, n) - h'_i(w)) - \sum_{i=1}^{k-t-|R|+2} (h_i(d_R(w), n-|R|) - h'_i(d_R(w))). \quad (18)$$

As an example of h_i and h'_i in practice, consider $\sigma = 4123$. Here $g(\pi)$ counts the number of copies of the vincular pattern 412-3. Then $a = 1$ and $b = 3$. Let π be any one of the $4!$ permutations of length 9 such that $w = \pi_1 \cdots \pi_5 = 86913$. Since $\pi_3\pi_4\pi_5 = 913 \sim 413 = \sigma_1\sigma_2\sigma_3$, we see that $h_3(\pi, 9) = \pi_3 - \pi_5 - 1 = 5$. Also $h'_3(\pi) = 2$, since both 6 and 8 appear before $\pi_3\pi_4\pi_5$ and have values which lie between $\pi_5 = 3$ and $\pi_3 = 9$. Therefore $g_3(\pi) = h_3(\pi) - h'_3(\pi) = 3$, and indeed there are 3 copies of 412-3 starting at π_3 (specifically, these copies are 9134, 9135, and 9137, since $\{4, 5, 7\} \subseteq \{\pi_6, \dots, \pi_9\}$). In this example $g_1(\pi) = g_2(\pi) = 0$. If $R = \{2, 3\}$, which respects the needed margin of 2 for a length-5 prefix, we get that $\pi' = d_R(\pi)$ has prefix $d_R(86913) = 713$. Deleting the 6 and 9 from π deletes the three copies of 412-3 described above, but creates three new copies since $g_1(\pi') = h_1(713, 7) - h'_1(713) = 3 - 0 = 3$ (specifically, the new copies are witnessed by 7134, 7135, and 7136). Hence in this case $\delta_R^g(\pi) = 0$.

Comparing Theorems 11 and 12, one might guess the trend continues, i.e., that patterns of the form $(\sigma, [t-3])$ for $\sigma \in \mathfrak{S}_t$ are ES-compatible with margin $t-3$. An extension involving partially-ordered generalized patterns, as introduced by Kitaev in [17], provides the proper generalization. For example, a copy of the pattern 124-3'-3'' would be witnessed by a copy of either 125-3-4 or 125-4-3. Such a statistic of the form

$$\sigma_1 \dots \sigma_{t-1} - \sigma'_t - \sigma''_t - \dots - \sigma'''_t$$

can be seen to be ES-compatible with margin $t - 2$ (one less than the length of the consecutive portion).

3.2.3 Right-to-left statistics

A letter w_i of word w is a right-to-left maximum [resp., minimum] if $w_i > w_j$ [resp., $w_i < w_j$] for all $j > i$. Let $\text{rtlmax}(w)$ be the number of right-to-left maxima in w and $\text{rtlmin}(w)$ be the number of right-to-left minima in word w . For example, if $\pi = 28674153$, we see $\text{rtlmax}(\pi) = 4$ (for π_2, π_4, π_7 and π_8) and $\text{rtlmin}(\pi) = 2$ (for π_6 and π_3).

In this subsection we prove the following theorem:

Theorem 13. *The statistics rtlmin and rtlmax are ES-compatible with margin 0.*

It will be useful to have the following characterization of the right-to-left minima and maxima for a permutation, which are based solely on prefixes. The proof follows directly from the definition above and is omitted.

Lemma 14. *Let $\pi = \pi_1 \cdots \pi_n$ be a permutation. Then,*

1. π_i is a right-to-left minimum of π if and only if $\{1, 2, \dots, \pi_i - 1\} \subseteq \{\pi_1, \pi_2, \dots, \pi_{i-1}\}$ (i.e., all numbers less than π_i lie to the left of π_i).
2. π_i is a right-to-left maximum of π if and only if $\{\pi_i + 1, \pi_i + 2, \dots, n\} \subseteq \{\pi_1, \pi_2, \dots, \pi_{i-1}\}$ (i.e., all numbers greater than π_i lie to the left of π_i).

For the remainder of the section, we will restrict ourselves to the proof that rtlmax is ES-compatible. The proof that rtlmin is ES-compatible is analogous.

For integers a and b let $\text{rtlmax}_{[a,b]}(\pi)$ be the number of right-to-left maxima π_i of π such that i is in the closed interval $[a, b]$. For example, $\text{rtlmax}_{[1,5]}(2867153) = 2$ (for π_2 and π_5). Note that while π_5 is a right-to-left maximum of $\pi_1 \cdots \pi_5 = 28671$, it is not counted since this function only counts those letters which are right-to-left maxima in the overall permutation. We may decompose $\text{rtlmax}(\pi)$ for any $\pi \in \mathfrak{S}_n$ and $1 \leq t \leq n$, by $\text{rtlmax}(\pi) = \text{rtlmax}_{[1,t]}(\pi) + \text{rtlmax}_{[t+1,n]}(\pi)$. Thus it follows from the first half of Definition 4 that if $\pi \in \mathfrak{S}_n$, $R \subseteq [t]$ and $\pi' := d_R(\pi) \in \mathfrak{S}_{n-|R|}$,

$$\begin{aligned} \delta_R^{\text{rtlmax}}(\pi) &= \text{rtlmax}(\pi) - \text{rtlmax}(\pi') \\ &= (\text{rtlmax}_{[1,t]}(\pi) + \text{rtlmax}_{[t+1,n]}(\pi)) - (\text{rtlmax}_{[1,t-|R|]}(\pi') + \text{rtlmax}_{[t-|R|+1,n-|R|]}(\pi')) \end{aligned} \tag{19}$$

Rearranging terms leaves us with

$$\delta_R^{\text{rtlmax}}(\pi) = (\text{rtlmax}_{[1,t]}(\pi) - \text{rtlmax}_{[1,t-|R|]}(\pi')) + (\text{rtlmax}_{[t+1,n]}(\pi) - \text{rtlmax}_{[t-|R|+1,n-|R|]}(\pi'))$$

By the original definition of rtlmax , it is clear the $\text{rtlmax}_{[a,n]}(\pi) = \text{rtlmax}(\pi_a \pi_{a+1} \cdots \pi_n)$ for any a . Since $\pi'_{t-|R|+1} \pi'_{t-|R|+2} \cdots \pi'_{n-|R|} \sim \pi_{t+1} \pi_{t+2} \cdots \pi_n$,

it follows that $\text{rtlmax}_{[t+1,n]}(\pi) - \text{rtlmax}_{[t-|R|+1,n-|R|]}(\pi') = 0$. Thus we see that

$$\delta_R^{\text{rtlmax}}(\pi) = \text{rtlmax}_{[1,t]}(\pi) - \text{rtlmax}_{[1,t-|R|]}(\pi') \quad (20)$$

Let w be a word in $[n]^k$ without repeated letters, and define

$$\text{rtlmax}^*(w, n) := |\{w_i : \{w_i + 1, w_i + 2, \dots, n\} \subseteq \{w_1, w_2, \dots, w_{i-1}\}\}|. \quad (21)$$

By Lemma 14 above immediately see for any $\pi \in \mathfrak{S}_n$

$$\text{rtlmax}_{[1,t]}(\pi) = \text{rtlmax}^*(\pi_1\pi_2 \cdots \pi_t, n).$$

Hence equation (20) becomes

$$\delta_R^{\text{rtlmax}}(\pi) = \text{rtlmax}^*(\pi_1\pi_2 \cdots \pi_t, n) - \text{rtlmax}^*(\pi'_1\pi'_2 \cdots \pi'_{t-|R|}, n - |R|). \quad (22)$$

Therefore if $R \subseteq [t]$, then $\delta_R^{\text{rtlmax}}(\pi)$ depends only the values of $\pi_1 \cdots \pi_t$. Thus by Definition 4 we see that rtlmax is ES-compatible with margin 0, and the proof is complete. As mentioned previously, the proof that rtlmin is ES-compatible with 0 proceeds analogously, where $\text{rtlmin}^*(w, n) := |\{w_i : \{1, 2, \dots, w_i - 1\} \subseteq \{w_1, w_2, \dots, w_{i-1}\}\}|$.

In [8] the author generalizes right-to-left maxima with the *right-to-left maximal copy* of a consecutive pattern σ . For a permutation π and consecutive pattern σ , the subfactor $\pi_i\pi_{i+1} \cdots \pi_{i+k-1}$ is a *right-to-left maximal copy* of σ if the following criteria are satisfied:

1. $\pi_i\pi_{i+1} \cdots \pi_{i+k-1} \sim \sigma$, and
2. if $j > i$ and $\pi_j\pi_{j+1} \cdots \pi_{j+k-1} \sim \sigma$ and $\sigma_m = \min(\sigma_1, \dots, \sigma_k)$, then $\pi_{j+m-1} < \pi_{i+m-1}$. In other words, the minimal letter of $\pi_i\pi_{i+1} \cdots \pi_{i+k-1}$ is greater than the minimal letter of any other copy of σ to starting the right of π_i .

For example, the permutation 31856742 has four copies of the consecutive pattern 21 (namely, 31, 85, 74, and 42), but only three of these copies (all but 31) are right-to-left maximal. The classical right-to-left maxima can be viewed as right-to-left maximal copies of the pattern 1. A straightforward generalization of the above argument proves that the statistic counting the number of right-to-left maximal copies of a consecutive pattern σ of length t is ES-compatible with margin $t - 1$.

Before closing this section, it should be noted that the number of left-to-right maxima and left-to-right minima are *not* ES-compatible. For example, if $n = 4$, $w = 12$, and $R = \{1, 2\}$ then $\delta_R^{\text{ltrmin}}(1234) = 0$ while $\delta_R^{\text{ltrmin}}(1243) = -1$. For $n = 4$, $w = 43$, and $R = \{1, 2\}$ we also see $\delta_R^{\text{ltrmax}}(4321) = 0$ while $\delta_R^{\text{ltrmax}}(4312) = -1$.

4 Deepening Schemes

Suppose that pattern set B admits a finite enumeration scheme. The question remains whether one can find a finite enumeration scheme for B with clearance sufficient to accomodate a given ES-compatible statistic with margin m . The algorithms from [9] can be altered to ensure that any constructed scheme has clearance c if such a scheme exists. The existence of such a scheme is guaranteed in the following theorem.

Theorem 15. *Suppose the pattern set B has a finite enumeration scheme E . Then for any $c \geq 0$ there is a finite enumeration scheme E' with clearance c .*

To prove Theorem 15, we will first prove a lemma regarding reversibly deletable sets.

Lemma 16. *Suppose that R is a reversibly deletable set for prefix $p \in \mathfrak{S}_k$ with respect to B . Then R is also reversibly deletable for any permutation $p' \in \mathfrak{S}_{k'}$ such that $p'_1 \cdots p'_k \sim p$, where $k' > k$.*

Proof. If $R \subseteq [k]$ is reversibly deletable for $p \in \mathfrak{S}_k$, then $d_R : \mathfrak{S}_n(B)[p; w] \rightarrow \mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)]$ is a bijection whenever $\mathfrak{S}_n(B)[p; w] \neq \emptyset$. Let $k' > k$ and let $p' \in \mathfrak{S}_{k'}$ such that $\text{red}(p'_1 \cdots p'_k) = p$. Let $w' \in [n]^{k'}$ so that $\mathfrak{S}_n(B)[p'; w'] \neq \emptyset$. We must show that $d_R : \mathfrak{S}_n(B)[p'; w'] \rightarrow \mathfrak{S}_{n-|R|}(B)[d_R(p'); d_R(w')]$ is a bijection.

First note that $\mathfrak{S}_n(B)[p'; w'] = \mathfrak{S}_n(B)[p; w] \cap \mathfrak{S}_n(\emptyset)[p'; w']$, where $w = w'_1 \cdots w'_k$. Since $\mathfrak{S}_n(B)[p'; w'] \neq \emptyset$ and $\mathfrak{S}_n(B)[p'; w'] \subseteq \mathfrak{S}_n(B)[p; w]$, we know that $\mathfrak{S}_n(B)[p; w] \neq \emptyset$. Since R is reversibly deletable for p , the map $d_R : \mathfrak{S}_n(B)[p; w] \rightarrow \mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)]$ is bijective. For the unrestricted case, d_R is a bijection from $\mathfrak{S}_n(\emptyset)[p'; w']$ to $\mathfrak{S}_{n-|R|}(\emptyset)[d_R(p'); d_R(w')]$. Therefore d_R is a bijection from $\mathfrak{S}_n(B)[p'; w']$ to $\mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)] \cap \mathfrak{S}_{n-|R|}(\emptyset)[d_R(p'); d_R(w')]$. Now it remains to show $\mathfrak{S}_{n-|R|}(B)[d_R(p'); d_R(w')] = \mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)] \cap \mathfrak{S}_{n-|R|}(\emptyset)[d_R(p'); d_R(w')]$ to complete our proof. By the action of d_R , the word formed by the first $k - |R|$ letters of $d_R(w')$ is exactly $d_R(w)$, and so $\mathfrak{S}_{n-|R|}(B)[d_R(p'); d_R(w')] \subseteq \mathfrak{S}_{n-|R|}(B)[d_R(p); d_R(w)]$, and the remaining inclusions are obvious from the definitions. Thus R is reversibly deletable for p' . \square

Note that the resulting set R in Lemma 16 is not necessarily a maximal reversibly-deletable set for p . For example, recall from Equation (3) that $\{2\}$ is reversibly deletable for the prefix 12 with respect to $B = \{1-2-3\}$. Lemma 16 implies that $\{2\}$ is reversibly deletable for the prefix 231, although the larger set $\{1, 2\}$ is also reversibly deletable for 231.

We are now ready to prove Theorem 15.

Proof of Theorem 15. Let E be a finite enumeration scheme for B , with depth K . We will construct a scheme E' with clearance $c \geq 1$. If $c = 0$, then E will suffice since any enumeration scheme has clearance 0.

We will construct a (finite) set E' by creating a triple $(p, G(p), R(p)) \in E'$ for each $p \in \bigcup_{k=0}^{K+c} \mathfrak{S}_k$. For $p = \epsilon$, let $G(\epsilon) = \emptyset$ and $R(\epsilon) = \emptyset$, so we see E' satisfies criterion 1 in Definition 3. For $p \neq \epsilon$, let $G(p)$ be a basis of gap vectors for p with respect to B , which may be constructed according to the algorithm described in [9].

We now construct $R(p)$. If $|p| < K+c$, then let $R(p) = \emptyset$. If $|p| = K+c$, then let p' be the longest prefix $p' = \text{red}(p_1 \cdots p_s)$ such that there is a triple (p', G', R') in the original scheme E . Then $|p'| \leq K$, and since no child of p' has a triple in E (by maximality of p') we know R' is nonempty. By Lemma 16, R' is also a reversibly deletable set for p , so we let $R(p) = R'$. Furthermore, $R(p) = R' \subseteq [K]$, so $|p| - \max R(p) \geq c$ and so we see that E' has clearance c .

We now verify that E' satisfies the criteria to be an enumeration scheme for B , as outlined in Definition 3. Each triple $(p, G(p), R(p)) \in E'$ is constructed so that $G(p)$ is a basis of gap vectors for p with respect to B and so that $R(p)$ is a reversibly deletable set for p with respect to B . As previously mentioned, E' satisfies criterion 1 since $(\epsilon, \emptyset, \emptyset) \in E'$. If $R(p) = \emptyset$, then $|p| < K + c$ and so E' contains a triple $(p', G(p'), R(p'))$ for each child p' since E' contains a triple for *every* permutation of length $|p| + 1$. Therefore E' satisfies criterion 2a. If $R(p) \neq \emptyset$, then $|p| = K + c$ and E' contains a triple $(\hat{p}, G(\hat{p}), R(\hat{p}))$ for $\hat{p} = d_{R(p)}(p)$ since E' contains a triple for *every* permutation with length less than $K + c$. Therefore E' satisfies criterion 2b. \square

It perhaps goes without saying that E' from the proof of Theorem 15 is not usually minimal, neither in terms of number of triples nor the encoded recurrence. For example, the proof constructs the following scheme E' with clearance 1 for $B = \{1-2-3\}$ based on the scheme E in (3):

$$\begin{aligned} E' = \{ & (\epsilon, \emptyset, \emptyset), (1, \emptyset, \emptyset), (12, \{\langle 0, 0, 1 \rangle\}, \emptyset), (21, \emptyset, \emptyset), \\ & (123, \{\langle 0, 0, 0, 0 \rangle\}, \{2\}), (132, \{\langle 0, 0, 1, 0 \rangle, \langle 0, 0, 0, 1 \rangle\}, \{2\}), (231, \{\langle 0, 0, 0, 1 \rangle\}, \{2\}), \\ & (213, \{\langle 0, 0, 0, 1 \rangle\}, \{1\}), (312, \{\langle 0, 0, 1, 0 \rangle, \langle 0, 0, 0, 1 \rangle\}, \{1\}), (321, \emptyset, \{1\}) \}. \end{aligned} \tag{23}$$

In practice, therefore, it is better to alter the automated-discovery algorithms from [9] to construct *ab initio* a reversibly deletable set for each prefix p so that only sets $R \subseteq [|p| - c]$ are considered. This change guarantees the clearance criterion holds, and Theorem 15 guarantees that the algorithm will succeed in finding a finite scheme whenever the algorithms would succeed without the clearance conditions. Such an approach yields the following scheme with clearance 1 for $B = \{1-2-3\}$.

$$\begin{aligned} E' = \{ & (\epsilon, \emptyset, \emptyset), (1, \emptyset, \emptyset), (12, \{\langle 0, 0, 1 \rangle\}, \emptyset), (21, \emptyset, \{1\}), \\ & (123, \{\langle 0, 0, 0, 0 \rangle\}, \{1, 2\}), (132, \{\langle 0, 0, 1, 0 \rangle, \langle 0, 0, 0, 1 \rangle\}, \{2\}), (231, \{\langle 0, 0, 0, 1 \rangle\}, \{1, 2\}) \}. \end{aligned} \tag{24}$$

Note in particular that the children of 21 do not need to appear in a scheme with clearance 1, since 21 has a nonempty reversibly deletable set which respects the clearance requirement.

We close this section with a corollary combining Theorem 15 and Corollary 9.

Corollary 17. *If f is a ES-compatible permutation statistic of margin m and E is an enumeration scheme for pattern set B , then $F(\mathfrak{S}_n(B), f, q)$ may be computed in polynomial time.*

5 Applications

We wish to take some time in this section to highlight some of the less obvious applications of the results above. Studying statistics over sets $\mathfrak{S}_n(B)$ is relatively new, so much of what follows only scratches the surface. In what follows we use the common notation that $(\sigma)(\pi)$ denotes the number of copies of vincular pattern σ in permutation π .

5.1 Implementation

The above algorithms have been implemented in a Maple package **Statter**. This package supercedes the package **gVatter** accompanying [9] and can perform the following tasks.

1. Build an enumeration scheme for a given pattern set B with given clearance c . (This also requires search parameters for the maximum size of gap vectors and maximum depth of prefixes.)
2. Read a given enumeration scheme to get the distribution over $\mathfrak{S}_n(B)$ for given n of a statistic based on the number of copies of a consecutive pattern, a pattern of type $\sigma_1 \dots \sigma_{t-1} \sigma_t$, right-to-left maxima, or right-to-left minima.
3. Read a given enumeration scheme to get the distribution of multi-statistics given above.

For example, **Statter** gave the the enumeration scheme for pattern set $B = \{1-2-3\}$ with clearance 1 shown above in (24). From that, **Statter** can quickly give the distribution of the descent statistic over, say, $\mathfrak{S}_{10}(1-2-3)$:

$$F(\mathfrak{S}_{10}(1-2-3), \text{des}, q) = 42q^4 + 1770q^5 + 7515q^6 + 6455q^7 + 1013q^8 + q^9.$$

A fuller investigation of the descent statistic over $\mathfrak{S}_n(1-2-3)$, particularly its connection to Dyck paths, is given in [5].

The package **Statter** is available for download from the author's homepage.

5.2 Peaks and Valleys

A *peak* of a permutation π is a letter π_i such that $\pi_{i-1} < \pi_i > \pi_{i+1}$. Let $\text{peak}(\pi)$ be the number of peaks of π . Therefore $\text{peak}(\pi)$ is the total number of copies of the consecutive patterns 132 and 231 in π . Likewise a *valley* is a letter π_i such that $\pi_{i-1} > \pi_i < \pi_{i+1}$. If $\text{vall}(\pi)$ is the number of valleys of π , then again we see that $\text{vall}(\pi)$ is the total number of copies of the consecutive patterns 213 and 312 in π . Therefore we see that $\text{peak}(\pi)$ and $\text{vall}(\pi)$ are ES-compatible statistics.

While exploring the distributions of the peak and vall statistics over sets $\mathfrak{S}_n(B)$, a few interesting confluences appeared:

Theorem 18. *The following three distributions are equivalent:*

- *the distribution of peaks over 1-2-3-avoiding permutations,*
- *the distribution of valleys over 1-2-3-avoiding permutations, and*
- *the distribution of valleys over 1-3-2-avoiding permutations.*

In the notation above, $F(\mathfrak{S}_n(1-2-3), \text{peak}, q) = F(\mathfrak{S}_n(1-2-3), \text{vall}, q) = F(\mathfrak{S}_n(1-3-2), \text{vall}, q)$.

Remark. *In [15], Dokos et al. started investigating statistic-Wilf-equivalence, where two pattern sets B and B' are f -Wilf-equivalent for permutation statistic f if $F(\mathfrak{S}_n(B), f, q) = F(\mathfrak{S}_n(B'), f, q)$ for all n . This equivalence*

is denoted $B \stackrel{f}{\cong} B'$. In this notation, the second half of Theorem 18 can be written as

$$1-2-3 \stackrel{\text{vall}}{\cong} 1-3-2.$$

Proof. The equivlence $F(\mathfrak{S}_n(1-2-3), \text{peak}, q) = F(\mathfrak{S}_n(1-2-3), \text{vall}, q)$ follows directly from symmetry. Permutation π avoids 1-2-3 if and only if its reverse-complement π^{rc} avoids 1-2-3. Next, the peaks in π correspond to valleys in π^{rc} and vice versa. Thus it follows that $F(\mathfrak{S}_n(1-2-3), \text{peak}, q) = F(\mathfrak{S}_n(1-2-3), \text{vall}, q)$.

We next prove $F(\mathfrak{S}_n(1-2-3), \text{vall}, q) = F(\mathfrak{S}_n(1-3-2), \text{vall}, q)$. We will begin by proving the following claim: *For any $\pi \in \mathfrak{S}_n(1-2-3) \cup \mathfrak{S}_n(1-3-2)$, π_i is a valley of π if and only if $i > 1$, π_i is a left-to-right minimum of π and π_{i+1} is not an left-to-right minimum of π .* Suppose that π_i is a valley but that there is some π_j such that $\pi_j < \pi_i$ and $j < i$. Then we see $\pi_j \pi_{j-1} \pi_i \sim 132$ and $\pi_j \pi_i \pi_{i+1} \sim 123$ and so π would contain 1-3-2 and 1-2-3. Thus if π avoids 1-3-2 or 1-2-3 then π_i must be a left-to-right minimum. Since $\pi_{i+1} > \pi_i$ we see that π_{i+1} is not a left-to-right minimum. Conversely, suppose that π_i is a left-to-right minimum of π for $i > 1$ but π_{i+1} is not. Then $\pi_{i+1} > \pi_i$, since otherwise π_{i+1} would be a left-to-right minimum, and also $\pi_{i-1} > \pi_i$, since otherwise π_i would not be an left-to-right minimum. Therefore π_i must be a valley of π . Hence the claim is proven.

The Simion-Schmidt bijection in [23] provides a map $\mathfrak{S}_n(1-3-2) \rightarrow \mathfrak{S}_n(1-2-3)$ which preserves the left-to-right minima. Therefore by the claim above, the same bijection also preserves the valleys of the permutations. Thus we see that $F(\mathfrak{S}_n(1-2-3), \text{vall}, q) = F(\mathfrak{S}_n(1-3-2), \text{vall}, q)$. \square

Remark. *The proof above implies that vall is equally distributed over $\mathfrak{S}_n(1-2-3)$ and $\mathfrak{S}_n(1-3-2)$ even when restricting further to those permutations with a given set of left-to-right minima. Letting $\text{LRmax}(\pi)$ be the set of indices which are left-to-right minima for π , then for any $S \subseteq [n]$.*

$$\sum_{\substack{\pi \in \mathfrak{S}_n(1-2-3) \\ \text{LRmax}(\pi)=S}} q^{\text{vall}(\pi)} = \sum_{\substack{\pi \in \mathfrak{S}_n(1-3-2) \\ \text{LRmax}(\pi)=S}} q^{\text{vall}(\pi)} \quad (25)$$

For completeness we will comment that $F(\mathfrak{S}_n(1-3-2), \text{peak}, q)$ appears in OEIS as A091894, suggesting the following correspondence. A Dyck path of semilength n is a lattice path from $(0,0)$ to $(n,0)$ composed of steps $U = (1,1)$ and $D = (1,-1)$ which never goes below the x -axis. We will write Dyck paths as words $U^{a_1} D^{b_1} U^{a_2} D^{b_2} \dots U^{a_k} D^{b_k}$.

Theorem 19. *The number of permutations in $\mathfrak{S}_n(1-3-2)$ with k peaks equals the number of Dyck paths of semilength n with k occurrences of the subfactor DDU .*

Proof. Krattenthaler provides a bijection, Φ , in [18] from $\mathfrak{S}_n(1-3-2)$ to the set of Dyck paths of semilength n . In that bijection, a permutation with k valleys maps to a Dyck path with k subfactors DDU . The remainder of this proof outlines this bijection.

Suppose that $\pi \in \mathfrak{S}_n(1-3-2)$ with $\text{ltrmin}(\pi) = k$. Then let j_1, j_2, \dots, j_k be the indices of the left-to-right minima and let $m_i = \pi_{j_i}$ be the values

of the left-to-right minima for $1 \leq i \leq k$. Further, let $m_0 = n + 1$ and $j_{k+1} = n + 1$. Then $\Phi(\pi)$ is the Dyck path $U^{a_1} D^{b_1} U^{a_2} D^{b_2} \dots U^{a_k} D^{b_k}$ where $a_i = m_{i-1} - m_i$ and $b_i = j_{i+1} - j_i$. For Φ^{-1} , observe that the left-to-right minima of $\Phi^{-1}(U^{a_1} D^{b_1} U^{a_2} D^{b_2} \dots U^{a_k} D^{b_k})$ occur at indices $1, 1 + b_1, 1 + b_1 + b_2, \dots, 1 + b_1 + b_2 + \dots + b_{k-1}$ and have values $n + 1 - m_1, n + 1 - (m_1 + m_2), \dots, n + 1 - (m_1 + m_2 + \dots + m_k)$. It is well-known that a 1-3-2-avoiding permutation is uniquely determined by the indices and values of its left-to-right minima, and thus Φ is bijective.

Observe that if $\text{ltrmin}(\pi) = k$, then $\Phi(\pi) = U^{a_1} D^{b_1} U^{a_2} D^{b_2} \dots U^{a_k} D^{b_k}$ for $a_i > 0$ and $b_i > 0$. Therefore each left-to-right minima beyond π_1 corresponds to a DU subfactor in $\Phi(\pi)$. Furthermore if adjacent letters π_i and π_{i+1} are both left-to-right minima, then the corresponding string of D 's in the image is only a single D . Therefore DDU subfactors correspond to non-adjacent left-to-right minima, which by the claim from Theorem 18 correspond to valleys in the permutation. \square

The distributions from Theorems 18 and 19 are given in Tables 1 and 2. This data was generated by **Statter**.

n	$F(\mathfrak{S}_n(1-2-3), \text{peak}, q)$
1	1
2	2
3	$3 + 2q$
4	$4 + 10q$
5	$5 + 32q + 5q^2$
6	$6 + 84q + 42q^2$
7	$7 + 198q + 210q^2 + 14q^3$
8	$8 + 438q + 816q^2 + 168q^3$
9	$9 + 932q + 2727q^2 + 1152q^3 + 42q^4$
10	$10 + 1936q + 8250q^2 + 5940q^3 + 660q^4$

Table 1: The distributions from Theorem 18 for $1 \leq n \leq 10$.

n	$F(\mathfrak{S}_n(1-3-2), \text{peak}, q)$
1	1
2	2
3	$4 + q$
4	$8 + 6q$
5	$16 + 24q + 2q^2$
6	$32 + 80q + 20q^2$
7	$64 + 240q + 120q^2 + 5q^3$
8	$128 + 672q + 560q^2 + 70q^3$
9	$256 + 1792q + 2240q^2 + 560q^3 + 14q^4$
10	$512 + 4608q + 8064q^2 + 3360q^3 + 252q^4$

Table 2: The distributions of peaks over 1-3-2-avoiding permutations. Corresponds to OEIS sequence A091894 [19].

A complete classification of the classical patterns according to peak-Wilf-equivalence is forthcoming in [10].

5.3 The major index

The major index is defined by $\text{maj}(\pi) := \sum_{i:\pi_i > \pi_{i+1}} i$. As shown in [2], the major index can be decomposed into the sum of four vincular pattern functions:

$$\text{maj}(\pi) = (3-21)(\pi) + (2-31)(\pi) + (1-32)(\pi) + (21)(\pi). \quad (26)$$

where $(\sigma)(\pi)$ is the number of copies of σ in permutation π . While these patterns do not conform to the structure of those in Theorem 12, their reverses do. Thus we define

$$\text{rmaj}(\pi) := \text{maj}(\pi^r) = (12-3)(\pi) + (13-2)(\pi) + (23-1)(\pi) + (12)(\pi).$$

Therefore Corollary 10, Theorem 12 imply the following special case:

Corollary 20. *If B is a set of patterns such that $B^r := \{\tau^r : \tau \in B\}$ admits a finite enumeration scheme, then $F(\mathfrak{S}_n(B), \text{maj}, q)$ can be computed via enumeration scheme.*

Distributions of the major index over avoidance sets $\mathfrak{S}_n(\tau)$ for classical patterns $\tau \in \mathfrak{S}_3$ are studied by Dokos et al. in [15].

To illustrate the method with a new example, consider the set of classical patterns

$$B_k = \{2-1-3, 1-2-\dots-(k-1)-k\}$$

for any $k \geq 2$. A discussion of $B_3^c = \{2-3-1, 3-2-1\}$ appears in [15], but we will consider the more general family. It was shown in [25] that $B_k^r = \{3-1-2, k-(k-1)-\dots-2-1\}$ admits a finite enumeration scheme of depth 2:

$$\{(\epsilon, \emptyset, \emptyset), (1, \emptyset, \emptyset), (12, \emptyset, \{1\}), (21, \{\langle 0, 1, 0 \rangle, \langle k-2, 0, 0 \rangle\}, \{1\})\} \quad (27)$$

The scheme in (27) derives from general-purpose algorithms of [25], and so is not necessarily optimal for a special case. There are a few missed gap vector criteria which will simplify the resulting recurrence. Observe that if π is a permutation with $\pi_1 \geq k$, then $2, 3, \dots, k-1$ lie among the remaining letters. Either these letters appear in decreasing order, in which case π contains a $k-(k-1)-\dots-2-1$, or at least two of the letters appear in increasing order, in which case π contains $3-1-2$. Therefore $\langle k-1, 0 \rangle$ is a gap vector for prefix pattern 1, and in turn this implies that $\{\langle i, j, 0 \rangle : i + j = k-1\}$ are gap vectors for prefix 12. Thus we arrive at the following enumeration scheme for B_k^r :

$$\begin{aligned} &\{(\epsilon, \emptyset, \emptyset), (1, \{\langle k-1, 0 \rangle\}, \emptyset), (12, \{\langle k-1, 0, 0 \rangle, \langle k-2, 1, 0 \rangle, \dots, \langle 0, k-1, 0 \rangle\}, \{1\}), \\ &\quad (21, \{\langle 0, 1, 0 \rangle, \langle k-2, 0, 0 \rangle\}, \{1\})\} \end{aligned} \quad (28)$$

We will consider the following analogue of the classic Euler-Mahonian distribution, restricted to the B_k -avoiding permutations:

$$\begin{aligned} G_n(p; w) &:= F(\mathfrak{S}_n(B_k)[p; w], \langle \text{maj}, \text{des} \rangle, \langle q, t \rangle) \\ &= F(\mathfrak{S}_n(B_k^r)[p; w], \langle \text{rmaj}, (12) \rangle, \langle q, t \rangle) \end{aligned}$$

This scheme in (28) translates into the following recurrences for $n \geq 2$:

$$\begin{aligned}
\sum_{\pi \in \mathfrak{S}_n(B_k)} q^{\text{maj}(\pi)} t^{\text{des}(\pi)} &= \sum_{\pi \in \mathfrak{S}_n(B_k^r)} q^{\text{rmaj}(\pi)} t^{(12)(\pi)} \\
&= G_n(\epsilon; \epsilon) = \sum_{a=1}^n G_n(1, a) \\
&= \sum_{a=1}^{k-1} G_n(1, a) \text{ (by gap vector criteria)} \\
G_n(1, a) &= \sum_{b=1}^{a-1} G_n(21; ab) + \sum_{b=a+1}^n G_n(12; ab) \\
&= G_n(21; a(a-1)) + \sum_{b=a+1}^k G_n(12; ab) \text{ (by gap vector criteria)} \\
G_n(21; a(a-1)) &= t^0 q^0 G_{n-1}(1; a-1) \\
G_n(12; ab) &= t^1 q^{n-1} G_{n-1}(1; b-1)
\end{aligned} \tag{29}$$

The reader is left to verify the following values:

- $\Delta_{\{1\}}^{(12)}(a(a-1), n) = 0$
- $\Delta_{\{1\}}^{\text{rmaj}}(a(a-1), n) = 0$
- $\Delta_{\{1\}}^{(12)}(ab, n) = 1$ for $ab \sim 12$
- $\Delta_{\{1\}}^{\text{rmaj}}(ab, n) = n-1$ for $ab \sim 12$

Combining the recurrence relations above yields the following recurrence for $G_n(1; a)$, the distribution of (maj, des) over B_k -avoiding permutations:

$$G_n(1; a) = \begin{cases} t q^{n-1} \sum_{b=1}^{k-1} G_{n-1}(1; b), & a = 1 \\ G_{n-1}(1; a-1) + t q^{n-1} \sum_{b=a}^{k-1} G_{n-1}(1; b), & 2 \leq a \leq k-1 \\ 0, & a \geq k. \end{cases} \tag{30}$$

6 Conclusion and Future Work

The techniques above face the same limitations as enumeration schemes. In short, the recurrences produced are often complicated and do not translate nicely into generating functions. The methods discussed in Chapter 5 of [7] make some progress toward converting schemes to generating functions, but cannot account for the full range of recurrences that schemes can produce.

Further, not all sets of vincular patterns B admit a finite enumeration scheme, and there is no full characterization predicting whether a given B will admit a finite scheme. Data on how many sets B do admit a small scheme are available in [9].

We note that it should be possible to adapt the insertion encodings from [1, 26] toward the purpose of computing $F(\mathfrak{S}_n(B), f, q)$ for permutation statistics f based on counting copies of consecutive patterns. The insertion encoding offers two advantages over enumeration schemes: (1) the recurrences developed lead directly to generating functions, and (2) there are more [sets of] classical patterns which admit regular insertion encodings than finite enumeration schemes. The current state of insertion encodings, however, cannot handle vincular patterns, however. Such tools could be very helpful in classification of patterns under statistic-Wilf-equivalence for various statistics.

References

- [1] Michael H. Albert, Steve Linton, and Nik Ruškuc. The insertion encoding of permutations. *Electron. J. Combin.*, 12:Research Paper 47, 31 pp. (electronic), 2005.
- [2] Eric Babson and Einar Steingrímsson. Generalized permutation patterns and a classification of the Mahonian statistics. *Sém. Lothar. Combin.*, 44:Art. B44b, 18 pp. (electronic), 2000.
- [3] E. Barucci, A. Del Lungo, E. Pergola, and R. Pinzani. Some permutations with forbidden subsequences and their inversion number. *Discrete Math.*, 234(1-3):1–15, 2001.
- [4] Marilena Barnabei, Flavio Bonetti, and Matteo Silimbani. The descent statistic on 123-avoiding permutations. *Sém. Lothar. Combin.*, 63:Art. B63a, 8, 2010.
- [5] Marilena Barnabei, Flavio Bonetti, and Matteo Silimbani. The joint distribution of consecutive patterns and descents in permutations avoiding 3-1-2. *European J. Combin.*, 31(5):1360–1371, 2010.
- [6] Andrew Baxter. Refining enumeration schemes to count according to the inversion number. *Pure Mathematics and Applications*, 21(2):137–160, 2010.
- [7] Andrew Baxter. *Algorithms for Permutation Statistics*. PhD thesis, Rutgers, The State University of New Jersey, 2011.
- [8] Andrew M. Baxter. Shape-Wilf-equivalences for vincular patterns. *Adv. in Appl. Math.*, 50(5):723–736, 2013.
- [9] Andrew M. Baxter and Lara K. Pudwell. Enumeration schemes for vincular patterns. *Discrete Math.*, 312(10):1699–1712, 2012.
- [10] Andrew M. Baxter and Charles Walker. Peak-wilf-equivalence. 2014. In preparation.
- [11] Miklós Bóna. The absence of a pattern and the occurrences of another. *Discrete Math. Theor. Comput. Sci.*, 12(2):89–102, 2010.

- [12] Miklós Bóna. Surprising symmetries in objects counted by catalan numbers. *Electron. J. Combin.*, 19(1):P62, 2012.
- [13] A. Burstein and S. Elizalde. Total occurrence statistics on restricted permutations. *ArXiv e-prints*, May 2013.
- [14] Anders Claesson. Generalized pattern avoidance. *European J. Combin.*, 22(7):961–971, 2001.
- [15] Theodore Dokos, Tim Dwyer, Bryan P. Johnson, Bruce E. Sagan, and Kimberly Selsor. Permutation patterns and statistics. *Discrete Math.*, 312(18):2760–2775, 2012.
- [16] Cheyne Homberger. Expected patterns in permutation classes. *Electron. J. Combin.*, 19(3):Paper 43, 12, 2012.
- [17] Sergey Kitaev. Partially ordered generalized patterns. *Discrete Math.*, 298(1-3):212–229, 2005.
- [18] C. Krattenthaler. Permutations with restricted patterns and Dyck paths. *Adv. in Appl. Math.*, 27(2-3):510–530, 2001. Special issue in honor of Dominique Foata’s 65th birthday (Philadelphia, PA, 2000).
- [19] OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences. <http://oeis.org>, 2011.
- [20] Lara Pudwell. Enumeration schemes for words avoiding patterns with repeated letters. *Integers*, 8:A40, 19, 2008.
- [21] Lara Pudwell. Enumeration schemes for permutations avoiding barred patterns. *Electron. J. Combin.*, 17(1):Research Paper 29, 27 pp., 2010.
- [22] Lara Pudwell. Enumeration schemes for words avoiding permutations. In *Permutation patterns*, volume 376 of *London Math. Soc. Lecture Note Ser.*, pages 193–211. Cambridge Univ. Press, Cambridge, 2010.
- [23] Rodica Simion and Frank W. Schmidt. Restricted permutations. *European J. Combin.*, 6(4):383–406, 1985.
- [24] Einar Steingrímsson. Generalized permutation patterns — a short survey. In Steve Linton, Nik Ruškuc, and Vincent Vatter, editors, *Permutation Patterns, St Andrews 2007*, volume 376 of *London Mathematical Society Lecture Note Series*, pages 193–211. Cambridge University Press, 2010.
- [25] Vincent Vatter. Enumeration schemes for restricted permutations. *Combin. Probab. Comput.*, 17(1):137–159, 2008.
- [26] Vincent Vatter. Finding regular insertion encodings for permutation classes. *J. Symbolic Comput.*, 47(3):259–265, 2012.
- [27] Doron Zeilberger. Enumeration schemes and, more importantly, their automatic generation. *Ann. Comb.*, 2(2):185–195, 1998.
- [28] Doron Zeilberger. On Vince Vatter’s brilliant extension of Doron Zeilberger’s enumeration schemes for Herb Wilf’s classes. Published in *The Personal Journal of Ekhad and Zeilberger*, 2006.