

CONSECUTIVE PRIMES AND LEGENDRE SYMBOLS

HAO PAN AND ZHI-WEI SUN

Department of Mathematics, Nanjing University
 Nanjing 210093, People's Republic of China
 haopan1979@gmail.com zwsun@nju.edu.cn

ABSTRACT. Let m be any positive integer and let $\delta_1, \delta_2 \in \{1, -1\}$. We show that for some constant $C_m > 0$ there are infinitely many integers $n > 1$ with $p_{n+m} - p_n \leq C_m$ such that

$$\left(\frac{p_{n+i}}{p_{n+j}}\right) = \delta_1 \quad \text{and} \quad \left(\frac{p_{n+j}}{p_{n+i}}\right) = \delta_2$$

for all $0 \leq i < j \leq m$, where p_k denotes the k -th prime, and $\left(\frac{\cdot}{p}\right)$ denotes the Legendre symbol for any odd prime p . We also prove that under the Generalized Riemann Hypothesis there are infinitely many positive integers n such that p_{n+i} is a primitive root modulo p_{n+j} for any distinct i and j among $0, 1, \dots, m$.

1. INTRODUCTION

For $n \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$ let p_n denote the n -th prime. The famous twin prime conjecture asserts that $p_{n+1} - p_n = 2$ for infinitely many $n \in \mathbb{Z}^+$. Although this remains open, recently Y. Zhang [Z] was able to prove that

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) \leq 7 \times 10^7.$$

The upper bound 7×10^7 was later reduced to 4680 by the Polymath team [Po] led by T. Tao, and 600 by J. Maynard [M], and 270 again by the Polymath team [Po]. Moreover, J. Maynard [M], as well as T. Tao, established the following deep result.

Theorem 1.1 (Maynard-Tao). *For any positive integer m , we have*

$$\liminf_{n \rightarrow \infty} (p_{n+m} - p_n) \leq C m^3 e^{4m},$$

where $C > 0$ is an absolutely constant.

Earlier than this work, in 2000 D.K.L. Shiu [S] proved the following nice theorem.

2010 *Mathematics Subject Classification.* Primary 11A41; Secondary 11A07, 11A15, 11N99.

Keywords. Consecutive primes, Legendre symbols, quadratic residues.

The second author is supported by the National Natural Science Foundation (Grant No. 11171140) of China.

Theorem 1.2 (Shiu). *Let $a \in \mathbb{Z}$ and $q \in \mathbb{Z}^+$ be relatively prime. Then, for any $m \in \mathbb{Z}^+$ there is a positive integer n such that*

$$p_n \equiv p_{n+1} \equiv \cdots \equiv p_{n+m} \equiv a \pmod{q}.$$

This was recently re-deduced in [BFTB] via the Maynard-Tao method.

In this paper we mainly establish the following new result on consecutive primes and Legendre symbols.

Theorem 1.3. *Let m be any positive integer and let $\delta_1, \delta_2 \in \{1, -1\}$. For some constant $C_m > 0$ depending only on m , there are infinitely many integers $n > 1$ with $p_{n+m} - p_n \leq C_m$ such that for any $0 \leq i < j \leq m$ we have*

$$\left(\frac{p_{n+i}}{p_{n+j}}\right) = \delta_1 \quad \text{and} \quad \left(\frac{p_{n+j}}{p_{n+i}}\right) = \delta_2. \quad (1.1)$$

Remark 1.1. (a) Instead of (1.1) in Theorem 1.3, actually we may require both (1.1) and the following property:

$$p_{ij} \parallel (p_{n+i} - p_{n+j}) \quad \text{for some prime } p_{ij} > 2m + 1. \quad (1.2)$$

(As usual, for a prime p and an integer a , by $p \parallel a$ we mean $p \mid a$ but $p^2 \nmid a$.)

(b) We conjecture the following extension of Theorem 1.3: For any $m \in \mathbb{Z}^+$, $\delta \in \{1, -1\}$ and $\delta_{ij} \in \{1, -1\}$ with $0 \leq i < j \leq m$, there are infinitely many integers $n > 1$ such that

$$\left(\frac{p_{n+i}}{p_{n+j}}\right) = \delta_{ij} = \delta \left(\frac{p_{n+j}}{p_{n+i}}\right)$$

for all $0 \leq i < j \leq m$.

Example 1.1. The smallest integer $n > 1$ with

$$\left(\frac{p_{n+i}}{p_{n+j}}\right) = 1 \quad \text{for all } i, j = 0, \dots, 6 \text{ with } i \neq j$$

is 176833, and a list of the first 200 such values of n is available from [Su2]. The 7 consecutive primes $p_{176833}, p_{176834}, \dots, p_{178639}$ have concrete values

$$2434589, 2434609, 2434613, 2434657, 2434669, 2434673, 2434681$$

respectively.

Example 1.2. The smallest integer $n > 1$ with

$$\left(\frac{p_{n+i}}{p_{n+j}}\right) = -1 \quad \text{for all } i, j = 0, \dots, 5 \text{ with } i \neq j$$

is 2066981, and the 6 consecutive primes $p_{2066981}, p_{2066982}, \dots, p_{2066986}$ have concrete values

$$33611561, 33611573, 33611603, 33611621, 33611629, 33611653$$

respectively.

Example 1.3. The smallest integer $n > 1$ with

$$-\left(\frac{p_{n+i}}{p_{n+j}}\right) = 1 = \left(\frac{p_{n+j}}{p_{n+i}}\right) \quad \text{for all } 0 \leq i < j \leq 6$$

is 7455790, and the 7 consecutive primes $p_{7455790}, p_{7455791}, \dots, p_{7455796}$ have concrete values

$$131449631, 131449639, 131449679, 131449691, 131449727, 131449739, 131449751$$

respectively.

Example 1.4. The smallest integer $n > 1$ with

$$\left(\frac{p_{n+i}}{p_{n+j}}\right) = 1 = -\left(\frac{p_{n+j}}{p_{n+i}}\right) \quad \text{for all } 0 \leq i < j \leq 5$$

is 59753753, and the 6 consecutive primes $p_{59753753}, p_{59753754}, \dots, p_{59753758}$ have concrete values

$$1185350899, 1185350939, 1185350983, 1185351031, 1185351059, 1185351091$$

respectively.

Actually Theorem 1.3 is motivated by the second author's following conjecture.

Conjecture 1.1 (Sun [Su1, Su2]). *For any positive integer m , there are infinitely many $n \in \mathbb{Z}^+$ such that for any distinct i and j among $0, 1, \dots, m$ the prime p_{n+i} is a primitive root modulo p_{n+j} .*

Example 1.5. The least $n \in \mathbb{Z}^+$ with p_{n+i} a primitive root modulo p_{n+j} for any distinct i and j among $0, 1, 2, 3$ is 8560, and a list of the first 50 such values of n is available from [Su2, A243839]. Note that

$$p_{8560} = 88259, p_{8561} = 88261 \text{ and } p_{8562} = 88289.$$

Our second result is the following theorem.

Theorem 1.4. *Conjecture 1.1 holds under the Generalized Riemann Hypothesis.*

We will prove Theorem 1.3 in the next section with the help of the Maynard-Tao work, and show Theorem 1.4 in Section 3 by combining our method with a recent result of P. Pollack [P] motivated by the Maynard-Tao work on bounded gaps of primes and Artin's conjecture on primitive roots modulo primes.

Throughout this paper, p always represents a prime. For two integers a and b , their greatest common divisor is denoted by $\gcd(a, b)$.

2. PROOF OF THEOREM 1.3

Let h_1, h_2, \dots, h_k be distinct positive integers. If $\bigcup_{j=1}^k h_i \pmod{p} \neq \mathbb{Z}$ for any prime p (where $a \pmod{p}$ denotes the residue class $a + p\mathbb{Z}$), then we call $\{h_i : i = 1, \dots, k\}$ an admissible set. Hardy and Littlewood conjectured that if $\mathcal{H} = \{h_i : i = 1, \dots, k\}$ is admissible then there are infinitely many $n \in \mathbb{Z}^+$ such that $n + h_1, n + h_2, \dots, n + h_k$ are all prime. We need the following result in this direction.

Lemma 2.1 (Maynard-Tao). *Let m be any positive integer. Then there is an integer $k > m$ depending only on m such that if $\mathcal{H} = \{h_i : i = 1, \dots, k\}$ is an admissible set of cardinality k and $W = q_0 \prod_{p \leq w} p$ (with $q_0 \in \mathbb{Z}^+$) is relatively prime to $\prod_{i=1}^k h_i$ with $w = \log \log \log x$ large enough, then for some integer $n \in [x, 2x]$ with $W \mid n$ there are more than m primes among $n + h_1, n + h_2, \dots, n + h_k$.*

Lemma 2.2. *Let $k > 1$ be an integer. Then there is an admissible set $\mathcal{H} = \{h_1, \dots, h_k\}$ with $h_1 = 0 < h_2 < \dots < h_k$ which has the following properties:*

- (i) *All those h_1, h_2, \dots, h_k are multiples of $K = 4 \prod_{p < 2k} p$.*
- (ii) *Each $h_i - h_j$ with $1 \leq i < j \leq k$ has a prime divisor $p > 2k$ with $h_i \not\equiv h_j \pmod{p^2}$.*
- (iii) *If $1 \leq i < j \leq k$, $1 \leq s < t \leq k$ and $\{i, j\} \neq \{s, t\}$, then no prime $p > 2k$ divides both $h_i - h_j$ and $h_s - h_t$.*

Proof. Set $h_1 = 0$ and let $1 \leq r < k$. Suppose that we have found nonnegative integers $h_1 < \dots < h_r$ divisible by K such that each $h_i - h_j$ with $1 \leq i < j \leq r$ has a prime divisor $p > 2k$ with $h_i \not\equiv h_j \pmod{p^2}$, and that no prime $p > 2k$ divides both $h_i - h_j$ and $h_s - h_t$ if $1 \leq i < j \leq r$, $1 \leq s < t \leq r$ and $\{i, j\} \neq \{s, t\}$. Let

$$X_r = \{p > 2k : p \mid h_s - h_t \text{ for some } 1 \leq s < t \leq r\}.$$

As K is relatively prime to $\prod_{p \in X_r} p$, for each $i = 1, \dots, r$ there is an integer b_i with $Kb_i \equiv h_i \pmod{\prod_{p \in X_r} p}$. For each $p \in X_r$, as $r < k < p$ there is an integer $a_p \not\equiv b_i \pmod{p}$ for all $i = 1, \dots, r$. Choose distinct primes q_1, \dots, q_r which are greater than $2k$ but not in the set X_r . For any $i = 1, \dots, r$, there is an integer c_i with $Kc_i \equiv h_i \pmod{q_i^2}$ since K is relatively prime to q_i^2 . By the Chinese Remainder Theorem, there is an integer $b > h_r/K$ such that $b \equiv a_p \pmod{p}$ for all $p \in X_r$, and $b \equiv c_i + q_i \pmod{q_i^2}$ for all $i = 1, \dots, r$.

Set $h_{r+1} = Kb > h_r$. If $1 \leq s \leq r$, then

$$h_{r+1} - h_s \equiv Kb - Kc_s = K(b - c_s) \equiv Kq_s \pmod{q_s^2},$$

hence $q_s > 2k$ is a prime divisor of $h_{r+1} - h_s$ but $h_{r+1} \not\equiv h_s \pmod{q_s^2}$.

For $s, t \in \{1, \dots, r\}$ with $s \neq t$, clearly

$$\gcd(h_{r+1} - h_s, h_{r+1} - h_t) = \gcd(h_{r+1} - h_s, h_s - h_t).$$

Let $1 \leq i < j \leq r$ and $1 \leq s \leq r$. If a prime $p > 2k$ divides $h_i - h_j$, then $p \in X_r$ and hence

$$h_{r+1} - h_s \equiv Ka_p - Kb_s = K(a_p - b_s) \not\equiv 0 \pmod{p}.$$

So $\gcd(h_{r+1} - h_s, h_i - h_j)$ has no prime divisor greater than $2k$.

In view of the above, we have constructed nonnegative integers $h_1 < h_2 < \dots < h_k$ satisfying (i)-(iii) in Lemma 2.2. Note that $\bigcup_{i=1}^k h_i \pmod{p} \neq \mathbb{Z}$ if $p > k$. For each $p \leq k$, clearly $h_i \equiv 0 \not\equiv 1 \pmod{p}$ for any $i = 1, \dots, k$. Therefore the set $\mathcal{H} = \{h_1, h_2, \dots, h_k\}$ is admissible. This concludes the proof. \square

Proof of Theorem 1.3. By Lemma 2.1, there is an integer $k = k_m > m$ depending on m such that for any admissible set $\mathcal{H} = \{h_1, \dots, h_k\}$ of cardinality k if x is sufficiently large and $\prod_{i=1}^k h_i$ is relatively prime to $W = 4 \prod_{p \leq w} p$ then for some integer $n \in [x/W, 2x/W]$ there are more than m primes among $Wn + h_1, Wn + h_2, \dots, Wn + h_k$, where $w = \log \log \log x$.

Let $\mathcal{H} = \{h_1, \dots, h_k\}$ with $h_1 = 0 < h_2 < \dots < h_k$ be an admissible set satisfying the conditions (i)-(iii) in Lemma 2.2. Clearly $K = 4 \prod_{p \leq 2k} p \equiv 0 \pmod{8}$. Let x be sufficiently large with the interval $(h_k, w]$ containing more than $h_k - k$ primes. Note that $8 \mid W$ since $w \geq 2$.

Let $\delta := \delta_1 \delta_2$. For any integer $b \equiv \delta \pmod{K}$ and each prime $p < 2k$, clearly $b + h_i \equiv \delta + 0 \pmod{p}$ and hence $\gcd(b + h_i, p) = 1$ for all $i = 1, \dots, k$.

For any $1 \leq i < j \leq k$, the number $h_i - h_j$ has a prime divisor $p_{ij} > 2k$ with $h_i \not\equiv h_j \pmod{p_{ij}^2}$. Suppose that $p > 2k$ is a prime dividing $\prod_{1 \leq i < j \leq k} (h_i - h_j)$, then there is a unique pair $\{i, j\}$ with $1 \leq i < j \leq k$ such that $h_i \equiv h_j \pmod{p}$. Note that $p \leq h_k$. All the $k - 2 < (p - 3)/2$ numbers $h_i - h_s$ with $1 \leq s \leq k$ and $s \neq i, j$ are relatively prime to p , so there is an integer $r_p \not\equiv h_i - h_s \pmod{p}$ for all $s = 1, \dots, k$ such that

$$\left(\frac{r_p \delta}{p} \right) = \begin{cases} \delta_2 & \text{if } p = p_{ij}, \\ 1 & \text{otherwise.} \end{cases}$$

So, for any integer $b \equiv r_p - h_i \pmod{p}$, we have $b + h_s \not\equiv 0 \pmod{p}$ for all $s = 1, \dots, k$.

Assume that $S = \{h_1, h_1 + 1, \dots, h_k\} \setminus \mathcal{H}$ is a set $\{a_i : i = 1, \dots, t\}$ of cardinality $t > 0$. Clearly $t \leq h_k - k + 1$ and hence we may choose t distinct primes $q_1, \dots, q_t \in (h_k, w]$. If $b \equiv -a_i \pmod{q_i}$, then $b + h_s \equiv h_s - a_i \not\equiv 0 \pmod{q_i}$ for all $s = 1, \dots, k$ since $0 < |h_s - a_i| < h_k < q_i$.

Let

$$Q = \left\{ p \in (2k, w] : p \nmid \prod_{1 \leq i < j \leq k} (h_i - h_j) \right\} \setminus \{q_i : i = 1, \dots, t\}.$$

For any prime $q \in Q$, there is an integer $r_q \not\equiv -h_i \pmod{q}$ for all $i = 1, \dots, k$ since \mathcal{H} is admissible.

By the Chinese Remainder Theorem, there is an integer b satisfying the following (1)-(4).

- (1) $b \equiv \delta = \delta_1 \delta_2 \pmod{K}$.
- (2) $b \equiv r_p - h_i \equiv r_p - h_j \pmod{p}$ if $p > 2k$ is a prime dividing $h_i - h_j$ with $1 \leq i < j \leq k$.
- (3) $b \equiv -a_i \pmod{q_i}$ for all $i = 1, \dots, t$.
- (4) $b \equiv r_q \pmod{q}$ for all $q \in Q$.

By the above analysis, $\prod_{s=1}^k (b + h_s)$ is relatively prime to W . As $\mathcal{H}' = \{b + h_s : s = 1, \dots, k\}$ is also an admissible set of cardinality k , for large x there is an integer $n \in [x/W, 2x/W]$ such that there are more than m primes among $Wn + b + h_s$ ($s = 1, \dots, k$). For $a_i \in S$, we have

$$Wn + b + a_i \equiv 0 - a_i + a_i = 0 \pmod{q_i}$$

and hence $Wn + b + a_i$ is composite since $W > q_i$. Therefore, there are consecutive primes $p_N, p_{N+1}, \dots, p_{N+m}$ with $p_{N+i} = Wn + b + h_{s(i)}$ for all $i = 0, \dots, m$, where $1 \leq s(0) < s(1) < \dots < s(m) \leq k$. Note that

$$p_{N+m} - p_N = (Wn + b + h_{s(m)}) - (Wn + b + h_{s(0)}) = h_{s(m)} - h_{s(0)} \leq h_k.$$

For each $s = 1, \dots, k$, clearly $Wn + b + h_s \equiv 0 + \delta + 0 = \delta \pmod{8}$ and hence

$$\left(\frac{-1}{Wn + b + h_s} \right) = \delta \quad \text{and} \quad \left(\frac{2}{Wn + b + h_s} \right) = 1.$$

As $p_{N+i} = Wn + b + h_{s(i)} \equiv \delta \pmod{8}$ for all $i = 0, \dots, m$, by the Quadratic Reciprocal Law we have

$$\left(\frac{p_{N+j}}{p_{N+i}} \right) = \delta \left(\frac{p_{N+i}}{p_{N+j}} \right) \quad \text{for all } 0 \leq i < j \leq m.$$

Let $0 \leq i < j \leq m$. Then

$$\left(\frac{p_{N+i}}{p_{N+j}} \right) = \left(\frac{Wn + b + h_{s(i)}}{Wn + b + h_{s(j)}} \right) = \left(\frac{h_{s(i)} - h_{s(j)}}{Wn + b + h_{s(j)}} \right) = \delta \left(\frac{h_{ij}}{Wn + b + h_{s(j)}} \right),$$

where h_{ij} is the odd part of $h_{s(j)} - h_{s(i)}$. For any prime divisor p of h_{ij} , clearly $p \leq h_k \leq w$ and

$$\left(\frac{p}{Wn + b + h_{s(j)}} \right) = \delta^{(p-1)/2} \left(\frac{Wn + b + h_{s(j)}}{p} \right) = \delta^{(p-1)/2} \left(\frac{b + h_{s(j)}}{p} \right).$$

If $p < 2k$, then $p \mid K$, hence $b + h_j \equiv \delta + 0 \pmod{p}$ and thus

$$\left(\frac{p}{Wn + b + h_{s(j)}} \right) = \delta^{(p-1)/2} \left(\frac{b + h_{s(j)}}{p} \right) = \delta^{(p-1)/2} \left(\frac{\delta}{p} \right) = 1.$$

If $p > 2k$, then by the choice of b we have

$$\begin{aligned} \left(\frac{p}{Wn + b + h_{s(j)}} \right) &= \delta^{(p-1)/2} \left(\frac{b + h_{s(j)}}{p} \right) = \delta^{(p-1)/2} \left(\frac{r_p}{p} \right) \\ &= \left(\frac{r_p \delta}{p} \right) = \begin{cases} \delta_2 & \text{if } p = p_{s(i), s(j)}, \\ 1 & \text{otherwise.} \end{cases} \end{aligned}$$

Recall that $p_{s(i), s(j)} \parallel h_{ij}$. Therefore,

$$\left(\frac{p_{N+i}}{p_{N+j}} \right) = \delta \left(\frac{h_{ij}}{Wn + b + h_{s(j)}} \right) = \delta \delta_2 = \delta_1$$

and

$$\left(\frac{p_{N+j}}{p_{N+i}} \right) = \delta \left(\frac{p_{N+i}}{p_{N+j}} \right) = \delta_2.$$

This concludes the proof. \square

3. PROOF OF THEOREM 1.4

The following lemma is a slight modification of Lemma 2.2 which can be proved in a similar way.

Lemma 3.1. *Let $k > 1$ be an integer. Then there is an admissible set $\mathcal{H} = \{h_1, \dots, h_k\}$ with $h_1 = 0 < h_2 < \dots < h_k$ which has the following properties:*

- (i) All those h_1, h_2, \dots, h_k are multiples of $K = 4 \prod_{p < 4k} p$.
- (ii) Each $h_i - h_j$ with $1 \leq i < j \leq k$ has a prime divisor $p > 4k$ with $h_i \not\equiv h_j \pmod{p^2}$.
- (iii) If $1 \leq i < j \leq k$, $1 \leq s < t \leq k$ and $\{i, j\} \neq \{s, t\}$, then no prime $p > 4k$ divides both $h_i - h_j$ and $h_s - h_t$.

Lemma 3.2. *Let $k > 1$ be an integer, and let $\mathcal{H} = \{h_1, \dots, h_k\}$ with $h_1 = 0 < h_2 < \dots < h_k$ be an admissible set satisfying (i)-(iii) in Lemma 3.1. Then there is a positive integer b with all of the following properties:*

- (i) $\prod_{i=1}^k (b + h_i)$ is relatively prime to the least common multiple W of those $h_j - h_i$ with $1 \leq i < j \leq k$ and $\prod_{2 < p \leq w} p$ if w is large enough.
- (ii) $\prod_{i=1}^k (b + h_i - 1)$ is relatively prime to $\prod_{2 < p \leq w} p$ if w is large enough.
- (iii) For any $i, j \in \{1, \dots, k\}$ with $i \neq j$, we have

$$\left(\frac{h_i - h_j}{b + h_j} \right) = -1.$$

- (iv) If $n > b$, $n \equiv b \pmod{W}$ and $a \in \{h_1, h_1 + 1, \dots, h_k\} \setminus \mathcal{H}$, then $n + a$ is not prime.

Proof. For any $1 \leq i < j \leq k$, the number $h_i - h_j$ has a prime divisor $p_{ij} > 4k$ with $h_i \not\equiv h_j \pmod{p_{ij}^2}$. Suppose that $p > 4k$ is a prime dividing

$\prod_{1 \leq i < j \leq k} (h_i - h_j)$, then there is a unique pair $\{i, j\}$ with $1 \leq i < j \leq k$ such that $h_i \equiv h_j \pmod{p}$. Note that $p \leq h_k$. As $h_i - h_j \equiv h_i - h_i \pmod{p}$ and $2(k-1) < (p-1)/2$, there is an integer $r_p \not\equiv h_i - h_s, h_i - h_s + 1 \pmod{p}$ for all $s = 1, \dots, k$ such that

$$\left(\frac{r_p}{p}\right) = \begin{cases} -\left(\frac{3}{p}\right)^{\text{ord}_3(h_j - h_i)} & \text{if } p = p_{ij}, \\ 1 & \text{otherwise.} \end{cases}$$

So, for any integer $b \equiv r_p - h_i \pmod{p}$, we have $b + h_s \not\equiv 0, 1 \pmod{p}$.

Assume that

$$S = \{h_1 < a < h_k : a \neq h_s, h_s - 1 \text{ for all } s = 1, \dots, k\} = \{a_i : i = 1, \dots, t\}.$$

Clearly $t \leq h_k - k$ and hence we may choose t distinct primes $q_1, \dots, q_t \in (h_k, w]$ if w is large enough. If $b \equiv -a_i \pmod{q_i}$, then $b + h_s \equiv h_s - a_i \not\equiv 0, 1 \pmod{q_i}$ for all $s = 1, \dots, k$ since $|h_s - a_i| < h_k < q_i$.

Let

$$Q = \left\{ p \in (4k, w] : p \nmid \prod_{1 \leq i < j \leq k} (h_i - h_j) \right\} \setminus \{q_i : i = 1, \dots, t\}.$$

For any prime $q \in Q$, there is an integer $r_q \not\equiv -h_i, -h_i + 1 \pmod{q}$ for all $i = 1, \dots, k$ since $q > 2k$.

By the Chinese Remainder Theorem, there is a positive integer b satisfying the following (1)-(4).

- (1) $b \equiv 17 \pmod{24}$, and $b \equiv 4 \pmod{p}$ for all primes $p \in [5, 4k]$.
- (2) $b \equiv r_p - h_i \equiv r_p - h_j \pmod{p}$ if $p > 4k$ is a prime dividing $h_i - h_j$ with $1 \leq i < j \leq k$.
- (3) $b \equiv -a_i \pmod{q_i}$ for all $i = 1, \dots, t$.
- (4) $b \equiv r_q \pmod{q}$ for all $q \in Q$.

By the above analysis, $\prod_{s=1}^k (b + h_s)(b + h_s - 1)$ is relatively prime to $\prod_{2 < p \leq w} p$. Note that $b + h_i \equiv 17 + 0 \pmod{24}$ for all $i = 1, \dots, k$. If $w \geq h_k$, then any prime divisor of W does not exceed w . So both (i) and (ii) holds.

For each $s = 1, \dots, k$, clearly $b + h_s \equiv 17 + 0 \equiv 1 \pmod{8}$ and hence

$$\left(\frac{-1}{b + h_s}\right) = \left(\frac{2}{b + h_s}\right) = 1.$$

Let $i, j \in \{0, \dots, m\}$ with $i \neq j$. Then

$$\left(\frac{h_i - h_j}{b + h_j}\right) = \left(\frac{h_{ij}}{b + h_j}\right),$$

where h_{ij} is the odd part of $|h_i - h_j|$. For any prime divisor p of h_{ij} , clearly $p \leq h_k \leq w$ and

$$\left(\frac{p}{b + h_j}\right) = \left(\frac{b + h_j}{p}\right).$$

If $3 < p < 4k$, then $p \mid K$, hence $b + h_j \equiv 4 + 0 \pmod{p}$ and thus

$$\left(\frac{p}{b+h_j}\right) = \left(\frac{b+h_j}{p}\right) = \left(\frac{4}{p}\right) = 1.$$

If $p > 4k$, then by the choice of b we have

$$\begin{aligned} \left(\frac{p}{b+h_j}\right) &= \left(\frac{b+h_j}{p}\right) = \left(\frac{r_p}{p}\right) \\ &= \begin{cases} -\left(\frac{3}{p}\right)^{\text{ord}_3(h_j-h_i)} & \text{if } p = p_{\min\{i,j\}, \max\{i,j\}}, \\ 1 & \text{otherwise.} \end{cases} \end{aligned}$$

Recall that $p_{ij} \parallel h_{ij}$. Therefore,

$$\left(\frac{h_i-h_j}{b+h_j}\right) = \left(\frac{h_{ij}}{b+h_j}\right) = -1.$$

So (iii) in Lemma 3.2 also holds.

Now suppose that $n > b$ is an integer with $n \equiv b \pmod{W}$, and that $a \in \{h_1, h_1 + 1, \dots, h_k\} \setminus \mathcal{H}$. If $a = h_s - 1$ for some $1 \leq s \leq k$, then $n + a \equiv b + h_s - 1 \equiv 0 \pmod{4}$ and hence $n + a$ is not prime. If $a \neq h_s - 1$ for all $s = 1, \dots, k$, then $a = a_i$ for some $1 \leq i \leq t$, hence $n + a \equiv b + a_i \equiv 0 \pmod{q_i}$ and thus $n + a$ is not prime. (Note that $n + a > W > w \geq q_i$.) Thus (iv) of Lemma 3.2 also holds.

In view of the above, we have completed the proof of Lemma 3.2. \square

Proof of Theorem 1.4. Choose k (depending on m) as in Pollack [P] in the spirit of Mynard-Tao's work. Let $\mathcal{H} = \{h_1, h_2, \dots, h_k\}$ be an admissible set constructed in Lemma 3.1 and choose an integer b as in Lemma 3.2. Let x be sufficiently large, and let W be the least common multiple of those $h_j - h_i$ ($1 \leq i < j \leq k$) and $\prod_{2 < p \leq \log \log \log x} p$. Then we have an analogue of Lemma 3.3 of Pollack [P]. When $n + h_i$ and $n + h_j$ ($i \neq j$) are both prime with $n \equiv b \pmod{W}$, $n + h_i$ is a primitive root modulo $n + h_j$ if and only if $|h_i - h_j|$ is a primitive root modulo $n + h_j$ since $n + h_j \equiv 1 \pmod{4}$.

Let P be the set of all primes. For $j = 1 \dots, k$, set

$$P_j := \{p \in P : |h_i - h_j| \text{ is a primitive root modulo } p \text{ for any } i \neq j\}.$$

Define the weight function $w(n)$ as in [M, Proposition 4.1] or [P, Proposition 3.1], and let $\chi_A(x)$ be the characteristic function of the set A . We need to show that

$$\sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \left(\sum_{j=1}^k \chi_{P_j}(n + h_j) \right) w(n) \sim \sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \left(\sum_{j=1}^k \chi_P(n + h_j) \right) w(n). \quad (3.1)$$

For a prime q and an integer g , define

$$P_q(g) = \{p \in P : p \equiv 1 \pmod{q} \text{ and } g^{(p-1)/q} \equiv 1 \pmod{p}\}$$

and

$$\mathcal{P}_q(g) = P_q(g) \setminus \bigcup_{q' < q} P_{q'}(g).$$

Pollack [P, the estimations of $\Sigma_1 - \Sigma_4$] showed that if $(\frac{g}{b+h_j}) = -1$ then

$$\sum_{q \in P} \sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \chi_{\mathcal{P}_q(g)}(n+h_j)w(n) = o\left(\frac{\varphi(W)^k}{W^{k+1}}x(\log x)^k\right).$$

Note that if $n \in P \setminus P_j$ then $n \in \mathcal{P}_q(|h_i - h_j|)$ for some $i \neq j$ and some prime q . Hence

$$\begin{aligned} & \sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \left(\sum_{j=1}^k \chi_P(n+h_j) \right) w(n) - \sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \left(\sum_{j=1}^k \chi_{P_j}(n+h_j) \right) w(n) \\ & \leq \sum_{j=1}^k \sum_{\substack{i=1 \\ i \neq j}}^k \sum_{q \in P} \sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \chi_{\mathcal{P}_q(|h_i - h_j|)}(n+h_j)w(n) = o\left(\frac{\varphi(W)^k}{W^{k+1}}x(\log x)^k\right). \end{aligned}$$

Maynard and Tao (cf. [M]) have proved

$$\sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} w(n) \sim \frac{\alpha \phi(W)^k}{W^{k+1}} x(\log x)^k \quad (3.2)$$

and

$$\sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \left(\sum_{j=1}^k \chi_P(n+h_j) \right) w(n) \sim \frac{\beta k \phi(W)^k}{W^{k+1}} x(\log x)^k, \quad (3.3)$$

where α and β are positive constants only depending on k and w . It follows from (3.1) and (3.3) that

$$\sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \left(\sum_{j=1}^k \chi_{P_j}(n+h_j) \right) w(n) \sim \frac{\beta k \phi(W)^k}{W^{k+1}} x(\log x)^k.$$

Furthermore, in view of [M], we may choose a sufficiently large integer k and a suitable weight function w such that

$$\beta k > m\alpha,$$

i.e.,

$$\sum_{\substack{x \leq n \leq 2x \\ n \equiv b \pmod{W}}} \left(\sum_{j=1}^k \chi_{P_j}(n + h_j) - m\alpha \right) w(n) > 0.$$

Since $w(n)$ is non-negative, for some $n \in [x, 2x]$ with $n \equiv b \pmod{W}$, $\{n + h_1, \dots, n + h_k\}$ contains at least $m + 1$ primes $n + h_j$ ($j \in J$) with $|J| > m$ and $n + h_j \in P_j$ for $j \in J$. According to the construction of b and Lemma 3.2 (iv), for each $j = 1, \dots, k$, the interval $(n + h_j, n + h_{j+1})$ contains no prime. So those primes in $\{n + h_1, \dots, n + h_k\}$ are consecutive primes. For any $i, j \in J$ with $i \neq j$, the number $h_i - h_j$, as well as the prime $n + h_i$, is a primitive root modulo the prime $n + h_j$. This concludes the proof. \square

REFERENCES

- [BFTB] W.D. Banks, T. Freiberg and C.L. Turnage-Butterbaugh, *Consecutive primes in tuples*, Acta Arith. **167** (2015), 261–266.
- [P] P. Pollack, *Bounded gaps between primes with a given primitive root*, preprint, [arXiv:1404.4007](https://arxiv.org/abs/1404.4007).
- [Po] D.H.J. Polymath, *Bounded gaps between primes*, Polymath 8 Project. Available from http://michaelnielsen.org/polymath1/index.php?title=Bounded_gaps_between_primes.
- [M] J. Maynard, *Small gaps between primes*, Annals of Math. **181** (2015), 383–413.
- [S] D.K.L. Shiu, *Strings of congruent primes*, J. London Math. Soc. **61** (2000), 359–373.
- [Su1] Z.-W. Sun, *New observations on primitive roots modulo primes*, preprint, [arXiv:1405.0290](https://arxiv.org/abs/1405.0290).
- [Su2] Z.-W. Sun, Sequences A243839 and A243901 in OEIS (On-Line Encyclopedia of Integer Sequences), <http://oeis.org>.
- [Z] Y. Zhang, *Bounded gaps between primes*, Annals of Math. **179** (2014), 1121–1174.