

Vertex and edge orbits of Fibonacci and Lucas cubes

Ali Reza Ashrafi ^a Jernej Azarija ^b Khadijeh Fathalikhani ^a
 Sandi Klavžar ^{c,d,b} Marko Petkovšek ^{c,b}

^a Department of Pure Mathematics, Faculty of Mathematical Sciences, University of Kashan, Kashan, Iran

^b Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

^c Faculty of Mathematics and Physics, University of Ljubljana, Slovenia

^d Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

Abstract

The Fibonacci cube Γ_n is obtained from the n -cube Q_n by removing all the vertices that contain two consecutive 1s. If, in addition, the vertices that start and end with 1 are removed, the Lucas cube Λ_n is obtained. The number of vertex and edge orbits, the sets of the sizes of the orbits, and the number of orbits of each size, are determined for the Fibonacci cubes and the Lucas cubes under the action of the automorphism group. In particular, the set of the sizes of the vertex orbits of Λ_n is $\{k \geq 1; k|n\} \cup \{k \geq 18; k|2n\}$, the number of the vertex orbits of Λ_n of size k , where k is odd and divides n , is equal to $\sum_{d|k} \mu\left(\frac{k}{d}\right) F_{\lfloor \frac{d}{2} \rfloor + 2}$, and the number of the edge orbits of Λ_n is equal to the number of the vertex orbits of Γ_{n-3} when $n \geq 5$. Primitive strings, dihedral transformations and asymmetric strings are essential tools to prove these results.

Key words: Fibonacci cube; Lucas cube; dihedral transformation; primitive string, vertex orbit; edge orbit

AMS Subj. Class: 68R15, 05C30

1 Introduction

Fibonacci cubes Γ_n [5] and the closely related Lucas cubes Λ_n [13] have been investigated from many points of view, let us briefly overview some recent achievements. Formulas for the number of vertices of a given degree as well as the corresponding

generating functions were determined in [9], while the domination number and the 2-packing number of these cubes were studied in [1, 16]. Motivated by the structure of Γ_n and Λ_n as interconnection networks, Mollard [12] characterized maximal induced hypercubes in these cubes and also determined the number of such hypercubes. From the metric graph theory point of view, eccentricity sequences were obtained in [2], the Wiener index and the Hosoya polynomial were determined in [7], while in [8] the asymptotic average eccentricity was determined. In the latter paper it is also proved that the eccentricity of a vertex of a given Fibonacci cube is equal to the depth of the associated leaf in the corresponding Fibonacci tree. For a connection between Fibonacci/Lucas cubes and Hasse diagrams (of the independent subsets of powers of paths and cycles) see [4]. From the chemical graph theory perspective we point out that Lucas cubes turned out to be precisely the so-called resonance graphs of cyclic fibonacenes [24]. Very recently several advances were made also from the algorithmic point of view. Linear recognition algorithm for Fibonacci cubes and for Lucas cubes were developed by Vesel [21] and Taranenko [19], respectively, while Ramras [17] studied off-line routing of linear permutations on these cubes. For additional information on Fibonacci cubes see the survey [6].

There are several reasons for this wide interest. These cubes are induced subgraphs of hypercubes that inherit many of the fine properties of the latter class. The main tool to derive such properties for Fibonacci cubes is the so-called fundamental decomposition that decomposes Γ_n into Γ_{n-1} and Γ_{n-2} , similarly as the n -cube decomposes into two $(n-1)$ -cubes via the Cartesian product operation. (There is also a similar decomposition for Lucas cubes.) On the other hand, the order of Fibonacci/Lucas cubes grows much slower than that of hypercubes, a property important for interconnection networks. A strong source of interest for these cubes also comes from theoretical chemistry, where Fibonacci cubes are precisely the so-called resonance graphs of fibonacenes [10] (see [22] for a generalization of this result), while for the role of Lucas cubes in chemistry, besides the already mentioned paper [25], see also [23, 25]. Fibonacci cubes also led to the notion of the Fibonacci dimension of a graph [3, 20].

When it comes to symmetries, it seems that only the automorphism groups of Fibonacci and Lucas cubes have been determined so far [1]. Hence, in this paper we take a closer look at their symmetries, more precisely at the orbits under the action of the automorphism group. We proceed as follows. The next two sections are of preliminary nature. In the first of them we introduce concepts and notations needed, and recall or prove some related results. In the subsequent section we investigate some properties of dihedral transformations of nonempty strings defined over a finite alphabet. In Section 4, we determine the number of vertex and edge orbits, the sets of the sizes of the orbits, and the number of orbits of each size of Fibonacci cubes, as well as give a combinatorial interpretation for the number of vertex orbits. In the last section we prove parallel results for Lucas cubes. Contrary to Fibonacci cubes, where there are only orbits of size 1 and 2, the situation with Lucas cubes is more

intriguing and complex.

2 Preliminaries

In this section we formally introduce the cubes studied here, list some notation, and prove an identity on the Lucas numbers. The n -cube Q_n , $n \geq 0$, is the graph whose vertex set is the set of all binary strings of length n , two vertices being adjacent if they differ in exactly one position. The n -dimensional *Fibonacci cube* Γ_n is the subgraph of Q_n induced by the set of all vertices that have no two consecutive 1s. Strings with no two consecutive 1s are called *Fibonacci strings*. The n -dimensional *Lucas cube* Λ_n is obtained from Γ_n by removing all the vertices that begin and end with 1. The vertices of Lucas cubes are called *Lucas strings*. The Fibonacci numbers F_n are defined by $F_0 = 0$, $F_1 = 1$, and $F_n = F_{n-1} + F_{n-2}$, $n \geq 2$, and the Lucas numbers L_n by $L_0 = 2$, $L_1 = 1$, and $L_n = L_{n-1} + L_{n-2}$, $n \geq 2$. We will use the following well-known facts about F_n , L_n , Γ_n , and Λ_n without special mention:

1. $L_n = F_{n-1} + F_{n+1}$ for $n \geq 1$,
2. $|V(\Gamma_n)| = F_{n+2}$ for $n \geq 0$,
3. $|E(\Gamma_n)| = (nF_{n+1} + 2(n+1)F_n)/5$ for $n \geq 0$,
4. $|V(\Lambda_n)| = L_n$ for $n \geq 1$, $|V(\Lambda_0)| = 1$,
5. $|E(\Lambda_n)| = nF_{n-1}$ for $n \geq 0$.

Proposition 2.1

$$F_{n+1} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} \quad \text{for } n \geq -1, \quad (1)$$

$$L_n = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n}{n-k} \binom{n-k}{k} \quad \text{for } n \geq 1, \quad (2)$$

$$\sum_{i=0}^n F_i L_{n-i} = (n+1)F_n \quad \text{for } n \geq 0. \quad (3)$$

Proof. Identities (1) and (3) are well known. To prove (2), note that by using (1) twice and shifting the index of summation in the first sum,

$$\begin{aligned}
L_n &= F_{n-1} + F_{n+1} = \sum_{k=0}^{\lfloor \frac{n-2}{2} \rfloor} \binom{n-k-2}{k} + \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} \\
&= \sum_{k=1}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k-1}{k-1} + \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} \\
&= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \left(\frac{k}{n-k} + 1 \right) \binom{n-k}{k} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n}{n-k} \binom{n-k}{k}. \quad \square
\end{aligned}$$

As usual, the automorphism group of a graph $G = (V, E)$ will be denoted by $\text{Aut}(G)$. The sets of orbits of $\text{Aut}(G)$ acting on V resp. E will be denoted by $\mathcal{O}_V(G)$ resp. $\mathcal{O}_E(G)$, and their cardinalities by $o_V(G) = |\mathcal{O}_V(G)|$ and $o_E(G) = |\mathcal{O}_E(G)|$. In the latter case, we consider the natural action of $\text{Aut}(G)$ on E , that is, for $g \in \text{Aut}(G)$ the edge $\{u, v\}$ is mapped to $\{g(u), g(v)\}$. We denote the orbit of $u \in V$ resp. $e \in E$ under the action of $\text{Aut}(G)$ by \bar{u} resp. \bar{e} , and its cardinality by $|\bar{u}|$ resp. $|\bar{e}|$. In addition, we denote the number of orbits of size k by

$$\begin{aligned}
o_V(G, k) &= |\{X \in \mathcal{O}_V(G); |X| = k\}|, \\
o_E(G, k) &= |\{Y \in \mathcal{O}_E(G); |Y| = k\}|,
\end{aligned}$$

so that the following identities hold:

$$\sum_k k o_V(G, k) = |V(G)|, \quad \sum_k o_V(G, k) = o_V(G), \quad (4)$$

$$\sum_k k o_E(G, k) = |E(G)|, \quad \sum_k o_E(G, k) = o_E(G). \quad (5)$$

We denote the dihedral group of order $2n$ by D_n , and the set of fixed points of a group element g acting on a set A , resp. its cardinality, by

$$\text{Fix}_A(g) = \{u \in A; g(u) = u\}, \quad \text{fix}_A(g) = |\text{Fix}_A(g)|.$$

Finally, \mathbb{N} is the set of all positive integers $\{1, 2, 3, \dots\}$. For $n \in \mathbb{N}$, $[n]$ denotes the set $\{1, 2, \dots, n\}$, and $[n]_0$ denotes the set $\{0, 1, \dots, n-1\}$.

3 Dihedral transformations of strings

As a preparation for what follows we investigate here some properties of dihedral transformations of nonempty strings defined over a finite alphabet.

Let Σ be an alphabet such that $0 \in \Sigma$. As usual, Σ^n denotes the set of all strings of length n over Σ and $\Sigma^+ = \bigcup_{n=1}^{\infty} \Sigma^n$ denotes the set of all nonempty strings over Σ . If $u, v \in \Sigma^+$ and $k \in \mathbb{N}$, we write uv for the concatenation of u and v , and u^k for the concatenation of k copies of u . For $u = u_1u_2 \cdots u_n \in \Sigma^n$, we define its *length* $|u|$, *weight* $w(u) \in [n] \cup \{0\}$, *cyclic shift* $\alpha(u) \in \Sigma^n$, and *reversal* $\beta(u) \in \Sigma^n$ by

$$\begin{aligned} |u| &= n, \\ w(u) &= |\{i \in [n]; u_i \neq 0\}|, \\ \alpha(u) &= u_n u_1 \cdots u_{n-1}, \\ \beta(u) &= u_n u_{n-1} \cdots u_1. \end{aligned}$$

It is straightforward to verify that

$$x = \alpha^j(u) \iff x_i = u_{(i-j) \bmod n} \iff u_i = x_{(i+j) \bmod n}, \quad (6)$$

$$x = \beta(u) \iff x_i = u_{(1-i) \bmod n} \iff u_i = x_{(1-i) \bmod n}, \quad (7)$$

$$x = \alpha^j \beta(u) \iff x_i = u_{(1-i+j) \bmod n} \iff u_i = x_{(1-i+j) \bmod n}, \quad (8)$$

where $a \bmod n$ is the unique $i \in [n]$ such that $a \equiv i \pmod{n}$. Since $\alpha^n = \beta^2 = \text{id}$ and $\alpha\beta = \beta\alpha^{-1}$, the group generated by α and β represents the action of D_n on Σ^n . We denote the orbit of $u \in \Sigma^n$ under this action by \bar{u} , and its cardinality by $|\bar{u}|$.

Lemma 3.1 *For all $u \in \Sigma^+$, $j \in \mathbb{Z}$, and $k \in \mathbb{N}$, we have*

$$(i) \quad \alpha^j(u^k) = (\alpha^j(u))^k,$$

$$(ii) \quad \beta(u^k) = \beta(u)^k.$$

Proof. This follows from (6), (7), and the fact that $(u^k)_i = u_{i \bmod |u|}$. \square

For $u \in \Sigma^+$ define its *period* $p(u)$ and *exponent* $\ell(u)$ by

$$\begin{aligned} p(u) &= \min\{k > 0; \alpha^k(u) = u\}, \\ \ell(u) &= \max\{k > 0; \exists v \in \Sigma^+ : v^k = u\}. \end{aligned}$$

If $\ell(u) = 1$ then u is *primitive*. It is well known [11, Cor. 4.2] that for each u there is a unique primitive string $\tau(u) \in \Sigma^+$ (called the *root* of u) such that $\tau(u)^{\ell(u)} = u$.

As an example, consider $\Sigma = \{0, 1\}$ and the following strings $u \in \Sigma^4$:

- $u = 0000 = 0^4$: $p(u) = 1$, $\tau(u) = 0$, $\ell(u) = 4$, u is not primitive,
- $u = 0101 = (01)^2$: $p(u) = 2$, $\tau(u) = 01$, $\ell(u) = 2$, u is not primitive,
- $u = 0011 = (0011)^1$: $p(u) = 4$, $\tau(u) = 0011$, $\ell(u) = 1$, u is primitive.

Proposition 3.2 *If $u \in \Sigma^n$, then*

- (i) $p(u) = |\{u, \alpha(u), \alpha^2(u), \dots, \alpha^{n-1}(u)\}|$,
- (ii) $p(u^k) = p(u)$ for all $k \in \mathbb{N}$,
- (iii) $p(u) \mid n$,
- (iv) $p(u) = p(\tau(u)) = |\tau(u)|$ and $\ell(u)p(u) = n$.

Proof.

- (i) Immediate from the definition of $p(u)$.
- (ii) Since $u^k = v^k$ if and only if $u = v$, this follows from (i) and Lemma 3.1(ii).
- (iii) By (i), $p(u)$ is the size of the orbit of u under the action of the cyclic group of order n generated by α , hence $p(u)$ divides n .
- (iv) Let $u = u_1u_2 \cdots u_n$ and $v = u_1u_2 \cdots u_{p(u)}$. From $\alpha^{p(u)}(u) = u$ it follows by (6) that $u_i = u_{i+p(u)}$ for $i = 1, 2, \dots, n - p(u)$ (with indices taken mod n). By induction on k , $u_i = u_{i+kp(u)}$ for $i = 1, 2, \dots, n - kp(u)$ and $k = 0, 1, \dots, \frac{n}{p(u)} - 1$. Hence $v = u_{1+kp(u)}u_{2+kp(u)} \cdots u_{p(u)+kp(u)}$ for $k = 0, 1, \dots, \frac{n}{p(u)} - 1$, therefore $u = v^{n/p(u)}$. If u is primitive, this implies that $|u|/p(u) = 1$ and so $p(u) = |u|$. For arbitrary u we then have, by (ii),

$$p(u) = p(\tau(u)^{\ell(u)}) = p(\tau(u)) = |\tau(u)|,$$

$$\text{so } \ell(u)p(u) = \ell(u)|\tau(u)| = |\tau(u)^{\ell(u)}| = |u| = n. \quad \square$$

Proposition 3.3 *For each $u \in \Sigma^+$,*

$$|\bar{u}| = \begin{cases} p(u), & \exists j : \beta(u) = \alpha^j(u), \\ 2p(u), & \forall j : \beta(u) \neq \alpha^j(u). \end{cases}$$

Proof. Denote

$$\begin{aligned} A(u) &= \{u, \alpha(u), \alpha^2(u), \dots, \alpha^{n-1}(u)\}, \\ B(u) &= \{\beta(u), \alpha\beta(u), \alpha^2\beta(u), \dots, \alpha^{n-1}\beta(u)\}. \end{aligned}$$

Then $\bar{u} = A(u) \cup B(u)$. If $\beta(u) = \alpha^j(u)$ for some j , it follows from $\alpha^n = \text{id}$ that $B(u) = A(u)$, hence $\bar{u} = A(u)$ and, by Proposition 3.2(i), $|\bar{u}| = p(u)$.

Otherwise, if $\beta(u) \neq \alpha^j(u)$ for all j , it follows from $\alpha^n = \text{id}$ that $B(u) \cap A(u) = \emptyset$ and $|B(u)| = |A(u)|$, so, by Proposition 3.2(i), $|\bar{u}| = 2|A(u)| = 2p(u)$. \square

In order to determine the orbit size $|\bar{u}|$ more precisely, we distinguish between symmetric and asymmetric strings, where for $n \in \mathbb{N}$ a string $u \in \Sigma^n$ will be called:

- *symmetric* if $|\bar{u}| < 2n$,
- *asymmetric* if $|\bar{u}| = 2n$.

Proposition 3.4 *Every asymmetric string is primitive.*

Proof. Let $u \in \Sigma^n$ be asymmetric. Then $|\bar{u}| = 2n$, hence by Proposition 3.3, $p(u) = n$. Then by Proposition 3.2(iv), $\ell(u) = 1$ and u is primitive. \square

As an example, consider $\Sigma = \{0, 1\}$ and the following strings $u \in \Sigma^6$:

- $u = 000000$ is symmetric and not primitive,
- $u = 001100$ is symmetric and primitive,
- $u = 010011$ is asymmetric (and hence primitive).

Theorem 3.5 *For each $u \in \Sigma^+$,*

$$|\bar{u}| = \begin{cases} p(u), & \tau(u) \text{ symmetric,} \\ 2p(u), & \tau(u) \text{ asymmetric.} \end{cases}$$

Proof. Using Propositions 3.3, 3.2(iv) and Lemma 3.1 repeatedly we obtain

$$\begin{aligned} |\bar{u}| = 2p(u) &\iff \forall j : \beta(u) \neq \alpha^j(u) \iff \forall j : \beta(\tau(u)^{\ell(u)}) \neq \alpha^j(\tau(u)^{\ell(u)}) \\ &\iff \forall j : \beta(\tau(u))^{\ell(u)} \neq \alpha^j(\tau(u))^{\ell(u)} \\ &\iff \forall j : \beta(\tau(u)) \neq \alpha^j(\tau(u)) \iff |\overline{\tau(u)}| = 2p(\tau(u)) \\ &\iff |\overline{\tau(u)}| = 2|\tau(u)| \iff \tau(u) \text{ asymmetric.} \end{aligned}$$

Together with Proposition 3.3 this proves the claim. \square

Corollary 3.6 *A string $u \in \Sigma^n$ is primitive symmetric if and only if it is primitive and $\alpha^j \beta(u) = u$ for some $j \in [n]_0$.*

Proof. By Theorem 3.5 and Proposition 3.3, we have

$$\begin{aligned} u \text{ primitive symmetric} &\iff u \text{ primitive} \wedge |\bar{u}| = p(u) \\ &\iff u \text{ primitive} \wedge \exists j \in [n]_0 : \beta(u) = \alpha^j(u) \\ &\iff u \text{ primitive} \wedge \exists j \in [n]_0 : \alpha^j \beta(u) = u. \quad \square \end{aligned}$$

4 Orbits of Fibonacci cubes

According to [1], for $n \geq 1$ the Fibonacci cube Γ_n admits exactly one non-trivial automorphism, hence $|\text{Aut}(\Gamma_n)| = 2$ and the only orbit sizes are 1 and 2. For $n \geq 2$, the non-trivial automorphism coincides with the reversal map $\beta : V(\Gamma_n) \rightarrow V(\Gamma_n)$. We denote the set of Fibonacci strings of length n which start with 0 resp. 1 by $V_0(\Gamma_n)$ resp. $V_1(\Gamma_n)$. We begin by enumerating Fibonacci palindromes.

Proposition 4.1 *For $k \in \mathbb{N}$,*

$$\begin{aligned} \text{fix}_{V(\Gamma_{2k})}(\beta) &= F_{k+1}, & \text{fix}_{V(\Gamma_{2k+1})}(\beta) &= F_{k+3}, \\ \text{fix}_{V_0(\Gamma_{2k})}(\beta) &= F_k, & \text{fix}_{V_0(\Gamma_{2k+1})}(\beta) &= F_{k+2}, \\ \text{fix}_{V_1(\Gamma_{2k})}(\beta) &= F_{k-1}, & \text{fix}_{V_1(\Gamma_{2k+1})}(\beta) &= F_{k+1}. \end{aligned}$$

Proof. Let $u \in \text{fix}_{V(\Gamma_n)}(\beta)$. We distinguish four cases:

1. $n = 2k$

(a) u starts with 0: $u = 0v00\beta(v)0$ with $v \in V(\Gamma_{k-2})$

(b) u starts with 1: $u = 10v00\beta(v)01$ with $v \in V(\Gamma_{k-3})$

2. $n = 2k + 1$

(a) u starts with 0: $u = 0v0\beta(v)0$ with $v \in V(\Gamma_{k-1})$, or $u = 0v010\beta(v)0$ with $v \in V(\Gamma_{k-2})$

(b) u starts with 1: $u = 10v0\beta(v)01$ with $v \in V(\Gamma_{k-2})$, or $u = 10v010\beta(v)01$ with $v \in V(\Gamma_{k-3})$

The stated equalities now follow from $|V(\Gamma_k)| = F_{k+2}$ and $F_k + F_{k+1} = F_{k+2}$. \square

4.1 Vertex orbits

Theorem 4.2 *Let $n \geq 2$. Then*

$$\begin{aligned} o_V(\Gamma_n, 1) &= F_{\lfloor \frac{n-(-1)^n}{2} \rfloor + 2}, \\ o_V(\Gamma_n, 2) &= \frac{1}{2} \left(F_{n+2} - F_{\lfloor \frac{n-(-1)^n}{2} \rfloor + 2} \right), \\ o_V(\Gamma_n) &= \frac{1}{2} \left(F_{n+2} + F_{\lfloor \frac{n-(-1)^n}{2} \rfloor + 2} \right). \end{aligned}$$

Proof. The orbits of size 1 correspond to the fixed points of β , i.e., to Fibonacci palindromes of length n , hence by Proposition 4.1,

$$o_V(\Gamma_n, 1) = \begin{cases} F_{k+1}, & \text{if } n = 2k, \\ F_{k+3}, & \text{if } n = 2k + 1, \end{cases}$$

which can be combined into $F_{\lfloor \frac{n-(-1)^n}{2} \rfloor + 2}$. The remaining two equalities now follow from (4) which in this case transforms into

$$\begin{aligned} o_V(\Gamma_n, 1) + 2o_V(\Gamma_n, 2) &= F_{n+2}, \\ o_V(\Gamma_n, 1) + o_V(\Gamma_n, 2) &= o_V(\Gamma_n). \quad \square \end{aligned}$$

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$ V(\Gamma_n) $	2	3	5	8	13	21	34	55	89	144	233	377	610	987	1597
$o_V(\Gamma_n)$	1	2	4	5	9	12	21	30	51	76	127	195	322	504	826
$o_V(\Gamma_n, 1)$	0	1	3	2	5	3	8	5	13	8	21	13	34	21	55
$o_V(\Gamma_n, 2)$	1	1	1	3	4	9	13	25	38	68	106	182	288	483	771

Table 1: The numbers of vertices, all orbits, orbits of size 1, and orbits of size 2 in $V(\Gamma_n)$ for $n \leq 15$

We remark that the numbers $o_V(\Gamma_n)$ appear as solutions of other combinatorial enumeration problems as well. For instance, in [14], the following problem is posed and solved: in how many ways can a $2 \times (n+1)$ rectangle be tiled with dominoes (i.e., rectangles of sizes 2×1 and 1×2)? More precisely, the problem asks for the number of *distinct* tilings, where two tilings are considered distinct if one cannot be obtained from the other by reflections and rotations. As it turns out, the answer is given by $o_V(\Gamma_n)$ (see also [18, sequence A001224]). For $n \in \{0, 1\}$, this can be checked directly (note that the 2×2 square has a single distinct tiling, due to the 90° rotation). For $n \geq 2$, we present here a bijective proof of this fact:

Let $u \in V(\Gamma_n)$ and $v = u0 \in V(\Gamma_{n+1})$. Assign to v a tiling of the $2 \times (n+1)$ rectangle with dominoes as follows. Going through v from left to right, assign to each 0 a vertical domino and to each 10 a pair of horizontal dominoes. Conversely, to each tiling of the $2 \times (n+1)$ rectangle assign $v \in V(\Gamma_{n+1})$ by going through the tiling from left to right, coding vertical dominoes with 0 and pairs of horizontal dominoes with 10. Then v ends with 0; let $u \in V(\Gamma_n)$ be v without the final 0. This establishes a bijection between $V(\Gamma_n)$ and the set of all tilings of the $2 \times (n+1)$ rectangle which preserves palindromes in both directions. Hence it gives rise to a bijection between $\mathcal{O}_V(\Gamma_n)$ and the set of all distinct tilings of the $2 \times (n+1)$ rectangle.

Another family of combinatorial objects enumerated by $o_V(\Gamma_n)$ are ordered integer partitions of $n+1$ with parts taken from the set $\{1, 2\}$ where two partitions are considered distinct if one cannot be obtained from the other by reflection. Such partitions are obviously in bijection with distinct domino tilings of the $2 \times (n+1)$ rectangle: to each part 1 in the partition assign a vertical domino, to each part 2 in the partition assign a pair of horizontal dominoes, and vice versa.

4.2 Edge orbits

Theorem 4.3 For all $n \geq 0$,

$$\begin{aligned} o_E(\Gamma_n, 1) &= \frac{1-(-1)^n}{2} F_{\lfloor \frac{n+1}{2} \rfloor}, \\ o_E(\Gamma_n, 2) &= \frac{1}{10} (nF_{n+1} + 2(n+1)F_n) - \frac{1-(-1)^n}{4} F_{\lfloor \frac{n+1}{2} \rfloor}, \\ o_E(\Gamma_n) &= \frac{1}{10} (nF_{n+1} + 2(n+1)F_n) + \frac{1-(-1)^n}{4} F_{\lfloor \frac{n+1}{2} \rfloor}. \end{aligned}$$

Proof. For $n \in \{0, 1\}$, this can be checked directly. Let $n \geq 2$. The orbits of size 1 correspond to the fixed points of β , i.e., to pairs $\{u, v\} \in E(\Gamma_n)$ such that $\{\beta(u), \beta(v)\} = \{u, v\}$. Since u and v differ in weight while β preserves it, this is only possible if $\beta(u) = u$ and $\beta(v) = v$. Hence u and v are Fibonacci palindromes of length n differing in a single position. This is only possible if $n = 2k + 1$ and

$$\begin{aligned} u &= x000\beta(x), \\ v &= x010\beta(x) \end{aligned}$$

or vice versa, for some $x \in V(\Gamma_{k-1})$. Hence

$$o_E(\Gamma_n, 1) = \begin{cases} |V(\Gamma_{k-1})| = F_{k+1}, & \text{if } n = 2k + 1, \\ 0, & \text{if } n = 2k, \end{cases} \quad (9)$$

which can be written as $\frac{1-(-1)^n}{2} F_{\lfloor \frac{n+1}{2} \rfloor}$. The remaining two equalities now follow from (5) which in this case transforms into

$$\begin{aligned} o_E(\Gamma_n, 1) + 2o_E(\Gamma_n, 2) &= \frac{1}{5} (nF_{n+1} + 2(n+1)F_n), \\ o_E(\Gamma_n, 1) + o_E(\Gamma_n, 2) &= o_E(\Gamma_n). \quad \square \end{aligned}$$

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14
$ E(\Gamma_n) $	1	2	5	10	20	38	71	130	235	420	744	1308	2285	3970
$o_E(\Gamma_n)$	1	1	3	5	11	19	37	65	120	210	376	654	1149	1985
$o_E(\Gamma_n, 1)$	1	0	1	0	2	0	3	0	5	0	8	0	13	0
$o_E(\Gamma_n, 2)$	0	1	2	5	9	19	34	65	115	210	368	654	1136	1985

Table 2: The numbers of edges, all orbits, orbits of size 1, and orbits of size 2 in $E(\Gamma_n)$ for $n \leq 14$

5 Orbits of Lucas cubes

Since Lucas cubes can be viewed as a symmetrization of Fibonacci cubes, the former should possess larger automorphism groups than the latter. Indeed, as shown in [1], $\text{Aut}(\Lambda_n)$ is generated by the cyclic shift and reversal maps $\alpha, \beta : V(\Lambda_n) \rightarrow V(\Lambda_n)$. Hence $\text{Aut}(\Lambda_n) = \{\text{id}, \alpha, \alpha^2, \dots, \alpha^{n-1}, \beta, \alpha\beta, \alpha^2\beta, \dots, \alpha^{n-1}\beta\} \simeq D_n$ when $n \geq 3$.

5.1 Vertex orbits

Theorem 5.1 *For all $n \in \mathbb{N}$,*

$$o_V(\Lambda_n) = \frac{1}{2} \left(\frac{1}{n} \sum_{d|n} \varphi\left(\frac{n}{d}\right) L_d + \sum_{0 \leq a \leq \lfloor \frac{n}{2} \rfloor} \binom{\lfloor \frac{n}{2} \rfloor - \lfloor \frac{a+1}{2} \rfloor}{\lfloor \frac{a}{2} \rfloor} \right).$$

Proof. For $n \leq 2$ this can be checked directly. Now let $n \geq 3$. Each $u \in V(\Lambda_n)$ can be constructed uniquely by concatenating a copies of $01 \in V(\Lambda_2)$ and $n - 2a$ copies of $0 \in V(\Lambda_1)$ for some a between 0 and $n/2$. Together this gives $n - a$ objects of two ‘‘colors’’ (01 and 0) arranged in a circle. Hence our problem is equivalent to determining the total number of 2-colorings of $n - a$ objects under the action of D_{n-a} , with a objects of color 01 and $n - 2a$ objects of color 0 , for $0 \leq a \leq n/2$. From the well-known Redfield-Pólya Theorem we have

$$o_V(\Lambda_n) = \sum_{a=0}^{\lfloor \frac{n}{2} \rfloor} [x_1^a x_2^{n-2a}] Z_{D_{n-a}}(x_1 + x_2, x_1^2 + x_2^2, \dots, x_1^n + x_2^n) \quad (10)$$

where

$$Z_{D_n}(y_1, y_2, \dots, y_n) = \frac{1}{2n} \sum_{d|n} \varphi(d) y_d^{n/d} + \frac{1}{4} \begin{cases} 2y_1 y_2^{(n-1)/2}, & n \text{ odd}, \\ y_1^2 y_2^{(n-2)/2} + y_2^{n/2}, & n \text{ even}, \end{cases} \quad (11)$$

is the cycle index polynomial of the natural action of D_n , and $[x_1^i x_2^j] p(x_1, x_2)$ denotes the coefficient of $x_1^i x_2^j$ in the polynomial $p(x_1, x_2)$. From (10) and (11) it follows by several applications of the Binomial Theorem that

$$o_V(\Lambda_n) = \frac{1}{2} \left(c(n) + \sum_{0 \leq a \leq \lfloor \frac{n}{2} \rfloor} \binom{\lfloor \frac{n}{2} \rfloor - \lfloor \frac{a+1}{2} \rfloor}{\lfloor \frac{a}{2} \rfloor} \right)$$

where

$$c(n) = \sum_{a=0}^{\lfloor \frac{n}{2} \rfloor} \frac{1}{n-a} \sum_{d|\gcd(n,a)} \varphi(d) \binom{\frac{n-a}{d}}{\frac{a}{d}}. \quad (12)$$

By changing the order of summation on the right side of (12), writing $a = kd$, and replacing d by n/d , we obtain

$$c(n) = \sum_{d|n} \varphi(d) \sum_{k=0}^{\lfloor \frac{n}{2d} \rfloor} \frac{1}{n - kd} \binom{\frac{n}{d} - k}{k} = \frac{1}{n} \sum_{d|n} \varphi\left(\frac{n}{d}\right) \sum_{k=0}^{\lfloor \frac{d}{2} \rfloor} \frac{d}{d - k} \binom{d - k}{k}.$$

From Proposition 2.1 it now follows that

$$c(n) = \frac{1}{n} \sum_{d|n} \varphi\left(\frac{n}{d}\right) L_d,$$

proving the claim. \square

We remark that there is a natural bijection between the set of Lucas strings of length n and the set of circular strings of n black and white beads with no two black beads adjacent. Hence $o_V(\Lambda_n)$ counts black-and-white bracelets with no consecutive black beads. These numbers are known (cf. [18, sequence A129526]), as are the numbers $c(n)$ which count black-and-white necklaces with no consecutive black beads (cf. [18, sequence A000358]).

Now we determine the sizes of orbits. They must divide $2n$, but the following result shows that not all divisors of $2n$ appear as orbit sizes.

Theorem 5.2 *For all $n \in \mathbb{N}$,*

$$\{|X|; X \in \mathcal{O}_V(\Lambda_n)\} = \{k \geq 1; k | n\} \cup \{k \geq 18; k | 2n\}.$$

To prove the theorem, we first show:

Lemma 5.3 *$V(\Lambda_n)$ contains an asymmetric string if and only if $n \geq 9$.*

Proof. To prove that all $u \in V(\Lambda_n)$ with $n < 9$ are symmetric we order strings according to their weight $w(u)$. Since conjugate strings are either all symmetric or all asymmetric, it suffices to consider a single representative from each orbit:

- $w(u) = 0$: $u = 0^n$ is symmetric since $\beta(u) = u$.
- $w(u) = 1$: $u = 10^{n-1}$ is symmetric since $\beta(u) = 0^{n-1}1 = \alpha^{-1}(u)$.
- $w(u) = 2$: $u = 10^a 10^b$ where $a, b \geq 1$ is symmetric since $\beta(u) = 0^b 10^a 1 = \alpha^b(u)$.
- $w(u) = 3$: $u = 10^a 10^b 10^c$ where $a, b, c \geq 1$. If any two of a, b, c are equal, then u is symmetric since
 - if $a = b$: $u = 10^a 10^a 10^c$, $\beta(u) = 0^c 10^a 10^a 1 = \alpha^c(u)$,

- if $a = c$: $u = 10^a 10^b 10^a$, $\beta(u) = 0^a 10^b 10^a 1 = \alpha^{-1}(u)$,
- if $b = c$: $u = 10^a 10^b 10^b$, $\beta(u) = 0^b 10^b 10^a 1 = \alpha^{-a-2}(u)$.

Hence in any asymmetric u of this form we have $a + b + c \geq 1 + 2 + 3 = 6$ and $n = a + b + c + 3 \geq 9$.

- $w(u) = 4$: $u = 10^a 10^b 10^c 10^d$ where $a, b, c, d \geq 1$. If $a = b = c = d = 1$ then $u = 10101010$ is symmetric since $\beta(u) = \alpha(u)$. Otherwise $a + b + c + d \geq 5$ and $n = a + b + c + d + 4 \geq 9$.
- $w(u) \geq 5$: any $u \in V(\Lambda_n)$ with $w(u) \geq 5$ contains at least 5 zeros, so in this case $n \geq w(u) + 5 \geq 10$.

Conversely, assume that $n \geq 9$ and let $u = 1010010^{n-6}$. Then $u \in V(\Lambda_n)$, u is obviously primitive, and $\beta(u) = 0^{n-6}100101$. Since $n - 6 \geq 3$, the only $j \in [n - 1]$ such that $\alpha^j(u)$ starts with 0^{n-6} is $n - 6$, but $\alpha^{n-6}(u) = 0^{n-6}101001 \neq \beta(u)$. Hence u is asymmetric. \square

Proof of Theorem 5.2. For $n \in \{0, 1, 2\}$, this can be checked directly. Let $n \geq 3$. Which divisors k of $2n$ appear as orbit sizes in the action of $\text{Aut}(\Lambda_n)$ on $V(\Lambda_n)$? We distinguish two cases:

Case 1: $k \mid n$. If $k = 1$, take $u = 0^n$; then $u \in V(\Lambda_n)$ and $|\bar{u}| = 1$. For $k \geq 2$, take $u = (10^{k-1})^{n/k} \in V(\Lambda_n)$. Since 10^{k-1} is primitive, $\ell(u) = n/k$ and $p(u) = k$, by Proposition 3.2(iv). Since $\beta(u) = (0^{k-1}1)^{n/k} = \alpha^{k-1}(u)$, it follows from Proposition 3.3 that $|\bar{u}| = p(u) = k$. This shows that $\{k \geq 1; k \mid n\} \subseteq \{|\bar{u}|; u \in V(\Lambda_n)\}$.

Case 2: $k \mid 2n$, but $k \nmid n$. Here k is even. Note that it suffices to prove that there is an orbit of size k in $V(\Lambda_n)$ if and only if $k \geq 18$.

Assume first that $u \in V(\Lambda_n)$ is such that $|\bar{u}| = k$. Then $k \in \{p(u), 2p(u)\}$ by Proposition 3.3. Since $p(u) \mid n$ by Proposition 3.2(iii) but $k \nmid n$, we conclude that $k = 2p(u)$. Theorem 3.5 now implies that $\tau(u)$ is asymmetric. By Lemma 5.3, $|\tau(u)| \geq 9$, hence by Proposition 3.2(iv), $p(u) \geq 9$ as well, so $k \geq 18$.

Conversely, assume that $k \geq 18$. By Lemma 5.3, there is an asymmetric $v \in V(\Lambda_{k/2})$. Let $u = v^s$ where $s = 2n/k$. Then $u \in V(\Lambda_n)$ and by Proposition 3.4, $v = \tau(u)$. By Theorem 3.5 and Proposition 3.2(iv), $|\bar{u}| = 2p(u) = 2|v| = k$. \square

In order to determine the number of orbits of size k , we enumerate primitive symmetric and asymmetric Lucas strings. We denote the set of primitive Lucas strings of length n by $V_p(\Lambda_n)$. For $n \in \mathbb{N}$ we define

$$\begin{aligned}
p_n &= |V_p(\Lambda_n)| = |\{u \in V(\Lambda_n); u \text{ primitive}\}|, \\
s_n &= |\{u \in V(\Lambda_n); u \text{ primitive symmetric}\}|, \\
t_n &= \sum_{j=0}^{n-1} \text{fix}_{V_p(\Lambda_n)}(\alpha^j \beta), \\
a_n &= |\{u \in V(\Lambda_n); u \text{ asymmetric}\}|.
\end{aligned}$$

Lemma 5.4 *Let $u \in \Sigma^n$ and $\alpha^j \beta(u) = \alpha^k \beta(u)$ for some $j, k \in [n]_0$ with $j < k$. Then u is not primitive.*

Proof. Since $k - j > 0$, we have by Lemma 3.1(ii),

$$\begin{aligned} \alpha^j \beta(u) = \alpha^k \beta(u) &\implies \alpha^{k-j} \beta(u) = \beta(u) &\implies p(\beta(u)) \leq k - j < n \\ &\implies \beta(u) \text{ not primitive} &\implies u \text{ not primitive.} \quad \square \end{aligned}$$

Corollary 5.5 (i) *The sets $\text{Fix}_{V_p(\Lambda_n)}(\alpha^j \beta)$, for $j \in [n]_0$, are pairwise disjoint.*

(ii) *For all $n \in \mathbb{N}$, $s_n = t_n$.*

Proof.

(i) Assume that $u \in \text{Fix}_{V_p(\Lambda_n)}(\alpha^j \beta) \cap \text{Fix}_{V_p(\Lambda_n)}(\alpha^k \beta)$. Then $\alpha^j \beta(u) = u = \alpha^k \beta(u)$ and u is primitive, hence Lemma 5.4 implies that $j = k$, proving (i).

(ii) Using (i) and Corollary 3.6, we obtain

$$t_n = \sum_{j=0}^{n-1} |\text{Fix}_{V_p(\Lambda_n)}(\alpha^j \beta)| = \left| \bigcup_{j=0}^{n-1} \text{Fix}_{V_p(\Lambda_n)}(\alpha^j \beta) \right| = s_n. \quad \square$$

Lemma 5.6 *Let $u \in \Sigma^n$ and $0 \leq j < n$. Then $\alpha^j \beta(u) = u$ if and only if there are $x \in \Sigma^j$ and $y \in \Sigma^{n-j}$ such that $x = \beta(x)$, $y = \beta(y)$, and $xy = u$.*

Proof. Assuming that $\alpha^j \beta(u) = u$, let $x = u_1 \cdots u_j$ and $y = u_{j+1} \cdots u_n$. Then $xy = u = \alpha^j \beta(u) = \alpha^j \beta(xy) = \alpha^j(\beta(y)\beta(x)) = \beta(x)\beta(y)$, hence $\beta(x) = x$ and $\beta(y) = y$.

Conversely, assuming that $x \in \Sigma^j$ and $y \in \Sigma^{n-j}$ are such that $x = \beta(x)$, $y = \beta(y)$, and $xy = u$, we have $\alpha^j \beta(u) = \alpha^j \beta(xy) = \alpha^j(\beta(y)\beta(x)) = \beta(x)\beta(y) = xy = u$. \square

In the next theorem, we express the numbers of primitive, primitive symmetric, and asymmetric Lucas strings of length n by means of Möbius function.

Theorem 5.7 *For all $n \in \mathbb{N}$,*

$$(i) \quad p_n = \sum_{d|n} \mu\left(\frac{n}{d}\right) L_d,$$

$$(ii) \quad s_n = n \sum_{d|n} \mu\left(\frac{n}{d}\right) F_{\lfloor \frac{d}{2} \rfloor + 2},$$

$$(iii) \quad a_n = \sum_{d|n} \mu\left(\frac{n}{d}\right) (L_d - n F_{\lfloor \frac{d}{2} \rfloor + 2}).$$

Proof. If u is a Lucas string and $k \in \mathbb{N}$, then $\tau(u)$ and u^k are Lucas strings as well. Hence we can enumerate $u \in V(\Lambda_n)$ by $|\tau(u)|$ which is a divisor of n . This yields

$$L_n = |V(\Lambda_n)| = \sum_{d|n} p_d,$$

from which (i) follows by Möbius inversion.

To derive (ii), we compute t_n and invoke Corollary 5.5(ii). By Lemma 3.1,

$$\begin{aligned} \text{fix}_{V(\Lambda_n)}(\alpha^j \beta) &= |\{u \in V(\Lambda_n); \alpha^j \beta(u) = u\}| \\ &= \left| \bigcup_{d|n} \{u \in V(\Lambda_n); |\tau(u)| = d \wedge \alpha^j \beta(\tau(u)) = \tau(u)\} \right| \\ &= \left| \bigcup_{d|n} \{v \in V(\Lambda_d); v \text{ primitive} \wedge \alpha^j \beta(v) = v\} \right| \\ &= \left| \bigcup_{d|n} \text{Fix}_{V_p(\Lambda_d)}(\alpha^j \beta) \right| = \sum_{d|n} \text{fix}_{V_p(\Lambda_d)}(\alpha^j \beta), \end{aligned}$$

hence by Möbius inversion,

$$\text{fix}_{V_p(\Lambda_n)}(\alpha^j \beta) = \sum_{d|n} \mu\left(\frac{n}{d}\right) \text{fix}_{V(\Lambda_d)}(\alpha^j \beta),$$

and, by definition of t_n ,

$$\begin{aligned} t_n &= \sum_{j=0}^{n-1} \sum_{d|n} \mu\left(\frac{n}{d}\right) \text{fix}_{V(\Lambda_d)}(\alpha^j \beta) = \sum_{d|n} \mu\left(\frac{n}{d}\right) \sum_{j=0}^{n-1} \text{fix}_{V(\Lambda_d)}(\alpha^j \beta) \\ &= \sum_{d|n} \mu\left(\frac{n}{d}\right) \frac{n}{d} \sum_{j=0}^{d-1} \text{fix}_{V(\Lambda_d)}(\alpha^j \beta). \end{aligned} \tag{13}$$

To evaluate $\sum_{j=0}^{d-1} \text{fix}_{V(\Lambda_d)}(\alpha^j \beta)$, we use Lemma 5.6 according to which we need to count strings $xy \in V(\Lambda_d)$ where $x \in \text{Fix}_{V(\Gamma_j)}(\beta)$ and $y \in \text{Fix}_{V(\Gamma_{d-j})}(\beta)$ are Fibonacci palindromes. We distinguish four cases according to the parities of d and j , computing the numbers of such strings in each case by means of Proposition 4.1:

1. $d = 2k$

(a) $j = 2i$

- $x \in \text{Fix}_{V_0(\Gamma_{2i})}(\beta)$, $y \in \text{Fix}_{V_0(\Gamma_{2(k-i)})}(\beta)$: $F_i F_{k-i}$ strings
- $x \in \text{Fix}_{V_0(\Gamma_{2i})}(\beta)$, $y \in \text{Fix}_{V_1(\Gamma_{2(k-i)})}(\beta)$: $F_i F_{k-i-1}$ strings
- $x \in \text{Fix}_{V_1(\Gamma_{2i})}(\beta)$, $y \in \text{Fix}_{V_0(\Gamma_{2(k-i)})}(\beta)$: $F_{i-1} F_{k-i}$ strings

in all: $F_i F_{k-i+1} + F_{i-1} F_{k-i} = \frac{1}{2}(F_i L_{k-i} + L_i F_{k-i})$ strings

(b) $j = 2i + 1$

- $x \in \text{Fix}_{V_0(\Gamma_{2i+1})}(\beta), y \in \text{Fix}_{V_0(\Gamma_{2(k-i-1)+1})}(\beta)$: $F_{i+2}F_{k-i+1}$ strings
- $x \in \text{Fix}_{V_0(\Gamma_{2i+1})}(\beta), y \in \text{Fix}_{V_1(\Gamma_{2(k-i-1)+1})}(\beta)$: $F_{i+2}F_{k-i}$ strings
- $x \in \text{Fix}_{V_1(\Gamma_{2i+1})}(\beta), y \in \text{Fix}_{V_0(\Gamma_{2(k-i-1)+1})}(\beta)$: $F_{i+1}F_{k-i+1}$ strings

in all: $F_{i+2}F_{k-i+2} + F_{i+1}F_{k-i+1} = \frac{1}{2}(F_{i+2}L_{k-i+1} + L_{i+2}F_{k-i+1})$ strings

2. $d = 2k + 1$

(a) $j = 2i$

- $x \in \text{Fix}_{V_0(\Gamma_{2i})}(\beta), y \in \text{Fix}_{V_0(\Gamma_{2(k-i)+1})}(\beta)$: F_iF_{k-i+2} strings
- $x \in \text{Fix}_{V_0(\Gamma_{2i})}(\beta), y \in \text{Fix}_{V_1(\Gamma_{2(k-i)+1})}(\beta)$: F_iF_{k-i+1} strings
- $x \in \text{Fix}_{V_1(\Gamma_{2i})}(\beta), y \in \text{Fix}_{V_0(\Gamma_{2(k-i)+1})}(\beta)$: $F_{i-1}F_{k-i+2}$ strings

in all: $F_iF_{k-i+3} + F_{i-1}F_{k-i+2} = \frac{1}{2}(F_iL_{k-i+2} + L_iF_{k-i+2})$ strings

(b) $j = 2i + 1$

- $x \in \text{Fix}_{V_0(\Gamma_{2i+1})}(\beta), y \in \text{Fix}_{V_0(\Gamma_{2(k-i)})}(\beta)$: $F_{i+2}F_{k-i}$ strings
- $x \in \text{Fix}_{V_0(\Gamma_{2i+1})}(\beta), y \in \text{Fix}_{V_1(\Gamma_{2(k-i)})}(\beta)$: $F_{i+2}F_{k-i-1}$ strings
- $x \in \text{Fix}_{V_1(\Gamma_{2i+1})}(\beta), y \in \text{Fix}_{V_0(\Gamma_{2(k-i)})}(\beta)$: $F_{i+1}F_{k-i}$ strings

in all: $F_{i+2}F_{k-i+1} + F_{i+1}F_{k-i} = \frac{1}{2}(F_{i+2}L_{k-i} + L_{i+2}F_{k-i})$ strings

Again we distinguish two cases according to the parity of d , and split the sum into two according to the parity of the summation index:

1. $d = 2k$:

$$\begin{aligned}
\sum_{j=0}^{d-1} \text{fix}_{V(\Lambda_d)}(\alpha^j \beta) &= \sum_{i=0}^{k-1} \text{fix}_{V(\Lambda_d)}(\alpha^{2i} \beta) + \sum_{i=0}^{k-1} \text{fix}_{V(\Lambda_d)}(\alpha^{2i+1} \beta) \\
&= \frac{1}{2} \left(\sum_{i=0}^{k-1} F_i L_{k-i} + \sum_{i=0}^{k-1} L_i F_{k-i} + \sum_{i=0}^{k-1} F_{i+2} L_{k-i+1} + \sum_{i=0}^{k-1} L_{i+2} F_{k-i+1} \right) \\
&= 2k F_{k+2} = d F_{\lfloor \frac{d}{2} \rfloor + 2},
\end{aligned}$$

by shifting summation indices and applying (3) to each of the four sums.

2. $d = 2k + 1$:

$$\begin{aligned}
\sum_{j=0}^{d-1} \text{fix}_{V(\Lambda_d)}(\alpha^j \beta) &= \sum_{i=0}^k \text{fix}_{V(\Lambda_d)}(\alpha^{2i} \beta) + \sum_{i=0}^{k-1} \text{fix}_{V(\Lambda_d)}(\alpha^{2i+1} \beta) \\
&= \frac{1}{2} \left(\sum_{i=0}^k F_i L_{k-i+2} + \sum_{i=0}^k L_i F_{k-i+2} + \sum_{i=0}^{k-1} F_{i+2} L_{k-i} + \sum_{i=0}^{k-1} L_{i+2} F_{k-i} \right) \\
&= (2k + 1) F_{k+2} = d F_{\lfloor \frac{d}{2} \rfloor + 2},
\end{aligned}$$

by shifting summation indices and applying (3) to each of the four sums.

The final expression is the same in both cases, so by (13) we obtain

$$s_n = t_n = \sum_{d|n} \mu\left(\frac{n}{d}\right) \frac{n}{d} \sum_{j=0}^{d-1} \text{fix}_{V(\Lambda_d)}(\alpha^j \beta) = n \sum_{d|n} \mu\left(\frac{n}{d}\right) F_{\lfloor \frac{d}{2} \rfloor + 2},$$

proving (ii).

Finally, (iii) follows from (i) and (ii) by noting that $a_n = p_n - s_n$. \square

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
L_n	1	3	4	7	11	18	29	47	76	123	199	322	521	843	1364	2207
p_n	1	2	3	4	10	12	28	40	72	110	198	300	520	812	1350	2160
s_n	1	2	3	4	10	12	28	40	54	90	132	180	260	392	450	752
a_n	0	0	0	0	0	0	0	0	18	20	66	120	260	420	900	1408

Table 3: The numbers of all Lucas strings, primitive Lucas strings, symmetric primitive Lucas strings, and asymmetric Lucas strings of length $n \leq 16$

Theorem 5.8 For all $n \in \mathbb{N}$ and $k | 2n$,

$$o_V(\Lambda_n, k) = \begin{cases} \sum_{d|k} \mu\left(\frac{k}{d}\right) F_{\lfloor \frac{d}{2} \rfloor + 2}, & k | n \wedge k \text{ odd} \\ \sum_{d|k} \mu\left(\frac{k}{d}\right) F_{\lfloor \frac{d}{2} \rfloor + 2} + \frac{1}{k} \sum_{d|\frac{k}{2}} \mu\left(\frac{k}{2d}\right) (L_d - \frac{k}{2} F_{\lfloor \frac{d}{2} \rfloor + 2}), & k | n \wedge k \text{ even} \\ \frac{1}{k} \sum_{d|\frac{k}{2}} \mu\left(\frac{k}{2d}\right) (L_d - \frac{k}{2} F_{\lfloor \frac{d}{2} \rfloor + 2}), & k | 2n \wedge k \nmid n \end{cases}$$

Proof. By Theorem 3.5,

$$|\bar{u}| = k \iff (|\tau(u)| = k \wedge \tau(u) \text{ symmetric}) \vee (2|\tau(u)| = k \wedge \tau(u) \text{ asymmetric}).$$

Since $\tau(u)$ is primitive and $|\tau(u)|$ divides n , it follows that

$$|\bar{u}| = k \iff \left(k | n \wedge \exists v \in V(\Lambda_k): (v \text{ symmetric primitive} \wedge u = v^{n/k}) \right) \vee \left(k | 2n \wedge k \text{ even} \wedge \exists v \in V(\Lambda_{k/2}): (v \text{ asymmetric} \wedge u = v^{2n/k}) \right),$$

hence

$$o_V(\Lambda_n, k) = \frac{1}{k} \begin{cases} s_k, & k | n \wedge k \text{ odd}, \\ s_k + a_{k/2}, & k | n \wedge k \text{ even}, \\ a_{k/2}, & k | 2n \wedge k \not| n. \end{cases} \quad (14)$$

Together with Proposition 5.7, this yields the statement of the theorem. \square

At first sight, it seems surprising that the summation formulas in Theorem 5.8 are free of n , therefore we try to explain this phenomenon here. Since both α and β commute with taking powers, the size of the orbit of u equals the size of the orbit of u^k , which means that $|\bar{u}|$ depends only on $|\tau(u)|$. Primitive strings of length n are of two types: symmetric (with orbit size n) and asymmetric (of orbit size $2n$). Hence in order to enumerate orbits of size k in $V(\Lambda_n)$, we need to enumerate primitive symmetric Lucas strings of length k and asymmetric Lucas strings of length $k/2$, with the latter possibility applicable only to even k . Furthermore, we wish to obtain a string of length n as a power of our primitive string, therefore n/k (in the first case) resp. $2n/k$ (in the second case) must be an integer. In summary, if k is even, only the first case applies, hence we obtain s_k such strings. If k is even and divides n , both cases apply and there are $s_k + a_{k/2}$ such strings. If k is even but does not divide n (however it must divide $2n$), only the second case applies, yielding $a_{k/2}$ strings. Finally, since we count orbits rather than individual strings, we divide by the orbit size k and obtain (14).

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
$o_V(\Lambda_n)$	1	2	2	3	3	5	5	8	9	14	16	26	31	49	64	99	133	209
$o_V(\Lambda_n, n)$	1	1	1	1	2	2	4	5	6	9	12	15	20	28	30	47	54	79
$o_V(\Lambda_n, 2n)$	0	0	0	0	0	0	0	0	1	1	3	5	10	15	30	44	78	119

Table 4: The numbers of all orbits, orbits of size n , and orbits of size $2n$ in $V(\Lambda_n)$ for $n \leq 18$

5.2 Edge orbits

We now present a surprising relationship between $o_E(\Lambda_n)$ and $o_V(\Gamma_n)$, namely that the former sequence is just a shift of the latter.

Theorem 5.9 (i) For $n \geq 5$, $o_E(\Lambda_n) = o_V(\Gamma_{n-3})$.

(ii) For all $n \in \mathbb{N}$,

$$o_E(\Lambda_n) = \frac{1}{2} \left(F_{n-1} + F_{\lfloor \frac{n+1+(-1)^n}{2} \rfloor} \right).$$

Proof. To prove (i), assume that $n \geq 5$. Then $\text{Aut}(\Gamma_{n-3}) = \{\text{id}, \beta\}$ and $\text{Aut}(\Lambda_n) = \{\text{id}, \alpha, \dots, \alpha^{n-1}, \beta, \alpha\beta, \dots, \alpha^{n-1}\beta\} \simeq D_n$. Define $s: E(\Lambda_n) \rightarrow V(\Gamma_{n-3})$ as follows (all indices will be taken mod n): For $e = \{u, v\} \in E(\Lambda_n)$, let $i \in [n]$ be such that $u_i \neq v_i$ where $u = u_1u_2 \cdots u_n$, $v = v_1v_2 \cdots v_n$. Then $\{u_i, v_i\} = \{0, 1\}$ and $u_{i-1} = v_{i-1} = u_{i+1} = v_{i+1} = 0$. Now set

$$s(e) = u_{i+2}u_{i+3} \cdots u_{i+n-2}.$$

Note that $s(e)$ may contain u_nu_1 as a substring, but as u is a Lucas string, $u_nu_1 \neq 11$, hence $s(e) \in V(\Gamma_{n-3})$ as desired. Let $f = \{x, y\} \in E(\Lambda_n)$ be such that $\bar{e} = \bar{f}$. We claim that $\overline{s(e)} = \overline{s(f)}$. To prove this, we distinguish two cases:

Case 1: $f = \alpha^j(e) = \{\alpha^j(u), \alpha^j(v)\}$ for some j . Without loss of generality assume that $x = \alpha^j(u)$ and $y = \alpha^j(v)$. By (6), $x_{i+j} \neq y_{i+j}$, hence

$$s(f) = x_{i+j+2}x_{i+j+3} \cdots x_{i+j+n-2} = u_{i+2}u_{i+3} \cdots u_{i+n-2} = s(e).$$

Case 2: $f = \alpha^j\beta(e) = \{\alpha^j\beta(u), \alpha^j\beta(v)\}$ for some j . Without loss of generality assume that $x = \alpha^j\beta(u)$ and $y = \alpha^j\beta(v)$. By (8), $x_{1-i+j} \neq y_{1-i+j}$, hence

$$\begin{aligned} s(f) &= x_{3-i+j}x_{4-i+j} \cdots x_{n-1-i+j} = u_{i-2}u_{i-3} \cdots u_{i-n+2} \\ &= \beta(u_{i-n+2} \cdots u_{i-3}u_{i-2}) = \beta(u_{i+2} \cdots u_{i+n-3}u_{i+n-2}) = \beta(s(e)). \end{aligned}$$

In either case, $\overline{s(e)} = \overline{s(f)}$ as claimed. This implies that s induces a well-defined mapping $\tilde{s}: \mathcal{O}_E(\Lambda_n) \rightarrow \mathcal{O}_V(\Gamma_{n-3})$ which satisfies $\tilde{s}(\bar{e}) = \overline{s(e)}$ for all $e \in E(\Lambda_n)$.

Assume that $\tilde{s}(\bar{e}) = \tilde{s}(\bar{f})$ for some $e, f \in E(\Lambda_n)$ where $e = \{u, v\}$ and $f = \{x, y\}$. Let $i, j \in [n]$ be such that $u_i \neq v_i$ and $x_j \neq y_j$. Then $\{u_i, v_i\} = \{x_j, y_j\} = \{0, 1\}$ and $u_{i-1} = v_{i-1} = u_{i+1} = v_{i+1} = x_{j-1} = y_{j-1} = x_{j+1} = y_{j+1} = 0$. Without loss of generality assume $u_i = x_j = 1$. Since $\overline{s(e)} = \overline{s(f)}$, we distinguish two cases:

Case 1: If $s(e) = s(f)$ then $u_{i+2}u_{i+3} \cdots u_{i+n-2} = x_{j+2}x_{j+3} \cdots x_{j+n-2}$, hence

$$\begin{aligned} \alpha^{2-i}(u) &= u_{i-1}u_iu_{i+1}u_{i+2} \cdots u_{i+n-2} = 010u_{i+2} \cdots u_{i+n-2} \\ &= x_{j-1}x_jx_{j+1}x_{j+2}x_{j+3} \cdots x_{j+n-2} = \alpha^{2-j}(x), \end{aligned}$$

and similarly, $\alpha^{2-i}(v) = \alpha^{2-j}(y)$. Consequently, $\alpha^{2-i}(e) = \alpha^{2-j}(f)$.

Case 2: If $s(e) = \beta s(f)$ then $u_{i+2}u_{i+3} \cdots u_{i+n-2} = x_{j+n-2} \cdots x_{j+3}x_{j+2}$, hence

$$\begin{aligned} \alpha^{2-i}(u) &= u_{i-1}u_iu_{i+1}u_{i+2} \cdots u_{i+n-2} = 010u_{i+2} \cdots u_{i+n-2} \\ &= x_{j+1}x_jx_{j-1}x_{j+n-2} \cdots x_{j+3}x_{j+2} \\ &= \beta(x_{j+2}x_{j+3} \cdots x_{j+n-2}x_{j-1}x_jx_{j+1}) = \beta\alpha^{-j-1}(x) \end{aligned}$$

by (6), and similarly, $\alpha^{2-i}(v) = \beta\alpha^{-j-1}(y)$. Consequently, $\alpha^{2-i}(e) = \beta\alpha^{-j-1}(f)$.

In either case, $\tilde{s}(\bar{e}) = \tilde{s}(\bar{f})$ implies that $\bar{e} = \bar{f}$, hence \tilde{s} is injective.

Now let $u \in V(\Gamma_{n-3})$. Then $e = \{010u, 000u\} \in E(\Lambda_n)$ and $s(e) = u$, implying that s and \tilde{s} are surjective. Thus \tilde{s} is bijective and $o_E(\Lambda_n) = o_V(\Gamma_{n-3})$.

Finally, (ii) follows from (i) and Theorem 4.2 for $n \geq 3$, and can be verified directly for $n \leq 2$. \square

Theorem 5.10 For all $n \in \mathbb{N}$,

$$\{|X|; X \in \mathcal{O}_E(\Lambda_n)\} \subseteq \{n, 2n\},$$

with equality if and only if $n \geq 5$.

Proof. For $n \leq 4$ this can be checked directly. Now assume that $n \geq 5$ and let $e = \{u, v\} \in E(\Lambda_n)$. Then by [15, Corollary 2 (ii)], at least one of the strings u and v is primitive, say u . By Proposition 3.3, $|\bar{u}| = p(u)$ or $|\bar{u}| = 2p(u)$. Moreover, because u is primitive, $p(u) = |u| = n$. We conclude that $|\bar{u}| \in \{n, 2n\}$.

Denote by S_u resp. S_e the stabilizer of u resp. e under the action of D_n on $V(\Lambda_n)$ resp. $E(\Lambda_n)$. Let $g \in S_e$. Then $\{g(u), g(v)\} = \{u, v\}$. Since $w(u) \neq w(v)$ and g preserves weight, it follows that $g(u) = u$, so $g \in S_u$. Consequently $S_e \subseteq S_u$, hence $S_e \leq S_u$ and $|S_e|$ divides $|S_u|$. From $|\bar{e}| |S_e| = |D_n| = |\bar{u}| |S_u|$ it now follows that $|\bar{u}|$ divides $|\bar{e}|$, so n divides $|\bar{e}|$ as well. But $|\bar{e}|$ divides $2n$, hence $|\bar{e}| \in \{n, 2n\}$.

To see that both n and $2n$ indeed appear as orbit sizes, consider the edges $e = \{0^n, 10^{n-1}\}$ and $f = \{10^{n-1}, 1010^{n-3}\}$. Then it is straightforward to check that $|\bar{e}| = n$ and $|\bar{f}| = 2n$. \square

Corollary 5.11 For all $n \in \mathbb{N}$,

$$\begin{aligned} o_E(\Lambda_n, n) &= F_{\lfloor \frac{n+1+(-1)^n}{2} \rfloor}, \\ o_E(\Lambda_n, 2n) &= \frac{1}{2} \left(F_{n-1} - F_{\lfloor \frac{n+1+(-1)^n}{2} \rfloor} \right). \end{aligned}$$

Proof. This follows from (5) which in this case transforms into

$$\begin{aligned} n o_E(\Lambda_n, n) + 2n o_E(\Lambda_n, 2n) &= n F_{n-1}, \\ o_E(\Lambda_n, n) + o_E(\Lambda_n, 2n) &= \frac{1}{2} \left(F_{n-1} + F_{\lfloor \frac{n+1+(-1)^n}{2} \rfloor} \right). \end{aligned} \quad \square$$

Acknowledgements

The research of Ali Reza Ashrafi and Khadijeh Fathalikhani was partially supported by the University of Kashan under the grant no 364988/9. The research of Sandi Klavžar and Marko Petkovšek was partially supported by the Ministry of Science of Slovenia under the grants P1-0297 and P1-0294.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$o_E(\Lambda_n)$	0	1	1	2	2	4	5	9	12	21	30	51	76	127	195	322
$o_E(\Lambda_n, n)$	0	1	1	2	1	3	2	5	3	8	5	13	8	21	13	34
$o_E(\Lambda_n, 2n)$	0	0	0	0	1	1	3	4	9	13	25	38	68	106	182	288

Table 5: The numbers of all orbits, orbits of size n , and orbits of size $2n$ in $E(\Lambda_n)$ for $n \leq 16$

References

- [1] A. Castro, S. Klavžar, M. Mollard, Y. Rho, On the domination number and the 2-packing number of Fibonacci cubes and Lucas cubes, *Comput. Math. Appl.* 61 (2011) 2655–2660.
- [2] A. Castro, M. Mollard, The eccentricity sequences of Fibonacci and Lucas cubes, *Discrete Math.* 312 (2012) 1025–1037.
- [3] S. Cabello, D. Eppstein, S. Klavžar, The Fibonacci dimension of a graph, *Electron. J. Combin.* 18 (2011) Paper 55, 23 pp.
- [4] P. Codara, O. M. D’Antona, Independent subsets of powers of paths, and Fibonacci cubes, *Electron. Notes Discrete Math.* 40 (2013) 65–69.
- [5] W.-J. Hsu, Fibonacci cubes—a new interconnection topology, *IEEE Trans. Parallel Distrib. Syst.* 4 (1993) 3–12.
- [6] S. Klavžar, Structure of Fibonacci cubes: a survey, *J. Comb. Optim.* 25 (2013) 505–522.
- [7] S. Klavžar, M. Mollard, Wiener index and Hosoya polynomial of Fibonacci and Lucas cubes, *MATCH Commun. Math. Comput. Chem.* 68 (2012) 311–324.
- [8] S. Klavžar, M. Mollard, Asymptotic properties of Fibonacci cubes and Lucas cubes, *Ann. Comb.*, in press, DOI 10.1007/s00026-014-0233-x.
- [9] S. Klavžar, M. Mollard, M. Petkovšek, The degree sequence of Fibonacci and Lucas cubes, *Discrete Math.* 311 (2011) 1310–1322.
- [10] S. Klavžar, P. Žigert, Fibonacci cubes are the resonance graphs of fibonaccenes, *Fibonacci Quart.* 43 (2005) 269–276.
- [11] R. C. Lyndon, M. P. Schützenberger, The equation $a^M = b^N c^P$ in a free group, *Michigan Math. J.* 9 (1962) 289–298.

- [12] M. Mollard, Maximal hypercubes in Fibonacci and Lucas cubes, *Discrete Appl. Math.* 160 (2012) 2479–2483.
- [13] E. Munarini, C. Perelli Cippo, N. Zagaglia Salvi, On the Lucas cubes, *Fibonacci Quart.* 39 (2001) 12–21.
- [14] W. E. Patten, S. W. Golomb, Problem E1470, *Amer. Math. Monthly* 69 (1962) 61–62.
- [15] G. Păun, N. Santean, G. Thierrin, S. Yu, On the robustness of primitive words, *Discrete Appl. Math.* 117 (2002) 239–252.
- [16] D. A. Pike, Y. Zou, The domination number of Fibonacci cubes, *J. Combin. Math. Combin. Comput.* 80 (2012) 433–444.
- [17] M. Ramras, Congestion-free routing of linear permutations on Fibonacci and Lucas cubes, *Australas. J. Combin.* 60 (2014) 1–10.
- [18] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, published electronically at <http://oeis.org>, 2014.
- [19] A. Taranenko, A new characterization and a recognition algorithm of Lucas cubes, *Discrete Math. Theor. Comput. Sci.* 15 (2013) 31–39.
- [20] A. Vesel, Fibonacci dimension of the resonance graphs of catacondensed benzenoid graphs, *Discrete Appl. Math.* 161 (2013) 2158–2168.
- [21] A. Vesel, Linear recognition and embedding of Fibonacci cubes, to appear in *Algorithmica*. DOI 10.1007/s00453-013-9839-3.
- [22] H. Zhang, L. Ou, H. Yao, Fibonacci-like cubes as Z-transformation graphs, *Discrete Math.* 309 (2009) 1284–1293.
- [23] P. Žigert, M. Berlič, Lucas cubes and resonance graphs of cyclic polyphenantrenes, *MATCH Commun. Math. Comput. Chem.* 68 (2012) 79–90.
- [24] P. Žigert Pleteršek, M. Berlič, The structure of Lucas cubes and maximal resonant sets of cyclic fibonacenes, *MATCH Commun. Math. Comput. Chem.* 69 (2013) 707–720.
- [25] P. Žigert Pleteršek, M. Berlič, Resonance graphs of armchair nanotubes cyclic polypyrenes and amalgams of Lucas cubes, *MATCH Commun. Math. Comput. Chem.* 70 (2013) 533–543.