

Bounds for variables with few occurrences in conjunctive normal forms

Oliver Kullmann^a, Xishun Zhao^{b,1}

^aComputer Science Department, Swansea University, UK

^bInstitute of Logic and Cognition, Sun Yat-sen University, Guangzhou, P.R.C.

Abstract

We investigate connections between SAT (the propositional satisfiability problem) and combinatorics, around the minimum degree of variables in various forms of redundancy-free boolean conjunctive normal forms (clause-sets). Extensive introductions, overviews, conclusions, examples and open problems are provided.

Let $\mu\text{vd}(F) \in \mathbb{N}$ for a clause-set F denote the *minimum variable-degree*, the minimal number of occurrences of a variable. A central result is the upper bound $\sigma(F) + 1 \leq \mu\text{vd}(F) \leq \text{nM}(\sigma(F)) \leq \sigma(F) + 1 + \log_2(\sigma(F))$ for *lean clause-sets* $F \in \mathcal{LEAN}$ in dependency on the *surplus* $\sigma(F) \in \mathbb{N}$. Lean clause-sets are defined by having no non-trivial autarkies (partial assignments satisfying some clauses and not touching the other clauses), and generalise *minimally unsatisfiable clause-sets* (which become satisfiable iff any single clause is removed), i.e., $\mathcal{LEAN} \supset \mathcal{MU}$. For the surplus we have $\sigma(F) \leq \delta(F) = c(F) - n(F)$, using the deficiency $\delta(F)$ of clause-sets, the difference between the number $c(F)$ of clauses and the number $n(F)$ of variables. And $\text{nM}(k) \in \mathbb{N}$ is the k -th “non-Mersenne” number, skipping in the sequence of natural numbers all numbers of the form $2^n - 1$. As an application of the upper bound we obtain, that clause-sets F violating $\mu\text{vd}(F) \leq \text{nM}(\sigma(F))$ must have a non-trivial autarky, and thus clauses can be removed satisfiability-equivalently. We obtain a polynomial time autarky reduction (removing the clauses), but where it is open whether the autarky itself can be found in polynomial time.

We show that the upper bound on μvd is sharp, indeed already on $\mathcal{MU} \subset \mathcal{VMU} \subset \mathcal{LEAN}$, the class of *variable-minimally unsatisfiable clause-sets* (these become satisfiable iff all clauses containing any single variable are removed). That is, $\mu\text{vd}(\mathcal{VMU}_{\delta=k}) = \text{nM}(k)$ for all deficiencies $k \in \mathbb{N}$, where $\mu\text{vd}(\mathcal{VMU}_{\delta=k})$ is the maximum of $\mu\text{vd}(F)$ over $F \in \mathcal{VMU}$ with $\delta(F) = k$. The determination of $\mu\text{vd}(\mathcal{MU}_{\delta=k}) =: \text{VDM}(k)$ seems to be a much more involved question. We show that for $k \leq 5$ we have $\text{VDM}(k) = \text{nM}(k)$, but for $k = 6$ we have $\text{VDM}(k) = \text{nM}(k) - 1$. Moreover this correction by -1 causes further corrections by -1 for infinitely many other deficiencies, resulting in the upper-bound function $\text{nM}_1 : \mathbb{N} \rightarrow \mathbb{N}$, an instance of a *generalised non-Mersenne function*, found by a novel recursion scheme.

We also show various auxiliary results, especially concerning \mathcal{VMU} .

Keywords: conjunctive normal form, deficiency, minimally unsatisfiable, variable minimal unsatisfiable, saturated minimal unsatisfiable, hitting clause-set, disjoint tautology, orthogonal tautology, lean clause-set, autarky, surplus, matching-lean, L-matrix, minimally sign-central matrix, fixed-parameter tractable, polynomial time, SAT decision, non-Mersenne number, minimum variable degree, variable occurrences, singular DP-reduction (variable elimination), subsumption resolution

URL: <http://cs.swan.ac.uk/~csoliver> (Oliver Kullmann),

<http://logic.sysu.edu.cn/faculty/zhaoxishun/en/> (Xishun Zhao)

¹Supported by Grants NSFC 61272059, NSSFC 13&ZD186 and MOE 11JJD7200020.

Contents

1	Introduction	3
1.1	Deficiency as the main structural parameter	4
1.2	Refining deficiency by surplus	5
1.3	Some basic intuitions about the upper bound nM	6
1.4	Related work on \mathcal{MU}	7
1.4.1	MUS	8
1.4.2	Tovey's problem (uniform clause-sets)	8
1.5	Autarkies	10
1.6	Connections to combinatorics	11
1.6.1	Hypergraph colouring	12
1.6.2	Hypergraph transversals	13
1.6.3	Autarkies for hypergraphs	14
1.6.4	Qualitative matrix analysis (QMA)	16
1.6.5	Biclique partitions of (multi-)graphs, and algebraic graph theory	17
1.7	Overview on results	19
2	Preliminaries	21
2.1	Clause-sets	21
2.2	Semantics	23
2.3	Resolution	24
2.4	Multi-clause-sets and restrictions	24
2.5	Degrees	26
2.6	Autarkies	27
2.7	The surplus	28
3	Minimally unsatisfiable clause-sets	29
3.1	\mathcal{MU} and subclasses	29
3.2	Saturation	30
3.3	Splitting	33
4	Variable-minimal unsatisfiability	34
5	Eliminating and creating singularity	36
5.1	Singular DP-reduction	36
5.2	Singular DP-extensions	38
5.3	Unit clauses	39
6	Full subsumption resolution / extension	42
6.1	Basic definitions	43
6.2	Extensions to full clause-sets	45
7	Non-Mersenne numbers	47
7.1	Basic properties	48
7.2	Characterising the jumps	50
7.3	Applications	54
8	The upper bound for VDM	56

9	The min-var-degree upper bound for \mathcal{LEAN}	58
9.1	Clause-sets with extremal surplus	58
9.2	The generalised upper bound	61
9.3	Sharpness of the bound for \mathcal{VMU}	62
10	Algorithmic implications	63
10.1	Autarky reduction	63
10.2	On finding the autarky	65
10.3	Final remarks on the surplus	66
11	Matching lean clause-sets	68
12	Lower bounds for VDM	69
12.1	Some precise values for VDM	70
12.2	On the number of full clauses	71
13	A method for improving upper bounds for VDM	72
13.1	Analysing splitting-situations	72
13.2	Recursion on potential degree-pairs	75
14	Strengthening the upper bound for VDM	77
14.1	Deficiencies 1, . . . , 6	77
14.2	Sharpening the bound	79
15	Conclusion and open problems	81
15.1	Conjectures and questions	81
15.2	Improved upper bounds for VDM	82
15.3	Determining FCM	83
15.4	Generalisation to non-boolean clause-sets	83
15.5	Classification of \mathcal{MU}	85

1. Introduction

In this work we aim at bringing together some aspects of combinatorics with the developing theory of SAT. We concentrate on degree considerations in “clause-sets” (conjunctive normal forms as set-systems), which can be considered as generalised hypergraphs, namely hypergraphs with “polarities”. The general goal is to develop an *understanding* of propositional (un)satisfiability, which corresponds for hypergraphs to an understanding of (non-)2-colourability.

SAT, the prototypical NP-complete problem ([1]), took a strong development in the past two decades also regarding (industrial) applications (see the handbook [2] for a recent overview). It is often mainly considered as belonging to complexity theory, algorithms and heuristics (with [3, 4] the basic papers here), and finally implementations and experimentation (“SAT solvers”). “Understanding” SAT in a precise sense is considered to be impossible, and only various investigations on random and approximation structures (including “islands of tractability”) in general are deemed fruitful. We want to challenge this view, starting to build a new bridge, towards an *understanding of unsatisfiability*. We note here that understanding unsatisfiability seems easier than to understand satisfiability, since unsatisfiability means a form of completion, *all* assignments have been excluded as potential satisfying assignments (“models”), while satisfiability means a lack of such completion. More precisely, we aim at understanding *minimal unsatisfiability*, the building blocks of unsatisfiability — similar to critical colourability, here removal of any clause renders the clause-set satisfiable.

A fundamental question, the subject of this study, is the existence of “simple” variables in clause-sets. “Simple” here means a variable occurring not very often. A major use of the existence of such variables is in inductive proofs of properties of minimally unsatisfiable clause-sets, using splitting on a variable to reduce n , the number of variables, to $n - 1$: here it is vital that we have control over the changes imposed by the substitution, and so we want to split on a variable occurring as few times as possible. “Splitting” of a clause-set F on variable v means the consideration of the two clause-sets $\langle v \rightarrow 0 \rangle * F$, $\langle v \rightarrow 1 \rangle * F$, that is, instantiating variable v by both truth values 0, 1. A feature of clause-sets is the closure under splitting, and splitting is a major tool for investigations into minimal unsatisfiability.

In the remainder of the introduction we give an overview on the results of this report and their context. In Subsection 1.1 we introduce minimal unsatisfiability and the main complexity parameter, the deficiency, and discuss the first main result, the upper bound for the minimum variable degree for minimally unsatisfiable clause-sets in the deficiency. In Subsection 1.2 we discuss the generalisation to “lean” clause-sets and their “surplus”. Back to the situation for minimal unsatisfiability, in Subsection 1.3 some basic intuition is developed regarding the number-theoretical upper-bound function. Related work on minimal unsatisfiability is reviewed in Subsection 1.4. The generalisation to lean clause-sets is based on “autarkies”, which are reviewed in Subsection 1.5. In Subsection 1.6 then the connections to combinatorics are discussed: hypergraph colouring, hypergraph transversals, combinatorial matrix theory, and biclique partitions of (multi)graphs, where always SAT is treated in disguise, and autarkies play an important role. The introduction is concluded by going through all main results in Subsection 1.7.

According to the goal of bringing different communities together, we provide and explain much of the relevant background, so that this report is mostly self-contained, and the results cited from the literature can be treated as black-boxes.

1.1. Deficiency as the main structural parameter

The definition of the class \mathcal{CLS} of “clause-sets”, and of the class $\mathcal{MU} \subset \mathcal{CLS}$ of “minimally unsatisfiable clause-sets”, can be quickly (and precisely) given as follows, using (just) natural numbers as “variables”:

A “literal” x is an element of $\mathbb{Z} \setminus \{0\}$. A “clause” C is a finite set of literals, such that there is no $x \in C$ with $-x \in C$. Using $-L := \{-x : x \in L\}$ for sets L of literals, the “clash-freeness” condition for C becomes $C \cap -C = \emptyset$. A “clause-set” F is a finite set of clauses, the set of all clause-sets is denoted by \mathcal{CLS} . The most basic measurements for $F \in \mathcal{CLS}$ are:

- the number $c(F) := |F| \in \mathbb{N}_0$ of clauses of F ;
- the number $n(F) := |\text{var}(F)| \in \mathbb{N}_0$ of variables of F , where $\text{var}(F)$ is the set of $v \in \mathbb{N}$ (variables as positive integers) with $\{v, -v\} \cap \bigcup F \neq \emptyset$;
- The “deficiency” $\delta(F) := c(F) - n(F) \in \mathbb{Z}$. This parameter is only informative when certain (weak) assumptions are made for F , and for general F the “maximal deficiency” $\delta^*(F) := \max_{F' \subseteq F} \delta(F') \in \mathbb{N}_0$ is to be used.

A clause-set F is “satisfiable” if there exists a partial assignment φ , which here in this introduction is just a clause, such that $\varphi \cap D \neq \emptyset$ for all $D \in F$.² The set of all satisfiable clause-sets is $\mathcal{SAT} \subset \mathcal{CLS}$, the set of all unsatisfiable clause-sets is $\mathcal{USAT} := \mathcal{CLS} \setminus \mathcal{SAT}$. Finally $\mathcal{MU} \subset \mathcal{USAT}$ is the set of $F \in \mathcal{USAT}$ such that for

²The clause φ is the set of satisfied literals of the corresponding “partial assignment”. This definition of “satisfying assignments”, via clauses intersecting every clause of F , generalises transversals of hypergraphs, by taking complementation into account (φ does not contain clashes).

all $C \in F$ we have $F \setminus \{C\} \in \mathcal{SAT}$. The background for the investigations of this report is the enterprise of classifying $F \in \mathcal{MU}$ in dependency on $\delta(F)$. The basic facts are $\delta^*(F) = \delta(F)$ (as will be discussed in Subsection 1.5), and the well-known $\delta(F) \geq 1$, as first shown in [5]. For $\delta(F) = 1$ the structure is completely known ([5, 6, 7]; see Example 3.2), for $\delta(F) = 2$ the structure after reduction of singular variables (occurring in one sign only once) is known ([8]; see Example 3.3), while for $\delta(F) \in \{3, 4\}$ only basic cases have been classified ([9]).

The starting point of our investigation is [7, Lemma C.2], where it is shown that $F \in \mathcal{MU}$ with $n(F) > 0$ must have a variable $v \in \text{var}(F)$ with at most $\delta(F)$ positive and at most $\delta(F)$ negative occurrences; we write this as $\text{ld}_F(v) \leq \delta(F)$ and $\text{ld}_F(-v) \leq \delta(F)$, using the notion of *literal degrees* (the number of occurrences of the literal), where for a literal x its degree is

$$\text{ld}_F(x) := |\{C \in F : x \in C\}| \in \mathbb{N}_0.$$

Thus we have $\text{vd}_F(v) \leq 2\delta(F)$, using the *variable degree*

$$\text{vd}_F(v) := \text{ld}_F(v) + \text{ld}_F(-v) \in \mathbb{N}_0.$$

Using the *minimum variable degree* (min-var-degree)

$$\mu\text{vd}(F) := \min_{v \in \text{var}(F)} \text{vd}_F(v) \in \mathbb{N}$$

of F with $n(F) > 0$, the upper bound becomes $\mu\text{vd}(F) \leq 2\delta(F)$. A main theme of this report is the consideration of $\mu\text{vd}(\mathcal{MU}_{\delta=k}) \in \mathbb{N}$ for $k \in \mathbb{N}$, the maximum of $\mu\text{vd}(F)$ for $F \in \mathcal{MU}$ with $\delta(F) = k$. The upper bound now becomes $\text{VDM}(k) := \mu\text{vd}(\mathcal{MU}_{\delta=k}) \leq 2k$. We show a sharper bound on $\mu\text{vd}(F)$, namely we show that the worst-cases $\text{ld}_F(v), \text{ld}_F(-v) \leq \delta(F)$ can not occur at the same time (for a suitable variable), but actually $\text{ld}_F(v) + \text{ld}_F(\bar{v}) - \delta(F)$ only grows logarithmically in $\delta(F)$. The really interesting aspect here is the precise determination of $\text{VDM}(k)$, and we investigate the (elementary) number-theoretic function $\text{nM}(k)$, which yields the upper bound $\text{VDM}(k) \leq \text{nM}(k)$ for all $k \in \mathbb{N}$, where the function $\text{nM} : \mathbb{N} \rightarrow \mathbb{N}$ fulfils $k + \lfloor \log_2(k+1) \rfloor \leq \text{nM}(k) \leq k + 1 + \lfloor \log_2(k) \rfloor$ for $k \in \mathbb{N}$.

1.2. Refining deficiency by surplus

After having settled this basic min-var-degree upper bound for $\mathcal{MU}_{\delta=k}$, we show a sharper bound on $\mu\text{vd}(F)$ for a larger class of clause-sets F :

- The larger class of clause-sets considered is the class \mathcal{LEAN} of *lean clause-sets* (introduced in [10]), which are clause-sets having no non-trivial autarky. For an overview on the theory of minimally unsatisfiable clause-sets and on the theory of autarkies see [11]. $\mathcal{LEAN} \subset \mathcal{CLS}$ is the set of $F \in \mathcal{CLS}$ such that there is no partial assignment φ (a “non-trivial autarky”) with the properties
 - for every clause $D \in F$ with $-\varphi \cap D \neq \emptyset$ we have $\varphi \cap D \neq \emptyset$ (note that this generalises the satisfaction criterion);
 - there exists $v \in \text{var}(F)$ with $\{v, -v\} \cap \varphi \neq \emptyset$.

Note $\mathcal{LEAN} \cap \mathcal{SAT} = \{\top\}$, where $\top := \emptyset \in \mathcal{CLS}$ is the empty clause-set (the standard satisfiable clause-set).

- The deficiency $\delta(F) \in \mathbb{Z}$ is strengthened by the *surplus* $\sigma(F) \in \mathbb{Z}$, defined in case of $n(F) > 0$ as follows.

Consider the bipartite clause-variable graph of F (generalising the incidence graph of a hypergraph), with the clauses $C \in F$ on one side of the bipartition, and the variables $v \in \text{var}(F)$ on the other side, and an edge between

v and C if $\{v, -v\} \cap C \neq \emptyset$. The “expansion” of a set $\emptyset \neq V \subseteq \text{var}(F)$ of variables is $|\Gamma(V)| - |V|$, where $\Gamma(V)$ is the set of neighbours of V (incident clauses), and the surplus then is the minimum expansion, i.e., $\sigma(F) = \min_{\emptyset \neq V \subseteq \text{var}(F)} |\Gamma(V)| - |V|$.

In the terminology of [12, Section 1.3], $\delta^*(F)$ is the deficiency of the bipartite clause-variable graph (with bipartition $(F, \text{var}(F))$), while $\sigma(F)$ is the surplus of the bipartite variable-clause graph (with bipartition $(\text{var}(F), F)$).

Note that by considering $V = \text{var}(F)$ we have $\sigma(F) \leq \delta(F)$, and by considering $V = \{v\}$ for $v \in \text{var}(F)$ we get $\sigma(F) \leq \mu\text{vd}(F) - 1$.

We have $\sigma(F) \geq 1$ for $F \in \mathcal{LEAN}$ with $n(F) > 0$ ([13, Lemma 7.7]), generalising the basic fact $\delta(F) \geq 1$ for $F \in \mathcal{MU}$.

Now a central result of this report (Theorem 9.10) is

$$\mu\text{vd}(F) \leq \text{nM}(\sigma(F))$$

for $F \in \mathcal{LEAN}$ with $n(F) > 0$. As an application we obtain (Theorem 10.2), that via removing satisfiability-equivalently some clauses (via some autarky), we can reduce every (multi-)clause-set F in polynomial time to a (multi-)clause-set F' containing a variable occurring with degree at most $\sigma(F') + 1 + \log_2(\sigma(F'))$. It is an open problem whether such an autarky can be found in polynomial time (for arbitrary clause-sets F); we conjecture (Conjecture 10.3) that this is possible.

We also show sharpness of the upper bound, i.e., $\mu\text{vd}(\mathcal{LEAN}_{\delta=k}) = \text{nM}(k)$ for all $k \in \mathbb{N}$, in Corollary 9.15 (proving Conjecture 23 from the conference version [14]), which indeed holds for every class of clause-sets between \mathcal{VMU} , i.e., “variable-minimally unsatisfiable clause-sets” as introduced in [15], and \mathcal{LEAN} ; the definition of \mathcal{VMU} is as the set of $F \in \mathcal{USAT}$ such that for all $F' \subseteq F$ with $\text{var}(F') \subset \text{var}(F)$ we have $F' \in \mathcal{SAT}$.

We then come back to the special case of minimal unsatisfiability. Here things are much more complicated, and the numbers $\text{VDM}(k)$, the guaranteed minimum variable degrees for minimally unsatisfiable clause-sets of deficiency k , are very interesting quantities. We prove the sharpened bound $\text{VDM}(k) \leq \text{nM}_1(k)$, which improves on $\text{nM}(k)$ for infinitely many k .

1.3. Some basic intuitions about the upper bound nM

As already mentioned, the function $\text{nM} : \mathbb{N} \rightarrow \mathbb{N}$ is strictly increasing with range

$$\text{nM}(\mathbb{N}) = \mathbb{N} \setminus \{2^n - 1 : n \in \mathbb{N}\} = \{\underline{2}, 4, 5, \underline{6}, 8, \dots, \underline{14}, 16, 17, \dots\}.$$

We show $\mu\text{vd}(\mathcal{LEAN}_{\delta=k}) = \text{nM}(k)$ for deficiencies $k \in \mathbb{N}$, that is, every lean clause-set F with $n(F) > 0$ contains a variable $v \in \text{var}(F)$ with $\text{vd}_F(v) \leq \text{nM}(\delta(F))$, and for every deficiency $k \geq 1$ there are lean clause-sets F with $\mu\text{vd}(F) = \text{nM}(\delta(F))$.

The underlined values $2, 6, 14, \dots$, which have the form $2^n - 2$ for $n \in \mathbb{N}$, are the function values at the “jump positions” $1, 4, 11, \dots$, which are of the form $2^n - n - 1$ for $n \geq 2$ (where the function values changes by $+2$, while otherwise it changes by $+1$ for an increment of the argument). This basic structure of nM can be motivated by the following constructions of $F \in \mathcal{MU}$ with “high” min-var-degree. Indeed these considerations only concern the lower bounds, given by appropriate constructions, while the arithmetic nature of the upper bound $\text{nM}(k)$ rests on different considerations, but for the deficiencies considered here, lower and upper bounds are equal, and the lower bounds are easier to understand here.

The basic clause-sets are the A_n for $n \in \mathbb{N}_0$, which consist of all 2^n sets (clauses) of numbers $\pm 1, \dots, \pm n$, using the natural numbers $1, \dots, n$ as variables. So $A_0 = \{\emptyset\}$, $A_1 = \{\{-1\}, \{1\}\}$, $A_2 = \{\{1, 2\}, \{-1, 2\}, \{1, -2\}, \{-1, -2\}\}$ and so on. It is

easy to see that we have $A_n \in \mathcal{MU}$ with $n(A_n) = n$, $c(A_n) = 2^n = \mu\text{vd}(A_n)$, and $\delta(A_n) = 2^n - n$. We will see that the A_n have the largest possible min-var-degree 2^n for given deficiency $2^n - n$, and we also have $\text{nM}(2^n - n) = 2^n$ for $n \in \mathbb{N}$. These deficiencies $k = 2^n - n$ (numerical values are 1, 2, 5, 12, ...) are the positions directly after the jump positions (excluding deficiency $k = 1$ as a special case).

How can we obtain from that more clause-sets in \mathcal{MU} with high min-var-degree? Consider A_3 : we have e.g. $\{1, 2, 3\}, \{1, 2, -3\} \in A_3$; now logically these two clauses are equivalent to the single clause $\{1, 2\}$ (i.e., we have the same satisfying assignments; technically, a “strict full subsumption resolution” is performed), and we obtain $A'_3 := (A_3 \setminus \{\{1, 2, 3\}, \{1, 2, -3\}\}) \cup \{\{1, 2\}\} \in \mathcal{MU}$. Performing this process in general, using $\{1, \dots, n\}, \{1, \dots, n-1, -n\} \in A_n$, yields $A'_n \in \mathcal{MU}$ for $n \geq 2$, with $n(A'_n) = n$, $c(A'_n) = 2^n - 1$, $\delta(A'_n) = 2^n - n - 1$, and $\mu\text{vd}(A'_n) = 2^n - 2$ (the (single) variable with minimum occurrences is n). These deficiencies are precisely the jump positions $2^n - n - 1$, and accordingly we have $\text{nM}(2^n - n - 1) = 2^n - 2$.

Performing the same trick again to A'_3 , we can replace $\{-1, 2, 3\}, \{-1, -2, 3\} \in A'_3$ by $\{-1, 3\}$, obtaining $A''_3 \in \mathcal{MU}$. Again for general $n \geq 3$ we get $A''_n \in \mathcal{MU}$, $n(A''_n) = n$, $c(A''_n) = 2^n - 2$, $\delta(A''_n) = 2^n - n - 2$, and $\mu\text{vd}(A''_n) = 2^n - 3$; note here the crucial difference, that the min-var-degree has only been changed by -1 . The reason is that there are two variables now with minimum occurrences, namely $n-1, n$, where the degree of variable n changed first by -2 , then by -1 , while for variable $n-1$ the degree first changed by -1 , and then by -2 (and for the other variables $1, \dots, n-2$ we have degree changes by $-1, -1$).

Now one might imagine this process of “strict full subsumption resolutions”/continuing until deficiency $2^{n-1} - (n-1) + 1$, always with change of the min-var-degree by -1 , just before the deficiency of the previous A_{n-1} — this would yield the function nM . However the combinatorial reality is more complicated, and as we prove in this report (Section 14), at least we can not get until $2^{n-1} - (n-1) + 1 = 2^{n-1} - n + 2$ for $n \geq 4$ (in effect), that is, at these deficiencies $k = 6, 13, 28, \dots$ we have $\text{VDM}(k) \leq \text{nM}_1(k) = \text{nM}(k) - 1$.³ We will develop the lower bound method outlined above somewhat further in Subsection 12.1. In general the determination of $\text{VDM}(k)$ seems complicated, however the construction of examples showing $\mu\text{vd}(\mathcal{LEAN}_{\delta=k}) \geq \text{nM}(k)$ is rather easy (see Lemma 9.13).

After having developed some intuition on the main results of this report, we turn now to the discussion of the background and context.

1.4. Related work on \mathcal{MU}

A general overview on minimally unsatisfiable clause-sets (also “minimal unsatisfiable clause-sets/formulas”, or “MU”) is [11]; later developments are in [16, 17] (generalisations to non-boolean clause-sets) and in [18, 19] (studying “singular DP-reduction”, the elimination of variables which occur in one sign only once).

Two early papers on the complexity aspects are [20, 21], who introduced the complexity class D^P and showed that the decision “ $F \in \mathcal{MU}$?” with input $F \in \mathcal{CLS}$ is complete for this class. Another important early paper is [5], which showed $\delta(F) \geq 1$ for $F \in \mathcal{MU}$, where the notion of “deficiency” was introduced by [22]. Furthermore [5] showed polytime-decision of the sub-class $\mathcal{SMU}_{\delta=1} \subset \mathcal{MU}_{\delta=1}$ (called “strongly minimal unsatisfiable” there), where $\mathcal{SMU} \subset \mathcal{MU}$ is the set of $F \in \mathcal{USAT}$ such that for all $C \in F$ and all $x \in \mathbb{Z} \setminus \{0\}$ with $\{x, -x\} \cap C = \emptyset$ holds $(F \setminus \{C\}) \cup \{C \cup \{x\}\} \in \mathcal{SAT}$, that is, adding any literal to any clause renders the clause-set satisfiable. We use the terminology “saturated minimally unsatisfiable” introduced in [23], where the important connection to splitting was introduced, and

³We do not apply the method outlined in this Subsection for obtaining lower bounds as far as we can, but only as needed in this report; see the end of Subsection 12.1 for some further remarks.

a simpler proof of $\delta(F) \geq 1$ for $F \in \mathcal{MU}$ was given. Just for this introduction we handle “partial assignments” via clauses φ (containing the satisfied literals; thus $-\varphi$ is the set of falsified literals), so for a literal x the partial assignment $\langle x \rightarrow 0 \rangle$ is given by $\{-x\}$, while $\langle x \rightarrow 1 \rangle$ is given by $\{x\}$. The application of φ to $F \in \mathcal{CLS}$ is defined as

$$\varphi * F := \{C \setminus -\varphi : C \in F \wedge \varphi \cap C = \emptyset\} \in \mathcal{CLS},$$

that is, removing first the satisfied clauses from F , and then the falsified literals from the remaining clauses. Now for $F \in \mathcal{CLS}$ holds $F \in \mathcal{SMU}$ iff for all $x \in \mathbb{Z} \setminus \{0\}$ holds $\langle x \rightarrow 1 \rangle * F \in \mathcal{MU}$ (the “only if”-direction was shown in [23], the “if”-direction in [7]). Due to this property plus the property, that every $F \in \mathcal{MU}$ can be “saturated” by adding literals to clauses, the class \mathcal{SMU} is an important helping class for investigations into \mathcal{MU} via the splitting method, splitting up $F \in \mathcal{MU}$ into $\langle v \rightarrow 0 \rangle * F$ and $\langle v \rightarrow 1 \rangle * F$ for selected variables v .

We have already mentioned the literature concerned with characterising the classes $\mathcal{MU}_{\delta=k}$ (and subclasses) for small deficiencies $k \leq 4$. Less ambitious is the goal of polytime decision of these classes: the problem was raised in [24], and has been solved via two independent approaches in [7] and [25] (indeed establishing polytime SAT decision for inputs $F \in \mathcal{CLS}$ and fixed $\delta^*(F)$), later strengthened in [26] (showing that SAT decision is even fixed-parameter tractable in $\delta^*(F)$); see also [16] for generalisations and simplifications).

1.4.1. MUS

As we have already mentioned, we consider \mathcal{MU} as the “primal” building block for understanding unsatisfiability. In general an unsatisfiable clause-set can contain many minimally unsatisfiable sub-clause-sets, called “MUSs”. The task of enumerating all of them or at least some “good” ones is also of practical importance, to extract more information on the “causes” of unsatisfiability. A recent overview is [27], while a clean approach to enumerate all MUSs, via hypergraph transversals, is in [28] (the earliest appearance of the underlying observation seems [29, Theorem 2]; compare also [17, Subsection 4.3] for generalisations of the fundamental approach). See also [30] for a reflection on various types of such sub-clause-sets, and on the connection to autarky theory (compare Subsection 1.6.3).

1.4.2. Tovey’s problem (uniform clause-sets)

This report appears to be the first systematic study of the problem of *minimum* variable occurrences / degrees in minimally unsatisfiable clause-sets and generalisations, in dependency on the deficiency — asking for the existence of a variable occurring “infrequently” in general, or for extremal examples where all variables occur not infrequently. The “dual” problem is to consider *maximum* variable occurrences / degrees — asking for the existence of a variable occurring frequently in general, or for extremal examples where all variables occur not frequently. More precisely, the *maximum variable degree* is

$$\nu\text{vd}(F) := \max_{v \in \text{var}(F)} \text{vd}_F(v) \in \mathbb{N},$$

for $n(F) > 0$, while for a class $\mathcal{C} \subseteq \mathcal{CLS}$ of clause-sets, the quantity $\nu\text{vd}(\mathcal{C})$ (to be studied) is the minimum of $\nu\text{vd}(F)$ for $F \in \mathcal{C}$.

This problem has been well-studied for *p-uniform* minimally unsatisfiable clause-sets, starting with [31, 32, 33].⁴ We denote by $p\text{-}\mathcal{CLS} \subseteq \mathcal{CLS}$ for $p \in \mathbb{N}_0$ the set of

⁴We remark that typically in the literature the connections to minimally unsatisfiable clause-sets are not emphasised, but it is clear that when considering (uniform) unsatisfiable clause-sets with a maximum variable degree as small as possible, then one can restrict attention to (uniform) minimally unsatisfiable clause-sets (as worst-cases).

all $F \in \mathcal{CLS}$ with $\forall C \in \mathcal{CLS} : |C| \leq p$, while by $\mathcal{UCLS} \subset \mathcal{CLS}$ we denote the set of all uniform clause-sets, i.e., those $F \in \mathcal{CLS}$ such that for $C, D \in F$, $C \neq D$, holds $|C| = |D|$. Finally $p\text{-}\mathcal{UCLS} := p\text{-}\mathcal{CLS} \cap \mathcal{UCLS}$ and $p\text{-}\mathcal{UMU} := p\text{-}\mathcal{UCLS} \cap \mathcal{MU}$. Now the basic fact is

$$\nu\text{vd}(p\text{-}\mathcal{UMU}) \geq p + 1$$

for $p \in \mathbb{N}$ ([31], generalised in [17, Corollary 7.3]). Trivially $\nu\text{vd}(1\text{-}\mathcal{UMU}) = 2$, and easily one sees $\nu\text{vd}(2\text{-}\mathcal{UMU}) = 3$, while by [31] holds $\nu\text{vd}(3\text{-}\mathcal{UMU}) = 4$. As reported in [34], we have $\nu\text{vd}(4\text{-}\mathcal{UMU}) = 5$, and these are all known precise values of $\nu\text{vd}(p\text{-}\mathcal{UMU})$ (where the notation $f(p) := \nu\text{vd}(p\text{-}\mathcal{UMU}) - 1$ was introduced in [33]). In [34] it was observed that extremal examples might be found in $\mathcal{MU}_{\delta=1}$, and this work was recently extended in [35], establishing the asymptotically tight bound $\lim_{p \rightarrow \infty} \frac{2}{e} \frac{2^p}{p} / \nu\text{vd}(p\text{-}\mathcal{UMU}) = 1$ (where indeed $p\text{-}\mathcal{UMU} \cap \mathcal{MU}_{\delta=1}$ is considered).

In our setting, studying the classes $\mathcal{MU}_{\delta=k}$, the max-var-degree is not very relevant, since we have $\nu\text{vd}(\mathcal{MU}_{\delta=1}) = 2$, while $\nu\text{vd}(\mathcal{MU}_{\delta=k}) = 3$ for $k \geq 2$. This can be seen as follows: As already noticed in [31], there is a poly-time transformation from \mathcal{CLS} to the class $\mathcal{CLS}(1, 2) \subset \mathcal{CLS}$, consisting of those $F \in \mathcal{CLS}$ where for every variable $v \in \text{var}(F)$ we have $\text{ld}_F(v) = 1$ and $\text{ld}_F(-v) \leq 2$. Namely if there is a literal x and two clauses $C, D \in F$ with $x \in C \cap D$, then we can introduce a new variable v , replace x in C, D by v , and add the new clause $\{-v, x\}$, obtaining F' . We study such extensions under the name of “singular DP-extension”, but it is also easy to see directly that F' is satisfiable iff F is, that F' is minimally unsatisfiable iff F is, and that $\delta(F') = \delta(F)$. By repeating this transformation, we obtain $t^{1,2} : \mathcal{CLS} \rightarrow \mathcal{CLS}(1, 2)$. So for $F \in \mathcal{MU}$ we get $t^{1,2}(F) \in \mathcal{MU} \cap \mathcal{CLS}(1, 2)$ with $\delta(t(F)) = \delta(F)$. Whence for all $k \in \mathbb{N}$ we have $\nu\text{vd}(\mathcal{MU}_{\delta=k}) \leq 3$. Now trivially $\nu\text{vd}(\mathcal{MU}_{\delta=1}) = 2$ due to $\{\{1\}, \{-1\}\} \in \mathcal{MU}_{\delta=1}$. On the other hand, if for $F \in \mathcal{MU}$ holds $\nu\text{vd}(F) \leq 2$ (thus $\nu\text{vd}(F) = 2$), then via so-called singular DP-reduction this clause-set can be reduced to $\{\perp\}$, whence $F \in \mathcal{MU}_{\delta=1}$ (this is well-known; compare Example 3.2 later).

So for the study of the max-var-degree, the uniformity restriction seems essential. This is similar to many investigations into (colour-)critical hypergraphs (discussed in Subsection 1.6.1 below), where uniformity is a crucial assumption, and the clause-length p is the main parameter. For investigations into the case of uniform (general) clause-sets, where clauses share at most one variable, see [36, 37]. The number of clauses in $F \in p\text{-}\mathcal{UMU}$ has been studied in [38], showing that for $p = 2$ holds $c(F) \leq 4n - 2$, while for $p \geq 3$ there are F with $c(F) = \Omega(n(F)^p)$. Finally, the number of conflicts (clashes) in $F \in p\text{-}\mathcal{UMU}$ is considered in [39], and for a review of the use of the Lovász Local Lemma in this context see [40].

In contrast, for the study of the minimum variable degree as in this report, in dependency on the deficiency, the restriction to uniformity seems not interesting, and is also not needed, but unrestricted clause-sets are considered. We remark that for every $p \in \mathbb{N}$, $p \geq 3$, there is a polytime translation $t_p : \mathcal{CLS} \rightarrow p\text{-}\mathcal{UCLS}$, such that $t_p(F)$ is satisfiable iff F is, $t_p(F)$ is minimally unsatisfiable iff F is, and $\delta(t_p(F)) = \delta(F)$. This works by replacing clauses C with $|C| < p$ by clauses $C \cup \{v\}, C \cup \{-v\}$ for some new variable v (in the MU-case we will call this a “non-strict full subsumption extension”), and by replacing clauses C with $|C| > p$ by clauses $C' \cup \{v\}, C'' \cup \{-v\}$ for some new variable v and choosing clauses C', C'' with $C = C' \cup C''$ and $|C'| = p - 1, |C''| \geq p - 1$ (in the MU-case again we have a singular DP-extension). But the transformation t_p appear to be useless, since it completely garbles the structure of F .

We conclude these remarks on p -uniform clause-sets by the observation, that for $p \geq 4$ the instances involved above become quickly very big, and only methods from random analysis are available (which by nature are very rough). It seems that these considerations do not have practical relevance. In contrast, we consider

all minimally unsatisfiable clause-sets (and more), that is, the deficiency does not filter out clause-sets, but only organises them in layers. And for a wide range of deficiency values, say, $k = 1, \dots, 10000$, there are interesting and relevant examples.

1.5. Autarkies

An important tool, used in this report to go beyond MU , is the theory of autarkies, which also provides links to various areas of combinatorics; the relations to hypergraph colouring will be discussed in Subsection 1.6.3. Recall that a partial assignment φ is an autarky for $F \in \mathcal{CLS}$ iff every clause $C \in F$ touched by φ (i.e., $\varphi \cap (C \cup -C) \neq \emptyset$) satisfies C (i.e., $\varphi \cap C \neq \emptyset$), which is equivalent to $\forall F' \subseteq F : \varphi * F' \subseteq F'$. Autarkies were introduced in [41] for improved worst-case upper bounds for SAT decision, applying that obviously $\varphi * F$ is sat-equivalent to F ($\varphi * F \in \mathcal{SAT} \Leftrightarrow F \in \mathcal{SAT}$) for an autarky φ . For a recent overview see [11].

Autarky reduction. The reduction of $F \in \mathcal{CLS}$ to $\varphi * F \in \mathcal{CLS}$ for a non-trivial autarky φ , is an essential concept, algorithmically as well as for theoretical understanding; see [11, Subsection 11.10] for an overview on finding autarkies. If we reduce all autarkies, then we obtain the (unique) *lean kernel* of F . If there are no non-trivial autarkies, then we have a lean clause-sets, i.e., $F \in \mathcal{LEAN}$, as already mentioned in Subsection 1.2; this concept was introduced in [10], and [11, Subsection 11.8.3] contains more information. The lean kernel of F is the largest lean sub-clause-set of a clause-set, that is, $\bigcup\{F' \subseteq F : F' \in \mathcal{LEAN}\}$; for recent work on the computation of the lean kernel see [42, 43]. The decision of leanness is coNP-complete, and so consideration of special autarkies is of interest; actually, these considerations are not just “algorithmic hacks”, but in a sense represent various areas of combinatorics (for example matching theory) via “autarky systems”.

Autarky systems. The notion of an “autarky system”, as a selection of special autarkies with similar good properties as general autarkies, was introduced in [13], partially further expanded in [44], and overviewed in [11, Subsection 11.11].

The starting point for an autarky system is to single out a restricted notion of autarky. This restricted autarky notion implies a restricted satisfiability notion, namely clause-sets satisfiable via (iterated) autarky reduction, using only these special autarkies. This is indeed equivalent for “normal autarky systems” to satisfiability by a single such special autarky.⁵ For these normal systems reversely the general autarkies of the system can be derived from those special autarkies, which are also satisfying assignments. For arbitrary autarky systems also the notions “minimal unsatisfiability” and “lean” are defined, and are central properties.

Balanced autarkies yield an example of a rather general autarky system, the basis of autarkies for hypergraph colouring; here for an autarky, touched clauses need not only have some satisfied literal, but also some falsified literal. The corresponding satisfiability notion is “NAE-satisfiability”, and will be further discussed in Subsection 1.6.3.

Matching autarkies. The autarky system especially of importance in this report, besides the full system, is that of *matching autarkies*; for a short introduction see [11, Subsection 11.11.2]. They yield the set $\mathcal{MLEAN} \supset \mathcal{LEAN}$ of *matching-lean clause-sets*, and the set $\mathcal{MSAT} \subset \mathcal{SAT}$ of *matching-satisfiable clause-sets* (called “matched clause-sets” in [22]):

- A matching autarky for $F \in \mathcal{CLS}$ is an autarky φ for F such that for all $C \in F$ touched by φ one can select $x_C \in C$ with $x \in \varphi$ such that the underlying variables $\text{var}(x_C)$ are pairwise different.

⁵“Normal autarky systems” were called “strong autarky systems” in [13, Section 8].

- We have $F \in \mathcal{MSAT} \Leftrightarrow \forall F' \subseteq F : \delta(F) \leq 0$, i.e., $\delta^*(F) = 0$.
- And $F \in \mathcal{MLEAN} \Leftrightarrow \forall F' \subset F : \delta(F') < \delta(F)$.
- Thus for $F \in \mathcal{MLEAN}$ holds $\delta^*(F) = \delta(F)$, and for $F \neq \top$ holds $\delta(F) \geq 1$ (note $\delta(\top) = 0$), a vast generalisation of this fact for \mathcal{MU} .
- More strongly, we have for $F \neq \top$ that $F \in \mathcal{MLEAN} \Leftrightarrow \sigma(F) \geq 1$.
- Every $F \in \mathcal{CLS}$ has a largest matching-lean sub-clause-set, the *matching-lean kernel*, namely $\bigcup\{F' \subseteq F : F' \in \mathcal{MLEAN}\}$, computable in polynomial time (for example via reduction by matching autarkies).

Linear autarkies. A stronger autarky system than matching autarkies is given by “linear autarkies”; we will not use them for the results of this report, but they are an important link to combinatorics, and so we discuss them here; see [11, Subsection 11.11.3] for a more elaborated introduction. “Simple linear autarkies” for $F \in \mathcal{CLS}$ have been introduced in [10], based on linear programming. For $F \in \mathcal{CLS}$ we consider the clause-variable matrix $M(F)$, which is a $c(F) \times n(F)$ matrix over \mathbb{R} (or over \mathbb{Q} for computational purposes), which encodes in the rows the clauses and in the columns the variables, by using 0 for absence of the variable, and ± 1 for positive resp. negative sign. Now the *simple linear autarkies* φ are obtained from solutions $\vec{x} \in \mathbb{R}^{n(F)}$ of $M(F) \cdot \vec{x} \geq 0$, by translating the values \vec{x}_i , where the indices i correspond to the variables of F , into “unassigned” for $\vec{x}_i = 0$, “true” (i.e., 1) for $\vec{x}_i > 0$, and “false” (i.e., 0) for $\vec{x}_i < 0$. It is an easy exercise to see that this yields indeed autarkies. We have a non-trivial simple linear autarky iff $M(F) \cdot \vec{x} \geq 0$ has a non-trivial solution. We obtain the classes

- \mathcal{LLEAN} of “linearly lean clause-sets” (not having a non-trivial simple linear autarky), with $\mathcal{LEAN} \subset \mathcal{LLEAN} \subset \mathcal{MLEAN}$;
- \mathcal{LSAT} of “linearly satisfiable clause-sets” (satisfiable by a sequence of simple linear autarkies), with $\mathcal{MSAT} \subset \mathcal{LSAT} \subset \mathcal{SAT}$.

Linear autarkies, as introduced in [13], are obtained from simple linear autarkies by composition, corresponding to iterated reduction by simple linear autarkies; simple linear autarkies yield an autarky system, while linear autarkies yield a normal autarky system. The main point here is, that the reduction to the linearly-lean kernel can be done by a single linear autarky, and linearly satisfiable clause-sets are satisfiable by a single linear autarky. In Subsection 1.6.3 we discuss the special case of “balanced linear autarkies”.

1.6. Connections to combinatorics

We now discuss the connections between SAT and combinatorics in a wider context than the degree considerations of this report, concentrating on aspects related to minimal unsatisfiability and autarkies (if one is only interested in the results of this report, then these discussions may be ignored). A general source on SAT is the handbook [2]; a classical connection to combinatorics, random satisfiability, is discussed in Chapter 8 ([45]) there, and of further general interest to combinatorics is Chapter 10 ([46]) on symmetry (group theory), Chapter 13 ([47]) on fixed-parameter tractable problems (for example treewidth and related notions), and Chapter 17 ([48]) on the handling of various combinatorial designs to SAT solving, for example from Ramsey theory. Ramsey theory has strong connections to hypergraph colouring, which we discuss next; we mention, that applying SAT solving to hypergraph colouring problems is a powerful tool, and a recent overview can be found in [49], where especially van-der-Waerden numbers are discussed, while a recent success concerning colouring Pythagorean triples can be found in [50].

1.6.1. Hypergraph colouring

Hypergraph-colouring, especially 2-colouring, and SAT are closely connected; see [51, Section 5] for a general introduction and overview on hypergraph colouring (from the combinatorial point of view), while a monograph is given with [52]. An overview especially on the question of the minimum number of hyperedges for a given number of vertices in non- k -colourable hypergraphs is given in [53].

Hypergraphs. For this introduction, a hypergraph G is a finite set of finite subsets of \mathbb{Z} ; so G itself is the set of hyperedges, i.e., $E(G) := G$, while $\bigcup G$ is the set of vertices, i.e., $V(G) := \bigcup G$. The set of all hypergraphs is denoted by $\mathcal{HY}\mathcal{P}$. Let the deficiency be $\delta_{\mathbb{H}}(G) := |E(G)| - |V(G)|$. Note that clause-sets are special hypergraphs ($\mathcal{CLS} \subset \mathcal{HY}\mathcal{P}$), but their deficiency is defined differently. Hypergraphs G with $\delta_{\mathbb{H}}(G) = 0$ are called *square hypergraphs*. Special hypergraphs are the positive clause-sets, and the set of all positive clause-sets is denoted by $\mathcal{PCLS} := \{F \in \mathcal{CLS} : \bigcup F \subset \mathbb{N}\} = \{G \in \mathcal{HY}\mathcal{P} : V(G) \subset \mathbb{N}\}$. For $F \in \mathcal{PCLS}$ we have $\delta(F) = \delta_{\mathbb{H}}(F)$; obviously every hypergraph can be renamed to a positive clause-set. From general clause-sets $F \in \mathcal{CLS}$ we obtain two hypergraphs:

- F itself is a hypergraph (breaking the link between positive and negative literals, which are now just unrelated vertices).

We note that we could have allowed $\mathcal{CLS} = \mathcal{HY}\mathcal{P}$, by allowing tautological clauses (i.e., clauses containing clashing literals) and self-complementary literals ($-0 = 0$). In certain contexts allowing such degenerations has advantages, but in our context it seems best to ban them (for example so we have a direct correspondence between clauses and partial assignments).

- The “variable-hypergraph” of F is $\{\text{var}(C) : C \in F\} \in \mathcal{PCLS}$. This formation for example is important to apply methods from matching theory.

For positive clause-sets both formations collapse to the identity, and we treat positive clause-sets as representing (general) hypergraphs by (special) clause-sets.

Colouring. A k -colouring for $k \in \mathbb{N}_0$ of G is a map $f : V(G) \rightarrow \{0, \dots, k-1\}$ such that for all $H \in G$ there are $x, y \in H$ with $f(x) \neq f(y)$; G is called k -colourable if there exists a k -colouring of G . Note that if there are $H \in G$ with $|H| \leq 1$, then G is not k -colourable for any k . A hypergraph G is *critically k -colourable* if G is k -colourable and not $(k-1)$ -colourable, but for all $H \in G$ the hypergraph $G \setminus \{H\}$ is $(k-1)$ -colourable. In the SAT-context there is no need to discard hyperedges containing at most one vertex, and then *minimally non- k -colourability* is more appropriate, that is G is not k -colourable (possibly not colourable at all), while after removal of any hyperedge G becomes k -colourable. The set of all minimally non- k -colourable hypergraphs is denoted by $\mathcal{MNC}[k] \subset \mathcal{HY}\mathcal{P}$ for $k \in \mathbb{N}_0$. We have $\{\emptyset\}, \{\{x\}\} \in \mathcal{MNC}[k]$ for all $k \in \mathbb{N}_0$ and $x \in \mathbb{Z}$.

We are especially interested in $\mathcal{MNC}[2]$. For $G \in \mathcal{MNC}[2]$ holds $\delta_{\mathbb{H}}(G) \geq 0$, as shown in [54], and so we can consider the sets $\mathcal{MNC}[2]_{\delta_{\mathbb{H}}=k}$ for deficiencies $k \in \mathbb{N}_0$ (all minimally non-2-colourable hypergraphs of deficiency (exactly) k).⁶ The famous problem of deciding in polynomial time, whether a directed graph contains an even cycle, is equivalent to the problem of deciding “ $F \in \mathcal{MNC}[2]_{\delta_{\mathbb{H}}=0}$?” for $F \in \mathcal{HY}\mathcal{P}$ (via simple transformations), and this problem was finally solved in [56, 57]. It was conjectured in [44], that for all $k \in \mathbb{N}_0$ the classes $\mathcal{MNC}[2]_{\delta_{\mathbb{H}}=k}$ are decidable

⁶Indeed in [55, Corollary 8.2] it is shown $\delta_{\mathbb{H}}(G) \geq 0$ for all $G \in \mathcal{MNC}[k]$ and all $k \geq 2$, as a simple application of the autarky method; note that for $G := \{\{1, \dots, n\}\} \in \mathcal{MNC}[k]$ for $k \leq 1$ and $n \geq 2$ holds $\delta_{\mathbb{H}}(G) = 1 - n < 0$.

in polynomial time (see also [11, Conjecture 11.12.1]). More on this in Subsection 1.6.4. In [58] one finds more information on vertex degrees in uniform elements of $\mathcal{MNC}[2]_{\delta_H=0}$ (i.e., where all hyperedges have the same length).

Translating hypergraphs into clause-sets. For a positive hypergraph $G \in \mathcal{PCLS}$ we obtain the translation of 2-colouring to satisfiability via

$$F_2(G) := G \cup \{-H : H \in G\} \in \mathcal{CLS}.$$

For a general discussion of such translations, also considering more colours, see [17, Subsection 1.2]. A hypergraph $G \in \mathcal{PCLS}$ is 2-colourable iff $F_2(G)$ is satisfiable, and G is minimally non-2-colourable iff $F_2(G)$ is minimally unsatisfiable, i.e., $F_2(G) \in \mathcal{MU} \Leftrightarrow G \in \mathcal{MNC}[2]$ (this is easy to prove, and a special case of [55, Lemma 8.1]). Regarding the deficiency we have $\delta(F_2(G)) = \delta_H(G) + |E(G)|$ for $\emptyset \notin G$, and thus e.g. $F_2(\mathcal{MNC}[2]_{\delta_H=0} \cap \mathcal{PCLS})$ is not contained in any $\mathcal{MU}_{\delta=k}$ for some $k \in \mathbb{N}$.

A slight generalisation of the image $F_2(\mathcal{PCLS})$ under this translation is the class of complementation-invariant clause-sets $F \in \mathcal{CLS}$, characterised by $C \in F \Leftrightarrow -C \in F$ for clauses C , as introduced in [44] (see also [11, Subsection 11.4.5]): The image $F_2(\mathcal{PCLS})$ is the set of complementation-invariant PN-clause-sets, that is, clause-sets F where every clause $C \in F$ is positive (i.e., $C \subset \mathbb{N}$) or negative ($-C \subset \mathbb{N}$). See Subsection 1.6.4 for how autarkies, considered for $F_2(G)$, can help understanding G .

Translating clause-sets into hypergraphs. In the other direction a translation was provided in [59]. For $F \in \mathcal{CLS}$ let

$$e(F) := \{C \cup \{0\} : C \in F\} \cup \{v, -v : v \in \text{var}(F)\} \in \mathcal{HYP}.$$

The hypergraph $e(F)$ is 2-colourable iff F is satisfiable, and F is minimally unsatisfiable iff $e(F)$ is minimally non-2-colourable, i.e., $e(F) \in \mathcal{MNC}[2] \Leftrightarrow F \in \mathcal{MU}$ (the direction “ \Leftarrow ” of the latter statement is stated in the proof of Theorem 3 in [5], the other direction is (also) very easy). Furthermore $\delta_H(e(F)) = \delta(F) - 1$. Thus e embeds the classes $\mathcal{MU}_{\delta=k}$ into $\mathcal{MNC}[2]_{\delta_H=k-1}$, which motivates the conjecture, that all $\mathcal{MNC}[2]_{\delta_H=k}$ for $k \in \mathbb{N}_0$ are polytime decidable, as a strengthening of the polytime decision of the $\mathcal{MU}_{\delta=k}$ for $k \in \mathbb{N}$ (recall Subsection 1.4).

We remark that via this embedding e we obtain a proof of $\delta(F) \geq 1$ for $F \in \mathcal{MU}$ from $\delta_H(G) \geq 0$ for $G \in \mathcal{MNC}$ (this is one of the proofs given in [5]). In [5] also an alternative proof of $\delta_H(G) \geq 0$ is given, based on matching theory, plus one further proof of $\delta(F) \geq 1$, using linear algebra, as in [54]. In Subsection 1.6.3 we will further comment on these proofs, as they are unfolded in the theory of autarkies.

We also remark, that the hypergraph class $e(\mathcal{MU}_{\delta=1}) \subset \mathcal{MNC}[2]_{\delta_H=0}$ has the property, that every hypergraph in it different from $\{\{0\}\}$ has a vertex of degree 2 (since every $F \in \mathcal{MU}_{\delta=1}$ different from $\{\emptyset\}$ has a variable of degree 2). More generally, for all $k \in \mathbb{N}$ every hypergraph in $e(\mathcal{MU}_{\delta=k}) \setminus \{\{\{0\}\}\} \subset \mathcal{MNC}[2]_{\delta_H=k-1}$ has a vertex of degree at most $k + 1$. We do not know whether the minimum vertex-degrees of general $G \in \mathcal{MNC}[2]_{\delta_H=k}$ for any (fixed) $k \in \mathbb{N}_0$ are bounded.

1.6.2. Hypergraph transversals

For $G \in \mathcal{HYP}$ let $\text{Tr}(G) \in \mathcal{HYP}$, the *transversal hypergraph* of G , be defined as the set of all minimal $T \subseteq V(G)$ such that $T \cap H \neq \emptyset$ for all $H \in G$. The *Transversal Hypergraph Problem* is the computational problem, given $G, G' \in \mathcal{HYP}$, to decide whether $\text{Tr}(G) = G'$ holds. Equivalently, the input is $G \in \mathcal{HYP}$, and it is to be decided whether $G = \text{Tr}(G)$ holds (obviously this is a special case of the Transversal Hypergraph Problem, and by a polynomial-time translation the general case can be reduced to it). For an overview on this important problem and its many guises

see [60]. It is known that the problem is solvable in quasi-polynomial time, and the long outstanding problem is whether it can be solved in polynomial time.

An *intersecting hypergraph* is a hypergraph $G \in \mathcal{HY}\mathcal{P}$, such that for $H, H' \in G$ with $H \neq H'$ holds $H \cap H' \neq \emptyset$, the class of all intersecting hypergraphs is denoted by $\mathcal{IHY}\mathcal{P} \subset \mathcal{HY}\mathcal{P}$. By definition we have $G \subseteq \text{Tr}(G)$ for $G \in \mathcal{IHY}\mathcal{P}$, and it is not hard to see that for $G \in \mathcal{IHY}\mathcal{P}$ holds $G \in \mathcal{MNC}[2]$ iff $\text{Tr}(G) = G$. Thus the Transversal Hypergraph Problem is equivalent to the problem of deciding, whether an intersecting hypergraph is minimally non-2-colourable. The natural question arises for the decision of the classes $(\mathcal{MNC} \cap \mathcal{IHY}\mathcal{P})_{\delta_H=k}$ for $k \in \mathbb{N}_0$. The case $k = 0$ has been handled in [54], indeed not just deciding the class in polynomial time, but efficiently classifying the elements. The cases $k \geq 1$ appear to be open, and whether decision is possible in polynomial time for fixed k , or is even fixed-parameter tractable (fpt) in k , is an interesting test case for the general Hypergraph Transversal Problem, as well as it is relevant for the understanding of minimally non-2-colourable hypergraphs.

The translation of intersecting hypergraphs $G \in \mathcal{IHY}\mathcal{P}$ into clause-sets $F_2(G) \in \mathcal{CLS}$ yields also a natural and interesting class of clause-sets. *Bihitting clause-sets*, introduced in [61, Subsection 4.2], are those $F \in \mathcal{CLS}$ where $F', F'' \subseteq F$ with $F' \cup F'' = F$, $F' \cap F'' = \emptyset$ exist, such that for all $C' \in F', C'' \in F''$ holds $C' \cap -C'' \neq \emptyset$, while F', F'' itself are clash-free (i.e., $(\bigcup F') \cap -(\bigcup F') = \emptyset$, and $(\bigcup F'') \cap -(\bigcup F'') = \emptyset$). Obviously, the images under F_2 of intersecting hypergraphs are precisely the bihitting complementation-invariant PN-clause-sets (i.e., the set of bihitting clause-sets in the image of F_2), and deciding their minimal unsatisfiability is thus another manifestation of the Hypergraph Transversal Problem, directly related to the decision “ $\text{Tr}(G) = G?$ ”. And another one is to decide SAT for general bihitting clause-sets (as can be easily seen, and is discussed in [61, Subsection 4.3]), this time directly related to the decision “ $\text{Tr}(G) = G?$ ”.

In [55, Theorem 8.14] (the first 6 sections are covered by [16, 17]) the characterisation of [54] (the intersecting $G \in \mathcal{MNC}[2]_{\delta_H=0}$) is translated into the \mathcal{CLS} -language.

1.6.3. Autarkies for hypergraphs

We discuss here now two autarky systems (recall Subsection 1.5 for a general introduction), which are especially relevant for hypergraph colouring.

Balanced autarkies. [44] introduced *balanced autarkies* for $F \in \mathcal{CLS}$, partial assignments φ , which in every clause of F they touch satisfy as well as falsify at least one literal (that is, for $C \in F$ with $C \cap (\varphi \cup -\varphi) \neq \emptyset$ holds $C \cap \varphi \neq \emptyset$ as well as $C \cap -\varphi \neq \emptyset$); [11, Subsection 11.11.4] provides an introduction. This is a normal autarky system, and thus we basically have all the good property general autarkies have. Balanced autarkies are closely related to hypergraph colouring. The balanced autarkies for F are precisely the autarkies of $F \cup \{-C : C \in F\}$, and every autarky for a complementation-invariant clause-set is automatically balanced. A clause-set is balanced-satisfiable, i.e., can be satisfied by a balanced autarky, iff it is NAE-satisfiable (“not-all-equal”; see [62] for basic results).

Balanced autarkies provide the general autarky form for \mathcal{PCLS} (whose elements are all trivially satisfiable, and thus unrestricted autarkies are not of interest here), which represents hypergraphs for the 2-colouring problem: an $F \in \mathcal{PCLS}$ is 2-colourable iff it is balanced-satisfiable, and F is minimally non-2-colourable iff it is minimally balanced-unsatisfiable. Finally we have *balanced lean* clause-sets (i.e., having no non-trivial balanced autarkies), and this is the appropriate notion of “leanness” for hypergraphs, as represented by the class \mathcal{PCLS} ; more precisely, a hypergraph G is lean iff for an isomorphic $F \in \mathcal{PCLS}$ (isomorphic as hypergraph)

we have that F is balanced lean. For lean hypergraphs G we have $\delta_{\mathbb{H}}(G) \geq 0$, and this is properly treated by “balanced linear autarkies”.

Balanced linear autarkies. The special case of “balanced linear autarkies” was introduced in [13, Section 6]; these are the simple linear autarkies for $F \cup \{-C : C \in F\}$ (recall Subsection 1.5).⁷ Equivalently, the balanced linear autarkies φ for $F \in \mathcal{CLS}$ are obtained from solutions $\vec{x} \in \mathbb{R}^{n(F)}$ of $M(F) \cdot \vec{x} = 0$, by translating the values \vec{x}_i as discussed before (it is an easy exercise to see that this yields indeed balanced autarkies). We have a non-trivial balanced linear autarky iff $M(F) \cdot \vec{x} = 0$ has a non-trivial solution, and so, in other words, F is balanced linearly lean iff the columns of $M(F)$ are linearly independent (iff $\text{rank}(M(F)) = n(F)$). Thus if $F \in \mathcal{CLS}$ is balanced linearly-lean, then $\delta(F) \geq 0$ holds; furthermore, as shown in [55, Lemma 7.2], there is then a matching in the clause-variable graph covering all variable nodes, and thus even $\delta^*(F) = \delta(F)$ holds. By noting that $F \in \mathcal{CLS}$ is balanced linearly lean iff $F \cup \{-C : C \in F\}$ is linearly lean, and considering \mathcal{PCLS} , we obtain that for lean hypergraphs G (especially, minimally non-2-colourable) we have $\delta_{\mathbb{H}}(G) \geq 0$. To say the argument again explicitly: Consider a hypergraph $G \in \mathcal{PCLS}$; then G (as a clause-set) is balanced linearly lean iff the variable-clause matrix has linearly independent rows, iff $F_2(G)$ is linearly lean (again, as a clause-set), which is implied by $F_2(G)$ being minimally unsatisfiable (or weaker, being lean), which in turn is equivalent to G (as a hypergraph) being minimally-non-2-colourable. This conclusion “The rows of the incidence matrix [our variable-clause matrix] of a minimally-non-2-colourable hypergraph are linearly independent over \mathbb{R} .” is shown in [54]; see [63, Lemma 4.7] for this and related results, while the conclusion “ $\delta_{\mathbb{H}}(G) \geq 0$ ” is discussed as Principle 2.1 in [63]. For properties of minimally balanced linearly unsatisfiable clause-sets see [44, Section 4].

Fundamental inequalities. We have yet seen two fundamental inequalities, namely $\delta(F) \geq 1$ for $F \in \mathcal{MLEAN}$, as first shown in [5] for minimally unsatisfiable clause-sets, and $\delta(F) \geq 0$ for balanced linearly lean clause-sets, first shown in [54] (as $\delta_{\mathbb{H}}(G) \geq 0$ for minimally non-2-colourable hypergraphs).⁸ Autarky theory shows the general structure of the arguments: We find “obstructions”, which prevent these bounds from holding, where such obstructions are given by a subset $F' \subseteq F$ where there is a partial assignment φ with $\varphi * F' = \top$, while $\text{var}(\varphi) \cap \text{var}(F' \setminus F) = \emptyset$. Now minimally unsatisfiable F do not have such F' , and thus the envisaged bound holds for them, and this is the argumentation in e.g. [54, 5].

But one can go beyond this, exploiting autarky *reduction* $F \rightsquigarrow F \setminus F'$ (as long as possible). Note that the above φ is precisely an autarky, and furthermore possibly one of a special structure. If we just look at general autarkies, then we obtain the first generalisation, to lean clause-sets (having no non-trivial autarkies) or balanced lean clause-sets (having no non-trivial balanced autarkies, covering the hypergraph cases). However often, due to the special structure, these special autarkies can be found in polynomial time, and their application yields some $F_0 \subseteq F$, such that the bound holds for F_0 (while for $F \in \mathcal{MU}$ we just have $F_0 = F$). If we have even an “autarky system”, then F_0 is uniquely determined, that is, does not rely on the choice of the autarkies in the reduction process. The case of main importance for this report is $\delta(F) \geq 1$, where the autarkies are matching autarkies, and the reduced

⁷More precisely one should speak of “balanced simple linear autarkies”, but for convenience “simple” is dropped. We note that “balanced linear autarkies” are balanced and linear autarkies, but in general a balanced and linear autarky need not be a balanced linear autarky, and thus one should speak of “balanced-linear autarkies”; again we abuse language, motivated by the fact that linear autarkies which are also balanced are apparently too general a concept to be useful.

⁸An application yielding Fisher’s inequality (design theory) is discussed in [55, Subsection 7.4] (while Seymour’s inequality is discussed there in Subsection 8.2).

F_0 is the matching-lean kernel of F , while those F with $F_0 = F$ are precisely the $F \in \mathcal{ML}\mathcal{E}\mathcal{AN}$. On the other hand, for hypergraph colouring the fundamental fact is $\delta(F) \geq 0$ for balanced linearly-lean clause-sets, where the autarkies are balanced linear autarkies, and the reduced F_0 is the balanced-linearly-lean kernel of F . In fact, via autarky reduction we obtain a general method to study *decompositions*, which we will discuss next, in the context of matrix analysis.

1.6.4. Qualitative matrix analysis (QMA)

QMA can be understood as the analysis of matrices M over the real numbers in abstraction of the absolute value of the entries, but only their signs count. More precisely, one considers the *qualitative class* $\mathfrak{Q}(M)$, which consists of all matrices with the same dimensions as M , which have entry-wise the same signs as M (positive, zero, negative), and investigates when a property of M holds for all $M' \in \mathfrak{Q}(M)$. For example, a matrix M , such that all $M' \in \mathfrak{Q}(M)$ have linearly independent rows, is called an *L-matrix*. The monograph [64] is an excellent source on QMA until the 1990's, while a more recent overview is given in [65].⁹

Indeed [66] introduces “weak satisfiability”, which is *precisely* the existence of a non-trivial autarky. It is shown ([66, Theorem 5]), that weak satisfiability is NP-complete, which is the earliest known proof of $\mathcal{LE}\mathcal{AN}$ being coNP-complete. However these connections to SAT have not been pursued further.

Starting from [67], which exploits Farkas’ lemma to understand (un)satisfiability, the connections to QMA have been explored in [13, Sections 3, 5]; see [11, Subsection 11.12.1] for a more substantial introduction. It is shown in [13, Remark 5 in Section 5], that *L*-matrices correspond (nearly) precisely (up to transposition and handling of zero-rows/columns and repeated rows/columns) to balanced lean clause-sets, while lean clause-sets correspond (nearly) precisely to so-called *L⁺-matrices* (as investigated in [68]). The square *L*-matrices are called *SNS-matrices*; SNS-matrices are at the heart of the poly-time decision for $\mathcal{MNC}[2]_{\delta_H=0}$ (recall Subsection 1.6.1), and the connections to autarky theory are explored in [44]; see [11, Subsection 11.12.2] for an overview.

Further in the translation of terms, now regarding unsatisfiability: unsatisfiable clause-sets correspond to *sign-central matrices*, minimally unsatisfiable clause-sets correspond to *minimally sign-central matrices*. So [64, Theorem 5.4.3] is yet another proof of $\delta(F) \geq 1$ for $F \in \mathcal{MU}$. The variable-degree, as studied in the current report, corresponds to the number of non-zero entries in the rows of the matrices (while the deficiency is the difference of the number of columns and the number of rows). The elements of $\mathcal{MU}_{\delta=1}$ correspond to *S-matrices*, the elements of $\mathcal{SMU}_{\delta=1}$ correspond to *maximal S-matrices*.

So our Theorem “ $F \in \mathcal{LE}\mathcal{AN} \Rightarrow \mu\text{vd}(F) \leq nM(\delta(F))$ ” is equivalent to the property of *L⁺-matrices* of dimension $m \times n$ (thus $n > m$), that there always exists a row with at most $nM(n - m)$ many non-zero entries.

As mentioned, autarky systems \mathcal{A} (like balanced autarkies, matching autarkies, etc.) also yield a framework for decomposition theorems. The basic decomposition is into \mathcal{A} -lean and \mathcal{A} -satisfiable sub-clause-sets, as given in [13, Theorem 8.5] (for normal autarky systems), which corresponds to a certain unique decomposition of the clause-variable matrix into a triangular shape with two blocks on the diagonal, and generalises various matrix decompositions in QMA, as discussed in [13, Footnote 7, Page 246]. \mathcal{A} -lean clause-sets itself can be further decomposed, and the main result is [44, Lemma 6] (reviewed in [11, Subsection 11.11.5]), generalising [64,

⁹For us the original notion of *L*-matrix, before [64], with (qualitatively) independent *columns*, would be more convenient (then $M(F)$ could be used without transposition), but we stick to the (important) monograph [64].

Theorem 2.2.5], stating that a clause-set $F \in \mathcal{CLS}$ is minimally \mathcal{A} -unsatisfiable iff F is barely \mathcal{A} -lean (it is lean, but removal of any single clause destroys this) and \mathcal{A} -indecomposable (no triangular decomposition into \mathcal{A} -lean blocks is possible for the clause-variable matrix).

1.6.5. *Biclique partitions of (multi-)graphs, and algebraic graph theory*

We finish this overview on related themes in combinatorics by a field of graph theory, which, like QMA, can be understood as a study of clause-sets from a special angle, focusing on the conflict-structure of clauses.

Certain aspects of algebraic graph theory. The starting point is [69], where the problem of “addressing a graph” was introduced. One considers a symmetric matrix D of dimension $m \in \mathbb{N}$ over \mathbb{N}_0 , with a zero-diagonal, where the entries are interpreted as “distances” (in [69] the $D_{i,j}$ are the distances between the nodes of some graph), and asks for the smallest $N \in \mathbb{N}_0$, such that there are m codewords $c_1, \dots, c_m \in \{0, 1, *\}^N$ with the property, that the modified Hamming distance between c_i and c_j , which simply ignores positions with $*$, is $D_{i,j}$. See [70, Chapter 9] for an introduction.

A basic result of [69] is that if D has all entries outside the diagonal equal to 1, then $N = m - 1$ (see also [63, Lemma 6.6] for a discussion in the context of eigenvalue methods; for a direct combinatorial proof see [71]). This follows from the general result $N \geq \max(n_+(D), n_-(D))$ of [69] (the “Lemma of Witsenhausen”), where $n_+(D)$ resp. $n_-(D)$ is the number of positive/negative eigenvalues of D . For the general case in [72] it is shown, that if the distances $D_{i,j}$ are indeed the distances between the nodes of some graph, then we have $N \leq m - 1$.

Actually, the Lemma of Witsenhausen works for arbitrary symmetric matrices D over \mathbb{N}_0 with zero diagonal, and a shift of perspective is useful. From our point of view, a codeword over $\{0, 1, *\}^N$ is nothing else than a clause over the variables $1, \dots, N$, while the modified Hamming distance is the number of clashes (conflicts). So the question is about the existence of clauses C_i for $i \in \{1, \dots, m\}$ over variables $1, \dots, N$, such that $D_{i,j} = |C_i \cap -C_j|$ for $i, j \in \{1, \dots, m\}$. The above parameter $N = N(D)$ is the minimal number N of variables in a (multi-)clause-set representing D (it is an easy exercise to see that N is finite, i.e., a representation is always possible). Considering D as the adjacency matrix of some multigraph (where parallel edges are allowed), we see that N is also equal to the minimum number of bicliques into which the edge-set of that multigraph can be partitioned, and N is therefore denoted by $\text{bcp}(A) \in \mathbb{N}_0$ (the “biclique partition number” of A resp. the corresponding multigraph).¹⁰

Clause-sets as biclique partitions. The essential observation is now that we can go back and forth between biclique partitions of multigraphs and clause-sets. In one direction we can understand clause-sets F as representations of biclique partitions of multigraphs, where for each vertex we get a clause, and from each biclique we obtain a variable, where the two sides of the biclique are the positive and negative occurrences of the variable. So we can understand a multigraph together with a biclique partition as a clause-set, and we can use tools from clause-set-logic to analyse the pair multigraph with biclique-partition. The deficiency then becomes the difference between the number of nodes and the number of bicliques. Satisfiability means that it is possible to select from each biclique one side such that all vertices are covered, and minimal unsatisfiability means that such a covering is not possible, but becomes possible as soon as any vertex is removed. The upper

¹⁰In [73] we used “ $n_s(A)$ ” instead, the “symmetric conflict number”.

bound $\text{VDM}(k) \leq nM(k)$ thus is equivalent to the theorem, that for every biclique partition with n bicliques of a multigraph with m vertices and with this minimal non-coverability property, there exists a biclique with at most $nM(m - n)$ vertices.

For example $A_0 = \{\perp\}$ represents the K_1 , the graph with one vertex, and $A_1 = \{\{1\}, \{-1\}\}$ represents K_2 , consisting of one edge (and two vertices), with this edge constituting the biclique partition. More generally, $\{\{1, \dots, n\}, \{-1\}, \dots, \{-n\}\}$ represents the star with n satellites, where each edge is taken as a biclique; if all the edges shall be taken as one biclique, then we have to use the multi-clause-set $\{\{1\}, n * \{-1\}\}$. To conclude these simple examples, $\{\{1, \dots, n\}, \{-1, \dots, -n\}\}$ represents the dipole D_n , consisting of two vertices and n (parallel) edges, together with the biclique partition with n bicliques.

In the other direction we can understand biclique partitions of multigraphs as representations of clause-sets F , namely the nodes of the *conflict multigraph* $\text{cmg}(F)$ are given by the clauses, while the edges are the conflicts (clashing literal occurrences $x, -x$), and the bicliques are given by the variables (their positive and negative occurrences). In this way we can analyse the influence of the “conflict structure” on properties of clause-sets; the basic notions, as introduced in [73] with underlying report [74], are as follows.

Conflict analysis. The notion “hermitian rank” has been introduced and studied in [75] for arbitrary hermitian matrices A (square matrices with complex numbers as entries, such that transposing the matrix and taking the complex conjugate of each entry yields back the original matrix), denoted by $h(A) := \max(n_+(A), n_-(A)) \in \mathbb{N}_0$. So the Lemma of Witsenhausen takes the form, that for symmetric matrices A over \mathbb{N}_0 with zero diagonal holds $\text{bcp}(A) \geq h(A)$.

For $F \in \mathcal{CLS}$ let $\text{CM}(F)$ (the *conflict matrix*) be the square matrix of dimension $c(F)$ over \mathbb{N}_0 , with entries $|C \cap -D|$ for $C, D \in F$ (thus with zero diagonal), i.e., $\text{CM}(F)$ is the adjacency matrix of $\text{cmg}(F)$. So we can use the hermitian rank as a measure $h : \mathcal{CLS} \rightarrow \mathbb{N}_0$ (as first done in [73, Subsection 3.2]), namely

$$h(F) := h(\text{CM}(F));$$

see Points 1, 3 in [61, Section 2] for various equivalent characterisations.¹¹ By definition we have $\text{bcp}(F) := \text{bcp}(\text{CM}(F)) \leq n(F)$, and thus $h(F) \leq \text{bcp}(F) \leq n(F)$. Since for a principal submatrix A' of a hermitian matrix A holds $h(A') \leq h(A)$ (this follows by “interlacing”; see [76, Theorem 9.1.1]), we get $h(\varphi * F) \leq h(F)$ for all partial assignments φ , and also $h(F') \leq h(F)$ for all $F' \subseteq F$, which gives motivation to consider $h(F)$ as a *complexity measure* for $F \in \mathcal{CLS}$.

In [73] also the *hermitian defect* $\delta_h : \mathcal{CLS} \rightarrow \mathbb{N}_0$ has been introduced as

$$\delta_h(F) := c(F) - h(F),$$

and thus $\delta(F) \leq \delta_h(F)$; see Point 2 in [61, Section 2] for a geometric characterisation (as the “Witt index” of the quadratic form associated with $\text{CM}(F)$). Actually $\delta^*(F) \leq \delta_h(F)$ holds and even stronger properties (see [73, Subsection 3.3]). An important property is (again) $\delta_h(\varphi * F) \leq \delta_h(F)$ for all $F \in \mathcal{CLS}$ and partial assignments φ , together with $\delta_h(F') \leq \delta_h(F)$ for all $F' \subseteq F$, by [73, Corollary 9], and so we might consider the hermitian defect as a stabilised version of the maximal defect (both are also complexity measures; recall that we have fixed-parameter tractable SAT decision for input $F \in \mathcal{CLS}$ in the parameter $\delta^*(F)$). Note that in general we can have $\delta^*(\varphi * F) > \delta^*(F)$, for example $F := \{\{1\}\}$ has $\delta^*(F) = \delta(F) = 0$, while for $F' := \langle 1 \rightarrow 0 \rangle * F$ we get $F' = \{\perp\}$, and thus $\delta^*(F') = \delta(F') = 1$.

¹¹For Point 1(c) there it must be a “diagonal matrix A' ”.

See [7, Subsection 3.3] and [16, Subsection 11.2] for more information on $\delta^*(\varphi * F)$; splitting on a single(!) variable is very important for this report, with the basic fact $\delta^*(\langle x \rightarrow 1 \rangle * F) \leq \delta(F)$ for $F \in \mathcal{MU}$ and any literal x .

Applications. The first direct applications applied the fact $\delta(F) \leq \delta_h(F)$ for $F \in \mathcal{CLS}$, namely that for a hitting clause-set $F \in \mathcal{HIT}$ (equivalently, all entries of $\text{CM}(F)$ outside the diagonal are non-zero) with a regular conflict multigraph (i.e., all entries of $\text{CM}(F)$ outside the diagonal are equal) we have $\delta_h(F) \leq 1$, and thus $\delta(F) \leq 1$ ([73, Theorem 33]).¹² We get that $\mathcal{SMU}_{\delta=1} = \mathcal{UHIT}_{\delta=1}$ is (precisely) the class of unsatisfiable hitting clause-sets with regular conflict multigraph ([73, Corollary 34]; a combinatorial proof of this was independently found in [77, Lemma 11]), and is also (precisely) the class of unsatisfiable clause-sets F with $\delta_h(F) \leq 1$ ([73, Theorem 26]).

A clause-set $F \in \mathcal{CLS}$ is called *exact* ([73, Subsection 3.4]) if $\text{bcp}(F) = n(F)$, that is, F is optimal in realising $\text{cmg}(F)$ with respect to the number of variables. Deciding exactness is coNP-complete, while the special class of *eigensharp clause-sets*, defined by $h(F) = n(F)$, or, equivalently, $\delta_h(F) = \delta(F)$, is decidable in polynomial time. With [73, Theorem 14] every eigensharp clause-set is matching lean. This leads to [73, Conjecture 16], “Every exact clause-set, whose conflict-matrix is the distance matrix of some connected graph, is matching lean.”, which generalises the already mentioned main result of [72] (the proof of the “squashed cube conjecture”).

As already mentioned, we consider $h(F)$ for $F \in \mathcal{CLS}$ as some form of complexity measure, measuring the complexity of representing the conflicts of F via simple matrices. In [61] polytime SAT decision in case $h(F) \leq 1$ was shown, while the cases $h(F) \leq k$ for fixed $k \geq 2$ are open; an interesting stepping stone would be to show polytime SAT decision for $F \in \mathcal{CLS}$ with $\text{bcp}(F) \leq k$ (recall $\mathcal{CLS}_{\text{bcp} \leq k} \subseteq \mathcal{CLS}_{h \leq k}$). The notion of *blocked clauses*, a special type of clauses which can be removed sat-equivalently, introduced in [78], is important here, and [61, Theorem 3] shows, that from $F \in \mathcal{CLS}_{h \leq 1}$ after elimination of all blocked clauses (which yields a unique sub-clause-set) we obtain $F' \subseteq F$ with $\text{bcp}(F') \leq 1$. We recall from Subsection 1.6.2, that SAT-decision for F' is now a special case of the Transversal Hypergraph Problem, namely, as shown in [61, Lemma 11], the problem is exactly the *Exact Transversal Hypergraph Problem*, where every transversal must be “exact”, that is, must intersect every hyperedge in exactly one vertex; this problem is decidable in polynomial time by [79], and thus we get SAT-decision for $\mathcal{CLS}_{h \leq 1}$ in polynomial time. The characterisation of $F \in \mathcal{MU}$ with $\text{bcp}(F) \leq 1$ is an open problem (while we have polytime membership decision for $\mathcal{MU}_{\text{bcp} \leq 1}$), and by [61, Conjecture 16] they would have a very simple structure.

We conclude by mentioning that in [17, Section 6] the above basic facts are generalised to non-boolean clause-sets, and that by extending the reduction of multiclique partitions to biclique partitions from [80] a new and interesting translation from non-boolean to boolean clause-sets was obtained.

1.7. Overview on results

Sections 2 to 6 provide foundations for the main results in the later sections. In Section 2 basic notions and concepts regarding clause-sets and autarkies are reviewed. In Section 3 we discuss minimal unsatisfiability, with some auxiliary results on saturation (adding literal occurrences to clauses, to make minimal unsatisfiability robust against splitting) and splitting. Section 4 reviews the concept of “variable-minimal unsatisfiability”, as introduced in [15], i.e., the class $\mathcal{MU} \subset \mathcal{VMU} \subset \mathcal{USAT}$. There are mistakes in this paper, and we rectify them:

¹²In [73] unfortunately the term “uniform” was (mis)used instead of “regular”.

- we show that $\mathcal{VMU} \subset \mathcal{LEAN}$ holds (Lemma 4.3);
- we provide a corrected characterisation of \mathcal{VMU} (Lemma 4.5);
- and we give a corrected proof of polytime decision of $\mathcal{VMU}_{\delta=k}$ for fixed k in Theorem 4.7, where we also obtain the stronger result, that decision of \mathcal{VMU} is fixed-parameter tractable in the deficiency $\delta(F)$.

Section 5 is then concerned with singular variables, eliminating them via “singular DP-reduction”, and creating them via “singular extensions”. An important auxiliary result is Lemma 5.4, showing that eliminating singular variables is harmless for upper bounds on the minimum variable-degree; we also show various auxiliary results on unit-clauses in minimally unsatisfiable clause-sets. This block of preparatory sections is concluded by Section 6 on “full subsumption resolution”, the ubiquitous reduction $C \vee x, C \vee \bar{x} \rightsquigarrow C$, which becomes an “extension” in the other direction; as an application, in Theorem 6.13 we can determine precisely the possible $n(F)$ and $c(F)$ for $F \in \mathcal{MU}_{\delta=k}$.

The first main results (but still on the preparation side) one finds in Section 7, which introduces the numbers $\text{nM}(k) \in \mathbb{N}$ and proves exact formulas and sharp lower and upper bounds; the point here is that the introduction of $\text{nM}(k)$ happens via a recursion which is tailor-made for our application in Section 8, but which makes it somewhat difficult to determine the numbers in a global way. A main technical result is Theorem 7.15, while Theorem 7.21 proves the general formula.

In Section 8 then we find a basic central result of this report, the upper bound $\text{VDM}(k) = \mu\text{vd}(\mathcal{MU}_{\delta=k}) \leq \text{nM}(k)$ (Theorem 8.3). Section 9 is concerned with generalising this upper bound. An interesting auxiliary class $\mathcal{SED} \subset \mathcal{CLS}$, clause-sets where deficiency and surplus coincide, is introduced in Subsection 9.1. The main characterisation of \mathcal{SED} is given in Theorem 9.5 (removal of all clauses containing any given variable yields a matching-satisfiable sub-clause-set), and we obtain Corollary 9.6, that unsatisfiable elements of \mathcal{SED} are in fact in \mathcal{VMU} . In Subsection 9.2 the upper bound for \mathcal{MU} then is lifted to lean clause-sets in Theorem 9.10, and also sharpened via replacing the deficiency δ by the surplus σ . Theorem 9.14 shows that the upper bound is sharp for any class between $\mathcal{VMU} \cap \mathcal{SED}$ and \mathcal{LEAN} .

Section 10 concerns algorithmic applications. A corollary of Theorem 9.10 is, that if the asserted upper bound on the minimum variable degree is not fulfilled, then a non-trivial autarky must exist (Lemma 10.1). Since the variable-set of such a non-trivial autarky is polytime computable, we show in Theorem 10.2 that we can indeed establish the upper bound shown for lean clause-sets also for general clause-sets, after a polytime autarky-reduction. In Subsection 10.2 then the problem of finding such autarky (that is, finding the assignment) is discussed, with Conjecture 10.3 making precise our believe that one can find such autarkies efficiently. Theorem 10.9 pinpoints the “critical” class $\mathcal{MLCR} \subset \mathcal{SAT} \cap \mathcal{SED}$, which is polytime decidable, and where we know that these clause-sets are satisfiable, but we even don’t know how to find any non-trivial autarky efficiently. This block on generalisations of the min-var-degree upper bound is concluded by Section 11, where we discuss the possibilities to generalise it to matching-lean clause-sets (where only the absence of special (non-trivial) autarkies is guaranteed).

In Section 12 we then turn to the study of the numbers $\text{VDM}(k)$, looking now for improved upper bounds and matching lower bounds. We present two infinite classes of deficiencies k with $\text{VDM}(k) = \text{nM}(k)$, and present a general method of obtaining lower bounds for $\text{VDM}(k)$, via counting full clauses (clauses containing all variables — these clause are strong structural anchors). In Section 13 we introduce a general recursive method to obtain upper bounds like $\text{nM}(k)$, via the “non-Mersenne operator” $\text{NM}(f)$, which takes a “valid bounds function” f , that is, some partial information on $\text{VDM}(k)$, and improves it (Definition 13.12). Theorem 13.10 shows

that this indeed yields a valid method for improving upper bounds on $\text{VDM}(k)$, while in Theorem 13.15 we demonstrate how this method recovers $\text{nM}(k)$, by just starting with the information $\text{VDM}(1) = 2$. In Section 14 we harvest (first) fruits of these methods. First in Theorem 14.1 we show $\text{VDM}(k) = \text{nM}(k)$ for $k \leq 5$. Then in Theorem 14.4 we prove $\text{VDM}(6) = \text{nM}(6) - 1$ (using a variety of structural results on \mathcal{MU} provided in this report). Plugging this information on VDM into our machinery, we obtain the improved upper bound $\text{VDM} \leq \text{nM}_1$ in Theorem 14.6, while in Theorem 14.6 we determine $\text{nM}_1(k)$ numerically.

Finally, in Section 15 open problems are stated, thoroughly discussing research perspectives, including nine conjectures. Subsection 15.2 discusses improved upper bounds for $\text{VDM}(k)$ from the forthcoming work [81]. Subsection 15.3 is about improved lower bounds, via counting full clauses. In Lemma 15.2 we present the lower bound via the ‘‘Smarandache primitive function’’ $S_2(k)$ from [82], yielding the first-order asymptotic determination of $\text{VDM}(k) \sim k$ (Corollary 15.4), where now the open question is about the asymptotic determination of $\text{VDM}(k) - k$. In Subsection 15.4 we discuss generalisations to non-boolean clause-sets.

The central Conjecture 15.6 of the project of ‘‘understanding \mathcal{MU} ’’, on the finitely many ‘‘characteristic patterns’’ for each $\mathcal{MU}_{\delta=k}$, is discussed in Subsection 15.5. An important special case is Conjecture 15.7 (now a fully precise statement), about the classification of unsatisfiable hitting clause-sets (or ‘‘disjoint/orthogonal tautologies’’ in the terminology of DNFs). In Lemma 15.9 we show how two of the conjectures together yield computability of $\text{nM}(k)$.

This report is a substantial extension of the conference paper [14]: Section 3 there has been extended to Section 7 here, with considerable more details and examples on non-Mersenne numbers. Section 4 there is covered by Sections 8, 9 and 10, with various additional results (for example showing sharpness of the upper bound for \mathcal{LEAN}). And the results for Section 5 there are contained in Subsection 11 here. All other sections in this report are new.

2. Preliminaries

We follow the general notations and definitions as outlined in [11], where also further background on autarkies and minimal unsatisfiability can be found. We use $\mathbb{N} = \{n \in \mathbb{Z} : n \geq 1\}$ and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For the binary logarithm we use $\text{ld}(x) := \log_2(x) \in \mathbb{R}$ (‘‘logarithm dualis’’) for $x \in \mathbb{R}_{>0}$, and $\text{fld}(x) := \lfloor \text{ld}(x) \rfloor \in \mathbb{Z}$ for the integral part (e.g., $\text{fld}(1) = 0$, $\text{fld}(2) = \text{fld}(3) = 1$, $\text{fld}(4) = \text{fld}(5) = 2$).

We apply standard set-theoretic concepts, like that of a map as a set of pairs, and standard set-theoretic notations, like $f(S) = \{f(x) : x \in S\}$ for maps f and $S \subseteq \text{dom}(f)$, and ‘‘ \subset ’’ for the strict subset-relation. We use also the less-known notation ‘‘ $A \cup B$ ’’ for union in case A, B are disjoint, that is, $A \cup B := A \cup B$ is only defined for $A \cap B = \emptyset$. For maps f, g with the same domain X we use $f \leq g := \Leftrightarrow \forall x \in X : f(x) \leq g(x)$ (i.e., pointwise comparison), while $f < g := \Leftrightarrow \forall x \in X : f(x) < g(x)$.

2.1. Clause-sets

The basic structure is a set \mathcal{LIT} , the elements called ‘‘literals’’, together with a fixed-point free involution called ‘‘complementation’’, written $x \in \mathcal{LIT} \mapsto \bar{x} \in \mathcal{LIT}$; so the laws are $\bar{\bar{x}} = x$ and $\bar{x} = x$, which imply $x \neq y \Rightarrow \bar{x} \neq \bar{y}$, for $x, y \in \mathcal{LIT}$. We assume $\mathbb{Z} \setminus \{0\} \subseteq \mathcal{LIT}$, with $\bar{x} = -x$ for $x \in \mathbb{Z} \setminus \{0\}$. For a set L of literals we define $\bar{L} := \{\bar{x} : x \in L\}$. Furthermore a set $\mathbb{N} \subseteq \mathcal{VA} \subset \mathcal{LIT}$, the elements called ‘‘variables’’, is given, with $\mathcal{LIT} = \mathcal{VA} \cup \bar{\mathcal{VA}}$. Variables are also called ‘‘positive literals’’, while literals \bar{v} for $v \in \mathcal{VA}$ are called ‘‘negative literals’’. The ‘‘underlying variable’’ of a literal is given by the operation $\text{var} : \mathcal{LIT} \rightarrow \mathcal{VA}$ (‘‘forgetting complementation’’), with $\text{var}(v) := v$ and $\text{var}(\bar{v}) := v$ for $v \in \mathcal{VA}$.

Example 2.1 We can thus write e.g. $1, 6$ for two (different) variables, and $1, 5, -1$ for three (different) literals. In examples we will also use v, w, a, b, c and such letters for variables (as it is customary), and accordingly \bar{v} etc. for literals, and in this context (only) it is then understood that these variables are pairwise different. So $\{v, w, x, \bar{x}\}$, when given in an example (without further specification), denotes a set of literals with $|\{v, w, x, \bar{x}\}| = 4$ and $|\{v, w, x, \bar{x}\} \cap \mathcal{VA}| = 3$.

Without restriction we could assume $\mathcal{LIT} = \mathbb{Z} \setminus \{0\}$ (as we did in the Introduction), but it is often convenient to use arbitrary mathematical objects as variables. All our objects build from literals are finite, and thus, because of the infinite supply of variables, there will always be “new variables” (that’s the mathematical point of having natural numbers as variables — we won’t use the arithmetical structure).

A **clause** C is a finite and clash-free set of literals (i.e., $C \cap \bar{C} = \emptyset$), the set of all clauses is \mathcal{CL} . A **clause-set** is a finite set of clauses, the set of all clause-sets is \mathcal{CLS} . The simplest clause is the *empty clause* $\perp := \emptyset \in \mathcal{CL}$, the simplest clause-set is the *empty clause-set* $\top := \emptyset \in \mathcal{CLS}$. The set of all **hitting clause-sets** is denoted by $\mathcal{HIT} \subset \mathcal{CLS}$, those $F \in \mathcal{CLS}$ such that two different clauses $C, D \in F$, $C \neq D$, have at least one clash, i.e., $C \cap \bar{D} \neq \emptyset$. In the language of DNF, hitting clause-sets are known as “orthogonal” or “disjoint” DNF’s; see [83, Chapter 7].

Example 2.2 We have e.g. $\{1, 2, -3\} \in \mathcal{CL}$, while $\{-1, 1\} \notin \mathcal{CL}$. The only clause-set in \mathcal{HIT} containing the empty clause is $\{\perp\} \in \mathcal{HIT}$. An example of a non-hitting clause-set is $\{\{1, 2\}, \{-1, 2\}, \{3\}\} \in \mathcal{CLS} \setminus \mathcal{HIT}$, where we obtain an element of \mathcal{HIT} if we add literal -2 to the third clause.

We use $\text{var}(F) := \bigcup_{C \in F} \text{var}(C) \subset \mathcal{VA}$ for the set of variables of $F \in \mathcal{CLS}$, where $\text{var}(C) := \{\text{var}(x) : x \in C\} \subset \mathcal{VA}$ is the set of variables of clause $C \in \mathcal{CL}$. The possible literals for a clause-set F are given by $\text{lit}(F) := \text{var}(F) \cup \overline{\text{var}(F)} \subset \mathcal{LIT}$, while the actually occurring literals are just given by $\bigcup F \subset \mathcal{LIT}$ (the union of the clauses of F). A literal x is **pure** for F if $\bar{x} \notin \bigcup F$. For a clause-set F we use the following measurements:

- $n(F) := |\text{var}(F)| \in \mathbb{N}_0$ is the number of variables,
- $c(F) := |F| \in \mathbb{N}_0$ is the number of clauses,
- $\delta(F) := c(F) - n(F) \in \mathbb{Z}$ is the **deficiency** (the difference of the number of clauses and the number of variables),
- $\ell(F) := \sum_{C \in F} |C| \in \mathbb{N}_0$ is the number of literal occurrences.

We call a clause C **full** for a clause-set F if $C \in F$ and $\text{var}(C) = \text{var}(F)$, while a clause-set F is called full if every clause is full. For a finite set V of variables let

$$\mathbf{A}(V) := \{C \in \mathcal{CL} : \text{var}(C) = V\} \in \mathcal{CLS}.$$

Obviously $\mathbf{A}(V) \in \mathcal{HIT}$ is the set of all $2^{|V|}$ full clauses over V , and $F \in \mathcal{CLS}$ is full iff $F \subseteq \mathbf{A}(\text{var}(F))$. We use $\mathbf{A}_n := \mathbf{A}(\{1, \dots, n\})$ for $n \in \mathbb{N}_0$. Dually, a variable $v \in \mathcal{VA}$ is called **full** for a clause-set F if for all $C \in F$ holds $v \in \text{var}(C)$. A clause-set is full iff every $v \in \text{var}(F)$ is full.

Example 2.3 For $F := \{\perp, \{1\}, \{-1, 2\}\}$ we have:

1. $\text{var}(F) = \{1, 2\}$, $\text{lit}(F) = \{-1, 1, -2, 2\}$, $\bigcup F = \{-1, 1, 2\}$.
2. Literal 2 is pure for F (the other literals in $\text{lit}(F)$ are not pure).
3. $n(F) = 2$, $c(F) = 3$, $\delta(F) = 1$, $\ell(F) = 3$.

4. $\{-1, 2\}$ is a full clause of F , while the two other clauses are not full.
5. F has no full variable, while $F \setminus \{\perp\}$ has the (single) full variable 1.

The standard “complete” full clause-sets are $A_0 = \{\perp\}$, $A_1 = \{\{-1\}, \{1\}\}$,

$$A_2 = \{\{-1, -2\}, \{-1, 2\}, \{1, -2\}, \{1, 2\}\},$$

and so on.

We often define a class of clause-sets via some measure μ as follows:

Definition 2.4 Consider a class $\mathcal{C} \subseteq \mathcal{CLS}$ and a measure $\mu : \mathcal{CLS} \rightarrow \mathbb{R}$. For $a \in \mathbb{R}$ we use $\mathbf{C}_{\mu=a} := \{F \in \mathcal{C} : \mu(F) = a\}$, and similarly we use $\mathbf{C}_{\mu < a}$ and analogous notations.

When we use the form “ $\mathbf{C}_{\mu \square a}$ ”, then μ stands for a measure (e.g., $\mu = \delta$ or $\mu = n$).

Example 2.5 $\mathcal{CLS}_{n=0} = \mathcal{CLS}_{\ell=0} = \{\top, \{\perp\}\}$, $\mathcal{CLS}_{c=0} = \{\top\}$, and $\mathcal{CLS}_{n < 0} = \emptyset$.

2.2. Semantics

A **partial assignment** is a map $\varphi : V \rightarrow \{0, 1\}$ for some finite (possibly empty) set $V \subset \mathcal{VA}$ of variables, where $\text{var}(\varphi) := V$ and $\text{lit}(\varphi) := \text{var}(\varphi) \cup \text{var}(\varphi)$. The set of all partial assignments is denoted by \mathcal{PASS} . For a literal $x \in \text{lit}(\varphi)$ we also define $\varphi(x) \in \{0, 1\}$, via $\varphi(\bar{v}) := 1 - \varphi(v)$ for $v \in \text{var}(\varphi)$. Via a small abuse of language we define $\varphi^{-1}(\varepsilon) := \{x \in \text{lit}(\varphi) : \varphi(x) = \varepsilon\} \in \mathcal{CL}$ for $\varepsilon \in \{0, 1\}$. Special partial assignments are the *empty partial assignment* $\langle \rangle := \emptyset$, and for literals $x \in \mathcal{LIT}$ and $\varepsilon \in \{0, 1\}$ the partial assignment $\langle x \rightarrow \varepsilon \rangle \in \mathcal{PASS}$, with $\text{var}(\langle x \rightarrow \varepsilon \rangle) = \{\text{var}(x)\}$ and $\langle x \rightarrow \varepsilon \rangle(x) = \varepsilon$.

The application of a partial assignment $\varphi \in \mathcal{PASS}$ to a clause-set F is denoted by $\varphi * F$, which yields the clause-set obtained from F by removing all satisfied clauses (which have at least one literal set to 1), and removing all falsified literals from the remaining clauses:

$$\varphi * F := \{C \setminus \varphi^{-1}(0) : C \in F \wedge C \cap \varphi^{-1}(1) = \emptyset\} \in \mathcal{CLS}.$$

This definition is motivated by the default interpretation of a clause-set as a “conjunctive normal form” (CNF), where a clause is understood as a disjunction of literals (thus is satisfied iff at least one literal in it is satisfied), while a clause-set is understood as a conjunction of its clauses (thus is satisfied iff all clauses are satisfied). A clause-set F is *satisfiable* iff there is a partial assignment φ with $\varphi * F = \top$, otherwise F is *unsatisfiable*. The set of satisfiable clause-sets is denoted by $\mathbf{SAT} \subset \mathcal{CLS}$, while $\mathbf{USAT} := \mathcal{CLS} \setminus \mathbf{SAT}$ denotes the set of all unsatisfiable clause-sets.

Example 2.6 If $F \in \mathbf{USAT}$ and for $F' \in \mathcal{CLS}$ holds $F \subseteq F'$, then also $F' \in \mathbf{USAT}$ (satisfying a clause-sets gets harder the more clauses there are).

By definition we have $\varphi * F = \top$ iff $\forall D \in F : \varphi^{-1}(1) \cap D \neq \emptyset$; thus $F \in \mathbf{SAT}$ iff there is a clause $C \in \mathcal{CL}$ with $C \cap D \neq \emptyset$ for all $D \in F$. (We could write “ $C \cap D \neq \perp$ ” here, but it appears somewhat more natural to use “ \emptyset ” here.)

The unsatisfiable hitting clause-sets are denoted by $\mathbf{UHIT} := \mathbf{USAT} \cap \mathbf{HIT}$.

Example 2.7 $\top \in \text{SAT} \cap \text{HIT}$ and $\{\perp\} \in \text{UHIT}$. In general a full clause-set F is unsatisfiable iff $F = A(\text{var}(F))$, and thus $A(V) \in \text{UHIT}$ for all finite $V \subset \mathcal{VA}$.

The fundamental property for $F \in \text{HIT}$ is that two different clauses do not have a common falsifying assignment. More precisely, consider $\varphi, \psi \in \text{PASS}$, such that there are $C, D \in F$, $C \neq D$, with $\varphi * \{C\} = \psi * \{D\} = \{\perp\}$ (that is, $\perp \in \varphi * F \cap \psi * F$, where there are different falsified clauses for these two partial assignments). Then φ, ψ are incompatible, i.e., there is $v \in \text{var}(\varphi) \cap \text{var}(\psi)$ with $\varphi(v) \neq \psi(v)$.

It follows easily that for $F \in \text{HIT}$ holds $F \in \text{USAT} \Leftrightarrow \sum_{C \in F} 2^{-|C|} = 1$.

A nice exercise is to show $\text{UHIT}_{\delta \leq 0} = \emptyset$ (in Section 2.6 a more general result is stated).

Finally, the *semantical implication* $F \models C$ for $F \in \text{CLS}$ and clauses $C \in \text{CL}$ holds iff $\forall \varphi \in \text{PASS} : \varphi * F = \top \Rightarrow \varphi * \{C\} = \top$. We have $F \in \text{USAT} \Leftrightarrow F \models \perp$.

2.3. Resolution

Two clauses $C, D \in \text{CL}$ are **resolvable** if $|C \cap \overline{D}| = 1$, i.e., they clash in exactly one variable (called the resolution variable $\text{var}(x)$, while x is called the resolution literal). For two resolvable clauses C and D , the **resolvent** $C \diamond D := (C \cup D) \setminus \{x, \overline{x}\} \in \text{CL}$ for $C \cap \overline{D} = \{x\}$ is the union of the two clauses minus the resolution literal and its complement. As it is well-known (the earliest source is [84, 85]), a clause-set F is unsatisfiable iff via resolution (i.e., closing F under addition of resolvents) we can derive \perp , and, more generally, we have $F \models C$ iff from F via resolution a clause $C' \subseteq C$ is derivable.

An important reduction for clause-sets $F \in \text{CLS}$ and variables $v \in \mathcal{VA}$, resulting in a clause-set satisfiability-equivalent to F (satisfiable iff F is; sometimes called “**equi-satisfiable**”) and with variable v eliminated, is **DP-reduction**

$$\text{DP}_v(F) := \{C \in F : v \notin \text{var}(C)\} \cup \{C \diamond D : C, D \in F \wedge C \cap \overline{D} = \{v\}\} \in \text{CLS}$$

(also called “variable elimination”), obtained from F by removing all clauses containing variable v from F , and replacing them by their resolvents on v . See [19] for a fundamental study of DP-reduction. The satisfying assignments φ of $\text{DP}_v(F)$ (i.e., $\varphi * \text{DP}_v(F) = \top$) with $\text{var}(\varphi) = \text{var}(F) \setminus \{v\}$ are precisely the satisfying assignments φ of F with $\text{var}(\varphi) = \text{var}(F)$, when restricting φ to $\text{var}(F) \setminus \{v\}$. Logically, $\text{DP}_v(F)$ is equivalent to $\exists v : F$, the existential quantification of v for F (but we do not use quantifiers in this report, so this remark might be ignored here).

2.4. Multi-clause-sets and restrictions

These notions are generalised to **multi-clause-sets**, which are maps $F : \text{CL} \rightarrow \mathbb{N}_0$, such that the underlying set of clauses $\{C \in \text{CL} : F(C) \neq 0\}$ is finite, and so we speak of the **underlying clause-set**; the set of all multi-clause-sets is denoted by $\underline{\text{CLS}} := \{F : \text{CL} \rightarrow \mathbb{N}_0 \mid \text{CL} \setminus F^{-1}(0) \text{ is finite}\}$.¹³ Clause-sets are implicitly promoted to multi-clause-sets, if needed, by using their characteristic functions, and multi-clause-sets are implicitly cast down, if needed, to clause-sets by considering the underlying clause-set; “if needed” refers to operations which either require multi-clause-sets or clause-sets. If however we want to make explicit these operations, we use $\text{cls} : \underline{\text{CLS}} \rightarrow \text{CLS}$ (with $\text{cls}(F) := \text{CL} \setminus F^{-1}(0)$) and $\underline{\text{cls}} : \text{CLS} \rightarrow \underline{\text{CLS}}$ (with $\underline{\text{cls}}(F)(C) := 1$ if $C \in F$, and $\underline{\text{cls}}(F)(C) := 0$ otherwise). For $F \in \underline{\text{CLS}}$ we extend the basic operations in the obvious way:

- $\text{var}(F) := \text{var}(\text{cls}(F))$, $\text{lit}(F) := \text{lit}(\text{cls}(F))$, $\bigcup F := \bigcup \text{cls}(F)$.

¹³In earlier papers we used “*MCLS*” instead of “*CLS*”.

- $n(F) := n(\text{cls}(F)) \in \mathbb{N}_0$, $c(F) := \sum_{C \in \mathcal{CL}} F(C) \in \mathbb{N}_0$, $\delta(F) := c(F) - n(F) \in \mathbb{Z}$, $\ell(F) := \sum_{C \in F} F(C) \cdot |C| \in \mathbb{N}_0$.

The application of partial assignments $\varphi \in \mathcal{PASS}$ to a multi-clause-set $F \in \mathcal{CL}\mathcal{S}$ yields a *multi-clause-set* $\varphi * F \in \mathcal{CL}\mathcal{S}$, where the multiplicity of a clause $C \in \mathcal{CL}$ in $\varphi * F$ is the sum of all multiplicities of clauses $D \in F$ (i.e., $D \in \text{cls}(F)$) which are shortened to C by φ :

$$(\varphi * F)(C) := \sum_{D \in F, D \cap \varphi^{-1}(1) = \emptyset, D \setminus \varphi^{-1}(0) = C} F(D).$$

Example 2.8 If φ is a total assignment for F (assigns all variables of F , that is, $\text{var}(\varphi) = \text{var}(F)$), then $\varphi * F$ is $\{m * \perp\}$, denoting the multiplicity of a clause by a (formal) factor, with $m = \sum_{C \in F, C \cap \varphi^{-1}(1) = \emptyset} F(C) \in \mathbb{N}_0$ (so $m = 0 \Leftrightarrow \varphi * F = \top$).

For us the clause-sets are the objects of interests, while multi-clause-sets are only auxiliary devices, created by the operation of “restriction” defined next (Definition 2.9). However we have to take care of the details, and thus together with introducing a class $\mathcal{C} \subseteq \mathcal{CL}\mathcal{S}$ we also introduce the corresponding class $\underline{\mathcal{C}} \subseteq \mathcal{CL}\mathcal{S}$ of multi-clause-sets, using the generalised definition of \mathcal{C} , and where $\mathcal{C} = \{\text{cls}(F) : F \in \underline{\mathcal{C}}\} = \text{cls}(\underline{\mathcal{C}})$ must hold. We have two main forms of relations between \mathcal{C} and $\underline{\mathcal{C}}$.

To start with, the classes $\underline{\mathcal{SAT}}$ and $\underline{\mathcal{USAT}}$ are **invariant under multiplicities**, that is, a multi-clause-set is in it iff the underlying clause-set is in the underlying class of clause-sets (\mathcal{SAT} resp. \mathcal{USAT}). In general $\underline{\mathcal{C}}$ is invariant under multiplicities iff for all $F, F' \in \mathcal{CL}\mathcal{S}$ with $\text{cls}(F) = \text{cls}(F')$ either $F, F' \in \underline{\mathcal{C}}$ or $F, F' \notin \underline{\mathcal{C}}$ holds. The other extreme we have with the class $\underline{\mathcal{HIT}}$ of multi-hitting-clause-sets, which **disallows multiplicities**, that is, all multiplicities must be 1 (since clauses can not clash with themselves, by definition of clauses), and thus up to the canonical identification the classes $\underline{\mathcal{HIT}}$ and $\underline{\mathcal{HIT}}$ are identical. In general $\underline{\mathcal{C}}$ disallows multiplicities iff for all $F \in \underline{\mathcal{C}}$ holds $F(\mathcal{CL}) \subseteq \{0, 1\}$ (in other words, $\underline{\mathcal{C}} = \underline{\text{cls}}(\mathcal{C})$).

An important operation with multi-clause-set is the “restriction” to a set of variables (see [16, Subsection 3.5] for more information):

Definition 2.9 For a set $V \subseteq \mathcal{VA}$ of variables and a multi-clause-set $F \in \mathcal{CL}\mathcal{S}$ by $F[V] \in \mathcal{CL}\mathcal{S}$ the **restriction** of F to V is denoted, which is the multi-clause-set obtained by removing clauses from F which have no variables in common with V , and removing from the remaining clauses all literals where the underlying variable is not in V :

$$F[V](C) := \sum_{D \in F, \text{var}(D) \cap V \neq \emptyset, D \cap (V \cup \overline{V}) = C} F(D).$$

Here it is essential that $F[V]$ is a multi-clause-set, and if previously unequal clauses become equal, then accordingly the multiplicity is increased.

Example 2.10 $\{\{a\}, \{a, b\}, \{b\}, \{\bar{a}, \bar{b}\}\}[\{a\}] = \{2 * \{a\}, \{\bar{a}\}\}$.

Simple properties of this operation are (for multi-clause-sets F and $V, V' \subseteq \mathcal{VA}$):

1. $F[\emptyset] = \top$, $F[V] = F \setminus \{\perp\}$ for $\text{var}(F) \subseteq V$ (where $F \setminus F'$ for a clause-set F' means that all occurrences of clauses from F' are removed from F).
2. $(F[V])[V'] = F[V \cap V']$.
3. $c(F[V]) = \sum_{C \in F, \text{var}(C) \cap V \neq \emptyset} F(C)$ (the number of clauses containing some variable from V).

Clause-sets F, G are called **isomorphic**, if the variables of F can be renamed and potentially flipped so that F is turned into G . More precisely, an isomorphism α from F to G is a bijection $\alpha : \text{lit}(F) \rightarrow \text{lit}(G)$ which preserves complementation ($\alpha(\bar{x}) = \overline{\alpha(x)}$), and which maps the clauses of F precisely to the clauses of G ; when considering multi-clause-sets, then the isomorphism must preserve the multiplicity of clauses (that is, $G(\alpha(C)) = F(C)$ for all $C \in \mathcal{CL}$).

2.5. Degrees

For the number of occurrences of a literal $x \in \mathcal{LIT}$ in a (multi-)clause-set $F \in \mathcal{CLS}$ we write

$$\text{ld}_F(x) := \sum_{C \in F, x \in C} F(C) \in \mathbb{N}_0,$$

called the **literal-degree**; note $\text{ld}_F(x) = 0 \Leftrightarrow x \notin \bigcup F$. The **variable-degree** of a variable v is defined as $\text{vd}_F(v) := \text{ld}_F(v) + \text{ld}_F(\bar{v}) \in \mathbb{N}_0$; note $\text{vd}_F(v) = 0 \Leftrightarrow v \notin \text{var}(F)$ and $\text{vd}_F(v) = c(F[\{v\}])$. We remark that the literal-degree $\text{ld}_F(x)$ (for $x \in \mathcal{LIT}$) always has the index F , the (multi-)clause-set in which the literal-degree is counted, and thus there is little danger of confusion with the binary logarithm $\text{ld}(x)$ (for $x \in \mathbb{R}$), which does not have an index.

A (multi-)clause-set F is called **variable-regular** if all variables $v \in \text{var}(F)$ have the same degree, or, stronger, **literal-regular**, if all literals $x \in \text{lit}(F)$ have the same degree. A **singular variable** in a (multi-)clause-set F is a non-pure variable occurring in one sign only once, that is, $\min(\text{ld}_F(v), \text{ld}_F(\bar{v})) = 1$, while F is called **nonsingular** if it does not have singular variables, i.e., iff $\forall x \in \bigcup F : \text{ld}_F(\bar{x}) \neq 0 \Rightarrow \text{ld}_F(x) \geq 2$. The central concept for this report is the minimum degree of a variable in a clause-set:

Definition 2.11 We define the **minimum variable degree** $\mu\text{vd} : \mathcal{CLS} \rightarrow \mathbb{N} \cup \{+\infty\}$ (“*min-var-degree*” for short) as follows:

- For $F \in \mathcal{CLS}$ with $n(F) \neq 0$ we let $\mu\text{vd}(F) := \min_{v \in \text{var}(F)} \text{vd}_F(v) \in \mathbb{N}$.
- While for $n(F) = 0$ we set $\mu\text{vd}(F) := +\infty$.

For a class $\mathcal{C} \subseteq \mathcal{CLS}$ of (multi-)clause-sets let $\mu\text{vd}(\mathcal{C}) \in \mathbb{N}_0 \cup \{+\infty\}$ be the supremum of $\mu\text{vd}(F)$ for $F \in \mathcal{C}$ with $n(F) > 0$, where we set $\mu\text{vd}(\mathcal{C}) := 0$ if there is no such F (while otherwise we have $\mu\text{vd}(\mathcal{C}) \geq 1$).

We have $\mu\text{vd}(F) = +\infty$ iff $n(F) = 0$ (otherwise $\mu\text{vd}(F) \in \mathbb{N}$), and the motivation for this setting is, that so for all $F \in \mathcal{CLS}$ and $K \in \mathbb{R}$ holds $\mu\text{vd}(F) \geq K$ iff $\forall v \in \text{var}(F) : \text{vd}_F(v) \geq K$. By definition we have $\mu\text{vd}(\mathcal{C}) \leq \mu\text{vd}(\mathcal{C}')$ for $\mathcal{C} \subseteq \mathcal{C}' \subseteq \mathcal{CLS}$, and furthermore for $K \in \mathbb{R}_{>0}$ we have $\mu\text{vd}(\mathcal{C}) \geq K$ iff there is $F \in \mathcal{C}$ with $n(F) > 0$ and $\forall v \in \text{var}(F) : \text{vd}_F(v) \geq K$, while for $K \in \mathbb{R}_{\geq 0}$ we have $\mu\text{vd}(\mathcal{C}) \leq K$ iff for all $F \in \mathcal{C}$ with $n(F) > 0$ there is $v \in \text{var}(F)$ with $\text{vd}_F(v) \leq K$.

Example 2.12 For $F := \{2 * \{a, b\}, \{\bar{a}, b\}, \{\bar{b}, c\}\} \in \mathcal{CLS}$ we have:

- $\text{ld}_F(a) = 2, \text{ld}_F(\bar{a}) = 1, \text{ld}_F(b) = 3, \text{ld}_F(\bar{b}) = 1, \text{ld}_F(c) = 1, \text{ld}_F(\bar{c}) = 0$.
- $\text{vd}_F(a) = 3, \text{vd}_F(b) = 4, \text{vd}_F(c) = 1$.
- $\mu\text{vd}(F) = 1$.

Every full clause-set is variable-regular (but in general not literal-regular). Examples for $\mu\text{vd}(\mathcal{C})$ are:

- $\mu\text{vd}(\emptyset) = 0$.

- $\mu\text{vd}(\mathcal{CLS}) = +\infty$.
- $\mu\text{vd}(\{\top, \{\perp\}, \{\{v\}, \{\bar{v}\}\}, F) = 2$.

The simplest but relevant class of clause-sets for us is given by the $A(V)$, the unsatisfiable full clause-sets; these are the simplest unsatisfiable clause-sets:

Lemma 2.13 *For $n \in \mathbb{N}_0$ we have*

1. $n(A_n) = n$, $c(A_n) = 2^n$, $\delta(A_n) = 2^n - n$.
2. A_n is full and unsatisfiable, and thus $A_n \in \mathcal{UHLT}_{\delta=2^n-n}$.
3. A_n is literal-regular (thus variable-regular).
4. $\mu\text{vd}(A_n) = 2^n$ for $n \geq 1$.

Further properties of unsatisfiable full clause-sets one finds in Example 2.20, Lemma 3.10, Corollary 3.12, Lemmas 6.9, 6.10, Corollaries 6.11, 6.12, and Examples 9.2, 9.12. Properties of satisfiable full clause-sets are found in Example 10.7.

2.6. Autarkies

Besides algorithmic considerations, which were always present since the introduction of the notion of an “autarky” in [41], also a kind of a “combinatorial SAT theory” has been developed around this notion of generalised satisfying assignments. A general overview is given in [11], with recent additions and generalisations in [16]. An **autarky** (see [11, Section 11.8]) for a clause-set $F \in \mathcal{CLS}$ is a partial assignment $\varphi \in \mathcal{PASS}$ which satisfies every clause $C \in F$ it touches, i.e., for all $C \in F$ with $\text{var}(\varphi) \cap \text{var}(C) \neq \emptyset$ holds $\varphi * \{C\} = \top$; equivalently, for all $C \in F$ holds $C \cap \varphi^{-1}(0) \neq \emptyset \Rightarrow C \cap \varphi^{-1}(1) \neq \emptyset$. The simplest examples for autarkies are as follows:

Example 2.14 *The empty partial assignment $\langle \rangle$ is an autarky for every $F \in \mathcal{CLS}$ (no clause is touched), and more generally all $\varphi \in \mathcal{PASS}$ with $\text{var}(\varphi) \cap \text{var}(F) = \emptyset$ are autarkies for F , the **trivial autarkies**. On the other end of the spectrum every satisfying assignment for F (i.e., $\varphi * F = \top$) is an autarky for F (every clause is satisfied). A literal $x \in \mathcal{LIT}$ is a pure literal for F iff $\langle x \rightarrow 1 \rangle$ is an autarky for F .*

If φ is an autarky for F , then $\varphi * F \subseteq F$ holds, and thus $\varphi * F$ is satisfiability-equivalent to F . Autarkies mark redundancies, and the corresponding notion of clause-sets without such redundancies was introduced in [10], namely a clause-set F is **lean** if there is no non-trivial autarky for F , and the set of all lean clause-sets is denoted by $\mathcal{LEAN} \subset \mathcal{USAT} \cup \{\top\}$. The class $\underline{\mathcal{LEAN}}$ of lean multi-clause-sets is invariant under multiplicities.

Example 2.15 *Some simple examples:*

1. $\top, \{\perp\}, \{\{v\}, \{\bar{v}\}\}, \{\{v\}, \{\bar{v}\}, \{w\}, \{\bar{w}\}\} \in \mathcal{LEAN}$.
2. If $F, F' \in \mathcal{LEAN}$, then $F \cup F' \in \mathcal{LEAN}$.
3. If $F \in \mathcal{LEAN}$ and $F' \in \mathcal{CLS}$ with $\text{var}(F') \subseteq \text{var}(F)$, then $F \cup F' \in \mathcal{LEAN}$.
4. $\{\{v\}, \{\bar{v}\}, \{w\}\} \notin \mathcal{LEAN}$.

A weakening is the notion of a **matching-lean** clause-set F (introduced in [13, Section 7]; see [11, Section 11.11] for an overview), which has no non-trivial **matching autarky**, which are special autarkies given by a matching condition: for every clause touched, a satisfied literal with unique underlying variable must be selectable; the class of all matching-lean clause-sets is denoted by $\mathcal{MLEAN} \supset \mathcal{LEAN}$.

Example 2.16 *A clause-set $F \in \mathcal{CLS}$ is matching-lean ($F \in \mathcal{MLEAN}$) iff for all $F' \subset F$ holds $\delta(F') < \delta(F)$ ([13, Theorem 7.5]). Thus for every matching-lean multi-clause-set $F \neq \top$ we have $\delta(F) \geq 1$ ([13], generalising [5]).*

$F := \{\{1, 3\}, \{2, -3\}, \{3\}, \{-3\}\}$ has the matching autarky $\langle 1 \rightarrow 1, 2 \rightarrow 1 \rangle$, while for $F' := F \cup \{\{1, 2\}\}$ we have $F' \in \mathcal{MLEAN}$. Note $\delta(F) = 1 = \delta(\{\{3\}, \{-3\}\})$, while $\delta(F') = 2$.

It is decidable in polynomial time whether $F \in \mathcal{MLEAN}$ holds (which follows for example by the characterisation of \mathcal{MLEAN} via the surplus below). The class $\mathcal{MLEAN} \supset \mathcal{LEAN}$ of matching-lean multi-clause-sets is not invariant under multiplicities: For a multi-clause-set F holds $\text{cls}(F) \in \mathcal{MLEAN} \Rightarrow F \in \mathcal{MLEAN}$, but not the other way around:

Example 2.17 *$\{\{v\}\} \notin \mathcal{MLEAN}$, but $\{2 * \{v\}\} \in \mathcal{MLEAN}$, and more generally $\{\{x_1, \dots, x_n\}\} \notin \mathcal{MLEAN}$ for $n \geq 1$, while $\{(n+1) * \{x_1, \dots, x_n\}\} \in \mathcal{MLEAN}$. Indeed it is easy to see that for every $F \in \mathcal{CLS}$ there is $F' \in \mathcal{CLS}$ with $\text{cls}(F') = F$ and $F' \in \mathcal{MLEAN}$.*

The process of applying autarkies as long as possible to a clause-set $F \in \mathcal{CLS}$ is confluent, yielding the **lean kernel** of F (the largest lean sub-clause-set of F , that is, $\bigcup\{F' \subseteq F : F' \in \mathcal{LEAN}\}$; see [10, Section 3]). Computation of the lean kernel is NP-hard, since the lean kernel of satisfiable clause-sets is \top . But the **matching-lean kernel** of F (the largest matching-lean sub-clause-set of F , that is, $\bigcup\{F' \subseteq F : F' \in \mathcal{MLEAN}\}$; see [13, Section 3]), now obtained by applying *matching* autarkies as long as possible, which again is a confluent process, is computable in polynomial time. Note that a clause-set F is lean resp. matching lean iff the lean resp. matching-lean kernel is F itself. Due to the polytime computability of the matching-lean kernel, which is a sub-clause-set obtained by removing clauses redundant in a strong sense, “w.l.o.g.” for the purpose of SAT-decision one might consider the inputs as matching-lean.

Example 2.18 *For inputs $F \in \mathcal{MLEAN}$ by [26, Theorem 4] we have SAT-decision in time $O(2^{\delta(F)} \cdot n(F)^3)$ (see [16] for generalisations), and thus SAT-decision for inputs $F \in \mathcal{MLEAN}$ is fixed-parameter tractability (fpt) in the parameter $\delta(F)$.*

We note here (though we won't use it in this report), that for inputs $F \in \mathcal{MLEAN}_{\delta=k}$ the computation of the lean kernel can be done in polynomial time for fixed k ([16, Theorem 10.3]; this computational problem appears not to be fpt).

2.7. The surplus

A multi-clause-set $F \neq \top$ is matching lean iff for the **surplus** we have $\sigma(F) \geq 1$ ([13, Lemma 7.7]), which is defined as follows (see [16, Subsection 11.1] for more information; in [26] a clause-set has “ q -expansion” iff $\sigma(F) \geq q$):

Definition 2.19 *For a multi-clause-set F let $\sigma(F) \in \mathbb{Z}$ be defined as the minimum of $\delta(F[V])$ (recall Definition 2.9) over all $\emptyset \neq V \subseteq \text{var}(F)$ if $n(F) > 0$, while $\sigma(F) := 0$ in case of $n(F) = 0$.*

So for $F \in \underline{\mathcal{CLS}}$, $n(F) > 0$, and $K \in \mathbb{R}$ holds $\sigma(F) \geq K$ iff for all $\emptyset \neq V \subseteq \text{var}(F)$ the number of clauses of F containing some variable of V is at least $K + |V|$. The special case of $n(F) = 0$ is handled so that we always have $-n(F) \leq \sigma(F) \leq c(F)$. Note that for $\emptyset \neq V \subseteq \text{var}(F)$ we have

$$\delta(F[V]) = c(F[V]) - |V| = \left(\sum_{C \in F, \text{var}(C) \cap V \neq \emptyset} F(C) \right) - |V|.$$

The surplus is computable in polynomial time. Some basic properties of the surplus are (for multi-clause-sets F):

1. $\sigma(F)$ is independent of $F(\perp)$ (the number of occurrences of the empty clause).
2. $\sigma(F) \leq \delta(F[\text{var}(F)]) = \delta(F \setminus \{\perp\}) \leq \delta(F) \leq c(F)$.
3. For every $\emptyset \subset V \subseteq \text{var}(F)$ holds $\sigma(F[V]) \geq \sigma(F)$.
4. For $F' \leq F$ with $\text{var}(F') = \text{var}(F)$ we have $\sigma(F') \leq \sigma(F)$.
5. $-n(F) \leq \sigma(F)$, and for $n(F) > 0$ holds $1 - n(F) \leq \sigma(F) \leq c(F) - 1$.

Example 2.20 $\sigma(A_0) = 0$, while $\sigma(A_n) = 2^n - n = \delta(A_n)$ for $n \in \mathbb{N}$ (since every variable occurs in every clause). If we take $F \in \mathcal{CLS}$ and some $v \in \mathcal{VA} \setminus \text{var}(F)$, then $\sigma(F \cup \{\{v\}\}) \leq 0$.

The basic intuition is that $\sigma(F)$ gives us the “easiest subinstances” of F , “easiest” in the sense of deficiency, “subinstance” in the sense of restriction. The theory of the surplus of clause-sets is further developed in this report in Section 9, especially Subsection 9.1, and Section 10, especially Subsection 10.3.

3. Minimally unsatisfiable clause-sets

In this section we review minimally unsatisfiable clause-sets; see [11] for an overview, while [17, 19] contain recent developments. First the basic definitions and examples are given in Subsection 3.1. In Subsection 3.2 we consider in some detail the fundamental process of “saturation”, which is about adding “missing literal occurrences” to minimally unsatisfiable clause-sets. Saturation repairs the problem that splitting of $F \in \mathcal{MU}$ into $\langle v \rightarrow 0 \rangle * F$ and $\langle v \rightarrow 1 \rangle * F$ may destroy minimal unsatisfiability, i.e., $\langle v \rightarrow 0 \rangle * F \notin \mathcal{MU}$ or $\langle v \rightarrow 1 \rangle * F \notin \mathcal{MU}$ might hold, due to some clauses missed to be deleted by the partial assignment, and this process of splitting is considered in Subsection 3.3.

3.1. \mathcal{MU} and subclasses

An unsatisfiable clause-set F is called **minimally unsatisfiable**, if for every clause $C \in F$ the clause-set $F \setminus \{C\}$ is satisfiable, and the set of minimally unsatisfiable clause-sets is denoted by $\mathcal{MU} \subset \mathcal{USAT}$. A clause-set $F \in \mathcal{MU}$ is called **saturated**, if replacing any $C \in F$ by any super-clause $C' \supset C$ yields a satisfiable clause-set, and the set of saturated minimally unsatisfiable clause-sets is denoted by $\mathcal{SMU} \subset \mathcal{MU}$.

Example 3.1 The simplest element of $\mathcal{USAT} \setminus \mathcal{MU}$ is $\{\perp, \{1\}\}$, while the simplest element of $\mathcal{MU} \setminus \mathcal{SMU}$ is $\{\{1, 2\}, \{-1\}, \{-2\}\}$ (see Example 3.6 for a “saturation”).

Unsatisfiable hitting clause-sets fulfil $\mathcal{UHIT} \subset \mathcal{SMU}$ (see [19, Lemma 2] for the proof). The subsets of nonsingular elements (i.e., there is no literal occurring only once) are denoted by $\mathcal{MU}' \subset \mathcal{MU}$, $\mathcal{SMU}' \subset \mathcal{SMU}$, and $\mathcal{UHIT}' \subset \mathcal{UHIT}$.

Example 3.2 By [6] holds $\mathcal{MU}'_{\delta=1} = \mathcal{SMU}'_{\delta=1} = \mathcal{UHIT}'_{\delta=1} = \{\{\perp\}\}$, while for the characterisation of $\mathcal{MU}_{\delta=1} \supset \mathcal{SMU}_{\delta=1} = \mathcal{UHIT}_{\delta=1}$ see also [5, 7]. As shown in [6], for $F \in \mathcal{MU}_{\delta=1}$ with $n(F) > 0$ holds $\mu\text{vd}(F) = 2$.

We consider the “reasons” for unsatisfiability as given by the elements of $\mathcal{MU}_{\delta=1}$ as “noise”, only “masking” the pure contradiction of the only element of $\mathcal{MU}'_{\delta=1} = \{\{\perp\}\}$ (in Section 5 the elimination of singular variables will be discussed). “Real reasoning” starts with deficiency 2:

Example 3.3 By [8], the elements of $\mathcal{MU}'_{\delta=2}$ are up to isomorphism precisely the \mathcal{F}_n for $n \geq 2$, where

$$\mathcal{F}_n := \{\{1, \dots, n\}, \{-1, \dots, -n\}, \{-1, 2\}, \dots, \{-(n-1), n\}, \{-n, 1\}\}.$$

All \mathcal{F}_n are literal-regular, with $\mu\text{vd}(\mathcal{F}_n) = 4$. It is easy to see that all \mathcal{F}_n are saturated, and thus $\mathcal{MU}'_{\delta=2} = \mathcal{SMU}'_{\delta=2}$. The only hitting clause-sets amongst the \mathcal{F}_n are for $n = 2, 3$, and thus up to isomorphism the elements of $\mathcal{UHIT}'_{\delta=2}$ are $\mathcal{F}_2, \mathcal{F}_3$, with $\mathcal{F}_2 = A_2$ and

$$\mathcal{F}_3 = \{\{1, 2, 3\}, \{-1, -2, -3\}, \{-1, 2\}, \{-2, 3\}, \{-3, 1\}\}.$$

We have $\sigma(\mathcal{F}_n) = 2 = \delta(\mathcal{F}_n)$, since any $m \leq n$ variables occur at least in m different binary clauses plus in the two “long clauses” (this is also obtained from the later Lemma 10.10, Part 1). Further properties of the \mathcal{F}_n we have in Examples 3.14, 6.2, 8.2, 9.2. See Section 7 in [19] for more information.

As shown in [19, Theorem 74], for every $F \in \mathcal{MU}_{\delta=2}$ there is a unique $n \geq 2$ such that \mathcal{F}_n “embeds” into F , and this n is called the “nonsingularity type” of F . So for $\mathcal{MU}_{\delta=2}$ we have identified the (in a sense) unique reason of unsatisfiability, the (possibly hidden) presence of a cycle $v_1 \rightarrow \dots \rightarrow v_n \rightarrow v_1$ together with the assertions, that one v_i must be true and one must be false (only the n is unique in general, not the v_i). We will come back to the theme of classifying $\mathcal{MU}_{\delta=k}$ in the Conclusion, Subsection 15.5.

By definition, \mathcal{MU} disallows multiplicities (since a duplicated clause is the trivial logical redundancy), and this also holds for the subclasses \mathcal{SMU} and \mathcal{UHIT} (as well as for all other subclasses of \mathcal{MU} considered here). A fundamental fact is $\delta(F) \geq 1$ for all $F \in \mathcal{MU}$ (note that every minimally unsatisfiable clause-set is lean), which motivates the investigation of the layers $\mathcal{MU}_{\delta=1}, \mathcal{MU}_{\delta=2}, \dots$. Special elements of \mathcal{UHIT} are the $A(V)$ for finite sets V of variables (recall Lemma 2.13), which are the minimally unsatisfiable clause-sets with maximal deficiency for a given number of variables, as we will see in Corollary 6.11.

Finally, the two main quantities studied in this report need dedicated names:

Definition 3.4 For $k \in \mathbb{N}$ let

$$\begin{aligned} \mathbf{VDM}(\mathbf{k}) &:= \mu\text{vd}(\mathcal{MU}_{\delta=k}) \in \mathbb{N} \\ \mathbf{VDH}(\mathbf{k}) &:= \mu\text{vd}(\mathcal{UHIT}_{\delta=k}) \in \mathbb{N} \end{aligned}$$

(these numbers are finite due to $\mathbf{VDM}(\mathbf{k}) \leq 2k$ by [7, Lemma C.2]).

By definition holds $\mathbf{VDH} \leq \mathbf{VDM}$.

3.2. Saturation

We recall the fact ([23, 14]) that every minimally unsatisfiable clause-set $F \in \mathcal{MU}$ can be **saturated**, i.e., by adding literal occurrences to F we can obtain $F' \in \mathcal{SMU}$ with $\text{var}(F') = \text{var}(F)$ (there is then a bijection $\alpha : F \rightarrow F'$ with

$C \subseteq \alpha(C)$ for all $C \in F$. Since we consider saturation in many situations, we introduce some special notations for it from [19, Subsection 2.2]. First we introduce the notation $S(F, C, x)$ for adding a literal x to a clause C in a clause-set F :

Definition 3.5 ([19]) *The operation (adding literal x to clause C in F)*

$$\mathbf{S}(F, C, x) := (F \setminus \{C\}) \cup (C \cup \{x\}) \in \mathcal{CLS}$$

is defined if $F \in \mathcal{CLS}$, $C \in F$, and x is a literal with $\text{var}(x) \in \text{var}(F) \setminus \text{var}(C)$.

Some technical remarks:

1. $\text{var}(S(F, C, x)) = \text{var}(F)$.
2. If $C \cup \{x\} \notin F$, then $c(S(F, C, x)) = c(F)$, and thus also $\delta(S(F, C, x)) = \delta(F)$.
3. For $F \in \mathcal{MU}$ we have:
 - (a) $S(F, C, x) \in \mathcal{MU}$ iff $S(F, C, x)$ is unsatisfiable (since all what happened is that a clause has been weakened, i.e., extended).
 - (b) If $S(F, C, x) \in \mathcal{MU}$, then $c(S(F, C, x)) = c(F)$ (no subsumption here).
 - (c) F is saturated iff there are no C, x such that $S(F, C, x) \in \mathcal{USAT}$.

Example 3.6 For $F := \{\{a, b\}, \{\bar{a}\}, \{\bar{b}\}\} \in \mathcal{MU} \setminus \mathcal{SMU}$ we have $S(F, \{\bar{a}\}, b) = \{\{a, b\}, \{\bar{a}, b\}, \{\bar{b}\}\} \in \mathcal{SMU}$.

A “saturation” of a minimally unsatisfiable clause-set is obtained by adding literals to clauses as long as possible while maintaining unsatisfiability (which is the same as maintaining minimal unsatisfiability):

Definition 3.7 ([19]) *A **saturation** $F' \in \mathcal{SMU}$ of $F \in \mathcal{MU}$ is obtained by a **saturation sequence** $F = F_0, \dots, F_m = F'$, $m \in \mathbb{N}_0$, such that*

- (i) for $0 \leq i < m$ there are C_i, x_i with $F_{i+1} = S(F_i, C_i, x_i)$,
- (ii) for all $1 \leq i \leq m$ we have $F_i \in \mathcal{USAT}$,
- (iii) the sequence cannot be extended (without violating conditions (i) or (ii)).

Note that $n(F') = n(F)$ and $c(F') = c(F)$ holds (and thus $\delta(F') = \delta(F)$). If we drop requirement (iii), then we speak of a **partial saturation sequence**, while $F' \in \mathcal{MU}$ is a **partial saturation** of $F \in \mathcal{MU}$.

Some technical remarks:

1. $F' \in \mathcal{MU}$ is a partial saturation of $F \in \mathcal{MU}$ iff
 - $\text{var}(F') = \text{var}(F)$
 - there is a bijection $\alpha : F \rightarrow F'$ such that for all $C \in F$ we have $C \subseteq \alpha(C)$.
2. For a partial saturation sequence F_0, \dots, F_m we have $\ell(F_m) = \ell(F_0) + m$.
3. F' is a saturation of $F \in \mathcal{MU}$ iff F' is a partial saturation of F with $F' \in \mathcal{SMU}$.

Example 3.8 A saturation sequence for $F := \{\{a, b, c\}, \{\bar{a}\}, \{\bar{b}\}, \{\bar{c}\}\}$ with $m = 3$ is obtained by adding literals b, c to clause $\{\bar{a}\}$, and adding literal c to clause $\{\bar{b}\}$.

We can perform a partial saturation $F \rightsquigarrow S(F, C, x)$ iff F without C implies (logically) $C \cup \{\bar{x}\}$ (note that $C \cup \{x\}, C \cup \{\bar{x}\}$ implies C):

Lemma 3.9 *Consider $F \in \mathcal{MU}$, $C \in F$, and a literal x with $\text{var}(x) \in \text{var}(F) \setminus \text{var}(C)$. Then $S(F, C, x)$ is a partial saturation of F if and only if $F \setminus \{C\} \models C \cup \{\bar{x}\}$.*

Proof: First assume that $S(F, C, x)$ is a partial saturation of F , but $F \setminus \{C\} \not\models C \cup \{\bar{x}\}$. So there is a partial assignment φ with $\varphi * (F \setminus \{C\}) = \top$ but $\varphi * \{C \cup \{\bar{x}\}\} = \{\perp\}$ (whence $\varphi(x) = 1$). But then we have $\varphi * S(F, C, x) = \top$. Reversely assume $F \setminus \{C\} \models C \cup \{\bar{x}\}$, but that $S(F, C, x)$ is not a partial saturation of F . So $S(F, C, x)$ has a satisfying assignment φ ; due to $F \in \mathcal{USAT}$ we have $\varphi(x) = 1$ and $\varphi * \{C\} = \{\perp\}$. But this yields $\varphi * (F \setminus \{C\}) = \top$ and $\varphi * \{C \cup \{\bar{x}\}\} = \{\perp\}$. \square

See Lemma 6.5, Part 7, for another characterisation of partial saturations. The dual notion of “saturated” is “marginal”: $F \in \mathcal{MU}$ is **marginal** iff replacing any clause by a strict subclause yields a clause-set not in \mathcal{MU} . The decision “ F marginal minimally unsatisfiable?” for inputs $F \in \mathcal{CLS}$ is D^P -complete ([86, Theorem 2]). By [87, Theorem 8] however this decision is easy for inputs $F \in \mathcal{SMU}$, namely there is the following characterisation of minimally unsatisfiable clause-sets which are marginal and saturated at the same time:

Lemma 3.10 ([87]) *$F \in \mathcal{MU}$ is both marginal and saturated iff $F = A(\text{var}(F))$.*

Thus $F \in \mathcal{SMU}$ is marginal iff $F = A(\text{var}(F))$; so if $F \in \mathcal{SMU}$ is not full, then there is a literal occurrence which can be removed without destroying minimal unsatisfiability, that is, there exists a clause $C \in F$ and $x \in C$ such that $F' := (F \setminus \{C\}) \cup \{C \setminus \{x\}\} \in \mathcal{MU}$ (note that $F' \in \mathcal{USAT}$ in any case, but minimality in general is not maintained). And for inputs $F \in \mathcal{SMU}$ the existence of such C, x is decidable in linear time (namely they exist iff F is not full). But finding such C, x should be hard in general, that is, the decision problem, whether a concrete literal can be removed, even for inputs $F \in \mathcal{SMU}$ should be NP-complete:

Question 3.11 *Is the promise problem for input $F \in \mathcal{SMU}$, $C \in F$, $x \in C$, whether “ $(F \setminus \{C\}) \cup \{C \setminus \{x\}\} \in \mathcal{MU}$?”, NP-complete? (That is, is there a polytime computation $G \in \mathcal{CLS} \rightsquigarrow (F, C, x) \in \mathcal{SMU} \times \mathcal{CL} \times \mathcal{LIT}$, with $x \in C \in F$, such that $G \in \mathcal{SAT} \Leftrightarrow (F \setminus \{C\}) \cup \{C \setminus \{x\}\} \in \mathcal{MU}$?)*

And is the promise problem for input $F \in \mathcal{MU}$, whether F is marginal, NP-complete? (That is, is there a polytime computation $G \in \mathcal{CLS} \rightsquigarrow F \in \mathcal{MU}$, such that $G \in \mathcal{SAT}$ iff F is marginal?)

Back to saturation: precisely all saturated clause-sets except the $A(V)$ are obtained as non-trivial saturations of some minimally unsatisfiable clause-set:

Corollary 3.12 *Consider $F \in \mathcal{SMU}$.*

1. F is trivially the saturation of itself.
2. If $F = A(\text{var}(F))$, then this is also the only possibility for F being a saturation, that is, if F is the saturation of some $F' \in \mathcal{MU}$, then we have $F' = F$.
3. Otherwise F is a saturation of some clause-set other than itself, that is, if $F \neq A(\text{var}(F))$, then there is some $F' \in \mathcal{MU}$ with $F' \neq F$ such that F is a saturation of F' .

Proof: Part 1 is trivial. For Part 2 assume that $F = A(\text{var}(F))$, and we have $F = S(F', C, x)$ for some $F' \in \mathcal{MU}$: But since F is marginal, F' is not minimally unsatisfiable. Finally for Part 3 note, that if $F \neq A(\text{var}(F))$, then by Lemma 3.10 F is not marginal, and thus there is $C \in F$ and $x \in C$ such that for $C' := C \setminus \{x\}$ and $F' := (F \setminus \{C\}) \cup \{C'\}$ we have $F' \in \mathcal{MU}$. Now $F = S(F', C', x)$. \square

As discussed above, we expect the decision, whether for inputs $F \in \mathcal{SMU}$, $C \in F$ and $x \in C$ we have $(F \setminus \{C\}) \cup \{C'\} \in \mathcal{MU}$, to be NP-complete. But this decision is easy for $F \in \mathcal{UHIT}$, namely iff no “subsumption resolution” with another clause containing \bar{x} can be performed, that is, there is no $D \in F$ with $\bar{x} \in D$ and $C \setminus \{x\} \subseteq D$:

Lemma 3.13 *Consider $F \in \mathcal{UHIT}$, $C \in F$ and $x \in C$. Let $C' := C \setminus \{x\}$, and let $F' := (F \setminus \{C\}) \cup \{C'\}$. Then we have $F' \in \mathcal{MU}$ iff there is no $D \in F \setminus \{C\}$ with $C' \subset D$.*

Proof: If there is $D \in F \setminus \{C\}$ with $C' \subset D$, then $F' \notin \mathcal{MU}$. So assume there is no such D . Assume $F' \notin \mathcal{MU}$. Thus there is $E \in F'$ with $F' \setminus \{E\}$ unsatisfiable. We must have $E \neq C'$, since otherwise $F \setminus \{C\}$ would be unsatisfiable. Since F is hitting, E clashes with every clause of $F' \setminus \{C'\}$. It follows that $C' \subset E$ must hold (since the falsifying assignments for E are disjoint with those for any clause in $F' \setminus \{C'\}$), contradicting the minimal unsatisfiability of F . \square

Some examples on removable literal occurrences illustrate Lemma 3.13:

Example 3.14 *For $F := \{\{a, b\}, \{\bar{a}, b\}, \{\bar{b}\}\} \in \mathcal{UHIT}_{\delta=1}$ we can exactly remove one of the two literal-occurrences of b and still obtain a clause-set in \mathcal{MU} (of course not in \mathcal{UHIT} anymore; the resulting clause-sets are in fact marginally minimally unsatisfiable). For $\mathcal{F}_2 = A_2 = \{\{1, 2\}, \{-1, -2\}, \{-1, 2\}, \{-2, 1\}\} \in \mathcal{UHIT}'_{\delta=2}$ we can not remove any literal occurrence without leaving \mathcal{MU} (i.e., \mathcal{F}_2 is marginal).*

3.3. Splitting

An important (equivalent) characterisation of saturation for $F \in \mathcal{CLS}$, as shown in [17], is that splitting on any variable v will yield minimally unsatisfiable clause-sets $\langle v \rightarrow 0 \rangle * F$, $\langle v \rightarrow 1 \rangle * F$. This enables induction on the number of variables, which is a central method for this report; see Lemma 8.1 for the basic example. We also have that if for some variable both splitting results are minimally unsatisfiable resp. saturated, then this can be lifted to the original clause-set, provided that no contraction took place (different clauses became equal as a result of the assignment). These basic facts are collected in the following lemma.

Lemma 3.15 *For all clause-sets $F \in \mathcal{CLS}$ we have:*

1. $F \in \mathcal{SMU}$ iff for all $v \in \text{var}(F)$ and all $\varepsilon \in \{0, 1\}$ we have $\langle v \rightarrow \varepsilon \rangle * F \in \mathcal{MU}$.
2. If for some variable v holds $\langle v \rightarrow 0 \rangle * F \in \mathcal{MU}$ and $\langle v \rightarrow 1 \rangle * F \in \mathcal{MU}$, and if for all $C \in F$ with $v \in \text{var}(C)$ we have $C \setminus \{v, \bar{v}\} \notin F$, then $F \in \mathcal{MU}$.
3. If for some variable v holds $\langle v \rightarrow 0 \rangle * F \in \mathcal{SMU}$ and $\langle v \rightarrow 1 \rangle * F \in \mathcal{SMU}$, and if for all $C \in F$ with $v \in \text{var}(C)$ we have $C \setminus \{v, \bar{v}\} \notin F$, then $F \in \mathcal{SMU}$.

Proof: Part 1 is Corollary 5.3 in [17]. For Part 2 assume $F \notin \mathcal{MU}$; thus there is $C \in F$ with $F' := F \setminus \{C\} \in \mathcal{USAT}$. We consider three cases:

1. $v \notin \text{var}(C)$: Due to the assumption on subsumption-freeness we have $C \cup \{v\} \notin F'$. Now $C \in \langle v \rightarrow 0 \rangle * F$, while $(\langle v \rightarrow 0 \rangle * F) \setminus \{C\} = \langle v \rightarrow 0 \rangle * F' \in \mathcal{USAT}$, contradicting $\langle v \rightarrow 0 \rangle * F \in \mathcal{MU}$.
2. $v \in C$: By assumption holds $C' := C \setminus \{v\} \notin F'$. Now $C' \in \langle v \rightarrow 0 \rangle * F$, while $(\langle v \rightarrow 0 \rangle * F) \setminus \{C'\} = \langle v \rightarrow 0 \rangle * F' \in \mathcal{USAT}$, contradicting $\langle v \rightarrow 0 \rangle * F \in \mathcal{MU}$.
3. $\bar{v} \in C$: By assumption holds $C' := C \setminus \{\bar{v}\} \notin F'$. Now $C' \in \langle v \rightarrow 1 \rangle * F$, while $(\langle v \rightarrow 1 \rangle * F) \setminus \{C'\} = \langle v \rightarrow 1 \rangle * F' \in \mathcal{USAT}$, contradicting $\langle v \rightarrow 1 \rangle * F \in \mathcal{MU}$.

Finally consider Part 3. By Part 2 we already know that $F \in \mathcal{MU}$ holds. Assume that $F \notin \mathcal{SMU}$; thus there is $C \in F$ and a literal x with $F' := S(F, C, x) \in \mathcal{USAT}$. So by Lemma 3.9 we have $F \setminus \{C\} \models C' := C \cup \{\bar{x}\}$. There exists at least one $\varepsilon \in \{0, 1\}$ with $\langle v \rightarrow \varepsilon \rangle * \{C'\} \neq \top$, and then $\langle v \rightarrow \varepsilon \rangle * (F \setminus \{C\}) \models \langle v \rightarrow \varepsilon \rangle * C'$. If $\text{var}(x) = v$, then this contradicts minimal unsatisfiability of $\langle v \rightarrow \varepsilon \rangle * F$. And if $\text{var}(x) \neq v$, then $\langle v \rightarrow \varepsilon \rangle * F \setminus \langle v \rightarrow \varepsilon \rangle * \{C\} \models (\langle v \rightarrow \varepsilon \rangle * C) \cup \{\bar{x}\}$, contradicting saturatedness of $\langle v \rightarrow \varepsilon \rangle * F$ by Lemma 3.9. \square

The additional assumption $C \setminus \{v, \bar{v}\} \notin F$ for Parts 2, 3 is equivalent to saying that no contraction takes place when applying $\langle v \rightarrow 0 \rangle, \langle v \rightarrow 1 \rangle$. An alternative way of stating this would be to use multi-clause-sets, since then no contractions would be performed, and the doubled clauses would destroy minimal unsatisfiability. In [14, Lemma 1] (and in the underlying report [88, Lemma 2.1]) that additional assumption for Parts 2, 3 is missing by mistake:

Example 3.16 *An example for $\langle v \rightarrow 0 \rangle * F \in \mathcal{UHLT}$ and $\langle v \rightarrow 1 \rangle * F \in \mathcal{UHLT}$, but $F \notin \mathcal{MU}$ is trivially given by $F := \{\perp, \{v\}, \{\bar{v}\}\}$, where for both $\varepsilon \in \{0, 1\}$ holds $\langle v \rightarrow \varepsilon \rangle * F = \{\perp\}$ (note that F is a set — for a multi-clause-set F the contraction would not occur).*

4. Variable-minimal unsatisfiability

In [15] the generalisation of minimal unsatisfiability to “variable-minimal unsatisfiability” has been introduced, and the class of all such clause-sets is denoted by \mathcal{VMU} , the set of clause-sets $F \in \mathcal{USAT}$ such that for every $F' \subseteq F$ with $F' \in \mathcal{USAT}$ holds $\text{var}(F') = \text{var}(F)$; see [89] for related algorithms for computing $\text{var}(F')$ for some $F' \subseteq F$ with $F' \in \mathcal{MU}$ for input $F \in \mathcal{USAT}$. The corresponding class $\underline{\mathcal{VMU}}$ of multi-clause-sets is invariant under multiplicities. Thus, as with \mathcal{LEAN} (and different from \mathcal{MU}), regarding variable-minimal unsatisfiability w.l.o.g. multi-clause-sets can be cast down to clause-sets. The basic (trivial) characterisation of \mathcal{VMU} is:

Lemma 4.1 *For $F \in \mathcal{CLS}$ holds $F \in \mathcal{VMU}$ if and only if $F \in \mathcal{USAT}$ and for all $v \in \text{var}(F)$ holds $\{C \in F : v \notin \text{var}(C)\} \in \mathcal{SAT}$.*

By definition we have $\mathcal{MU} \subset \mathcal{VMU}$, moreover, as shown in [15, Lemma 6], for every deficiency $k \geq 2$ we have $\mathcal{MU}_{\delta=k} \subset \mathcal{VMU}_{\delta=k}$ (for example, for every $F \in \mathcal{MU}_{\delta=k}$, $k \in \mathbb{N}$, and every non-full clause $C \in F$, i.e., $\text{var}(C) \subset \text{var}(F)$, we can add to F a full clause subsumed by C , obtaining $F' \in \mathcal{VMU}_{\delta=k+1} \setminus \mathcal{MU}_{\delta=k+1}$).

In [15] there is the false statement “ $\mathcal{VMU} \not\subseteq \mathcal{LEAN}$ ”, based on the following erroneous example:

Example 4.2 [15, Page 266] gives the example $F_4 := \{\{a\}, \{\bar{b}\}, \{\bar{a}, b\}, \{a, \bar{b}\}\}$ with the assertion “ $F_4 \in \mathcal{VMU} \setminus \mathcal{LEAN}$ ”. Obviously we have $F_4 \in \mathcal{VMU}$, but we also have $F_4 \in \mathcal{LEAN}$. Using the characterisation from [10] (which is the only

characterisation used in [15]), that $F \in \mathcal{LEAN}$ holds iff every clause of F can be used in a tree-resolution refutation of F , we see this as follows: the sole subset of F_4 in \mathcal{MU} is $\{\{a\}, \{\bar{b}\}, \{\bar{a}, b\}\}$, while the clause $\{a, \bar{b}\}$ can(!) also be used in a tree-resolution refutation — it is obviously superfluous, but nevertheless there is a tree-resolution refutation using it, namely via $(\{a\} \diamond \{\bar{a}, b\}) \diamond \{a, \bar{b}\} = \{a\}$.

Based on the characterisation of lean clause-sets via autarkies, it is easy to show that \mathcal{VMU} consists of special lean clause-sets (thus Figure 1 in [15] needs to be corrected, showing instead that \mathcal{LEAN} is indeed a superclass of \mathcal{VMU}):

Lemma 4.3 $\mathcal{VMU} \subset \mathcal{LEAN} \setminus \{\top\}$.

Proof: While in [15] the characterisation of \mathcal{LEAN} via variables usable in resolution refutation was (only) used, here we need to use the equivalent characterisation via autarkies, shown in [10, Theorem 3.16], and used as our definition in Subsection 2.6, namely that for $F \in \mathcal{CLS}$ holds $F \in \mathcal{LEAN}$ iff there is no autarky φ for F with $\text{var}(\varphi) \cap \text{var}(F) \neq \emptyset$: If we had such an autarky for $F \in \mathcal{VMU}$, then $\varphi * F \in \mathcal{USAT}$ with $\varphi * F \subset F$ and $\text{var}(\varphi * F) \subseteq \text{var}(F) \setminus \text{var}(\varphi)$, contradicting $F \in \mathcal{VMU}$. That indeed \mathcal{VMU} is a strict subset of $\mathcal{LEAN} \setminus \{\top\}$, can for example be seen by [10, Lemma 3.2], which shows that if we extend a minimally unsatisfiable clause-set via Extended Resolution, then we always stay in \mathcal{LEAN} , while just adding new clauses, each containing a new variable, and thus leaving \mathcal{VMU} ; another example is clause-set F_3 from [15, Page 266]. \square

Thus it follows $\mathcal{VMU}_{\delta=1} = \mathcal{MU}_{\delta=1}$ (shown in [15, Lemma 6]), since by [10, Corollary 5.7] holds $\mathcal{LEAN}_{\delta=1} \cap \mathcal{USAT} = \mathcal{MU}_{\delta=1}$.

For $F \in \mathcal{VMU}$ obviously there is some $F' \subseteq F$ with $\text{var}(F') = \text{var}(F)$ and $F' \in \mathcal{MU}$; [15, Lemma 5] asserts the converse, but this is false, as the following simple example shows:

Example 4.4 Consider $F := \{\perp, \{v\}, \{\bar{v}\}\}$ and $F' := \{\{v\}, \{\bar{v}\}\}$; we have $F' \in \mathcal{MU}$ and $\text{var}(F') = \text{var}(F)$, but $F \notin \mathcal{VMU}$, since $\{\perp\} \in \mathcal{USAT}$. If we don't want to use the empty clause, then we can consider any $F' \in \mathcal{MU}$ with $v \in \text{var}(F')$ and $\{\{v\}, \{\bar{v}\}\} \cap F' = \emptyset$, and let $F := F' \cup \{\{v\}, \{\bar{v}\}\}$ — again we have $F' \in \mathcal{MU}$ and $\text{var}(F') = \text{var}(F)$, but $F \notin \mathcal{VMU}$.

The corrected version of [15, Lemma 5] is as follows:

Lemma 4.5 For $F \in \mathcal{CLS}$ let \mathbb{U}_F be the set of $F' \subseteq F$ with $\text{var}(F') = \text{var}(F)$, $\delta(F') \geq 1$, and $F' \in \mathcal{USAT}$. Then $F \in \mathcal{VMU}$ if and only if $F \in \mathbb{U}_F$ and all minimal elements of \mathbb{U}_F w.r.t. the subset-relation are minimally unsatisfiable.

Proof: The condition is necessary, since if $F \in \mathcal{VMU}$, then on the one hand we have $F \in \mathcal{LEAN} \setminus \{\top\}$, and thus $\delta(F) \geq 1$ by [10] (or use [15, Lemma 3]); and on the other hand if there would be a minimal element $F' \in \mathbb{U}_F$ which wouldn't be minimally unsatisfiable, then there would be some $F'' \subset F'$ with $F'' \in \mathcal{MU}$, whence by definition of \mathbb{U}_F we get $\text{var}(F'') \subset \text{var}(F)$ contradicting $F \in \mathcal{VMU}$.

For the other direction assume, that we have \mathbb{U}_F as specified, and we have to show $F \in \mathcal{VMU}$. Since $F \in \mathbb{U}_F$, we have $F \in \mathcal{USAT}$. Consider now some $F' \subseteq F$ with $F' \in \mathcal{USAT}$, and assume $\text{var}(F') \subset \text{var}(F)$. Consider some minimal $F'' \in \mathbb{U}_F$ (regarding inclusion) with $F' \subset F'' \subseteq F$. Furthermore consider a minimal element $G \in \mathbb{U}_F$ with $G \subseteq F''$; by assumption $G \in \mathcal{MU}$, and since $F' \subset F''$, we have $G \subset F''$. If for $C \in F''$ we had $\text{var}(F'' \setminus \{C\}) \subset \text{var}(F'')$, then there would be $x \in C$ such that x or \bar{x} is pure in F'' , thus also pure in G , whence $C \notin G$ (since $G \in \mathcal{MU}$), contradicting $\text{var}(G) = \text{var}(F)$. Now choose some $C \in F'' \setminus F'$ (we

have $\text{var}(F'' \setminus \{C\}) = \text{var}(F'')$; by minimality of F'' we now have $\delta(F'' \setminus \{C\}) \leq 0$ (otherwise all conditions for \mathbb{U}_F are fulfilled for $F'' \setminus \{C\}$), whence $\delta(F'') = 1$. But then due to $\text{var}(F'') = \text{var}(G)$ and $G \subset F''$ it follows $\delta(G) \leq 0$, contradicting $G \in \mathcal{MU}$. \square

The following examples show applications of Lemma 4.5:

Example 4.6 Consider the two (non-)examples from Example 4.4:

1. For $F = \{\perp, \{v\}, \{\bar{v}\}\}$ we have the minimal element $\{\perp, \{v\}\}$ of \mathbb{U}_F which is not minimally unsatisfiable.
2. For $F = F' \cup \{\{v\}, \{\bar{v}\}\}$ (note that from the assumptions follows $\text{var}(F') \supset \{v\}$) consider a minimal $F'' \subseteq F'$ with $\text{var}(G) = \text{var}(F)$ and $\delta(G) \geq 1$ for $G := F'' \cup \{\{v\}, \{\bar{v}\}\}$ (note $\top \subset F'' \subset F$): now G is a minimal element of \mathbb{U}_F which is not minimally unsatisfiable.

Based on [15, Lemma 5], also the proof of Theorem 3 in [15] is false (the procedure goes astray on the clause-sets of Example 4.4). Fortunately we can give a simple proof of the assertion, which even shows fixed-parameter tractability (fpt) of the decision problem “ $F \in \mathcal{VMU}_{\delta=k}$?” in the parameter k :

Theorem 4.7 Membership decision “ $F \in \mathcal{VMU}_{\delta=k}$?” for input $F \in \mathcal{CLS}$ is fpt in the parameter $k \in \mathbb{Z}$.

Proof: If $F \notin \mathcal{MLEN}$, then $F \notin \mathcal{VMU}$ (by Lemma 4.3). So we can assume now $F \in \mathcal{MLEN}$, and thus we have $\delta(F') < \delta(F)$ for all $F' \subset F$. Now the decisions of Lemma 4.1, as discussed in Example 2.18, are fpt in k . \square

[15, Theorem 1] shows that decision of “ $F \in \mathcal{VMU}$?” is D^P -complete.

5. Eliminating and creating singularity

In this section we continue the study of the handling of singular variables in minimally unsatisfiable clause-sets, as initiated in [18, 19]. In Section 5.1 we study the reduction process, eliminating singular variables. A main insight is Lemma 5.4, showing that the elimination is harmless concerning the minimum variable degree. In Subsection 5.2 we introduce the inverse elimination (“extension”); the main point here is the precise statement of the various conditions. Finally in Subsection 5.3 we consider a special case of singularity, namely unit-clauses.

5.1. Singular DP-reduction

In [19] the process of “singular DP-reduction” has been studied for minimally unsatisfiable clause-sets. By it we can reduce the case of arbitrary $F \in \mathcal{MU}$ to (nonsingular) $F' \in \mathcal{MU}'$ (that is, for every $v \in \text{var}(F')$ we have $\text{ld}_{F'}(v), \text{ld}_{F'}(\bar{v}) \geq 2$). The definition is as follows (see [19, Definition 8]):

Definition 5.1 ([19]) The relation $\mathbf{F} \xrightarrow{sDP} \mathbf{F}'$ (singular DP-reduction) holds for clause-sets $F, F' \in \mathcal{CLS}$, if there is a singular variable v in F , such that F' is obtained from F by DP-reduction on v , that is, $F' = \text{DP}_v(F)$. The reflexive-transitive closure of this relation is denoted by $\mathbf{F} \xrightarrow{sDP}_* \mathbf{F}'$.

By $\text{sDP}(F) \subset \mathcal{MU}$ for $F \in \mathcal{MU}$ the set of nonsingular $F' \in \mathcal{MU}$ with $F \xrightarrow{\text{sDP}}_* F'$ is denoted. For us the main property of $\text{sDP}(F)$ is that it is not empty. In [19] it is shown that for $S \in \mathcal{SMU}$ we have $|\text{sDP}(F)| = 1$, and that for arbitrary $F \in \mathcal{MU}$ and $F', F'' \in \text{sDP}(F)$ we have $n(F') = n(F'')$.

Example 5.2 In [19] the following is shown for $F \in \mathcal{MU}$:

1. For $\delta(F) = 1$ we have $\text{sDP}(F) = \{\perp\}$.
2. For $\delta(F) = 2$ all elements of $\text{sDP}(F)$ are pairwise isomorphic.
3. For $\delta(F) \geq 3$ in general there are non-isomorphic elements in $\text{sDP}(F)$.

By the results of Sections 3.1, 3.2 in [19] we have the following basic preservation properties:

Lemma 5.3 ([19]) For $F, F' \in \mathcal{MU}$ with $F \xrightarrow{\text{sDP}}_* F'$ we have:

1. $\delta(F') = \delta(F)$.
2. $F \in \mathcal{MU} \Rightarrow F' \in \mathcal{MU}$.
3. $F \in \mathcal{SMU} \Rightarrow F' \in \mathcal{SMU}$.
4. $F \in \mathcal{UHLT} \Rightarrow F' \in \mathcal{UHLT}$.

Although singular DP-reduction can reduce the variable-degree of some variables, it can not decrease the *minimum* variable-degree:

Lemma 5.4 For $F, F' \in \mathcal{MU}$ with $F \xrightarrow{\text{sDP}}_* F'$ we have $\mu\text{vd}(F') \geq \mu\text{vd}(F)$.

Proof: It is sufficient to consider the case $F' = \text{DP}_v(F)$ for a singular variable v . Assume $\mu\text{vd}(F') < \mu\text{vd}(F)$; thus $\text{var}(F') \neq \emptyset$ (otherwise we have $\mu\text{vd}(F') = +\infty$), and we consider $w \in \text{var}(F')$ with $\text{vd}_{F'}(w) = \mu\text{vd}(F')$. So we have $\text{vd}_{F'}(w) < \text{vd}_F(w)$, and thus by [19, Lemma 24], for all clauses $C \in F$ with $v \in \text{var}(C)$ we have $w \in \text{var}(C)$. But then $\mu\text{vd}(F') = \text{vd}_{F'}(w) \geq \text{vd}_F(v) \geq \mu\text{vd}(F) > \mu\text{vd}(F')$, a contradiction. \square

Thus in order to determine the minimum variable-degree for minimally unsatisfiable clause-sets in dependency on the deficiency, w.l.o.g. one can restrict attention to saturated and nonsingular instances:

Corollary 5.5 For all $k \in \mathbb{N}$, $k \geq 2$, holds:

1. $\text{VDM}(k) = \mu\text{vd}(\mathcal{MU}_{\delta=k}) = \mu\text{vd}(\mathcal{SMU}'_{\delta=k})$.
2. $\text{VDH}(k) = \mu\text{vd}(\mathcal{UHLT}_{\delta=k}) = \mu\text{vd}(\mathcal{UHLT}'_{\delta=k})$.
3. $\text{VDM}(1) = \text{VDH}(1) = \mu\text{vd}(\{\{1\}, \{-1\}\}) = 2$.

Proof: For Part 1 we note that by Lemma 5.4 for every $F \in \mathcal{MU}_{\delta=k}$ we can find $F' \in \mathcal{SMU}'_{\delta=k}$ with $\mu\text{vd}(F') \geq \mu\text{vd}(F)$, and thus $\mu\text{vd}(\mathcal{MU}_{\delta=k}) \leq \mu\text{vd}(\mathcal{SMU}'_{\delta=k})$, while $\mu\text{vd}(\mathcal{MU}_{\delta=k}) \geq \mu\text{vd}(\mathcal{SMU}'_{\delta=k})$ holds due to $\mathcal{SMU}'_{\delta=k} \subseteq \mathcal{MU}_{\delta=k}$. The same reasoning applies for Part 2. We used that by singular DP-reduction not all variables can vanish here, since $\delta(\{\perp\}) = 1$. For $F \in \mathcal{MU}_{\delta=1}$ however we can reach $\{\perp\}$ (recall Example 3.2), and thus we stop one step before, which proves Part 3. \square

See Lemma 15.9 for some conditions under which $k \in \mathbb{N} \mapsto \text{VDH}(k)$ and $k \in \mathbb{N} \mapsto \text{VDM}(k)$ would be computable (which is part of Conjecture 15.1, while Conjecture 12.1 states that both maps are indeed equal).

5.2. Singular DP-extensions

We consider now the reverse direction of singular DP-reduction, from $\text{DP}_v(F)$ to F , as a *singular extension*, and also generalise it to arbitrary clause-sets. This process was mentioned in [19, Examples 15,19,54] for minimally unsatisfiable $\text{DP}_v(F)$, called “inverse singular DP-reduction” there:

Definition 5.6 Consider a clause-set $G \in \mathcal{CLS}$, a variable $v \in \mathcal{VA} \setminus \text{var}(G)$, and $m \in \mathbb{N}$ with $m \leq c(G)$. A **singular m -extension of G with v** is a clause-set $F \in \mathcal{CLS}$ obtained as follows (employing four choice steps):

- (i) m different clauses $D_1, \dots, D_m \in G$ are chosen.
- (ii) A subset $C \subseteq \bigcap_{i=1}^m D_i$ is chosen.
- (iii) Clauses $D'_i \in \mathcal{CL}$ for $i \in \{1, \dots, m\}$ with $(D_i \setminus C) \subseteq D'_i \subseteq D_i$ are chosen.
- (iv) A literal x with $\text{var}(x) = v$ is chosen.

Let $C' := C \cup \{x\}$, and let $D''_i := D'_i \cup \{\bar{x}\}$ for $i \in \{1, \dots, m\}$. F is obtained from G by adding C' and replacing the D_i with D''_i :

$$F := (G \setminus \{D_1, \dots, D_m\}) \cup \{C', D''_1, \dots, D''_m\}.$$

Note that the order of the clauses D_i does not matter at all (Step (i)), so in reality an m -element subset of G is chosen, and that the choice of x (Step (iv)), either $x = v$ or $x = \bar{v}$, is inessential (the resulting F just differ by flipping variable v). We also note that the clauses D'_1, \dots, D'_m are pairwise different. Obviously the four choices are always possible (only $1 \leq m \leq c(G)$ is needed for that).

Example 5.7 Consider $G := \{\{a, b, c\}, \{a, b, \bar{c}\}\}$, $m := 2$, and so $\{D_1, D_2\} = G$ and $\bigcap_{i=1}^m D_i = \{a, b\}$, and the choices $C := \{a\}$, $D'_1 := \{b, c\}$, $D'_2 := \{a, b, \bar{c}\}$, and $x := v$. Then the 2-extension F of G is $F = \{\{v, a\}, \{\bar{v}, b, c\}, \{\bar{v}, a, b, \bar{c}\}\}$.

By definition we have for an m -extension F of $G \in \mathcal{CLS}$ with v the following simple properties:

1. $c(F) = c(G) + 1$, $n(F) = n(G) + 1$, $\delta(F) = \delta(G)$.
2. v is singular for F , $\text{vd}_F(v) = m + 1$.
3. $\text{DP}_v(F) = G$.

We now show that indeed the process of Definition 5.6 is precisely the inversion of singular DP-reduction:

Lemma 5.8 Consider $m \in \mathbb{N}$, $G, F \in \mathcal{CLS}$ and $v \in \mathcal{VA}$. Then F is an m -extension of G by v iff the following conditions are fulfilled:

1. v is singular for F ;
2. $\text{vd}_F(v) = m + 1$;
3. $\text{DP}_v(F) = G$, $c(G) = c(F) - 1$.

Proof: If F is an m -extension of G by v , then the three properties hold, as we have already mentioned. So assume these three properties hold. Now let the m clauses D_1, \dots, D_m be the result of singular DP-reduction on v for F ; they must be pairwise different, and all m resolutions must be possible, otherwise $c(G) < c(F) - 1$. And let C be singular occurrence of v minus the variable v . Now all properties of a singular m -extension are easily checked. \square

Singular extensions behave well regarding minimal unsatisfiability:

Lemma 5.9 *Consider $m \in \mathbb{N}$, $G \in \mathcal{CLS}$ and an m -extension F of G by $v \in \mathcal{VA}$. Then $F \in \mathcal{MU} \Leftrightarrow G \in \mathcal{MU}$.*

Proof: This follows by Lemma 5.8 together with [19, Lemma 9, Parts 1, 2]. \square

In the situation of Lemma 5.9, regarding saturatedness we only have the direction $F \in \mathcal{SMU} \Rightarrow G \in \mathcal{SMU}$, while for the other direction the conditions of [19, Lemma 12] need to be observed (this would yield “saturated extensions”, which however we do not need here).

5.3. Unit clauses

We conclude this section by considering unit-clauses in minimally unsatisfiable clause-sets. The following (fundamental, simple) lemma is [19, Lemma 14]; in [19, Subsection 3.3] one finds further information on unit-clauses in minimally unsatisfiable clause-sets.

Lemma 5.10 ([19]) *Consider $F \in \mathcal{MU}$.*

1. *If v is full and singular in F , then we have $\{v\} \in F$ or $\{\bar{v}\} \in F$.*
2. *If $\{x\} \in F$, then $v := \text{var}(x)$ is singular in F (with $\text{ld}_F(x) = 1$). If here F is saturated, then v is also full in F .*

So unit-clauses in minimally unsatisfiable clause-sets are strong cases of singular variables. They can obviously be removed by singular DP-reduction, while on the other hand singular ≥ 2 -extensions can *not* remove all unit-clauses:

Lemma 5.11 *Consider a clause-set $F \in \mathcal{MU}$ containing at least one unit-clause, and obtain F' from F by a singular m -extension, where $m \geq 2$. Then also F' must contain at least one unit-clause.*

Proof: For a unit-clause $\{x\} \in F$ to be removed in F' , it needs to be one of the D_i (using the terminology of Definition 5.6). Then the intersection C must be empty (otherwise any other D_i needed to contain x , and since $m \geq 2$ this would mean a subsumption in F). Thus the extension introduces the new unit-clause C' . \square

The following examples show that the assumptions $F \in \mathcal{MU}$ and $m \geq 2$ in Lemma 5.11 are needed:

Example 5.12 *For the condition “ $m \geq 2$ ” consider $F := \{\{a\}, \{\bar{a}, b\}, \{\bar{a}, \bar{b}\}\} \in \mathcal{MU}_{\delta=1}$. Via a 1-singular extension we obtain $F' := \{\{v, a\}, \{\bar{v}, a\}, \{\bar{a}, b\}, \{\bar{a}, \bar{b}\}\}$, which has no unit-clauses. On the other hand, using the notation from Definition 5.6, any ≥ 2 -singular extension of F which touches $\{a\}$ (i.e., $\{a\}$ is one of the D_i) has intersection $C = \perp$, and thus C' is a new unit-clause, while if $\{a\}$ is not touched, then this unit-clause is simply maintained.*

For the condition “ $F \in \mathcal{MU}$ ” consider $F := \{\{a\}, \{a, b\}\} \in \mathcal{CLS} \setminus \mathcal{MU}$, where $F' := \{\{v, a\}, \{\bar{v}, a\}, \{\bar{v}, b\}\}$ is a 2-extension without unit-clauses.

For certain $F \in \mathcal{MU}_{\delta=2}$ the existence of a unit-clause is actually necessary for singularity:

Lemma 5.13 *Consider $F \in \mathcal{MU}_{\delta=2}$ with $\mu\text{vd}(F) \geq 4$. Then F is singular if and only if F contains a unit-clause.*

Proof: That if F contains a unit-clause, then F must be singular, follows by Lemma 5.10, Part 2. So assume now that F is singular, and we have to show that F contains a unit-clause. Consider a reduction sequence $F = F_0 \xrightarrow{\text{sDP}} F_1 \xrightarrow{\text{sDP}} \dots \xrightarrow{\text{sDP}} F_m$, where F_m is nonsingular (note $m \geq 1$). So there exists $n \geq 2$ such that F_m is isomorphic to \mathcal{F}_n (recall Example 3.3), and thus every variable of F_m has degree 4. So by Lemma 5.4 we know $\mu\text{vd}(F_i) = 4$ for $i \in \{0, \dots, m\}$. We show by induction on m that F contains a unit-clause. If $m = 1$, then in order to obtain the min-var-degree of at least 4, at least 3 side-clauses $D_1, \dots, D_3 \in \mathcal{F}_n$ for the singular extension have to be chosen (using Definition 5.6), but every literal occurs precisely twice in \mathcal{F}_n (because of variable-degree 4 and nonsingularity), and thus the intersection C has to be empty, and the new clause introduced by the singular extension is a unit-clause, whence F contains a unit-clause. Finally assume $m > 1$. So by induction hypothesis, F_1 contains a unit-clause, and thus by Lemma 5.11 also F_0 contains a unit-clause. \square

We will later see (Theorem 8.3) that the condition $\mu\text{vd}(F) \geq 4$ in Lemma 5.13 is equivalent to $\mu\text{vd}(F) = 4$; the following examples show that this condition can not be improved:

Example 5.14 *A 1-singular extension of A_2 is*

$$F_1 := \{\{1, 2, 3\}, \{1, 2, -3\}, \{-1, 2\}, \{1, -2\}, \{-1, -2\}\} \in \mathcal{MU}_{\delta=2},$$

where F_1 is singular, has no unit-clause, and $\mu\text{vd}(F_1) = 2$. While a 2-singular extension of A_2 is

$$F_2 := \{\{1, 3\}, \{1, 2, -3\}, \{1, -2, -3\}, \{-1, 2\}, \{-1, -2\}\} \in \mathcal{MU}_{\delta=2},$$

where F_2 is singular, has no unit-clause, and $\mu\text{vd}(F_2) = 3$.

We conclude with a simple form of adding a new variable, by adding it in one sign as unit-clause, and adding it in the other sign to all given clauses (so that we obtain a full variable, occurring in all clauses):

Definition 5.15 *A full singular unit-extension of a clause-set $F \in \mathcal{CLS}$ (by unit-clause $\{x\}$) is a clause-set $F' \in \mathcal{CLS}$ obtained from F by adding a unit-clause $\{x\}$ with $\text{var}(x) \notin \text{var}(F)$, and by adding literal \bar{x} to all clauses of F , i.e., $F' := \{\{x\}\} \cup \{C \cup \{\bar{x}\} : C \in F\}$ for some $x \in \mathcal{LIT} \setminus \text{lit}(F)$.*

A full singular unit-extension F' of $F \neq \top$ by $\{x\}$ is a special case of a singular $c(F)$ -extension of F with $\text{var}(x)$ (recall Definition 5.6), and thus $F' \xrightarrow{\text{sDP}} F$.¹⁴

Example 5.16 *Starting with $\{\perp\}$, the first full singular unit-extension is $\{\{v\}, \{\bar{v}\}\}$ (up to the choice of the new literal), the second is $\{\{w\}, \{v, \bar{w}\}, \{\bar{v}, w\}\}$. In this way we get special examples of $\mathcal{SMU}_{\delta=1}$ (since we started with $\{\perp\} \in \mathcal{SMU}_{\delta=1}$).*

¹⁴The case $m = 0$ is excluded in Definition 5.6, since it is not needed, and would only complicate the formulation.

If we start with \top instead, then first we get $\{\{v\}\}$, and then $\{\{w\}, \{v, \bar{w}\}\}$.

[19, Example 15, Part 1] contains two example of “inverse unit elimination”, where Example (a) there is an example of a full singular unit-extension, while Example (b) there would be a non-full singular unit-extension (where the new variable is not full; this is not used in the present report).

The process of full singular unit-extension of a clause-set F maintains many properties of F , and we list here those we use:

Lemma 5.17 *Consider a full singular unit-extension F' of F (by $\{v\}$):*

1. $n(F') = n(F) + 1$ and $c(F') = c(F) + 1$.
2. $\delta(F') = \delta(F)$.
3. $\sigma(F') = \sigma(F)$ for $F \neq \{\perp\}$.
4. $\mu\text{vd}(F') = \mu\text{vd}(F)$ for $n(F) > 0$.
5. F' is satisfiable iff F is satisfiable.
6. For $F \neq \top$: F' is lean iff F is lean.
7. F' is (saturated) minimally unsatisfiable iff F is (saturated) minimally unsatisfiable.
8. F' is hitting iff F is hitting.

Proof: Parts 1, 2 follow directly by definition. For Part 3 we notice that for $F = \top$ we have $\sigma(F') = \sigma(F) = 0$, while for $n(F) > 0$ consider $\emptyset \subset V \subseteq \text{var}(F')$: if $v \notin V$, then $F'[V] = F[V]$, and thus the minimisation for $\sigma(F)$ is included in $\sigma(F')$, and if $v \in V$, then $\delta(F'[V]) = c(F') - |V| \geq \delta(F') = \delta(F) \geq \sigma(F)$, and thus these V do not contribute to the minimisation.

For Part 4 we just note that the variables of F keep their degrees in F' , while the new variable has degree $\text{vd}_{F'}(v) = c(F') > c(F)$, and thus does not contribute to the min-var degree. Part 5 is trivial, and follows also by the satisfiability-equivalence of $\text{DP}_v(F)$ and F . For Part 6 we note, that an autarky for F' involving v must be a satisfying assignment for F' , while the autarkies for F' not involving v are the same as the autarkies for F . Part 7 concerning (just) minimal unsatisfiability follows with [19, Lemma 9], while regarding saturatedness we can use [19, Lemma 12] (both assertions also follow easily by direct reasoning). Part 8 is trivial. \square

So our fundamental classes are respected by full singular unit-extension:

Corollary 5.18 *If $F \in \mathcal{MU}_{\delta=k}$ ($k \in \mathbb{N}$), then every full singular unit-extension is also in $\mathcal{MU}_{\delta=k}$. If furthermore F is saturated resp. hitting, then every full singular unit-extension is also saturated resp. hitting.*

Obviously, full singular unit-extension is unique up to isomorphism:

Lemma 5.19 *Consider a clause-set $F \in \mathcal{CLS}$ and clause-sets $F', F'' \in \mathcal{CLS}$ obtained from F by repeated full singular unit-extensions. Then F', F'' are isomorphic if and only if $n(F') = n(F'')$.*

Proof: The number of repeated full singular unit-extensions leading to F' resp. F'' is the number of variables in these clause-sets with degree strictly greater than $c(F)$, and sorting these variables by increasing degree yields the sequence of extensions. Thus just from knowing the number of variables in F', F'' we can reconstruct

them up to isomorphism (using that a full singular unit-extension of F by $\{x\}$ is isomorphic to one by $\{y\}$, for arbitrary literals x, y with new variables). \square

6. Full subsumption resolution / extension

In this section we investigate the second reduction concept for this report, “full subsumption resolution”. As with singular DP-reduction from Section 5, in general this reduction uncovers hidden structure, while the inverse process, “full subsumption extension”, serves as a generator for minimally unsatisfiable clause-sets with various properties. In this report, unlike with singular DP-reduction, we will not consider full subsumption resolution for arbitrary $F \in \mathcal{MU}$, but only starting from some $A(V)$, while a deeper use will be important for [82, 90]. Subsection 6.1 discusses the basic definitions (there are various technicalities one needs to be aware of), and first applications are given in Subsection 6.2.

The basic idea is, for a clause-set F containing two clauses $R \cup \{v\}, R \cup \{\bar{v}\} \in F$, to replace these two clauses by the clause R , i.e., we consider the case where the resolvent R of parent clauses C, D subsumes both parent clauses (thus the name). This is a very old procedure, based on the trivial observation that $(R \vee v) \wedge (R \vee \neg v)$ is logically equivalent to R . If we perform this in the inverse direction, as an “extension”, then every clause-set $F \in \mathcal{CLS}$ can be transformed into its (equivalent) “distinguished” or “canonical” CNF $F' \subseteq A(\text{var}(F))$ (just expand every non-full clause), which is uniquely determined, namely F' is the set of $C \in A(\text{var}(F))$ such that there is $D \in F$ with $D \subseteq C$.

We however have to be more careful about deficiency and membership in \mathcal{MU} , and thus will consider only “full subsumption resolution”, where the resolvent must not be present already, while for the “strict” form additionally the resolution variable v must occur also in other clauses. Then from $A(V)$ by strict full subsumption resolution we can obtain precisely the $F \in \mathcal{UHT}$ with $\text{var}(F) = V$ (Lemma 6.9). For the inverse forms we have to be equally carefully, making sure that none of the two parent clauses is already present (this prevents the above expansion of arbitrary $F \in \mathcal{CLS}$ to $A(\text{var}(F))$).

The (more general) well-known “subsumption resolution” is the reduction $F \rightsquigarrow (F \setminus \{C\}) \cup \{C \setminus \{x\}\}$ for $F \in \mathcal{CLS}$, that is the removal of a literal $x \in C$ from a clause $C \in F$, in case there exists $D \in F$ with $\bar{x} \in D$ and $D \setminus \{\bar{x}\} \subseteq C$ (note that $C \diamond D = C \setminus \{x\}$ subsumes C). An early use is in [91], under the name “replacement principle”, while the terminology “subsumption resolution” is used in [92] (for SAT solving). The earliest sources with a systematic treatment appear to be [93, Section 7] and [94, Section 7]. An experimental study of the practical importance of subsumption resolution in connection with DP-reductions $F \rightsquigarrow \text{DP}_v(F)$ (under suitable additional conditions to make DP-reduction feasible; see [19, Subsection 1.3] for an overview on such restrictions) is performed in [95], under the name of “self-subsuming resolution”, and continued in [96]. A theoretic (similar) use one finds in [97, Section 4], where a variable v is called “DP-simplicial” for $F \in \mathcal{CLS}$ iff all resolutions performed by the reduction $F \rightsquigarrow \text{DP}_v(F)$ are subsumption resolutions.

Our special form, where *both* parent clauses are subsumed, we call *full subsumption resolution*, namely the reduction $F \rightsquigarrow (F \setminus \{C, D\}) \cup \{C \diamond D\}$ in case of $C, D \in F$ such that $C \cap \bar{D} = \{x\}$ and $C \setminus \{x\} = D \setminus \{\bar{x}\}$. A main tool is Lemma 6.5, where especially Part 7 is somewhat subtle, and can be easily overlooked. Via this tool we have a controlled way of transforming $F \in \mathcal{MU}$ resp. $F \in \mathcal{UHT}$ into $A(\text{var}(F))$, and in Theorem 6.13 this yields the determination of the possible numbers of variables and clauses in minimally unsatisfiable clause-sets of a given deficiency.

6.1. Basic definitions

Before defining “full subsumption reduction” $F \rightsquigarrow (F \setminus \{R \cup \{v\}, R \cup \{\bar{v}\}\}) \cup \{R\}$ in Definition 6.7 (so R is new and the two clauses $R \cup \{v\}, R \cup \{\bar{v}\}$ vanish), we introduce the “strict” form, which is more important to us, and which has the additional condition that v must still occur (in other clauses of F ; the “non-strict” form on the other hand guarantees that v vanishes (see Definition 6.7)):

Definition 6.1 For clause-sets $F, F' \in \mathcal{CLS}$ by $F \xrightarrow{sfsR} F'$ we denote that F' is obtained from F by one step of **strict full subsumption resolution**, that is,

- there is a clause $R \in F'$
- and a literal x with $\text{var}(x) \notin R$
- such that for the clauses $C := R \cup \{x\}$ and $D := R \cup \{\bar{x}\}$
- we have $F = (F' \setminus \{R\}) \cup \{C, D\}$;
- we furthermore require $\text{var}(x) \in \text{var}(F')$.
- As usual, the literals x, \bar{x} are the resolution literals, $\text{var}(x)$ is the resolution variable, C, D are the parent clauses, and R is the resolvent.

We write $F \xrightarrow{sfsR}_k F'$ for $k \in \mathbb{N}_0$ if exactly k steps have been performed, while we write $F \xrightarrow{sfsR}_* F'$ for an arbitrary number of steps (including zero).

We require $R \notin F$, that is, the (full subsumption) resolvent is not already present in the original clause-set. This is of course satisfied if $F \in \mathcal{MU}$. We also require that the variable v does not vanish, for the sake of keeping control on the deficiency.

Example 6.2 Some simple examples are:

1. $\mathcal{F}_2 = \{\{1, 2\}, \{-1, -2\}, \{-1, 2\}, \{-2, 1\}\} \xrightarrow{sfsR} \{\{2\}, \{-1, -2\}, \{-2, 1\}\}$, and no further reduction is possible (note that the only possibility is blocked, since variable 1 would vanish).
2. $\{\{v\}, \{\bar{v}\}\} \not\xrightarrow{sfsR} \{\perp\}$, as v vanishes, while $\{\{v\}, \{\bar{v}\}, \{v, x\}\} \xrightarrow{sfsR} \{\perp, \{v, x\}\}$.
3. $\{\{v, w\}, \{\bar{v}, w\}, \{v, \bar{w}\}\} \not\xrightarrow{sfsR} \{\{v, w\}, \{\{w\}, \{v, \bar{w}\}\}\}$, as one parent clause would be kept, while $\{\{v, w\}, \{\bar{v}, w\}, \{v, \bar{w}\}\} \xrightarrow{sfsR} \{\{\{w\}, \{v, \bar{w}\}\}\}$.
4. $\{\{v, w\}, \{\bar{v}, w\}, \{v, \bar{w}\}, \{v\}, \{w\}\}$ can not be reduced by strict-full-subsumption resolution, since all possible resolvents are already there.

The expansion of a clause R to two clauses $R \cup \{v\}, R \cup \{\bar{v}\}$ under the above requirements is called “extension”:

Definition 6.3 For clause-sets $F, F' \in \mathcal{CLS}$ we say that F is obtained from F' by **strict full subsumption extension** if $F \xrightarrow{sfsR} F'$. And for $k \in \mathbb{N}_0$ we say that F is obtained from F' by **strict full subsumption extension with k steps** if $F \xrightarrow{sfsR}_k F'$.

So one step of strict full subsumption extension for a clause-set F uses a non-full clause $R \in F$ and a variable $v \in \text{var}(F) \setminus \text{var}(R)$, and replaces R by the two clauses $R \cup \{v\}, R \cup \{\bar{v}\}$, where none of them is already present.

Example 6.4 From $\{\{a\}, \{b\}\}$ by one step of strict full subsumption extension we can obtain $\{\{a, b\}, \{a, \bar{b}\}, \{b\}\}$ or $\{\{a\}, \{a, b\}, \{\bar{a}, b\}\}$; note that no new variable has been introduced, that the original clause ($\{a\}$ resp. $\{b\}$) vanished, and that its replacement clauses were not already present. For $\{\{a, b\}, \{a\}\}$ no strict full subsumption extension is possible. Further examples are obtained by “reading Example 6.2 backwards”.

The basic properties of strict full subsumption resolution are collected in the following lemma.

Lemma 6.5 For clause-sets $F, F' \in \mathcal{CLS}$ with $F \xrightarrow{sfsR}_k F'$ ($k \in \mathbb{N}_0$) we have:

1. $\text{var}(F') = \text{var}(F)$.
2. $c(F') = c(F) - k$, $\delta(F') = \delta(F) - k$.
3. F' is logically equivalent to F .
4. $\mu\text{vd}(F) \geq \mu\text{vd}(F')$.
5. $F \in \mathcal{MU} \Rightarrow F' \in \mathcal{MU}$.
6. $F \in \mathcal{SMU} \Rightarrow F' \in \mathcal{SMU}$.
7. Assume $k = 1$ with resolution variable v and resolvent R , and assume $F' \in \mathcal{MU}$. Then exactly one of the following three possibilities holds:
 - (a) $S(F', R, v)$ is a partial saturation of F' (recall Definition 3.7).
 - (b) $S(F', R, \bar{v})$ is a partial saturation of F' .
 - (c) $F \in \mathcal{MU}$.
8. $F' \in \mathcal{SMU} \Rightarrow F \in \mathcal{MU}$.
9. $F \in \mathcal{HIT} \Leftrightarrow F' \in \mathcal{HIT}$, $F \in \mathcal{UHIT} \Leftrightarrow F' \in \mathcal{UHIT}$.

Proof: Parts 1, 2, 3, 4 follow directly from the definition. Part 5 holds since we strengthen two clauses into one, which is logically equivalent to its parent clauses, and for Part 6 additionally note that a saturation of F' by one literal can also be done on F (if the resolvent is involved, then the resolution variable can not be v , since $F \in \mathcal{MU}$). Now consider Part 7. That the two possibilities for partial saturation exclude each other follows by Lemma 3.9 (and $F' \setminus \{R\} \not\equiv R$). And that each possibility for partial saturation excludes $F \in \mathcal{MU}$ follows by definition. Finally, that the negation of the two partial saturation possibilities implies $F \in \mathcal{MU}$ follows again by Lemma 3.9. Finally Part 8 follows by Part 7, while Part 9 follows by trivial combinatorics. \square

Part 7 of Lemma 6.5 handles a subtle source for errors: One could easily think that for $F' \in \mathcal{MU}$ a strict full subsumption extension yields another $F \in \mathcal{MU}$, but this is not so, as there are three possible cases to be considered here, illustrated by the following examples:

Example 6.6 Consider $F := \{\{v, a\}, \{\bar{v}, a\}, \{\bar{v}\}, \{v, \bar{a}\}\}$. So $F \xrightarrow{sfsR}_1 F'$ for $F' := \{\{a\}, \{\bar{v}\}, \{v, \bar{a}\}\}$. We have $F' \in \mathcal{MU}$, but $F \notin \mathcal{MU}$, and indeed $S(F', R, v) = \{\{a, v\}, \{\bar{v}\}, \{v, \bar{a}\}\}$ is a partial saturation of F' (while $S(F', R, \bar{v})$ isn't one).

An example that Part 8 can not be strengthened to “ $F' \in SMU \Rightarrow F \in SMU$ ” is obtained from $F' := \mathcal{F}_4$ by one strict full subsumption extension on $\{-1, 2\}$ with resolution variable 3, obtaining

$$F = \{\{1, 2, 3, 4\}, \{-1, -2, -3, -4\}, \\ \{-1, 2, 3\}, \{-1, 2, -3\}, \{-2, 3\}, \{-3, 4\}, \{-4, 1\}\} \in \mathcal{MU}'_{\delta=3} \setminus SMU,$$

where $\langle 4 \rightarrow 0 \rangle * F = \{\{1, 2, 3\}, \{-1, 2, 3\}, \{-1, 2, -3\}, \{-2, 3\}, \{-3\}\} \notin \mathcal{MU}$.

The condition on the resolution variable for strict full subsumption resolution (that it must not vanish) is exactly needed for Parts 1, 2 of Lemma 6.5. If this condition is dropped, then we speak of *full subsumption resolution*:

Definition 6.7 *Full subsumption resolution* is defined as strict full subsumption resolution, but now the resolution variable is allowed to vanish. If the resolution variable definitely vanishes, then we speak of *non-strict full subsumption resolution*. In the other direction we speak of *full subsumption extension* resp. *non-strict full subsumption extension*.

So if F' is obtained from F by one step of non-strict full subsumption extension, then we have $c(F') = c(F) + 1$, $n(F') = n(F) + 1$ and $\delta(F') = \delta(F)$. As mentioned, all parts of Lemma 6.5 are true for full subsumption resolution except of Parts 1, 2.

Example 6.8 *Considering the non-examples from Example 6.2:*

1. $\{\{v\}, \{\bar{v}\}\} \not\xrightarrow{sfsR} \{\perp\}$, but by full subsumption resolution we obtain $\{\perp\}$.
2. $\{\{v, w\}, \{\bar{v}, w\}, \{v, \bar{w}\}\} \not\xrightarrow{sfsR} \{\{v, w\}, \{\{w\}, \{v, \bar{w}\}\}\}$, and the transition is also not possible by full subsumption resolution.
3. $\{\{v, w\}, \{\bar{v}, w\}, \{v, \bar{w}\}, \{v\}, \{w\}\}$ is irreducible by full subsumption resolution.

As follows from the characterisation of $SMU_{\delta=1} = \mathcal{UHIT}_{\delta=1}$ in [7], a clause-set $F \in \mathcal{CLS}$ can be reduced by a series of non-strict full subsumption resolutions to $\{\perp\}$ iff $F \in SMU_{\delta=1} = \mathcal{UHIT}_{\delta=1}$.

6.2. Extensions to full clause-sets

If we start with the full clause-sets $A(V)$, then by strict full subsumption resolution we obtain exactly all unsatisfiable hitting clause-sets:

Lemma 6.9 *If for some finite $V \subset \mathcal{VA}$ we have $A(V) \xrightarrow{sfsR}_* F$, then $F \in \mathcal{UHIT}$ holds. And for $F \in \mathcal{UHIT}$ we have $A(\text{var}(F)) \xrightarrow{sfsR}_* F$.*

Proof: The first part follows by Lemma 6.5, Part 9 (and $A(V) \in \mathcal{UHIT}$). And for the second part note, that if $F \in \mathcal{UHIT}$ has a non-full clause, then a strict full subsumption extension step can be applied, where the result is still in \mathcal{UHIT} (again by Lemma 6.5, Part 9; if F has only full clauses, then $F = A(\text{var}(F))$). \square

Recall that in Example 6.6 we have seen, that strict full subsumption extension does not maintain minimal unsatisfiability in general. Now we show that from arbitrary minimally unsatisfiable F we can indeed go all the way up to $A(\text{var}(F))$, while staying in \mathcal{MU} , when we additionally allow partial saturation:

Lemma 6.10 *For $F \in \mathcal{MU}$ we can obtain $A(\text{var}(F))$ from F by a series of strict full subsumption extensions in combination with partial saturations, where all intermediate results stay in \mathcal{MU} .*

Proof: If $F \in \mathcal{MU}$ has a non-full clause, and if strict full subsumption extension can not be applied in order to obtain $F' \in \mathcal{MU}$, then by Lemma 6.5, Part 7, a partial saturation is possible. \square

We obtain sharp upper bounds on deficiency and number of clauses in terms of the number of variables, showing that for $F \in \mathcal{MU}$ with a given number $n = n(F)$ of variables, the worst-case concerning deficiency and number of clauses is reached by $A_n \in \mathcal{UHLT}$:

Corollary 6.11 *For $F \in \mathcal{MU}$ holds:*

1. $\delta(F) \leq 2^{n(F)} - n(F)$.
2. $c(F) \leq 2^{n(F)}$.

In both cases we have equality iff F is full (i.e., $F = A(\text{var}(F))$).

Proof: For Part 1 note that by Lemma 6.10 we can transform F into $A(\text{var}(F))$ by a series of steps not decreasing the deficiency. Thus $\delta(F) \leq \delta(A(\text{var}(F))) = 2^{n(F)} - n(F)$. For Part 2 note $c(F) = \delta(F) + n(F) \leq 2^{n(F)}$ (by Part 1). For $F = A(\text{var}(F))$ these inequalities are indeed equalities. If we had $\delta(F) = 2^{n(F)} - n(F)$ for some non-full F , then some strict full subsumption extension must be possible, contradicting the upper bound of Part 1. And if we have $c(F) \leq 2^{n(F)}$ for some non-full F , then again some strict full subsumption extension must be possible, contradicting the upper bound of Part 2. \square

We explicitly state the instructive reformulation, that the A_n are the minimally unsatisfiable clause-sets of maximal deficiency for given number m of variables:

Corollary 6.12 *Consider $m \in \mathbb{N}_0$ and $F \in \mathcal{MU}_{n=m}$ such that $\delta(F)$ is maximal.¹⁵ Then $F = A(\text{var}(F))$. Thus the maximal deficiency for $F \in \mathcal{MU}_{n=m}$ is $2^m - m$ (realised by $A_m \in \mathcal{UHLT}_{n=m} \cap \mathcal{UHLT}_{\delta=2^m-m}$).*

So for $m = 0, 1, 2, 3, 4, 5, 6$ variables the maximal deficiency of minimally unsatisfiable clause-sets is 1, 1, 2, 5, 12, 27, 58; in general the deficiencies of the form $2^m - m$ are central for our investigations (note that the function $m \in \mathbb{N}_0 \mapsto 2^m - m \in \mathbb{N}$ is monotonically increasing). We are now able to determine the numbers of variables and numbers of clauses possible for minimally unsatisfiable clause-sets with a given deficiency:

Theorem 6.13 *For $k \in \mathbb{N}$ let $\text{nA}(k) \in \mathbb{N}_0$ be the smallest $n \in \mathbb{N}_0$ with $2^n - n \geq k$.*

1. $\{n(F) : F \in \mathcal{UHLT}_{\delta=k}\} = \{n(F) : F \in \mathcal{MU}_{\delta=k}\} = \{n \in \mathbb{N}_0 : n \geq \text{nA}(k)\}$.
2. $\{c(F) : F \in \mathcal{UHLT}_{\delta=k}\} = \{c(F) : F \in \mathcal{MU}_{\delta=k}\} = \{n \in \mathbb{N} : n \geq \text{nA}(k) + k\}$.

Proof: Part 2 follows by Part 1, so it remains to show Part 1. By Corollary 6.11 we see $\{n(F) : F \in \mathcal{MU}_{\delta=k}\} \subseteq \{n \in \mathbb{N}_0 : n \geq \text{nA}(k)\}$, and it remains to show $\{n \in \mathbb{N}_0 : n \geq \text{nA}(k)\} \subseteq \{c(F) : F \in \mathcal{UHLT}_{\delta=k}\}$.

We first note that increasing the number of variables while keeping the deficiency constant is achieved by one non-strict full subsumption extension step. It remains to show the existence of $F \in \mathcal{UHLT}_{\delta=k}$ with $n(F) = \text{nA}(k)$. For $k = 1$ we have $F = \{\perp\}$, so assume $k > 1$. Let $F_0 := A(\text{nA}(k) - 1)$ (so $\delta(F_0) = k - 1$; note

¹⁵That is, $F \in \mathcal{MU}$, $n(F) = m$, and for all $F' \in \mathcal{MU}$ with $n(F') = m$ we have $\delta(F') \leq \delta(F)$.

$\text{nA}(k) - 1 \geq 1$). Add a variable by one step of non-strict full subsumption extension, obtaining $F_1 \in \mathcal{UHT}_{\delta=k-1}$ with one new variable, and then take a clause in F_1 without that new variable and perform one step of strict full subsumption extension (on that new variable), obtaining F_2 with $n(F_2) = n(F_1) = \text{nA}(k)$ and $\delta(F_2) = \delta(F_1) + 1 = k$. \square

$\text{nA}(k)$ by definition is the smallest $n \geq 0$ with $\delta(A_n) \geq k$ (thus the notation “ nA ”), and by Theorem 6.13 it is the smallest $n \geq 0$ such that there is $F \in \mathcal{MU}_{\delta=k}$ with $n(F) = n$. We have $\text{nA}(1) = 0$, $\text{nA}(2) = 2$, $\text{nA}(3) = \dots = \text{nA}(5) = 3$, $\text{nA}(6) = \dots = \text{nA}(12) = 4$ and $\text{nA}(13) = \dots = \text{nA}(27) = 5$. Except for the first term (note the anomaly that both A_0 and A_1 have deficiency 1), the sequence $(\text{nA}(k))_{k \in \mathbb{N}}$ is sequence <http://oeis.org/A103586> in the “On-Line Encyclopedia of Integer Sequences” ([98]); as noted there we have $\text{nA}(k) = 1 + \text{fld}(k + \text{fld}(k))$ for $k \geq 2$ (recall $\text{fld}(k) = \lfloor \log_2 k \rfloor$, that is, $\text{fld}(k)$ is the largest $n \in \mathbb{N}_0$ with $2^n \leq k$).

7. Non-Mersenne numbers

In this section we study the function $\text{nM} : \mathbb{N} \rightarrow \mathbb{N}$ via a recursive definition (Definition 7.1; see Table 1). The understanding of this recursion is the underlying topic of this section. This recursion is naturally obtained from splitting on variables with minimum occurrence in minimally unsatisfiable clause-sets, and will be used in Theorem 8.3 later. The sequence nM is sequence <http://oeis.org/A062289> in the “On-Line Encyclopedia of Integer Sequences”:

- It can be defined as the enumeration of those natural numbers containing “10” in their binary representation; in other words, exactly the numbers whose binary representation contain only 1’s are skipped.
- Thus the sequence leaves out exactly the numbers of the form $2^n - 1$ for $n \in \mathbb{N}$ (that is, 1, 3, 7, 15, 31, \dots ; sequence <http://oeis.org/A000225>), which are often called “Mersenne numbers”.
- The sequence nM consists of arithmetic progressions of slope 1 and length again the Mersenne numbers, each such progression separated by an additional step of +1.

k	1	2	3	4	5	\dots	11	12	\dots	26	27	\dots	57	58
$\text{nM}(k)$	2	4	5	6	8	\dots	14	16	\dots	30	32	\dots	62	64

Table 1: Values for $\text{nM}(k)$, $k \in \{1, \dots, 58\}$

The key deficiencies (values of k) in Table 1 are the following two classes:

1. The k -values $k = 1, 2, 5, 12, 27, 58, \dots$ (sequence <http://oeis.org/A000325>) are the deficiencies $k = 2^n - n$ of the clause-sets A_n , $n \in \mathbb{N}$, while the corresponding values $\text{nM}(k) = 2^k$ are the minimum variable-degree of the clause-sets A_n (see Lemma 2.13), as explained in Subsection 1.3.
2. The k -values 1, 4, 11, 26, 57, \dots (<http://oeis.org/A000295>) are the positions just before these deficiencies, as also discussed in Subsection 1.3; we call them “jump positions”, since precisely at these positions the function value increases by 2 for the next argument (compare Definition 7.12).

The recursion in Definition 7.1 is new, and so we can not use these characterisations, but must directly prove the basic properties; the deficiencies $k = 2^n - n$ will be handled in Corollary 7.24, while the jump positions are handled in Lemma 7.20. Later we will obtain two further alternative characterisations of nM :

- Combinatorial characterisations are obtained in Corollary 9.15, where we will see that $\text{nM}(k)$ for $k \in \mathbb{N}$ is the maximal min-var-degree for lean clause-sets or variable-minimal unsatisfiable clause-sets with deficiency k .
- In Subsection 13.2 we will develop a general recursion scheme, which has the function nM “built-in”, as shown in Theorem 13.15.

Definition 7.1 For $k \in \mathbb{N}$ let $\text{nM}(k) := 2$ if $k = 1$, while else

$$\text{nM}(k) := \max_{i \in \{2, \dots, k\}} \min(2 \cdot i, \text{nM}(k - i + 1) + i).$$

The intuition underlying Definition 7.1 of $\text{nM}(k)$, as later unfolded in Theorem 8.3, is that we want to get an upper bound on the min-var-degree of an $F \in \mathcal{MU}_{\delta=k}$ (recall Definition 2.11), and for that we consider a variable $v \in \text{var}(F)$ of minimum var-degree, set it to 0, 1, and infer an upper bound on $\text{vd}_F(v)$ from the two splitting results. The index i runs over the possible literal-degrees of v (thus we have to maximise over it), where i actually is the maximum degree over both signs, and thus we can take the minimum with $i + i = 2i$ for the var-degree. In the splitting results $\langle v \rightarrow \varepsilon \rangle * F$ ($\varepsilon \in \{0, 1\}$) the deficiency is reduced by $i - 1$, since i occurrences (i.e., clauses) and one variable are lost, and we apply recursively the lower bound $\text{nM}(k - (i - 1))$, where then the i cancelled occurrences have to be re-added.

Example 7.2 Computing $\text{nM}(k)$ for $2 \leq k \leq 5$:

1. $\text{nM}(2) = \min(2 \cdot 2, \text{nM}(2 - 2 + 1) + 2) = \min(4, 4) = 4.$
2. $\text{nM}(3) = \max(\min(2 \cdot 2, \text{nM}(3 - 2 + 1) + 2), \min(2 \cdot 3, \text{nM}(3 - 3 + 1) + 3)) = \max(\min(4, 6), \min(6, 5)) = 5.$
3. $\text{nM}(4) = \max(\min(2 \cdot 2, \text{nM}(4 - 2 + 1) + 2), \min(2 \cdot 3, \text{nM}(4 - 3 + 1) + 3), \min(2 \cdot 4, \text{nM}(4 - 4 + 1) + 4)) = \max(\min(4, 7), \min(6, 7), \min(8, 6)) = 6.$
4. $\text{nM}(5) = \max(\min(2 \cdot 2, \text{nM}(5 - 2 + 1) + 2), \min(2 \cdot 3, \text{nM}(5 - 3 + 1) + 3), \min(2 \cdot 4, \text{nM}(5 - 4 + 1) + 4), \min(2 \cdot 5, \text{nM}(5 - 5 + 1) + 5)) = \max(\min(4, 8), \min(6, 8), \min(8, 8), \min(10, 7)) = 8.$

7.1. Basic properties

We begin our investigations into $\text{nM}(k)$ by some simple bounds:

Lemma 7.3 Consider $k \in \mathbb{N}$.

1. $k + 1 \leq \text{nM}(k) \leq 2 \cdot k$ for $k \in \mathbb{N}$.
2. For $k \geq 2$ we have $\text{nM}(k) \geq 4$.

Proof: The upper bound of Part 1 follows directly from the definition (by the min-component $2i$). The lower bounds follows by induction: $\text{nM}(1) = 2 \geq 1 + 1$, while for $k > 1$ we have $\text{nM}(k) \geq \min(2k, \text{nM}(k - k + 1) + k) = \min(2k, 2 + k) = k + 2$. Part 1 follows by Part 1 and $\text{nM}(2) = 4$. \square

A basic tool for investigating sequences is the Delta-operator, which measures the differences in values between two neighbouring arguments:

Definition 7.4 For a sequence $a : I \rightarrow \mathbb{R}$, where $I \subseteq \mathbb{Z}$ is stable under increment ($n \in I \Rightarrow n + 1 \in I$), the sequence $\Delta a : I \rightarrow \mathbb{R}$ is defined for $k \in I$ by $\Delta a(k) := a(k + 1) - a(k)$ (i.e., the step in the value of the sequence from k to $k + 1$).

A few obvious properties of the Delta-operator are as follows:

1. $\Delta : \mathbb{R}^I \rightarrow \mathbb{R}^I$ is linear: $\Delta(\lambda \cdot a + \mu \cdot b) = \lambda \cdot \Delta(a) + \mu \cdot \Delta(b)$.
2. $a \in \mathbb{R}^I$ is constant iff $\Delta a = (0)$.
3. a is increasing iff $\Delta a \geq 0$, while a is strictly increasing iff $\Delta a > 0$. Here for sequences $a, b : \mathbb{R}^I \rightarrow \mathbb{R}^I$ of real numbers we use $a \leq b \Leftrightarrow \forall n \in I : a_n \leq b_n$, and $a < b \Leftrightarrow \forall n \in I : a_n < b_n$.

The first key insight is, that the next number in the sequence of non-Mersenne numbers is obtained by adding 1 or 2 to the previous number:

Lemma 7.5 *For $k \in \mathbb{N}$ holds $\Delta \text{nM}(k) \in \{1, 2\}$.*

Proof: For $k = 1$ we get $\Delta \text{nM}(1) = 2$. Now consider $k \geq 2$. We have

$$\begin{aligned} \text{nM}(k+1) &= \max(\min(4, \text{nM}(k) + 2), \max_{i \in \{3, \dots, k+1\}} \min(2i, \text{nM}(k-i+2) + i)) = \\ &= \max_{i \in \{3, \dots, k+1\}} \min(2i, \text{nM}(k-i+2) + i) = \\ &= \max_{i \in \{2, \dots, k\}} \min(2(i+1), \text{nM}(k-(i+1)+2) + (i+1)) = \\ &= \max_{i \in \{2, \dots, k\}} \min(2i+2, \text{nM}(k-i+1) + i + 1) = 1 + \max_{i \in \{2, \dots, k\}} \min(2i+1, \text{nM}(k-i+1) + i). \end{aligned}$$

Thus on the one hand we have $\text{nM}(k+1) \geq 1 + \max_{i \in \{2, \dots, k\}} \min(2i, \text{nM}(k-i+1) + i) = 1 + \text{nM}(k)$, and on the other hand $\text{nM}(k+1) \leq 1 + \max_{i \in \{2, \dots, k\}} \min(2i+1, \text{nM}(k-i+1) + i + 1) = 2 + \text{nM}(k)$. \square

Thus increasing the deficiency k by 1 increases $\text{nM}(k)$ at least by 1:

Corollary 7.6 *$\text{nM} : \mathbb{N} \rightarrow \mathbb{N}$ is strictly increasing.*

And changing $\text{nM}(a+b)$ to $\text{nM}(a) + b$ can not increase the value:

Corollary 7.7 *We have $\text{nM}(a+b) \geq \text{nM}(a) + b$ for $a \in \mathbb{N}$ and $b \in \mathbb{N}_0$, and thus $\text{nM}(a-b) \leq \text{nM}(a) - b$ for $b < a$.*

Proof: We have $\text{nM}(a+b) - \text{nM}(a) = \sum_{i=0}^{b-1} \Delta \text{nM}(a+i) \geq b \cdot 1$, whence the first inequality. Applying it yields $\text{nM}(a-b) + b \leq \text{nM}(a-b+b) = \text{nM}(a)$. \square

Instead of considering the maximum over $k-1$ cases $i \in \{2, \dots, k\}$ to compute $\text{nM}(k)$ (according to Definition 7.1), we can now simplify the recursion to only one case $i_{\text{nM}}(k) \in \{2, \dots, k\}$, and for that case also consideration of the minimum is dispensable. $i_{\text{nM}}(k)$ is the first index i in Definition 7.1, where the minimum is attained by the nM -term, that is, where $2i \geq \text{nM}(k-i+1) + i$:

Definition 7.8 *Let $i_{\text{nM}} : \mathbb{N}_{\geq 2} \rightarrow \mathbb{N}$ be defined for $k \in \mathbb{N}$, $k \geq 2$, by $i_{\text{nM}}(k) := i$ for the smallest $i \in \{2, \dots, k\}$ with $i \geq \text{nM}(k-i+1)$ (note that $k \geq \text{nM}(k-k+1) = 2$, and thus $i_{\text{nM}}(k)$ is well-defined).*

Example 7.9 *We have*

1. $i_{\text{nM}}(2) = 2$.
2. $i_{\text{nM}}(3) = 3$, since $\text{nM}(3-2+1) = 4$, $\text{nM}(3-3+1) = 2$.

3. $i_{\text{nM}}(4) = 4$, since $\text{nM}(4 - 3 + 1) = 4$, $\text{nM}(4 - 4 + 1) = 2$.
4. $i_{\text{nM}}(5) = 4$, since $\text{nM}(5 - 3 + 1) = 5$, $\text{nM}(5 - 4 + 1) = 4$.
5. $i_{\text{nM}}(6) = 5$, since $\text{nM}(6 - 4 + 1) = 5$, $\text{nM}(6 - 5 + 1) = 4$.

As promised, from $i_{\text{nM}}(k)$ we can compute $\text{nM}(k)$ by one recursive call of nM :

Lemma 7.10 *For $k \in \mathbb{N}$, $k \geq 2$, we have:*

1. $0 \leq i_{\text{nM}}(k) - \text{nM}(k - i_{\text{nM}}(k) + 1) \leq 2$.
2. $\Delta i_{\text{nM}}(k) \in \{0, 1\}$.
3. $\text{nM}(k) = \text{nM}(k - i_{\text{nM}}(k) + 1) + i_{\text{nM}}(k)$.

Proof: For Part 1 we consider the sequence $i \mapsto f_k(i) := i - \text{nM}(k - i + 1)$; this sequence starts with $f_k(2) = 2 - \text{nM}(k - 1) \leq 0$, and finishes with $f_k(k) = k - \text{nM}(1) \geq 2$, and $i_{\text{nM}}(k)$ is the smallest i with $f_k(i) \geq 0$. By Lemma 7.5 we have $\Delta f_k(i) = \Delta i(i) - \Delta \text{nM}(k - i + 1)(i) \in \{1 + 1, 1 + 2\} = \{2, 3\}$. So for $i_{\text{nM}}(k) - \text{nM}(k - i_{\text{nM}}(k) + 1) = f_k(i_{\text{nM}}(k))$ by definition we have $f_k(i_{\text{nM}}(k)) \geq 0$, while $f_k(i_{\text{nM}}(k)) \leq 2$ due to $\Delta f_k(i_{\text{nM}}(k)) \leq 3$ (otherwise $i_{\text{nM}}(k)$ wouldn't be minimal).

For Part 2 we consider the sequence $k \mapsto g_i(k) := i - \text{nM}(k - i + 1)$. Again by Lemma 7.5 we get $\Delta g_i(k) \in \{-1, -2\}$. It follows immediately $\Delta i_{\text{nM}}(k) \geq 0$. Now assume $\Delta i_{\text{nM}}(k) \geq 1$; thus $-2 \leq g_{i_{\text{nM}}(k)}(k + 1) < 0$, whence, as shown before, $g_{i_{\text{nM}}(k)+1}(k + 1) \geq -2 + 2 = 0$, and thus $\Delta i_{\text{nM}}(k) = 1$.

For Part 3 we consider the sequence $i \mapsto h_k(i) := \text{nM}(k - i + 1) + i$; by Lemma 7.5 we have $\Delta h_k(i) \in \{-1 + 1, -2 + 1\} = \{0, -1\}$. Thus, and by definition of $i_{\text{nM}}(k)$, we get $\text{nM}(k) = \max(2 \cdot 1, \dots, 2 \cdot (i_{\text{nM}}(k) - 1), h_k(i_{\text{nM}}(k))) = \max(2 i_{\text{nM}}(k) - 2, h_k(i_{\text{nM}}(k)))$. Finally $h_k(i_{\text{nM}}(k)) \geq 2 i_{\text{nM}}(k) - 2 \Leftrightarrow \text{nM}(k - i_{\text{nM}}(k) + 1) + 2 \geq i_{\text{nM}}(k)$, which holds by Part 1. \square

We obtain an alternative, functional characterisation of $i_{\text{nM}}(k)$:

Corollary 7.11 *For $k \in \mathbb{N}$, $k \geq 2$, the value $i_{\text{nM}}(k)$ is the unique $i \in \{2, \dots, k\}$ fulfilling the two inequalities*

$$\text{nM}(k - i + 1) \leq i \leq \text{nM}(k - i + 2).$$

Proof: As shown in the first part of the proof of Lemma 7.10, the sequence $i \in \{1, \dots, k\} \mapsto f_k(i) := i - \text{nM}(k - i + 1) \in \mathbb{Z}$ is strictly increasing, while the two inequalities are equivalent to $f_k(i) \geq 0$, $f_k(i - 1) \leq -1$, and so they determine i as the smallest $i \in \{2, \dots, k\}$ with $f_k(i) \geq 0$, which is the definition of $i_{\text{nM}}(k)$. \square

7.2. Characterising the jumps

After these preparations we are able to characterise the ‘‘jump positions’’, which are defined as those k where the function nM increases by 2:

Definition 7.12 *Let $J := \{k \in \mathbb{N} : \Delta \text{nM}(k) = 2\}$ be the set of **jump positions**.*

Thus $\Delta \text{nM}(k) = 1$ iff $k \notin J$, and by Table 1 we see $J = \{1, 4, 11, 26, 57, \dots\}$. Note that $\text{nM}(k) = 1 + k + |\{k' \in J : k' < k\}|$. It is useful to define two auxiliary functions:

Definition 7.13 *Let $i', h : \mathbb{N}_{\geq 2} \rightarrow \mathbb{N}$ be defined by $i'(k) := k - i_{\text{nM}}(k) + 1 \in \mathbb{N}$ for $k \in \mathbb{N}$, $k \geq 2$, while $h(k) := \text{nM}(i'(k)) \in \mathbb{N}$.*

Some basic properties:

1. We have $\Delta i'(k) = 1 - \Delta i_{nM}(k)$.
2. Thus by Lemma 7.10, Part 2, holds $\Delta i'(k) \in \{0, 1\}$.
3. By Lemma 7.10, Part 3, we have $nM(k) = h(k) + i_{nM}(k)$.
4. Thus $\Delta h(k) = \Delta nM(k) - \Delta i_{nM}(k)$.
5. By Lemmas 7.5 and 7.10, Part 2 we get $\Delta h(k) \in \{0, 1, 2\}$.
6. By Lemma 7.10, Part 1 we have $i_{nM}(k) - h(k) \in \{0, 1, 2\}$.
7. By Corollary 7.11 we have $h(k) = nM(i'(k)) \leq i_{nM}(k) \leq nM(i'(k) + 1)$.

It is instructive to consider initial values of the auxiliary functions in Table 2; this table is constructed as follows (in Example 7.18 we'll see how to construct this table row by row):

- The values for $nM(k)$ are from Table 1.
- Columns i' and i_{nM} are connected via $i_{nM} + i' = k + 1$.
- The columns i' , h duplicate the columns k , nM (but with repetitions).
- Columns nM , i_{nM} and h are connected via $nM = i_{nM} + h$.
- The values of column i_{nM} are determined according to Corollary 7.11 by the condition $h \leq i_{nM} \leq h'$, where h uses i_{nM} via i' , and h' is the next nM -value.

k	nM	ΔnM	i_{nM}	Δi_{nM}	i'	$\Delta i'$	h	Δh	$i_{nM} - h$
1	2	2	-	-	-	-	-	-	-
2	4	1	2	1	1	0	2	0	0
3	5	1	3	1	1	0	2	0	1
4	6	2	4	0	1	1	2	2	2
5	8	1	4	1	2	0	4	0	0
6	9	1	5	0	2	1	4	1	1
7	10	1	5	1	3	0	5	0	0
8	11	1	6	0	3	1	5	1	1
9	12	1	6	1	4	0	6	0	0
10	13	1	7	1	4	0	6	0	1
11	14	2	8	0	4	1	6	2	2
12	16	1	8	1	5	0	8	0	0

Table 2: Values of auxiliary functions; underlined the jump positions

First we show some further simple properties of the auxiliary functions:

Lemma 7.14 Consider $k \geq 2$.

1. If $\Delta i_{nM}(k) = 0$, then:
 - (a) $\Delta i_{nM}(k + 1) = 1$.
 - (b) $i_{nM}(k) - h(k) \in \{1, 2\}$.
 - (c) $i_{nM}(k + 1) = h(k + 1)$.
2. $\Delta i_{nM}(k) = 1 \Leftrightarrow \Delta i'(k) = 0 \Leftrightarrow \Delta h(k) = 0$.

3. If $\Delta i_{\text{nM}}(k) = 1$, then:

- (a) $k \notin J$.
- (b) $i_{\text{nM}}(k) - h(k) \in \{0, 1\}$.

Proof: For Part 1a assume $\Delta i_{\text{nM}}(k+1) = 0$ (and thus $\Delta i'(k+1) = 1$ due to $\Delta i' = 1 - \Delta i_{\text{nM}}$). Because of $\Delta h = \Delta \text{nM} - \Delta i_{\text{nM}}$ we obtain $\Delta h(k+1) \geq 1$. Thus $i_{\text{nM}}(k) = i_{\text{nM}}(k+2) \geq h(k+2) \geq h(k+1) + 1 = \text{nM}(i'(k+1)) + 1 = \text{nM}(i'(k) + 1) + 1$, contradicting $i_{\text{nM}}(k) \leq \text{nM}(i'(k) + 1)$. For the remainder of Part 1 note $\Delta h(k) = \Delta \text{nM}(k) \geq 1$.

For Part 1b note $i_{\text{nM}}(k) = i_{\text{nM}}(k+1) \geq h(k+1) \geq h(k) + 1$.

For Part 1c assume $i_{\text{nM}}(k+1) > h(k+1)$. Thus $i_{\text{nM}}(k) = i_{\text{nM}}(k+1) \geq h(k+1) + 1 \geq h(k) + 2$, whence $i_{\text{nM}}(k) = h(k) + 2$. If we would have $\Delta h(k) = 2$, then $i_{\text{nM}}(k) = i_{\text{nM}}(k+1) > h(k+1) = h(k) + 2$; thus $h(k+1) = h(k) + 1$. Now $i_{\text{nM}}(k) = h(k) + 2 = h(k+1) + 1 = \text{nM}(i'(k+1)) + 1 = \text{nM}(i'(k) + 1) + 1$, a contradiction.

Part 2 is obvious, and Part 3a follows. Finally, Part 3b follows by $i_{\text{nM}}(k+1) \leq h(k+1) + 2$ and $i_{\text{nM}}(k+1) = i_{\text{nM}}(k) + 1$, while $h(k+1) = h(k)$ due to Part 2, whence $i_{\text{nM}}(k) \leq h(k) + 1$. \square

We obtain the main characterisation of the jump positions via the auxiliary functions:

Theorem 7.15 *For $k \geq 2$ the following conditions are equivalent:*

1. $k \in J$
2. $\Delta h(k) = 2$
3. $i_{\text{nM}}(k) = h(k) + 2$
4. $\Delta i_{\text{nM}}(k-1) = 1$ and $i_{\text{nM}}(k-1) = h(k-1) + 1$
5. $\Delta i_{\text{nM}}(k-2) = \Delta i_{\text{nM}}(k-1) = 1$ (yielding various equivalent forms via Lemma 7.14, Part 2).

Proof: Condition 1 implies Condition 2 due to $\Delta i_{\text{nM}}(k) = 0$ in case of $k \in J$ by Lemma 7.14, Part 3a. Condition 2 implies Condition 3, since $\Delta h(k) = 2$ implies $\Delta i_{\text{nM}}(k) = 0$, and so by Lemma 7.14, Part 1c we have $i_{\text{nM}}(k) = i_{\text{nM}}(k+1) = h(k+1)$, while the assumption says $h(k+1) = h(k) + 2$. In turn Condition 3 implies Condition 1, since by Lemma 7.14, Part 3b we get $\Delta i_{\text{nM}}(k) = 0$, and thus $\Delta \text{nM}(k) = \Delta h(k)$, where in case of $\Delta h(k) \leq 1$ we would have $h(k) + 2 = i_{\text{nM}}(k) \leq \text{nM}(i'(k) + 1) = \text{nM}(i'(k+1)) = h(k+1) \leq h(k) + 1$. So now we can freely use the equivalence of these three conditions.

Condition 3 implies Condition 4, since we have $\Delta i_{\text{nM}}(k) = 0$, and thus $\Delta i_{\text{nM}}(k-1) = 1$ with Lemma 7.14, Part 1a, from which we furthermore get $i_{\text{nM}}(k) = i_{\text{nM}}(k-1) + 1$ and $h(k-1) = h(k)$, and so $i_{\text{nM}}(k-1) = i_{\text{nM}}(k) - 1 = h(k) + 1 = h(k-1) + 1$. Condition 4 implies Condition 5, since in case of $\Delta i_{\text{nM}}(k-2) = 0$ we had $i_{\text{nM}}(k-1) = h(k-1)$ with Lemma 7.14, Part 1c. In turn Condition 5 implies Condition 3, since $i_{\text{nM}}(k) = i_{\text{nM}}(k-1) + 1 = i_{\text{nM}}(k-2) + 2$, while $h(k) = h(k-1) = h(k-2)$, where by definition $i_{\text{nM}}(k-2) \geq h(k-2)$ holds, whence $i_{\text{nM}}(k) \geq h(k) + 2$, which implies $i_{\text{nM}}(k) = h(k) + 2$. \square

We understand now the shape of the four Δ -sequences:

Corollary 7.16 *By definition the sequence $(\Delta \text{nM}(k))_{k \in \mathbb{N}}$ is 1 except at the jump positions k , where it is 2. The other three Δ -sequences are shaped as follows:*

1. The sequence $(\Delta i_{\text{nM}}(k))_{k \in \mathbb{N}, k \geq 2}$ consists of alternating 0, 1's except the two positions $k - 2, k - 1$ before a jump position $k \in J$, where we have two consecutive 1's (while at the jump position we have 0).
2. The sequence $(\Delta i'(k))_{k \in \mathbb{N}, k \geq 2}$ consists of alternating 0, 1's except two positions before a jump position k , where we have two consecutive 0's.
3. The sequence $(\Delta h(k))_{k \in \mathbb{N}, k \geq 2}$ consists of alternating 0, 1's except two positions before a jump position k , where we have two consecutive 0's, followed by a 2 at the jump position k , which is followed by 0.

Proof: Part 1: By Lemma 7.14, 1a we have $\Delta i_{\text{nM}}(k) = 0 \Rightarrow \Delta i_{\text{nM}}(k + 1) = 1$, while by Theorem 7.15, Part 5 we have $\Delta i_{\text{nM}}(k) = \Delta i_{\text{nM}}(k + 1) = 1 \Rightarrow k + 2 \in J$, and by Lemma 7.14, Part 3a we have $k \in J \Rightarrow \Delta i_{\text{nM}}(k) = 0$.

Part 2 follows from Part 1 by $\Delta i' = 1 - \Delta i_{\text{nM}}$.

Part 3: By Lemma 7.14, Part 2 the 0's in the sequence Δh are precisely the 1's in the sequence Δi_{nM} , while a 0 of Δi_{nM} translates into a 2 precisely at the jump positions by Theorem 7.15, Part 2. The assertion follows now by Part 1. \square

Especially instructive is understanding of the i' -sequence:

Corollary 7.17 *The i' -sequence $(i'(k))_{k \in \mathbb{N}, k \geq 2}$ consists of doublets m, m for consecutive $m = 1, 2, \dots$, except for $k \in J \setminus \{1\}$, where we have at positions $k - 2, k - 1, k$ a triplet m, m, m . These triplet-values occur exactly when $m \in J$.*

Proof: The doublet/triplet structure follows by Corollary 7.16, Part 2. Now consider a triplet $i'(k - 2) = i'(k - 1) = i'(k) = m$ for $k \in J \setminus \{1\}$, $m \in \mathbb{N}$. By definition we have $\Delta nM(m) = \Delta h(k)$ (due to $h(k) = h(i'(k)) = nM(m)$, $h(k + 1) = nM(i'(k + 1)) = nM(i'(k) + 1) = nM(m + 1)$). By Theorem 7.15, Part 2 we have thus have $\Delta nM(m) = 2$, i.e., $m \in J$. The triplets do not leave out some jump-value in J , since for $m \in J$ and for the last position k with $i'(k) = m$ we have $\Delta nM(m) = \Delta h(k)$. \square

Example 7.18 *We see now how we can built up the three columns k, nM, i' of Table 2 together with an enumeration of the set J , which is built up as the set I (I is an initial part of J , which in the limit becomes J):*

1. We start with the first row $k := 1$, initialising the value n of $nM(1) = n$ to $n := 2$, while i_{nM} is undefined; $k = 1$ is the first jump position, that is, $I := \{1\}$.
2. We go to the second row, $k := k + 1$. We update $n := n + 2$ and initialise the running value of $i'(k) = m$ to $m := 1$.
3. We repeat the following steps ad infinitum:
 - (a) If $m \in I$, then three rows are created:
 - i. $i_{\text{nM}}(k) = n$, $i'(k) = m$, $k := k + 1$, $n := n + 1$
 - ii. $i_{\text{nM}}(k) = n$, $i'(k) = m$, $k := k + 1$, $n := n + 1$
 - iii. $i_{\text{nM}}(k) = n$, $i'(k) = m$, $I := I \cup \{k\}$, $k := k + 1$, $n := n + 2$, $m := m + 1$.
 - (b) If $m \notin I$, then two rows are created:
 - i. $i_{\text{nM}}(k) = n$, $i'(k) = m$, $k := k + 1$, $n := n + 1$
 - ii. $i_{\text{nM}}(k) = n$, $i'(k) = m$, $k := k + 1$, $n := n + 1$, $m := m + 1$.

Next we show that $i'(k)$ for jump positions is the previous jump position:

Lemma 7.19 *For $k \in J$, $k \geq 2$, holds $i'(k) = \max\{k' \in J : k' < k\}$.*

Proof: We prove the assertion by induction on k (regarding the enumeration of J). We have $i'(4) = 1$, and so the induction holds for $k = 4$, the smallest jump position $k \geq 2$. Now assume that the assertion holds for all elements of $J \cap \{1, \dots, k-1\}$, where $k > 4$, and we have to show the assertion for k . By Corollary 7.17 we know $i'(k) \in J$, where $2 \leq i'(k) < k$. Assume there is $m \in J$ with $i'(k) < m < k$. By induction hypothesis we get $i'(k) \leq i'(m) < m$. However by Lemma 7.14 we get $\Delta i'(m) = 1$, and thus $i'(k) > i'(m)$ (since $k > m$). \square

We obtain the promised characterisation of the jump positions:

Lemma 7.20 *We have $J = \{2^{m+1} - (m+1) - 1 : m \in \mathbb{N}\}$.*

Proof: Let k_m for $m \in \mathbb{N}$ be the m th element of J ; so the assertion is $k_m = 2^{m+1} - m - 2$. We have $k_1 = 4 - 1 - 2 = 1 = \min J$; in the remainder assume $m \geq 2$. We prove the assertion by induction, in parallel with $i_{\text{nM}}(k_m) = 2^{m+1} - 2^m$. For $m = 2$ we have $k_2 = 8 - 2 - 2 = 4 = \min J \setminus \{1\}$, while $i_{\text{nM}}(4)$ is the smallest $i \in \{2, 3, 4\}$ with $i \geq \text{nM}(5 - i)$, which yields $i_{\text{nM}}(4) = 4 = 2^3 - 2^2$. Now we consider the induction step, from $m-1$ to m . The induction hypothesis yields $k_{m-1} = 2^m - m - 1$ and $i_{\text{nM}}(k_{m-1}) = 2^m - 2^{m-1}$. Lemma 7.19 yields $i'(k_m) = k_{m-1}$, from which by $i'(k_m) = k_m - i_{\text{nM}}(k_m) + 1$ follows

$$k_m = 2^m - m - 2 + i_{\text{nM}}(k_m).$$

Via a telescoping series we get

$$i_{\text{nM}}(k_m) = \Delta i_{\text{nM}}(k_m - 1) + \dots + \Delta i_{\text{nM}}(k_{m-1}) + i_{\text{nM}}(k_{m-1}).$$

By Corollary 7.16, Part 1 the sequence $\Delta i_{\text{nM}}(k_{m-1}), \dots, \Delta i_{\text{nM}}(k_m - 1)$ has the form $0, 1, 0, 1, \dots, 0, 1, 1$, and thus their sum has the value $\frac{1}{2}(k_m - k_{m-1} - 1) + 1$. So we get

$$\begin{aligned} i_{\text{nM}}(k_m) &= \frac{1}{2}(k_m - k_{m-1} - 1) + 1 + i_{\text{nM}}(k_{m-1}) = \\ &= \frac{1}{2}(2^m - m - 2 + i_{\text{nM}}(k_m) - 2^m + m + 1 - 1) + 1 + 2^m - 2^{m-1} = \\ &= \frac{1}{2}i_{\text{nM}}(k_m) - 1 + 1 + 2^m - 2^{m-1}, \end{aligned}$$

from which $i_{\text{nM}}(k_m) = 2^{m+1} - 2^m$ follows. Finally $k_m = 2^m - m - 2 + 2^{m+1} - 2^m = 2^{m+1} - m - 2$. \square

7.3. Applications

Now the closed formula for $\text{nM}(k)$ can be proven (recall $\text{fld}(x) = \lfloor \log_2(x) \rfloor$):

Theorem 7.21 *For $k \in \mathbb{N}$ holds $\text{nM}(k) = k + \text{fld}(k+1) + \text{fld}(k+1)$.*

Proof: Let $g(k) := \text{fld}(k+1) + \text{fld}(k+1)$ and $f(k) := k + g(k)$ (so $\text{nM}(k) = f(k)$ is to be shown, for $k \geq 1$). We have $f(1) = 1 + \text{fld}(2) + \text{fld}(2) = 1 + \text{fld}(3) = 2 = \text{nM}(1)$. We will now prove that the function $g(k)$ changes values exactly at the transitions $k \mapsto k+1$ for $k \in J$, that is, for indices $k = k_m := 2^{m+1} - m - 2$ (using Lemma

7.20) with $m \in \mathbb{N}$ we have $\Delta g(k_m) = 1$, while otherwise we have $\Delta g(k_m) = 0$, from which the assertion follows (by the definition of J).

We have $g(1) = 1$ and $g(2) = 2$. Now consider $m \in \mathbb{N}$ and $k_m + 1 \leq k \leq k_{m+1}$. We show $g(k) = m + 1$, which proves the claim. Note that $g(k)$ is monotonically increasing. Now $g(k) \geq g(k_m + 1) = \lfloor \text{ld}(2^{m+1} - m + \lfloor \text{ld}(2^{m+1} - m) \rfloor) \rfloor = \lfloor \text{ld}(2^{m+1} - m + m) \rfloor = m + 1$ and $g(k) \leq g(k_{m+1}) = \lfloor \text{ld}(2^{m+2} - m - 2 + \lfloor \text{ld}(2^{m+2} - m - 2) \rfloor) \rfloor \leq \lfloor \text{ld}(2^{m+2} - m - 2 + m + 1) \rfloor = \lfloor \text{ld}(2^{m+2} - 1) \rfloor = m + 1$. \square

Using $nA : \mathbb{N} \rightarrow \mathbb{N}_0$ as introduced in Theorem 6.13, we get:

Corollary 7.22 *For $k \in \mathbb{N}$ holds $nM(k) = k - 1 + nA(k + 1)$.*

From Theorem 7.21 we obtain very precise bounds for $nM(k)$:

Corollary 7.23 *$k + \text{fld}(k + 1) \leq nM(k) \leq k + 1 + \text{fld}(k)$ holds for $k \in \mathbb{N}$.*

Proof: The lower bound follows trivially. The upper bound holds (with equality) for $k \leq 2$, so assume $k \geq 3$. We have to show $g(k) = \text{fld}(k + 1 + \text{fld}(k + 1)) \leq 1 + \text{fld}(k)$, which follows from $\text{ld}(k + 1 + \text{fld}(k + 1)) \leq 1 + \text{ld}(k)$. Now $\text{ld}(k + 1 + \text{fld}(k + 1)) \leq \text{ld}(k + 1 + \text{ld}(k + 1)) \leq \text{ld}(k + k) = 1 + \text{ld}(k)$. \square

Note that $(k + 1 + \text{fld}(k)) - (k + \text{fld}(k + 1)) \in \{0, 1\}$, where this difference is zero iff $k + 1$ is a power of 2. Finally we can prove the already mentioned characterisation, which motivates the terminology of “non-Mersenne numbers”, namely that $(nM(k))_{k \in \mathbb{N}}$ enumerates $\mathbb{N} \setminus \{2^n - 1 : n \in \mathbb{N}\}$.¹⁶ For that we consider the positions directly after the jump positions, which by Lemma 7.20 are the positions $2^n - n$ for $n \geq 2$. From that position on until the next jump position, which is $2^{n+1} - n - 2$, the nM -values increase constantly by 1 per step. So we just need to understand the values of $nM(2^n - n)$, to understand all of nM , which is achieved as follows (note that $(2^{n+1} - n - 2) - (2^n - n) = 2^n - 2$):

Corollary 7.24 *Consider $n \in \mathbb{N}$, $k := 2^n - n$, and $m \in \mathbb{N}_0$ with $m \leq 2^n - 1$.*

1. $nM(k) = 2^n$.
2. *More generally for $m < 2^n - 1$ holds $nM(k + m) = 2^n + m$.*
3. *For $m = 2^n - 1$ we have $k + m = 2^{n+1} - (n + 1)$, and thus $nM(k + m) = 2^{n+1}$.*

Proof: By Theorem 7.21, Part 1 follows with $nM(2^n - n) = 2^n - n + \text{fld}(2^n - n + 1 + \text{fld}(2^n - n + 1)) = 2^n - n + \text{fld}(2^n - n + 1 + (n - 1)) = 2^n - n + \text{fld}(2^n) = 2^n$. Part 2 follows by Lemma 7.20, and Part 3 follows by Part 1. \square

Besides $nM(2^n - n) = 2^n$ also the following special value is of importance:

Corollary 7.25 *For $n \in \mathbb{N}$, $n \geq 2$, we have $nM(2^n - n - 1) = 2^n - 2$.*

It is also useful to have simple formulas for the $i_{nM}(k)$ -values around the jump positions:

Corollary 7.26 *For $n \in \mathbb{N}$, $n \geq 3$ the values of $i_{nM}(2^n - n + m)$ are as follows, using $p := 2^{n-1}$ (where for $m = -4$ we need $n \geq 4$):*

m	-4	-3	-2	-1	0	1	2	3	4
i_{nM}	$p - 2$	$p - 2$	$p - 1$	p	p	$p + 1$	$p + 1$	$p + 2$	$p + 2$

¹⁶Note that we are not speaking of “non-Mersenne primes”.

Proof: We have $i_{\text{nM}}(2^n - n) = 2^{n-1}$ by Corollary 7.11:

$$\begin{aligned} 2^{n-1} &\geq \text{nM}(2^n - n - 2^{n-1} + 1) = \text{nM}(2^{n-1} - (n-1)) = 2^{n-1} \\ 2^{n-1} &\leq \text{nM}(2^n - n - 2^{n-1} + 2) = \text{nM}(2^{n-1} - (n-1) + 1) = 2^{n-1} + 1. \end{aligned}$$

The remaining values follow by Corollary 7.16, Part 1. \square

We conclude with an alternative characterisation of the jump-set J :

Corollary 7.27 *For $k \in \mathbb{N}$ the following conditions are equivalent:*

1. $\text{nM}(k) < 2 \cdot i_{\text{nM}}(k) - 1$.
2. $\text{nM}(k) = 2 \cdot i_{\text{nM}}(k) - 2$.
3. $k \in J$, that is, $k = 2^{m+1} - m - 2$ for some $m \in \mathbb{N}$.

Proof: If $k \in J$, then by Theorem 7.15, Part 3, we have $\text{nM}(k) = 2 \cdot i_{\text{nM}}(k) - 2 < 2 \cdot i_{\text{nM}}(k) - 1$. And if $k \notin J$, then by the same lemma we have $i_{\text{nM}}(k) \leq h(k) + 1$, and thus $\text{nM}(k) = h(k) + i_{\text{nM}}(k) \geq 2 \cdot i_{\text{nM}}(k) - 1$. \square

8. The upper bound for VDM

In a sense the main auxiliary lemma of this report is the following Lemma 8.1 (especially Part 3) on the deficiencies obtained when splitting a saturated minimally unsatisfiable clause-set. This receives its importance from the fact that every minimally unsatisfiable clause-set can be saturated (recall Subsection 3.2; this method was first applied in this context in [7]).

The first task here is to determine the deficiency after assigning a variable. In general we have for $F \in \mathcal{CLS}$, $v \in \text{var}(F)$ and $\varepsilon \in \{0, 1\}$, that $\delta(\langle v \rightarrow \varepsilon \rangle * F) = \delta(F) - m_\varepsilon + 1 + r_\varepsilon$, where m_ε is the number of clauses satisfied by the assignment, that is, the clauses containing \bar{v} for $\varepsilon = 0$ resp. the clauses containing v for $v = 1$, while r_ε is the number of variables in F which occur only in these clauses, that is, $r_0 = |\text{var}(\{C \in F : \bar{v} \in C\}) \setminus \text{var}(\{C \in F : \bar{v} \notin C\})|$ and $r_1 = |\text{var}(\{C \in F : v \in C\}) \setminus \text{var}(\{C \in F : v \notin C\})|$. We want $r_\varepsilon = 0$, and this is guaranteed in Lemma 8.1 by choosing v of minimal degree, where v occurs in both signs.

If now $F \in \mathcal{SMU}$ holds, then $\langle v \rightarrow \varepsilon \rangle * F \in \mathcal{MU}$ by Lemma 3.15, Part 1. Since this is an essential insight (which the first author got from [23]), we prove this here for the sake of completeness, where w.l.o.g. $\varepsilon = 1$ and $F' := \langle v \rightarrow 1 \rangle * F$. If there were $C \in F'$ with $F' \setminus \{C\} \in \mathcal{USAT}$, then we have to consider two cases:

- If $C \in F$, then we had the partial saturation $S(F, C, v) \in \mathcal{USAT}$ (recall Definitions 3.5, 3.7, and note that $C \cup \{v\} \notin F$, using $F \in \mathcal{MU}$), due to $\langle v \rightarrow 0 \rangle * S(F, C, v) = \langle v \rightarrow 0 \rangle * F \in \mathcal{USAT}$ and $\langle v \rightarrow 1 \rangle * S(F, C, v) = F' \setminus \{C\} \in \mathcal{USAT}$, contradicting saturation of F .
- So we must have that $C' := C \cup \{\bar{v}\} \in F$. But now $G := F \setminus \{C'\} \in \mathcal{USAT}$, due to $\langle v \rightarrow 0 \rangle * G = \langle v \rightarrow 0 \rangle * F \in \mathcal{USAT}$ and $\langle v \rightarrow 1 \rangle * G = F' \in \mathcal{USAT}$, contradicting the minimality of F , which concludes the proof.

Lemma 8.1 *Consider $F \in \mathcal{CLS}$ and $v \in \text{var}(F)$, and let $m_0 := \text{ld}_F(\bar{v})$ and $m_1 := \text{ld}_F(v)$. Assume v is not pure in F (i.e., $m_0, m_1 \geq 1$), and that v realises the minimum var-degree (i.e., $\text{vd}_F(v) = \mu\text{vd}(F)$). Consider $\varepsilon \in \{0, 1\}$.*

1. $\text{var}(\langle v \rightarrow \varepsilon \rangle * F) = \text{var}(F) \setminus \{v\}$ and $n(\langle v \rightarrow \varepsilon \rangle * F) = n(F) - 1$.

2. $\delta(\langle v \rightarrow \varepsilon \rangle * F) = \delta(F) - m_\varepsilon + 1$, where $\delta(F) - m_\varepsilon + 1 < \delta(F)$ iff $m_\varepsilon \geq 2$.
3. If $F \in \mathcal{SMU}_{\delta=k}$ for $k \in \mathbb{N}$, then $\langle v \rightarrow \varepsilon \rangle * F \in \mathcal{MU}_{\delta=k-m_\varepsilon+1}$, where $m_\varepsilon \leq k$.

Proof: Part 1 follows by the fact, that due to degree-minimality and non-pureness of v , no variable can have all its occurrences only in clauses containing v resp. \bar{v} . Part 2 follows by Part 1 and the general fact $c(\langle v \rightarrow \varepsilon \rangle * F) = c(F) - m_\varepsilon$. Part 3 follows by Lemma 3.15, Part 1 (as proven above) together with Part 2, where $m_\varepsilon \leq k$ follows by the general fact $\delta(F) \geq 1$ for $F \in \mathcal{MU}$. \square

If in the situation of Lemma 8.1, Part 3 (the essential part for us) the value of m_ε is minimal, i.e., $m_\varepsilon = 1$, then we have $\delta(\langle v \rightarrow \varepsilon \rangle * F) = \delta(F) = k$, while if m_ε is maximal, i.e., $m_\varepsilon = k$, then we have $\delta(\langle v \rightarrow \varepsilon \rangle * F) = 1$. The deficiency is strictly decreased for both splitting results $\langle v \rightarrow \varepsilon \rangle * F$ iff v is nonsingular.

Example 8.2 For $\mathcal{F}_n \in \mathcal{SMU}'_{\delta=2}$ ($n \geq 2$; recall Example 3.3) and each $v \in \text{var}(\mathcal{F}_n)$, $\varepsilon \in \{0, 1\}$, holds $\langle v \rightarrow \varepsilon \rangle * \mathcal{F}_n \in \mathcal{MU}_{\delta=1}$.

The definition of $\text{nM}(k)$ (recall Definition 7.1) matches the recursion-structure of Lemma 8.1, and we obtain an upper bound on the min-var-degree for minimally unsatisfiable clause-sets:

Theorem 8.3 For all $k \in \mathbb{N}$ and $F \in \mathcal{MU}_{\delta \leq k} \setminus \{\{\perp\}\}$ we have $\mu\text{vd}(F) \leq \text{nM}(k)$. More precisely, there exists a variable $v \in \text{var}(F)$ with $\text{vd}_F(v) \leq \text{nM}(k)$ and $\text{ld}_F(v), \text{ld}_F(\bar{v}) \leq k$.

Proof: The assertion is known for $k = 1$, so assume $k > 1$, and we apply induction on k . Assume $\delta(F) = k$ (due to $k > 1$ we have $n(F) > 1$). Saturate F and obtain F' . Consider a variable $v \in \text{var}(F')$ realising the min-var-degree of F' . If $\text{vd}_{F'}(v) = 2$ then we are done, so assume $\text{vd}_{F'}(v) \geq 3$. Let $i := \max(\text{ld}_{F'}(v), \text{ld}_{F'}(\bar{v}))$; so $\text{vd}_{F'}(v) \leq 2i$. W.l.o.g. assume that $i = \text{ld}_{F'}(v)$. By Lemma 8.1 we get $2 \leq i \leq k$. Applying the induction hypothesis and Lemma 8.1 we obtain a variable $w \in \text{var}(G)$ for $G := \langle v \rightarrow 1 \rangle * F$ with $\text{vd}_G(w) \leq \text{nM}(k - i + 1)$. By definition we have $\text{vd}_{F'}(w) \leq \text{vd}_G(w) + \text{ld}_{F'}(v)$. Altogether $\mu\text{vd}(F) \leq \min(2i, \text{nM}(k - i + 1) + i) \leq \text{nM}(k)$. \square

The upper bound on the minimum variable degree of Theorem 8.3 is not sharp, and will be further investigated from Section 12 on. However the bound is attained for infinitely many deficiencies, and we demonstrate now that the jump positions (the set J ; recall Definition 7.12) are such deficiencies. Moreover, to investigate the remaining deficiencies, we show that they always have at least two variables realising the bound (if the bound is attained at all); this will be used to prove Theorem 14.4. So we consider “extremal” $F \in \mathcal{MU}_{\delta=k}$ with $\mu\text{vd}(F) = \text{VDM}(k)$, and we show that such extremal clause-sets have at least two different variables of minimal degree, if $k \notin J$. First, it is useful to have a notation for the set of variables of minimal degree:

Definition 8.4 For $F \in \mathcal{CLS}$ let $\text{var}_{\mu\text{vd}}(F) \subseteq \text{var}(F)$ be the set of variables of minimal degree, that is, $\text{var}_{\mu\text{vd}}(F) := \{v \in \text{var}(F) : \text{vd}_F(v) = \mu\text{vd}(F)\}$.

Obviously $\text{var}_{\mu\text{vd}}(F) \neq \emptyset$ iff $n(F) > 0$, and $\text{var}_{\mu\text{vd}}(F) = \text{var}(F)$ holds iff F is variable-regular.

Lemma 8.5 Consider $k \in \mathbb{N}$.

1. For $k \notin J$ and $F \in \mathcal{MU}_{\delta=k}$ with $\mu\text{vd}(F) = \text{nM}(k)$ we have $|\text{var}_{\mu\text{vd}}(F)| \geq 2$.

2. For $k \in J$ there is $F \in \mathcal{UHLT}_{\delta=k}$ with $\mu\text{vd}(F) = \text{nM}(k)$ and $|\text{var}_{\mu\text{vd}}(F)| = 1$.

Proof: First assume $k \notin J$; we have to show the existence of different $v, w \in \text{var}_{\mu\text{vd}}(F)$. W.l.o.g. F is saturated. Consider $v \in \mu\text{vd}(F)$. By Corollary 7.27 we have $\text{nM}(k) \geq 2 \cdot \text{i}_{\text{nM}}(k) - 1$. Because of $\text{ld}_F(v) + \text{ld}_F(\bar{v}) = \text{nM}(k)$ thus w.l.o.g. $e_1 := \text{ld}_F(v) \geq \text{i}_{\text{nM}}(k)$. Let $F' := \langle v \rightarrow 1 \rangle * F$. So $\delta(F') = k - e_1 + 1$. Recall $\text{nM}(k) = \text{nM}(k - \text{i}_{\text{nM}}(k) + 1) + \text{i}_{\text{nM}}(k)$ (Lemma 7.10, Part 3), and thus $\text{nM}(k) \geq \text{nM}(k - e_1 + 1) + e_1$. Since $n(F) \geq 2$, we can consider $w \in \text{var}_{\mu\text{vd}}(F')$. We have $\text{vd}_{F'}(w) \leq \text{nM}(k - e_1 + 1)$ and $\text{vd}_F(w) = \text{vd}_{F'}(w) + e_1$. Thus $w \in \text{var}_{\mu\text{vd}}(F)$.

Now assume $k \in J$, that is, $k = 2^{m+1} - m - 2$ for $m \geq 1$. For $k = 1$ we have the example $\{1, -1\}$, so assume $k \geq 2$. Then we have $\text{nM}(k) = 2^{m+1} - 2$. Now we obtain an example from A_{m+1} by performing one strict full subsumption resolution: The resolution variable occurs $2^{m+1} - 2$ times, the other $m - 1$ variables occur $2^{m+1} - 1$ times. \square

In Lemma 12.12 we formulate the sharpness of the upper bound of Theorem 8.3 for these cases.

9. The min-var-degree upper bound for \mathcal{LEAN}

In this section we prove Theorem 9.10, the upper bound $\text{nM}(k)$ on the min-var-degree for lean clause-sets of surplus k , and the sharpness of this bound for any class between \mathcal{VMU} and \mathcal{LEAN} in Theorem 9.14. The proof consists in lifting Theorem 8.3 to the general case in Subsection 9.2, while sharpness of the upper bound is considered in Subsection 9.3.

As a preparation, in Subsection 9.1 we introduce the class \mathcal{SED} of clause-sets, where deficiency and surplus coincide; this class is also interesting in its own right. Theorem 9.5 characterises the elements of \mathcal{SED} as those clause-sets, where after removal of all clauses containing any given variable we obtain a matching-satisfiable clause-set. It follows (Corollary 9.6) that unsatisfiable elements of \mathcal{SED} are variable-minimally unsatisfiable.

9.1. Clause-sets with extremal surplus

We consider the task of generalising Theorem 8.3 to $F \in \mathcal{LEAN}$. Consider an arbitrary (multi-)clause-set F . Consider a set of variables $\emptyset \neq V \subseteq \text{var}(F)$ realising the surplus of F , i.e., such that $\delta(F[V])$ is minimal (recall Definition 2.19). If $F[V]$ would be satisfiable, then a satisfying assignment would give a non-trivial autarky for F . Assuming that F is lean thus yields that $F[V]$ must be unsatisfiable. So there exists a minimally unsatisfiable $F' \subseteq F[V]$. If now $\text{var}(F') \neq \text{var}(F[V]) = V$ would be the case, then we would loose control over the deficiency of F' . But this can not happen, as we will show in Corollary 9.7, namely $F[V]$ here is *variable-minimally unsatisfiable*. We conclude this from the fact that $F[V]$ has a special structure, namely it belongs to the following class of clause-sets with maximal surplus (relative to the deficiency), whose basic theory we develop first:

Definition 9.1 Let the class $\mathcal{SED} \subset \mathcal{CLS}$ (“*surplus equal deficiency*”) consist of those clause-sets $F \in \mathcal{CLS}$ with $\sigma(F) = \delta(F)$.

So for $F \in \mathcal{CLS}$ with $n(F) > 0$ we have $F \in \mathcal{SED} = \{F \in \mathcal{CLS} : \sigma(F) = \delta(F)\}$ iff for all $\emptyset \subset V \subseteq \text{var}(F)$ holds $\delta(F[V]) \geq \delta(F)$. It seems the class \mathcal{SED} crosses the classes considered in this report in interesting extremal cases.

Example 9.2 Some basic examples:

1. We have $\top \in \mathcal{SED}$ and $\{\perp\} \notin \mathcal{SED}$, and more generally, for every $F \in \mathcal{SED}$ we have $\perp \notin F$.
2. For $F := \{\{1\}, \{2\}\}$ we have $\sigma(F) = \delta(F) = 0$, and thus $F \in \mathcal{SED}$. On the other hand, for the multi-clause-set $F' := \{2 * \{1\}, \{2\}\}$ we have $\delta(F') = 1$, while still $\sigma(F') = 0$, and thus $F' \notin \mathcal{SED}$.
3. For every $F \in \mathcal{CLS}$ with $c(F) \leq 2$ and $\perp \notin F$ holds $F \in \mathcal{SED}$.
4. $A_n \in \mathcal{SED}$ for $n \geq 1$ (Example 2.20).
5. $\mathcal{MU}_{\delta=1} \setminus \{\{\perp\}\} \subset \mathcal{SED}$ (since for $F \in \mathcal{MU} \setminus \{\{\perp\}\}$ holds $\sigma(F) \geq 1$).
6. $\mathcal{F}_n \in \mathcal{SED}$ for $n \geq 2$ (Example 3.3).
7. For $F := \{\{1, 2, 3\}, \{1, 2, -3\}, \{1, -2\}, \{-1, 2\}, \{-1, -2\}\}$ we have $F \in \mathcal{MU}$ with $\delta(F) = 2$, but $\sigma(F) = 1$, and thus $F \notin \mathcal{SED}$.
8. In Definition 10.5 we introduce the class $\mathcal{MLCR} \subset \mathcal{SED} \cap \mathcal{SAT} \cap \mathcal{MLEAN}$, and Example 10.7 shows elements of this class.
9. See also Example 10.11 and Question 10.12.

Finally we note that $F \in \mathcal{SED}$ if $F' \in \mathcal{SED}$, where F' is the multi-clause-set obtained from F by forgetting all signs of the literals, i.e., replacing clauses $C \in F$ by $\text{var}(C)$ (since $\delta(F') = \delta(F)$ and $\sigma(F') = \sigma(F)$).

The class \mathcal{SED} is not invariant under multiplicities; consider a multi-clause-set F and the underlying clause-set F' :

1. If $F' \in \mathcal{SED}$, then in general we do not have $F \in \mathcal{SED}$ (Example 9.2).
2. However in general holds $F \in \mathcal{SED} \Rightarrow F' \in \mathcal{SED}$, since if we would have $F' \notin \mathcal{SED}$, then $\sigma(F') < \delta(F')$, and adding a duplicated clause to a multi-clause-set increases δ by +1, while σ is at most increased by +1 (it may also stay unchanged).

We proceed by showing in the following Lemma 9.3, that $F \in \mathcal{SED}$ iff for every strict subset V of variables of F , the number of clauses of F with all variables contained in V is at most $|V|$. This basic but fundamental characterisation also yields a stronger Corollary 9.4 than the above “ $F \in \mathcal{SED} \Rightarrow F' \in \mathcal{SED}$ ”. We use in this subsection for $F \in \mathcal{CLS}$ and $V \subseteq \mathcal{VA}$ the notation $F^V \in \mathcal{CLS}$, which is the sub-multi-clause-set of F (i.e., $F^V \leq F$) consisting of all $C \in F$ with $\text{var}(C) \subseteq V$ (with the same multiplicities), that is, for $C \in \mathcal{CL}$ with $\text{var}(C) \subseteq V$ we set $F^V(C) := F(C)$, while otherwise $F^V(C) := 0$. So we have $F(\emptyset) = F(\perp)$ and $F^{\text{var}(F)} = F$, and for $V \subseteq V'$ and $F \leq F'$ holds $F^V \leq F'^{V'}$; fundamental is

$$c(F[V]) + c(F^{\text{var}(F) \setminus V}) = c(F[\text{var}(F) \setminus V]) + c(F^V) = c(F).$$

We always have $\text{var}(F^V) \subseteq V$, but also for $V \subseteq \text{var}(F)$ we might have $\text{var}(F^V) \subset V$, for example for $F := \{\{1, 2\}, \{1, 3\}\}$ we have $F^{\{2,3\}} = \top$ (so $\text{var}(F^{\{2,3\}}) = \emptyset$). Obviously we have $F^{\text{var}(F^V)} = F^V$.

Lemma 9.3 *For a multi-clause-set F the following three conditions are equivalent:*

- (i) $F \in \mathcal{SED}$.
- (ii) $\forall \emptyset \subseteq V \subset \text{var}(F) : c(F^V) \leq |V|$.

(iii) $\forall \emptyset \subseteq V \subseteq \text{var}(F) : \delta(F^V) \leq 0$.

Proof: First we show the equivalence of Conditions (i) and (ii). For $\emptyset \subset V' \subseteq \text{var}(F)$ let $V := \text{var}(F) \setminus V'$, which runs through all $\emptyset \subseteq V \subseteq \text{var}(F)$. We have $c(F[V']) = c(F) - c(F^V)$, and thus we get:

$$\begin{aligned} \delta(F[V']) \geq \delta(F) &\Leftrightarrow c(F[V']) - |V'| \geq c(F) - n(F) \Leftrightarrow \\ c(F) - c(F^V) - |V'| &\geq c(F) - n(F) \Leftrightarrow c(F^V) \leq n(F) - |V'| = |V|, \end{aligned}$$

and thus indeed (i) and (ii) are equivalent. Condition (iii) implies trivially (ii), since $\delta(F^V) = c(F^V) - n(F^V) \geq c(F^V) - |V|$. Finally assume that (ii) holds, and consider $\emptyset \subseteq V \subseteq \text{var}(F)$. For $V' := \text{var}(F^V)$ by (ii): $c(F^{V'}) \leq |V'|$, where $c(F^{V'}) = c(F^V)$, and thus $\delta(F^V) = c(F^V) - n(F^V) = c(F^{V'}) - |V'| \leq 0$. \square

We obtain, that decreasing multiplicities in $F \in \underline{\mathcal{SED}}$ does not leave this class (even if the multiplicity drops to zero):

Corollary 9.4 *For $F \in \underline{\mathcal{SED}}$ and $F' \leq F$ we have $F' \in \underline{\mathcal{SED}}$.*

Proof: For all $\emptyset \subseteq V \subseteq \text{var}(F')$ by Lemma 9.3 we have $c(F'^V) \leq c(F^V) \leq |V|$, and thus $F' \in \underline{\mathcal{SED}}$. \square

Even stronger, we can characterise the elements of $\underline{\mathcal{SED}}$ as clause-sets close to matching-satisfiable clause-sets:

Theorem 9.5 *For $F \in \underline{\mathcal{CLS}}$ holds $F \in \underline{\mathcal{SED}}$ if and only if for all $v \in \text{var}(F)$ holds $F^{\text{var}(F) \setminus \{v\}} \in \underline{\mathcal{MSAT}}$.*

Proof: Recall that by [17, Lemma 7.2] we have $G \in \underline{\mathcal{MSAT}}$ iff $\delta^*(G) = 0$ for $G \in \underline{\mathcal{CLS}}$, where $\delta^*(G) = \max_{G' \leq G} \delta(G')$. If $F \in \underline{\mathcal{SED}}$, but $F^{\text{var}(F) \setminus \{v\}} \notin \underline{\mathcal{MSAT}}$ for some $v \in \text{var}(F)$, then there would be $G \leq F^{\text{var}(F) \setminus \{v\}}$ with $\delta(G) > 0$, violating Lemma 9.3 by choosing $V := \text{var}(G)$ (namely $\delta(F^V) \leq 0$, but $\delta(G) = c(G) - n(G) \leq c(F^V) - n(G) = \delta(F^V)$). And if $F \notin \underline{\mathcal{SED}}$, then there is $V \subseteq \text{var}(F)$ with $\delta(F^V) \geq 1$, where $F^V \leq F^{\text{var}(F) \setminus \{v\}}$ for $v \in \text{var}(F) \setminus V$. \square

We note that in Example 9.2 we have seen satisfiable as well as unsatisfiable elements of $\underline{\mathcal{SED}}$, and for the satisfiable elements we have matching-satisfiable as well as matching-unsatisfiable ones. Recalling Lemma 4.1, where we have characterised the elements of \mathcal{VMU} as those $F \in \underline{\mathcal{USAT}}$, such that for all $v \in \text{var}(F)$ holds $F^{\text{var}(F) \setminus \{v\}} \in \underline{\mathcal{SAT}}$, we see that the unsatisfiable elements of $\underline{\mathcal{SED}}$ are variable-minimally unsatisfiable:

Corollary 9.6 $\underline{\mathcal{SED}} \cap \underline{\mathcal{USAT}} \subset \mathcal{VMU}$ (and thus $\underline{\mathcal{SED}} \cap \underline{\mathcal{USAT}} \subset \underline{\mathcal{VMU}}$).

Proof: $\underline{\mathcal{SED}} \cap \underline{\mathcal{USAT}} \subset \mathcal{VMU}$ follows from Theorem 9.5 by Lemma 4.1. An example of $F \in \mathcal{MU} \setminus \underline{\mathcal{SED}}$ is given in Example 9.2. And for $F \in \underline{\mathcal{SED}} \cap \underline{\mathcal{USAT}}$ and for the underlying clause-set F' holds $F' \in \underline{\mathcal{SED}} \cap \underline{\mathcal{USAT}}$, whence $F' \in \mathcal{VMU}$, and thus $F \in \underline{\mathcal{VMU}}$. \square

By definition follows that for $\emptyset \subset V \subseteq \text{var}(F)$ with minimal $\delta(F[V])$ we have $F[V] \in \underline{\mathcal{SED}}$, which is a central insight for generalising the upper bound in the subsequent Subsection 9.2:

Corollary 9.7 *Consider $F \in \underline{\mathcal{CLS}}$, $n(F) > 0$, and $\emptyset \subset V \subseteq \text{var}(F)$ with $\delta(F[V]) = \sigma(F)$. Then we have:*

1. $F[V] \in \underline{\mathcal{SED}}$.
2. If $F \in \mathcal{LEAN}$, then $F[V] \in \mathcal{VMU}$.

Proof: Part 1 holds since $F[V]$ has maximal deficiency amongst all $F[V']$ for $V' \neq \emptyset$, and for $V' \subseteq V$ holds $F[V][V'] = F[V']$. Part 2 follows by Part 1, Corollary 9.6, and the unsatisfiability of $F[V]$ (recall that a satisfying assignment for $F[V]$, when restricted to V , is an autarky for F). \square

In Part 2 of Corollary 9.7 we use \mathcal{LEAN} , \mathcal{VMU} instead of $\underline{\mathcal{LEAN}}$, $\underline{\mathcal{VMU}}$, since these classes are invariant under multiplicities (different from \mathcal{SED}).

We conclude this subsection by considering the complexity of SAT decision for $F \in \mathcal{SED}_{\delta=k}$ for fixed parameter $k \in \mathbb{N}$. By Corollary 9.6 we could use Theorem 4.7, however we can do better, namely we have $\mathcal{SED}_{\delta=k} \subset \mathcal{MLEAN}$ (due to $\sigma(F) = k \geq 1$ for $F \in \mathcal{SED}_{\delta=k}$), and so we can apply the fpt-result discussed in Example 2.18, obtaining that SAT decision for inputs in $\mathcal{SED}_{\delta=k}$ is fpt in k . The question is whether we actually have polytime SAT decision:

Question 9.8 *Can SAT decision for \mathcal{SED} be done in polynomial time? If so, can we also find a satisfying assignment quickly?*

In Subsection 10.2 we will see a (potential) application of Question 9.8.

9.2. The generalised upper bound

Back to the main task, we first show that from $F[V] \in \mathcal{VMU}$ we obtain variables of low degree for F itself:

Lemma 9.9 *Consider $F \in \underline{\mathcal{CLS}}$ and $\emptyset \subset V \subseteq \text{var}(F)$ such that $F[V] \in \mathcal{VMU}$. Then there exists $v \in V$ with $\text{vd}_F(v) \leq \text{nM}(\delta(F[V]))$ and $\text{ld}_F(v), \text{ld}_F(\bar{v}) \leq \delta(F[V])$.*

Proof: Let $F' := F[V]$ and consider some minimally unsatisfiable $F'' \subseteq F'$; by assumption we have $\text{var}(F'') = \text{var}(F')$. So we get $\delta(F'') = \delta(F') - (c(F') - c(F''))$. By Theorem 8.3 there is $v \in \text{var}(F'')$ with

$$\text{vd}_{F''}(v) \leq \text{nM}(\delta(F'')) = \text{nM}(\delta(F') - (c(F') - c(F''))) \leq \text{nM}(\delta(F')) - (c(F') - c(F''))$$

and $\text{ld}_{F''}(v), \text{ld}_{F''}(\bar{v}) \leq \delta(F'') = \delta(F') - (c(F') - c(F''))$. Finally we have $\text{vd}_F(v) \leq \text{vd}_{F''}(v) + (c(F') - c(F''))$ (note that all occurrences of v in F are also in F'), and similarly for the literal degrees. \square

We obtain the generalisation and strengthening of Theorem 8.3:

Theorem 9.10 *We have $\mu\text{vd}(F) \leq \text{nM}(\sigma(F))$ for a lean multi-clause-set F with $n(F) > 0$. More precisely, there exists a variable $v \in \text{var}(F)$ with $\text{vd}_F(v) \leq \text{nM}(\sigma(F))$ and $\text{ld}_F(v), \text{ld}_F(\bar{v}) \leq \sigma(F)$.*

Proof: By Corollary 9.7, Part 2, and Lemma 9.9. \square

Since for a variable $v \in \text{var}(F)$ for any $F \in \mathcal{CLS}$ holds

$$\delta(F[\{v\}]) = \text{vd}_F(v) - 1 \geq \mu\text{vd}(F) - 1 \geq \sigma(F),$$

and the surplus is a lower bound for the deficiency, we get:

Corollary 9.11 For a lean multi-clause-set F , $n(F) > 0$, we have

$$\begin{aligned}\sigma(F) + 1 &\leq \mu\text{vd}(F) \leq nM(\sigma(F)) \leq \sigma(F) + 1 + \text{fld}(\sigma(F)) \\ \mu\text{vd}(F) &\leq nM(\delta(F)) \leq \delta(F) + 1 + \text{fld}(\delta(F)).\end{aligned}$$

That the bounds from Corollary 9.11 are sharp in general, is shown by the following examples.

Example 9.12 First we consider any lean clause-set $F \neq \top$, and perform a non-strict full subsumption extension $F \rightsquigarrow F'$. Obviously F' is lean as well (with $\delta(F') = \delta(F)$). We have $\mu\text{vd}(F') = 2$ and $\sigma(F') = 1$, and thus

$$2 = \sigma(F') + 1 = \mu\text{vd}(F') = nM(\sigma(F')) = \sigma(F) + 1 + \text{fld}(\sigma(F)),$$

while $\delta(F')$ is unbounded. This construction will be taken up again in Lemma 10.13

Now we turn to the δ -upper bounds. For $n \geq 2$ consider A_n . We have $\sigma(A_n) = \delta(A_n) = 2^n - n$ by Example 2.20. Thus here the inequalities of Corollary 9.11 are

$$\begin{aligned}2^n - n + 1 = \sigma(A_n) + 1 &< \\ 2^n = \mu\text{vd}(A_n) = nM(\delta(A_n)) &= \delta(A_n) + 1 + \text{fld}(\delta(A_n))\end{aligned}$$

(using Corollary 7.24).

9.3. Sharpness of the bound for \mathcal{VMU}

We now show that for every deficiency k there are variable-minimally unsatisfiable clause-sets where the min-var degree is $nM(k)$ (strengthening Example 9.12). The examples are obtained as follows:

Lemma 9.13 For a clause-set $F \in \mathcal{CLS}$, $\perp \notin F$ and $n(F) > 0$, with at least one full clause, consider the following construction of $F' \in \mathcal{CLS}$:

1. Let C be a full clause of F .
2. Let F'' be a full singular unit-extension of F (recall Definition 5.15).
3. Let $F' := F'' \cup \{C\}$.

We have the following properties:

1. $\sigma(F') = \sigma(F) + 1$, $\delta(F') = \delta(F) + 1$, $\mu\text{vd}(F') = \mu\text{vd}(F) + 1$.
2. $F \in \mathcal{SED} \Rightarrow F' \in \mathcal{SED}$.
3. $F \in \mathcal{USAT} \Rightarrow F' \in \mathcal{USAT}$.

Proof: With Lemma 5.17 we get $\sigma(F'') = \sigma(F)$, $\delta(F'') = \delta(F)$, $\mu\text{vd}(F'') = \mu\text{vd}(F)$. Obviously $\delta(F') = \delta(F'') + 1$. Let $\text{var}(F'') \setminus \text{var}(F) = \{v\}$. To see $\mu\text{vd}(F') = \mu\text{vd}(F'') + 1$, we note that for $w \in \text{var}(F)$ we have $\text{vd}_{F'}(w) = \text{vd}_{F''}(w) + 1$, while $\text{vd}_{F'}(v) = c(F') - 1 = c(F'') \geq \text{vd}_{F''}(w) + 1$.

To prove $\sigma(F') = \sigma(F'') + 1$, we consider $\emptyset \subset V \subseteq \text{var}(F') = \text{var}(F'')$. If $v \notin V$, then $\delta(F'[V]) = \delta(F''[V]) + 1$, since C is full for F . If $V = \{v\}$, then $\delta(F'[V]) = \delta(F''[V]) = c(F'') \geq \sigma(F'') + 1$. Finally, if $V \supset \{v\}$, then $\delta(F'[V]) = c(F') - |V| \geq \delta(F') = \delta(F'') + 1 \geq \sigma(F'') + 1$.

The implication $F \in \mathcal{SED} \Rightarrow F' \in \mathcal{SED}$ follows now by definition of \mathcal{SED} , and $F \in \mathcal{USAT} \Rightarrow F' \in \mathcal{USAT}$ is trivial. \square

With the construction of Lemma 9.13 we now show that the general upper bound on the min-var degree of lean clause-sets is tight for variable-minimally unsatisfiable clause-sets:

Theorem 9.14 For a class $\mathcal{VMU} \cap \mathcal{SED} \subseteq \mathcal{C} \subseteq \mathcal{LEAN}$ and $k \in \mathbb{N}$ we have $\mu\text{vd}(\mathcal{C}_{\delta=k}) = \text{nM}(k)$.

Proof: By Theorem 9.10 it remains to show the lower bound $\mu\text{vd}(\mathcal{VMU}_{\delta=k} \cap \mathcal{SED}) \geq \text{nM}(k)$. For deficiencies $k = 2^n - n$, $n \in \mathbb{N}$ we have $\text{nM}(k) = 2^n$, and thus A_n serves as lower bound example (as shown in Example 9.12), while until the next jump position we can use Lemma 9.13 together with Corollary 9.6, where due to Corollary 7.24 in this range also nM increases only by 1 for $k \rightsquigarrow k + 1$. \square

Using Lemma 4.3, we can now determine the min-var degrees for the classes \mathcal{LEAN} and \mathcal{VMU} , separated into layers via deficiency or surplus:

Corollary 9.15 For $k \in \mathbb{N}$ holds $\text{nM}(k) = \mu\text{vd}(\mathcal{LEAN}_{\delta=k}) = \mu\text{vd}(\mathcal{LEAN}_{\sigma=k}) = \mu\text{vd}(\mathcal{VMU}_{\delta=k}) = \mu\text{vd}(\mathcal{VMU}_{\sigma=k})$.

10. Algorithmic implications

In Subsections 10.1, 10.2 we consider the algorithmic implications of Theorem 9.10 (the upper bound on the minimum var-degree for lean clause-sets). First in Theorem 10.2 we show that via a poly-time autarky-reduction every clause-set $F \in \mathcal{CLS}$ can be reduced to some $F' \subseteq F$, where F' fulfils the min-var-degree upper bound of Theorem 9.10, although F' might not be lean. For this autarky-reduction we do not know whether we can efficiently compute a certificate, the autarky, and we discuss Conjecture 10.3, that efficient computation is possible, in Subsection 10.2. We conclude with some remarks on the surplus in Subsection 10.3.

10.1. Autarky reduction

By Theorem 9.10 lean clause-sets fulfil a condition on the minimum variable-degree — if that condition is not fulfilled, then there exists an autarky. In this section we try to pinpoint these autarkies. We consider a vast generalisation of lean clause-sets, namely matching-lean clause-sets (recall Subsection 2.6, especially that a multi-clause-set F with $n(F) > 0$ is matching-lean iff $\sigma(F) \geq 1$). It is also useful to note here the observation that a (multi-)clause-set F has a non-trivial autarky (is not lean) iff there is $\emptyset \subset V \subseteq \text{var}(F)$ such that $F[V]$ is satisfiable, and the corresponding autarky reduction of F removes all clauses containing some variable of V ; note that to perform this autarky reduction the autarky itself (the satisfying assignment for $F[V]$) is not needed, only its set V of variables.

To start, we obtain a sufficient criterion for the existence of a non-trivial autarky by considering the converse of Theorem 9.10:

Lemma 10.1 Consider a matching-lean multi-clause-set F with $n(F) > 0$. If we have $\mu\text{vd}(F) > \text{nM}(\sigma(F))$, then for all $F' := F[V]$ with $\emptyset \subset V \subseteq \text{var}(F)$ and $\delta(F[V]) = \sigma(F)$ we have:

1. $\delta(F') = \sigma(F') = \sigma(F)$, and so $F' \in \underline{\mathcal{SED}} \cap \underline{\mathcal{MLLEAN}}$.
2. $\mu\text{vd}(F') > \text{nM}(\sigma(F'))$.
3. $F' \in \mathcal{SAT}$ (thus there is $\varphi \in \mathcal{PASS}$ with $\text{var}(\varphi) = V$ and $\varphi * F' = \top$; this φ is a non-trivial autarky for F).

Proof: Part 1 follows by Corollary 9.7, Part 1. For Part 2 note that $\mu\text{vd}(F') \leq \text{nM}(\sigma(F'))$ implies $\mu\text{vd}(F) \leq \mu\text{vd}(F') \leq \text{nM}(\sigma(F))$ contradicting the assumption. Finally for Part 3 assume $F' \in \mathcal{USAT}$. Then by Part 1 and Corollary 9.6 we obtain $F' \in \mathcal{VMU}$, which contradicts Part 2 by Lemma 9.9. \square

To better understand the background, we recall two fundamental facts regarding the surplus $\sigma(F)$ for multi-clause-set F with $n(F) > 0$:

1. $\sigma(F)$ together with some $\emptyset \subset V \subseteq \text{var}(F)$ with $\sigma(F) = \delta(F[V])$ can be computed in polynomial time (see [16, Subsection 11.1]).
2. If $\sigma(F) \leq 0$, then one can compute a non-trivial matching autarky for F in polynomial time (see [13, Section 7] or [16, Section 9]).

We see now that we can reach the conclusion of Theorem 9.10 for arbitrary inputs F in polynomial time, via some autarky reduction (maintaining satisfiability-equivalence):

Theorem 10.2 *Consider a multi-clause-set F . We can find in polynomial time a sub-clause-set $F' \subseteq F$ such that:*

1. *There exists an autarky φ for F with $F' = \varphi * F$.*
2. *If $n(F') > 0$, then $\sigma(F') \geq 1$ and $\mu\text{vd}(F') \leq nM(\sigma(F'))$.*

Proof: First F is reduced to the underlying clause-set (and all further computations only handle clause-sets), and if $\perp \in F$, then we reduce F to $\{\perp\}$ and are finished. Otherwise, the reduction process of F yielding the final F' consists of a loop with two steps:

1. First eliminate matching-autarkies (i.e., compute the matching-lean kernel), such that we reach $\sigma(F) \geq 1$.
2. Then apply the autarky-reduction according to Part 3 of Lemma 10.1, removing all clauses containing a variable of V , in case of $\mu\text{vd}(F) > nM(\sigma(F))$.

This loop is aborted if \top is reached or the criterion no longer applies. All autarkies are composed together (as shown in [10] for the general case, the composition of autarkies is again an autarky; here the autarkies are variable-disjoint, and thus we can just take their union), yielding the final φ . \square

In Theorem 10.2 we can only show the *existence* of an autarky φ for F with $F' = \varphi * F$, however we currently do not know how to compute it efficiently. We conjecture that it can be found in polynomial time:

Conjecture 10.3 *For $F \in \mathcal{CLS}_{\sigma \geq 1}$ there is a poly-time algorithm for computing a non-trivial autarky φ for F in case of $\mu\text{vd}(F) > nM(\sigma(F))$.*

Note that we ask only to find *some* autarky φ , not necessarily one given by Lemma 10.1 (i.e., with $\text{var}(\varphi) = V$ as in Part 3 of Lemma 10.1). That this is enough follows by the fact that the number of variables is reduced by such a reduction, and this by some autarky:

Lemma 10.4 *If Conjecture 10.3 is true, then for the algorithm from Theorem 10.2, which reduces a multi-clause-set F to some (satisfiability-equivalent) $F' \subseteq F$, we can also compute an autarky φ for F with $F' = \varphi * F$ in polynomial time.*

Proof: In the loop as given in the proof of Theorem 10.2, we can replace the autarky-reduction according to Part 3 of Lemma 10.1 by the reduction $F \rightsquigarrow \varphi * F$ according to a (non-trivial) autarky as given by Conjecture 10.3. \square

In the subsequent Subsection 10.2 we discuss what we know about Conjecture 10.3.

10.2. On finding the autarky

Consider a matching-lean multi-clause-set F with $n(F) > 0$, where Lemma 10.1 is applicable (recall that we have $\sigma(F) \geq 1$), that is, we have $\mu\text{vd}(F) > nM(\sigma(F))$. So we know that F has a non-trivial autarky. Conjecture 10.3 states that finding such a non-trivial autarky in this case can be done in polynomial time (recall that finding a non-trivial autarky in general is NP-complete, which was shown in [13]).

The task of actually finding the autarky can be considered as finding a satisfying assignment for the following class $\underline{\mathcal{MLCR}} \subset \underline{\mathcal{SAT}} \cap \underline{\mathcal{MLEAN}}$ of satisfiable(!) multi-clause-sets F , obtained by considering all $F[V]$ for minimal sets of variables V with $\delta(F[V]) = \sigma(F)$ (where ‘‘CR’’ stands for ‘‘critical’’):

Definition 10.5 *Let \mathcal{MLCR} be the class of clause-sets F fulfilling the following three conditions:*

1. $F \in \mathcal{MLEAN}$, $\perp \notin F$, $F \neq \top$.
2. For all $\emptyset \subset V \subset \text{var}(F)$ holds $\delta(F[V]) > \sigma(F)$.
3. $\mu\text{vd}(F) > nM(\sigma(F))$.

(The definition of $\underline{\mathcal{MLCR}}$ just uses $F \in \underline{\mathcal{MLEAN}}$ instead.)

The basic properties of this class are collected in the following lemma:

Lemma 10.6 *For $F \in \underline{\mathcal{MLCR}}$ holds:*

1. $\delta(F) = \sigma(F) \geq 1$ (whence $F \in \underline{\mathcal{SED}}$).
2. $F \in \underline{\mathcal{SAT}}$.

Proof: Since $F \in \underline{\mathcal{MLEAN}}$ and $n(F) > 0$, we have $\sigma(F) \geq 1$. By $\perp \notin F$ we get $F = F[\text{var}(F)]$, and thus $\sigma(F) = \delta(F[\text{var}(F)]) = \delta(F)$, while $F \in \underline{\mathcal{SAT}}$ follows by Lemma 10.1. \square

The examples we know for elements of \mathcal{MLCR} are as follows:

Example 10.7 *A simple example for $F \in \mathcal{MLCR}_{\delta=1} \cap \mathcal{HIT}$ is given by*

$$F := \{\{1, 2\}, \{-1, 2, -3\}, \{-2, 3\}, \{1, -2, -3\}\}.$$

We have $\delta(F) = 4 - 3 = 1$ and $\mu\text{vd}(F) = 3$; for $\sigma(F) = 1$ and Condition 2 of Definition 10.5 notice, that any two variables cover all four clauses, and thus the minimum of $\delta(F[V])$ is only attained for $V = \text{var}(F)$; finally by $\mu\text{vd}(F) = 3 > nM(1) = 2$ we get $F \in \mathcal{MLCR}$, while $F \in \mathcal{HIT}$ by definition (any two clauses have a clash).

This example also shows that $\underline{\mathcal{MLCR}}$ is not invariant under multiplicities: Obtain F' from F by duplicating the first clause. We still have $F' \in \underline{\mathcal{MLEAN}}$, but $\delta(F') = 2$, while $\delta(F'[\{3\}]) = 2$ as well, and thus $F' \notin \underline{\mathcal{MLCR}}$. So duplicating a clause can lead outside of $\underline{\mathcal{MLCR}}$. In the other direction, removing a duplication, we have the following simple (counter-)example: Let $F := \{2 * \{1\}\}$; trivially we have $F \in \underline{\mathcal{MLCR}}$, but for the underlying clause-set we have $\{\{1\}\} \notin \underline{\mathcal{MLEAN}}$, thus it is not in \mathcal{MLCR} .

A class of example is obtained by full clause-sets. Let F be a full clause-set and $n := n(F)$, $m := c(F)$. Then $F \in \mathcal{MLCR}$ iff $n < m < 2^n$:

1. We have $\delta(F) = m - n$ and thus $\delta(F) \geq 1 \Leftrightarrow m > n$.

2. Furthermore $F \in \mathcal{SAT} \Leftrightarrow m < 2^n$.
3. For $\emptyset \subset V \subset \text{var}(F)$ we have $\delta(F[V]) = m - |V|$. Thus $\sigma(F) = \delta(F)$, and Condition 2 of Definition 10.5 is fulfilled.
4. It remains to show the condition on the min-var degree: We have $\mu\text{vd}(F) = m$, while $\text{nM}(\sigma(F)) = \text{nM}(m - n)$. By Theorem 7.21 we obtain $\text{nM}(m - n) = m - n + \text{fld}(m - n + 1 + \text{fld}(m - n + 1))$. We obtain for $n \geq 1$:

$$\begin{aligned} \mu\text{vd}(F) > \text{nM}(\sigma(F)) &\Leftrightarrow m > m - n + \text{fld}(m - n + 1 + \text{fld}(m - n + 1)) \Leftrightarrow \\ &\text{fld}(m - n + 1 + \text{fld}(m - n + 1)) < n \Leftarrow \\ &\text{fld}(2^n - 1 - n + 1 + \text{fld}(2^n - 1 - n + 1)) < n \Leftrightarrow \\ &\text{fld}(2^n - n + \text{fld}(2^n - n)) = \text{fld}(2^n - n + (n - 1)) = \text{fld}(2^n - 1) < n. \end{aligned}$$

Finally we note that the class $\underline{\mathcal{MLCCR}}$ is invariant against changes of polarities of literal occurrences (if clauses become equal in this way, then their multiplicities have to be added), and thus for example replacing all clauses $C \in F \in \underline{\mathcal{MLCCR}}$ by their positive forms, $\text{var}(C)$, we obtain a positive (no complementations occur) multi-clause-set $F' \in \underline{\mathcal{MLCCR}}$ (with $c(F') = c(F)$ and $n(F') = n(F)$).

The importance of \mathcal{MLCCR} is, that it is sufficient to find a non-trivial autarky for this class of satisfiable clause-sets. In order to show this, we need to strengthen the polytime computation of $\sigma(F)$:

Lemma 10.8 For a multi-clause-set F with $n(F) > 0$ we can compute in polynomial time a minimal subset $\emptyset \subset V \subseteq \text{var}(F)$ with $\delta(F[V]) = \sigma(F)$.

Proof: Let $V := \text{var}(F)$. Check whether there is $v \in \text{var}(F)$ with $\sigma(F[V \setminus \{v\}]) = \sigma(F)$ — if yes, then $V := V \setminus \{v\}$ and repeat, if not, then V is the desired result. \square

We are ready to show that \mathcal{MLCCR} is really the “critical class” for the problem of finding the witness-autarky underlying the reduction $F \rightsquigarrow F'$ of Theorem 10.2:

Theorem 10.9 Consider $F \in \mathcal{CLS}$ with $\sigma(F) \geq 1$ and $\mu\text{vd}(F) > \text{nM}(\sigma(F))$.

1. For every minimal subset $\emptyset \subset V \subseteq \text{var}(F)$ with $\delta(F[V]) = \sigma(F)$ we have $F[V] \in \underline{\mathcal{MLCCR}}$.
2. We can compute in polytime some $\emptyset \subset V \subseteq \text{var}(F)$ with $F[V] \in \underline{\mathcal{MLCCR}}$.
3. Conjecture 10.3 is equivalent to the statement, that finding a non-trivial autarky for clause-sets in $\underline{\mathcal{MLCCR}}$ can be achieved in polynomial time.

Proof: Part 1 follows with Lemma 10.1. Part 2 follows from Part 1 with Lemma 10.8. Part 3 follows with Part 2 (note that every autarky for some $F[V]$ yields an autarky for F). \square

Since $\mathcal{MLCCR} \subset \mathcal{SED}$, if both questions of Question 9.8 have a yes-answer, then this would prove Conjecture 10.3.

10.3. Final remarks on the surplus

It is instructive to investigate the precise relationship between minimum variable-degree and surplus for lean clause-sets, which by Corollary 9.11 are indeed very close. Small values behave as follows:

Lemma 10.10 Consider $F \in \mathcal{LEAN}$, $n(F) > 0$ (so $\sigma(F) \geq 1$ and $\mu\text{vd}(F) \geq 2$).

1. $\sigma(F) = 1$ holds if and only if $\mu\text{vd}(F) = 2$ holds.
2. $\mu\text{vd}(F) = 3$ implies $\sigma(F) = 2$.
3. $\sigma(F) = 2$ implies $\mu\text{vd}(F) \in \{3, 4\}$.
4. $\mu\text{vd}(F) = 4$ implies $\sigma(F) \in \{2, 3\}$.

Proof: First consider Part 1. If $\sigma(F) = 1$ (so $n(F) > 0$), then by Theorem 9.10 we have $\mu\text{vd}(F) \leq \text{nM}(1) = 2$, while in case of $\mu\text{vd}(F) = 1$ there would be a matching autarky for F . If on the other hand $\mu\text{vd}(F) = 2$ holds, then by definition $\sigma(F) \leq 2 - 1 = 1$, while $\sigma(F) \geq 1$ holds since F is matching lean. For Part 2 note that due to $\sigma(F) + 1 \leq \mu\text{vd}(F)$ we have $\sigma(F) \leq 2$, and then the assertion follows by Part 1; Part 4 follows in the same way. Finally Part 3 follows by Part 1 and $\text{nM}(2) = 4$. \square

So for $F \in \mathcal{MU}' \setminus \{\{\perp\}\}$ we have $\sigma(F) \geq 2$; for a general polytime reduction of arbitrary (also non-boolean) clause-sets achieving surplus at least 2 see [16, Lemma 11.9]. We illustrate Lemma 10.10, using F with $\delta(F) = \sigma(F)$:

Example 10.11 *Examples for cases $\sigma(F) \in \{2, 3\}$ in Lemma 10.10:*

1. An example for $\mu\text{vd}(F) = 4$ in Part 3 with $F \in \mathcal{UHIT} \cap \mathcal{SED}$ is given by A_2 .
2. For $\{\{a, b, c\}, \{\bar{a}, b, c\}, \{a, \bar{b}, c\}, \{\bar{a}, \bar{b}, c\}, \{a, \bar{c}\}, \{\bar{a}, \bar{c}\}\} \in \mathcal{UHIT} \cap \mathcal{SED}$ we have $\mu\text{vd}(F) = 4$ and $\sigma(F) = 3$ (Part 4).

Question 10.12 *is there for every $k \in \mathbb{N}$ an $F \in \mathcal{UHIT} \cap \mathcal{SED}$ with $\sigma(F) = k$ and $\mu\text{vd}(F) = k + 1$?*

As we have for \mathcal{MU} the levels $\mathcal{MU}_{\delta=k}$ for $k = 1, 2, \dots$, we can consider for \mathcal{LEAN} the levels $\mathcal{LEAN}_{\sigma=k}$ for $k = 1, 2, \dots$. However, while the levels $\mathcal{MU}_{\delta=k}$ as well as $\mathcal{LEAN}_{\delta=k}$ all are decidable in polynomial time, already the first level $\mathcal{LEAN}_{\sigma=1}$ is NP-complete:

Lemma 10.13 *Consider the map $E : \mathcal{CLS} \rightarrow \mathcal{CLS}$, which has $E(\top) := \top$, while otherwise for $F \in \mathcal{CLS} \setminus \{\top\}$ it chooses (by some rule — it doesn't matter) a clause $C \in F$ and a variable $v \in \mathcal{VA} \setminus \text{var}(F)$, and replaces C by $C \cup \{v\}, C \cup \{\bar{v}\}$; in other words, a non-strict full subsumption extension $F \rightsquigarrow E(F)$ is performed, as in Example 9.12. Then we have for $F \in \mathcal{CLS}$:*

1. $F \in \mathcal{LEAN}$ iff $E(F) \in \mathcal{LEAN}$.
2. $F \in \mathcal{MU}$ iff $E(F) \in \mathcal{MU}$.
3. $\sigma(F) \leq 1$.

Thus $\mathcal{LEAN}_{\sigma=1}$ is coNP-complete, while $\mathcal{MU}_{\sigma=1}$ is D^P -complete.

Proof: The properties of the map E are trivial. The completeness-properties follow with the coNP-completeness of \mathcal{LEAN} ([13]) and the D^P -completeness of \mathcal{MU} ([21]). \square

With Lemma 10.13 we also get easy examples for minimally unsatisfiable clause-sets of surplus = 1 and arbitrary deficiency ≥ 1 .

11. Matching lean clause-sets

In this section, which concludes our considerations on generalisations (beyond \mathcal{MU}), we consider the question whether Theorem 9.10 can incorporate non-lean clause-sets. We consider the large class \mathcal{MLEAN} of matching lean clause-sets, which is natural, since a basic property of $F \in \mathcal{MU}$ used in the proof of Theorem 9.10 is $\delta(F) \geq 1$ for $F \neq \top$, and this actually holds for all $F \in \mathcal{MLEAN}$. We will construct for arbitrary deficiency $k \in \mathbb{N}$ and $K \in \mathbb{N}$ clause-sets $F \in \mathcal{MLEAN}$ of deficiency k , where every variable occurs positively at least K times. Thus neither the upper bound $\max(\text{ld}_F(v), \text{ld}_F(\bar{v})) \leq f(\delta(F))$ nor $\text{ld}_F(v) + \text{ld}_F(\bar{v}) = \text{vd}_F(v) \leq f(\delta(F))$ for some chosen variable v and for any function f does hold for \mathcal{MLEAN} .

An example for $F_{\text{sf}} \in \mathcal{HIT}_{\delta=1}$, $n(F) > 0$, with $\mu\text{ld}(F) := \min_{x \in \text{lit}(F)} \text{ld}_F(x) \geq 2$ (the minimal literal degree; and thus $\mu\text{vd}(F) \geq 4$) is given in [73, Section 5], a “star-free” clause-set F_{sf} , as discussed in Subsection 1.6.5 (furthermore between any two different clauses of F_{sf} there is exactly one conflict). In [16, Subsection 9.3] it is shown that $F_{\text{sf}} \in \mathcal{MLEAN}$ holds. “Star-freeness” in our context means, that there are no singular variables (occurring in one sign only once). The simpler construction of this section pushes the number of positive occurrences arbitrary high, but there are variables with only one negative occurrence (i.e., there are singular variables).

For a finite set V of variables let $M(V) \subseteq A(V)$ be the full clause-set over V containing all full clauses with at most one complementation; e.g. $M(\{1, 2\}) = \{\{1, 2\}, \{-1, 2\}, \{1, -2\}\}$:

1. Obviously $n(M(V)) = |V|$, $c(M(V)) = |V| + 1$ and $\delta(M(V)) = 1$ holds.
2. We have already seen that $M(V) \in \mathcal{MLEAN}$ (indeed $M(V) \in \mathcal{MLCR}$, as shown in Example 10.7).
3. By definition we have $\text{ld}_{M(V)}(v) = |V|$ and $\text{ld}_{M(V)}(\bar{v}) = 1$ for all $v \in V$.

Lemma 11.1 *For $k \in \mathbb{N}$ and $K \in \mathbb{N}$ there is $F \in \mathcal{MLEAN}_{\delta=k}$ such that for all variables $v \in \text{var}(F)$ we have $\text{ld}_F(v) \geq K$, and with $F \in \mathcal{USAT}$ for $k \geq 2$.*

Proof: For $k = 1$ we can set $F := M(\{v_1, \dots, v_K\})$; so assume $k \geq 2$. Consider any clause-set $G \in \mathcal{MU}_{\delta=k-1}$ with $n := n(G) \geq K$, and let $V := \text{var}(G)$. Consider a disjoint copy of V , that is a set V' of variables with $V' \cap V = \emptyset$ and $|V'| = |V|$, and consider two enumerations of the clauses $M(V) = \{C_1, \dots, C_{n+1}\}$, $M(V') = \{C'_1, \dots, C'_{n+1}\}$. Now

$$F := G \cup \{C_i \cup C'_i : i \in \{1, \dots, n+1\}\} \in \mathcal{USAT}$$

(with $\text{var}(F) = V \cup V'$) has no matching autarky: If φ is a matching autarky for F , then $\text{var}(\varphi) \cap V = \emptyset$ since G is matching lean, thus $\text{var}(\varphi) \cap V' = \emptyset$ since $M(V')$ is matching lean, and thus φ must be trivial. Furthermore we have $n(F) = 2n$ and $c(F) = c(G) + n + 1$, and thus $\delta(F) = c(G) + n + 1 - 2n = \delta(G) + 1 = k$. By definition for all variables $v \in \text{var}(F)$ we have $\text{ld}_F(v) \geq n$. \square

For $k = 1$ the examples of Lemma 11.1 for $K \geq 3$ are necessarily satisfiable, since $\mathcal{MLEAN}_{\delta=1} \cap \mathcal{USAT} = \mathcal{MU}_{\delta=1}$. It remains the questions whether the singular variables can be eliminated:

Question 11.2 *Are there examples for deficiency $k \in \mathbb{N}$ of $F \in \mathcal{MLEAN}_{\delta=k}$, $n(F) > 0$, with $\mu\text{ld}(F) \geq k + 1$?*

1. The above mentioned star-free F_{sf} shows that this is the case for $k = 1$.
2. By Theorem 9.10 for such examples we have $F \notin \mathcal{LEAN}$.
3. What about the stronger condition $\mu\text{ld}(F) \geq K$ for arbitrary $K \in \mathbb{N}$?

12. Lower bounds for VDM

We now return to minimally unsatisfiable clause-sets. By Theorem 8.3 we have $\text{VDM}(k) = \mu\text{vd}(\mathcal{MU}_{\delta=k}) \leq \text{nM}(k)$ for all $k \in \mathbb{N}$. The task of precisely determining $\text{VDM}(k)$ for all k seems a deep question, and is the subject of the remainder of this report. All our examples yielding lower bounds on $\text{VDM}(k)$ are actually (unsatisfiable) hitting clause-sets, and thus we believe (recall Definition 3.4)

Conjecture 12.1 *For all $k \in \mathbb{N}$ holds $\text{VDM}(k) = \text{VDH}(k)$.*

We will see in Theorem 14.4 that $\text{VDM} \neq \text{nM}$. We believe that VDM is a highly complicated function, but the true values deviate only at most by one from nM :

Conjecture 12.2 *For all $k \in \mathbb{N}$ we have $\text{VDM}(k) \geq \text{nM}(k) - 1$.*

By Corollary 7.23 we get:

Lemma 12.3 *If Conjecture 12.2 holds, then $k - 1 + \text{fld}(k + 1) \leq \text{VDM}(k) \leq k + 1 + \text{fld}(k)$ holds for $k \in \mathbb{N}$.*

Later in Lemma 13.1 we will see that $\text{VDM} : \mathbb{N} \rightarrow \mathbb{N}$ is monotonically increasing. In Theorem 14.6 we will (implicitly) construct a correction function $\gamma_1 : \mathbb{N} \rightarrow \{0, 1\}$ such that $\text{VDM} \leq \text{nM} - \gamma_1$, where we remark in the Conclusion (Section 15) that also $\text{VDM} \neq \text{nM} - \gamma_1$ holds. Note that Conjecture 12.2 says that there exists $\gamma : \mathbb{N} \rightarrow \{0, 1\}$ with $\text{VDM} = \text{nM} - \gamma$, while for every $\gamma : \mathbb{N} \rightarrow \{0, 1\}$ the function $\text{nM} - \gamma$ is still monotonically increasing (by Lemma 7.5), and is thus a possible candidate.

In Subsection 12.1 we provide a general method for obtaining *lower bounds*, via considering full clauses (while in Section 13 we turn to improved upper bounds):

Definition 12.4 *For a clause-set $F \in \mathcal{CLS}$ let $\text{fc}(F) \in \mathbb{N}_0$ be the number of full clauses, that is $\text{fc}(F) := |\{C \in F : \text{var}(C) = \text{var}(F)\}|$. And for a class $\mathcal{C} \subseteq \mathcal{CLS}$ of clause-sets we define $\text{fc}(\mathcal{C}) := \{\text{fc}(F) : F \in \mathcal{C}\} \subseteq \mathbb{N}_0$ as the set of all possible numbers of full clauses, while $\nu\text{fc}(\mathcal{C}) \in \mathbb{N}_0 \cup \{+\infty\}$ is the supremum of $\text{fc}(\mathcal{C})$.*

The maximum possible number of full clauses is an interesting quantity:

Definition 12.5 *For $k \in \mathbb{N}$ let*

$$\begin{aligned} \mathbf{FCM}(k) &:= \nu\text{fc}(\mathcal{MU}_{\delta=k}) \in \mathbb{N} \\ \mathbf{FCH}(k) &:= \nu\text{fc}(\mathcal{UHT}_{\delta=k}) \in \mathbb{N} \end{aligned}$$

(these numbers are finite due to $\text{FCH}(k) \leq \text{FCM}(k) \leq \text{VDM}(k)$).

According to our numerical investigations the number of full clauses is very close to VDM , and indeed to $\text{nM}(k)$:

Conjecture 12.6 *For all $k \in \mathbb{N}$ we have $\text{FCM}(k) \geq \text{nM}(k) - 1$.*

Conjecture 12.6 implies Conjecture 12.2; regarding $\text{FCH}(k)$, the maximal number of full clauses for unsatisfiable hitting clause-sets with deficiency k , we conjecture the difference $\text{FCM}(k) - \text{FCH}(k)$ is unbounded (see Subsection 15.3 for a discussion). The smallest deficiency k with $\text{FCM}(k) = \text{VDM}(k) - 1$ (and also $\text{FCH}(k) = \text{VDM}(k) - 1$) is $k = 3$, as shown in Lemma 12.18 (together with Theorem 14.4). We show for two infinite classes of deficiencies k that $\text{FCH}(k) = \text{VDH}(k) = \text{nM}(k)$ holds (Lemmas 12.11, 12.12). The main point here could be considered as (just) the equalities $\text{VDM}(k) = \text{nM}(k)$, for which in these two cases the proofs don't needed

to consider full clauses, and so the general method for computing lower bounds on $\text{FCM}(k)$, with the beginnings developed in Subsection 12.2, is not applied for these two lemmas. However in future work we will employ this method more fully (see Subsection 15.3), and, more important for the report at hand, we need for the proof of Theorem 14.4 (that $\text{VDM}(6) = \text{nM}(6) - 1 = 8$) the fact $\text{FCM}(3) \leq 4$, shown in Lemma 12.18.

12.1. Some precise values for VDM

A general lower-bound method for VDM is provided by the number $\text{fc}(F)$ of full clauses in a clause-set F . The supremum $\nu\text{fc}(\mathcal{MU}_{\delta=k}) = \text{FCM}(k)$ of this number over all elements of $\mathcal{MU}_{\delta=k}$ for fixed k is an interesting quantity in its own right, but in this report we only touch on this subject, providing the bare minimum of information needed in our context. See Subsection 15.3 for an outlook on the interesting properties of this quantity. Some simple examples:

Example 12.7 $\text{fc}(\top) = 0$, $\text{fc}(\{\perp\}) = 1$, and $\text{fc}(\{\{1\}, \{-1, 2\}\}) = 1$. While $\text{fc}(\emptyset) = \emptyset$, thus $\nu\text{fc}(\emptyset) = 0$, and $\text{fc}(\mathcal{CLS}) = \mathbb{N}_0$, thus $\nu\text{fc}(\mathcal{CLS}) = +\infty$.

By definition we have:

Lemma 12.8 $\text{fc}(F) \leq \mu\text{vd}(F)$ holds for every $F \in \mathcal{CLS}$ (since every variable in F has degree at least $\text{fc}(F)$), and thus $\nu\text{fc}(\mathcal{C}) \leq \mu\text{vd}(\mathcal{C})$ for every $\mathcal{C} \subseteq \mathcal{CLS}$.

We obtain that for lean clause-sets (especially minimally unsatisfiable clause-sets) of fixed deficiency the number of full clauses is bounded:

Corollary 12.9 A lean clause-set of deficiency k can have at most $\text{nM}(k)$ many full clauses; i.e., for all $k \in \mathbb{N}$ we have $\nu\text{fc}(\mathcal{LEAN}_{\delta=k}) \leq \text{nM}(k)$.

Question 12.10 Is the upper bound of Corollary 12.9 sharp?

Precise values for $\nu\text{fc}(\mathcal{MU}_{\delta=k}) = \text{FCM}(k)$ we show for two infinite classes of deficiencies. The simplest class are the deficiencies directly after the jumps (recall Lemma 7.20), the deficiencies of the A_n :

Lemma 12.11 For $n \in \mathbb{N}$ and $k := 2^n - n$ holds

$$\text{FCH}(k) = \text{FCM}(k) = \text{VDH}(k) = \text{VDM}(k) = \text{nM}(k) = 2^n.$$

Proof: We have $\nu\text{fc}(A_n) = 2^n$ (recall Lemma 2.13), and thus $\nu\text{fc}(\mathcal{UHT}_{\delta=k}) \geq 2^n$, while by Corollary 7.24 we have $\text{nM}(k) = 2^n$. \square

Also for the jumps themselves (recall Definition 7.12 and Lemma 7.20) the same conclusions hold, namely by Lemma 8.5, Part 2 (and the proof) we have:

Lemma 12.12 For all $k \in J$ holds

$$\text{FCH}(k) = \text{FCM}(k) = \text{VDH}(k) = \text{VDM}(k) = \text{nM}(k).$$

Note that for $k \in J$ there is $n \in \mathbb{N}$, $n \geq 2$, with $k = 2^n - n - 1$ and $\text{nM}(k) = 2^n - 2$. The underlying method of Lemmas 12.11, 12.12 is simple (as we already have explained in Subsection 1.3): start with A_n and apply strict full subsumption resolution to full clauses. Zero steps have been used in Lemma 12.11, one step in Lemma 12.12, and one example for two steps will be seen in the proof of Theorem 14.1 (clause-set F_3 there). The further development of this method we have to leave for future work:

Question 12.13 Explore the application of strict full subsumption resolution to A_n in order to obtain lower bounds for $\text{VDH}(k)$ (and also $\text{FCH}(k)$). (By Lemma 6.9 we have $A_n \xrightarrow{\text{sfsR}}_* F$ iff $F \in \mathcal{UHT}$ with $\text{var}(F) = \text{var}(A_n) = \{1, \dots, n\}$; we obtain the lower bounds $\text{VDH}(\delta(F)) \geq \mu\text{vd}(F)$ and $\text{FCH}(\delta(F)) \geq \text{fc}(F)$.)

12.2. On the number of full clauses

We have a special interest in those $F \in \mathcal{MU}$ where the lower bound $\text{fc}(F)$ meets the upper bound $\mu\text{vd}(F)$. In this case this number must be even, and we obtain another $F' \in \mathcal{MU}$ by resolving on any variable realising the minimum variable degree (where any such variable only occurs in full clauses here):

Lemma 12.14 Consider $F \in \mathcal{MU}$ with $\text{fc}(F) = \mu\text{vd}(F)$. Then $\text{fc}(F)$ is even.

Proof: Consider $v \in \text{var}(F)$ with $\text{vd}_F(v) = \mu\text{vd}(F)$. The occurrences of v are now exactly in the full clauses of F . Every full clause C must be resolvable on v with another full clause D , yielding $E := C \diamond D$, and thus the full clauses of F can be partitioned into pairs $\{v\} \cup E, \{\bar{v}\} \cup E$ (disjoint unions) for $\frac{\text{fc}(F)}{2}$ many clauses E (of length $n(F) - 1$; note that because of fullness for a given E the clauses C, D are uniquely determined up to order). \square

Thus, if lower and upper bound match, they must be even numbers:

Corollary 12.15 If $\text{FCM}(k) = \text{nM}(k)$ or $\text{FCM}(k) = \mu\text{vd}(\mathcal{MU}_{\delta=k})$ for some $k \in \mathbb{N}$, then $\text{FCM}(k)$ is even.

Another property of $\text{fc}(\mathcal{MU}_{\delta=k})$ related to evenness is that if m is a possible number of full clauses, then $2m$ is a possible number for $\delta = k + m - 1$:

Lemma 12.16 $2m \in \text{fc}(\mathcal{MU}_{\delta=k+m-1})$ for $k \in \mathbb{N}$ and $m \in \{1, \dots, \nu\text{fc}(\mathcal{MU}_{\delta=k})\}$.

Proof: Consider $F \in \mathcal{MU}_{\delta=k}$ with $\text{fc}(F) = \nu\text{fc}(\mathcal{MU}_{\delta=k})$. Choose m of the full clauses of F , and choose a new variable $v \notin \text{var}(F)$. Replace each of the chosen full clauses $C \in F$ by two clauses $C \cup \{v\}, C \cup \{\bar{v}\}$ (one non-strict and $m - 1$ strict full subsumption extensions), obtaining F' . We have $F' \in \mathcal{MU}_{\delta=k+m-1}$ and $\text{fc}(F') = 2m$. \square

As a special case we obtain that 2, 4 are always possible for the number of full clauses (except for $k = 1$):

Corollary 12.17 For $k \in \mathbb{N}$ holds $2 \in \text{fc}(\mathcal{MU}_{\delta=k})$, and if $k \geq 2$ then $4 \in \text{fc}(\mathcal{MU}_{\delta=k})$.

Proof: We show the assertion by induction over k , using Lemma 12.16, as follows: We have $\{\{1\}, \{-1\}\} \in \mathcal{MU}_{\delta=1}$, so consider $k \geq 2$. We know $2 \in \text{fc}(\mathcal{MU}_{\delta=k-1})$, thus $4 \in \text{fc}(\mathcal{MU}_{\delta=k-1+2-1=k})$. And once we have any $F \in \mathcal{MU}_{\delta=k}$ with a full clause, we get $F' \in \mathcal{MU}_{\delta=k}$ with $\text{fc}(F') = 2$ by performing a non-strict full subsumption extension on that full clause (i.e., $F' = E(F)$ as in Lemma 10.13, with C a full clause). \square

We now turn to the determination of $\nu\text{fc}(\mathcal{MU}_{\delta=k})$ for $k = 1, 2, 3$.

Lemma 12.18 We have:

1. $\text{FCH}(1) = \text{FCM}(1) = 2 = \text{nM}(1)$.
2. $\text{FCH}(2) = \text{FCM}(2) = 4 = \text{nM}(2)$.

3. $\text{FCM}(3) = 4 = \text{nM}(3) - 1$.

Proof: Part 1: By Corollary 12.9 we have $\nu\text{fc}(\mathcal{MU}_{\delta=1}) \leq 2$ (which can also be deduced from the fact, that between two clauses of $F \in \mathcal{MU}_{\delta=1}$ there is at most one conflict). Due to $A_1 \in \mathcal{UHT}_{\delta=1}$ we have $\nu\text{fc}(\mathcal{UHT}_{\delta=1}) \geq 2$.

Part 2: By $A_2 \in \mathcal{UHT}_{\delta=2}$ holds $\nu\text{fc}(\mathcal{UHT}_{\delta=2}) \geq 4$, by Corollary 12.9 we have $\nu\text{fc}(\mathcal{MU}_{\delta=2}) \leq 4$.

Part 3: By Corollary 12.17 we have $\nu\text{fc}(\mathcal{MU}_{\delta=3}) \geq 4$, by Corollary 12.15 we have $\nu\text{fc}(\mathcal{MU}_{\delta=3}) \leq 4$. \square

In Corollary 14.2 together with Theorem 14.4 these determinations will be completed for $k \leq 6$.

13. A method for improving upper bounds for VDM

We now present a framework for generalising the argumentation of Theorem 8.3 together with the analysis of the underlying recursion from Section 7. The idea is as follows:

1. We start with upper bounds $\text{VDM}(k) \leq a_k$ for $k = 1, \dots, p$, collected in a “valid bounds-function” f .
2. For deficiency $p + 1$ and an envisaged min-var-degree m we consider the set $\text{pp}_f(p + 1, m)$ of “possible” degree-pairs of variables (the degrees of the positive and negative literals) in an envisaged clause-set $F \in \mathcal{SMU}_{\delta=p+1}$ with $\mu\text{vd}(F) = m$.
3. If $\text{pp}_f(p + 1, m) = \emptyset$, then m is “inconsistent”, that is, impossible to realise, and thus $\text{VDM}(p + 1) < m$ (as shown in Theorem 13.10).
4. While in case of $\text{pp}_f(p + 1, m) \neq \emptyset$ there might exist such an F or not (the formal reasoning underlying the definition of $\text{pp}_f(p + 1, m)$ is not complete).

So here we generalise the approach of Section 7 for describing the function nM to a general recursion scheme, obtaining a general method for improved *upper bounds*. The applications in this report are as follows:

- In Theorem 13.15 we obtain an alternative description of $\text{nM}(k)$.
- In Section 14 we will first show that the smallest k , where we don’t have equality, is $k = 6$, namely $\mu\text{vd}(\mathcal{MU}_{\delta=6}) = 8 = \text{nM}(6) - 1$ (Theorem 14.4).
- By the general recursion scheme then follows from this improvement of the upper bound, that for all $k = 2^m - m + 1$ for $m \geq 3$ we have $\mu\text{vd}(\mathcal{MU}_{\delta=k}) \leq \text{nM}(k) - 1$. This improved upper bound is denoted by $\text{nM}_1 : \mathbb{N} \rightarrow \mathbb{N}$ (Theorem 14.6).

13.1. Analysing splitting-situations

“Valid bounds-functions” shall be monotonically increasing — we know that nM is (strictly) monotonically increasing, and we show that μvd is also monotonically increasing (not strictly, as we will later see in Theorem 14.4):

Lemma 13.1 *The map VDM is monotonically increasing ($\text{VDM}(k) \leq \text{VDM}(k+1)$) for $k \in \mathbb{N}$).*

Proof: For $F \in \mathcal{MU}_{\delta=k}$, $n(F) \neq 0$, we can construct $F' \in \mathcal{MU}_{\delta=k+1}$ with $\mu\text{vd}(F) \leq \mu\text{vd}(F')$ as follows:

1. If F is full, then obtain a non-full $F'' \in \mathcal{MU}_{\delta=k}$ with $\mu\text{vd}(F) = \mu\text{vd}(F'')$ by a full singular unit-extension (Lemma 5.17), and replace F by F'' .
2. If F is not full, then first saturate F , and then perform a strict full subsumption extension (Lemma 6.5), obtaining the desired F' . \square

We define now “valid bounds-functions”, which are sensible as upper bounds on VDM, and we also define how to obtain such a function from initial upper bounds $\text{VDM}(k) \leq a_k$ for $k = 1, \dots, p$:

Definition 13.2 *A **valid bounds-function** is a function $f : \mathbb{N} \rightarrow \mathbb{N} \cup \{+\infty\}$ fulfilling the following three conditions:*

1. $f(1) = 2$.
2. f is monotonically increasing (i.e., $\forall k, k' \in \mathbb{N} : k \leq k' \Rightarrow f(k) \leq f(k')$).
3. $f(k)$ is an upper bound for the minimal-variable degree of minimally unsatisfiable clause-sets of deficiency k (i.e., $\forall k \in \mathbb{N} : \text{VDM}(k) \leq f(k)$).

The set of all valid bounds-functions is denoted by $\mathcal{VB} \subset \mathbb{N}^{\mathbb{N}}$. And by $\mathcal{VB}^* := \{f \in \mathcal{VB} : f \leq \text{nM}\}$ we denote the set of valid bounds-functions (pointwise) less-or-equal than the non-Mersenne function.

For $a_1, \dots, a_p \in \mathbb{N}$, $p \in \mathbb{N}$. such that $a_1 = 2$, $a_i \leq a_j$ for $i \leq j$, and $a_i \geq \text{VDM}(i)$, we define $[a_1, \dots, a_p]$ as that $f \in \mathcal{VB}$ with $f(k) = a_k$ for $k \in \{1, \dots, p\}$, while $f(k) = \infty$ for $k > p$.

By Lemma 13.1, VDM is a valid bounds-function, namely the smallest possible one. By Theorem 8.3 and Corollary 7.6 also nM is a valid bounds-function. In Corollary 13.16 we will see, that the continuation of $[a_1, \dots, a_p]$ by just ∞ is harmless in the sense, that nM is automatically taken into account via the improvement of valid bounds-functions through the use of potential degree-pairs defined below.

Lemma 13.3 \mathcal{VB} as well as \mathcal{VB}^* , together with \leq , is a complete lattice, where infima resp. suprema are given by pointwise minimum resp. pointwise supremum. The smallest elements of both lattices is VDM, while the largest is $[2]$ resp. nM.

The following definition reflects the main method to analyse and improve a given upper bound f for $\text{VDM}(k)$, namely it determines the numerical possibilities compatible with f :

Definition 13.4 *Consider $k, m \in \mathbb{N}$ with $k \geq 2$ and $m \geq 4$, together with a valid bounds-function f . The **set of potential (variable-)degree-pairs w.r.t. f** for (deficiency) k and (minimum variable-degree) m , denoted by $\text{pp}_f(\mathbf{k}, \mathbf{m})$, is the set of pairs $(e_0, e_1) \in \mathbb{N}^2$ fulfilling the following conditions:*

- (i) $e_0, e_1 \geq 2$
- (ii) $e_0, e_1 \leq k$
- (iii) $e_0 + e_1 = m$
- (iv) $e_0 \leq e_1$
- (v) $\forall \varepsilon \in \{0, 1\} : f(k - e_\varepsilon + 1) + e_\varepsilon \geq m$.

We set $\text{pp}(\mathbf{k}, \mathbf{m}) := \text{pp}_{\text{VDM}}(\mathbf{k}, \mathbf{m})$.

The motivation for Definition 13.4 is to assume $F \in \mathcal{SMU}_{\delta=k}$ with $\mu\text{vd}(F) = m$ and $v \in \text{var}_{\mu\text{vd}}(F)$, and to determine the possible literal-degrees $e_0 = \text{ld}_F(\bar{v})$, $e_1 = \text{ld}_F(v)$, “possible” in a formal sense. “e” stands for “eliminated clauses”, namely e_ε is the number of clauses eliminated by $\langle v \rightarrow \varepsilon \rangle$. The “high” values of m (for fixed k) are of real interest; compare Lemma 13.7. The basic properties of pp_f are as follows:

1. For every valid f and $k \geq 2$ we have $\text{pp}_f(k, 4) = \{(2, 2)\}$ and $\text{pp}_f(k, m) = \emptyset$ for $m > 2k$.
2. Discussion of the five conditions (i) - (v) in Definition 13.4:
 - (i) Only nonsingular variables are considered, since only in this way the deficiency strictly decreases.
 - (ii) The deficiency of $F_\varepsilon := \langle v \rightarrow \varepsilon \rangle * F$ is $k_\varepsilon := k - e_\varepsilon + 1$ (assuming we split on a variable with minimal degree), and we must have $k_\varepsilon \geq 1$.
 - (iii) e_0, e_1 are the literal-degrees of \bar{v}, v , which sum up to the variable-degree m of v .
 - (iv) W.l.o.g. we can restrict attention to such degree-pairs, since F plays a role only up to isomorphism, and thus one can flip the sign of v in F .
 - (v) We have $F_\varepsilon \in \mathcal{MU}_{\delta=k-e_\varepsilon+1}$ (assuming F is saturated). And for $w \in \text{var}(F_\varepsilon)$ we have $\text{vd}_F(w) \leq \text{vd}_{F_\varepsilon}(w) + e_\varepsilon$. If for some $\varepsilon \in \{0, 1\}$ we would have $\mu\text{vd}(\mathcal{MU}_{\delta=k-e_\varepsilon+1}) + e_\varepsilon < m$, then for $w \in \text{var}_{\mu\text{vd}}(F_\varepsilon)$ we would have $\text{vd}_F(w) \leq \text{vd}_{F_\varepsilon}(w) + e_\varepsilon \leq \mu\text{vd}(\mathcal{MU}_{\delta=k-e_\varepsilon+1}) + e_\varepsilon < m$, but by assumption on w we have $\text{vd}_F(w) \geq m$.
3. An important special case of $\text{pp}_f(k, m)$ is $\text{pp}_{\text{nM}}(k, m)$; we have $\text{pp}(k, m) \subseteq \text{pp}_{\text{nM}}(k, m)$ (see Lemma 13.8 for a generalisation). The main point in using functions f is that the precise values of $\text{VDM}(k)$ might not be known.
4. To compute $\text{pp}_f(k, m)$ according to the definition, only the values $f(k')$ for $k' \in \{1, \dots, k-1\}$ are needed.

Example 13.5 Consider $f := [\text{nM}(1), \text{nM}(2), \text{nM}(3)] = [2, 4, 5]$. First we determine $\text{pp}_f(4, 7)$:

1. By Conditions (i) - (iv) only $\{(3, 4)\}$ remains.
2. Now $f(4 - 3 + 1) + 3 = f(2) + 3 = 7$, but $f(4 - 4 + 1) + 4 = f(1) + 4 = 6 < 7$.
3. Thus $\text{pp}_f(4, 7) = \emptyset$.

We will see in Theorem 13.10 that from this we can conclude $\text{VDM}(4) \leq 6$ (there is no “formal” possibility to reach the min-var degree of 7 for deficiency 4). Now we determine $\text{pp}_f(4, 6)$:

1. By Conditions (i) - (iv), $\{(2, 4), (3, 3)\}$ are the possibilities.
2. Checking Condition (v) for $(2, 4)$: $f(4 - 2 + 1) + 2 = f(3) + 2 = 7$, $f(4 - 4 + 1) + 4 = f(1) + 4 = 6$.
3. Checking Condition (v) for $(3, 3)$: $f(4 - 3 + 1) + 3 = f(2) + 3 = 7$.
4. Thus $\text{pp}_f(4, 6) = \{(2, 4), (3, 3)\}$.

The intuitive meaning of this is, that a min-var-degree of 6 can not be excluded by this type of formal reasoning, and 6 is the first new value according to this reasoning for $[2, 4, 5]$.

We invite the reader to compute the following special case of what we show later (in the proof of Theorem 14.7; it might also be useful to consider Table 2):

Example 13.6 $\text{pp}_{\text{nM}}(13, 17) = \{(8, 9)\}$, while for any valid bounds-function f with $f(k) = \text{nM}(k)$ for $k \in \{1, \dots, 5\}$ and $f(6) = \text{nM}(6) - 1 = 8$ holds $\text{pp}_f(13, 17) = \emptyset$.

If we have a potential degree-pair for m , then also for $m' \leq m$:

Lemma 13.7 Consider $k, m, m' \in \mathbb{N}$ with $k \geq 2$ and $4 \leq m' \leq m$, and consider a valid bounds-function f . If $\text{pp}_f(k, m) \neq \emptyset$, then also $\text{pp}_f(k, m') \neq \emptyset$.

Proof: Consider $(e_0, e_1) \in \text{pp}_f(k, m)$. Consider any $2 \leq e'_0 \leq e_0$ and $2 \leq e'_1 \leq e_1$ with $e'_0 \leq e'_1$ and $e'_0 + e'_1 = m'$. Now $f(k - e'_\varepsilon + 1) + e'_\varepsilon \geq f(k - e_\varepsilon + 1) + e'_\varepsilon = f(k - e_\varepsilon + 1) + e_\varepsilon - e_\varepsilon + e'_\varepsilon \geq m - e_\varepsilon + e'_\varepsilon = m - (m - e_\varepsilon) + (m' - e'_\varepsilon) = e_\varepsilon + m' - e'_\varepsilon \geq m'$ for $\varepsilon \in \{0, 1\}$, and thus $(e'_0, e'_1) \in \text{pp}_f(k, m') \neq \emptyset$. \square

Using a smaller bounds-function can not yield more potential degree-pairs, as is obvious from Definition 13.4:

Lemma 13.8 Consider $k, m \in \mathbb{N}$ with $k \geq 2$, $m \geq 4$, and valid bounds-functions f_1, f_2 with $f_1 \leq f_2$ (pointwise). Then $\text{pp}_{f_1}(k, m) \subseteq \text{pp}_{f_2}(k, m)$. Especially for any valid bounds-function f holds $\text{pp}(k, m) \subseteq \text{pp}_f(k, m)$.

Again directly by definition (using monotonicity of valid bounds functions) we get that increasing k while keeping m can not remove potential degree-pairs:

Lemma 13.9 Consider $k, m \in \mathbb{N}$ with $k \geq 2$, $m \geq 4$, and a valid bounds-function f . Then $\text{pp}_f(k, m) \subseteq \text{pp}_f(k + 1, m)$.

The main use of potential degree-pairs is to provide upper bounds on $\text{VDM}(k)$:

Theorem 13.10 Consider $k, m \in \mathbb{N}$ with $k \geq 2$, $m \geq 4$, and a valid bounds-function f . If $\text{pp}_f(k, m) = \emptyset$, then $\text{VDM}(k) < m$.

Proof: Assume $\mu\text{vd}(\mathcal{MU}_{\delta=k}) \geq m$. Then there is $F \in \mathcal{SMU}'_{\delta=k}$ with $\mu\text{vd}(F) \geq m$ (using Corollary 5.5). Consider $v \in \text{var}_{\mu\text{vd}}(F)$; if $\text{ld}_F(\bar{v}) \leq \text{ld}_F(v)$ holds, then let $e := (\text{ld}_F(\bar{v}), \text{ld}_F(v))$, while otherwise flip the components of this pair. Now we have $e \in \text{pp}(k, \mu\text{vd}(F))$ (using Remark 2 to Definition 13.4), and thus $\text{pp}_f(k, m) \neq \emptyset$ by Lemmas 13.7, 13.8, contradicting the assumption. \square

13.2. Recursion on potential degree-pairs

Via potential degree-pairs and Theorem 13.10, we obtain a method for improving valid bounds-functions:

Lemma 13.11 Consider $f \in \mathcal{VB}$. We obtain $f' \in \mathcal{VB}$ recursively as follows:

1. $f'(1) := 2$.
2. For $k > 1$ consider the largest $4 \leq m \leq 2k$ such that $\text{pp}_{f'}(k, m) \neq \emptyset$, using Remark 4 to Definition 13.4 (that we only need $f'(k')$ for $k' < k$).
3. Now $f'(k) := \min(m, f(k))$.

Proof: $f'(k)$ is well-defined for $k > p$ due to Remark 1 to Definition 13.4. That f' is valid follows by induction as follows. We have to show $f'(k) \leq f'(k+1)$ and $\text{VDM}(k) \leq f'(k)$ for all $k \in \mathbb{N}$. For $k = 1$ both properties are true by definition. And the induction step follows for monotonicity by Lemma 13.9, and for the upper-bound-condition by Theorem 13.10. \square

The mapping $f \in \mathcal{VB} \mapsto f' \in \mathcal{VB}$ we call the “non-Mersenne operator”:

Definition 13.12 For $f \in \mathcal{VB}$ let the $f' \in \mathcal{VB}$ according to Lemma 13.11 be denoted by $\text{NM}(f) := f'$ (defined via “recursion on potential degree-pairs”); we call $\text{NM} : \mathcal{VB} \rightarrow \mathcal{VB}$ the “non-Mersenne operator”.

The basic properties of the non-Mersenne operator are that of a kernel operator, which are order-theoretic properties as follows:

Lemma 13.13 The map $\text{NM} : \mathcal{VB} \rightarrow \mathcal{VB}$ is a kernel operator of the complete lattice \mathcal{VB} , that is, for all $f, g \in \mathcal{VB}$ holds:

1. $\text{NM}(f) \leq f$ (intensive)
2. $\text{NM}(\text{NM}(f)) = \text{NM}(f)$ (idempotent)
3. $f \leq g \Rightarrow \text{NM}(f) \leq \text{NM}(g)$ (monotonically increasing).

Proof: Intensivity follows by definition of NM (note that in Lemma 13.11 we have defined $f'(k)$ such that $f'(k) \leq f(k)$ holds). Also idempotence follows directly from the definition in Lemma 13.11, namely that $f'(k)$ for $k > 1$ already uses the improved values $f'(k')$ for $k' < k$. Monotonicity follows by Lemma 13.8. \square

By Lemma 13.13 we get that $\text{NM}(f)$ for $f \in \mathcal{VB}$ is the supremum of the set of $f' \leq f$ with $\text{NM}(f') = f'$. By Theorem 13.10 we get $\text{NM}(\text{VDM}) = \text{VDM}$. In order to show that the non-Mersenne operator at most reproduces nM , that is, for all $f \in \mathcal{VB}$ holds $\text{NM}(f) \leq \text{nM}$, we need to provide potential degree-pairs for nM :

Lemma 13.14 For $k \geq 2$ we have (recall Definition 7.13):

1. $(h(k), i_{\text{nM}}(k)) \in \text{pp}_{\text{nM}}(k, \text{nM}(k))$.
2. $\text{pp}_{\text{nM}}(k, \text{nM}(k) + 1) = \emptyset$.

Proof: Let $m := \text{nM}(k)$. For Part 1 let $e_0 := h(k)$, $e_1 := i_{\text{nM}}(k)$; so we have to show $(e_0, e_1) \in \text{pp}_{\text{nM}}(k, m)$. Consider the conditions (i) - (v) in Definition 13.4. We have $e_0 \geq 2$, since $\text{nM} \geq 2$ in general, and $e_1 \geq 2$ by Definition 7.8. As shown in Corollary 7.11 we have $e_0 \leq e_1$, where $e_1 \leq k$ by definition. Furthermore we have $e_0 + e_1 = m$ by Lemma 7.10, Part 3. Altogether we have now shown conditions (i) - (iv), and it remains to show that $\text{nM}(k - e_\varepsilon + 1) + e_\varepsilon \geq m$ holds for both $\varepsilon \in \{0, 1\}$; for $\varepsilon = 1$ we have equality, as already remarked, and it remains to show $\text{nM}(k - e_0 + 1) + e_0 \geq m$, which is equivalent to

$$\text{nM}(k - e_0 + 1) \geq i_{\text{nM}}(k).$$

By Definition 7.8 of $i_{\text{nM}}(k)$ (as the smallest i) this is implied by $\text{nM}(k - e_0 + 1) \geq \text{nM}(k - \text{nM}(k - e_0 + 1) + 1)$. By the monotonicity of nM this is implied by $e_0 \leq \text{nM}(k - e_0 + 1)$, i.e., $\text{nM}(k - i_{\text{nM}}(k) + 1) \leq \text{nM}(k - e_0 + 1)$. Again by monotonicity, this is implied by $i_{\text{nM}}(k) \geq e_0$, i.e., $e_1 \geq e_0$, which we have already shown.

For Part 2 we have to show $\text{pp}_{\text{nM}}(k, m + 1) = \emptyset$. Assume that we have $(e_0, e_1) \in \text{pp}_{\text{nM}}(k, m + 1)$ according to Definition 13.4. Thus we have $\text{nM}(k - e_1 + 1) + e_1 \geq$

$m+1$, where $2 \leq e_1 \leq k$. Because of $e_0+e_1 = m+1$ and $e_0 \leq e_1$ we get $e_1 \geq \frac{1}{2}(m+1)$, whence $\min(2 \cdot e_1, \text{nM}(k-e_1+1)+e_1) \geq m+1$, and thus $\text{nM}(k) \geq m+1$ by Definition 7.1. \square

We obtain an alternative recursion for $\text{nM}(k)$ (recall Definition 7.1):

Theorem 13.15 $\text{NM}([2]) = \text{NM}(\text{nM}) = \text{nM}$.

Proof: By Definition 13.12 and Lemma 13.14 we get $\text{NM}([2]) = \text{nM}$. Since NM is idempotent, we also get $\text{NM}(\text{nM}) = \text{nM}$. \square

So the non-Mersenne operator yields nM in the worst-case:

Corollary 13.16 $\text{NM} : \mathcal{VB} \rightarrow \mathcal{VB}^*$, that is, for every $f \in \mathcal{VB}$ holds $\text{NM}(f) \leq \text{nM}$.

14. Strengthening the upper bound for VDM

In this final section many techniques introduced in this report come together, and we give some initial sharpness results (considering small deficiencies), and some non-sharpness results in the form of improved bounds (improving nM for infinitely many deficiencies). In Subsection 14.1 we determine $\text{VDM}(k)$ for $1 \leq k \leq 6$ as values 2, 4, 5, 6, 8, 8, where the main achievement is Theorem 14.4, showing $\text{VDM}(6) = 8 = \text{nM}(6) - 1$. Applying the non-Mersenne operator, we obtain the improved upper bound $\text{VDM}(k) \leq \text{nM}_1(k)$ in Subsection 14.2, where nM_1 is like nM , but with a duplication after the jump positions, that is, $\Delta \text{nM}(k) = \Delta \text{nM}_1(k) = 2$ is followed by $\Delta \text{nM}_1(k+1) = \Delta \text{nM}(k+1) - 1 = 0$.

14.1. Deficiencies 1, ..., 6

We now show that the first deficiency k , for which the bound $\text{VDM}(k) \leq \text{nM}(k)$ is not sharp, is $k = 6$. First we show sharpness for the first five values:

Theorem 14.1 For $k \in \{1, \dots, 5\}$ we have $\text{VDM}(k) = \text{VDH}(k) = \text{nM}(k)$.

Proof: We have to give examples showing that the upper bound $\text{nM}(k)$ is attained for examples in $\mathcal{UHLT}_{\delta=k}$. Lemma 12.11 covers deficiencies $k = 1, 2, 5$, namely

1. $A_1 \in \mathcal{UHLT}_{\delta=1}$ has $\text{fc}(A_1) = 2 = \text{nM}(1)$ (recall Example 3.2).
2. $A_2 \in \mathcal{UHLT}_{\delta=2}$ has $\text{fc}(A_2) = 4 = \text{nM}(2)$ (recall Example 3.3).
3. $A_3 \in \mathcal{UHLT}_{\delta=5}$ has $\text{fc}(A_3) = 8 = \text{nM}(5)$.

Deficiency $k = 4$ is a jump position, and thus covered by Lemma 12.12, where the example is as follows:

4. For $F_4 := \{\{1, 2\}, \{-1, 2, 3\}, \{1, -2, 3\}, \{-1, -2, 3\}, \{-1, 2, -3\}, \{1, -2, -3\}, \{-1, -2, -3\}\}$ we have $F_4 \in \mathcal{UHLT}_{\delta=4}$ with $\text{fc}(F_4) = 6 = \text{nM}(4)$.

The remaining case $k = 3$ we obtain via strict full subsumption resolution from F_4 :

5. For $F_3 := \{\{1, 2\}, \{-1, 3\}, \{1, -2, 3\}, \{-1, 2, -3\}, \{1, -2, -3\}, \{-1, -2, -3\}\}$ we have $F_3 \in \mathcal{UHLT}_{\delta=3}$ with $\mu\text{vd}(F_3) = 5 = \text{nM}(3)$. \square

The examples in the proof of Theorem 14.1 together with Lemma 12.18 yield

Corollary 14.2 For $k \in \{1, \dots, 5\} \setminus \{3\}$ we have $\text{FCH}(k) = \text{FCM}(k) = \text{nM}(k)$, while $\text{FCH}(3) = \text{FCM}(3) = 4 = \text{VDM}(3) - 1 = \text{nM}(3) - 1$.

In the sequel of this subsection we consider $k = 6$. A computation shows that there is only one potential degree-pair for the min-var-degree as given by $\text{nM}(6) = 9$:

Lemma 14.3 $\text{pp}_{\text{nM}}(6, 9) = \{(4, 5)\}$.

Proof: Conditions (i) - (iv) of Definition 13.4 yield $\text{pp}_{\text{nM}}(6, 9) \subseteq \{(3, 6), (4, 5)\}$. Condition (v) excludes $(3, 6)$, since we have $\mu\text{vd}(\mathcal{MU}_{\delta=6-6+1}) + 6 = 8 \not\geq 9$, while $(4, 5)$ fulfils this condition due to $\text{nM}(6-4+1)+4 = 5+4 \geq 9$ and $\text{nM}(6-5+1)+5 = 4+5 \geq 9$. \square

However, the potential degree-pair of Lemma 14.3 actually can not be realised, and thus $\text{VDM}(6) < \text{nM}(6)$:

Theorem 14.4 $\text{FCH}(6) = \text{FCM}(6) = \text{VDH}(6) = \text{VDM}(6) = 8 = \text{nM}(6) - 1$.

Proof: $\nu\text{fc}(\mathcal{UHT}_{\delta=6}) \geq 8$ is confirmed by the variable-clause matrix

$$\begin{pmatrix} + & + & + & - & + & - & - & - & + & - \\ + & + & + & + & + & + & + & + & - & - \\ + & + & - & + & - & + & - & - & 0 & 0 \\ + & - & + & + & - & - & + & - & 0 & 0 \end{pmatrix}$$

(where unsatisfiability is given by $8 \cdot 2^{-4} + 2 \cdot 2^{-2} = 1$).

Assume now that there exists $F \in \mathcal{MU}_{\delta=6}$ with $\mu\text{vd}(F) = 9$. By Lemmas 5.3, 5.4 w.l.o.g. we can assume that F is saturated and nonsingular. By Theorem 6.13 we know $n(F) \geq 4$. Consider $v \in \text{var}(F)$ with $\text{vd}_F(v) = 9$. W.l.o.g. we assume $\text{ld}_F(v) \geq \text{ld}_F(\bar{v})$. By Lemma 14.3 we have $\text{ld}_F(v) = 5$, $\text{ld}_F(\bar{v}) = 4$, and $\delta(F_0) = 6 - 4 + 1 = 3$, $\delta(F_1) = 6 - 5 + 1 = 2$.

Let the 5 occurrences of v in F be $C_1, \dots, C_5 \in F$, and let $C'_i := C_i \setminus \{v\}$. And let the 4 occurrences of \bar{v} in F be $D_1, \dots, D_4 \in F$, and let $D'_i := D_i \setminus \{\bar{v}\}$. Using $G := \{C \in F : v \notin \text{var}(C)\} = F \setminus \{C_1, \dots, C_5, D_1, \dots, D_4\}$ we get

$$\begin{aligned} F_0 &= \{C'_1, \dots, C'_5\} \cup G \\ F_1 &= \{D'_1, \dots, D'_4\} \cup G \end{aligned}$$

where $c(F_0) = 5 + c(G) = c(F) - 4$ and $c(F_1) = 4 + c(G) = c(F) - 5$.

Consider first $F_0 \in \mathcal{MU}_{\delta=3}$. We have $\mu\text{vd}(F_0) \geq 9 - 4 = 5$, and thus $\mu\text{vd}(F_0) = 5$ (due to $\mu\text{vd}(F_0) \leq \text{nM}(3) = 5$). Every variable $w \in \text{var}(F_0)$ realising the min-var-degree of F_0 has at least 9 occurrences in F , from which at most 4 are eliminated, and thus actually such variables have $\text{vd}_F(w) = 9$, and furthermore $w \in \text{var}(D_i)$ for all $i \in \{1, \dots, 4\}$. By Lemma 8.5, Part 1, there exist two different variables $u_1, u_2 \in \text{var}(F_0)$ with $\text{vd}_{F_0}(u_i) = 5$ for $i \in \{1, 2\}$, and so we have $|D_i| \geq 3$ for all $i \in \{1, \dots, 4\}$.

Now consider $F_1 \in \mathcal{MU}_{\delta=2}$. We have $\mu\text{vd}(F_1) \geq 9 - 5 = 4$, thus $\mu\text{vd}(F_1) = 4$ (due to $\mu\text{vd}(F_1) \leq \text{nM}(2) = 4$), and thus by Lemma 5.13 F_1 is nonsingular iff F_1 does not contain unit-clauses. If F_1 would contain a unit-clause, then there would be a binary clause $\{\bar{v}, x\} \in F$, contradicting that all D_i contain at least three literals. So F_1 is nonsingular, and thus F_1 is isomorphic to some \mathcal{F}_m for some $m \geq 2$. So F_1 is 4-variable-regular, where all the variables of F_1 have at least 9 occurrences in F , and thus we have $\text{var}(F_1) \subseteq \text{var}(C'_i)$ for all $i \in \{1, \dots, 5\}$, which implies that actually $\text{var}(C'_i) = \text{var}(F_1) = \text{var}(F_0)$ holds.

Coming back to the structure of F_0 , we now know that F_0 has five full clauses C'_1, \dots, C'_5 , which contradicts Lemma 12.18, Part 3. \square

14.2. Sharpening the bound

Based on recursion on potential degree-pairs, we can improve the upper bound $\text{nM}(k)$ for $\mu\text{vd}(\mathcal{MU}_{\delta=k})$ for $k \geq 6$ (generalising Example 13.6):

Definition 14.5 Let $\text{nM}_1 : \mathbb{N} \rightarrow \mathbb{N}$ be defined as $\text{nM}_1 := \text{NM}([2, 4, 5, 6, 8, 8])$ (recall Definition 13.12).

By Lemma 13.11 together with Theorem 14.4 we get:

Theorem 14.6 For all $k \in \mathbb{N}$ we have $\text{VDM}(k) \leq \text{nM}_1(k)$.

It remains to determine nM_1 numerically:

Theorem 14.7 In Table 3 we find the values of $\text{nM}_1(k)$ for $k \leq 30$. We have $\text{nM}_1(k) = \text{nM}(k)$ for $k \notin \{2^m - m + 1 : m \in \mathbb{N}, m \geq 3\}$, while for $k = 2^m - m + 1$ we have $\text{nM}_1(k) = \text{nM}(k) - 1 = 2^m$.

k	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$\text{nM}_1(k)$	2	4	5	6	8	<u>8</u>	10	11	12	13	14	16	<u>16</u>	18	19
k	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
$\text{nM}_1(k)$	20	21	22	23	24	25	26	27	28	29	30	32	<u>32</u>	34	35

Table 3: Values of $\text{nM}_1(k)$ for $k \in \{1, \dots, 30\}$, in bold the jump-values (i.e., $k \in J$), and underlined the changed values compared to $\text{nM}(k)$; we see that directly after the jump we have stagnation, followed by a second jump.

Proof: We show the formula for $\text{nM}_1(k)$ via induction on k . Due to $2^3 - 3 + 1 = 6$ it holds for $k \leq 6$. So assume $k \geq 7$. We show by induction on $k \geq 7$ the following two properties (which imply the assertions of the theorem):

1. For $k = 2^m - m + 1$, $m \geq 4$, we have
 - (a) $\text{pp}_{\text{nM}_1}(k, \text{nM}(k)) = \emptyset$ and
 - (b) and $\text{pp}_{\text{nM}_1}(k, \text{nM}(k) - 1) \neq \emptyset$.
2. Otherwise $\text{pp}_{\text{nM}_1}(k, \text{nM}(k)) \neq \emptyset$.

By Lemma 13.13 we know $\text{nM}_1 \leq \text{nM}$.

Part 1. We consider $k = 2^m - m + 1$, $m \geq 4$. We have $\text{nM}(k) = 2^m + 1$.

Part (a) To show $\text{pp}_{\text{nM}_1}(2^m - m + 1, 2^m + 1) = \emptyset$, we assume $(e_0, e_1) \in \text{pp}_{\text{nM}_1}(2^m - m + 1, 2^m + 1)$. Thus we know $e_0, e_1 \geq 2$, $e_0, e_1 \leq 2^m - m + 1$, $e_0 + e_1 = 2^m + 1$, $e_0 \leq e_1$, whence $e_0 \leq 2^{m-1}$, and

$$\text{nM}_1(2^m - m + 1 - e_\varepsilon + 1) + e_\varepsilon \geq 2^m + 1 \quad (1)$$

for both $\varepsilon \in \{0, 1\}$.

Case (a.1). Assume $e_0 \leq 2^{m-1} - 1$, and thus $e_1 \geq 2^{m-1} + 2$.

From (1) we get $\text{nM}(2^m - m + 1 - e_1 + 1) + e_1 \geq 2^m + 1$, where (using Corollary 7.6):

$$\begin{aligned} \text{nM}(2^m - m + 1 - e_1 + 1) + e_1 &\geq 2^m + 1 \Rightarrow \\ \text{nM}(2^m - m + 1 - (2^{m-1} + 2) + 1) + 2^{m-1} + 2 &\geq 2^m + 1 \Leftrightarrow \\ \text{nM}(2^{m-1} - m) &\geq 2^{m-1} - 1, \end{aligned}$$

where by Corollary 7.25 we have $\text{nM}(2^{m-1} - m) = \text{nM}(2^{m-1} - (m-1) - 1) = 2^{m-1} - 2$, and we obtained a contradiction, finishing Case (a.1).

Case (a.2). It remains $e_0 = 2^{m-1}$. From (1) we get $\text{nM}_1(2^m - m + 1 - e_0 + 1) + e_0 \geq 2^m + 1$, where $2^m - m + 1 - e_0 + 1 = 2^m - m + 1 - 2^{m-1} + 1 = 2^{m-1} - (m-1) + 1$, and thus by induction hypothesis we get $\text{nM}_1(2^m - m + 1 - e_0 + 1) + e_0 = 2^{m-1} + 2^{m-1} = 2^m$, a contradiction. This concludes Part (a).

Part (b). We show $(2^{m-1}, 2^{m-1}) \in \text{pp}_{\text{nM}_1}(k, \text{nM}(k) - 1)$.¹⁷ For this it remains to show $\text{nM}_1(2^m - m + 1 - 2^{m-1} + 1) + 2^{m-1} \geq 2^m$, and indeed $\text{nM}_1(2^m - m + 1 - 2^{m-1} + 1) = \text{nM}_1(2^{m-1} - (m-1) + 1) = 2^{m-1}$ by induction hypothesis. This concludes Part 1.

Part 2. $k \neq 2^m - m + 1$ for any $m \geq 4$. We have to show $\text{pp}_{\text{nM}_1}(k, \text{nM}(k)) \neq \emptyset$.

Part (a). $k = 2^m - m + 2$; thus $\text{nM}(k) = 2^m + 2$.

We have $(2^{m-1}, 2^{m-1} + 2) \in \text{pp}_{\text{nM}_1}(k, 2^m + 2)$ ¹⁸, due to $\text{nM}_1(2^m - m + 2 - 2^{m-1} + 1) + 2^{m-1} = \text{nM}_1(2^{m-1} - (m-1) + 2) + 2^{m-1} = 2^{m-1} + 2 + 2^{m-1}$ and $\text{nM}_1(2^m - m + 2 - (2^{m-1} + 2) + 1) + 2^{m-1} + 2 = \text{nM}_1(2^{m-1} - (m-1)) + 2^{m-1} + 2 = 2^{m-1} + 2^{m-1} + 2$.

Part (b). $k = 2^m - m + 3$; thus $\text{nM}(k) = 2^m + 3$.

We have $(2^{m-1}, 2^{m-1} + 3) \in \text{pp}_{\text{nM}_1}(k, 2^m + 3)$ ¹⁹, due to $\text{nM}_1(2^m - m + 3 - 2^{m-1} + 1) + 2^{m-1} = \text{nM}_1(2^{m-1} - (m-1) + 3) + 2^{m-1} = 2^{m-1} + 3 + 2^{m-1}$ and $\text{nM}_1(2^m - m + 3 - (2^{m-1} + 3) + 1) + 2^{m-1} + 3 = \text{nM}_1(2^{m-1} - (m-1)) + 2^{m-1} + 3 = 2^{m-1} + 2^{m-1} + 3$.

For all remaining cases

$$2^m - m + 4 \leq k \leq 2^{m+1} - (m + 1)$$

we show $(e_0, e_1) := (h(k), i_{\text{nM}}(k)) \in \text{pp}_{\text{nM}_1}(k, \text{nM}(k))$, as in Lemma 13.14, Part 1. Recall Corollary 7.26 for the computation of $i_{\text{nM}}(k)$.

For the critical condition “ $\text{nM}_1(k - e_1 + 1) + e_1 \geq \text{nM}(k)$ ” we recall $k - i_{\text{nM}}(k) + 1 = i'(k)$ (recall Definition 7.13), where $i'(k)$ is monotonically increasing. We want to show that $i'(k)$ is never equal to some argument where $\text{nM}_1 \neq \text{nM}$, and thus we need to check the upper and the lower bounds:

For $k = 2^m - m + 4$ holds $i_{\text{nM}}(k) = 2^{m-1} + 2$, thus $i'(k) = 2^m - m + 4 - (2^{m-1} + 2) + 1 = 2^{m-1} - m + 3 > 2^{m-1} - (m-1) + 1$.

For $k = 2^{m+1} - (m+1)$ holds $i_{\text{nM}}(k) = 2^m$, thus $i'(k) = 2^{m+1} - (m+1) - 2^m + 1 = 2^m - m < 2^m - m + 1$.

So all critical conditions for e_1 are fulfilled, and it remains to check e_0 .

Part (c). $2^m - m + 4 \leq k \leq 2^{m+1} - (m + 1) - 2$.

Here for the critical conditions “ $\text{nM}_1(k - e_0 + 1) + e_0 \geq \text{nM}(k)$ ” the values $k - e_0 + 1$ are never equal to some argument where $\text{nM}_1 \neq \text{nM}$, since $\Delta(k - h(k) + 1) = 1 - \Delta h(k) \geq 0$ for $k \leq 2^{m+1} - (m + 1) - 2$ by Theorem 7.15, and thus $k - h(k) + 1$ is monotonically increasing for the k -range we consider.

Checking for the lower bound $k = 2^m - m + 4$: $i_{\text{nM}}(k) = 2^{m-1} + 2$, thus $h(k) = \text{nM}(k) - i_{\text{nM}}(k) = 2^m + 4 - 2^{m-1} - 2 = 2^{m-1} + 2$, which is the same as $i_{\text{nM}}(k)$, and doesn't need to be checked again.

Checking for the upper bound $k = 2^{m+1} - (m + 1) - 2$: $i_{\text{nM}}(k) = 2^m - 1$, and thus $h(k) = \text{nM}(k) - i_{\text{nM}}(k) = 2^{m+1} - 3 - 2^m + 1 = 2^m - 2$, and so we check $2^{m+1} - (m + 1) - 2 - (2^m - 2) + 1 = 2^m - m < 2^m - m + 1$.

Part (d). $k = 2^{m+1} - (m + 1) - 1$; thus $\text{nM}(k) = 2^{m+1} - 2$.

¹⁷We have $(2^{m-1}, 2^{m-1}) = (h(k), i_{\text{nM}}(k) - 1)$, but we don't need this here.

¹⁸We have $(2^{m-1}, 2^{m-1} + 2) = (h(k) - 1, i_{\text{nM}}(k) + 1)$.

¹⁹We have $(2^{m-1}, 2^{m-1} + 3) = (h(k) - 1, i_{\text{nM}}(k) + 1)$.

Now $i_{\text{nM}}(k) = 2^m$, and so $h(k) = \text{nM}(k) - i_{\text{nM}}(k) = 2^m - 2$, and so the critical argument is $2^{m+1} - (m+1) - 1 - (2^m - 2) + 1 = 2^m - m + 1$. It remains to check $\text{nM}_1(2^m - m + 1) + 2^m - 2 \geq 2^{m+1} - 2$, and indeed $\text{nM}_1(2^m - m + 1) + 2^m - 2 = 2^m + 2^m - 2$.

Part (e). $k = 2^{m+1} - (m+1)$; thus $\text{nM}(k) = 2^{m+1}$.

Now $h(k) = 2^m = i_{\text{nM}}(k)$, and no check is needed. \square

It is instructive to note the new Δ -values explicitly:

Corollary 14.8 *For $k \in \mathbb{N}$ holds $\Delta \text{nM}_1(k) \in \{0, 1, 2\}$, with*

1. $\Delta \text{nM}_1(k) = 0 \iff k = 2^m - m$ for some $m \in \mathbb{N}$, $m \geq 3$.
2. $\Delta \text{nM}_1(k) = 2 \iff k = 2^m - m \pm 1$ for some $m \in \mathbb{N}$, $m \geq 3$.

15. Conclusion and open problems

The main subject of this report can be seen in the study of $\mu\text{vd}(\mathcal{C}_{\delta=k})$ for classes $\text{UHIT} \subseteq \mathcal{C} \subseteq \text{MLEAN}$ and $k \in \mathbb{N}$, that is, the study of the maximal minimum variable-degree of classes of matching-lean clause-sets containing all minimally unsatisfiable clause-sets, parameterised by the deficiency. If $\mathcal{C} \subseteq \text{LEAN}$, then this quantity is bounded, and indeed we have shown $\mu\text{vd}(\text{LEAN}_{\delta=k}) = \text{nM}(k)$ (more generally this holds for every subclass of LEAN containing VMU). While for $\mathcal{C} = \text{MLEAN}$ this quantity is unbounded. For MU we have shown the improved bound $\mu\text{vd}(\text{MU}_{\delta=k}) \leq \text{nM}_1(k)$, where indeed also this bound is not sharp (as will be shown in [81]; see Subsection 15.2) — the question about the determination of $\text{VDM}(k) = \mu\text{vd}(\text{MU}_{\delta=k})$ is a major open research question.

For lean clause-sets we have shown the strengthened upper bound $\mu\text{vd}(F) \leq \text{nM}(\sigma(F))$, and indeed for every clause-set F we can satisfiability-equivalently remove some clauses in polynomial time such that this upper bound holds.

15.1. Conjectures and questions

We made the following conjectures:

1. Conjecture 10.3: If a clause-set violates the upper bound on the min-var-degree for lean clause-sets, then it must have a non-trivial autarky. As we have seen, we can determine the set of variables involved, but the determination of the autarky itself is open — the conjecture states that there is a poly-time algorithm for computing such an autarky. See Subsection 10.2 for more information on this topic.
2. Conjecture 12.1: the maximum min-var-degree for unsatisfiable hitting clause-sets is the same as for the larger class of minimally unsatisfiable clause-sets. In Conjecture 15.5 we generalise this to non-boolean clause-sets.
3. Conjecture 12.2: nM is not far away from the VDM, more precisely, $\text{nM} - 1 \leq \text{VDM} \leq \text{nM}$. The stronger Conjecture 12.6 says, that the same holds even for the maximal number of full clauses, that is, $\text{nM}(k) - 1 \leq \text{FCM}(k) \leq \text{nM}(k)$ for all $k \in \mathbb{N}$. In Lemma 15.3 we will state a weaker, but proven (in future work) lower bound.

Five more conjectures will be presented in this final section. We also asked the following questions:

1. Question 3.11 is about some complexity problems around the elimination of literal occurrences in minimally unsatisfiable clause-sets.

2. Question 9.8 is about the complexity of SAT decision for \mathcal{SED} . At first sight it might seem easy to translate every $F \in \mathcal{CLS}$ into some sat-equivalent element of \mathcal{SED} , and in fact to manipulate deficiency and surplus alone is rather easy, but we do not know how to handle them together.
3. Question 10.12 concerns the existence of unsatisfiable hitting clause-sets of arbitrary surplus equal deficiency and a min-var-degree as low as possible. An underlying question is to understand better the quantity “surplus”.
4. Question 11.2 is about strengthening the construction of Lemma 11.1, for finding matching-lean clause-sets of a given deficiency with high minimum literal-degree (perhaps completely different constructions are needed).
5. Questions 12.10 asks for the determination of $\nu\text{fc}(\mathcal{LEAN}_{\delta=k})$.
6. Question 12.13 is about lower bounds for $\text{VDH}(k)$, $\text{FCH}(k)$ via strict full subsumption resolution starting with A_n .

In the remainder we outline main research areas related to the topics of the report.

15.2. Improved upper bounds for VDM

We know $\text{VDM} \leq nM_1$, and we know $\text{VDM}(k)$ precisely for $k \in \{1, \dots, 6\}$ and for $k \in J, J + 1$. Also of high relevance here is to determine $\text{VDH}(k)$, which by Conjecture 12.1 is the same on $\text{VDM}(k)$. Another major conjecture is Conjecture 12.2, which says that VDM deviates at most by 1 from nM . Beyond the current report, we know the following improvements of the upper bound nM_1 :

- Generalising the ideas of Theorem 14.4, which is based on the improved upper bound for deficiency $2^3 - 3 + 1 = 6$, we can show also for deficiency $k = 2^4 - 4 + 2 = 14$ that we have $\text{VDM}(k) = nM(k) - 1$. Via the non-Mersenne operator, this yields the improved upper bound nM_2 .
- Altogether we obtain a sequence of improved upper bounds nM_{m-2} for $m \in \mathbb{N}$, $m \geq 3$, improving the upper bound at deficiency $k = 2^m - 2$ for nM_{m-3} and applying the non-Mersenne operator.
- The infimum of nM_1, nM_2, \dots is nM_ω . This will be developed in [81].
- However, this is not the end of it — also for deficiency $k = 15$ we have $\text{VDM}(k) = nM(k) - 1$, obtaining $nM_{\omega+1}$. This new improvement depends on new ideas — will there be an infinite chain of ever-increasing complexity of such improvements?

We believe that a closed “nice” formula for $\text{VDM}(k)$ is impossible, but that however computation is possible:

Conjecture 15.1 *The function $\text{VDM} : \mathbb{N} \rightarrow \mathbb{N}$ is “complex”, and for no finite tuple \vec{a} holds $\text{NM}(\vec{a}) = \text{VDM}$, but VDM is computable in doubly-exponential time.*

See Lemma 15.9 for some conditions which imply the computability-part of Conjecture 15.1.

15.3. Determining FCM

While Subsection 15.2 was about improving the upper bound, here now we turn to the lower bound. In Subsection 12.2 we provided only the minimum needed in this report for the measure $\text{fc}(F)$ of full clauses. In the forthcoming [82] we show the following lower bound, using $S_2 : \mathbb{N} \rightarrow \mathbb{N}$, the function for the ‘‘Smarandache Primitive Numbers’’ introduced in [99, Unsolved Problem 47], which for $k \in \mathbb{N}$ is defined as the minimal natural number $s \in \mathbb{N}$ such that 2^k divides $s!$.

Lemma 15.2 ([82]) *For all $k \in \mathbb{N}$ holds $\text{FCH}(k) \geq S_2(k)$.*

The basic structure of the proof of Lemma 15.2 is as follows: Full subsumption extension (strict and non-strict) yields a non-deterministic process to create elements of \mathcal{UHLT} with ‘‘many’’ full clauses. Maximising the number of full clauses for a given deficiency, a recursion is obtained, which is roughly similar to our recursion for $\text{nM}(k)$ (Definition 7.1). The analysis yields that the solution for this recursion is a *meta-Fibonacci sequence* as introduced in [100, Page 145], of a special form as analysed in [101]. We are then able to identify this special form as identical with $S_2(k)$ (as conjectured on the OEIS [98]).

Lemma 15.2 yields the interesting inequality $S_2 \leq \text{VDM} \leq \text{nM}$. This is relevant as the upper bound nM on S_2 as well as the lower bound S_2 on VDM . From [102] we get that $k + 1 \leq S_2(k)$ and thus by Corollary 7.23 we get

Lemma 15.3 ([82]) *$k + 1 \leq S_2(k) \leq \text{nM}(k) \leq k + 1 + \text{fld}(k)$ for $k \in \mathbb{N}$.*

Recall that in Lemma 12.3 we obtained a much sharper lower bound for $\text{VDM}(k)$ from Conjecture 12.2. For sequences $a, b : \mathbb{N} \rightarrow \mathbb{R}$ let asymptotic equality be denoted by $a \sim b : \Leftrightarrow \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1$. It is known that $S_2(k)$ and $\text{nM}(k)$ are asymptotically equal to $(k)_{k \in \mathbb{N}}$, and thus

Corollary 15.4 ([82]) *The six sequences S_2 , FCH , FCM , VDH , VDM , nM are asymptotically equal to $(k)_{k \in \mathbb{N}}$.*

In Figure 1 we show the six quantities from Corollary 15.4 and the relations between them (we do not mention the dependencies on the deficiency $k \in \mathbb{N}$ there). An arrow means a (proven) \leq -relation. If the arrow is labelled with ‘‘ -1 ’’, then we conjecture the difference is at most -1 (where in all three cases here we know examples where the difference is indeed equal to -1), while the label ‘‘ $=$ ’’ means that we conjecture equality, and the label ‘‘ ∞ ’’ means that we conjecture that the difference is unbounded.

For a more precise asymptotic determination of these six quantities from Corollary 15.4, calling them a_k , we need to consider the six sequences $(a_k - k)_{k \in \mathbb{N}}$. Currently we only know $\text{nM}(k) - k \sim \text{ld}(k)$.

15.4. Generalisation to non-boolean clause-sets

It is interesting to generalise Theorem 8.3 for generalised clause-sets; see [16, 17] for a systematic study, while the most general notion of generalised clause-sets, ‘‘signed clause-sets’’ are discussed in [103]. Generalised clause-sets F have literals (v, ε) , meaning ‘‘ $v \neq \varepsilon$ ’’, for variables v with non-empty finite domains D_v and values $\varepsilon \in D_v$. The deficiency is generalised by giving every variable a weight $|D_v| - 1 \in \mathbb{N}_0$ (which is 1 in the boolean case), i.e., $\delta(F) = c(F) - \sum_{v \in \text{var}(F)} (|D_v| - 1) = c(F) + n(F) - \sum_{v \in \text{var}(F)} |D_v|$; see [16, Subsection 7.2]. A partial assignment is a map φ with some finite set of variables as domain $\text{dom}(\varphi) =: \text{var}(\varphi)$, which maps $v \in \text{var}(\varphi)$ to $\varphi(v) \in D_v$. A partial assignment φ satisfies a clause-set F iff for every $C \in F$ there is $(v, \varepsilon) \in C$ with $v \in \text{var}(\varphi)$ and $\varphi(v) \neq \varepsilon$. Minimally




Figure 1: The four main combinatorial quantities, and the two numerical functions

unsatisfiable (generalised) clause-sets are defined as usual (they are unsatisfiable, while every strict subset is satisfiable). In [16, Corollary 9.9] it is shown that also all minimally unsatisfiable generalised clause-sets F fulfil $\delta(F) \geq 1$ (based, like in the boolean case, on matching autarkies).

The degree $\text{vd}_F(v)$ of a variable v in a clause-set F is the sum of the degrees of the literals (v, ε) for $\varepsilon \in D_v$, and thus $\text{vd}_F(v) = |C \in F : C \cap (\{v\} \times D_v) \neq \emptyset|$. For a given deficiency $k \in \mathbb{N}$, the basic question is to determine the supremum of $\mu\text{vd}(F)$ over all minimally unsatisfiable F with $\delta(F) = k$. The base case of deficiency $k = 1$ is handled in [17, Lemma 5.4], showing that for generalised minimally unsatisfiable clause-sets of deficiency 1 we have $\mu\text{vd}(F) \leq \max_{v \in \text{var}(F)} |D_v|$; actually all structural knowledge from [5, 6, 7] has been completely generalised in [17, Subsection 5.2].

But $k \geq 2$ requires more work, since here the basic method of saturation is not available for generalised clause-sets, as discussed in [17, Subsection 5.1]: saturated generalised clause-sets (i.e., unsatisfiable clause-sets, where no literal occurrence can be added without rendering the clause-set satisfiable) with deficiency at least 2 after splitting do not necessarily generate minimally unsatisfiable (generalised) clause-sets. Thus the proofs for the boolean case seem not to be generalisable for arbitrary minimally unsatisfiable (generalised) clause-sets.

In order to repair this, the “substitution stability parameter regarding irredundancy” $\text{sir}(F) \in \mathbb{Z}_{\geq -1} \cup \{+\infty\}$ is introduced in [17, Subsection 5.3], defined as the supremum of $k \in \mathbb{Z}_{\geq -1}$ such that for every partial assignment with $n(\varphi) := |\text{var}(\varphi)| \leq k$ the clause-set $\varphi * F$, obtained as usual by application of φ to F , is minimally unsatisfiable. So $\text{sir}(F) \geq 0$ iff F is minimally unsatisfiable, and as shown in [17, Corollary 4.8], $\text{sir}(F) = +\infty$ iff F is a hitting clause-set (i.e., for all $C, D \in F$, $C \neq D$, there are $x \in C$, $y \in D$ with $x = (v, \varepsilon)$ and $y = (v, \varepsilon')$ for some variable v and $\varepsilon, \varepsilon' \in D_v$ with $\varepsilon \neq \varepsilon'$). And $\text{sir}(F) \geq 1$ iff splitting on any variable yields always a minimally unsatisfiable clause-set. So for a boolean clause-sets F holds $\text{sir}(F) \geq 1$ iff F is saturated, but for generalised clause-sets we only have that $\text{sir}(F) \geq 1$ implies saturatedness ([17, Corollary 5.3]).

In [17, Corollary 5.10] one finds a generalisation of the basic bound $\mu\text{vd}(F) \leq 2\delta(F)$ for the boolean case. Namely $\mu\text{vd}(F) \leq \max_{v \in \text{var}(F)} |D_v| \cdot \delta(F)$ is shown for F with $\text{sir}(F) \geq 1$. Since for (generalised) saturated F with $\delta(F) = 1$ we have $\text{sir}(F) = \infty$ ([17, Corollary 5.6]), this covers the above mentioned result $\mu\text{vd}(F) \leq \max_{v \in \text{var}(F)} |D_v|$ for (arbitrary) minimally unsatisfiable F with $\delta(F) = 1$ (note that here saturation works as in the boolean case).

In [104] we concentrate on unsatisfiable hitting (generalised) clause-sets, and via generalised non-Mersenne numbers $nM^d(k)$ we are able to generalise Theorem 8.3 to generalised clause-sets. We believe that in general the minimum variable degree of minimally unsatisfiable clause-sets F with $\text{sir}(F) \geq 1$ for a given deficiency is always obtained by unsatisfiable hitting clause-sets (generalising Conjecture 12.2):

Conjecture 15.5 *Let $\mathcal{UHIT}_{\delta=k}^d$ denote the set of generalised unsatisfiable hitting clause-sets of deficiency $k \in \mathbb{N}$ and with domain-sizes at most $d \in \mathbb{N}$, and let $\mathcal{MU}_{\delta=k, \text{sir} \geq 1}^d$ be defined in the same way. Then we have for all $k, d \in \mathbb{N}$ that $\mu\text{vd}(\mathcal{UHIT}_{\delta=k}^d) = \mu\text{vd}(\mathcal{MU}_{\delta=k, \text{sir} \geq 1}^d)$.*

Furthermore, the “2” in $S_2(k)$ in Lemma 15.2 is comes from the boolean domain, and generalising the results of this report and from [82] in [104] to the non-boolean domain sheds light on $S_d(k)$ (the minimal $s \in \mathbb{N}$ such that d^k divides $s!$) for arbitrary prime numbers $d \in \mathbb{N}$, as introduced in [99, Unsolved Problem 49] (while for non-prime-numbers d the definition of S_d has to be generalised). See [105, Subsection III.1] for basic properties of $S_p(k)$.

15.5. Classification of \mathcal{MU}

As mentioned in the introduction, a major motivation for us is the project of the classification of minimally unsatisfiable clause-sets in the deficiency (recall Examples 3.2, 3.3), where the main conjecture is:

Conjecture 15.6 *For every deficiency $k \in \mathbb{N}$ there are finitely many “patterns” which determine the nonsingular elements of $\mathcal{MU}_{\delta=k}$, as well as the saturated and hitting cases amongst them. Especially for every k the isomorphism types of $\mathcal{MU}'_{\delta=k}$ can be efficiently enumerated (without repetitions), and for any given $F \in \mathcal{MU}'_{\delta=k}$ its isomorphism type can be determined in polynomial time.*

Conjecture 15.6 has been shown for $k \leq 2$ (recall Examples 3.2, 3.3). As we discussed in Subsection 1.6.1, the translation $e : \mathcal{CLS} \rightarrow \mathcal{HYP}$ has the property $F \in \mathcal{MU}_{\delta=k} \Leftrightarrow e(F) \in \mathcal{MNC}_{\delta_H=k-1}$ for $k \in \mathbb{N}$, and so the classification of $\mathcal{MU}_{\delta=k}$ can be seen as a subtask of the classification of $\mathcal{MNC}_{\delta_H=k}$ for $k \in \mathbb{N}_0$. The possibility of a characterisation of $\mathcal{MNC}_{\delta_H=0}$ was already raised in [5] (where concentration on the special case of saturated (“strong” there) minimally non-2-colourable hypergraphs was recommended), but is indeed still outstanding, which is understandable, given that polytime decision of $\mathcal{MU}_{\delta=1}$ is easy when compared with polytime decision of $\mathcal{MNC}_{\delta_H=0}$. In the other direction, the consideration of $\mathcal{MU}_{\delta_h=k} \subset \mathcal{MU}_{\delta \leq k}$ (recall the hermitian defect $\delta_h \geq \delta$ from Subsection 1.6.5) could be a stepping stone (recall $\mathcal{MU}_{\delta_h=1} = \mathcal{UHIT}_{\delta=1}$).

A major step towards Conjecture 15.6 should be the classification of unsatisfiable (nonsingular) hitting clause-sets in dependency on the deficiency, i.e., determining the isomorphism types of $\mathcal{UHIT}'_{\delta=k}$. We remark here that unsatisfiable hitting clause-sets do not seem to have a close correspondence in hypergraph colouring, due to the lack of complementation in hypergraphs. Here the main conjecture is (which should be a special case of Conjecture 15.6, once we found the precise formulation of “finitely many patterns”):

Conjecture 15.7 *For every deficiency $k \in \mathbb{N}$ there are only finitely many isomorphism types of nonsingular unsatisfiable hitting clause-sets, or equivalently, the number of variables of elements of $\mathcal{UHIT}'_{\delta=k}$ is bounded.*

For $k \leq 2$ finiteness has been established (Examples 3.2, 3.3), while recently we were able to prove it for $k = 3$ ([90]), exploiting “clause-factorisations”, which generalise

singular DP-reduction and full subsumption resolution, and where the irreducible case corresponds to “irreducible covering systems” as investigated in [106, 107]. Assuming Conjecture 15.7, the question arises about the computability of the function, which maps $k \in \mathbb{N}$ to the set of isomorphism types. Equivalently one can consider the computability of any function, which maps $k \in \mathbb{N}$ to an upper bound on the number of variables of elements of $\mathcal{UHT}'_{\delta=k}$. It is conceivable that such functions must grow so quickly that they are not computable, we however believe that actually a small bound holds, and we conjecture the following strengthened form of Conjecture 15.7:

Conjecture 15.8 *For every $k \geq 2$ holds $\max\{n(F) : F \in \mathcal{UHT}'_{\delta=k}\} = 4k - 5$.*

By Example 3.3 indeed for $k = 2$ the maximal number of variables is $4 \cdot 2 - 5 = 3$, and in the forthcoming [90] we prove Conjecture 15.8 for $k = 3$. We obtain computability of VDM as follows (using Corollary 5.5):

Lemma 15.9 *Assume that Conjecture 15.8 holds.*

1. *Then the map $k \in \mathbb{N} \mapsto \text{VDH}(k)$ is computable, by enumerating all possible clause-sets F with at most $4k - 5$ variables, checking whether they are in $\mathcal{UHT}'_{\delta=k}$, and if so, including $\text{mvd}(F)$ into the maximum-computation.*
2. *If also Conjecture 12.1 holds, then also the function VDM is computable.*

Conjecture 15.1 says additionally, that although VDM is computable, it should be “complex”.

References

- [1] S. A. Cook, The complexity of theorem-proving procedures, in: Proceedings 3rd Annual ACM Symposium on Theory of Computing (STOC '71), 1971, pp. 151–158. doi:10.1145/800157.805047.
- [2] A. Biere, M. J. Heule, H. van Maaren, T. Walsh (Eds.), Handbook of Satisfiability, Vol. 185 of Frontiers in Artificial Intelligence and Applications, IOS Press, 2009.
- [3] M. Davis, H. Putnam, A computing procedure for quantification theory, Journal of the ACM 7 (3) (1960) 201–215. doi:doi.acm.org/10.1145/321033.321034.
- [4] M. Davis, G. Logemann, D. Loveland, A machine program for theorem-proving, Communications of the ACM 5 (7) (1962) 394–397. doi:10.1145/368273.368557.
- [5] R. Aharoni, N. Linial, Minimal non-two-colorable hypergraphs and minimal unsatisfiable formulas, Journal of Combinatorial Theory, Series A 43 (2) (1986) 196–204. doi:10.1016/0097-3165(86)90060-9.
- [6] G. Davydov, I. Davydova, H. Kleine Büning, An efficient algorithm for the minimal unsatisfiability problem for a subclass of CNF, Annals of Mathematics and Artificial Intelligence 23 (3-4) (1998) 229–245. doi:10.1023/A:1018924526592.
- [7] O. Kullmann, An application of matroid theory to the SAT problem, in: Proceedings of the 15th Annual IEEE Conference on Computational Complexity, 2000, pp. 116–124. doi:10.1109/CCC.2000.856741.

- [8] H. Kleine Büning, On subclasses of minimal unsatisfiable formulas, *Discrete Applied Mathematics* 107 (1-3) (2000) 83–98. doi:10.1016/S0166-218X(00)00245-6.
- [9] Z. Xishun, D. Decheng, Two tractable subclasses of minimal unsatisfiable formulas, *Science in China Series A: Mathematics* 42 (7) (1999) 720–731. doi:10.1007/BF02878991.
- [10] O. Kullmann, Investigations on autark assignments, *Discrete Applied Mathematics* 107 (2000) 99–137. doi:10.1016/S0166-218X(00)00262-6.
- [11] H. Kleine Büning, O. Kullmann, Minimal unsatisfiability and autarkies, in: Biere et al. [2], Ch. 11, pp. 339–401. doi:10.3233/978-1-58603-929-5-339.
- [12] L. Lovász, M. D. Plummer, *Matching Theory*, AMS Chelsea Publishing, 2009, ISBN 978-0-8218-4759-6; first published 1986, here with an additional appendix and with errata.
- [13] O. Kullmann, Lean clause-sets: Generalizations of minimally unsatisfiable clause-sets, *Discrete Applied Mathematics* 130 (2003) 209–249. doi:10.1016/S0166-218X(02)00406-7.
- [14] O. Kullmann, X. Zhao, On variables with few occurrences in conjunctive normal forms, in: L. Simon, K. Sakallah (Eds.), *Theory and Applications of Satisfiability Testing - SAT 2011*, Vol. 6695 of *Lecture Notes in Computer Science*, Springer, 2011, pp. 33–46. doi:10.1007/978-3-642-21581-0_5.
- [15] Z. Chen, D. Ding, Variable minimal unsatisfiability, in: J.-Y. Cai, S. B. Cooper, A. Li (Eds.), *Theory and Applications of Models of Computation*, Vol. 3959 of *Lecture Notes in Computer Science*, Springer, 2006, pp. 262–273. doi:10.1007/11750321_25.
- [16] O. Kullmann, Constraint satisfaction problems in clausal form I: Autarkies and deficiency, *Fundamenta Informaticae* 109 (1) (2011) 27–81. doi:10.3233/FI-2011-428.
- [17] O. Kullmann, Constraint satisfaction problems in clausal form II: Minimal unsatisfiability and conflict structure, *Fundamenta Informaticae* 109 (1) (2011) 83–119. doi:10.3233/FI-2011-429.
- [18] O. Kullmann, X. Zhao, On Davis-Putnam reductions for minimally unsatisfiable clause-sets, in: Cimatti and Sebastiani [108], pp. 270–283. doi:10.1007/978-3-642-31612-8_21.
- [19] O. Kullmann, X. Zhao, On Davis-Putnam reductions for minimally unsatisfiable clause-sets, *Theoretical Computer Science* 492 (2013) 70–87. doi:10.1016/j.tcs.2013.04.020.
- [20] C. H. Papadimitriou, M. Yannakakis, The complexity of facets (and some facets of complexity), *Journal on Computer and System Sciences* 28 (2) (1984) 244–259. doi:10.1016/0022-0000(84)90068-0.
- [21] C. H. Papadimitriou, D. Wolfe, The complexity of facets resolved, *Journal of Computer and System Sciences* 37 (1) (1988) 2–13. doi:10.1016/0022-0000(88)90042-6.
- [22] J. Franco, A. V. Gelder, A perspective on certain polynomial-time solvable classes of satisfiability, *Discrete Applied Mathematics* 125 (2-3) (2003) 177–214. doi:10.1016/S0166-218X(01)00358-4.

- [23] T. Fliti, G. Reynaud, Sizes of minimally unsatisfiable conjunctive normal forms, faculté des Sciences de Luminy, Dpt. Mathématique-Informatique, 13288 Marseille, France (November 1994).
- [24] H. Kleine Büning, An upper bound for minimal resolution refutations, in: G. Gottlob, E. Grandjean, K. Seyr (Eds.), Computer Science Logic 12th International Workshop, CSL'98, Vol. 1584 of Lecture Notes in Computer Science, Springer, 1999, pp. 171–178. doi:10.1007/10703163_12.
- [25] H. Fleischner, O. Kullmann, S. Szeider, Polynomial-time recognition of minimal unsatisfiable formulas with fixed clause-variable difference, Theoretical Computer Science 289 (1) (2002) 503–516. doi:10.1016/S0304-3975(01)00337-1.
- [26] S. Szeider, Minimal unsatisfiable formulas with bounded clause-variable difference are fixed-parameter tractable, Journal of Computer and System Sciences 69 (4) (2004) 656–674. doi:10.1016/j.jcss.2004.04.009.
- [27] J. Marques-Silva, Computing minimally unsatisfiable subformulas: State of the art and future directions, Journal of Multiple-Valued Logic and Soft Computing 19 (1-3) (2012) 163–183.
URL <http://www.oldcitypublishing.com/journals/mvlsc-home/mvlsc-issue-contents/mvlsc-volume>
- [28] M. H. Liffiton, K. A. Sakallah, On finding all minimally unsatisfiable subformulas, in: Bacchus and Walsh [109], pp. 173–186. doi:10.1007/11499107_13.
- [29] R. Bruni, Approximating minimal unsatisfiable subformulae by means of adaptive core search, Discrete Applied Mathematics 130 (2) (2003) 85–100. doi:10.1016/S0166-218X(02)00399-2.
- [30] O. Kullmann, I. Lynce, J. Marques-Silva, Categorisation of clauses in conjunctive normal forms: Minimally unsatisfiable sub-clause-sets and the lean kernel, in: A. Biere, C. P. Gomes (Eds.), Theory and Applications of Satisfiability Testing - SAT 2006, Vol. 4121 of Lecture Notes in Computer Science, Springer, 2006, pp. 22–35. doi:10.1007/11814948_4.
- [31] C. A. Tovey, A simplified NP-complete satisfiability problem, Discrete Applied Mathematics 8 (1) (1984) 85–89. doi:10.1016/0166-218X(84)90081-7.
- [32] O. Dubois, On the r, s -satisfiability problem and a conjecture of Tovey, Discrete Applied Mathematics 26 (1) (1990) 51–60. doi:10.1016/0166-218X(90)90020-D.
- [33] J. Kratochvíl, P. Savický, Z. Tuza, One more occurrence of variables makes satisfiability jump from trivial to NP-complete, SIAM Journal on Computing 22 (1) (1993) 203–210. doi:10.1137/0222015.
- [34] S. Hoory, S. Szeider, Computing unsatisfiable k -SAT instances with few occurrences per variable, Theoretical Computer Science 337 (1-3) (2005) 347–359. doi:10.1016/j.tcs.2005.02.004.
- [35] H. Gebauer, T. Szabo, G. Tardos, The local lemma is tight for SAT, Tech. Rep. arXiv:1006.0744v2 [math.CO], arXiv.org (August 2013).
URL <http://arxiv.org/abs/1006.0744>
- [36] S. Porschen, E. Speckenmeyer, X. Zhao, Linear CNF formulas and satisfiability, Discrete Applied Mathematics 157 (5) (2009) 1046–1068. doi:10.1016/j.dam.2008.03.031.

- [37] D. Scheder, Unsatisfiable linear CNF formulas are large and complex, in: J.-Y. Marion, T. Schwentick (Eds.), 27th International Symposium on Theoretical Aspects of Computer Science (STACS 2010), 2010, pp. 621–632. doi:10.4230/LIPIcs.STACS.2010.2490.
- [38] C. Lee, On the size of minimal unsatisfiable formulas, The Electronic Journal of Combinatorics 16 (1), note #N3.
URL http://www.combinatorics.org/Volume_16/Abstracts/v16i1n3.html
- [39] D. Scheder, Unsatisfiable CNF formulas contain many conflicts, in: L. Cai, S.-W. Cheng, T.-W. Lam (Eds.), Algorithms and Computation — 24th International Symposium, ISAAC 2013, Vol. 8283 of Lecture Notes in Computer Science, Springer, 2013, pp. 273–283. doi:10.1007/978-3-642-45030-3_26.
- [40] H. Gebauer, R. A. Moser, D. Scheder, E. Welzl, The Lovász local lemma and satisfiability, in: S. Albers, H. Alt, S. Näher (Eds.), Efficient Algorithms — Essays Dedicated to Kurt Mehlhorn on the Occasion of His 60th Birthday, Vol. 5760 of Lecture Notes in Computer Science, Springer, 2009, pp. 30–54. doi:10.1007/978-3-642-03456-5_3.
- [41] B. Monien, E. Speckenmeyer, Solving satisfiability in less than 2^n steps, Discrete Applied Mathematics 10 (3) (1985) 287–295. doi:10.1016/0166-218X(85)90050-2.
- [42] J. Marques-Silva, A. Ignatiev, A. Morgado, V. Manquinho, I. Lynce, Efficient autarkies, in: T. Schaub, G. Friedrich, B. O’Sullivan (Eds.), 21st European Conference on Artificial Intelligence (ECAI 2014), Vol. 263 of Frontiers in Artificial Intelligence and Applications, IOS Press, 2014, pp. 603–608. doi:10.3233/978-1-61499-419-0-603.
- [43] O. Kullmann, J. Marques-Silva, Computing maximal autarkies with few and simple oracle queries, in: M. H. Heule, S. Weaver (Eds.), Theory and Applications of Satisfiability Testing - SAT 2015, Vol. 9340 of Lecture Notes in Computer Science, Springer, 2015, pp. 138–155. doi:10.1007/978-3-319-24318-4_11.
- [44] O. Kullmann, Polynomial time SAT decision for complementation-invariant clause-sets, and sign-non-singular matrices, in: J. Marques-Silva, K. A. Sakallah (Eds.), Theory and Applications of Satisfiability Testing - SAT 2007, Vol. 4501 of Lecture Notes in Computer Science, Springer, 2007, pp. 314–327. doi:10.1007/978-3-540-72788-0_30.
- [45] D. Achlioptas, Random satisfiability, in: Biere et al. [2], Ch. 8, pp. 245–270. doi:10.3233/978-1-58603-929-5-245.
- [46] K. A. Sakallah, Symmetry and satisfiability, in: Biere et al. [2], Ch. 10, pp. 289–338. doi:10.3233/978-1-58603-929-5-289.
- [47] M. Samer, S. Szeider, Fixed-parameter tractability, in: Biere et al. [2], Ch. 13, pp. 425–454. doi:10.3233/978-1-58603-929-5-425.
- [48] H. Zhang, Combinatorial designs by SAT solvers, in: Biere et al. [2], Ch. 17, pp. 533–568. doi:10.3233/978-1-58603-929-5-533.
- [49] T. Ahmed, O. Kullmann, H. Snevily, On the van der Waerden numbers $w(2; 3, t)$, Discrete Applied Mathematics 174 (2014) 27–51. doi:10.1016/j.dam.2014.05.007.

- [50] M. J. Heule, O. Kullmann, V. W. Marek, Solving and verifying the boolean Pythagorean Triples problem via Cube-and-Conquer, in: N. Creignou, D. L. Berre (Eds.), Theory and Applications of Satisfiability Testing - SAT 2016, Vol. ??? of Lecture Notes in Computer Science, Springer, 2016, p. ???
doi:???
- [51] P. Duchet, Hypergraphs, in: R. L. Graham, M. Grötschel, L. Lovász (Eds.), Handbook of Combinatorics, Volume 1, North-Holland, Amsterdam, 1995, Ch. 7, pp. 381–432, ISBN 0-444-82346-8.
- [52] T. R. Jensen, B. Toft, Graph Coloring Problems, Wiley-Interscience Series in Discrete Mathematics and Optimization, John Wiley & Sons, 1995, ISBN 0-471-02865-7, QA612.18J46; problem status updated at <http://www.imada.sdu.dk/Research/Graphcol>.
doi:10.1002/9781118032497.
- [53] A. Kostochka, Color-critical graphs and hypergraphs with few edges: A survey, in: E. Györi, G. O. Katona, L. Lovász (Eds.), More Sets, Graphs and Numbers, Vol. 15 of Bolyai Society Mathematical Studies, Springer, 2006, pp. 175–197. doi:10.1007/978-3-540-32439-3_9.
- [54] P. D. Seymour, On the two-colouring of hypergraphs, The Quarterly Journal of Mathematics (Oxford University Press) 25 (1974) 303–312.
doi:10.1093/qmath/25.1.303.
- [55] O. Kullmann, Constraint satisfaction problems in clausal form: Autarkies, minimal unsatisfiability, and applications to hypergraph inequalities, in: N. Creignou, P. Kolaitis, H. Vollmer (Eds.), Complexity of Constraints, no. 06401 in Dagstuhl Seminar Proceedings, Internationales Begegnungs- und Forschungszentrum fuer Informatik (IBFI), Schloss Dagstuhl, Germany, 2006, pp. 1–69.
URL <http://drops.dagstuhl.de/opus/volltexte/2006/803>
- [56] N. Robertson, P. D. Seymour, R. Thomas, Permanents, Pfaffian orientations, and even directed circuits, Annals of Mathematics 150 (3) (1999) 929–975.
doi:10.2307/121059.
- [57] W. McCuaig, Pólya’s permanent problem, The Electronic Journal of Combinatorics 11, #R79, 83 pages.
URL <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v11i1r79>
- [58] H. Abbott, D. Hare, Square critically 3-chromatic hypergraphs, Discrete Mathematics 197-198 (1999) 3–13. doi:10.1016/S0012-365X(99)90031-6.
- [59] N. Linial, M. Tarsi, Deciding hypergraph 2-colourability by H-resolution, Theoretical Computer Science 38 (1985) 343–347.
doi:10.1016/0304-3975(85)90227-0.
- [60] T. Eiter, K. Makino, G. Gottlob, Computational aspects of monotone dualization: A brief survey, Discrete Applied Mathematics 156 (11) (2008) 2035–2049. doi:10.1016/j.dam.2007.04.017.
- [61] N. Galesi, O. Kullmann, Polynomial time SAT decision, hypergraph transversals and the hermitian rank, in: H. H. Hoos, D. G. Mitchell (Eds.), Theory and Applications of Satisfiability Testing 2004, Vol. 3542 of Lecture Notes in Computer Science, Springer, Berlin, 2005, pp. 89–104. doi:10.1007/11527695_8.

- [62] S. Porschen, B. Randerath, E. Speckenmeyer, Linear time algorithms for some not-all-equal satisfiability problems, in: Giunchiglia and Tacchella [110], pp. 172–187. doi:10.1007/978-3-540-24605-3_14.
- [63] C. D. Godsil, Tools from linear algebra, in: R. L. Graham, M. Grötschel, L. Lovász (Eds.), Handbook of Combinatorics, Volume 2, North-Holland, Amsterdam, 1995, Ch. 31, pp. 1705–1748, iISBN 0-444-82351-4.
- [64] R. A. Brualdi, B. L. Shader, Matrices of sign-solvable linear systems, Vol. 116 of Cambridge Tracts in Mathematics, Cambridge University Press, 1995, iISBN 0-521-48296-8. doi:10.1017/CB09780511574733.
- [65] F. J. Hall, Z. Li, Sign pattern matrices, in: L. Hogben (Ed.), Handbook of Linear Algebra, Discrete Mathematics and Its Applications, Chapman & Hall/CRC, 2007, pp. 33(1)–33(21), iISBN 1-58488-510-6. doi:10.1201/9781420010572.ch33.
- [66] V. Klee, R. Ladner, Qualitative matrices: Strong sign-solvability and weak satisfiability, in: H. J. Greenberg, J. S. Maybee (Eds.), Computer-Assisted Analysis and Model Simplification, 1981, pp. 293–320, proceedings of the First Symposium on Computer-Assisted Analysis and Model Simplification, University of Colorado, Boulder, Colorado, March 28, 1980. doi:10.1016/B978-0-12-299680-1.50022-7.
- [67] G. Davydov, I. Davydova, Tautologies and positive solvability of linear homogeneous systems, Annals of Pure and Applied Logic 57 (1) (1992) 27–43. doi:10.1016/0168-0072(92)90060-D.
- [68] G.-Y. Lee, B. L. Shader, Sign-consistency and solvability of constrained linear systems, The Electronic Journal of Linear Algebra 4 (1998) 1–18.
URL <http://emis.matem.unam.mx/journals/ELA/ela-articles/4.html>
- [69] R. L. Graham, H. Pollak, On the addressing problem for loop switching, Bell System Technical Journal 50 (8) (1971) 2495–2519.
- [70] J. H. van Lint, R. M. Wilson, A Course in Combinatorics, 2nd Edition, Cambridge University Press, 2001, iISBN 0 521 00601 5; fifth printing with corrections 2006.
- [71] S. Vishwanathan, A counting proof of the Graham-Pollak theorem, Discrete Mathematics 313 (6) (2013) 765–766. doi:10.1016/j.disc.2012.12.017.
- [72] P. M. Winkler, Proof of the squashed cube conjecture, Combinatorica 3 (1) (1983) 135–139. doi:10.1007/BF02579350.
- [73] O. Kullmann, The combinatorics of conflicts between clauses, in: Giunchiglia and Tacchella [110], pp. 426–440, this is a slightly shorted version of [111]. doi:10.1007/978-3-540-24605-3_32.
- [74] O. Kullmann, On the conflict matrix of clause-sets, Tech. Rep. CSR 7-2003, University of Wales Swansea, Computer Science Report Series (March 2003).
URL <http://www.cs.swansea.ac.uk/reports/yr2003/CSR7-2003.pdf>
- [75] D. A. Gregory, V. L. Watts, B. L. Shader, Biclique decompositions and hermitian rank, Linear Algebra and its Applications 292 (1999) 267–280. doi:10.1016/S0024-3795(99)00042-7.

- [76] C. Godsil, G. Royle, Algebraic Graph Theory, Vol. 207 of Graduate Texts in Mathematics, Springer, New York, 2001, ISBN 0-387-95220-9, QA166 .G63 2001.
- [77] R. H. Sloan, B. Sörényi, G. Turán, On k -term DNF with the largest number of prime implicants, *SIAM Journal on Discrete Mathematics* 21 (4) (2007) 987–998. doi:10.1137/050632026.
- [78] O. Kullmann, New methods for 3-SAT decision and worst-case analysis, *Theoretical Computer Science* 223 (1-2) (1999) 1–72. doi:10.1016/S0304-3975(98)00017-6.
- [79] T. Eiter, Exact transversal hypergraphs and application to boolean μ -functions, *Journal of Symbolic Computation* 17 (3) (1994) 215–225. doi:10.1006/jscs.1994.1013.
- [80] D. A. Gregory, K. N. V. Meulen, Sharp bounds for decompositions of graphs into complete r -partite subgraphs, *Journal of Graph Theory* 21 (4) (1996) 393–400. doi:10.1002/(SICI)1097-0118(199604)21:4<393::AID-JGT4>3.3.CO;2-9.
- [81] O. Kullmann, X. Zhao, Sharper bounds on minimum variable-degrees for minimally unsatisfiable clause-sets, in preparation (April 2016).
- [82] O. Kullmann, X. Zhao, Parameters for minimal unsatisfiability: Smarandache primitive numbers and full clauses, Tech. Rep. arXiv:1505.02318v2 [cs.DM], arXiv (July 2015).
URL <http://arxiv.org/abs/1505.02318>
- [83] Y. Crama, P. L. Hammer, Boolean Functions: Theory, Algorithms, and Applications, Vol. 142 of Encyclopedia of Mathematics and Its Applications, Cambridge University Press, 2011, ISBN 978-0-521-84751-3.
- [84] A. Blake, Canonical expressions in boolean algebra, Ph.D. thesis, Chicago, see [85] (1937).
- [85] J. C. C. McKinsey, Archie Blake: Canonical expressions in Boolean algebra, Review, *The Journal of Symbolic Logic* (3) (1938).
- [86] H. Kleine Büning, X. Zhao, The complexity of some subclasses of minimal unsatisfiable formulas, *Journal on Satisfiability, Boolean Modeling and Computation* 3 (2007) 1–17.
URL <http://satassociation.org/jsat/index.php/jsat/article/view/30>
- [87] H. Kleine Büning, X. Zhao, On the structure of some classes of minimal unsatisfiable formulas, *Discrete Applied Mathematics* 130 (2) (2003) 185–207. doi:10.1016/S0166-218X(02)00405-5.
- [88] O. Kullmann, X. Zhao, On variables with few occurrences in conjunctive normal forms, Tech. Rep. arXiv:1010.5756v3 [cs.DM], arXiv (March 2011).
URL <http://arxiv.org/abs/1010.5756>
- [89] A. Belov, A. Ivrii, A. Matsliah, J. Marques-Silva, On efficient computation of Variable MUSes, in: Cimatti and Sebastiani [108], pp. 298–311. doi:10.1007/978-3-642-31612-8_23.
- [90] O. Kullmann, X. Zhao, Unsatisfiable hitting clause-sets with three more clauses than variables, Tech. Rep. arXiv:1604.01288v1 [cs.DM], arXiv (April 2016).
URL <http://arxiv.org/abs/1604.01288>

- [91] J. Robinson, A machine-oriented logic based on the resolution principle, *Journal of the ACM* 12 (1) (1965) 23–41. doi:10.1145/321250.321253.
- [92] A. V. Gelder, Y. K. Tsuji, Satisfiability testing with more reasoning and less guessing, in: D. S. Johnson, M. A. Trick (Eds.), *Cliques, Coloring, and Satisfiability*, Vol. 26 of DIMACS Series in Discrete Mathematics and Theoretical Computer Science, American Mathematical Society, 1996, pp. 559–586, the Second DIMACS Challenge.
- [93] O. Kullmann, H. Luckhardt, Deciding propositional tautologies: Algorithms and their complexity, preprint, 82 pages (January 1997).
URL <http://cs.swan.ac.uk/~csoliver/papers.html#Pre199799>
- [94] O. Kullmann, H. Luckhardt, Algorithms for SAT/TAUT decision based on various measures, preprint, 71 pages (February 1999).
URL <http://cs.swan.ac.uk/~csoliver/papers.html#Pre199799>
- [95] N. Eén, A. Biere, Effective preprocessing in SAT through variable and clause elimination, in: *Bacchus and Walsh* [109], pp. 61–75. doi:10.1007/11499107_5.
- [96] H. Han, F. Somenzi, On-the-fly clause improvement, in: O. Kullmann (Ed.), *Theory and Applications of Satisfiability Testing 2009*, Vol. 5584 of *Lecture Notes in Computer Science*, Springer, 2009, pp. 209–222. doi:10.1007/978-3-642-02777-2_21.
- [97] S. Ordyniak, D. Paulusma, S. Szeider, Satisfiability of acyclic and almost acyclic CNF formulas, *Theoretical Computer Science* 481 (2013) 85–99. doi:10.1016/j.tcs.2012.12.039.
- [98] N. J. Sloane, *The On-Line Encyclopedia of Integer Sequences (OEIS)* (2008).
URL <http://oeis.org/>
- [99] F. Smarandache, *Only problems, not solutions!*, 4th Edition, Xiquan Publishing House, 1993.
URL <http://www.gallup.unm.edu/~smarandache/OPNS.pdf>
- [100] D. R. Hofstadter, *Gödel, Escher, Bach: An eternal golden braid*, Basic Books, 1979, pdf version with 801 pages, md5sum="0cb32e8ea5dd2485f63842f5acffb3f0 GEBen.pdf".
URL <http://www.physixfan.com/wp-content/files/GEBen.pdf>
- [101] B. Conolly, Meta-Fibonacci sequences (a letter from B.W. Conolly), in: *Fibonacci and Lucas Numbers and the Golden Section*, Dover Publications, Inc., 2008, Ch. 12, pp. 127–138, unabridged republication of the work originally published in 1989 by Ellis Horwood Limited, Chichester, England.
- [102] Z. Wenpeng, L. Duansen, On the primitive numbers of power P and its asymptotic property, in: J. Allen, F. Liu, D. Constantinescu (Eds.), *Smarandache Notions*, American Research Press, Rehoboth, NM, USA, 2002, pp. 173–175.
URL <http://dl.acm.org/citation.cfm?id=773309>
- [103] B. Beckert, R. Hähnle, F. Manyà, The SAT problem of signed CNF formulas, in: D. Basin, M. D’Agostino, D. Gabbay, S. Matthews, L. Viganò (Eds.), *Labelled Deduction*, Vol. 17 of *Applied Logic Series*, Springer Netherlands, 2000, pp. 59–80. doi:10.1007/978-94-011-4040-9_3.

- [104] O. Kullmann, X. Zhao, Some connections of elementary number theory with the combinatorics of multivalued propositional logic, in preparation (April 2016).
- [105] H. Ibstedt, *Computer Analysis of Number Sequences*, American Research Press, 1998.
URL <http://www.gallup.unm.edu/~smarandache/Ibstedt-computer.pdf>
- [106] I. Korec, Irreducible disjoint covering systems, *Acta Arithmetica* 44 (4) (1984) 389–395.
URL <http://pldml.icm.edu.pl/pldml/element/bwmeta1.element.bwnjournal-article-aav44i4p389bw>
- [107] M. A. Berger, A. Felzenbaum, A. S. Fraenkel, Irreducible disjoint covering systems (with an application to boolean algebra), *Discrete Applied Mathematics* 29 (2-3) (1990) 143–164. doi:10.1016/0166-218X(90)90140-8.
- [108] A. Cimatti, R. Sebastiani (Eds.), *Theory and Applications of Satisfiability Testing - SAT 2012*, Vol. 7317 of *Lecture Notes in Computer Science*, Springer, 2012. doi:10.1007/978-3-642-31612-8.
- [109] F. Bacchus, T. Walsh (Eds.), *Theory and Applications of Satisfiability Testing 2005*, Vol. 3569 of *Lecture Notes in Computer Science*, Springer, Berlin, 2005. doi:10.1007/b137280.
- [110] E. Giunchiglia, A. Tacchella (Eds.), *Theory and Applications of Satisfiability Testing 2003*, Vol. 2919 of *Lecture Notes in Computer Science*, Springer, Berlin, 2004. doi:10.1007/b95238.
- [111] O. Kullmann, On some connections between linear algebra and the combinatorics of clause-sets, in: J. Franco, E. Giunchiglia, H. Kautz, H. Kleine Büning, H. van Maaren, B. Selman, E. Speckenmeyer (Eds.), *Sixth International Conference on Theory and Applications of Satisfiability Testing*, Santa Margherita Ligure – Portofino (Italy), 2003, pp. 45–59.