

Variations on an error sum function for the convergents of some powers of e

J.-P. Allouche*

CNRS, Institut de Mathématiques de Jussieu-PRG
Équipe Combinatoire et Optimisation
Université Pierre et Marie Curie, Case 247
4 Place Jussieu
F-75252 Paris Cedex 05 France
jean-paul.allouche@imj-prg.fr

Thomas Baruchel
Lycée naval
Avenue de l'École navale
F-29240 Brest Armées, France
baruchel@riseup.net

Abstract

Several years ago the second author playing with different “recognizers of real constants”, e.g., the LLL algorithm, the Plouffe inverter, etc. found the following formula empirically. Let p_n/q_n denote the n th convergent of the continued fraction of the constant e . Then

$$\sum_{n \geq 0} |q_n e - p_n| = \frac{e}{4} \left(-1 + 10 \sum_{n \geq 0} \frac{(-1)^n}{(n+1)!(2n^2 + 7n + 3)} \right).$$

The purpose of the present paper is to prove this formula and to give similar formulas for some powers of e .

Keywords: Continued fractions; convergents; approximation of real numbers; Hurwitzian continued fractions; error sum function.

MSC Classes: 11A55, 11J70, 11B83, 11B75.

*The author was partially supported by the ANR project “FAN” (Fractals et Numération), ANR-12-IS01-0002.

1 Introduction

Playing with the convergents of e , the second author discovered several years ago the formula

$$\sum_{n \geq 0} |q_n e - p_n| = \frac{e}{4} \left(-1 + 10 \sum_{n \geq 0} \frac{(-1)^n}{(n+1)!(2n^2 + 7n + 3)} \right). \quad (1)$$

While trying to prove the formula rigorously we began being interested in the following quantity. If α is a positive real number, and if p_n/q_n is the n th convergent of its continued fraction, the quantity $|q_n \alpha - p_n|$ tends rapidly to zero. Thus the series $\sum_{n \geq 0} |q_n \alpha - p_n|$ converges. This series measures in some sense the “global approximation” of α by its convergents. We then learned from J. Shallit that the quantity $\sum_{n \geq 0} |q_n \alpha - p_n| = \sum_{n \geq 0} (-1)^n (q_n \alpha - p_n)$ was investigated in several papers [5, 8, 9, 6], where the study of the quantity $\sum_{n \geq 0} (q_n \alpha - p_n)$ (first defined in [28]) can also be found.

It is natural to ask whether the sum of the series $\sum_{n \geq 0} |q_n \alpha - p_n|$ can be expressed in terms of α without explicitly using the convergents, in particular in the case where α has a “nice” continued fraction expansion, e.g., when α is quadratic or when $\alpha = e$.

2 Quadratic numbers

The case of quadratic numbers was addressed in [5] (also see [6]).

Theorem 1 (Elsner) *Let p_n/q_n be the n th convergent of the continued fraction of α . Then the series $\sum_{n \geq 0} (q_n \alpha - p_n) x^n$ converges absolutely at least for $|x| < \frac{1+\sqrt{5}}{2}$ and*

$$\sum_{n \geq 0} (q_n \alpha - p_n) x^n \in \mathbb{Q}[\alpha](x).$$

In particular (taking $x = -1$), $\sum_{n \geq 0} |q_n \alpha - p_n|$ belongs to $\mathbb{Q}[\alpha]$.

Example 1 (Elsner)

- $\sum_{n \geq 0} |q_n \sqrt{7} - p_n| = \frac{7+5\sqrt{7}}{14}$.
- For any integer $n \geq 1$ we have $\sum_{n \geq 0} |q_n (\frac{n+\sqrt{4+n^2}}{2}) - p_n| = \frac{1}{\frac{n+\sqrt{4+n^2}}{2} - 1}$.
- In particular $\sum_{n \geq 0} |q_n (\frac{1+\sqrt{5}}{2}) - p_n| = \frac{1+\sqrt{5}}{2}$.

3 Powers of e

Euler [10] proved that the continued fraction expansion of e is $[2, 1, 2, 1, 1, 4, 1, 1, 6, 1, 1, 8, \dots]$ (sometimes replaced by the not really regular expression $[1, 0, 1, 1, 2, 1, 1, 4, 1, 1, 6, 1, 1, 8, \dots]$). After Euler, a large number of papers contained the computation of continued fraction

expansions for some expressions containing e (typically certain powers of e possibly multiplied by some rational numbers, or numbers like $\frac{e^{2/k}-1}{e^{2/k}+1}$), see in particular [11, 19, 13, 20, 4, 24, 23, 27, 3, 26, 14, 22, 15, 17, 16, 21, 18, 12].

The fundamental theorem we will use here is due to Komatsu [17, Theorem 6, first part]. Komatsu's theorem contains several previous results.

Theorem 2 (Komatsu) *Let $\ell \geq 2$ and $s \geq 1$ be two integers. Let p_n/q_n be the n th convergent of the continued fraction of*

$$se^{1/(\ell s)} = [s, \ell - 1, 1, 2s - 1, 3\ell - 1, 1, 2s - 1, 5\ell - 1, 1, 2s - 1, \dots, (2k - 1)\ell - 1, 1, 2s - 1, \dots].$$

Then for $n \geq 0$

$$p_{3n} - se^{1/(\ell s)}q_{3n} = -\frac{1}{(\ell s)^{n+1}} \int_0^1 \frac{x^n(x-1)^n}{n!} se^{x/(\ell s)} dx$$

$$p_{3n+1} - se^{1/(\ell s)}q_{3n+1} = \frac{1}{s(\ell s)^{n+1}} \int_0^1 \frac{(x+s-1)x^n(x-1)^n}{n!} se^{x/(\ell s)} dx$$

$$p_{3n+2} - se^{1/(\ell s)}q_{3n+2} = \frac{1}{s(\ell s)^{n+1}} \int_0^1 \frac{x^n(x-1)^{n+1}}{n!} se^{x/(\ell s)} dx$$

Let $s \geq 1$ be an integer. Let p_n^*/q_n^* be the n th convergent of the continued fraction of

$$se^{1/s} = [s + 1, 2s - 1, 2, 1, 2s - 1, 4, 1, \dots, 2s - 1, 2k, 1, \dots].$$

Then $p_n^*/q_n^* = p_{n+2}/q_{n+2}$ with p_n/q_n is as above. More precisely for $n \geq 0$

$$p_{3n}^* - se^{1/s}q_{3n}^* = \frac{1}{s^{n+2}} \int_0^1 \frac{x^n(x-1)^{n+1}}{n!} se^{x/s} dx$$

$$p_{3n+1}^* - se^{1/s}q_{3n+1}^* = -\frac{1}{s^{n+2}} \int_0^1 \frac{x^{n+1}(x-1)^{n+1}}{(n+1)!} se^{x/s} dx$$

$$p_{3n+2}^* - se^{1/s}q_{3n+2}^* = \frac{1}{s^{n+3}} \int_0^1 \frac{(x+s-1)x^{n+1}(x-1)^{n+1}}{(n+1)!} se^{x/s} dx$$

Using Komatsu's result we can prove the following theorem. First recall that the "error function" erf is defined by $\text{erf}(x) := \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$.

Note The name "error sum function" (or "error-sum function") that goes back at least to [28] should not be confused with the name "error function". To (try to) avoid any ambiguity, we will always write for the latter "error function erf".

Theorem 3 Let $\ell \geq 2$ and $s \geq 1$ be two integers. Let p_n/q_n be the n th convergent of the continued fraction of

$$se^{1/(\ell s)} = [s, \ell - 1, 1, 2s - 1, 3\ell - 1, 1, 2s - 1, 5\ell - 1, 1, 2s - 1, \dots, (2k - 1)\ell - 1, 1, 2s - 1, \dots].$$

Then

$$\sum_{n \geq 0} |p_n - se^{1/(\ell s)} q_n| = e^{1/\ell s} \sqrt{\frac{\pi s}{\ell}} \operatorname{erf}(1/\sqrt{\ell s}).$$

Let $s \geq 1$ be an integer. Let p_n^*/q_n^* be the n th convergent of the continued fraction of

$$se^{1/s} = [s + 1, 2s - 1, 2, 1, 2s - 1, 4, 1, \dots, 2s - 1, 2k, 1, \dots].$$

Then

$$\sum_{n \geq 0} |p_n^* - se^{1/s} q_n^*| = e^{1/s} \sqrt{\pi s} \operatorname{erf}(1/\sqrt{s}) + s(1 - e^{1/s}) - 1.$$

Proof. It suffices to use Komatsu's theorem (Theorem 2 above) after writing

$$\sum_{n \geq 0} |p_n - se^{1/\ell s} q_n| = \sum_{0 \leq j \leq 2} \sum_{n \geq 0} |p_{3n+j} - se^{1/\ell s} q_{3n+j}|$$

respectively

$$\sum_{n \geq 0} |p_n^* - se^{1/s} q_n^*| = |p_0^* - se^{1/s} q_0| + \sum_{1 \leq j \leq 3} \sum_{n \geq 0} |p_{3n+j}^* - se^{1/s} q_{3n+j}^*|. \quad \square$$

We deduce the following corollary.

Corollary 1 Let p_n^*/q_n^* be the n th convergent of the continued fraction of e (recall that $e = [2, 1, 2, 1, 1, 4, 1, \dots, 1, 2n, 1, \dots]$). Then

$$\sum_{n \geq 0} |p_n^* - eq_n^*| = 2e \int_0^1 e^{-t^2} dt - e = e\sqrt{\pi} \operatorname{erf}(1) - e.$$

Let p_n/q_n be the n th convergent of the continued fraction of $e^{1/\ell}$ (with $\ell \geq 2$). Then

$$\sum_{n \geq 0} |p_n - e^{1/\ell} q_n| = e^{1/\ell} \sqrt{\frac{\pi}{\ell}} \operatorname{erf}(1/\sqrt{\ell}).$$

Remark 1 The first result in Corollary 1 above was already obtained by Elsner in [5, p. 2].

Now we prove Formula (1).

Corollary 2 Let $A(\ell, s)$ be defined for positive reals ℓ and s by

$$A(\ell, s) := \sum_{n \geq 0} \frac{(-1)^n}{(n+1)!(2n^2 + 7n + 3)(\ell s)^n}.$$

Then

$$A(\ell, s) = -\frac{3}{10}\ell s + \frac{1}{5}\ell s(2 - \ell s - \ell^2 s^2)e^{-1/\ell s} + \frac{4}{5} \int_0^1 e^{-t^2/\ell s} dt$$

In particular

$$A(1, 1) = -\frac{3}{10} + \frac{4}{5} \int_0^1 e^{-t^2} dt.$$

so that

$$\sum_{n \geq 0} |q_n e - p_n| = 2 \int_0^1 e^{-t^2} dt - e = \frac{e}{4}(-1 + 10A(1, 1))$$

Proof. The proof is easy. First write

$$\frac{1}{2n^2 + 7n + 3} = \frac{2}{5(2n + 1)} - \frac{1}{5(n + 3)}.$$

Then introduce the series

$$\sum_{n \geq 0} \frac{(-1)^{n+1} x^{2n+1}}{(n+1)!(\ell s)^{n+1}(2n+1)} \quad \text{and} \quad \sum_{n \geq 0} \frac{(-1)^{n+1} x^{n+3}}{(n+1)!(\ell s)^{n+1}(n+3)}.$$

The derivative of these series are easily computed. We then need their values at $x = 1$. \square

Remark 2 It is immediate to use the values of $A(\ell, s)$ to obtain similar formulas for $se^{\ell s}$. We also note we first thought that the quantity $(2n^2 + 7n + 3)$ was somehow crucial in Formula (1): there might ever have been (though it would have been quite surprising) a link with the number of independent parameters of the orthosymplectic group $\text{OSP}(3, 2n)$ which is precisely $(2n^2 + 7n + 3)$ (see, e.g., [2, p. 223]). But this quantity is not crucial; compare with Formula (2) given below which can be proved by using a step of the proof of Corollary 2 above with $s = \ell = 1$ and $x = 1$:

$$\sum_{n \geq 0} \frac{(-1)^{n+1}}{(n+1)!(2n+1)} = 1 - e^{-1} - 2 \int_0^1 e^{-t^2} dt.$$

This implies

$$\sum_{n \geq 0} |p_n^* - eq_n^*| = e \sum_{n \geq 0} \frac{(-1)^n}{(n+1)!(2n+1)} - 1. \quad (2)$$

Remark 3 The value of $\sum_{n \geq 0} |p_n - \alpha q_n|$ for α equal to one of the above real numbers can also be expressed as another kind of series. Namely a classical series for the error function erf (see, e.g., [1, 7.1.6, p. 297] reads

$$\text{erf}(z) = \frac{2}{\sqrt{\pi}} e^{-z^2} \sum_{n \geq 0} \frac{2^n}{1 \times 3 \times 5 \cdots \times (2n+1)} z^{2n+1}.$$

Using Corollary 1 and the notation therein, this gives in particular the following formulas

$$\sum_{n \geq 0} |p_n^* - e q_n^*| = \sum_{n \geq 0} \frac{2^{n+1}}{1 \times 3 \times 5 \cdots \times (2n+1)} - e = \sum_{n \geq 0} \frac{2^{2n+1} n!}{(2n+1)!} - e \quad (3)$$

and, for any integer $\ell \geq 2$,

$$\sum_{n \geq 0} |p_n - e^{1/\ell} q_n| = \sum_{n \geq 0} \frac{2^{n+1}}{\ell^{n+1} (1 \times 3 \times 5 \cdots \times (2n+1))} = \sum_{n \geq 0} \frac{2^{2n+1} n!}{\ell^{n+1} (2n+1)!}. \quad (4)$$

Note that the second author obtained Formula (4) empirically. Also note that the digits of the decimal expansion of the right side of Equation (3) (up to the $-e$ term) is given in [25] as A125961, and that the expansions of the right side of Equation (4) above for $\ell = 2$ and $\ell = 4$ are given in [25] as A060196 and A214869 respectively.

4 More fun with the error sum function

Formulas similar to the formulas in the previous section can be stated by using results on the convergents for continued fractions with “regular” patterns, in particular at least for (some of) the so-called Hurwitz continued fractions, sometimes also called (regular) continued fractions of Hurwitzian type, see, e.g., [22]. We simply list below results that can be used to yield nice formulas for the error sum function we considered. They give in terms of integrals for some reals α and their convergents p_n/q_n the quantity $p_{an+b} - \alpha q_{an+b}$ (for any b in a complete system of residues modulo a) and they are due to Komatsu.

- for $\alpha = \frac{e^{1/\ell s}}{s}$, with s and ℓ any two integers ≥ 2 , and $\alpha = \frac{e^{1/s}}{s}$, with $s \geq 2$, integral expressions for $p_{3n+j} - \alpha q_{3n+j}$ with $j \in \{0, 1, 2\}$ are given in [17, Theorem 3, second part].
- for $\alpha = e^{2/s}$, with $s \geq 3$ and odd, integral expressions for $p_{5n+j} - \alpha q_{5n+j}$ with $j \in \{0, 1, 2, 3, 4\}$ are given in [14];
- for $\alpha = \frac{e^{1/(3s+1)}}{3}$ (resp. $\alpha = \frac{e^{1/(3s+2)}}{3}$), integral expressions for $p_{9n+j} - \alpha q_{9n+j}$ with $j \in \{-6, -5, -4, -3, -2, -1, 0, 1, 2\}$ are given in [17];
- for $\alpha = \sqrt{\frac{v}{u}} \tanh \frac{1}{\sqrt{uv}}$, integral expressions for $p_{2n-1} - \alpha q_{2n-1}$ and $p_{2n} - \alpha q_{2n}$ are given in [16];
- for $\alpha = \sqrt{\frac{v}{u}} \tan \frac{1}{\sqrt{uv}}$, integral expressions for $p_{4n-j} - \alpha q_{4n-j}$ with $j \in \{0, 1, 2, 3\}$ are given in [16].

The last result we would like to cite here is a nice particular case of a theorem of Heteyi [12, Theorem 2.9] (also see [12, p. 21]) which could be used to compute the error sum function for $\alpha = \frac{4(11 \sin(1/2) - 6 \cos(1/2))}{53 \cos(1/2) - 97 \sin(1/2)}$.

Theorem 4 (Heteyi) *We have the following continued fraction expansion*

$$\frac{4(11 \sin(1/2) - 6 \cos(1/2))}{53 \cos(1/2) - 97 \sin(1/2)} = [4, 3, 4, 4, 4, 4, 5, 4, 6, 4, 7, 4, \dots].$$

5 Conclusion

The error sum function of some other continued fractions with “regular” patterns could probably be studied. Another appealing possibility is the definition and study of error sum functions similar continued fractions in the function field case (see in particular [29, 30, 31]). Finally we give a last relation that the second author discovered empirically: we did not locate it in the literature (yet) and did not prove it (yet)

$$\int_0^1 e^{-t^2} dt = 3/8 + \frac{5/4}{3 + \frac{9}{21 + \frac{288}{63 + \frac{\dots}{\dots + \frac{n(n+2)^2(2n-1)^2}{(2n+5)(n^2+n+1) + \dots}}}}$$

Acknowledgements We want to warmly thank J. Shallit who indicated to us the papers [5, 8, 9, 6].

Addendum After we posted the first version of this paper on ArXiv, C. Elsner kindly sent us the preprint [7], which was written a few months ago, and where the reader can find some results in another direction (limit formulas, differential equations, algebraic independence results, relations to Hall’s theorem) but also a proof of the result in Corollary 1

$$\forall \ell \geq 2, \sum_{n \geq 0} |p_n - e^{1/\ell} q_n| = e^{1/\ell} \sqrt{\frac{\pi}{\ell}} \operatorname{erf}(1/\sqrt{\ell}).$$

References

- [1] M. Abramowitz, I. A. Stegun, Handbook of Mathematical Functions With Formulas, Graphs, and Mathematical Tables, National Bureau of Standards Applied Mathematics Series **55**, 10th printing, 1972.
- [2] A. H. Chamseddine, Massive supergravity from spontaneously breaking orthosymplectic gauge symmetry, *Ann. Physics* **113** (1978) 219–234.
- [3] H. Cohn, A short proof of the simple continued fraction expansion of e , *Amer. Math. Monthly* **113** (2006) 57–62.

- [4] C. S. Davis, On some simple continued fractions connected with e , *J. London Math. Soc.* **20** (1945) 194–198.
- [5] C. Elsner, Series of error terms for rational approximations of irrational numbers, *J. Integer Seq.* **14** (2011), Article 11.1.4, 20 p.
- [6] C. Elsner, On error sums of square roots of positive integers with applications to Lucas and Pell numbers, *J. Integer Seq.* **17** (2014), Article 14.4.4, 21 p.
- [7] C. Elsner, A. Klauke, Errorsums for the values of the exponential function, *Bericht Nr. 02014/01*, 1–19; Technische Informationsbibliothek Hannover: RS 8153 (2014,1).
- [8] C. Elsner, M. Stein, On error sum functions formed by convergents of real numbers, *J. Integer Seq.* **14** (2011), Article 11.8.6, 14 p.
- [9] C. Elsner, M. Stein, On the value distribution of error sums for approximations with rational numbers, *Integers* **12** (2012), Article A66, 28 p.
- [10] L. Euler, De fractionibus continuis dissertatio, *Comm. Acad. Sci. Petropol.* **9** (1744) 98–137. Available at <http://eulerarchive.maa.org/docs/originals/E071.pdf> or at <http://www.math.dartmouth.edu/~euler/docs/originals/E071.pdf>
English translation: M. F. Wyman, B. F. Wyman, An essay on continued fractions, *Math. Systems Theory* **18** (1985) 295–328.
- [11] C. Hermite, Sur la fonction exponentielle, *C. R. Acad. Sci.* **77** (1873) 18–24, 74–79, 226–233, and 285–293; also in *Œuvres de Charles Hermite*, publiées sous les auspices de l’Académie des sciences par Émile Picard, vol. 3, Gauthier-Villars, Paris, 1912, pp. 150–181. Available at <https://archive.org/details/oeuvresdecharles03hermuoft>
- [12] G. Heteyi, Hurwitzian continued fractions containing a repeated constant and an arithmetic progression, *SIAM J. Discrete Math.* **28** (2014) 962–985.
- [13] A. Hurwitz, Über die Kettenbrüche, deren Teilnenner arithmetische Reihen bilden, *Zürich. Naturf. Ges.* **41** 2nd Part (1896) 34–64.
- [14] T. Komatsu, A proof of the continued fraction expansion of $e^{2/s}$, *Integers* **7** (2007), Article A30, 8 p.
- [15] T. Komatsu, More on Hurwitz and Tasojev continued fractions, *Sarajevo J. Math.* **4** (2008) 155–180.
- [16] T. Komatsu, Diophantine approximations of \tanh , \tan and linear forms of e in terms of integrals, *Rev. Roum. Math. Pures Appl.* **54** (2009) 223–242.
- [17] T. Komatsu, A diophantine approximation of $e^{1/s}$ in terms of integrals, *Tokyo J. Math.* **32** (2009) 159–176.
- [18] T. Komatsu, Some exact algebraic expressions for the tails of Tasojev continued fractions, *J. Aust. Math. Soc.* **92** (2012) 179–193.

- [19] J. H. Lambert, Mémoires sur quelques propriétés remarquables des quantités transcendentes, circulaires et logarithmiques, *Mémoires de l'Académie royale des sciences de Berlin* **17** (1761/1768) 265–322. Available at <http://www.kuttaka.org/~JHL/L1768b.pdf>
- [20] D. N. Lehmer, Arithmetical theory of certain Hurwitzian continued fractions, *Amer. J. Math.* **40** (1918) 375–390.
- [21] J. H. McCabe, On the Padé table for e^x and the simple continued fractions for e and $e^{L/M}$, *Ramanujan J.* **19** (2009) 95–105.
- [22] J. Mc Laughlin, Some new families of Tasoevian and Hurwitzian continued fractions, *Acta Arith.* **135** (2008) 247–268.
- [23] K. R. Matthews, R. F. C. Walters, Some properties of the continued fraction expansion of $(m/n)e^{1/q}$, *Proc. Cambridge Philos. Soc.* **67** (1970) 67–74.
- [24] C. D. Olds, The simple continued fraction expansion of e , *Amer. Math. Monthly* **77** (1970) 968–974.
- [25] On-Line Encyclopedia of Integer Sequences, available electronically at <http://oeis.org>
- [26] T. J. Osler, A proof of the continued fraction expansion of $e^{1/M}$, *Amer. Math. Monthly* **113** (2006) 62–66.
- [27] A. J. van der Poorten, Continued fraction expansions of values of the exponential function and related fun with continued fractions, *Nieuw Arch. Wisk.* **14** (1996) 221–230.
- [28] J. N. Ridley, G. Petruska, The error-sum function of continued fractions, *Indag. Math.* **11** (2000) 273–282.
- [29] D. S. Thakur, Continued fraction for the exponential for $\mathbf{F}_q[\mathbf{T}]$, *J. Number Theory* **41** (1992) 150–155.
- [30] D. S. Thakur, Exponential and continued fractions, *J. Number Theory* **59** (1996) 248–261.
- [31] D. S. Thakur, Patterns of continued fractions for the analogues of e and related numbers in the function field case, *J. Number Theory* **66** (1997) 129–147.