

INDEX- p ABELIANIZATION DATA OF p -CLASS TOWER GROUPS

DANIEL C. MAYER

ABSTRACT. Given a fixed prime number p , the multiplet of abelian type invariants of the p -class groups of all unramified cyclic degree p extensions of a number field K is called its IPAD (index- p abelianization data). These invariants have proved to be a valuable information for determining the Galois group G_p^2 of the second Hilbert p -class field and the p -capitulation type \varkappa of K . For $p = 3$ and a number field K with elementary p -class group of rank two, all possible IPADs are given in the complete form of several infinite sequences. Iterated IPADs of second order are used to identify the group G_p^∞ of the maximal unramified pro- p extension of K .

1. INTRODUCTION

After a thorough discussion of the terminology used in this article, the logarithmic and power form of abelian type invariants in § 2, and multilayered transfer target types (TTTs), ordered and accumulated index- p abelianization data (IPADs) up to the third order in § 3, we state the main results in § 3.1 on IPADs of exceptional form, and in § 3.2 on IPADs in parametrized infinite sequences. These main theorems give all possible IPADs of number fields K with 3-class group $\text{Cl}_3(K)$ of type $(3, 3)$.

Before we turn to applications in extreme computing, that is, squeezing the computational algebra systems PARI [33] and MAGMA [6, 7, 23] to their limits in § 5, where we show how to detect malformed IPADs in § 5.1, and how to complete partial p -capitulation types in § 5.2, we have to establish a componentwise correspondence between transfer kernel types (TKTs) and IPADs in § 4 by exploiting details of proofs which were given in [28].

Iterated IPADs of second order are used in § 6 for the indirect calculation of TKTs in § 6.1, and for determining the exact length $\ell_p(K)$ of the p -class tower of a number field K in § 6.2. This sophisticated technique proves $\ell_3(K) = 3$ for $K = \mathbb{Q}(\sqrt{d})$ with $d \in \{342\,664, 957\,013\}$ (the first real quadratic fields) and $d = -3\,896$ (the first tough complex quadratic field after the ‘easy’ $d = -9\,748$ [13]), which resisted all attempts up to now.

Finally, we emphasize that infinite p -class towers admit an unknown wealth of possible fine structure in § 7 on complex quadratic fields K having a 3-class group $\text{Cl}_3(K)$ of type $(3, 3, 3)$.

2. ABELIAN TYPE INVARIANTS

Let p be a prime number and A be a finite abelian p -group. According to the main theorem on finitely generated abelian groups, there exists a non-negative integer $r \geq 0$, the *rank* of A , and a sequence n_1, \dots, n_r of positive integers such that $n_1 \leq n_2 \leq \dots \leq n_r$ and

$$(2.1) \quad A \simeq \mathbb{Z}/p^{n_1}\mathbb{Z} \oplus \dots \oplus \mathbb{Z}/p^{n_r}\mathbb{Z}.$$

The powers $d_i := p^{n_i}$, $1 \leq i \leq r$, are known as the *elementary divisors* of A , since $d_i \mid d_{i+1}$ for each $1 \leq i \leq r - 1$. It is convenient to collect equal elementary divisors in formal powers with

Date: February 11, 2015.

2000 Mathematics Subject Classification. Primary 11R29, 11R37, 11R11; secondary 20D15.

Key words and phrases. p -class groups, p -principalization types, p -class field towers, quadratic fields, second p -class groups, p -class tower groups, coclass graphs.

Research supported by the Austrian Science Fund (FWF): P 26008-N25.

positive exponents r_1, \dots, r_s such that $r_1 + \dots + r_s = r$, $0 \leq s \leq r$, and

$$n_1 = \dots = n_{r_1} < n_{r_1+1} = \dots = n_{r_1+r_2} < \dots < n_{r_1+\dots+r_{s-1}+1} = \dots = n_{r_1+\dots+r_s}.$$

The cumbersome subscripts can be avoided by defining $m_j := n_{r_1+\dots+r_j}$ for each $1 \leq j \leq s$. Then

$$(2.2) \quad A \simeq (\mathbb{Z}/p^{m_1}\mathbb{Z})^{r_1} \oplus \dots \oplus (\mathbb{Z}/p^{m_s}\mathbb{Z})^{r_s}$$

and we can define:

Definition 2.1. The *abelian type invariants* (ATI) of A are given by the sequence

$$(2.3) \quad (m_1^{r_1}, \dots, m_s^{r_s})$$

of strictly increasing positive integers $m_1 < \dots < m_s$ with multiplicities r_1, \dots, r_s written as formal exponents indicating iteration.

Remark 2.1. The integers m_j are the p -logarithms of the elementary divisors d_i .

- (1) For abelian type invariants of high complexity, the *logarithmic form* in Definition 2.1 requires considerably less space (e.g. in tables) than the usual *power form*

$$(2.4) \quad (\overbrace{p^{m_1}, \dots, p^{m_1}}^{r_1}, \dots, \overbrace{p^{m_s}, \dots, p^{m_s}}^{r_s}).$$

- (2) For brevity, we can even omit the commas separating the entries of the logarithmic form of abelian type invariants, provided all the m_j remain smaller than 10.
- (3) A further advantage of the brief logarithmic notation is the independence of the prime p , in particular when p -groups with distinct p are being compared.
- (4) Finally, since our preference is to select generators of finite p -groups with decreasing orders, we agree to write abelian type invariants from the right to the left, in both forms.

Example 2.1. For instance, if $p = 3$, then the abelian type invariants (21^4) in logarithmic form correspond to the power form $(9, 3, 3, 3, 3)$ and $(2^2 1^2)$ corresponds to $(9, 9, 3, 3)$.

Now let G be an arbitrary finite p -group or infinite topological pro- p group with derived subgroup G' and finite abelianization $G^{\text{ab}} = G/G'$.

Definition 2.2. The abelian type invariants of the commutator quotient group G^{ab} are called the *abelian quotient invariants* (AQI) of G .

3. INDEX- p ABELIANIZATION DATA

Let p be a fixed prime number and K be a number field with p -class group $\text{Cl}_p(K)$ of order p^v , where $v \geq 0$ denotes a non-negative integer.

According to the Artin reciprocity law of class field theory [1], $\text{Cl}_p(K)$ is isomorphic to the commutator quotient group G/G' of the Galois group $G = \text{Gal}(\mathbb{F}_p^\infty(K) \mid K)$ of the maximal unramified pro- p extension $\mathbb{F}_p^\infty(K)$ of K . G is called the *p -tower group* of K . The fixed field of the commutator subgroup G' in $\mathbb{F}_p^\infty(K)$ is the maximal abelian unramified p -extension of K , that is the (first) Hilbert p -class field $\mathbb{F}_p^1(K)$ of K with Galois group $\text{Gal}(\mathbb{F}_p^1(K) \mid K) \simeq G/G'$. The derived subgroup G' is a closed (and open) subgroup of finite index $(G : G') = p^v$ in the topological pro- p group G .

Definition 3.1. For each integer $0 \leq n \leq v$, the system

$$(3.1) \quad \text{Lyr}_n(K) = \{K \leq L \leq \mathbb{F}_p^1(K) \mid [L : K] = p^n\}$$

of intermediate fields $K \leq L \leq \mathbb{F}_p^1(K)$ with relative degree $[L : K] = p^n$ is called the *n -th layer* of abelian unramified p -extensions of K . In particular, for $n = 0$, K forms the *bottom layer* $\text{Lyr}_0(K) = \{K\}$, and for $n = v$, $\mathbb{F}_p^1(K)$ forms the *top layer* $\text{Lyr}_v(K) = \{\mathbb{F}_p^1(K)\}$.

Now let $0 \leq n \leq v$ be a fixed integer and suppose that $K \leq L \leq F_p^1(K)$ belongs to the n -th layer. Then the Galois group $H = \text{Gal}(F_p^\infty(K) \mid L)$ is of finite index $(G : H) = [L : K] = p^n$ in the p -tower group G of K and the quotient $G/H \simeq \text{Gal}(L \mid K)$ is abelian, since H contains the commutator subgroup $G' = \text{Gal}(F_p^\infty(K) \mid F_p^1(K))$ of G .

Definition 3.2. For each integer $0 \leq n \leq v$, the system

$$(3.2) \quad \text{Lyr}_n(G) = \{G' \leq H \leq G \mid (G : H) = p^n\}$$

of intermediate groups $G' \leq H \leq G$ with index $(G : H) = p^n$ is called the n -th layer of normal subgroups of G with abelian quotients G/H . In particular, for $n = 0$, G forms the *top layer* $\text{Lyr}_0(G) = \{G\}$, and for $n = v$, G' forms the *bottom layer* $\text{Lyr}_v(G) = \{G'\}$.

A further application of Artin's reciprocity law [1] shows that

$$(3.3) \quad H/H' = \text{Gal}(F_p^\infty(K) \mid L) / \text{Gal}(F_p^\infty(K) \mid F_p^1(L)) \simeq \text{Gal}(F_p^1(L) \mid L) \simeq \text{Cl}_p(L),$$

for every subgroup $H \in \text{Lyr}_n(G)$ and its corresponding extension field $L \in \text{Lyr}_n(K)$, where $0 \leq n \leq v$ is fixed (but arbitrary).

Since the abelianization $H^{\text{ab}} = H/H'$ forms the target of the Artin transfer homomorphism $T_{G,H} : G \rightarrow H/H'$ from G to H , we introduced a preliminary instance of the following terminology in [27, Dfn.1.1, p.403].

Definition 3.3. For each integer $0 \leq n \leq v$, the multiplet $\tau_n(G) = (H/H')_{H \in \text{Lyr}_n(G)}$, where each member H/H' is interpreted rather as its abelian type invariants, is called the n -th layer of the *transfer target type* (TTT) of the pro- p group G ,

$$(3.4) \quad \tau(G) = [\tau_0(G); \dots; \tau_v(G)], \text{ where } \tau_n(G) = (H/H')_{H \in \text{Lyr}_n(G)} \text{ for each } 0 \leq n \leq v.$$

Similarly, the multiplet $\tau_n(K) = (\text{Cl}_p(L))_{L \in \text{Lyr}_n(K)}$, where each member $\text{Cl}_p(L)$ is interpreted rather as its abelian type invariants, is called the n -th layer of the *transfer target type* (TTT) of the number field K ,

$$(3.5) \quad \tau(K) = [\tau_0(K); \dots; \tau_v(K)], \text{ where } \tau_n(K) = (\text{Cl}_p(L))_{L \in \text{Lyr}_n(K)} \text{ for each } 0 \leq n \leq v.$$

Remark 3.1. (1) If it is necessary to specify the underlying prime number p , then the symbol $\tau(p, G)$, resp. $\tau(p, K)$, can be used for the TTT.

(2) Suppose that $0 < n < v$. If an ordering is defined for the elements of $\text{Lyr}_n(G)$, resp. $\text{Lyr}_n(K)$, then the same ordering is applied to the members of the layer $\tau_n(G)$, resp. $\tau_n(K)$, and the TTT layer is called *ordered*. Otherwise, the TTT layer is called *unordered* or *accumulated*, since equal components are collected in powers with formal exponents denoting iteration.

(3) In view of the considerations in Equation (3.3), it is clear that we have the equality

$$(3.6) \quad \tau(G) = \tau(K),$$

in the sense of componentwise isomorphisms.

Since it is increasingly difficult to compute the structure of the p -class groups $\text{Cl}_p(L)$ of extension fields $L \in \text{Lyr}_n(K)$ in higher layers with $n \geq 2$, it is frequently sufficient to make use of information in the first layer only, that is the layer of subgroups with index p . Therefore, Boston, Bush and Hajir [8] invented the following *first order approximation* of the TTT, a concept which had been used in earlier work already [9, 12, 3, 10, 31], without explicit terminology.

Definition 3.4. The restriction

$$(3.7) \quad \begin{aligned} \tau^{(1)}(G) &= [\tau_0(G); \tau_1(G)], \text{ resp.} \\ \tau^{(1)}(K) &= [\tau_0(K); \tau_1(K)], \end{aligned}$$

of the TTT $\tau(G)$, resp. $\tau(K)$, to the zeroth and first layer is called the *index- p abelianization data* (IPAD) of G , resp. K .

So, the complete TTT is an extension of the IPAD. However, there also exists another extension of the IPAD which is not covered by the TTT. It has also been used already in previous investigations by Boston, Bush and Nover [12, 10, 31] and is constructed from the usual IPAD $[\tau_0(K); \tau_1(K)]$ of K , firstly, by observing that $\tau_1(K) = (Cl_p(L))_{L \in \text{Ly}_{r_1}(K)}$ can be viewed as $\tau_1(K) = (\tau_0(L))_{L \in \text{Ly}_{r_1}(K)}$ and, secondly, by extending each $\tau_0(L)$ to the IPAD $[\tau_0(L); \tau_1(L)]$ of L .

Definition 3.5. The family

$$(3.8) \quad \begin{aligned} \tau^{(2)}(G) &= [\tau_0(G); ([\tau_0(H); \tau_1(H)])_{H \in \text{Ly}_{r_1}(G)}], \text{ resp.} \\ \tau^{(2)}(K) &= [\tau_0(K); ([\tau_0(L); \tau_1(L)])_{L \in \text{Ly}_{r_1}(K)}], \end{aligned}$$

is called the *iterated IPAD of second order* of G , resp. K .

The concept of iterated IPADs as given in Dfn. 3.5 is restricted to the second order and first layers, and thus is open for further generalization (higher orders and higher layers). Since it could be useful for 2-power extensions, whose absolute degrees increase moderately and remain manageable by MAGMA or PARI, we briefly indicate how the *iterated IPAD of third order* could be defined:

$$(3.9) \quad \begin{aligned} \tau^{(3)}(G) &= [\tau_0(G); ([\tau_0(H); ([\tau_0(I); \tau_1(I)])_{I \in \text{Ly}_{r_1}(H)}])_{H \in \text{Ly}_{r_1}(G)}], \text{ resp.} \\ \tau^{(3)}(K) &= [\tau_0(K); ([\tau_0(L); ([\tau_0(M); \tau_1(M)])_{M \in \text{Ly}_{r_1}(L)}])_{L \in \text{Ly}_{r_1}(K)}]. \end{aligned}$$

3.1. Sporadic IPADs. In the next two central theorems, we present complete specifications of all possible IPADs of pro- p groups G for $p = 3$ and the simplest case of an abelianization G/G' of type $(3, 3)$. We start with pro-3-groups G whose metabelianizations G/G'' are vertices on sporadic parts of coclass graphs outside of coclass trees.

Since the abelian type invariants of the members of TTT layers will depend on the parity of the nilpotency class c or coclass r , a more economic notation, avoiding the tedious distinction of the cases odd or even, is provided by the following definition.

Definition 3.6. For an integer $n \geq 2$, the *nearly homocyclic abelian 3-group* $A(3, n)$ of order 3^n is defined by its type invariants $(q+r, q) \hat{=} (3^{q+r}, 3^q)$, where the quotient $q \geq 1$ and the remainder $0 \leq r < 2$ are determined uniquely by the Euclidean division $n = 2q + r$. Two degenerate cases are included by putting $A(3, 1) = (1) \hat{=} (3)$ the cyclic group C_3 of order 3 and $A(3, 0) = (0) \hat{=} 1$ the trivial group of order 1.

Theorem 3.1. (*First Main Theorem on $p = 3$, $G/G' \simeq (3, 3)$, and G/G'' of small class*)

Let G be a pro-3 group having a transfer target type $\tau(G) = [\tau_0(G); \tau_1(G); \tau_2(G)]$ with top layer component $\tau_0(G) = 1^2$. Let $0 \leq k \leq 1$ denote the defect of commutativity [27, § 3.1.1, p.412, and § 3.3.2, p.429] of the metabelianization G/G'' of G . Then the ordered first layer $\tau_1(G)$ and the bottom layer $\tau_2(G)$ are given in the following way.

(1) If G/G'' is of coclass $\text{cc}(G/G'') = 1$ and nilpotency class $\text{cl}(G/G'') = c \leq 3$, then

$$(3.10) \quad \begin{aligned} \tau_1(G) &= (1)^4; \tau_2(G) = (0), \text{ if } c = 1, G \simeq \langle 9, 2 \rangle, \\ \tau_1(G) &= (1^2)^4; \tau_2(G) = (1), \text{ if } c = 2, G \simeq \langle 27, 3 \rangle, \\ \tau_1(G) &= (1^2, (\mathbf{2})^3); \tau_2(G) = (1), \text{ if } c = 2, G \simeq \langle 27, 4 \rangle, \\ \tau_1(G) &= (\mathbf{1}^3, (1^2)^3); \tau_2(G) = (1^2), \text{ if } c = 3, G \simeq \langle 81, 7 \rangle, \\ \tau_1(G) &= (21, (1^2)^3); \tau_2(G) = (1^2), \text{ if } c = 3, G \simeq \langle 81, 8|9|10 \rangle, \end{aligned}$$

where generally $G'' = 1$.

(2) If G/G'' is of coclass $\text{cc}(G/G'') = 2$ and nilpotency class $\text{cl}(G/G'') = c = 3$, then

$$(3.11) \quad \begin{aligned} \tau_1(G) &= ((21)^2, 1^3, 21); \tau_2(G) = (1^3), \text{ if } G \simeq \langle 243, 5 \rangle \text{ or } G/G'' \simeq \langle 243, 6 \rangle, \\ \tau_1(G) &= ((21)^2, (1^3)^2); \tau_2(G) = (1^3), \text{ if } G/G'' \simeq \langle 243, 3 \rangle, \\ \tau_1(G) &= (1^3, 21, 1^3, 21); \tau_2(G) = (1^3), \text{ if } G \simeq \langle 243, 7 \rangle, \\ \tau_1(G) &= ((\mathbf{1}^3)^2, 21, 1^3); \tau_2(G) = (1^3), \text{ if } G/G'' \simeq \langle 243, 4 \rangle, \\ \tau_1(G) &= (21)^4; \tau_2(G) = (1^3), \text{ if } G/G'' \simeq \langle 243, 8|9 \rangle, \end{aligned}$$

where $G'' = 1$ can be warranted for $G/G'' \simeq \langle 243, 5|7 \rangle$ only.

However, if $\text{cl}(G/G'') = c = 4$ with $k = 1$, then

$$(3.12) \quad \begin{aligned} \tau_1(G) &= ((21)^2, (1^3)^2); \tau_2(G) = (21^2), \text{ if } G/G'' \simeq \langle 729, 37|38|39 \rangle, \\ \tau_1(G) &= ((21)^2, (1^3)^2); \tau_2(G) = (\mathbf{1}^4), \text{ if } G/G'' \simeq \langle 729, 34|35|36 \rangle, \\ \tau_1(G) &= ((1^3)^2, 21, 1^3); \tau_2(G) = (21^2), \text{ if } G/G'' \simeq \langle 729, 44|45|46|47 \rangle, \\ \tau_1(G) &= (21)^4; \tau_2(G) = (\mathbf{1}^4), \text{ if } G/G'' \simeq \langle 729, 56|57 \rangle. \end{aligned}$$

(3) If G/G'' is of coclass $\text{cc}(G/G'') = r \geq 3$ and nilpotency class $\text{cl}(G/G'') = c = r + 1$, then

$$(3.13) \quad \tau_1(G) = (A(3, r + 1)^2, (1^2)^3); \tau_2(G) = A(3, r) \times A(3, r - 1) \text{ and } k = 0.$$

However, if $c = r + 2$, then

$$(3.14) \quad \begin{aligned} \tau_1(G) &= (A(3, r + 2), A(3, r + 1), (1^2)^3); \tau_2(G) = A(3, r + 1) \times A(3, r - 1), \text{ if } k = 0 \\ \tau_1(G) &= (A(3, r + 1)^2, (1^2)^3); \tau_2(G) = A(3, r + 1) \times A(3, r - 1), \text{ if } k = 1, \text{ regular case,} \\ \tau_1(G) &= (A(3, r + 1)^2, (1^2)^3); \tau_2(G) = \mathbf{A}(3, \mathbf{r}) \times \mathbf{A}(3, \mathbf{r}), \text{ if } k = 1, \text{ irregular case,} \end{aligned}$$

where the irregular case can only occur for even class and coclass $c = r + 2 \equiv 0 \pmod{2}$, positive defect of commutativity $k = 1$, and relational parameter $\rho = -1$ in [28, Eqn.(3.6), p.424] or [27, Eqn.(3.3), p.430].

Proof. Since this proof heavily relies on our earlier paper [28], it should be pointed out that, for a p -group G , the index of nilpotency $m = c + 1$ is used generally instead of the nilpotency class $\text{cl}(G) = c = m - 1$ and the invariant $e = r + 1$ frequently (but not always) replaces the coclass $\text{cc}(G) = r = e - 1$ in that paper.

- (1) Using the association between the identifier of G in the SmallGroups Library [4, 5] and the transfer kernel type (TKT) [26], which is visualized in [28, Fig.3.1, p.423], this claim follows from [28, Thm.4.1, p.427, and Tbl.4.1, p.429].
- (2) For $c = 3$, resp. $c = 4$ with $k = 1$, the statement is a consequence of [28, Thm.4.2 and Tbl.4.3, p.434], resp. [28, Thm.4.3 and Tbl.4.5, p.438], when the association between the identifier of G in the SmallGroups Database and the TKT is taken into consideration, as visualized in [28, Fig.4.1, p.433].
- (3) All the regular cases behave completely similar as the general case in Theorem 3.2, item (3), Equation (3.18). In the irregular case, only the bottom layer $\tau_2(G)$, consisting of the abelian quotient invariants G'/G'' of the derived subgroup G' , is exceptional and must be taken from [28, Appendix § 8, Thm.8.8, p.461].

□

3.2. Infinite IPAD sequences. Now we come to the IPADs of pro- p -groups G whose metabelianizations G/G'' are members of infinite periodic sequences, inclusively mainlines, of coclass trees.

Theorem 3.2. (Second Main Theorem on $p = 3$, $G/G' \simeq (3, 3)$, and G/G'' of large class)
 Let G be a pro-3 group having a transfer target type $\tau(G) = [\tau_0(G); \tau_1(G); \tau_2(G)]$ with top layer component $\tau_0(G) = 1^2$. Let $0 \leq k \leq 1$ denote the defect of commutativity [27, § 3.1.1, p.412, and § 3.3.2, p.429] of the metabelianization G/G'' of G . Then the ordered first layer $\tau_1(G)$ and the bottom layer $\tau_2(G)$ are given in the following way.

- (1) If G/G'' is of coclass $\text{cc}(G/G'') = 1$ and nilpotency class $\text{cl}(G/G'') = c \geq 4$, then

$$(3.15) \quad \begin{aligned} \tau_1(G) &= (A(3, c - k), (1^2)^3); \\ \tau_2(G) &= A(3, c - 1). \end{aligned}$$

- (2) If G/G'' is of coclass $\text{cc}(G/G'') = 2$ and nilpotency class $\text{cl}(G/G'') = c \geq 5$, or $c = 4$ with $k = 0$, then

$$(3.16) \quad \begin{aligned} \tau_1(G) &= (A(3, c - k), 21, (1^3)^2) \text{ or} \\ \tau_1(G) &= (A(3, c - k), 21, 1^3, 21) \text{ or} \\ \tau_1(G) &= (A(3, c - k), (21)^3), \end{aligned}$$

in dependence on the coclass tree $G/G'' \in \mathcal{T}^2(\langle 729, i \rangle)$, $i \in \{40, 49, 54\}$, but uniformly

$$(3.17) \quad \tau_2(G) = A(3, c - 1) \times A(3, 1).$$

- (3) If G/G'' is of coclass $\text{cc}(G/G'') = r \geq 3$ and nilpotency class $\text{cl}(G/G'') = c \geq r + 3$, or $c = r + 2$ with $k = 0$, then

$$(3.18) \quad \begin{aligned} \tau_1(G) &= (A(3, c - k), A(3, r + 1), (1^2)^3); \\ \tau_2(G) &= A(3, c - 1) \times A(3, r - 1). \end{aligned}$$

The first member H_1/H_1' of the ordered first layer $\tau_1(G)$ reveals a uni-polarization (dependence on the nilpotency class c) whereas the other three members H_i/H_i' , $2 \leq i \leq 4$, show a stabilization (independence of c) for fixed coclass r .

Proof. Again, we make use of [28], and we point out that, for a p -group G , the index of nilpotency $m = c + 1$ is used generally instead of the nilpotency class $\text{cl}(G) = c = m - 1$ and the invariant $e = r + 1$ frequently (but not always) replaces the coclass $\text{cc}(G) = r = e - 1$ in that paper.

- (1) All components of $\tau_1(G)$ are given in [28, § 3.1, Thm.3.1, Eqn.(3.4)–(3.5), p.421] when their ordering is defined by the special selection of generators [28, § 3.1, Eqn.(3.1)–(3.2), p.420]. There is only a unique coclass tree with 3-groups of coclass 1.
- (2) The first component of $\tau_1(G)$ is given in [28, § 3.2, Thm.3.2, Eqn.(3.7), p.424], and the last three components of $\tau_1(G)$ are given in [28, § 4.5, Thm.4.4, p.440] and [28, § 4.5, Tbl.4.7, p.441], when their ordering is defined by the special selection of generators [28, § 3.2, Eqn.(3.6), p.424]. The invariant $\varepsilon \in \{0, 1, 2\}$ [28], which counts IPAD components of rank 3, decides to which of the mentioned three coclass trees the group G belongs [27, Fig.3.6–3.7, pp.442–443].
- (3) The first two components of $\tau_1(G)$ are given in [28, § 3.2, Thm.3.2, Eqn.(3.7)–(3.8), p.424], and the last two components of $\tau_1(G)$ are given in [28, § 4.6, Thm.4.5, p.444], when their ordering is defined by the special selection of generators [28, § 3.2, Eqn.(3.6), p.424]. For coclass bigger than 2, it is irrelevant to which of the four (in the case of odd coclass r) or six (in the case of even coclass r) coclass trees the group G belongs. The IPAD is independent of this detailed information, provided that $c \geq r + 3$.

Finally, the bottom layer $\tau_2(G)$, consisting of the abelian quotient invariants G'/G'' of the derived subgroup G' , is generally taken from [28, Appendix § 8, Thm.8.8, p.461].

□

4. COMPONENTWISE CORRESPONDENCE OF IPAD AND TKT

Within this section, where generally $p = 3$, we employ some special terminology. We say a class of a base field K remains *resistant* if it does not capitulate in any unramified cyclic cubic extension $L|K$. When the 3-class group of K is of type $(3, 3)$ the next layer of unramified abelian extensions is already the top layer consisting of the Hilbert 3-class field $F_3^1(K)$, where the resistant class must capitulate, according to the Hilbert/Artin/Furtwängler principal ideal theorem.

Our desire is to show that the components of the ordered IPAD and TKT are in a strict correspondence to each other. For this purpose, we use details of the proofs given in [28], where generators of metabelian 3-groups G with $G/G' \simeq (3, 3)$ were selected in a canonical way, particularly adequate for theoretical aspects. Since we now prefer a more computational aspect, we translate the results into a form which is given by the computational algebra system MAGMA [23].

To be specific, we choose the vertices of two important coclass trees for illustrating these peculiar techniques. The vertices of depth (distance from the mainline) at most 1 of both coclass trees, with roots $\langle 243, 6 \rangle$ and $\langle 243, 8 \rangle$ [27, Fig.3.6–3.7, pp.442–443], are metabelian 3-groups G with order $|G| \geq 3^5$, nilpotency class $c = \text{cl}(G) \geq 3$, and fixed coclass $\text{cc}(G) = 2$.

4.1. The coclass tree $\mathcal{T}^2(\langle 243, 6 \rangle)$.

Remark 4.1. The first layers of the TTT and TKT of vertices of depth at most 1 of the coclass tree $\mathcal{T}^2(\langle 243, 6 \rangle)$ [27, Fig.3.6, p.442] consist of four components each, and share the following common properties with respect to MAGMA's selection of generators:

- (1) polarization (dependence on the class c) at the first component,
- (2) stabilization (independence of the class c) at the last three components,
- (3) rank 3 at the second TTT component ($\varepsilon = 1$ in [28]).

Using the class c , resp. an asterisk, as wildcard characters, these common properties can be summarized as follows, now including the details of the stabilization:

$$(4.1) \quad \tau_1(G) = [A(3, c), 1^3, (21)^2], \text{ and } \varkappa_1(G) = (*, 1, 2, 2).$$

However, to assure the general applicability of the theorems and corollaries in this section, we aim at independency of the selection of generators (and thus invariance under permutations).

Theorem 4.1. (*in field theoretic terminology*)

- (1) The class associated with the polarization becomes principal in the extension with rank 3.
- (2) The class associated with rank 3 becomes principal in both extensions of type (21), in particular, $\varkappa_1(G)$ cannot be a permutation and can have at most one fixed point.

Remark 4.2. Aside from the common properties, there also arise variations due to the polarization, which we first express with respect to MAGMA's selection of generators:

- (1) The TKT is E.6, $\varkappa_1(G) = (1, 1, 2, 2)$, if and only if the polarized extension reveals a *fixed point* principalization.
- (2) The TKT is E.14, $\varkappa_1(G) \in \{(3, 1, 2, 2), (4, 1, 2, 2)\}$, if and only if one of the classes associated with type (21) becomes principal in the polarized extension.
- (3) The TKT is H.4, $\varkappa_1(G) = (2, 1, 2, 2)$, if and only if the class associated with rank 3 becomes principal in the polarized extension.
- (4) The TKT is c.18, $\varkappa_1(G) = (0, 1, 2, 2)$, if and only if the polarized extension reveals a *total* principalization (indicated by 0).

Corollary 4.1. (*in field theoretic terminology*)

- (1) For the TKTs E.6 and H.4, both classes associated with type (21) remain resistant, for TKT E.14 only one of them.
- (2) All extensions satisfy Taussky's condition (B) [38], with the single exception of of the polarized extension in the case of TKT E.6 or c.18, which satisfies condition (A).
- (3) TKT E.6 has a single fixed point, E.14 contains a 3-cycle, and H.4 contains a 2-cycle.

Proof. (of Theorem 4.1 and Corollary 4.1)

Observe that in [28], the index of nilpotency $m = c + 1$ and the invariant $e = r + 1$ are used rather than the nilpotency class $c = m - 1$ and the coclass $r = e - 1$. The claims are a consequence of [28, § 4.5, Tbl.4.7, p.441], when we perform a permutation from the first layer TKT and TTT

$$\varkappa_1(G) = (*, 3, 1, 3), \quad \tau_1(G) = [A(3, c), 21, 1^3, 21],$$

with respect to the canonical generators, to the corresponding invariants

$$\varkappa_1(G) = (*, 1, 2, 2), \quad \tau_1(G) = [A(3, c), 1^3, (21)^2],$$

with respect to MAGMA's generators. □

4.2. The coclass tree $\mathcal{T}^2(\langle 243, 8 \rangle)$.

Remark 4.3. The first layer TTT and TKT of vertices of depth at most 1 of the coclass tree $\mathcal{T}^2(\langle 243, 8 \rangle)$ [27, Fig.3.7, p.443] consist of four components each, and share the following common properties with respect to MAGMA's choice of generators:

- (1) polarization (dependence on the class c) at the second component,
- (2) stabilization (independence of the class c) at the other three components,
- (3) rank 3 does not occur at any TTT component ($\varepsilon = 0$ in [28]).

Using the class c , resp. an asterisk, as wildcard characters, the common properties can be summarized as follows, now including details of the stabilization:

$$(4.2) \quad \tau_1(G) = [21, A(3, c), (21)^2], \quad \text{and} \quad \varkappa_1(G) = (2, *, 3, 4).$$

Again, we have to ensure the general applicability of the following theorem and corollary, which must be independent of the choice of generators (and thus invariant under permutations).

Theorem 4.2. (*in field theoretic terminology*)

- (1) Two extensions of type (21) reveal fixed point principalization satisfying condition (A) [38].
- (2) The remaining extension of type (21) satisfies condition (B), since the class associated with the polarization becomes principal there.

Remark 4.4. Next, we come to variations caused by the polarization, which we now express with respect to MAGMA's choice of generators:

- (1) The TKT is E.8, $\varkappa_1(G) = (2, 2, 3, 4)$, if and only if the polarized extension reveals a *fixed point* principalization.
- (2) The TKT is E.9, $\varkappa_1(G) \in \{(2, 3, 3, 4), (2, 4, 3, 4)\}$, if and only if one of the classes associated with fixed points becomes principal in the polarized extension.
- (3) The TKT is G.16, $\varkappa_1(G) = (2, 1, 3, 4)$, if and only if the class associated with type (21), satisfying condition (B), becomes principal in the polarized extension.
- (4) The TKT is c.21, $\varkappa_1(G) = (2, 0, 3, 4)$, if and only if the polarized extension reveals a *total* principalization (indicated by 0).

Corollary 4.2. (*in field theoretic terminology*)

- (1) For the TKTs E.8 and E.9, the class associated with the polarization remains resistant,
- (2) The polarized extension satisfies condition (B) [38] in the case of TKT E.9 or G.16, and it satisfies condition (A) in the case of TKT E.8 or c.21.
- (3) TKT G.16 is a permutation containing a 2-cycle, and TKT E.8 is the unique TKT possessing three fixed points.

Proof. (of Theorem 4.2 and Corollary 4.2)

In our paper [28], the index of nilpotency $m = c + 1$ and the invariant $e = r + 1$ are used rather than the nilpotency class $c = m - 1$ and the coclass $r = e - 1$. All claims are a consequence of [28, § 4.5, Tbl.4.7, p.441], provided we perform a transformation from the first layer TKT and TTT

$$\varkappa_1(G) = (*, 2, 3, 1), \quad \tau_1(G) = [A(3, c), (21)^3],$$

with respect to the canonical generators, to the corresponding invariants

$$\varkappa_1(G) = (2, *, 3, 4), \quad \tau_1(G) = [21, A(3, c), (21)^2],$$

with respect to MAGMA's generators. \square

5. APPLICATIONS IN EXTREME COMPUTING

5.1. Application 1: Sifting malformed IPADs.

Definition 5.1. An IPAD with bottom layer component $\tau_0(K) = (3, 3)$ is called *malformed* if it is not covered by Theorems 3.1 and 3.2.

To verify predicted asymptotic densities of maximal unramified pro-3 extensions in the article [8] numerically, the IPADs of all complex quadratic fields $K = \mathbb{Q}(\sqrt{d})$ with discriminants $-10^8 < d < 0$ and 3-class rank $r_3(K) = 2$ were computed with the aid of PARI/GP [33]. In particular, there occurred 276 375, resp. 122 444, such fields with 3-class group $\text{Cl}_3(K)$ of type $(3, 3)$, resp. $(9, 3)$.

Example 5.1. A check of all 276 375 IPADs for complex quadratic fields with type $(3, 3)$ in the range $-10^8 < d < 0$ of discriminants, for which Theorem 3.2 states that the 3-class groups of the 4 unramified cyclic cubic extensions can only have 3-rank 2, except for the unique type $(3, 3, 3)$, revealed that the following 5 IPADs were computed erroneously by the used version of PARI/GP [33] in [8]. The successful recomputation was done with MAGMA [23].

- (1) For $d = -96\,174\,803$, the erroneous IPAD $\tau^{(1)}(K) = [(3, 3); (3, 3, 3), (9, 3, 3, 3), (27, 9)^2]$ contained the malformed component $(9, 3, 3, 3)$ instead of the correct $(3, 3, 3, 3)$. The transfer kernel type (TKT) [26, 27] turned out to be F.12.
- (2) For $d = -77\,254\,244$, the erroneous IPAD $\tau^{(1)}(K) = [(3, 3); (3, 3, 3)^2, (3, 3, 3, 3), (9, 3)]$ contained the malformed component $(3, 3, 3, 3)$ instead of the correct $(3, 3, 3)$. Its TKT is H.4.
- (3) For $d = -73\,847\,683$, the erroneous IPAD $\tau^{(1)}(K) = [(3, 3); (3, 3, 3), (9, 3, 3), (9, 3)^2]$ contained the malformed component $(9, 3, 3)$ instead of the correct $(9, 3)$. The TKT is D.10.
- (4) For $d = -81\,412\,223$, the erroneous IPAD $\tau^{(1)}(K) = [(3, 3); (9, 3, 3), (9, 3)^2, (27, 9)]$ contained the malformed component $(9, 3, 3)$ instead of the correct $(9, 3)$. This could be a TKT E.8 or E.9 or G.16.
- (5) For $d = -82\,300\,871$, the erroneous IPAD $\tau^{(1)}(K) = [(3, 3); (3, 3, 3), (9, 3), (9, 9, 3), (27, 9)]$ contained the malformed component $(9, 9, 3)$ instead of the correct $(9, 3)$. This could be a TKT E.6 or E.14 or H.4.

For the last two cases, Magma failed to determine the TKT. Nevertheless, none of the discriminants

$$d \in \{-73\,847\,683, -77\,254\,244, -81\,412\,223, -82\,300\,871, -96\,174\,803\}$$

is particularly spectacular.

Example 5.2. We also checked all 122 444 IPADs for complex quadratic fields with type $(9, 3)$ in the range $-10^8 < d < 0$ of discriminants, Again, we found exactly 5 errors among these IPADs which had been computed by PARI/GP [33] in [8]. For the recomputation we used MAGMA [23]. The study of this extensive material was very helpful for the deeper understanding of 3-groups having abelianization of type $(9, 3)$. Systematic results in the style of Theorems 3.1 and 3.2 will be given in a forthcoming paper. The abbreviation pTKT means the *punctured* TKT.

- (1) For $d = -94\,304\,231$, the erroneous IPAD $\tau^{(1)}(K) = [(9, 3); (9, 3, 3), (27, 3), (9, 9, 9), (27, 9)]$ contained the malformed component $(27, 9)$ instead of the correct $(27, 3)$. This could be a homocyclic pTKT B.2 or C.4 or D.5.
- (2) For $d = -79\,749\,087$, the erroneous IPAD $\tau^{(1)}(K) = [(9, 3); (9, 3, 3)^2, (27, 3, 3), (27, 3)]$ contained the malformed component $(27, 3, 3)$ instead of the correct $(27, 3)$. It is a pTKT D.11.

- (3) For $d = -74\,771\,240$, the erroneous IPAD $\tau^{(1)}(K) = [(9, 3); (9, 3, 3), (27, 3, 3), (27, 3), (9, 9, 9)]$ contained the malformed component $(27, 3, 3)$ instead of the correct $(27, 3)$. It could be a homocyclic pTKT B.2 or C.4 or D.5.
- (4) For $d = -70\,204\,919$, the erroneous IPAD $\tau^{(1)}(K) = [(9, 3); (9, 3, 3), (27, 3)^2, (81, 27, 27)]$ contained the malformed component $(81, 27, 27)$ instead of the correct $(81, 27, 3)$. This could be a pTKT B.2 or C.4 or D.5 in the first excited state.
- (5) For $d = -86\,139\,199$, the erroneous IPAD $\tau^{(1)}(K) = [(9, 3); (81, 3, 3, 3), (9, 3, 3), (27, 3)^2]$ contained the malformed component $(81, 3, 3, 3)$ instead of the correct $(9, 3, 3)$. This is clearly a pTKT D.11.

Again, none of the corresponding discriminants

$$d \in \{-70\,204\,919, -74\,771\,240, -79\,749\,087, -86\,139\,199, -94\,304\,231\}$$

is particularly spectacular.

We emphasize that, in both Examples 5.1 and 5.2, the errors of PARI/GP [33] occurred in the upper limit range of absolute discriminants above 70 millions. This seems to be a critical region of extreme computing where current computational algebra systems become unstable. MAGMA [23] also often fails to compute the TKT in that range.

Fortunately, there appeared a single discriminant only for each of the 5 erroneous IPADs, in both examples. This indicates that the errors are not systematic but rather stochastic.

5.2. Application 2: Completing partial capitulation types.

Example 5.3. For the discriminant $d = -3\,849\,267$ of the complex quadratic field $K = \mathbb{Q}(\sqrt{d})$ with 3-class group of type $(3, 3)$, we constructed the four unramified cyclic cubic extensions $L_i|K$, $1 \leq i \leq 4$, and computed the IPAD $\tau^{(1)}(K) = [1^2; (54, 21, 1^3, 21)]$ with the aid of MAGMA [23].

According to Theorem 3.2, the second 3-class group G of K must be of coclass $\text{cc}(G) = 2$, and the polarized component 54 of the IPAD shows that $c - k = 5 + 4 = 9$ and thus the nilpotency class $c = \text{cl}(G)$ and the defect of commutativity k are given by either $c = 9$, $k = 0$, or $c = 10$, $k = 1$. Further, in view of the rank-3 component 1^3 of the IPAD, G must be a vertex of the coclass tree $\mathcal{T}^2(\langle 729, 49 \rangle)$.

When we tried to determine the 3-principalization type $\varkappa := \varkappa_1(3, K)$, MAGMA succeeded in calculating $\varkappa(1) = 3$ and $\varkappa(2) = 3$ but unfortunately failed to give $\varkappa(3)$ and $\varkappa(4)$. With respect to the complete IPAD, Theorem 4.1 enforces $\varkappa(3) = 1$ (item (1)) and $\varkappa(4) = 3$ (item (2)), and therefore the partial result $\varkappa = (3, 3, *, *)$ is completed to $\varkappa = (3, 3, 1, 3)$. According to item (3) of Remark 4.2 or item (3) of Corollary 4.1, K is of TKT H.4. Our experience suggests that this TKT compels the arrangement $c = 10$, $k = 1$, expressed by the *weak leaf conjecture* [27, Cnj.3.1, p.423].

Example 5.4. For the discriminant $d = -4\,928\,155$ of the complex quadratic field $K = \mathbb{Q}(\sqrt{d})$ with 3-class group of type $(3, 3)$, we constructed the four unramified cyclic cubic extensions $L_i|K$, $1 \leq i \leq 4$, and computed the IPAD $\tau^{(1)}(K) = [1^2; (21, 54, (21)^2)]$ with the aid of MAGMA [23].

According to Theorem 3.2, the second 3-class group G of K must be of coclass $\text{cc}(G) = 2$, and the polarized component 54 of the IPAD shows that $c - k = 5 + 4 = 9$ and thus the nilpotency class $c = \text{cl}(G)$ and the defect of commutativity k are given by either $c = 9$, $k = 0$, or $c = 10$, $k = 1$. Further, due to the lack of a rank-3 component 1^3 in the IPAD, G must be a vertex of the coclass tree $\mathcal{T}^2(\langle 729, 54 \rangle)$.

Next, we tried to determine the 3-principalization type $\varkappa := \varkappa_1(3, K)$. MAGMA succeeded in calculating two fixed points $\varkappa(1) = 1$ and $\varkappa(2) = 2$ but unfortunately failed to give $\varkappa(3)$ and $\varkappa(4)$. With respect to the complete IPAD, Theorem 4.2 enforces $\varkappa(3) = 3$ or $\varkappa(4) = 4$ (item (1)), and $\varkappa(4) = 2$ or $\varkappa(3) = 2$ (item (2)), and therefore the partial result $\varkappa = (1, 2, *, *)$ is completed to $\varkappa = (1, 2, 3, 2)$ or $\varkappa = (1, 2, 2, 4)$. According to item (1) of Remark 4.4 or item (3) of Corollary 4.2, K is of TKT E.8, and this TKT enforces the arrangement $c = 9$, $k = 0$, since $k = 1$ is impossible.

Example 5.5. For the discriminant $d = -65\,433\,643$ of the complex quadratic field $K = \mathbb{Q}(\sqrt{d})$ with 3-class group of type $(3, 3)$, we constructed the four unramified cyclic cubic extensions $L_i|K$, $1 \leq i \leq 4$, and computed the IPAD $\tau^{(1)}(K) = [1^2; (65, 1^3, (21)^2)]$ with the aid of MAGMA [23].

According to Theorem 3.2, the second 3-class group G of K must be of coclass $\text{cc}(G) = 2$, and the polarized component 65 of the IPAD shows that $c - k = 6 + 5 = 11$ and thus the nilpotency class $c = \text{cl}(G)$ and the defect of commutativity k are given by either $c = 11$, $k = 0$, or $c = 12$, $k = 1$. Further, in view of the rank-3 component 1^3 of the IPAD, G must be a vertex of the coclass tree $\mathcal{T}^2(\langle 729, 49 \rangle)$.

Then we tried to determine the 3-principalization type $\varkappa := \varkappa_1(3, K)$. MAGMA succeeded in calculating $\varkappa(1) = 4$ and $\varkappa(2) = 1$ but unfortunately failed to give $\varkappa(3)$ and $\varkappa(4)$. With respect to the complete IPAD, Theorem 4.1 enforces $\varkappa(3) = 2$ and $\varkappa(4) = 2$ (item (2)), whereas the claim in item (1) is confirmed, and therefore the partial result $\varkappa = (4, 1, *, *)$ is completed to $\varkappa = (4, 1, 2, 2)$. According to item (2) of Remark 4.2 or item (3) of Corollary 4.1, K is of TKT E.14, and this TKT enforces the arrangement $c = 11$, $k = 0$, since $k = 1$ is impossible.

6. ITERATED IPADS OF SECOND ORDER

6.1. p -capitulation type. By means of the following theorem, the exact 3-principalization type \varkappa of real quadratic fields $K = \mathbb{Q}(\sqrt{d})$, $d > 0$, can be determined indirectly with the aid of information on the structure of 3-class groups of number fields of absolute degree $6 \cdot 3 = 18$.

Theorem 6.1. (*Indirect computation of the p -capitulation type*)

Suppose that $p = 3$ and let K be a number field with 3-class group $\text{Cl}_3(K)$ of type $(3, 3)$ and 3-tower group G .

(1) If the IPAD of K is given by

$$\tau^{(1)}(K) = [1^2; (21; (1^2)^3)],$$

then

$$G'' = 1, \quad G \simeq G/G'', \quad \text{cc}(G) = 1, \quad \text{and } G \in \{\langle 81, 8 \rangle, \langle 81, 9 \rangle, \langle 81, 10 \rangle\},$$

in particular, the length of the 3-class tower of K is given by $\ell_3(K) = 2$.

(2) If the first layer $\text{Ly}_{\Gamma_1}(K)$ of abelian unramified extensions of K consists of L_1, \dots, L_4 , then the iterated IPAD of second order

$$\tau^{(2)}(K) = [\tau_0(K); (\tau_0(L_i); \tau_1(L_i))_{1 \leq i \leq 4}], \quad \text{with } \tau_0(K) = 1^2,$$

admits a sharp decision about the group G and the first layer of the transfer kernel type

$$\varkappa(K) = [\varkappa_0(K); \varkappa_1(K); \varkappa_2(K); \dots] \quad \text{where trivially } \varkappa_0(K) = 1, \quad \varkappa_2(K) = 0.$$

$$(6.1) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1)] &= [21; (1^2, (2)^3)], \\ [\tau_0(L_i); \tau_1(L_i)] &= [1^2; (1^2, (2)^3)], \quad \text{for } 2 \leq i \leq 4, \end{aligned}$$

implies $G \simeq \langle 81, 10 \rangle$ and thus $\varkappa_1(K) = (1, 0, 0, 0)$,

$$(6.2) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1)] &= [21; (1^2, (2)^3)], \\ [\tau_0(L_2); \tau_1(L_2)] &= [1^2; (1^2)^4], \\ [\tau_0(L_i); \tau_1(L_i)] &= [1^2; (1^2, (2)^3)], \quad \text{for } 3 \leq i \leq 4, \end{aligned}$$

implies $G \simeq \langle 81, 8 \rangle$ and thus $\varkappa_1(K) = (2, 0, 0, 0)$, and

$$(6.3) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1)] &= [21; (1^2, (2)^3)], \\ [\tau_0(L_i); \tau_1(L_i)] &= [1^2; (1^2)^4], \quad \text{for } 2 \leq i \leq 4, \end{aligned}$$

implies $G \simeq \langle 81, 9 \rangle$ and thus $\varkappa_1(K) = (0, 0, 0, 0)$.

Example 6.1. A possible future application of Theorem 6.1 could, for instance, be the separation of the capitulation types a.2, $\varkappa_1(K) = (1, 0, 0, 0)$, and a.3, $\varkappa_1(K) = (2, 0, 0, 0)$, among the 1386 real quadratic fields $K = \mathbb{Q}(\sqrt{d})$, $0 < d < 10^7$, with 3-class group $\text{Cl}_3(K)$ of type $(3, 3)$ and IPAD $\tau^{(1)}(K) = [1^2; (21; (1^2)^3)]$, which was outside of our reach in all investigations of [25, Tbl.2, p.496], [28, Tbl.6.1, p.451] and [27, Fig.3.2, p.422]. The reason why we expect this enterprise to be promising is that our experience with Magma [23] shows that computing class groups can become slow but remains sound and stable for huge discriminants d , whereas the calculation of capitulation kernels frequently fails.

6.2. Length of the p -class tower. In this section, we use the iterated IPAD of second order $\tau^{(2)}(K) = [\tau_0(K); (\tau_0(L); \tau_1(L))_{L \in \text{Ly}_{\Gamma_1}(K)}]$ for the indirect computation of the length $\ell_p(K)$ of the p -class tower of a number field K , where p denotes a fixed prime.

Theorem 6.2. (*Length $\ell_3(K)$ of the 3-class tower for $G/G'' \in \mathcal{T}^2(\langle 243, 6 \rangle)$*)

Suppose that $p = 3$ and let K be a number field with 3-class group $\text{Cl}_3(K)$ of type $(3, 3)$ and 3-tower group G .

- (1) *If the IPAD of K is given by*

$$\tau^{(1)}(K) = [1^2; (32; 1^3, (21)^2)],$$

and the first layer TKT $\varkappa_1(K)$ neither contains a total principalization nor a 2-cycle, then there are two possibilities $\ell_3(K) \in \{2, 3\}$ for the length of the 3-class tower of K .

- (2) *If the first layer $\text{Ly}_{\Gamma_1}(K)$ of abelian unramified extensions of K consists of L_1, \dots, L_4 , then the iterated IPAD of second order*

$$\tau^{(2)}(K) = [\tau_0(K); (\tau_0(L_i); \tau_1(L_i))_{1 \leq i \leq 4}], \text{ with } \tau_0(K) = 1^2,$$

admits a sharp decision about the length $\ell_3(K)$:

$$(6.4) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1)] &= [32; (2^2 1, (31^2)^3)], \\ [\tau_0(L_2); \tau_1(L_2)] &= [1^3; (2^2 1, (\mathbf{1}^3)^3, (1^2)^9)], \\ [\tau_0(L_i); \tau_1(L_i)] &= [21; (2^2 1, (\mathbf{21})^3)], \text{ for } 3 \leq i \leq 4, \end{aligned}$$

if and only if $\ell_3(K) = 2$, and

$$(6.5) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1)] &= [32; (2^2 1, (31^2)^3)], \\ [\tau_0(L_2); \tau_1(L_2)] &= [1^3; (2^2 1, (\mathbf{21}^2)^3, (1^2)^9)], \\ [\tau_0(L_i); \tau_1(L_i)] &= [21; (2^2 1, (\mathbf{31})^3)], \text{ for } 3 \leq i \leq 4, \end{aligned}$$

if and only if $\ell_3(K) = 3$.

Proof. According to Theorem 3.2, an IPAD of the form $\tau^{(1)}(K) = [1^2; (32; 1^3, (21)^2)]$ indicates that the metabelianization of the group G belongs to the coclass tree $\mathcal{T}^2(\langle 243, 6 \rangle)$ [27, Fig.3.6, p.442] and has nilpotency class $3 + 2 = 5$, due to the polarization.

According to § 4.1, the lack of a total principalization excludes the TKT c.18 and the absence of a 2-cycle discourages the TKT H.4, whence the group G must be of TKT E.6 or E.14.

By means of the techniques described in [13], a search in the complete descendant tree $\mathcal{T}(\langle 243, 6 \rangle)$, not restricted to groups of coclass 2, yields exactly six candidates for the group G : three metabelian groups $\langle 2187, i \rangle$ with $i \in \{288, 289, 290\}$, and three groups of derived length 3 and order 3^8 with generalized identifiers $\langle 729, 49 \rangle - \#2; i, i \in \{4, 5, 6\}$. There cannot exist adequate groups of bigger orders. The former three groups are characterized by Equations (6.4) the latter three groups (see [29, § 20.2, Fig.8]) by Equations (6.5).

Finally, we have $\ell_3(K) = \text{dl}(G)$. □

Theorem 6.3. (*Length $\ell_p(K)$ of the 3-class tower for $G/G'' \in \mathcal{T}^2(\langle 243, 8 \rangle)$*)

Suppose that $p = 3$ and let K be a number field with 3-class group $\text{Cl}_3(K)$ of type $(3, 3)$ and 3-tower group G .

(1) If the IPAD of K is given by

$$\tau^{(1)}(K) = [1^2; (32; (21)^3)],$$

and the first layer TKT $\varkappa_1(K)$ neither contains a total principalization nor a 2-cycle, then there are two possibilities $\ell_3(K) \in \{2, 3\}$ for the length of the 3-class tower of K .

(2) If the first layer $\text{Ly}\Gamma_1(K)$ of abelian unramified extensions of K consists of L_1, \dots, L_4 , then the iterated IPAD of second order

$$\tau^{(2)}(K) = [\tau_0(K); (\tau_0(L_i); \tau_1(L_i))_{1 \leq i \leq 4}], \text{ with } \tau_0(K) = 1^2,$$

admits a sharp decision about the length $\ell_3(K)$:

$$(6.6) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1)] &= [32; (2^2 1, (31^2)^3)], \\ [\tau_0(L_i); \tau_1(L_i)] &= [21; (2^2 1, (\mathbf{21})^3)], \text{ for } 2 \leq i \leq 4, \end{aligned}$$

if and only if $\ell_3(K) = 2$, and

$$(6.7) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1)] &= [32; (2^2 1, (31^2)^3)], \\ [\tau_0(L_i); \tau_1(L_i)] &= [21; (2^2 1, (\mathbf{31})^3)], \text{ for } 2 \leq i \leq 4, \end{aligned}$$

if and only if $\ell_3(K) = 3$.

Proof. According to Theorem 3.2, an IPAD of the form $\tau^{(1)}(K) = [1^2; (32; (21)^3)]$ indicates that the metabelianization of the group G belongs to the coclass tree $\mathcal{T}^2(\langle 243, 8 \rangle)$ [27, Fig.3.7, p.443] and has nilpotency class $3 + 2 = 5$, due to the polarization.

According to § 4.2, the lack of a total principalization excludes the TKT c.21 and the absence of a 2-cycle discourages the TKT G.16, whence the group G must be of TKT E.8 or E.9.

As we have shown in detail in [13], a search in the complete descendant tree $\mathcal{T}(\langle 243, 8 \rangle)$, not restricted to groups of coclass 2, yields exactly six candidates for the group G : three metabelian groups $\langle 2187, i \rangle$ with $i \in \{302, 304, 306\}$, and three groups of derived length 3 and order 3^8 with generalized identifiers $\langle 729, 54 \rangle - \#2; i, i \in \{2, 4, 6\}$. There cannot exist adequate groups of bigger orders. The former three groups are characterized by Equations (6.6) the latter three groups (see [29, § 20.2, Fig.9]) by Equations (6.7).

Eventually, the 3-tower length of K , $\ell_3(K) = \text{dl}(G)$, coincides with the derived length of G . \square

Example 6.2. In June 2006, we discovered the smallest discriminant $d = 342\,664$ of a real quadratic field $K = \mathbb{Q}(\sqrt{d})$ with 3-class group of type $(3, 3)$ whose 3-tower group G possesses the transfer kernel type E.9, $\varkappa = (2, 3, 3, 4)$.

The complex quadratic analogue $k = \mathbb{Q}(\sqrt{-9\,748})$ was known since 1934 by the famous paper of Scholz and Taussky [34]. However, it required almost 80 years until M.R. Bush and ourselves [13] succeeded in providing the first faultless proof that k has a 3-class tower of exact length $\ell_3(k) = 3$ with 3-tower group G one of the two Schur σ -groups $\langle 729, 54 \rangle - \#2; i, i \in \{2, 6\}$, of order 3^8 .

For $K = \mathbb{Q}(\sqrt{342\,664})$, the methods in [13] do not admit a final decision about the length $\ell_3(K)$. They only yield four possible 3-tower groups of K , namely either the two unbalanced groups $\langle 2187, i \rangle$ with $i \in \{302, 306\}$ and relation rank $r = 3$ bigger than the generator rank $d = 2$ or the two Schur σ -groups $\langle 729, 54 \rangle - \#2; i$ with $i \in \{2, 6\}$ and $r = 2$ equal to $d = 2$.

In October 2014, we succeeded in proving that three of the unramified cyclic cubic extensions $L_i|K$ reveal the critical IPAD component $\tau_1(L_i) = (2^2 1, (31)^3)$ in Equation (6.7) of Theorem 6.3, item (2), whence $\ell_3(K) = 3$. This was done by computing 3-class groups of number fields of absolute degree $6 \cdot 3 = 18$ with the aid of MAGMA [23].

L. Bartholdi and M.R. Bush [3] have shown that the unbalanced metabelian 3-group $G = \langle 729, 45 \rangle$ possesses an infinite balanced cover $\text{cov}_*(G)$ [29, Dfn.21.2], which implies that the length $\ell_3(K)$ of the 3-class tower of a complex quadratic field K with IPAD $\tau(K) = [1^2; ((1^3)^3, 21)]$ can take any value bigger than 2 or even ∞ . The group theoretic reason for this remarkable extravagance is that G is not coclass-settled and gives rise to a descendant tree $\mathcal{T}(G)$ which contains infinitely many periodic bifurcations [29, § 21].

As a final coronation of this section, we show that our new IPAD strategies are powerful enough to enable the determination of the length $\ell_3(K)$ with the aid of information on the structure of 3-class groups of number fields of absolute degree $6 \cdot 9 = 54$.

For this purpose, we extend the concept of iterated IPADs of second order

$$\tau^{(1)}(K) = [\tau_0(K); (\tau^{(1)}(L))_{L \in \text{Ly}_{\Gamma_1}(K)}] = [\tau_0(K); (\tau_0(L); \tau_1(L))_{L \in \text{Ly}_{\Gamma_1}(K)}]$$

once more by adding the second layers $\tau_2(L)$ of all IPADs $\tau^{(1)}(L)$ of unramified degree- p extensions $L|K$. The resulting *iterated multi-layered IPAD of second order* is indicated by an asterisk

$$\tau_*^{(1)}(K) = [\tau_0(K); (\tau_0(L); \tau_1(L); \tau_2(L))_{L \in \text{Ly}_{\Gamma_1}(K)}].$$

Theorem 6.4. (*Length $\ell_p(K)$ of the 3-class tower for $G/G'' \in \mathcal{T}(\langle 243, 4 \rangle)$)*)

Suppose that $p = 3$ and let K be a number field with 3-class group $\text{Cl}_3(K)$ of type $(3, 3)$ and 3-tower group G .

(1) *If the IPAD of K is given by*

$$\tau^{(1)}(K) = [1^2; ((1^3)^3, 21)],$$

then the first layer TKT is $\varkappa_1(K) = (4, 1, 1, 1)$ and there exist infinitely many possibilities $\ell_3(K) \geq 2$ for the length of the 3-class tower of K .

(2) *If the first layer $\text{Ly}_{\Gamma_1}(K)$ of abelian unramified extensions of K consists of L_1, \dots, L_4 , then the iterated multi-layered IPAD of second order*

$$\tau_*^{(2)}(K) = [\tau_0(K); (\tau_0(L_i); \tau_1(L_i); \tau_2(L_i))_{1 \leq i \leq 4}], \text{ with } \tau_0(K) = 1^2,$$

admits certain partial decisions about the length $\ell_3(K)$:

$$(6.8) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1); \tau_2(L_1)] &= [1^3; ((1^3)^4, (1^2)^9); (1^2)^{13}], \\ [\tau_0(L_i); \tau_1(L_i); \tau_2(L_i)] &= [1^3; (1^3, (21)^3, (1^2)^9); ((1^2)^4, (2)^9)], \text{ for } 2 \leq i \leq 3, \\ [\tau_0(L_4); \tau_1(L_4); \tau_2(L_4)] &= [21; (1^3, (21)^3); (1^2)^4] \end{aligned}$$

implies $G \simeq \langle 243, 4 \rangle$ and $\ell_3(K) = 2$.

$$(6.9) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1); \tau_2(L_1)] &= [1^3; (\mathbf{21}^2, (1^3)^3, (1^2)^9); (1^3, (21)^3, (1^2)^9)], \\ [\tau_0(L_i); \tau_1(L_i); \tau_2(L_i)] &= [1^3; (\mathbf{21}^2, (\mathbf{21})^{12}); (1^3, (21)^{12})], \text{ for } 2 \leq i \leq 3, \\ [\tau_0(L_4); \tau_1(L_4); \tau_2(L_4)] &= [21; (\mathbf{21}^2, (21)^3); (1^3, (21)^3)] \end{aligned}$$

implies $G \simeq \langle 729, 45 \rangle$ and $\ell_3(K) = 2$.

$$(6.10) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1); \tau_2(L_1)] &= [1^3; (21^2, (1^3)^3, (1^2)^9); (21^2, (21)^3, (1^2)^9)], \\ [\tau_0(L_2); \tau_1(L_2); \tau_2(L_2)] &= [1^3; (21^2, (21)^{12}); (21^2, (21)^{12})], \\ [\tau_0(L_3); \tau_1(L_3); \tau_2(L_3)] &= [1^3; ((\mathbf{21}^2)^4, (\mathbf{2}^2)^9); (21^2)^{13}], \\ [\tau_0(L_4); \tau_1(L_4); \tau_2(L_4)] &= [21; (21^2, (21)^3); (21^2, (21)^3)] \end{aligned}$$

implies $G \simeq \langle 2187, 273 \rangle$ and $\ell_3(K) = 3$.

$$(6.11) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1); \tau_2(L_1)] &= [1^3; ((21^2)^4, (1^2)^9); (2^2 1, (1^3)^3, (\mathbf{2}^2)^3, (21)^6)], \\ [\tau_0(L_i); \tau_1(L_i); \tau_2(L_i)] &= [1^3; ((21^2)^4, (2^2)^9); (2^2 1, (\mathbf{21}^2)^{12})], \text{ for } 2 \leq i \leq 3, \\ [\tau_0(L_4); \tau_1(L_4); \tau_2(L_4)] &= [21; (21^2, (31)^3); (2^2 1, (\mathbf{2}^2)^3)] \end{aligned}$$

implies $G \simeq \langle 729, 45 \rangle - \#2; 2$ of order 3^8 and $\ell_3(K) = 3$.

$$(6.12) \quad \begin{aligned} [\tau_0(L_1); \tau_1(L_1); \tau_2(L_1)] &= [1^3; ((21^2)^4, (1^2)^9); (2^2 1, (1^3)^3, (\mathbf{32})^3, (21)^6)], \\ [\tau_0(L_i); \tau_1(L_i); \tau_2(L_i)] &= [1^3; ((21^2)^4, (2^2)^9); ((\mathbf{2}^2 \mathbf{1})^4, (\mathbf{31}^2)^9)], \text{ for } 2 \leq i \leq 3, \\ [\tau_0(L_4); \tau_1(L_4); \tau_2(L_4)] &= [21; (21^2, (31)^3); (2^2 1, (\mathbf{32})^3)] \end{aligned}$$

implies either

$G \simeq \langle 729, 45 \rangle (-\#2; 1 - \#1; 2)^j - \#2; 2, 1 \leq j \leq 2$, of order 3^{8+3j} and $\ell_3(K) = 3$
or $G \simeq \langle 729, 45 \rangle (-\#2; 1 - \#1; 2)^3 - \#2; 2$ of order 3^{17} and $\ell_3(K) = 4$.

Example 6.3. In December 2009, we discovered the smallest discriminant $d = 957\,013$ of a real quadratic field $K = \mathbb{Q}(\sqrt{d})$ with 3-class group of type $(3, 3)$ whose 3-tower group G possesses the transfer kernel type H.4, $\varkappa = (4, 1, 1, 1)$. The complex quadratic analogue $K = \mathbb{Q}(\sqrt{-3\,896})$ was known since 1982 by the paper of Heider and Schmithals [20]. Both fields share the same IPAD $\tau^{(1)}(K) = [1^2; ((1^3)^3, 21)]$.

In February 2015, we succeeded in proving that the unramified cyclic cubic extensions $L_i|K$, for $d = -3\,896$, resp. $d = 957\,013$, reveal the critical (first and) second layer IPAD components

$$\begin{aligned} \tau_2(L_1) &= (2^2 1, (1^3)^3, (\mathbf{2}^2)^3, (21)^6), \\ \tau_2(L_i) &= (2^2 1, (\mathbf{21}^2)^{12}), \text{ for } 2 \leq i \leq 3, \text{ and} \\ \tau_2(L_4) &= (2^2 1, (\mathbf{2}^2)^3) \end{aligned}$$

in Equation (6.11), resp.

$$\begin{aligned} [\tau_1(L_1); \tau_2(L_1)] &= [(21^2, (1^3)^3, (1^2)^9); (21^2, (21)^3, (1^2)^9)], \\ [\tau_1(L_2); \tau_2(L_2)] &= [(21^2, (21)^{12}); (21^2, (21)^{12})], \\ [\tau_1(L_3); \tau_2(L_3)] &= [((\mathbf{21}^2)^4, (\mathbf{2}^2)^9); (21^2)^{13}], \\ [\tau_1(L_4); \tau_2(L_4)] &= [(21^2, (21)^3); (21^2, (21)^3)] \end{aligned}$$

in Equation (6.10), of Theorem 6.4, item (2), whence $\ell_3(K) = 3$, for both fields.

However, the 3-class tower groups are different:

$K = \mathbb{Q}(\sqrt{-3\,896})$ has the Schur σ -group $G \simeq \langle 729, 45 \rangle - \#2; 2$ of order 3^8 , and
 $K = \mathbb{Q}(\sqrt{957\,013})$ has the unbalanced group $G \simeq \langle 2187, 273 \rangle$.

This was done by computing 3-class groups of number fields of absolute degree $6 \cdot 9 = 54$ with the aid of MAGMA [23].

7. COMPLEX QUADRATIC FIELDS OF 3-RANK THREE

In this concluding section we present another impressive application of IPADs.

Due to Koch and Venkov [22], it is known that a complex quadratic field K with 3-class rank $r_3(K) \geq 3$ has an infinite 3-class field tower $K < F_3^1(K) < F_3^2(K) < \dots < F_3^\infty(K)$ of length $\ell_3(K) = \infty$. In the time between 1973 and 1978, Diaz y Diaz [14, 15] and Buell [11] have determined the smallest absolute discriminants $|d|$ of such fields. Recently, we have launched a computational project which aims at verifying these classical results and adding sophisticated arithmetical details. Below the bound 10^7 there exist 25 discriminants d of this kind, and 14 of the corresponding fields K have a 3-class group $\text{Cl}_3(K)$ of elementary abelian type $(3, 3, 3)$. For each of these 14 fields, we determine the type of 3-principalization $\varkappa := \varkappa_1(3, K)$ in the thirteen unramified cyclic cubic extensions L_1, \dots, L_{13} of K , and the structure of the 3-class groups $\text{Cl}_3(L_i)$ of these extensions, i.e., the IPAD of K . We characterize the metabelian Galois group $G = G_3^2(K) = \text{Gal}(F_3^2(K)|K)$ of the second Hilbert 3-class field $F_3^2(K)$ by means of kernels and targets of its Artin transfer homomorphisms [2] to maximal subgroups. We provide evidence of a wealth of structure in the set of infinite topological 3-class field tower groups $G_3^\infty(K) = \text{Gal}(F_3^\infty(K)|K)$ by showing that the 14 groups G are pairwise non-isomorphic.

We summarize our results and their obvious conclusion in the following theorem.

Theorem 7.1. *There exist exactly 14 complex quadratic number fields $K = \mathbb{Q}(\sqrt{d})$ with 3-class groups $\text{Cl}_3(K)$ of type $(3, 3, 3)$ and discriminants in the range $-10^7 < d < 0$. They have pairwise non-isomorphic*

- (1) *second and higher 3-class groups $\text{Gal}(F_3^n(K)|K)$, $n \geq 2$,*
- (2) *infinite topological 3-class field tower groups $\text{Gal}(F_3^\infty(K)|K)$.*

TABLE 1. Data collection for $\text{Cl}_3(K) \simeq (3, 3, 3)$ and $-10^7 < d$

No.	discriminant d	$\text{Cl}_3(K)$	$\text{Cl}(K)$
1	-3 321 607	(9, 3, 3)	(63, 3, 3)
2	-3 640 387	(9, 3, 3)	(18, 3, 3)
3	-4 019 207	(9, 3, 3)	(207, 3, 3)
4	-4 447 704	(3, 3, 3)	(24, 6, 6)
5	-4 472 360	(3, 3, 3)	(30, 6, 6)
6	-4 818 916	(3, 3, 3)	(48, 3, 3)
7	-4 897 363	(3, 3, 3)	(33, 3, 3)
8	-5 048 347	(9, 3, 3)	(18, 6, 3)
9	-5 067 967	(3, 3, 3)	(69, 3, 3)
10	-5 153 431	(27, 3, 3)	(216, 3, 3)
11	-5 288 968	(9, 3, 3)	(72, 3, 3)
12	-5 769 988	(3, 3, 3)	(12, 6, 6)
13	-6 562 327	(9, 3, 3)	(126, 3, 3)
14	-7 016 747	(9, 3, 3)	(99, 3, 3)
15	-7 060 148	(3, 3, 3)	(60, 6, 3)
16	-7 503 391	(9, 3, 3)	(90, 6, 3)
17	-7 546 164	(9, 3, 3)	(18, 6, 6, 2)
18	-8 124 503	(9, 3, 3)	(261, 3, 3)
19	-8 180 671	(3, 3, 3)	(159, 3, 3)
20	-8 721 735	(3, 3, 3)	(60, 6, 6)
21	-8 819 519	(3, 3, 3)	(276, 3, 3)
22	-8 992 363	(3, 3, 3)	(48, 3, 3)
23	-9 379 703	(3, 3, 3)	(210, 3, 3)
24	-9 487 991	(3, 3, 3)	(381, 3, 3)
25	-9 778 603	(3, 3, 3)	(48, 3, 3)

Before we come to the proof of Theorem 7.1 in § 7.3, we collect basic numerical data concerning fields with $r_3(K) = 3$ in § 7.1, and we completely determine sophisticated arithmetical invariants in § 7.2 for all fields with $\text{Cl}_3(K)$ of type $(3, 3, 3)$. The first attempt to do so for the smallest absolute discriminant $|d| = 3\,321\,607$ with $r_3(K) = 3$ is due to Heider and Schmithals in [20, § 4, Tbl.2, p.18], but it resulted in partial success only.

7.1. Discriminants $-10^7 < d < 0$ of fields $K = \mathbb{Q}(\sqrt{d})$ with rank $r_3(K) = 3$. Since one of our aims is to investigate tendencies for the coclass of second and higher p -class groups $G_p^n(K) = \text{Gal}(\mathbb{F}_p^n(K)|K)$, $n \geq 2$, [25, 27] of a series of algebraic number fields K with infinite p -class field tower, for an odd prime $p \geq 3$, the most obvious choice which suggests itself is to take the smallest possible prime $p = 3$ and to select complex quadratic fields $K = \mathbb{Q}(\sqrt{d})$, $d < 0$, having the simplest possible 3-class group $\text{Cl}_3(K)$ of rank 3, that is, of elementary abelian type $(3, 3, 3)$.

The reason is that Koch and Venkov [22] have improved the lower bound of Golod, Shafarevich [35, 19] and Vinberg [39] for the p -class rank $r_p(K)$, which ensures an infinite p -class tower of a complex quadratic field K , from 4 to 3.

However, quadratic fields with 3-rank $r_3(K) = 3$ are sparse. Diaz y Diaz and Buell [14, 36, 11, 15] have determined the minimal absolute discriminant of such fields to be $|d| = 3\,321\,607$.

To provide an independent verification, we used the computational algebra system Magma [6, 7, 23] for compiling a list of all quadratic fundamental discriminants $-10^7 < d < 0$ of fields $K = \mathbb{Q}(\sqrt{d})$ with 3-class rank $r_3(K) = 3$. In 16 hours of CPU time we obtained the 25 desired discriminants and the abelian type invariants (here written in 3-power form) of the corresponding 3-class groups $\text{Cl}_3(K)$, and also of the complete class groups $\text{Cl}(K)$, as given in Table 1. There

appeared only one discriminant $d = -7503391$ (No. 16) which is not contained in [15, Appendix 1, p.68] already.

There are 14 discriminants, starting with $d = -4447704$, such that $\text{Cl}_3(K)$ is elementary abelian of type $(3, 3, 3)$, and 10 discriminants, starting with -3321607 , such that $\text{Cl}_3(K)$ is of non-elementary type $(9, 3, 3)$. For the single discriminant $d = -5153431$, we have a 3-class group of type $(27, 3, 3)$. We have published this information in the Online Encyclopedia of Integer Sequences (OEIS) [37], sequences A244574 and A244575.

TABLE 2. Pattern recognition via ordered IPADs

No.	discriminant												
i	1	2	3	4	5	6	7	8	9	10	11	12	13
1	$d = -4447704$												
\varkappa	8	1	8	8	10	8	6	13	8	2	10	8	9
$o(\varkappa)$	1	1	0	0	0	1	0	6	1	2	0	0	1
τ	$2^2 1^2$	21^4	$2^2 1^2$	$32^2 1$	$2^2 1^2$	21^4	$2^2 1^2$	$2^2 1^2$	21^4	21^4	21^4	$2^2 1^2$	$2^2 1^2$
τ^0	1^2	1^2	1^2	21	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2
2	$d = -4472360$												
\varkappa	1	12	6	13	6	10	4	13	10	10	1	8	4
$o(\varkappa)$	2	0	0	2	0	2	0	1	0	3	0	1	2
τ	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	21^4	$2^2 1^2$	21^4	21^4	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	$32^2 1$	$2^2 1^2$	21^4
τ^0	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	21	1^2	1^2
3	$d = -4818916$												
\varkappa	6	9	13	1	5	6	9	4	11	7	1	3	4
$o(\varkappa)$	2	0	1	2	1	2	1	0	2	0	1	0	1
τ	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	21^4	21^4	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	431^3	$2^2 1^2$	21^4	$32^2 1$	$2^2 1^2$
τ^0	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	31	1^2	1^2	21	1^2
4	$d = -4897363$												
\varkappa	3	8	11	2	6	6	12	7	2	2	9	13	6
$o(\varkappa)$	0	3	1	0	0	3	1	1	1	0	1	1	1
τ	$2^2 1^2$	321^3	431^3	$2^2 1^2$	21^4	$2^2 1^2$	$32^2 1$	21^4	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$
τ^0	1^2	21	31	1^2	1^2	1^2	21	1^2	1^2	1^2	1^2	1^2	1^2
5	$d = -5067967$												
\varkappa	8	6	9	2	3	7	12	7	1	4	3	9	4
$o(\varkappa)$	1	1	2	2	0	1	2	1	2	0	0	1	0
τ	21^4	21^4	$2^2 1^2$	21^4	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	$32^2 1$	$2^2 1^2$	21^4	21^4	$2^2 1^2$	$2^2 1^2$
τ^0	1^2	1^2	1^2	1^2	1^2	1^2	1^2	21	1^2	1^2	1^2	1^2	1^2
6	$d = -5769988$												
\varkappa	12	11	7	6	1	1	10	10	9	6	4	3	13
$o(\varkappa)$	2	0	1	1	0	2	1	0	1	2	1	1	1
τ	321^3	$2^2 1^2$	21^4	321^3	$2^2 1^2$	21^4	21^4	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	$2^2 1^2$	21^4	$32^2 1$
τ^0	21	1^2	1^2	21	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	21

7.2. Arithmetic invariants of fields $K = \mathbb{Q}(\sqrt{d})$ with $\text{Cl}_3(K) \simeq (3, 3, 3)$. After the preliminary data collection in section § 7.1, we restrict ourselves to the 14 cases with elementary abelian 3-class group of type $(3, 3, 3)$. The complex quadratic field $K = \mathbb{Q}(\sqrt{d})$ possesses 13 unramified cyclic cubic extensions L_1, \dots, L_{13} with dihedral absolute Galois group $\text{Gal}(L_i|\mathbb{Q})$ of order six [25]. Based on Fieker's technique [16], we use the computational algebra system Magma [7, 23] to construct these extensions and to calculate their arithmetical invariants. In Table 2, which is continued in Table 3 on the following page, we present the kernel \varkappa_i of the 3-principalization of K in

TABLE 3. Pattern recognition (continued)

No.	discriminant												
i	1	2	3	4	5	6	7	8	9	10	11	12	13
7	$d = -7060148$												
\varkappa	2	4	4	9	4	10	8	10	10	1	6	8	3
$o(\varkappa)$	1	1	1	3	0	1	0	2	1	3	0	0	0
τ	21^4	21^4	32^21	2^21^2	32^21	21^4	2^21^2	321^3	21^4	21^4	2^21^2	2^21^2	321^3
τ^0	1^2	1^2	21	1^2	21	1^2	1^2	21	1^2	1^2	1^2	1^2	21
8	$d = -8180671$												
\varkappa	12	9	2	6	10	6	8	2	10	10	9	11	4
$o(\varkappa)$	0	2	0	1	0	2	0	1	2	3	1	1	0
τ	321^3	2^21^2	21^4	2^21^2	2^21^2	2^21^2	2^21^2	21^4	2^21^2	2^21^2	2^21^2	21^4	2^21^2
τ^0	21	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2
9	$d = -8721735$												
\varkappa	5	2	5	1	10	13	4	7	11	3	9	8	8
$o(\varkappa)$	1	1	1	1	2	0	1	2	1	1	1	0	1
τ	2^21^2	21^4	21^4	21^4	321^3	2^21^2	21^4	2^21^2	32^21	32^21	21^4	32^21	2^21^2
τ^0	1^2	1^2	1^2	1^2	21	1^2	1^2	1^2	21	21	1^2	21	1^2
10	$d = -8819519$												
\varkappa	2	7	8	12	4	12	9	5	5	3	10	6	10
$o(\varkappa)$	0	1	1	1	2	1	1	1	1	2	0	2	0
τ	2^21^2	21^4	32^21	2^21^2	2^21^2	2^21^2	2^21^2	2^21^2	21^4	2^21^2	2^21^2	2^21^2	1^6
τ^0	1^2	1^2	21	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2
11	$d = -8992363$												
\varkappa	12	10	2	12	9	5	10	10	2	12	6	9	7
$o(\varkappa)$	0	2	0	0	1	1	1	0	2	3	0	3	0
τ	2^21^2	21^4	2^21^2	2^21^2	32^21	2^21^2	21^4	21^4	21^4	2^21^2	2^21^2	21^4	2^21^2
τ^0	1^2	1^2	1^2	1^2	21	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2
12	$d = -9379703$												
\varkappa	8	11	8	13	9	5	6	1	2	13	4	12	3
$o(\varkappa)$	1	1	1	1	1	1	0	2	1	0	1	1	2
τ	21^4	2^21^2	321^3	2^21^2	21^4	21^4	2^21^2	21^4	21^4	2^21^2	2^21^2	2^21^2	2^21^2
τ^0	1^2	1^2	21	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2
13	$d = -9487991$												
\varkappa	4	2	2	11	13	9	12	9	8	1	1	12	3
$o(\varkappa)$	2	2	1	1	0	0	0	1	2	0	1	2	1
τ	2^21^2	2^21^2	2^21^2	2^21^2	321^3	2^21^2	2^21^2	21^4	2^21^2	21^4	2^21^2	2^21^2	2^21^2
τ^0	1^2	1^2	1^2	1^2	21	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2
14	$d = -9778603$												
\varkappa	10	6	6	9	9	10	8	10	13	5	12	6	10
$o(\varkappa)$	0	0	0	0	1	3	0	1	2	4	0	1	1
τ	2^21^2	321^3	21^4	32^21	32^21	2^21^2	2^21^2	2^21^2	21^4	21^4	2^21^2	2^21^2	2^21^2
τ^0	1^2	21	1^2	21	21	1^2	1^2	1^2	1^2	1^2	1^2	1^2	1^2

L_i [24, 25], the occupation numbers $o(\varkappa)_i$ of the principalization kernels [26], and the abelian type invariants τ_i , resp. τ_i^0 , of the 3-class group $\text{Cl}_3(L_i)$, resp. $\text{Cl}_3(K_i)$, for each $1 \leq i \leq 13$ [28, 27]. Here, we denote by K_i the unique real non-Galois absolutely cubic subfield of L_i . For brevity, we

give 3-logarithms of abelian type invariants and we denote iteration by formal exponents. Note that the multiplets \varkappa and τ are *ordered* and in componentwise mutual correspondence, in the sense of § 4.

TABLE 4. Accumulative (unordered) form of IPADs

No.	discriminant d	$2^2 1^2$	$2 1^4$	1^6	$3 2^2 1$	$3 2 1^3$	$4 3 1^3$	polarization	state
1	-4 447 704	7	5	0	1	0	0	uni	ground
2	-4 472 360	8	4	0	1	0	0	uni	ground
3	-4 818 916	8	3	0	1	0	1	bi	excited
4	-4 897 363	8	2	0	1	1	1	tri	excited
5	-5 067 967	7	5	0	1	0	0	uni	ground
6	-5 769 988	6	4	0	1	2	0	tri	ground
7	-7 060 148	4	5	0	2	2	0	tetra	ground
8	-8 180 671	9	3	0	0	1	0	uni	ground
9	-8 721 735	4	5	0	3	1	0	tetra	ground
10	-8 819 519	9	2	1	1	0	0	uni	ground
11	-8 992 363	7	5	0	1	0	0	uni	ground
12	-9 379 703	7	5	0	0	1	0	uni	ground
13	-9 487 991	10	2	0	0	1	0	uni	ground
14	-9 778 603	7	3	0	2	1	0	tri	ground

In Table 4, we classify each of the 14 complex quadratic fields $K = \mathbb{Q}(\sqrt{d})$ of type $(3, 3, 3)$ according to the occupation numbers of the abelian type invariants of the 3-class groups $\text{Cl}_3(L_i)$ of the 13 unramified cyclic cubic extensions L_i , that is the *accumulated* (unordered) form of the IPAD of K . Whereas the dominant part of these groups is of order $3^6 = 729$, there always exist(s) at least one and at most four distinguished groups of bigger order, usually $3^8 = 6\,561$ and occasionally even $3^{10} = 59\,049$. According to the number of distinguished groups, we speak about *uni*-, *bi*-, *tri*- or *tetra*-polarization. If the maximal value of the order is 3^8 , then we have a *ground* state, otherwise an *excited* state.

7.3. Proof of Theorem 7.1.

Proof. According to [27, Thm.1.1 and Dfn.1.1, pp.402–403], the information given in Table 4 consists of isomorphism invariants of the metabelian Galois group $G = \text{Gal}(\mathbb{F}_3^2(K)|K)$ of the second Hilbert 3-class field of K [25]. Consequently, with respect to the 13 abelian type invariants of the 3-class groups $\text{Cl}_3(L_i)$ alone, only the groups G for $d \in \{-4\,447\,704, -5\,067\,967, -8\,992\,363\}$ could be isomorphic. However, Tables 2 and 3 show that these three groups differ with respect to another isomorphism invariant, the 3-principalization type \varkappa [24, 26], since the corresponding maximal occupation numbers of the multiplot $o(\varkappa)$ are 6, 2, 3, respectively. \square

7.4. Final remark. We would like to emphasize that Theorem 7.1 provides evidence for *a wealth of structure* in the set of infinite 3-class field towers, which was unknown up to now, since the common practice is to consider a 3-class field tower as “done” when some criterion in the style of Golod-Shafarevich-Vinberg [35, 19, 39] or Koch-Venkov [22] ensures just its infinity. However, this perspective is very coarse and our result proves that it can be refined considerably.

It would be interesting to extend the range of discriminants $-10^7 < d < 0$ and to find the first examples of isomorphic infinite 3-class field towers.

Another very difficult remaining open problem is the actual identification of the metabelianizations of the 3-tower groups G of the 14 fields. The complexity of this task is due to unmanageable descendant numbers of certain vertices, e.g. $\langle 243, 37 \rangle$ and $\langle 729, 122 \rangle$, in the tree with root $\langle 27, 5 \rangle$.

REFERENCES

- [1] E. Artin, *Beweis des allgemeinen Reziprozitätsgesetzes*, Abh. Math. Sem. Univ. Hamburg **5** (1927), 353–363.
- [2] E. Artin, *Idealklassen in Oberkörpern und allgemeines Reziprozitätsgesetz*, Abh. Math. Sem. Univ. Hamburg **7** (1929), 46–51.
- [3] L. Bartholdi and M. R. Bush, *Maximal unramified 3-extensions of imaginary quadratic fields and $SL_2\mathbb{Z}_3$* , J. Number Theory **124** (2007), 159–166.
- [4] H. U. Besche, B. Eick, and E. A. O’Brien, *A millennium project: constructing small groups*, Int. J. Algebra Comput. **12** (2002), 623–644.
- [5] H. U. Besche, B. Eick, and E. A. O’Brien, *The SmallGroups Library — a Library of Groups of Small Order*, 2005, an accepted and refereed GAP 4 package, available also in MAGMA.
- [6] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235–265.
- [7] W. Bosma, J. J. Cannon, C. Fieker, and A. Steels (eds.), *Handbook of Magma functions* (Edition 2.21, Sydney, 2014).
- [8] N. Boston, M. R. Bush and F. Hajir, *Heuristics for p -class towers of imaginary quadratic fields*, to appear in Math. Annalen, 2015. (arXiv: 1111.4679v1 [math.NT] 20 Nov 2011.)
- [9] N. Boston, C. Leedham-Green, *Explicit computation of Galois p -groups unramified at p* , J. Algebra **256** (2002), 402–413.
- [10] N. Boston, H. Nover, *Computing pro- p Galois groups*, Proceedings of ANTS 2006, Lecture Notes in Computer Science **4076**, 1–10, Springer-Verlag Berlin Heidelberg, 2006.
- [11] D. A. Buell, *Class groups of quadratic fields*, Math. Comp. **30** (1976), no. 135, 610–623.
- [12] M. R. Bush, *Computation of Galois groups associated to the 2-class towers of some quadratic fields*, J. Number Theory **100** (2003), 313–325.
- [13] M. R. Bush and D. C. Mayer, *3-class field towers of exact length 3*, J. Number Theory **147** (2015), 766–777, DOI 10.1016/j.jnt.2014.08.010. (arXiv: 1312.0251v1 [math.NT] 1 Dec 2013.)
- [14] F. Diaz y Diaz, *Sur les corps quadratiques imaginaires dont le 3-rang du groupe des classes est supérieur à 1*, Séminaire Delange-Pisot-Poitou, 1973/74, no. G15.
- [15] F. Diaz y Diaz, *Sur le 3-rang des corps quadratiques*, No. 78-11, Univ. Paris-Sud, 1978.
- [16] C. Fieker, *Computing class fields via the Artin map*, Math. Comp. **70** (2001), no. 235, 1293–1303.
- [17] The GAP Group, *GAP – Groups, Algorithms, and Programming — a System for Computational Discrete Algebra*, Version 4.7.5, Aachen, Braunschweig, Fort Collins, St. Andrews, 2014, (`\protect\vrule width0pt\protect\href{http://www.gap-system.org}{http://www.gap-system.org}`).
- [18] G. Gamble, W. Nickel, and E. A. O’Brien, *ANU p -Quotient — p -Quotient and p -Group Generation Algorithms*, 2006, an accepted GAP 4 package, available also in MAGMA.
- [19] E. S. Golod and I. R. Shafarevich, *On class field towers* (Russian), Izv. Akad. Nauk SSSR, Ser. Mat. **28** (1964), no. 2, 261–272. (English transl. in Amer. Math. Soc. Transl. (2) **48** (1965), 91–102.)
- [20] F.-P. Heider und B. Schmithals, *Zur Kapitulation der Idealklassen in unverzweigten primzyklischen Erweiterungen*, J. Reine Angew. Math. **336** (1982), 1–25.
- [21] D. F. Holt, B. Eick, and E. A. O’Brien, *Handbook of computational group theory*, Discrete mathematics and its applications, Chapman and Hall/CRC Press, 2005.
- [22] H. Koch und B. B. Venkov, *Über den p -Klassenkörperturm eines imaginär-quadratischen Zahlkörpers*, Astérisque **24–25** (1975), 57–67.
- [23] The MAGMA Group, *MAGMA Computational Algebra System*, Version 2.21-1, Sydney, 2014, (`\protect\vrule width0pt\protect\href{http://magma.maths.usyd.edu.au}{http://magma.maths.usyd.edu.au}`).
- [24] D. C. Mayer, *Principalization in complex S_3 -fields*, Congressus Numerantium **80** (1991), 73–87. (Proceedings of the Twentieth Manitoba Conference on Numerical Mathematics and Computing, Univ. of Manitoba, Winnipeg, Canada, 1990.)
- [25] D. C. Mayer, *The second p -class group of a number field*, Int. J. Number Theory **8** (2012), no. 2, 471–505, DOI 10.1142/S179304211250025X.
- [26] D. C. Mayer, *Transfers of metabelian p -groups*, Monatsh. Math. **166** (2012), no. 3–4, 467–495, DOI 10.1007/s00605-010-0277-x.
- [27] D. C. Mayer, *The distribution of second p -class groups on coclass graphs*, J. Théor. Nombres Bordeaux **25** (2013), no. 2, 401–456, DOI 10.5802/jtnb842. (27th Journées Arithmétiques, Faculty of Mathematics and Informatics, Univ. of Vilnius, Lithuania, 2011.)
- [28] D. C. Mayer, *Principalization algorithm via class group structure*, J. Théor. Nombres Bordeaux **26** (2014), no. 2, 415–464.
- [29] D. C. Mayer, *Periodic bifurcations in descendant trees of finite p -groups*, to appear in Adv. Pure Math., Special Issue on Group Theory, 2015.
- [30] D. C. Mayer, *Index- p abelianization data of p -class field tower groups*, 29th Journées Arithmétiques, Univ. of Debrecen, Hungary, 2015.
- [31] H. Nover, *Computation of Galois Groups of 2-Class Towers*, Ph.D. Thesis, University of Wisconsin, Madison, 2009.
- [32] E. A. O’Brien, *The p -group generation algorithm*, J. Symbolic Comput. **9** (1990), 677–698.

- [33] The PARI Group, PARI/GP, Version 2.7.2, Bordeaux, 2014, ([\protect\vrule width0pt\protect\href{http://pari.math.u-bordeaux.fr}](http://pari.math.u-bordeaux.fr)).
- [34] A. Scholz und O. Taussky, *Die Hauptideale der kubischen Klassenkörper imaginär quadratischer Zahlkörper: ihre rechnerische Bestimmung und ihr Einfluß auf den Klassenkörperturm*, J. Reine Angew. Math. **171** (1934), 19–41.
- [35] I. R. Shafarevich, *Extensions with prescribed ramification points* (Russian), Publ. Math., Inst. Hautes Études Sci. **18** (1964), 71–95. (English transl. by J. W. S. Cassels in Amer. Math. Soc. Transl., II. Ser., **59** (1966), 128–149.)
- [36] D. Shanks, *Class groups of the quadratic fields found by Diaz y Diaz*, Math. Comp. **30** (1976), 173–178.
- [37] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences* (OEIS), The OEIS Foundation Inc., 2014, ([\protect\vrule width0pt\protect\href{http://oeis.org/}](http://oeis.org/){<http://oeis.org/>}).
- [38] O. Taussky, *A remark concerning Hilbert's Theorem 94*, J. Reine Angew. Math. **239/240** (1970), 435–438.
- [39] E. B. Vinberg, *On a theorem concerning the infinite-dimensionality of an associative algebra* (Russian), Izv. Akad. Nauk SSSR, Ser. Mat. **29** (1965), 209–214. (English transl. in Amer. Math. Soc. Transl. (2) **82** (1969), 237–242.)

NAGLERGASSE 53, 8010 GRAZ, AUSTRIA

E-mail address: algebraic.number.theory@algebra.at

URL: <http://www.algebra.at>