

# Counting toroidal binary arrays, II

S. N. Ethier\* and Jiyeon Lee†

## Abstract

We derive formulas for (i) the number of toroidal  $n \times n$  binary arrays, allowing rotation of rows and/or columns as well as matrix transposition, and (ii) the number of toroidal  $n \times n$  binary arrays, allowing rotation and/or reflection of rows and/or columns as well as matrix transposition.

2010 *Mathematics Subject Classification*: Primary 05A05.

*Keywords*: toroidal array, Euler's phi function, group action, orbit, Pólya's enumeration theorem.

## 1 Introduction

A previous paper [1] found the number of (distinct) toroidal  $m \times n$  binary arrays, allowing rotation of rows and/or columns, to be

$$a(m, n) := \frac{1}{mn} \sum_{c|m} \sum_{d|n} \varphi(c)\varphi(d) 2^{mn/\text{lcm}(c,d)}, \quad (1)$$

where  $\varphi$  is Euler's phi function and lcm stands for least common multiple. This is [A184271](#) in the *On-Line Encyclopedia of Integer Sequences* [2]. The main diagonal is [A179043](#). It was also shown that, allowing rotation and/or reflection of rows and/or columns, the number becomes

$$b(m, n) := b_1(m, n) + b_2(m, n) + b_3(m, n) + b_4(m, n), \quad (2)$$

where

$$b_1(m, n) := \frac{1}{4mn} \sum_{c|m} \sum_{d|n} \varphi(c)\varphi(d) 2^{mn/\text{lcm}(c,d)},$$

---

\*Department of Mathematics, University of Utah, 155 South 1400 East, Salt Lake City, UT 84112 USA. [ethier@math.utah.edu](mailto:ethier@math.utah.edu). Partially supported by a grant from the Simons Foundation (209632).

†Department of Statistics, Yeungnam University, 214-1 Daedong, Kyeongsan, Kyeongbuk 712-749, South Korea. [leejy@yu.ac.kr](mailto:leejy@yu.ac.kr). Supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (No. 2013R1A1A3A04007670).

$$\begin{aligned}
b_2(m, n) &:= \frac{1}{4n} \sum_{d|n} \varphi(d) 2^{mn/d} \\
&+ \begin{cases} (4n)^{-1} \sum' \varphi(d) (2^{(m+1)n/(2d)} - 2^{mn/d}), & \text{if } m \text{ is odd;} \\ (8n)^{-1} \sum' \varphi(d) (2^{mn/(2d)} + 2^{(m+2)n/(2d)} - 2 \cdot 2^{mn/d}), & \text{if } m \text{ is even,} \end{cases}
\end{aligned}$$

with  $\sum' := \sum_{d|n: d \text{ is odd}}$ ,

$$b_3(m, n) := b_2(n, m),$$

and

$$b_4(m, n) := \begin{cases} 2^{(mn-3)/2}, & \text{if } m \text{ and } n \text{ are odd;} \\ 3 \cdot 2^{mn/2-3}, & \text{if } m \text{ and } n \text{ have opposite parity;} \\ 7 \cdot 2^{mn/2-4}, & \text{if } m \text{ and } n \text{ are even.} \end{cases}$$

(The formula for  $b_2(m, n)$  given in [1] is simplified here.) This is [A222188](#) in the *OEIS* [2]. The main diagonal is [A209251](#).

Our aim here is to derive the corresponding formulas when  $m = n$  and we allow matrix transposition as well. More precisely, we show that the number of (distinct) toroidal  $n \times n$  binary arrays, allowing rotation of rows and/or columns as well as matrix transposition, is

$$\alpha(n) = \frac{1}{2} a(n, n) + \frac{1}{2n} \sum_{d|n} \varphi(d) 2^{n(n+d-2\lfloor d/2 \rfloor)/(2d)}, \quad (3)$$

where  $a(n, n)$  is from (1). When we allow rotation and/or reflection of rows and/or columns as well as matrix transposition, the number becomes

$$\begin{aligned}
\beta(n) &= \frac{1}{2} b(n, n) + \frac{1}{4n} \sum_{d|n} \varphi(d) 2^{n(n+d-2\lfloor d/2 \rfloor)/(2d)} \\
&+ \begin{cases} 2^{(n^2-5)/4}, & \text{if } n \text{ is odd;} \\ 5 \cdot 2^{n^2/4-3}, & \text{if } n \text{ is even,} \end{cases} \quad (4)
\end{aligned}$$

where  $b(n, n)$  is from (2). At the time of writing, sequences (3) and (4) were not in the *OEIS*.

For an alternative description, we could define a group action on the set of  $n \times n$  binary arrays, which has  $2^{n^2}$  elements. If the group is generated by  $\sigma$  (row rotation) and  $\tau$  (column rotation), then the number of orbits is given by  $a(n, n)$ ; see [1]. If the group is generated by  $\sigma$ ,  $\tau$ , and  $\zeta$  (matrix transposition), then the number of orbits is given by  $\alpha(n)$ ; see Theorem 1 below. If the group is generated by  $\sigma$ ,  $\tau$ ,  $\rho$  (row reflection), and  $\theta$  (column reflection), then the number of orbits is given by  $b(n, n)$ ; see [1]. If the group is generated by  $\sigma$ ,  $\tau$ ,  $\rho$ ,  $\theta$ , and  $\zeta$ , then the number of orbits is given by  $\beta(n)$ ; see Theorem 2 below.

Both theorems are proved using Pólya's enumeration theorem (actually, the simplified unweighted version; see, e.g., van Lint and Wilson [3, Theorem 37.1, p. 524]).

To help clarify the distinction between the various group actions, we consider the case of  $3 \times 3$  binary arrays as in [1]. When the group is generated by  $\sigma$  and  $\tau$  (allowing rotation of rows and/or columns), there are 64 orbits, which were listed in [1]. When the group is generated by  $\sigma$ ,  $\tau$ , and  $\zeta$  (allowing rotation of rows and/or columns as well as matrix transposition), there are 44 orbits, which are listed in Table 1 below. When the group is generated by  $\sigma$ ,  $\tau$ ,  $\rho$ , and  $\theta$  (allowing rotation and/or reflection of rows and/or columns), there are 36 orbits, which were listed in [1]. When the group is generated by  $\sigma$ ,  $\tau$ ,  $\rho$ ,  $\theta$ , and  $\zeta$  (allowing rotation and/or reflection of rows and/or columns as well as matrix transposition), there are 26 orbits, which are listed in Table 2 below.

Table 1: A list of the 44 orbits of the group action in which the group generated by  $\sigma$ ,  $\tau$ , and  $\zeta$  acts on the set of  $3 \times 3$  binary arrays. (Rows and/or columns can be rotated and matrices can be transposed.) Each orbit is represented by its minimal element in 9-bit binary form. Subscripts indicate orbit size. Bars separate different numbers of 1s.

$$\begin{aligned}
& \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right)_1 \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{array} \right)_9 \mid \\
& \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 1 & 1 \end{array} \right)_6 \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 0 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{array} \right)_3 \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{array} \right)_3 \mid \\
& \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{array} \right)_9 \mid \\
& \left( \begin{array}{ccc} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{array} \right)_9 \mid \\
& \left( \begin{array}{ccc} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right)_9 \mid \\
& \left( \begin{array}{ccc} 0 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right)_6 \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{array} \right)_3 \mid \left( \begin{array}{ccc} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{array} \right)_3 \mid \\
& \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right)_{18} \mid \left( \begin{array}{ccc} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right)_9 \mid \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right)_1
\end{aligned}$$

Table 2: A list of the 26 orbits of the group action in which the group generated by  $\sigma, \tau, \rho, \theta$ , and  $\zeta$  acts on the set of  $3 \times 3$  binary arrays. (Rows and/or columns can be rotated and/or reflected and matrices can be transposed.) Each orbit is represented by its minimal element in 9-bit binary form. Subscripts indicate orbit size. Bars separate different numbers of 1s.

$$\begin{aligned}
 & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}_1 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}_6 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}_{36} \\
 & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}_6 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}_{36} \mid \\
 & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix}_9 \mid \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_6 \mid \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{36} \\
 & \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}_{36} \mid \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}_6 \mid \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}_{18} \mid \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_9 \mid \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}_1
 \end{aligned}$$

Table 3 provides numerical values for  $\alpha(n)$  and  $\beta(n)$  for small  $n$ .

Table 3: The values of  $\alpha(n)$  and  $\beta(n)$  for  $n = 1, 2, \dots, 12$ .

$n$	$\alpha(n)$	$\beta(n)$
1		2
2		6
3		44
4		2209
5		674384
6		954623404
7		5744406453840
8		144115192471496836
9		14925010120653819583840
10		6338253001142965335834871200
11	10985355337065423791175013899922368	1584563250300891724601560272
12	77433143050453552587418968170813573149024	2746338834266358751489231123956672
		19358285762613388352671214587818634041520

We take this opportunity to correct a small gap in the proof of Theorem 2 in [1]. The proof assumed implicitly that  $m, n \geq 3$ . The theorem is correct as stated for  $m, n \geq 1$ , so the proof is incomplete if  $m$  or  $n$  is 1 or 2. Following the proof of Theorem 2 below, we supply the missing steps.

## 2 Rotation of rows and columns, and matrix transposition

Let  $X_n := \{0, 1\}^{\{0, 1, \dots, n-1\}^2}$  be the set of  $n \times n$  matrices of 0s and 1s, which has  $2^{n^2}$  elements. Let  $\alpha(n)$  denote the number of orbits of the group action on  $X_n$  by the group of order  $2n^2$  generated by  $\sigma$  (row rotation),  $\tau$  (column rotation), and  $\zeta$  (matrix transposition). (Exception: If  $n = 1$ , the group is of order 1.)

Informally,  $\alpha(n)$  is the number of (distinct) toroidal  $n \times n$  binary arrays, allowing rotation of rows and/or columns as well as matrix transposition.

**Theorem 1.** *With  $a(n, n)$  defined using (1),  $\alpha(n)$  is given by (3).*

*Proof.* Let us assume that  $n \geq 2$ . By Pólya's enumeration theorem,

$$\alpha(n) = \frac{1}{2n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{A_{ij}} + 2^{E_{ij}}), \quad (5)$$

where  $A_{ij}$  (resp.,  $E_{ij}$ ) is the number of cycles in the permutation  $\sigma^i \tau^j$  (resp.,  $\sigma^i \tau^j \zeta$ ); here  $\sigma$  rotates the rows (row 0 becomes row 1, row 1 becomes row 2, ..., row  $n-1$  becomes row 0),  $\tau$  rotates the columns, and  $\zeta$  transposes the matrix. We know from [1] that

$$a(n, n) = \frac{1}{n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{A_{ij}}, \quad (6)$$

so it remains to find  $E_{ij}$ . The permutation  $\zeta$  has  $n$  fixed points and  $\binom{n}{2}$  transpositions, so  $E_{00} = n(n+1)/2$ .

Notice that  $\sigma$  and  $\tau$  commute, whereas  $\sigma\zeta = \zeta\tau$  and  $\tau\zeta = \zeta\sigma$ . Let  $(i, j) \in \{0, 1, \dots, n-1\}^2 - \{(0, 0)\}$  be arbitrary. Then

$$(\sigma^i \tau^j \zeta)^2 = (\sigma^i \tau^j \zeta)(\zeta \tau^i \sigma^j) = \sigma^{i+j} \tau^{i+j},$$

hence

$$\begin{aligned} (\sigma^i \tau^j \zeta)^{2d} &= \sigma^{(i+j)d} \tau^{(i+j)d} = ((\sigma\tau)^{i+j})^d, \\ (\sigma^i \tau^j \zeta)^{2d+1} &= \sigma^{(i+j)d+i} \tau^{(i+j)d+j} \zeta. \end{aligned}$$

Clearly,  $(\sigma^i \tau^j \zeta)^{2d+1}$  cannot be the identity permutation, so  $\sigma^i \tau^j \zeta$  is of even order. Using the fact that, in the cyclic group  $\{a, a^2, \dots, a^{n-1}, a^n = e\}$  of order  $n$ ,  $a^k$  is of order  $n/\gcd(k, n)$ , we find that the permutation  $\sigma^i \tau^j \zeta$  is of order  $2d$ , where  $d := n/\gcd(i+j, n)$ . Therefore, every cycle of this permutation must have length that divides  $2d$ .

We claim that all cycles have length  $d$  or  $2d$ . Accepting that for now, let us determine how many cycles have length  $d$ . A cycle that includes entry  $(k, l)$  has length  $d$  if  $(k, l)$  is a fixed point of  $(\sigma^i \tau^j \zeta)^d$ . For this to hold we must have

$d$  odd (otherwise there would be no fixed points because we have excluded the case  $i = j = 0$  and  $(i + j)d/2 = \text{lcm}(i + j, n)/2$  is not a multiple of  $n$ ). Since

$$(\sigma^i \tau^j \zeta)^d = \sigma^{(i+j)(d-1)/2+i} \tau^{(i+j)(d-1)/2+j} \zeta,$$

we must also have

$$(k, l) = ([l + (i + j)(d - 1)/2 + j], [k + (i + j)(d - 1)/2 + i]), \quad (7)$$

where  $d := n / \gcd(i + j, n)$  and, for simplicity,  $[r] := (r \bmod n) \in \{0, 1, \dots, n - 1\}$ . For each  $k \in \{0, 1, \dots, n - 1\}$ , there is a unique  $l$  (namely,  $l := [k + (i + j)(d - 1)/2 + i]$ ) such that (7) holds; indeed,

$$\begin{aligned} & [l + (i + j)(d - 1)/2 + j] \\ &= [[k + (i + j)(d - 1)/2 + i] + (i + j)(d - 1)/2 + j] \\ &= [k + (i + j)(d - 1)/2 + i + (i + j)(d - 1)/2 + j] \\ &= [k + (i + j)d] \\ &= [k + (i + j)(n / \gcd(i + j, n))] \\ &= [k + \text{lcm}(i + j, n)] \\ &= k. \end{aligned}$$

This shows that there are  $n$  fixed points of  $(\sigma^i \tau^j \zeta)^d$ . Each cycle of length  $d$  of  $\sigma^i \tau^j \zeta$  will account for  $d$  such fixed points, hence there are  $n/d$  such cycles. All remaining cycles will have length  $2d$ , and so there are  $n(n - 1)/(2d)$  of these. The total number of cycles is therefore  $n(n + 1)/(2d)$ .

The other possibility is that  $d$  is even and all cycles have the same length,  $2d$ , so there are  $n^2/(2d)$  of them. Notice that  $d$  is a divisor of  $n$ , so the contribution to

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{E_{ij}}$$

from odd  $d$  is

$$\sum_{d | n: d \text{ is odd}} n\varphi(d)2^{n(n+1)/(2d)} \quad (8)$$

and from even  $d$  is

$$\sum_{d | n: d \text{ is even}} n\varphi(d)2^{n^2/(2d)}. \quad (9)$$

The reason for the coefficient  $n\varphi(d)$  is that, if  $d | n$ , then the number of elements of the cyclic group  $\{e, \sigma\tau, (\sigma\tau)^2, \dots, (\sigma\tau)^{n-1}\}$  that are of order  $d$  is  $\varphi(d)$ . And for a given  $(i, j) \in \{0, 1, \dots, n - 1\}^2$ , there are  $n$  pairs  $(k, l) \in \{0, 1, \dots, n - 1\}^2$  such that  $[k + l] = [i + j]$ . Putting (8) and (9) together, we obtain

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{E_{ij}} = \sum_{d | n} n\varphi(d)2^{n(n+d-2\lfloor d/2 \rfloor)/(2d)}, \quad (10)$$

which, together with (5) and (6), yields (3).  $\square$

It remains to prove our claim that, for  $(i, j) \in \{0, 1, \dots, n-1\}^2 - \{(0, 0)\}$ , the permutation  $\sigma^i \tau^j \zeta$  cannot have any cycles whose length is a proper divisor of  $d := n / \gcd(i+j, n)$ . Let  $c | d$  with  $1 \leq c < d$ . We must show that  $(\sigma^i \tau^j \zeta)^c$  has no fixed points. We can argue as above with  $c$  in place of  $d$ . For  $(k, l)$  to be a fixed point of  $(\sigma^i \tau^j \zeta)^c$  we must have  $(i+j)c$  a multiple of  $n$ . But  $d := n / \gcd(i+j, n)$  is the smallest integer  $c$  such that  $(i+j)c$  is a multiple of  $n$  because  $(i+j)n / \gcd(i+j, n) = \text{lcm}(i+j, n)$ .

Finally, we excluded the case  $n = 1$  at the beginning of the proof, but we notice that the formula (3) gives  $\alpha(1) = 2$ , which is correct.

### 3 Rotation and reflection of rows and columns, and matrix transposition

Let  $X_n := \{0, 1\}^{\{0, 1, \dots, n-1\}^2}$  be the set of  $n \times n$  matrices of 0s and 1s, which has  $2^{n^2}$  elements. Let  $\beta(n)$  denote the number of orbits of the group action on  $X_n$  by the group of order  $8n^2$  generated by  $\sigma$  (row rotation),  $\tau$  (column rotation),  $\rho$  (row reflection),  $\theta$  (column reflection), and  $\zeta$  (matrix transposition). (Exceptions: If  $n = 2$ , the group is of order 8; if  $n = 1$ , the group is of order 1.)

Informally,  $\beta(n)$  is the number of (distinct) toroidal  $n \times n$  binary arrays, allowing rotation and/or reflection of rows and/or columns as well as matrix transposition.

**Theorem 2.** *With  $b(n, n)$  defined using (2),  $\beta(n)$  is given by (4).*

*Proof.* Let us assume that  $n \geq 3$ . (We will treat the cases  $n = 1$  and  $n = 2$  later.) By Pólya's enumeration theorem,

$$\beta(n) = \frac{1}{8n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{A_{ij}} + 2^{B_{ij}} + 2^{C_{ij}} + 2^{D_{ij}} + 2^{E_{ij}} + 2^{F_{ij}} + 2^{G_{ij}} + 2^{H_{ij}}),$$

where  $A_{ij}$  (resp.,  $B_{ij}$ ,  $C_{ij}$ ,  $D_{ij}$ ,  $E_{ij}$ ,  $F_{ij}$ ,  $G_{ij}$ ,  $H_{ij}$ ) is the number of cycles in the permutation  $\sigma^i \tau^j$  (resp.,  $\sigma^i \tau^j \rho$ ,  $\sigma^i \tau^j \theta$ ,  $\sigma^i \tau^j \rho \theta$ ,  $\sigma^i \tau^j \zeta$ ,  $\sigma^i \tau^j \rho \zeta$ ,  $\sigma^i \tau^j \theta \zeta$ ,  $\sigma^i \tau^j \rho \theta \zeta$ ); here  $\sigma$  rotates the rows (row 0 becomes row 1, row 1 becomes row 2, ..., row  $n-1$  becomes row 0),  $\tau$  rotates the columns,  $\rho$  reflects the rows (rows 0 and  $n-1$  are interchanged, rows 1 and  $n-2$  are interchanged, ..., rows  $\lfloor n/2 \rfloor - 1$  and  $n - \lfloor n/2 \rfloor$  are interchanged),  $\theta$  reflects the columns, and  $\zeta$  transposes the matrix. The order of the group generated by  $\sigma$ ,  $\tau$ ,  $\rho$ ,  $\theta$ , and  $\zeta$  is  $8n^2$ , using the assumption that  $n \geq 3$ .

We have already evaluated

$$a(n, n) = \frac{1}{n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{A_{ij}},$$

$$\alpha(n) = \frac{1}{2n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{A_{ij}} + 2^{E_{ij}}),$$

and

$$b(n, n) = \frac{1}{4n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{A_{ij}} + 2^{B_{ij}} + 2^{C_{ij}} + 2^{D_{ij}}),$$

so

$$\beta(n) = \frac{1}{2} b(n, n) + \frac{1}{4} \left( \alpha(n) - \frac{1}{2} a(n, n) \right) + \frac{1}{8n^2} \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} (2^{F_{ij}} + 2^{G_{ij}} + 2^{H_{ij}}). \quad (11)$$

Let us begin with

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{H_{ij}}.$$

Here we are concerned with the permutations  $\sigma^i \tau^j \rho \theta \zeta$  for  $(i, j) \in \{0, 1, \dots, n-1\}^2$ . We will need some multiplication rules for the permutations  $\sigma, \tau, \rho, \theta$ , and  $\zeta$ , specifically

$$\sigma\tau = \tau\sigma, \quad \sigma\theta = \theta\sigma, \quad \tau\rho = \rho\tau, \quad \rho\theta = \theta\rho, \quad \sigma\rho = \rho\sigma^{-1}, \quad \tau\theta = \theta\tau^{-1},$$

and

$$\sigma\zeta = \zeta\tau, \quad \tau\zeta = \zeta\sigma, \quad \rho\zeta = \zeta\theta, \quad \theta\zeta = \zeta\rho.$$

It follows that (with  $\tau^{-i} := (\tau^{-1})^i$ )

$$\sigma^i \tau^j \rho \theta \zeta = \sigma^i \tau^j \zeta \theta \rho = \zeta \tau^i \sigma^j \theta \rho = \zeta \theta \tau^{-i} \sigma^j \rho = \zeta \theta \rho \tau^{-i} \sigma^{-j},$$

and hence

$$(\sigma^i \tau^j \rho \theta \zeta)^2 = (\sigma^i \tau^j \rho \theta \zeta)(\zeta \theta \rho \tau^{-i} \sigma^{-j}) = \sigma^{i-j} \tau^{-i+j} = (\sigma \tau^{-1})^{i-j} = (\sigma^{-1} \tau)^{-i+j}. \quad (12)$$

In particular, if  $i \in \{0, 1, \dots, n-1\}$ , then the permutation  $\sigma^i \tau^i \rho \theta \zeta$  is of order 2. Furthermore, under this permutation, the entry in position  $(k, l)$  moves to position  $(n-1-[l+i], n-1-[k+i])$ , where, as before,  $[r] := (r \bmod n) \in \{0, 1, \dots, n-1\}$ . Thus,  $(k, l)$  is a fixed point if and only if

$$(k, l) = (n-1-[l+i], n-1-[k+i]). \quad (13)$$

For each  $k \in \{0, 1, \dots, n-1\}$  there is a unique  $l \in \{0, 1, \dots, n-1\}$  (namely  $l := n-1-[k+i]$ ) such that (13) holds; indeed,

$$\begin{aligned} n-1-[l+i] &= n-1-[n-1-[k+i]+i] = n-1-[n-1-(k+i)+i] \\ &= n-1-[n-1-k] = n-1-(n-1-k) = k. \end{aligned}$$

Thus,  $\sigma^i \tau^i \rho \theta \zeta$  with  $i \in \{0, 1, \dots, n-1\}$  is of order 2 and has exactly  $n$  fixed points, hence  $\binom{n}{2}$  transpositions. This implies that  $H_{ii} = n(n+1)/2$  for such  $i$ .

Now we let  $(i, j) \in \{0, 1, \dots, n-1\}^2$  be arbitrary but with  $i \neq j$ . Let us generalize (12) to

$$(\sigma^i \tau^j \rho \theta \zeta)^{2d} = \sigma^{(i-j)d} \tau^{(-i+j)d} = ((\sigma \tau^{-1})^{i-j})^d = ((\sigma^{-1} \tau)^{-i+j})^d,$$



$$(\sigma^i \tau^j \rho \theta \zeta)^{2d+1} = \sigma^{(i-j)d+i} \tau^{(-i+j)d+j} \rho \theta \zeta.$$

The proof proceeds much like the proof of Theorem 1. Specifically,  $\sigma^i \tau^j \rho \theta \zeta$  is of order  $2d$ , where  $d := n / \gcd(|i-j|, n)$ . All cycles have length  $d$  or  $2d$ . In fact, if  $d$  is odd, there are  $n/d$  cycles of length  $d$  and  $n(n-1)/(2d)$  cycles of length  $2d$ . If  $d$  is even, there are  $n^2/(2d)$  cycles, all of length  $2d$ . And for a given  $(i, j) \in \{0, 1, \dots, n-1\}^2$ , there are  $n$  pairs  $(k, l) \in \{0, 1, \dots, n-1\}^2$  such that  $[k-l] = [|i-j|]$ . We arrive at the conclusion that

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{H_{ij}} = \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{E_{ij}}. \quad (14)$$

Next we evaluate

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{F_{ij}} = \sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{G_{ij}}, \quad (15)$$

where the equality holds by symmetry. We consider the permutations  $\sigma^i \tau^j \rho \zeta$  for  $(i, j) \in \{0, 1, \dots, n-1\}^2$ . From the multiplication rules, it follows that

$$\sigma^i \tau^j \rho \zeta = \zeta \theta \tau^{-i} \sigma^j$$

and hence

$$(\sigma^i \tau^j \rho \zeta)^2 = (\sigma^i \tau^j \rho \zeta)(\zeta \theta \tau^{-i} \sigma^j) = \sigma^i \tau^j \rho \theta \tau^{-i} \sigma^j = \sigma^{i-j} \tau^{i+j} \rho \theta = \theta \rho \tau^{-i-j} \sigma^{-i+j}, \quad (16)$$

which implies

$$(\sigma^i \tau^j \rho \zeta)^4 = (\sigma^{i-j} \tau^{i+j} \rho \theta)(\theta \rho \tau^{-i-j} \sigma^{-i+j}) = e.$$

So the permutation  $\sigma^i \tau^j \rho \zeta$  is of order 4. The entry in position  $(k, l)$  moves to position  $([l+j], n-1-[k+i])$  under this permutation. Thus,  $(k, l) \in \{0, 1, \dots, n-1\}^2$  is a fixed point of  $\sigma^i \tau^j \rho \zeta$  if and only if

$$(k, l) = ([l+j], n-1-[k+i]).$$

There is a solution  $(k, l)$  if and only if there exists  $l \in \{0, 1, \dots, n-1\}$  such that, with  $k := [l+j]$ , we have  $n-1-[k+i] = l$  or, equivalently,

$$[l+i+j] = n-1-l. \quad (17)$$

When  $i+j \leq n-1$ , (17) is equivalent to

$$l+i+j = n-1-l \quad \text{or} \quad l+i+j-n = n-1-l$$

or to

$$l = (n-1-i-j)/2 \quad \text{or} \quad l = (2n-1-i-j)/2.$$

If  $n$  is odd and  $i+j$  is odd, then there is one fixed point,  $(k, l) = ((2n-1-i-j)/2, [(2n-1-i-j)/2])$ . If  $n$  is odd and  $i+j$  is even, then there is one fixed

point,  $(k, l) = ([\frac{n-1-i+j}{2}], [\frac{n-1-i-j}{2}])$ . If  $n$  is even and  $i+j$  is odd, then there are two fixed points, namely

$$\begin{aligned}(k, l) &= ([\frac{n-1-i+j}{2}], [\frac{n-1-i-j}{2}]), \\ (k, l) &= ([\frac{2n-1-i+j}{2}], [\frac{2n-1-i-j}{2}]).\end{aligned}$$

Finally, if  $n$  is even and  $i+j$  is even, then there is no fixed point.

When  $i+j \geq n$ , (17) is equivalent to

$$l+i+j-n = n-1-l \quad \text{or} \quad l+i+j-2n = n-1-l$$

or to

$$l = (2n-1-i-j)/2 \quad \text{or} \quad l = (3n-1-i-j)/2.$$

If  $n$  is odd and  $i+j$  is odd, then there is one fixed point,  $(k, l) = ([\frac{2n-1-i+j}{2}], [\frac{2n-1-i-j}{2}])$ . If  $n$  is odd and  $i+j$  is even, then there is one fixed point,  $(k, l) = ([\frac{n-1-i+j}{2}], [\frac{3n-1-i-j}{2}]) = ([\frac{n-1-i+j}{2}], [\frac{n-1-i-j}{2}])$ . If  $n$  is even and  $i+j$  is odd, then there are two fixed points, namely

$$\begin{aligned}(k, l) &= ([\frac{2n-1-i+j}{2}], [\frac{2n-1-i-j}{2}]), \\ (k, l) &= ([\frac{n-1-i+j}{2}], [\frac{n-1-i-j}{2}]).\end{aligned}$$

Finally, if  $n$  is even and  $i+j$  is even, then there is no fixed point. Notice that the results are the same for  $i+j \geq n$  as for  $i+j \leq n-1$ .

Using (16), under the permutation  $(\sigma^i \tau^j \rho \zeta)^2$ , the entry in position  $(k, l)$  moves to position  $(n-1-[k+i-j], n-1-[l+i+j])$ . Thus,  $(k, l) \in \{0, 1, \dots, n-1\}^2$  is a fixed point of  $(\sigma^i \tau^j \rho \zeta)^2$  if and only if

$$(k, l) = (n-1-[k+i-j], n-1-[l+i+j]).$$

A necessary and sufficient condition on  $(k, l)$  is (17) together with  $[k+i-j] = n-1-k$ . Solutions have  $l$  as before. On the other hand,  $k$  must satisfy

$$k+i-j-n = n-1-k, \quad k+i-j = n-1-k, \quad \text{or} \quad k+i-j+n = n-1-k,$$

or equivalently,

$$k = [(n-1-i+j)/2] \quad \text{or} \quad k = [(2n-1-i+j)/2].$$

If  $n$  is odd, the only fixed points of  $(\sigma^i \tau^j \rho \zeta)^2$  are those already shown to be fixed points of  $\sigma^i \tau^j \rho \zeta$ . If  $n$  is even and  $i+j$  is odd, there are two fixed points of  $(\sigma^i \tau^j \rho \zeta)^2$  that are not fixed points of  $\sigma^i \tau^j \rho \zeta$ , namely

$$\begin{aligned}(k, l) &= ([\frac{n-1-i+j}{2}], [\frac{2n-1-i-j}{2}]), \\ (k, l) &= ([\frac{2n-1-i+j}{2}], [\frac{n-1-i-j}{2}]).\end{aligned}$$

Finally, there are no fixed points when  $n$  is even and  $i+j$  is even.

Consequently, if  $n$  is odd, then the permutation  $\sigma^i \tau^j \rho \zeta$ , which is of order 4, has only one fixed point. Therefore, it has one cycle of length 1 and  $(n^2 - 1)/4$  cycles of length 4. Thus,

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{F_{ij}} = n^2 2^{(n^2+3)/4}.$$

For even  $n$ , if  $i + j$  is odd, then the permutation  $\sigma^i \tau^j \rho \zeta$  has two cycles of length 1 and one cycle of length 2, and the remaining cycles are of length 4. If  $i + j$  is even, then all cycles of the permutation  $\sigma^i \tau^j \rho \zeta$  are of length 4, hence there are  $n^2/4$  of them. Thus,

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n-1} 2^{F_{ij}} = \frac{1}{2} n^2 2^{(n^2-4)/4+3} + \frac{1}{2} n^2 2^{n^2/4} = 5n^2 2^{n^2/4-1}.$$

These results, together with (3), (10), (11), (14), and (15), yield (4).

Finally, recall that we have assumed that  $n \geq 3$ . We notice that the formula (4) gives  $\beta(1) = 2$  and  $\beta(2) = 6$ , which are correct, as we can see by direct enumeration.  $\square$

In the derivation of (2) in [1], the proof requires  $m, n \geq 3$  because the group  $D_m \times D_n$  used in the application of Pólya's enumeration theorem ( $D_m$  being the dihedral group of order  $2m$ ), is incorrect if  $m$  or  $n$  is 1 or 2. If  $m = 2$ , row rotation and row reflection are the same, so the latter is redundant. Thus,  $D_2$  should be replaced by  $C_2$ , the cyclic group of order 2. The reason (2) is still valid is that  $b_1(2, n) = b_2(2, n)$  and  $b_3(2, n) = b_4(2, n)$ , as is easily verified. If  $m = 1$ , again row reflection is redundant, so  $D_1$  should be replaced by  $C_1$ . Here (2) remains valid because  $b_1(1, n) = b_2(1, n)$  and  $b_3(1, n) = b_4(1, n)$ . A similar remark applies to  $n = 2$  and  $n = 1$ , except that here  $b_1(m, 2) = b_3(m, 2)$ ,  $b_2(m, 2) = b_4(m, 2)$ ,  $b_1(m, 1) = b_3(m, 1)$ , and  $b_2(m, 1) = b_4(m, 1)$ .

## References

- [1] S. N. Ethier, Counting toroidal binary arrays, *J. Integer Sequences* **16** (2013), (13.4.7) 1–8.
- [2] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, <http://oeis.org/>, 2015.
- [3] J. H. van Lint and R. M. Wilson, *A Course in Combinatorics*, Second Ed., Cambridge University Press, 2001.