

ACTION GRAPHS AND CATALAN NUMBERS

GERARDO ALVAREZ, JULIA E. BERGNER AND RUBEN LOPEZ

ABSTRACT. We introduce an inductively defined sequence of directed graphs and prove that the number of edges added at step k is equal to the k th Catalan number. Furthermore, we establish an isomorphism between the set of edges adjoined at step k and the set of planar rooted trees with k edges.

1. INTRODUCTION

In a recent paper, the second-named author and Hackney introduced certain inductively defined directed graphs with the goal of understanding the structure of a rooted category action on another category [1]. While these graphs were developed in such a way that they encoded the desired structure, the question remained what kinds of patterns could be found in these inductively defined sequences of graphs. In this paper, we look at the most basic of these graph sequences. We begin with the trivial graph with one vertex and no edges, and we inductively add new vertices and edges depending on the number of paths in the previous graph. We prove that at the k th step, the number of vertices and edges added is given by the k th Catalan number.

The Catalan numbers give a well-known sequence of natural numbers, arising in many contexts in combinatorics [3, A000108]. A long list of ways to obtain the Catalan numbers is found in Stanley's book [4] and subsequent online addendum [5]. Here we use that the 0th Catalan number is $C_0 = 1$, and, for any $k \geq 1$, the $(k + 1)$ st Catalan number C_{k+1} is given by the formula

$$C_{k+1} = \sum_{i=0}^k C_i C_{k-i}.$$

However, it is often of interest to determine a direct comparison with one of the other ways of obtaining the Catalan numbers. To this end, we establish a direct isomorphism between the set of edges added at step k in the action graph construction and the set of planar rooted trees with k edges, also known to have C_k elements.

2. ACTION GRAPHS

We begin by recalling a few basic definitions.

Definition 2.1. A *directed graph* is a pair $G = (V, E)$ where V is a set whose elements are called *vertices* and E is a set of ordered pairs of vertices, called *edges*. Given an edge $e = (v, w)$, we call v the *source* of e , denoted by $v = s(e)$, and call w the *target* of e , denoted by $w = t(e)$.

2010 *Mathematics Subject Classification.* 05A19, 05C05.
Key words and phrases. Catalan number, directed graph.

labeled by 1 pass through the vertex labeled by 0. Therefore, edges with source at the vertex labeled by 1 are adjoined exactly in the same way as edges with source 0, but one step later. \square

Proof of Theorem 2.4. We use induction and Lemma 2.5. For the base case we know that A_0 has one vertex labeled by 0 and $C_0 = 1$.

For the inductive step, suppose we know A_k has C_k vertices labeled by k . There is always a unique path from 0 to any vertex, so there are C_k paths from the vertex labeled 0 to a vertex labeled by k . Hence, by considering paths with source the 0 vertex, we create $C_k = C_0 C_k$ new vertices in A_{k+1} . Lemma 2.5 tells us that there are C_{k-1} paths from the single vertex labeled by 1 to vertices labeled by k . Thus, we must add $C_{k-1} = C_1 C_{k-1}$ new vertices in A_{k+1} . Applying Lemma 2.5 twice, we similarly see that there are C_{k-2} paths from any vertex labeled 2 (of which there are C_2) to any vertex labeled C_k , for a total of $C_2 C_{k-1}$ new vertices added. We continue this process, concluding by using the inductive hypothesis that there are C_k vertices labeled by k , from each of which there is only one (trivial) path, from which we produce $C_k = C_k C_0$ new vertices. Taking the sum, we obtain that the total number of paths in A_k ending at vertices labeled by k is

$$C_0 C_k + C_1 C_{k-1} + \cdots + C_k C_0 = C_{k+1}.$$

Since a new vertex is added for each such path, and a new edge for each such vertex, we have completed the proof. \square

3. A COMPARISON WITH PLANAR ROOTED TREES

In this section, we give an explicit one-to-one correspondence between the set of leaves of A_{k+1} and the set of planar rooted trees with $k+1$ edges. It is proved in that the latter set has C_{k+1} elements [2, 8.4].


We begin with the necessary definitions. Here, we work with graphs which are no longer assumed to be directed. A graph is *connected* if there exists a path between any two vertices.

Definition 3.1. A *tree* is a connected graph with no loops or multiple edges. A *rooted tree* is a tree with a specified vertex called the *root*.


Definition 3.2. A *leaf* of a rooted tree is a vertex of valence 1 which is not the root. (In the case of a single vertex with no edges, we take this root vertex also to be a leaf.) A *branch* of a rooted tree is a path from either the root or a vertex of valence greater than 2 to a leaf.

Note in particular that the action graphs are (directed) trees, and that the vertices of A_{k+1} which are not in A_k are precisely the leaves of A_{k+1} .

Here we want to consider *planar* rooted trees. Thus, if we view the bottom vertex as the root, we regard the following two trees as different:



To aid in our comparison with action graphs, we define a means of labeling the vertices of a planar rooted tree. The root vertex is always given the label 0. Given a representative of the tree with the root at the bottom, label the vertices by successive natural numbers, moving upward from the root and from left to right. For example, we have the labelings



Theorem 3.3. *There is an isomorphism between the set of leaves of the action graph A_k and the set of planar rooted trees with k edges.*

Proof. We prove this theorem inductively. When $k = 0$, both sets have only one element, since both A_0 and the only planar rooted tree with no edges consist of a single vertex and no edges.

Thus, suppose that we have proved the result for $k \geq 0$. Consider the set of planar rooted trees with $k + 1$ edges. As described above, assign a labeling of the vertices by the natural numbers $0, 1, \dots, k + 1$, using each number exactly once. Assign a direction to each edge, so that the label of the source vertex is less than the label of the target vertex.

Observe that the set of all such directed trees can be partially ordered by the length of the unique path from the vertex labeled by 0 to the vertex labeled by $k + 1$. Beginning with the tree with longest such path length, namely, the tree with a path of length $k + 1$, for each directed tree, identify the edge from 0 to 1, and any branches that do not end in the vertex $k + 1$, with vertices and edges already present in the graph. Adjoin new vertices and edges to the graph corresponding to the branch containing the vertex $k + 1$. Repeating for all planar rooted trees with $k + 1$ edges, we claim that the resulting directed graph is precisely the action graph A_{k+1} .

By our inductive hypothesis, the trees with k vertices assemble to produce the action graph A_k . But these trees can be obtained from those with $k + 1$ vertices by removing the leaf labeled by $k + 1$. Indeed, the (possibly multiple) ways of obtaining a rooted tree with $k + 1$ edges from one with k edges correspond to the ways we adjoin new leaves to produce A_{k+1} from A_k . In particular, there is precisely one vertex labeled by $k + 1$ coming from each planar rooted tree.

Conversely, regard A_{k+1} as a labeled rooted tree. We claim that the subtrees of A_{k+1} containing exactly one vertex labeled by i for each $0 \leq i \leq k + 1$ correspond exactly to the planar rooted trees with $k + 1$ edges. Again, assume this fact is true for $k \geq 0$. Given every such subtree with k edges, and vertices labeled from 0 to k , the choices for adjoining an extra edge containing a vertex labeled by $k + 1$ correspond exactly to the vertices on the path from the vertex labeled by 0 to the vertex labeled by k . But, these choices coincide with the ways to obtain a planar rooted tree with $k + 1$ edges from a given planar rooted tree with k edges. \square

4. ACKNOWLEDGMENTS

The first- and third-named authors were participants in the RISE program at UC Riverside in Summer 2013, with support from the HSI-STEM and CAMP programs, respectively. The second-named author was partially supported by NSF grant DMS-1105766 and CAREER award DMS-1352298.

REFERENCES

- [1] Julia E. Bergner and Philip Hackney, Reedy categories which encode the notion of category actions, to appear in *Fund. Math.*, preprint available at math.AT/1207.3467.
- [2] Thomas Koshy, *Catalan Numbers with Applications*, Oxford University Press, 2009.
- [3] Online Encyclopedia of Integer Sequences, <http://oeis.org/>.
- [4] Richard P. Stanley, *Enumerative Combinatorics. Vol. 2*. Cambridge Studies in Advanced Mathematics, 62. Cambridge University Press, Cambridge, 1999.
- [5] Richard P. Stanley, Catalan Addendum, <http://www-math.mit.edu/~rstan/ec/catadd.pdf>.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, CA 92521

E-mail address: `galva012@ucr.edu`

E-mail address: `bergnerj@member.ams.org`

E-mail address: `rlope015@ucr.edu`