

On One-Parameter Catalan Arrays

José Agapito ¹, Ângela Mestre ², and Maria M. Torres
Centro de Análise Funcional, Estruturas Lineares e Aplicações
Grupo de Estruturas Algébricas, Lineares e Combinatórias
Departamento de Matemática
Faculdade de Ciências, Universidade de Lisboa
1749-016 Lisboa
Portugal

jose.agapito@gmail.com

amestre@fc.ul.pt

mmtorres@fc.ul.pt

Pasquale Petrullo
Dipartimento di Matematica e Informatica
Università degli Studi della Basilicata
Potenza
Italy

p.petrullo@gmail.com

Abstract

We present a parametric family of Riordan arrays, which are obtained by multiplying any Riordan array with a generalized Pascal array. In particular, we focus on some interesting properties of one-parameter Catalan triangles. We obtain several combinatorial identities that involve two special Catalan matrices, the Chebyshev polynomials of the second kind, some periodic sequences, and the Fibonacci numbers.

1 Introduction

There are two infinite lower triangular matrices in the mathematical literature, that are both called Catalan triangles. Let us denote them by \mathbf{C} and \mathbf{B} , respectively. By way of

¹ Corresponding author.

² Work performed within the activities of Centro de Análise Funcional, Estruturas Lineares e Aplicações (Faculdade de Ciências, Universidade de Lisboa) and supported by the fellowship SFRH/BPD/48223/2008 provided by the Portuguese Foundation for Science and Technology (FCT).

illustration, their first rows are shown below:

$$\mathbf{C} = \begin{pmatrix} 1 & & & & & & & & \\ 1 & 1 & & & & & & & \\ 2 & 2 & 1 & & & & & & \\ 5 & 5 & 3 & 1 & & & & & \\ 14 & 14 & 9 & 4 & 1 & & & & \\ 42 & 42 & 28 & 14 & 5 & 1 & & & \\ 132 & 132 & 90 & 48 & 20 & 6 & 1 & & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 1 & & & & & & & & \\ 2 & 1 & & & & & & & \\ 5 & 4 & 1 & & & & & & \\ 14 & 14 & 6 & 1 & & & & & \\ 42 & 48 & 27 & 8 & 1 & & & & \\ 132 & 165 & 110 & 44 & 10 & 1 & & & \\ 429 & 572 & 429 & 208 & 65 & 12 & 1 & & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \end{pmatrix}.$$

Note that the 0th column of \mathbf{C} is given by the sequence of Catalan numbers $(C_n)_{n \geq 0}$, where $C_n = \frac{1}{n+1} \binom{2n}{n}$. In turn, the 0th column of \mathbf{B} is given by $(C_{n+1})_{n \geq 0}$. This similarity explains why they are called Catalan triangles.

According to the online encyclopedia of integer sequences (OEIS) [11], the non-zero entries of \mathbf{C} (sequence [A033184](#)) are given by the formula $C_{n,k} = \frac{k+1}{n+1} \binom{2n-k}{n}$, for $n \geq k \geq 0$. They are known as the (ordinary) ballot numbers. Aigner [2] used these numbers (and generalizations of them) to enumerate various combinatorial instances. Furthermore, Ferrari and Pinzani [8] gave an interpretation of \mathbf{C} , using the ECO method, and a suitable change of basis in the vector space of one-variable polynomials. On the other hand, the non-zero entries of \mathbf{B} ([A039598](#)) are given by the formula $B_{n,k} = \frac{k+1}{n+1} \binom{2n+2}{n-k}$, for $n \geq k \geq 0$. The numbers $B_{n,k}$ first appeared in the work of Shapiro [17, 19], in problems connected with non-intersecting paths, random walks, Eulerian numbers, runs, slides, and moments.³ We will refer from now on to \mathbf{C} as the Aigner array, and to \mathbf{B} as the Shapiro array.

The aim of this paper is to discuss some interesting features regarding the product matrix $\mathbf{R}(r) = \mathbf{R}\mathbf{P}(r)$, where \mathbf{R} is a given Riordan array, and the parameter r is any real or complex number. The notation $\mathbf{P}(r)$ stands for a generalized Pascal array, such that $\mathbf{P}(0) = \mathbf{I}$ is the identity matrix, and $\mathbf{P}(1) = \mathbf{P}$ is the classical Pascal array of binomial numbers, that comes arranged as a lower triangular matrix. We elaborate, in particular, on the product matrix $\mathbf{C}(r) = \mathbf{C}\mathbf{P}(r)$. Clearly, we have $\mathbf{C}(0) = \mathbf{C}$. It turns out that $\mathbf{C}(1) = \mathbf{B}$. The factorization $\mathbf{B} = \mathbf{C}\mathbf{P}$ resembles Barry's definition [3, see Sections 6.3 and 6.5] of the generalized ballot array given by $\mathbf{Bal} = \mathbf{Cat} \cdot \mathbf{Bin}$; where $\mathbf{Bin} = \mathbf{P}$, and $\mathbf{Cat} = \text{diag}(1, \mathbf{C})$.

Recently, Yang [23] introduced a generalized Catalan matrix, given by

$$C[a, b; r] = \left(\left(\frac{1 - \sqrt{1 - 4rz}}{2rz} \right)^a, z \left(\frac{1 - \sqrt{1 - 4rz}}{2rz} \right)^b \right),$$

where a, b are integer numbers, and r is an arbitrary parameter. Such a generalization provides a unified way of presenting the matrices \mathbf{C} and \mathbf{B} , along with many other matrices related to Catalan numbers. One can check that the 0th column of $C[a, b; r]$ forms a sequence

³Actually, Shapiro used the formula $B_{n,k} = \frac{k}{n} \binom{2n}{n+k}$, for $n, k \geq 1$. His formula is just a shifted version of the formula shown previously for $B_{n,k}$.

of degree n monomials in r , whose coefficients are polynomials in a and b , that have rational coefficients. The one-parameter Catalan triangles that we will study are instead given by

$$\mathbf{C}(r) = \left(\frac{1 - 2rz - \sqrt{1 - 4z}}{2z(1 - r + r^2z)}, \frac{1 - 2rz - \sqrt{1 - 4z}}{2(1 - r + r^2z)} \right).$$

We shall show explicitly that the 0^{th} column of $\mathbf{C}(r)$ forms a sequence $(\mathbf{C}(r)_{n,0})_{n \geq 0}$ of degree n monic polynomials in r . These polynomials have positive integer coefficients. In particular, the constant term of $\mathbf{C}(r)_{n,0}$ is the Catalan number C_n . For this reason, we shall refer to $\mathbf{C}(r)$ as a Catalan triangle too.

It is also worth mentioning that He [9] introduced recently, a family of Catalan triangles which depend on two parameters. His (c, r) -Catalan triangles are based on the sequence characterization of Bell-type Riordan arrays. We emphasize that they are also different from the Catalan triangles $\mathbf{C}(r)$. One way to see this difference is to compare, once again, the 0^{th} column of both types of matrices. On the one hand, one can check that the 0^{th} column of a (c, r) -Catalan triangle forms a sequence of degree n homogeneous polynomials in c and r , whose coefficients are the well-known Narayana numbers. On the other hand, the 0^{th} column of a Catalan triangle $\mathbf{C}(r)$ forms a sequence of one-variable polynomials of degree n , whose coefficients are the ordinary ballot numbers. Recall that both the Narayana and the ballot numbers are known to refine the Catalan numbers.

The fundamental theorem of Riordan arrays, and the isomorphism between Riordan arrays and Sheffer polynomials (the former being the coefficient matrices for the latter), allow us to interpret $\mathbf{C}(r)$ as a change of basis matrix in the space of polynomials in one variable. Thus, for instance, the Shapiro array \mathbf{B} is seen as the change of basis between the Chebyshev polynomials of the second kind, and the standard polynomial sequence of powers of one variable. In this way, by specifying values for the one-variable of the corresponding polynomials, several formulas related to periodic sequences, the sequence of natural numbers, and the sequence of Fibonacci numbers, can be stated and generalized.

2 General setting

The literature on Riordan arrays is vast and constantly growing (see, for instance, [1, 4, 5, 7, 9, 10, 12, 13, 14, 15, 16, 18, 20, 21, 22]). In this section we recall the fundamental theorem of Riordan arrays, and discuss some general facts regarding the generalized Sheffer polynomial sequence associated with any Riordan array. Then we define a one-parameter family of Riordan arrays, and focus our attention on Catalan arrays in Section 3.

2.1 Riordan Arrays

Let $d(z) = d_0 + d_1z + d_2z^2 + \dots$, and let $h(z) = h_0 + h_1z + h_2z^2 + \dots$ be two formal power series in z with coefficients in a given integral domain R , with unit 1. Assume that

$d_0 = d(0) \neq 0$, and $h_0 = h(0) = 0$. A *Riordan array* \mathbf{R} is an infinite lower triangular matrix, whose entries are given by

$$\mathbf{R}_{n,k} = [z^n]d(z)h(z)^k. \quad (1)$$

The operator $[z^n]$ acts on a formal power series $f(z) = \sum_{k \geq 0} f_k z^k$ by extracting its n^{th} coefficient; that is, we have $[z^n]f(z) = f_n$. It is customary to denote $\mathbf{R} = (d(z), h(z))$. In addition, the set of Riordan arrays for which $h'(0) \neq 0$, and that is equipped with the multiplication given by

$$(d_1(z), h_1(z))(d_2(z), h_2(z)) = (d_1(z)d_2(h_1(z)), h_2(h_1(z))), \quad (2)$$

forms a group. We denote this group by \mathfrak{Rio} . The group identity is the usual identity matrix $\mathbf{I} = (1, z)$. Furthermore, for any $\mathbf{R} \in \mathfrak{Rio}$, its inverse is given by $\mathbf{R}^{-1} = (1/d(\bar{h}(z)), \bar{h}(z))$, where $\bar{h}(z)$ denotes the compositional inverse of $h(z)$.

The *order* of a formal power series $g(z) = \sum_{k \geq 0} g_k z^k$ is the minimal index k such that $g_k \neq 0$. Since $R[[z]]$ denotes the ring of formal power series in z with coefficients in R , we will write $R_r[[z]]$ to denote the set of formal power series of order r . Therefore, if $\mathbf{R} = (d(z), h(z)) \in \mathfrak{Rio}$, then $d(z) \in R_0[[z]]$, and $h(z) \in R_1[[z]]$. For simplicity, and without loss of generality, we further assume that $d_0 = 1$, and $h_1 = h'(0) = 1$, so that $\mathbf{R}_{n,n} = h_1^n = 1$, for all $n \geq 0$.

Now, let us define an action $*$ of the group \mathfrak{Rio} on $R_0[[z]]$ by

$$(d(z), h(z)) * g(z) = d(z)g(h(z)). \quad (3)$$

Formula (3) represents the traditional matrix-column multiplication. Therefore, it is also equivalent to saying that, for any $n \geq 0$, the following identity holds

$$\sum_{k=0}^n \mathbf{R}_{n,k} g_k = [z^n]d(z)g(h(z)). \quad (4)$$

Equation (4) is known as the fundamental theorem of Riordan arrays. It is usually used to deal with sums and combinatorial identities. In particular, let $g(z) = 1 + xz + x^2z^2 + \dots = \frac{1}{1-xz}$. We write

$$p_n^{\mathbf{R}}(x) = \sum_{k=0}^n \mathbf{R}_{n,k} x^k = [z^n] \frac{d(z)}{1-xh(z)}. \quad (5)$$

We call $(p_n^{\mathbf{R}}(x))_{n \geq 0}$ the *Sheffer polynomial sequence* associated with \mathbf{R} . Formula (5) implies that the generating function of $p_n^{\mathbf{R}}(x)$ is given by

$$\sum_{n \geq 0} p_n^{\mathbf{R}}(x) z^n = \frac{d(z)}{1-xh(z)}. \quad (6)$$

Likewise, the generating function for the Sheffer polynomial associated with the inverse array $\mathbf{R}^{-1} = (1/d(\bar{h}(z)), \bar{h}(z))$ is

$$\sum_{n \geq 0} p_n^{\mathbf{R}^{-1}}(x) z^n = \frac{1}{d(\bar{h}(z))} \frac{1}{1-x\bar{h}(z)}. \quad (7)$$

Since \mathbf{R} is invertible, and the sequence $(x^n)_{n \geq 0}$ is a basis for the linear space $\mathbb{C}[x]$, the sequence $(p_n^{\mathbf{R}}(x))_{n \geq 0}$ is also a basis for $\mathbb{C}[x]$. Therefore, any $\mathbf{R} \in \mathfrak{Rio}$ can be interpreted as the change of basis matrix between the standard basis $(x^n)_{n \geq 0}$, and the basis of polynomials $(p_n^{\mathbf{R}}(x))_{n \geq 0}$. Hence, we have

$$\sum_{k=0}^n \mathbf{R}_{n,k} p_k^{\mathbf{R}^{-1}}(x) = x^n. \quad (8)$$

We will use Equation (8) in Section 4 to deduce some interesting combinatorial identities involving the Riordan arrays \mathbf{C} and \mathbf{B} .

2.2 One-parameter Riordan arrays

The classical Pascal array \mathbf{P} of binomial numbers $\binom{n}{k}$ is a Riordan array, since we can write $\mathbf{P} = (p(z), zp(z))$, where $p(z) = \frac{1}{1-z}$. Given any real number r , the Riordan array $\mathbf{P}(r) = (p(rz), zp(rz)) = \left(\frac{1}{1-rz}, \frac{z}{1-rz}\right)$ is known as the *generalized Pascal array of parameter r* . Its entries are given by $\mathbf{P}(r)_{n,k} = \binom{n}{k} r^{n-k}$.

Definition 1. Let \mathbf{R} be in \mathfrak{Rio} . The *r -Riordan array $\mathbf{R}(r)$* is defined by $\mathbf{R}(r) = \mathbf{R}\mathbf{P}(r)$.

Set $\mathbf{R} = (d(z), h(z))$. Then, by Equation (2), we have $\mathbf{R}(r) = \left(\frac{d(z)}{1-rh(z)}, \frac{h(z)}{1-rh(z)}\right)$. Using Equation (1), it is easy to check that the entries of $\mathbf{R}(r)$ are given by

$$\mathbf{R}(r)_{n,k} = [z^n] \frac{d(z)}{1-rh(z)} \left(\frac{h(z)}{1-rh(z)}\right)^k = [z^n] \frac{d(z)h(z)^k}{(1-rh(z))^{k+1}}.$$

Lemma 2. For any real number r , the Sheffer polynomial sequence $(p_n^{\mathbf{R}(r)}(x))_{n \geq 0}$ satisfies the identity

$$p_n^{\mathbf{R}(r)}(x) = p_n^{\mathbf{R}}(x+r).$$

Proof. By Equation (5), we have

$$p_n^{\mathbf{R}(r)}(x) = \sum_{k=0}^n \mathbf{R}(r)_{n,k} x^k = [z^n] \frac{\frac{d(z)}{1-rh(z)}}{1-x\frac{h(z)}{1-rh(z)}} = [z^n] \frac{d(z)}{1-(x+r)h(z)} = p_n^{\mathbf{R}}(x+r). \quad \square$$

Corollary 3. Let n be a non-negative integer. The constant term and the sum of coefficients of $p_n^{\mathbf{R}(r)}(x)$ are given by

$$p_n^{\mathbf{R}(r)}(0) = \mathbf{R}(r)_{n,0} = \sum_{k=0}^n \mathbf{R}_{n,k} r^k. \quad (9)$$

$$p_n^{\mathbf{R}(r)}(1) = \sum_{k=0}^n \mathbf{R}(r)_{n,k} = \mathbf{R}(r+1)_{n,0}. \quad (10)$$

This is a well-known result. In this sense, we say that the polynomial sequence $(\mathbf{C}(r)_{n,0})_{n \geq 0}$ is a refinement of the shifted Catalan sequence $(C_{n+1})_{n \geq 0}$.

Now, set $h(z) = zc(z)$. We have $z = \bar{h}(z)c(\bar{h}(z))$. It is easy to check that $\bar{h}(z) = z - z^2$. Hence, we have $\mathbf{C}^{-1} = (1/c(\overline{zc(z)}), \overline{zc(z)}) = (1 - z, z - z^2)$. Since $\mathbf{C}(r)^{-1} = \mathbf{P}(r)^{-1}\mathbf{C}^{-1}$, it follows that

$$\mathbf{C}(r)^{-1} = \left(\frac{1}{1+rz}, \frac{z}{1+rz} \right) (1 - z, z - z^2) = \left(\frac{1 + (r-1)z}{(1+rz)^2}, \frac{z + (r-1)z^2}{(1+rz)^2} \right). \quad (13)$$

Lemma 6. *The entries of $\mathbf{C}(r)^{-1}$ are given by*

$$\mathbf{C}(r)_{n,k}^{-1} = [z^{n-k}] \left(\frac{1 + (r-1)z}{(1+rz)^2} \right)^{k+1} = (-1)^{n-k} \sum_{i=0}^{n-k} \binom{i+k+1}{n-i-k} \binom{i+k}{k} r^i. \quad (14)$$

Lemma 5 and Lemma 6 show that both $\mathbf{C}(r)_{n,k}$ and $(-1)^{n-k}\mathbf{C}(r)_{n,k}^{-1}$ are polynomials in r with positive integer coefficients. In addition, note that the polynomial $\mathbf{C}(r)_{n+1,1}$ is of degree n , has constant term C_{n+1} , and its leading coefficient is $n+1$. Both lemmas are useful to deduce several formulas and combinatorial identities. For instance, letting $r = 0$ in Equation (14), we obtain

$$\mathbf{C}_{n,k}^{-1} = [z^{n-k}] (1-z)^{k+1} = \binom{n-2k-2}{n-k} = (-1)^{n-k} \binom{k+1}{n-k}.$$

Similarly, letting $r = 1$ in Equation (14), we get

$$\begin{aligned} \mathbf{B}_{n,k}^{-1} &= [z^{n-k}] \left(\frac{1}{(1+z)^2} \right)^{k+1} = (-1)^{n-k} \binom{n+k+1}{n-k} \\ &= (-1)^{n-k} \sum_{i=0}^{n-k} \binom{i+k+1}{n-i-k} \binom{i+k}{k}. \end{aligned} \quad (15)$$

Thus, the following formula is immediate.

Lemma 7.

$$\binom{n+k+1}{n-k} = \sum_{i=0}^{n-k} \binom{i+k+1}{n-i-k} \binom{i+k}{k}.$$

The next non-trivial combinatorial identity follows directly from Equation (10) and Equation (12).

Lemma 8.

$$\sum_{k=0}^n \sum_{i=0}^{n-k} \frac{i+k+1}{n+1} \binom{2n-i-k}{n} \binom{i+k}{k} r^i = \sum_{i=0}^n \frac{i+1}{n+1} \binom{2n-i}{n} (r+1)^i.$$

As for the generating functions for the Sheffer polynomials associated with $\mathbf{C}(r)$ and $\mathbf{C}(r)^{-1}$, the next formulas are straightforward consequences of Equation (6) and Equation (7), using Formula (11) and Formula (13), respectively.

Proposition 9.

$$\begin{aligned} \sum_{n \geq 0} p_n^{\mathbf{C}(r)}(x)z^n &= \frac{c(z)}{1 - (x+r)zc(z)} \\ \sum_{n \geq 0} p_n^{\mathbf{C}(r)^{-1}}(x)z^n &= \frac{1 + (r-1)z}{1 + (2r-x)z + (r^2 - (r-1)x)z^2} \end{aligned} \quad (16)$$

In Section 4, we will use Proposition 9 to obtain more combinatorial identities.

4 Special identities

4.1 Identities involving the Aigner array

Set $r = 0$ and $x = -1$ in Equation (16). We obtain

$$\begin{aligned} 1 + \sum_{n \geq 1} p_n^{\mathbf{C}^{-1}}(-1)(-z)^n &= \frac{1+z}{1-z-z^2} \\ &= 1 + 2z + 3z^2 + 5z^3 + 8z^4 + 13z^5 + 21z^6 + 34z^7 + 55z^8 + \dots \\ &= F_2 + F_3z + F_4z^2 + F_5z^3 + \dots, \end{aligned}$$

where F_n is the n^{th} Fibonacci number, for $n \geq 2$. Hence, for $n \geq 0$, it holds that

$$(-1)^n p_n^{\mathbf{C}^{-1}}(-1) = F_{n+2}.$$

Since $p_n^{\mathbf{C}^{-1}}(x) = \sum_{k=0}^n \mathbf{C}_{n,k}^{-1} x^k$, we have

$$\mathbf{C}^{-1} \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \\ \vdots \end{pmatrix} = \begin{pmatrix} F_2 \\ -F_3 \\ F_4 \\ -F_5 \\ \vdots \end{pmatrix}, \quad \text{or equivalently,} \quad \mathbf{C} \begin{pmatrix} F_2 \\ -F_3 \\ F_4 \\ -F_5 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \\ \vdots \end{pmatrix}.$$

Then

$$\sum_{k=0}^n \binom{k+1}{n-k} = F_{n+2}, \quad \text{and} \quad \sum_{k=0}^n \frac{k+1}{n+1} \binom{2n-k}{n} (-1)^{n-k} F_{k+2} = 1.$$

4.2 Identities involving the Shapiro array

Set $r = 1$ in Equation (16). Shapiro, Woan, and Getu [19] observed that the Catalan triangle \mathbf{B} verified the following interesting identity:

$$\mathbf{B} \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \\ 4^2 \\ 4^3 \\ 4^4 \\ \vdots \end{pmatrix}. \quad (17)$$

The problem of finding a matrix identity that extends the sequence $(1, 4, 4^2, 4^3, \dots)$ to the sequence $(1, k, k^2, k^3, \dots)$ in Equation (17), was studied by Chen, Li, Shapiro and Yan [5]. They generalized the following known recurrence relation for \mathbf{B} ; namely,

$$\mathbf{B}_{n,k} = \mathbf{B}_{n-1,k-1} + 2\mathbf{B}_{n-1,k} + \mathbf{B}_{n-1,k+1},$$

by constructing a matrix \mathbf{M} that satisfies the identity

$$\mathbf{M} \begin{pmatrix} 1 \\ 1+t \\ 1+t+t^2 \\ 1+t+t^2+t^3 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ k \\ k^2 \\ k^3 \\ \vdots \end{pmatrix}.$$

The first column of \mathbf{M} (that is, its 0th column) was then interpreted in terms of weighted partial Motzkin paths. Here, we approach the problem of extending the sequence $(1, 4, 4^2, 4^3, \dots)$ to the sequence $(1, k, k^2, k^3, \dots)$, by keeping $\mathbf{C}(1) = \mathbf{B}$ fixed, and using Sheffer sequences. More explicitly, by Equation (16), we have

$$1 + \sum_{n \geq 1} p_n^{\mathbf{B}^{-1}}(x) z^n = \frac{1}{1 - (x-2)z + z^2}.$$

Furthermore, by Equation (8), we also have

$$\mathbf{B} \begin{pmatrix} 1 \\ p_1^{\mathbf{B}^{-1}}(x) \\ p_2^{\mathbf{B}^{-1}}(x) \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ x \\ x^2 \\ \vdots \end{pmatrix}, \quad \text{or equivalently,} \quad \mathbf{B}^{-1} \begin{pmatrix} 1 \\ x \\ x^2 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ p_1^{\mathbf{B}^{-1}}(x) \\ p_2^{\mathbf{B}^{-1}}(x) \\ \vdots \end{pmatrix}. \quad (18)$$

By Identity (15), the polynomial $p_n^{\mathbf{B}^{-1}}(x)$ is given by

$$p_n^{\mathbf{B}^{-1}}(x) = \sum_{k=0}^n \mathbf{B}_{n,k}^{-1} x^k = \sum_{k=0}^n (-1)^{n-k} \binom{n+k+1}{n-k} x^k. \quad (19)$$

Note that $p_n^{\mathbf{B}^{-1}}(x) = U_n\left(\frac{x-2}{2}\right)$, where $(U_n(x))_{n \geq 0}$ denotes the sequence of Chebyshev polynomials of the second kind [6]. Moreover, the sequence $(p_n^{\mathbf{B}^{-1}}(x))_{n \geq 0}$ satisfies the three-term recursion

$$p_n^{\mathbf{B}^{-1}}(x) = (x-2)p_{n-1}^{\mathbf{B}^{-1}}(x) - p_{n-2}^{\mathbf{B}^{-1}}(x), \quad n \geq 2,$$

with initial values $p_0^{\mathbf{B}^{-1}}(x) = 1$ and $p_1^{\mathbf{B}^{-1}}(x) = x-2$. Whenever $x \in \mathbb{Z}$, the polynomial sequence $(p_n^{\mathbf{B}^{-1}}(x))_{n \geq 0}$ reduces to an integer sequence. By specifying the values of x in Equation (18), we obtain many interesting relations.

Example 10 (Periodic sequences). When $x = 2$, the recursion $p_n^{\mathbf{B}^{-1}}(2) = -p_{n-2}^{\mathbf{B}^{-1}}(2)$, with initial values $p_0^{\mathbf{B}^{-1}}(2) = 1$ and $p_1^{\mathbf{B}^{-1}}(2) = 0$, is solved by the period-4 sequence $1, 0, -1, 0$. Hence, we have

$$\begin{pmatrix} 1 & & & & & & \\ 2 & 1 & & & & & \\ 5 & 4 & 1 & & & & \\ 14 & 14 & 6 & 1 & & & \\ 42 & 48 & 27 & 8 & 1 & & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \\ 1 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 2^2 \\ 2^3 \\ 2^4 \\ \vdots \end{pmatrix}.$$

When $x = 3$, the recursion $p_n^{\mathbf{B}^{-1}}(3) = p_{n-1}^{\mathbf{B}^{-1}}(3) - p_{n-2}^{\mathbf{B}^{-1}}(3)$, satisfying the initial conditions $p_0^{\mathbf{B}^{-1}}(3) = p_1^{\mathbf{B}^{-1}}(3) = 1$, is solved by the period-6 sequence $1, 1, 0, -1, -1, 0$. Thus, we have

$$\begin{pmatrix} 1 & & & & & & \\ 2 & 1 & & & & & \\ 5 & 4 & 1 & & & & \\ 14 & 14 & 6 & 1 & & & \\ 42 & 48 & 27 & 8 & 1 & & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 0 \\ -1 \\ -1 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ 3^2 \\ 3^3 \\ 3^4 \\ \vdots \end{pmatrix}.$$

These two cases are better understood by recalling that $U_n(\cos y) = \frac{\sin((n+1)y)}{\sin y}$. Hence, we have $p_n^{\mathbf{B}^{-1}}(2) = U_n\left(\cos \frac{\pi}{2}\right)$ and $p_n^{\mathbf{B}^{-1}}(3) = U_n\left(\cos \frac{\pi}{3}\right)$. These identities explain why we obtained periodic sequences.

Example 11 (Natural numbers). When $x = 4$, the recursion $p_n^{\mathbf{B}^{-1}}(4) = 2p_{n-1}^{\mathbf{B}^{-1}}(4) - p_{n-2}^{\mathbf{B}^{-1}}(4)$, with initial values $p_0^{\mathbf{B}^{-1}}(4) = 1$, $p_1^{\mathbf{B}^{-1}}(4) = 2$, is solved by the sequence $(p_n^{\mathbf{B}^{-1}}(4))_{n \geq 0} =$

$(n+1)_{n \geq 0}$. Thus, we obtain the Identity (17):

$$\begin{pmatrix} 1 \\ 2 & 1 \\ 5 & 4 & 1 \\ 14 & 14 & 6 & 1 \\ 42 & 48 & 27 & 8 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \\ 4^2 \\ 4^3 \\ 4^4 \\ \vdots \end{pmatrix}.$$

The previous matrix identity is equivalent to Chen's combinatorial formula [5]:

$$\sum_{k=0}^n \frac{(k+1)^2}{n+1} \binom{2n+2}{n-k} = 4^n.$$

Example 12 (Fibonacci numbers). When $x = 5$, the recursion $p_n^{\mathbf{B}^{-1}}(5) = 3p_{n-1}^{\mathbf{B}^{-1}}(5) - p_{n-2}^{\mathbf{B}^{-1}}(5)$, with initial conditions $p_0^{\mathbf{B}^{-1}}(5) = 1$ and $p_1^{\mathbf{B}^{-1}}(5) = 3$, is solved by the subsequence of Fibonacci numbers $(p_n^{\mathbf{B}^{-1}}(5))_{n \geq 0} = (F_{2n+2})_{n \geq 0}$. Thus, Shapiro's triangle \mathbf{B} , and the Fibonacci numbers of even index, are related by the following identity:

$$\begin{pmatrix} 1 \\ 2 & 1 \\ 5 & 4 & 1 \\ 14 & 14 & 6 & 1 \\ 42 & 48 & 27 & 8 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} 1 \\ 3 \\ 8 \\ 21 \\ 55 \\ \vdots \end{pmatrix} = \begin{pmatrix} 1 \\ 5 \\ 5^2 \\ 5^3 \\ 5^4 \\ \vdots \end{pmatrix}.$$

Equivalently, we have

$$\sum_{k=0}^n \frac{k+1}{n+1} \binom{2n+2}{n-k} F_{2k+2} = 5^n. \quad (20)$$

Using the well-known binomial recurrence relation $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$ in Equation (19), and then inverting Formula (20), we obtain

$$F_{2n+2} = (-1)^n \sum_{k=0}^n \left(\binom{n+k+2}{n-k} - \binom{n+k+1}{n-k-1} \right) (-5)^k.$$

Moreover, seeing as $F_{2n+3} = F_{2(n+1)+2} - F_{2n+2}$, we can deduce an analogous formula for the Fibonacci numbers of odd index; that is

$$F_{2n+1} = (-1)^n \sum_{k=0}^n \left(\binom{n+k+2}{n-k} - \binom{n+k}{n-k-2} \right) (-5)^k.$$

5 Acknowledgements

The authors wish to thank the editor and the referee for their helpful comments.

References

- [1] J. Agapito, Â. Mestre, P. Petruccio, and M. M. Torres, A symbolic treatment of Riordan arrays, *Linear Algebra Appl.* **439** (2013), 1700–1715.
- [2] M. Aigner, Enumeration via ballot numbers, *Discrete Math.* **308** (2008), 2544–2563.
- [3] P. Barry, A study of integer sequences, Riordan arrays, Pascal-like arrays and Hankel transforms, PhD thesis, University College Cork, 2009.
- [4] E. H. M. Brietzke, An identity of Andrews and a new method for the Riordan array proof of combinatorial identities, *Discrete Math.* **308** (2008), 4246–4262.
- [5] W. Y. C. Chen, N. Y. Li, L. W. Shapiro, and S. H. F. Yan, Matrix identities on weighted partial Motzkin paths, *European J. Combin.* **28** (2007), 1196–1207.
- [6] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Riedel, Dordrecht, 1974.
- [7] G. della Riccia, Riordan arrays, Sheffer sequences and orthogonal polynomials, *J. Integer Seq.* **11** (2008), Article 08.5.3.
- [8] L. Ferrari and R. Pinzani, Catalan-like numbers and succession rules, *Pure Math. Appl.* **16** (2005), 229–250.
- [9] T.-X. He, Parametric Catalan numbers and Catalan triangles, *Linear Algebra Appl.* **438** (2013), 1467–1484.
- [10] T.-X. He, L. C. Hsu, and P. J.-S. Shiue, The Sheffer group and the Riordan group, *Discrete Appl. Math.* **155** (2007), 1895–1909.
- [11] O.E.I.S. Foundation Inc., The Online Encyclopedia of Integer Sequences. <http://oeis.org>, 2011.
- [12] A. Luzón, D. Merlini, M. A. Morón, and R. Sprugnoli, Identities induced by Riordan arrays, *Linear Algebra Appl.* **436** (2012), 631–647.
- [13] A. Luzón and M. A. Morón, Recurrence relations for polynomial sequences via Riordan matrices, *Linear Algebra Appl.* **433** (2010), 1422–1446.
- [14] D. Merlini, D. G. Rogers, R. Sprugnoli, and M. C. Verri, On some alternative characterizations of Riordan arrays, *Canad. J. Math.* **49** (1997), 301–320.

- [15] D. Merlini and R. Sprugnoli, Algebraic aspects of some Riordan arrays related to binary words avoiding a pattern, *Theor. Comput. Sci.* **412** (2011), 2988–3001.
- [16] D. G. Rogers, Pascal triangles, Catalan numbers and renewal arrays, *Discrete Math.* **22** (1978), 301–310.
- [17] L. W. Shapiro, A Catalan triangle, *Discrete Math.* **14** (1976), 83–90.
- [18] L. W. Shapiro, S. Getu, W.-J. Woan, and L. C. Woodson, The Riordan group, *Discrete Appl. Math.* **34** (1991), 229–239.
- [19] L. W. Shapiro, W. Woan, and S. Getu, Runs, slides and moments, *SIAM J. Alg. Disc. Math.* **4** (1983), 459–466.
- [20] R. Sprugnoli, Riordan arrays and combinatorial sums, *Discrete Math.* **132** (1994), 267–290.
- [21] R. Sprugnoli, Riordan arrays and the Abel-Gould identity, *Discrete Math.* **142** (1995), 213–233.
- [22] W. Wang and T. Wang, Generalized Riordan arrays, *Discrete Math.* **308** (2008), 6466–6500.
- [23] S.-L. Yang, Some inverse relations determined by Catalan matrices, *Int. J. Comb.* (2013), Art. ID 528584.

2010 *Mathematics Subject Classification*: Primary 15B; Secondary 05A19, 11B37, 11B83, 11C.

Keywords: Riordan array, Pascal array, Catalan triangle, Sheffer polynomial, combinatorial identity, recurrence, Chebyshev polynomial, Fibonacci number.

(Concerned with sequence [A001700](#), [A008549](#), [A033184](#), [A035330](#), [A039598](#), [A045720](#), [A045894](#), [A049027](#), and [A076025](#).)
