Snow Leopard Permutations and their Even and Odd Threads

Eric S. Egge^{1*}

Kailee Rubin²

¹ Carleton College, Northfield, MN, USA

² Epic, Madison, WI, USA

received 24th Aug. 2015, revised 3rd May 2016, accepted 12th May 2016.

Caffrey, Egge, Michel, Rubin and Ver Steegh recently introduced snow leopard permutations, which are the anti-Baxter permutations that are compatible with the doubly alternating Baxter permutations. Among other things, they showed that these permutations preserve parity, and that the number of snow leopard permutations of length 2n - 1is the Catalan number C_n . In this paper we investigate the permutations that the snow leopard permutations induce on their even and odd entries; we call these the *even threads* and the *odd threads*, respectively. We give recursive bijections between these permutations and certain families of Catalan paths. We characterize the odd (resp. even) threads which form the other half of a snow leopard permutation whose even (resp. odd) thread is layered in terms of pattern avoidance, and we give a constructive bijection between the set of permutations of length n which are both even threads and odd threads and the set of peakless Motzkin paths of length n + 1.

Keywords: Baxter permutation, Catalan path, Motzkin path, snow leopard permutation

1 Introduction

A complete Baxter permutation π , as introduced by Baxter (1964) and characterized by Chung et al. (1978), is a permutation of length 2n + 1 such that, for all i with $1 \le i \le 2n + 1$,

- $\pi(i)$ is even if and only if *i* is even and
- if $\pi(x) = i$, $\pi(y) = i + 1$, and z is between x and y (that is, x < z < y or y < z < x), then $\pi(z) < i$ if i is odd and $\pi(z) > i + 1$ if i is even.

As Chung and her coauthors note, each complete Baxter permutation is uniquely determined by its odd entries. Accordingly, for each complete Baxter permutation π of length 2n+1, the associated *reduced Baxter permutation* of length n + 1 is the permutation π induces on its odd entries. For example, 981254367 is a complete Baxter permutation of length nine, whose associated reduced Baxter permutation is 51324. One can show that a permutation is a reduced Baxter permutation if and only if it avoids the vincular patterns 3142 and 2413. That is, if π has length n, then π is a reduced Baxter permutation whenever

ISSN 1365–8050 © 2016 by the author(s) Distributed under a Creative Commons Attribution 4.0 International License

^{*}Corresponding author.

there are no indices $1 \le i < j < j + 1 < k < n$ such that $\pi(j) < \pi(k) < \pi(i) < \pi(j + 1)$ (for $3\underline{142}$) or $\pi(j + 1) < \pi(i) < \pi(k) < \pi(j)$ (for $2\underline{413}$). For example, the complete Baxter permutation $3\ 2\ 1\ 4\ 13\ 12\ 7\ 8\ 11\ 10\ 9\ 6\ 5$ has reduced Baxter permutation 2174653, which has no instances of $3\underline{142}$ or $2\underline{413}$. In contrast, 4613752 is not a reduced Baxter permutation because the subsequence 6375 is an instance of $3\underline{142}$. We sometimes refer to reduced Baxter permutations as *Baxter permutations* for short.

For each complete Baxter permutation π of length 2n + 1, the associated *anti-Baxter permutation* is the permutation of length n that π induces on its even entries. Although complete Baxter permutations are determined by their associated Baxter permutations, they are not determined by their associated anti-Baxter permutations: many anti-Baxter permutations appear in several complete Baxter permutations. For example, 4123 is the anti-Baxter permutation associated with both of the complete Baxter permutations 981254367 and 983214765. On the other hand, anti-Baxter permutations are characterized by pattern avoidance conditions nearly identical to those which characterize the Baxter permutations: a permutation π is anti-Baxter if and only if it avoids the vincular patterns 3412 and 2143. That is, if π has length n, then π is anti-Baxter whenever there are no indices $1 \le i < j < j + 1 < k < n$ such that $\pi(j+1) < \pi(k) <$ $\pi(i) < \pi(j)$ (for 3412) or $\pi(j) < \pi(i) < \pi(k) < \pi(j + 1)$ (for 2143). We say a Baxter permutation π_1 and an anti-Baxter permutation π_2 are *compatible* whenever there is a complete Baxter permutation π such that π_1 and π_2 are the permutations induced on the odd and even entries of π , respectively. One can show that this definition of compatibility is equivalent to the definition of compatibility given by Caffrey et al. (2015).

Baxter first introduced the permutations that now bear his name in connection with a problem involving fixed points of commuting continuous functions on the closed interval [0, 1], but they have also appeared in a variety of other settings. One such setting involves the *Aztec diamond* of order n, which is an array of unit squares with 2i squares in row i for $1 \le i \le n$ and 2(2n - i + 1) squares in row i for $n < i \le 2n$, in which the squares are centered in each row. In Figure 1 we have the Aztec diamond of order three. The




Fig. 1: The Aztec diamond of order 3.

entries of our complete Baxter permutations will be at the vertices of an Aztec diamond. We prefer to arrange these vertices in rows and columns, so we will orient all of our Aztec diamonds as in Figure 2.

Aztec diamonds can be tiled by dominos, and Elkies et al. (1992) have described how to construct, for each such tiling of the Aztec diamond of order n, a pair of matrices SASM(T) and LASM(T) of sizes $n \times n$ and $(n + 1) \times (n + 1)$, respectively. Each of these matrices is an *alternating-sign matrix* (ASM), which is a matrix with entries in $\{0, 1, -1\}$ whose nonzero entries in each row and in each column alternate in sign and sum to one. (Note that SASM and LASM above are short for small ASM and large




Fig. 2: The Aztec diamond of order 3, reoriented.

ASM, respectively. For an introduction to ASMs and a variety of related combinatorial objects, see the work of Robbins (1991), Bressoud (1999), and Propp (2001).) To carry out this construction, first note that a tiling of an Aztec diamond with dominos gives rise to a graph whose vertices are lattice points, as in Figure 3. Each exterior vertex of this graph has the same degree in all domino tilings, so we disregard




Fig. 3: The graph of a domino tiling of the Aztec diamond of order 3.

these vertices. The remaining vertices fall naturally into two sets as in Figure 4: the black vertices form an $(n+1) \times (n+1)$ matrix while the white vertices form an $n \times n$ matrix. We construct LASM(T) on the black vertices by labeling each vertex of degree four with a 1, labeling each vertex of degree three with a 0, and labeling each vertex of degree two with a -1. We construct SASM(T) on the white vertices by labeling each vertex of degree two with a -1. We construct SASM(T) on the white vertices by labeling each vertex of degree four with a -1. We construct SASM(T) on the white vertices by labeling each vertex of degree four with a -1. The construct SASM(T) on the white vertices by labeling each vertex of degree four with a -1. The construct SASM(T) on the white vertices by labeling each vertex of degree four with a -1. The construct SASM(T) on the white vertices by labeling each vertex of degree four with a -1. The construct SASM(T) on the white vertices by labeling each vertex of degree four with a -1, labeling each vertex of degree three

$$LASM(T) = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$
$$SASM(T) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

and

Canary (2010) has shown that LASM(T) is a permutation matrix if and only if it is the matrix for a Baxter permutation. In this case, if we now introduce gray dots inside the dominos as in Figure 5, and




Fig. 4: The graph of a domino tiling of the Aztec diamond of order 3 with outer vertices removed.





Fig. 5: The entries of the associated complete Baxter permutation are at the black, white, and gray dots.

assign 0s to each of them, then the black, white, and gray dots together form the matrix for a complete Baxter permutation.

While Baxter permutations have been studied in several other combinatorial incarnations (see, for example, the papers of Asinowski et al. (2013) and Dilks (2014)), we focus on a certain subset of the Baxter permutations, and their compatible anti-Baxter permutations. Specifically, an *alternating permutation*, or an *up-down permutation*, is a permutation which begins with an ascent, and in which ascents and descents alternate. Similarly, a *doubly alternating permutation* is an alternating permutation whose inverse is also alternating. Guibert and Linusson (2000) have shown that the number of doubly alternating Baxter permutations of length 2n is the Catalan number $C_n = \frac{1}{n+1} \binom{2n}{n}$, as is the number of doubly alternating Baxter permutations.

Definition 1.1. A snow leopard permutation is an anti-Baxter permutation which is compatible with a doubly alternating Baxter permutation.

Caffrey et al. (2015) show that compatibility is a bijection between the set of doubly alternating Baxter permutations of length n and the set of snow leopard permutations of length n - 1. They also give a simple bijection between the set of snow leopard permutations of length 2n and the set of snow leopard permutations of length 2n - 1 (see Theorem 3.1), so we focus our attention on the snow leopard permutations of odd length. We write SL_n to denote the set of snow leopard permutations of length 2n - 1; in Table 1 we list the snow leopard permutations of lengths one, three, and five.



Tab. 1: The snow leopard permutations of lengths one, three, and five.

Like the complete Baxter permutations, and as Caffrey et al. (2015) show, the snow leopard permutations preserve parity. In this paper, we study the permutations the snow leopard permutations induce on their odd and even entries; we call these induced permutations odd threads and even threads, respectively. In Section 2 we recall some useful permutation tools. In Section 3 we give recursive decompositions of the even and odd threads, and we explore the extent to which these decompositions are unique. In Section 4 we use our decompositions to give recursive bijections between the set of even threads of length n and the set of Catalan paths of length n + 1 with no ascent of length two, and between the set of odd threads of length n and the set of Catalan paths of length n with no four consecutive steps NEEN. In Section 5 we begin to describe which even and odd threads are induced by the same snow leopard permutation. In particular, we show that the even threads which can be paired with the increasing odd thread to form a snow leopard permutation are exactly the 3412-avoiding involutions, we show that the odd threads which can be paired with the decreasing even thread to form a snow leopard permutation are exactly the complements of the 3412-avoiding involutions, and we extend these results in a natural way to layered even and odd threads. Finally, in Section 6 we give a constructive bijection between the set of permutations of length n which are both even threads and odd threads and the set of peakless Motzkin paths of length n+1.

2 Permutation Tools

Throughout we write S_n denote the set of all permutations of length n, written in one-line notation, and for any permutation π we write $|\pi|$ to denote the length of π . The complement operation on permutations will be useful for us, so we recall it next.

Definition 2.1. Following Kitaev (2011), for any permutation $\pi \in S_n$, we write $c(\pi)$ to denote the complement of π , which is the permutation in S_n with

$$c(\pi)(j) = n + 1 - \pi(j)$$

for all $j, 1 \leq j \leq n$.

We will also make extensive use of the following two ways of combining two permutations.

Definition 2.2. For permutations $\pi \in S_n$ and $\sigma \in S_m$ we write $\pi \oplus \sigma$ to denote the permutation in S_{n+m} with

$$(\pi \oplus \sigma)(j) = \begin{cases} \pi(j) & \text{if } 1 \le j \le n \\ n + \sigma(j - n) & \text{if } n + 1 \le j \le n + m \end{cases}$$

for all $j, 1 \leq j \leq n$, and we write $\pi \ominus \sigma$ to denote the permutation in S_{n+m} with

$$(\pi \ominus \sigma)(j) = \begin{cases} m + \pi(j) & \text{if } 1 \le j \le n \\ \sigma(j-n) & \text{if } n+1 \le j \le n+m \end{cases}$$

for all $j, 1 \leq j \leq n$.

Example 2.3. If $\pi = 14235$ and $\sigma = 312$ then $c(\pi) = 52431$, $\pi \oplus \sigma = 14235867$, and $\pi \oplus \sigma = 47568312$.

At times we will be working with permutations of length 0 or -1. For these, we use the following convention.

Definition 2.4. We write \emptyset to denote the empty permutation, which is the unique permutation of length 0, and we write @ to denote the antipermutation, which is the unique permutation of length -1. We have c(@) = @ and $1 \oplus @ = @ \oplus 1 = 1 \oplus @ = @ \oplus 1 = \emptyset$.

Note that 1 is a complete Baxter permutation, whose corresponding (reduced) Baxter permutation is also 1. Since 1 is also doubly alternating, \emptyset (its compatible anti-Baxter permutation) is a snow leopard permutation. Similarly, \emptyset is a complete Baxter permutation, whose corresponding Baxter permutation is also \emptyset . Since \emptyset is also doubly alternating and has length 0, its compatible anti-Baxter permutation is @. In particular, @ is a snow leopard permutation.

3 Even Threads, Odd Threads, and their Decompositions

Our starting point, and one of the main results of Caffrey et al. (2015), is the following recursive decomposition of the snow leopard permutations of odd positive length.

Theorem 3.1. (*Caffrey et al., 2015, Thm. 2.20 and Prop. 2.22*) For any permutation π of odd positive length, π is a snow leopard permutation if and only if there exist snow leopard permutations π_1 and π_2 of odd length such that $\pi = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2$. For any permutation σ of even length, σ is a snow leopard permutation if and only if $\sigma = 1 \oplus c(\sigma_1)$ for some snow leopard permutation σ_1 . In addition, these decompositions are uniquely determined by π and σ , respectively.

We use the term *connector* to refer to the entry in π corresponding to the final 1 in the decomposition of π . From the recursive decomposition of π in Theorem 3.1, we see that the connector of π will always equal $\pi(1) - 1$ if $\pi(1) > 1$. When $\pi(1) = 1$, the permutation will not have a connector. Note that when a snow leopard permutation π does have a connector, the corresponding entry in $c(\pi)$ will be a left-to-right maximum and a fixed point in $c(\pi)$.

Example 3.2. Let $\pi = 587694321 = (1 \oplus c(123) \oplus 1) \oplus 1 \oplus 321$. Since both 123 and 321 are snow leopard permutations, π must also be a snow leopard permutation, with connector $\pi(6) = 4$. Here, $c(\pi) = 523416789$ so $c(\pi)(6) = 6$ is both a left-to-right maximum and a fixed point in $c(\pi)$.

Note that for snow leopard permutations of lengths one and three we have

$$1 = 1 \oplus c(@) \oplus 1,$$
$$123 = 1 \oplus c(1) \oplus 1,$$

and

$$1 = (1 \oplus c(@) \oplus 1) \ominus 1 \ominus (1 \oplus c(@) \oplus 1),$$

so by using Theorem 3.1 inductively we get the following natural "block decomposition" for snow leopard permutations of odd positive length.

Snow Leopard Permutations and their Even and Odd Threads

Corollary 3.3. For any permutation π of odd positive length, π is a snow leopard permutation if and only if there is a sequence π_1, \ldots, π_k of snow leopard permutations of odd length such that

$$\pi = (1 \oplus c(\pi_1) \oplus 1) \ominus 1 \ominus (1 \oplus c(\pi_2) \oplus 1) \ominus 1 \ominus \cdots \ominus 1 \ominus (1 \oplus c(\pi_k) \oplus 1).$$

In addition, the sequence π_1, \ldots, π_k is uniquely determined by π .

Theorem 3.1 tells us that the set of snow leopard permutations of odd length is closed under a complicated operation, but we might hope it is also closed under some simpler operation. Unfortunately, this set is not closed under \oplus or \ominus : neither $321 \oplus 123 = 654123$ nor $321 \oplus 123 = 321456$ are snow leopard permutations, even though 321 and 123 are. However, we do have the following result.

Theorem 3.4. Suppose π and σ are snow leopard permutations of odd length. Then $\pi \ominus 1 \ominus \sigma$ is also a snow leopard permutation.

Proof: Suppose π and σ are snow leopard permutations of odd length with $|\pi| = 2n - 1$. We argue by induction on n.

When n = 0 we have $\pi = @$ and $\pi \ominus 1 \ominus \sigma = @ \ominus 1 \ominus \sigma = \sigma$, which is a snow leopard permutation. When n = 1 we have $\pi = 1$ and

$$\begin{aligned} \pi \ominus 1 \ominus \sigma &= 1 \ominus 1 \ominus \sigma \\ &= (1 \oplus @ \oplus 1) \ominus 1 \ominus \sigma, \end{aligned}$$

which is a snow leopard permutation by Theorem 3.1.

Now suppose $\pi \ominus 1 \ominus \sigma$ is a snow leopard permutation whenever $|\pi| < 2k - 1$ and let $|\pi| = 2k - 1$. Since π is a snow leopard permutation of odd positive length, by Theorem 3.1 there are snow leopard permutations π_1 and π_2 with

$$\pi \ominus 1 \ominus \sigma = ((1 \oplus c(\pi_1) \oplus 1) \ominus 1 \ominus \pi_2) \ominus 1 \ominus \sigma$$

= $(1 \oplus c(\pi_1) \oplus 1) \ominus 1 \ominus (\pi_2 \ominus 1 \ominus \sigma).$

Since $|\pi| = 2k - 1$ and $|(1 \oplus c(\pi_1) \oplus 1) \oplus 1| \ge 2$, we must have $|\pi_2| < 2k - 1$. Now, since π_2 and σ are snow leopard permutations with $|\pi_2| < 2k - 1$, $\pi_2 \oplus 1 \oplus \sigma$ is a snow leopard permutation by induction. Thus, $\pi \oplus 1 \oplus \sigma = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_3$, where π_3 is a snow leopard permutation, so $\pi \oplus 1 \oplus \sigma$ is a snow leopard permutation by Theorem 3.1.

Caffrey et al. (2015) show in their Corollary 2.24 that snow leopard permutations preserve parity. That is, for any snow leopard permutation π and for all j with $1 \le j \le |\pi|$, we know that $\pi(j)$ is even if and only if j is even. We can separate any parity-preserving permutation into two smaller permutations as follows.

Definition 3.5. For any permutation π which preserves parity, we write π^e to denote the permutation π induces on its even entries and we write π^o to denote the permutation π induces on its odd entries. Note that if $|\pi| = 2n + 1$ then $|\pi^e| = n$ and $|\pi^o| = n + 1$.

Although in general we will only consider snow leopard permutations of positive odd length, we include one special case. If $\pi = @$, then since $|\pi| = -1$, we must have $|\pi^o| = 0$ and $|\pi^e| = -1$. This gives us $\pi^o = \emptyset$ and $\pi^e = @$.

Definition 3.6. We say a permutation σ is an even thread (resp. odd thread) whenever there is a snow leopard permutation π of odd length such that $\pi^e = \sigma$ (resp. $\pi^o = \sigma$), and we say an even thread α and an odd thread β are entangled whenever there is a snow leopard permutation π with $\pi_e = \alpha$ and $\pi^o = \beta$. We write ET_n (resp. OT_n) to denote the set of even (resp. odd) threads of length n.

Example 3.7. The permutation $\pi = 587694321$ is a snow leopard permutation with $\pi^o = 34521$ and $\pi^e = 4321$, so 34521 is an odd thread, 4321 is an even thread, and these threads are entangled with each other.

The recursive decomposition for snow leopard permutations that we gave in Theorem 3.1 induces a similar decomposition on the odd and even threads.

Theorem 3.8. A permutation α is an even thread of nonnegative length if and only if there is an even thread α_1 and an odd thread β_1 with

$$\alpha = c(\beta_1) \ominus 1 \ominus \alpha_1. \tag{1}$$

Similarly, a permutation β is an odd thread of positive length if and only if there is an even thread α_2 and an odd thread β_2 with

$$\beta = (1 \oplus c(\alpha_2) \oplus 1) \ominus \beta_2. \tag{2}$$

Proof: (\Rightarrow) If α is an even thread of length n, then by definition there is a snow leopard permutation π of length 2n+1 such that $\pi_e = \alpha$. Since π is a snow leopard permutation of odd positive length, by Theorem 3.1 there are snow leopard permutations π_1 and π_2 of odd length such that $\pi = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2$. We see that $\pi^e = c(\pi_1^e) \oplus 1 \oplus \pi_2^e$.

If β is an odd thread of length n + 1, then by definition there is a snow leopard permutation π of length 2n + 1 such that $\pi^o = \beta$. Again, we can write $\pi = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2$. Here we see that $\pi^o = (1 \oplus c(\pi_1^e) \oplus 1) \oplus \pi_2^o$.

(\Leftarrow) Now suppose α is a permutation of length n and there are permutations $\beta_1 \in OT_k$ and $\alpha_1 \in ET_{n-k-1}$ such that $\alpha = c(\beta_1) \ominus 1 \ominus \alpha_1$. There must be snow leopard permutations π_1 and π_2 of lengths 2k - 1 and 2n - 2k - 1, respectively, such that $\pi_1^o = \beta_1$ and $\pi_2^e = \alpha_1$. By Theorem 3.1, $\pi = (1 \oplus c(\pi_1) \oplus 1) \ominus 1 \ominus \pi_2$ is a snow leopard permutation of length 2n + 1. Here, $\pi^e = c(\pi_1^o) \ominus 1 \ominus \pi_2^e = c(\beta_1) \ominus 1 \ominus \alpha_1 = \alpha$.

Finally, suppose β is a permutation of length n + 1 and there are permutations $\alpha_2 \in ET_k$ and $\beta_2 \in OT_{n-k-1}$ such that $\beta = (1 \oplus c(\alpha_2) \oplus 1) \oplus \beta_2$. Since α_2 is an even thread of length k, there is a snow leopard permutation π_1 of length 2k + 1 such that $\pi_1^e = \alpha_2$. Similarly, since β_2 is an odd thread of length n - k - 1, there is a snow leopard permutation π_2 of length 2n - 2k - 3 such that $\pi_2^o = \beta_2$. By Theorem 3.1, $\pi = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2$ is a snow leopard permutation of length 2n + 1. Here, $\pi^o = (1 \oplus c(\pi_1^e) \oplus 1) \oplus \pi_2^o = (1 \oplus c(\alpha_2) \oplus 1) \oplus \beta_2 = \beta$.

Note that the decomposition of an odd thread given in Theorem 3.8 is unique, while the decomposition of an even thread is not. That is, for a given odd thread β there is a unique even thread α_1 and a unique odd thread β_1 for which (2) holds. The first entry of β corresponds to the first 1 in the decomposition, the largest entry of β corresponds to the second 1, and then α_1 and β_1 are determined. But for a given even thread there can be more than one pair (α_2, β_2), consisting of an even thread and an odd thread, for which (1) holds.

Example 3.9. *The odd threads* 4657312 *and* 7243561 *can be decomposed as* $(1 \oplus c(12) \oplus 1) \oplus 312$ *and* $(1 \oplus @ \oplus 1) \oplus 243561$, *respectively*.

Example 3.10. The even thread $\alpha = 653421$ can be decomposed as $c(\emptyset) \ominus 1 \ominus 53421, c(1) \ominus 1 \ominus 3421$ or $c(12435) \ominus 1 \ominus \emptyset$.

In each of the three decompositions of α in Example 3.10, we see that the middle 1 corresponds to a left-to-right maximum and fixed point in $c(\alpha)$. It's not difficult to see that this will hold for all even threads. In particular, suppose α is an even thread, and α_1 and β_1 are an even thread and an odd thread, respectively, for which $\alpha = c(\beta_1) \oplus 1 \oplus \alpha_1$. Then $c(\alpha) = \beta_1 \oplus 1 \oplus c(\alpha_1)$, and there are exactly $k = |\beta_1|$ entries of $c(\alpha)$ preceding $c(\alpha)(k+1)$, all of which are less than $c(\alpha)(k+1)$. It follows that $c(\alpha)(k+1)$ is a left-to-right maximum and fixed point in $c(\alpha)$. Since the connector of a snow leopard permutation π is both a left-to-right maximum and fixed point in $c(\pi)$, and the 1 in the decomposition of an even thread corresponds to the connector of a snow leopard permutation, we will refer to an entry of an even thread α as an *eligible connector* whenever it corresponds to an entry of $c(\alpha)$ which is both a left-to-right maximum and a fixed point.

It is worth noting that the converse of the above statement does not hold: some even threads have eligible connectors which do not arise from a decomposition of the thread. For example, $\alpha = 354621$ is an even thread with eligible connectors 2 and 1. If 1 is to correspond to the 1 in a decomposition $c(\beta) \ominus 1 \ominus \alpha_1$, then $c(\beta) = 24351$ and $\beta = 42315$. However, 42315 is not an odd thread. We will prove that although not every eligible connector in an even thread α corresponds to a 1 in a decomposition of α , the leftmost eligible connector does always correspond to such a 1. We begin with two simple ways of constructing new threads from old threads, which arise from the fact that if π and σ are snow leopard permutations of odd length then $\pi \ominus 1 \ominus \sigma$ is, too.

Proposition 3.11. Suppose β_1 and β_2 are odd threads. Then $\beta_1 \ominus \beta_2$ is also an odd thread, and the permutations which are entangled with $\beta_1 \ominus \beta_2$ are exactly the permutations of the form $\alpha_1 \ominus 1 \ominus \alpha_2$, where α_1 and α_2 are entangled with β_1 and β_2 , respectively.

Proof: If β_1 and β_2 are odd threads, then there exist snow leopard permutations π_1 and π_2 such that $\pi_1^o = \beta_1$ and $\pi_2^o = \beta_2$. From Theorem 3.4, we know that $\pi = \pi_1 \ominus 1 \ominus \pi_2$ is a snow leopard permutation with $\pi^o = \pi_1^o \ominus \pi_2^o$. Thus, $\pi_1^o \ominus \pi_2^o = \beta_1 \ominus \beta_2$ is an odd thread.

To prove the rest of the Proposition, first suppose α_1 and α_2 are even threads which are entangled with the odd threads β_1 and β_2 , respectively. Then there are snow leopard permutations π_1 and π_2 such that $\pi_1^e = \alpha_1, \pi_1^o = \beta_1, \pi_2^e = \alpha_2$, and $\pi_2^o = \beta_2$. By Theorem 3.4, $\pi = \pi_1 \ominus 1 \ominus \pi_2$ is a snow leopard permutation, and we find $\pi^e = \alpha_1 \ominus 1 \ominus \alpha_2$ and $\pi^o = \beta_1 \ominus \beta_2$. Therefore, every permutation of the form $\alpha_1 \ominus 1 \ominus \alpha_2$, where α_1 and α_2 are entangled with β_1 and β_2 , respectively, is entangled with $\beta_1 \ominus \beta_2$.

To prove the reverse inclusion, suppose α is an even thread entangled with $\beta_1 \ominus \beta_2$, and π is a snow leopard permutation with $\pi^e = \alpha$ and $\pi^o = \beta_1 \ominus \beta_2$. By Corollary 3.3, there are snow leopard permutations π_1, \ldots, π_k of odd length such that

$$\pi = (1 \oplus c(\pi_1) \oplus 1) \ominus 1 \ominus (1 \oplus c(\pi_2) \oplus 1) \ominus 1 \ominus \cdots \ominus 1 \ominus (1 \oplus c(\pi_k) \oplus 1).$$

The 1s in each summand $1 \oplus c(\pi_j) \oplus 1$ are all in odd positions, so there must be some m for which $\beta_1 = \pi_1^o$ and $\beta_2 = \pi_2^o$, where

$$\pi_1 = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \cdots \oplus 1 \oplus (1 \oplus c(\pi_m) \oplus 1)$$

and

$$\pi_2 = (1 \oplus c(\pi_{m+1}) \oplus 1) \oplus 1 \oplus \cdots \oplus 1 \oplus (1 \oplus c(\pi_k) \oplus 1).$$

By Corollary 3.3, π_1 and π_2 are snow leopard permutations of odd length, so if $\alpha_1 = \pi_1^e$ and $\alpha_2 = \pi_2^e$, then we have $\alpha = \alpha_1 \ominus 1 \ominus \alpha_2$, and α_1 and α_2 are entangled with β_1 and β_2 , respectively.

It is worth noting that even though (a) is not an odd thread, it behaves like one in some circumstances. For instance, if (a) were an odd thread, then for any odd thread $1 \ominus \beta$ we could use Proposition 3.11 to conclude that $(a) \ominus 1 \ominus \beta = \beta$ is also an odd thread. And if (a) were an odd thread then we could use a similar argument to show that if $\beta \ominus 1$ is an odd thread then β is also an odd thread. Even though these arguments fail, the results still hold, as we show next.

Proposition 3.12. For any permutation β of nonnegative length, if $1 \ominus \beta$ or $\beta \ominus 1$ is an odd thread then β is an odd thread.

Proof: Suppose $1 \ominus \beta$ is an odd thread. Then by (2) there is an even thread α_2 and an odd thread β_2 such that $1 \ominus \beta = (1 \oplus c(\alpha_2) \oplus 1) \ominus \beta_2$, and it's not hard to see that this decomposition is unique. Comparing the largest elements on each side, we find $\alpha_2 = @$ and $\beta_2 = \beta$, so β is an odd thread.

Now suppose $\beta \ominus 1$ is an odd thread; we argue by induction on $|\beta \ominus 1|$. If $|\beta \ominus 1| = 1$ then $\beta \ominus 1 = 1$ and $\beta = \emptyset$, and the result holds. If $|\beta \ominus 1| = 2$ then $\beta \ominus 1 = 21$ and $\beta = 1$, and the result holds. If $|\beta \ominus 1| \ge 3$ then by (2) there is an even thread α_2 and an odd thread β_2 such that $\beta \ominus 1 = (1 \oplus c(\alpha_2) \oplus 1) \ominus \beta_2$. Since $|1 \oplus c(\alpha_2) \oplus 1| \ge 1$, we must have $|\beta_2| < |\beta \ominus 1|$. By construction there is a permutation β_1 of nonnegative length such that $\beta_2 = \beta_1 \ominus 1$, and by induction β_1 is an odd thread. Therefore, $(1 \oplus c(\alpha_2) \oplus 1) \ominus \beta_1 = \beta$ is also an odd thread, by Theorem 3.8.

We now turn our attention to constructing new even threads from old even threads.

Proposition 3.13. Suppose α_1 and α_2 are even threads. Then $\alpha_1 \ominus 1 \ominus \alpha_2$ is also an even thread, and if β_1 and β_2 are odd threads entangled with α_1 and α_2 , respectively, then $\beta_1 \ominus \beta_2$ is an odd thread entangled with $\alpha_1 \ominus 1 \ominus \alpha_2$.

Proof: If α_1 and α_2 are even threads, then there exist snow leopard permutations π_1 and π_2 such that $\pi_1^e = \alpha_1$ and $\pi_2^e = \alpha_2$. From Theorem 3.4, we know that $\pi = \pi_1 \ominus 1 \ominus \pi_2$ is a snow leopard permutation, with $\pi^e = \pi_1^e \ominus 1 \ominus \pi_2^e$. Thus, $\pi_1^e \ominus 1 \ominus \pi_2^e = \alpha_1 \ominus 1 \ominus \alpha_2$ is an even thread.

To prove the rest of the Proposition, suppose β_1 and β_2 are odd threads which are entangled with the even threads α_1 and α_2 , respectively. Then there are snow leopard permutations π_1 and π_2 such that $\pi_1^e = \alpha_1, \pi_1^o = \beta_1, \pi_2^e = \alpha_2$, and $\pi_2^o = \beta_2$. By Theorem 3.4, $\pi = \pi_1 \ominus 1 \ominus \pi_2$ is a snow leopard permutation, and we find $\pi^e = \alpha_1 \ominus 1 \ominus \alpha_2$ and $\pi^o = \beta_1 \ominus \beta_2$. In particular, $\beta_1 \ominus \beta_2$ is an odd thread which is entangled with $\alpha_1 \ominus 1 \ominus \alpha_2$.

Note that if α_1 and α_2 are even threads, then there may be an odd thread β entangled with $\alpha_1 \ominus 1 \ominus \alpha_2$ which does not have the form $\beta_1 \ominus \beta_2$ for odd threads β_1 and β_2 which are entangled with α_1 and α_2 , respectively. For example, if $\alpha_1 = \emptyset$ and $\alpha_2 = 1$ then $\alpha_1 \ominus 1 \ominus \alpha_2 = 21$, which is entangled with 123. But α_1 is only entangled with 1, α_2 is only entangled with 12 and 21, and neither $1 \ominus 12 = 312$ nor $1 \ominus 21 = 321$ is equal to 123.

The next step in proving that the leftmost eligible connector in an even thread always corresponds to a connector in some snow leopard permutation is to show that if any other eligible connector corresponds to a connector in a snow leopard permutation, then the even thread must begin with its largest element.

Lemma 3.14. Suppose α is an even thread of length n, the permutation π is a snow leopard permutation with $\pi^e = \alpha$, and the connector of π does not correspond with the leftmost eligible connector of α . Then $\alpha(1) = n$.

Proof: Let $\alpha(j)$ be the leftmost eligible connector in α . Since this eligible connector does not correspond to the connector in π , there must be an eligible connector in α to the right of $\alpha(j)$ which does correspond to the connector in π . Let this eligible connector be $\alpha(k)$, and note that k > j. Since α is an even thread, (1) there is an even thread α_1 and an odd thread β_1 such that $|\beta_1| = k - 1$ and

$$\alpha = c(\beta_1) \ominus 1 \ominus \alpha_1$$

In particular, $\alpha(j)$ is among $\beta_1(1), \ldots, \beta_1(k-1)$. Now by (2) there is an even thread α_2 and an odd thread β_2 such that

$$\beta_1 = (1 \oplus c(\alpha_2) \oplus 1) \ominus \beta_2. \tag{3}$$

We claim that $\beta_2 = \emptyset$.

To prove our claim, first note that $c(\beta_1)(j)$ is an eligible connector in α , so it is a left-to-right maximum in $c(\alpha)$, a left-to-right minimum in α , and a left-to-right minimum in $c(\beta_1)$. But $c(\alpha)(j)$ is also a fixed point in $c(\alpha)$, so the entries to its left in $c(\alpha)$ are exactly $1, 2, \ldots, j - 1$. This means the entries to the left of $\alpha(j)$ in α are $n, n - 1, \ldots, n - j + 2$, which are all of the entries of α which are larger than $\alpha(j)$. Therefore, the entries to the left of $c(\beta_1)(j)$ are $k - 1, k - 2, \ldots, k - j + 1$, the largest entries of $c(\beta_1)$, and $c(\beta_1)(j) = k - j$. But $c(\beta_1) = (1 \ominus \alpha_2 \ominus 1) \oplus \beta_2$, as in Figure 6. Now if $c(\beta_2) \neq \emptyset$, and $c(\beta_1)(j)$




Fig. 6: A decomposition of $c(\beta_1)$.

falls in $1 \ominus \alpha_2 \ominus 1$, then one of $k - 1, \ldots, k - j + 1$ is to the right of $c(\beta_1)(j)$, which is a contradiction. On the other hand, if $c(\beta_1)(j)$ falls in $c(\beta_2)$, then there is an entry smaller than $c(\beta_1)(j)$ to the left of $c(\beta_1)(j)$ (since $1 \ominus \alpha_2 \ominus 1$ must be nonempty), which is also a contradiction. Therefore, $\beta_2 = \emptyset$. Since $\beta_2 = \emptyset$, by (3) we have $\beta_1 = 1 \oplus c(\alpha_2) \oplus 1$, so $\beta_1(1) = 1$ and $\alpha(1) = n$, as desired.

Now that we have assembled all of the necessary tools, we are ready to show that the leftmost eligible connector in an even thread always corresponds to a connector in some snow leopard permutation.

Theorem 3.15. If α is an even thread of length n, then there exists a snow leopard permutation π with $\pi^e = \alpha$ whose connector corresponds to the leftmost eligible connector in α . In particular, if this entry is $c(\alpha)(n-j+1)$, then $\pi(1) = 2j + 1$. By convention, we say j = 0 if no such entry exists.

Proof: Suppose α is an even thread of length n, and π is a snow leopard permutation with $\pi^e = \alpha$. If the connector of π corresponds with the leftmost eligible connector of α , then the result holds. If not, then by Lemma 3.14 we have $\alpha(1) = n$. By (1) there is an even thread α and an odd thread β such that

$$\alpha = c(\beta_1) \ominus 1 \ominus \alpha_1. \tag{4}$$

We claim that $\alpha = 1 \ominus \alpha_4$ for some even thread α_4 .

Our claim follows from (4) if $\beta_1 = \emptyset$, so suppose $\beta_1 \neq \emptyset$. By (2) there is an odd thread β_2 and an even thread α_2 with

$$\beta_1 = (1 \oplus c(\alpha_2) \oplus 1) \ominus \beta_2,$$

which implies

$$\alpha = ((1 \ominus \alpha_2 \ominus 1) \oplus c(\beta_2)) \ominus 1 \ominus \alpha_1$$

Since $\alpha(1) = n$, we must have $\beta_2 = \emptyset$. Furthermore, $1 \ominus \alpha_1 = \emptyset \ominus 1 \ominus \alpha_1$ is an even thread by Proposition 3.13; writing α_3 to denote this thread, we have

$$\alpha = 1 \ominus \alpha_2 \ominus 1 \ominus \alpha_3.$$

Using Proposition 3.13 again, we see that $\alpha_2 \ominus 1 \ominus \alpha_3$ is also an even thread, which we denote by α_4 , and our claim follows.

To complete the proof, note that since α_4 is an even thread, there is a snow leopard permutation π with $\pi^e = \alpha_4$. By Theorem 3.4, $1 \ominus 1 \ominus \pi$ is a snow leopard permutation with even thread α , whose connector corresponds to $\alpha(1)$, the leftmost eligible connector of α .

Theorem 3.15 tells us that while a given even thread α may have several decompositions as in (1), it always has such a decomposition in which the 1 in (1) corresponds to the leftmost eligible connector in α . This decomposition is clearly unique, so we will call it the *leftmost decomposition* of α .

4 Even Threads, Odd Threads, and Restricted Catalan Paths

Now that we have recursive decompositions of the even and odd threads, and we have investigated the extent to which these decompositions are unique for a given thread, it's natural to consider the numbers of these threads of each length. In Table 2 we have the number of even and odd threads of length six or less.

n	-1	0	1	2	3	4	5	6
$ ET_n $	1	1	1	2	6	17	46	128
$ OT_n $	0	1	1	2	4	9	23	63

Tab. 2: The number of even threads and odd threads of length six or less.

It turns out that these numbers also count some more familiar combinatorial objects. To describe these objects, recall that a *Catalan path* of length n is a lattice path from (0,0) to (n,n) consisting of unit North

(0,1) and East (1,0) steps which does not pass below the line y = x. For convenience, we sometimes write a Catalan path as a sequence of Ns and Es, with N denoting a North step and E denoting an East step. For instance, the five Catalan paths of length three are NNNEEE, NNENEE, NNEENE, NEENE, NENNEE, and NENENE. If p is a Catalan path of length n, then we will write p^r to denote the *reverse* of p, which is the path obtained by reflecting p over the line x + y = n. In terms of Ns and Es, reversing a Catalan path is equivalent to reversing the corresponding string and then exchanging Ns and Es. For example, if p = NENNEE then $p^r = NNEENE$.

Sapounakis et al. (2007) show that the number of Catalan paths of length $n \ge 1$ with k occurrences of NEEN (in consecutive entries) is given by

$$a_{n,k} = \frac{1}{n} \binom{n}{k} \sum_{j=k}^{\lfloor \frac{n-1}{2} \rfloor} (-1)^{j-k} \binom{n-k}{j-k} \binom{2n-3j}{n-j+1}.$$

Notice that when k = 0 the first few terms of this sequence (beginning with n = 1) are 1, 2, 4, 9, and 23, which suggests that $|OT_n| = a_{n,0}$. On the other hand, the values of $|ET_n|$ appear to match OEIS sequence A102403, whose *n*th term is the number of Catalan paths of length *n* with no ascent of length exactly two. With this in mind, we introduce some notation for the sets of these paths of a given length.

Definition 4.1. For each $n \ge 0$, we write $ENNE_n$ to denote the set of Catalan paths of length n which have no ascent of length exactly two. Similarly, we write $NEEN_n$ to denote the set of Catalan paths of length n which do not contain the four consecutive steps NEEN.

Our data suggest that $|ET_n| = |ENNE_{n+1}|$ and $|OT_n| = |NEEN_n|$ for each nonnegative integer n. To prove these results, we introduce decompositions of the paths in $ENNE_n$ and $NEEN_n$ which mirror our decompositions of the even and odd threads, respectively.

Theorem 4.2. Suppose n is a positive integer. For each $p \in ENNE_n$, there are unique nonnegative integers k and l, and unique Catalan paths $a \in ENNE_k$ and $b \in NEEN_l$, such that n = k + l + 1, $p = aNb^r E$, and b does not end with NE. Conversely, for every $a \in ENNE_k$ and $b \in NEEN_l$ such that b does not end with NE, the path $aNb^r E$ is in $ENNE_n$, where n = k + l + 1.

Proof: (\Rightarrow) Suppose $p \in ENNE_n$. Since p is a Catalan path which begins at (0, 0) and ends at (n, n), it must return to the diagonal y = x at least once. Suppose the last time p touches the diagonal before (n, n) is at (k, k). Let a denote the subpath of p from (0, 0) to (k, k) and let b denote the subpath of p from (k, k + 1) to (n - 1, n). Then a and b are Catalan paths, and |a| = k and |b| = n - k - 1, so |a| + |b| = n - 1. Furthermore, if c and d are any Catalan paths with |c| = k and |d| = n - k - 1 then the Catalan path cNd^rE has its last return to the diagonal before (n, n) at (k, k). Therefore, a, b, k, and l = n - k - 1 are uniquely determined by p, and it remains only to show that $a \in ENNE_k$ and $b^r \in NEEN_{n-k-1}$.

Since a is a subpath of p and p contains no ascent of length exactly two, neither does a. Thus $a \in ENNE_k$.

To show $b^r \in NEEN_{n-k-1}$, first notice that since $NbE \in ENNE_{n-k}$, if b contains an ascent of length exactly two then this ascent must start the path. Equivalently, if b^r contains a plateau of length exactly two, then this plateau must end the path. Thus, b^r does not contain NEEN, so $b^r \in NEEN_{n-k-1}$. In addition, b^r cannot start with NE, for otherwise the Nb portion of p would contain an ascent of length exactly two. Thus, b^r does not end with an NE.

(\Leftarrow) Suppose $a \in ENNE_k$ and $b \in NEEN_l$, and b does not end with NE. We want to show that the path aNb^rE does not contain an ascent of length exactly two. Since $a \in ENNE_k$, if aNb^rE did contain such an ascent, it would have to occur in the Nb^rE portion of the path. Additionally, b contains no NEEN, so b^r contains no ENNE. Thus, the problematic ascent would have to begin with the N of Nb^rE . However, we assumed that b does not end with NE, so b^r does not begin with NE. Thus, there is no problematic ascent in aNb^rE , and this path of length n = k + l + 1 will be in $ENNE_n$.

Theorem 4.3. Suppose n is a positive integer. For each $p \in NEEN_n$ there are unique nonnegative integers k and l, and unique Catalan paths $a \in ENNE_k$ and $b \in NEEN_l$, such that n = k + l + 1 and $p = a^r NbE$. Conversely, for every $a \in ENNE_k$ and $b \in NEEN_l$, the path $a^r NbE$ is in $NEEN_n$, where n = k + l + 1.

Proof: (\Rightarrow) Suppose $p \in NEEN_n$. As in the proof of Theorem 4.2, let *a* denote the subpath of *p* from (0,0) to (k,k) and let *b* denote the subpath of *p* from (k,k+1) to (n-1,1), where (k,k) is the point at which *p* last touches the diagonal before (n,n). As before, for any Catalan paths *c* and *d* with |c| = k and |d| = n - k - 1 the Catalan path $c^r NdE$ has its last return to the diagonal at (k,k), so *a*, *b*, *k*, and l = n - k - 1 are uniquely determined by *p*.

Since NbE is a path in $NEEN_{n-k}$, we have $b \in NEEN_{n-k-1}$. Similarly, since a does not contain NEEN, the path a^r does not contain ENNE. Therefore, if a^r contains an ascent of length exactly two, then this ascent must come at the beginning of the path, in which case a^r begins with NNE. In this case, a would end with NEE, so p would contain NEEN. This contradiction shows $a^r \in ENNE_k$.

(\Leftarrow) Suppose $a \in ENNE_k$ and $b \in NEEN_l$, and let $p = a^r NbE$. We want to show that p does not contain NEEN. Since there is no NEEN in b, if p contains NEEN, then it must fall in the $a^r N$ portion of p. However, $a \in ENNE_k$ so a has no ascent of length exactly two, meaning a^r has no plateau of length exactly two. Thus, $a^r N$ also cannot contain NEEN, so $p \in NEEN_n$, where n = |a| + |b| + 1 = k + l + 1.

Now that we can decompose threads and Catalan paths in similar ways, we can recursively define functions from $ENNE_{n+1}$ to ET_n , and from $NEEN_n$ to OT_n ; these functions will turn out to be bijections.

Theorem 4.4. For each nonnegative integer n, there exist unique functions

 $H: ENNE_{n+1} \to ET_n$ $J: NEEN_n \to OT_n$

such that $H(\emptyset) = @$, $J(\emptyset) = \emptyset$, if $p = aNb^r E$ for Catalan paths a and b then

$$H(p) = c(J(b)) \ominus 1 \ominus H(a), \tag{5}$$

and if $p = a^r N b E$ for Catalan paths a and b then

$$J(p) = (1 \oplus c(H(a)) \oplus 1) \oplus J(b).$$
(6)

Proof: The result is clear for n = 0, so suppose $n \ge 1$.

Since every Catalan path p of positive length has unique decompositions into the forms aNb^rE and a^rNbE , some simple computations with lengths shows that there exist unique functions $H : ENNE_{n+1} \rightarrow S_n$ and $J : NEEN_n \rightarrow S_n$ satisfying (5) and (6). So it's sufficient to show that if $p \in ENNE_{n+1}$ then $H(p) \in ET_n$ and if $p \in NEEN_n$ then $J(p) \in OT_n$.

The result is true by construction for n = 0, and it's routine to check that $H(NE) = \emptyset \in ET_0$ and $J(NE) = 1 \in OT_1$, so suppose $n \ge 2$ and the result holds for all paths of length n - 1 or less. If $p \in ENNE_n$ then by Theorem 4.2 there are unique paths $a \in ENNE_k$ and $b \in NEEN_l$, such that n = k + l + 1, $p = aNb^rE$, and b does not end with NE. By induction $H(a) \in ET_{k-1}$ and $J(b) \in OT_l$, so by (5) and (1) the permutation H(p) is in ET_{n-1} . The proof that if $p \in NEEN_n$ then $J(p) \in OT_n$ is similar, using Theorem 4.3, along with (6) and (2).

In Table 3 we have the values of H and J on paths of length three or less. This data suggests H and J

path a	H(a)		
Ø	0	path a	J(a)
NE	Ø	Ø	\emptyset
NENE	1		1
NENENE	21		12
NNNEEE	12	NNEE	21
NNNNEEEE	123	NNNEEE	321
NNNENEEE	132	NNENEE	312
NNNEENEE	213	NENNEE	231
NNNEEENE	312	NENENE	123
NENNNEEE	231		I
NENENENE	321		

Tab. 3: Values of H(a) and J(a) for small |a|.

are bijections; to prove this, we construct their inverses.

Theorem 4.5. For each nonnegative integer n, there exist unique functions

$$F: ET_n \to ENNE_{n+1}$$
$$G: OT_n \to NEEN_n$$

such that $F(@) = \emptyset$, $G(\emptyset) = \emptyset$, if $\pi = c(\beta) \ominus 1 \ominus \alpha$ for an even thread α and an odd thread β is the leftmost decomposition of the even thread π , then

$$F(\pi) = F(\alpha)NG(\beta)^{r}E,$$
(7)

and if $\pi = (1 \oplus c(\alpha) \oplus 1) \ominus \beta$ for an even thread α and an odd thread β then

$$G(\pi) = F(\alpha)^r N G(\beta) E.$$
(8)

Proof: This is similar to the proof of Theorem 4.4.

Theorem 4.6. F and G are the inverse functions of H and J, respectively.

Proof: It's routine to check that $H(F(@)) = @, H(F(\emptyset)) = \emptyset, F(H(\emptyset)) = \emptyset, J(G(\emptyset)) = \emptyset$, and $G(J(\emptyset)) = \emptyset$, so the map $H \circ F$ is the identity on ET_{-1} and ET_0 , the map $F \circ H$ is the identity on $ENNE_0$, the map $J \circ G$ is the identity on OT_0 , and the map $G \circ J$ is the identity on $NEEN_0$. Now fix $n \ge 1$, and suppose by induction that for all k < n we have that $H \circ F$ is the identity on ET_k , $F \circ H$ is the identity on OT_k , and $G \circ J$ is the identity on $NEEN_k$. It turns out that the four facts we need to prove are not completely independent, so we start with $J \circ G$.

To show $J \circ G$ is the identity on OT_n , suppose we have $\beta \in OT_n$, and let $\beta = (1 \oplus c(\alpha_1) \oplus 1) \ominus \beta_1$ be the decomposition of β from (2). By (8) we have

$$G(\beta) = F(\alpha_1)^r N G(\beta_1) E.$$

Now when we decompose $G(\beta)$ in the form $a^r N b E$, we find $a = F(\alpha_1)$ and $b = G(\beta_1)$. Therefore, by (6) we have

$$J(G(\beta)) = (1 \oplus c(H(F(\alpha_1))) \oplus 1) \oplus J(G(\beta_1)).$$

Since $|\beta_1| < n$ and $|\alpha_1| < n$, by induction we have

$$J(G(\beta)) = (1 \oplus c(\alpha_1) \oplus 1) \ominus \beta_1$$

= β ,

as desired.

To show $H \circ F$ is the identity on ET_n , suppose we have $\alpha \in ET_n$, and let $\alpha = c(\beta_1) \ominus 1 \ominus \alpha_1$ be the leftmost decomposition of α . By (7) we have

$$F(\alpha) = F(\alpha_1) N G(\beta_1)^r E_{\cdot}$$

Now when we decompose $F(\alpha)$ in the form $aNb^r E$, we find $a = F(\alpha_1)$ and $b = G(\beta_1)$. Therefore, by (5) we have

$$H(F(\alpha)) = c(J(G(\beta_1))) \ominus 1 \ominus H(F(\alpha_1)).$$

Now if $\alpha_1 = @$ then $|\beta_1| = n$, and by induction and our previous case we have

$$H(F(\alpha)) = c(\beta_1) \ominus 1 \ominus \alpha_1$$

= α ,

as desired. On the other hand, if $\alpha_1 \neq 0$ then $|\alpha_1| < n$ and $|\beta_1| < n$, and the result follows by induction.

To show $F \circ H$ is the identity on $ENNE_n$, suppose we have $p \in ENNE_n$, and let $p = aNb^r E$ for Catalan paths a and b. By (5) we have

$$H(p) = c(J(b)) \ominus 1 \ominus H(a).$$
(9)

We claim this is the leftmost decomposition of H(p). To see this, first note that since J(b) is an odd thread and H(a) is an even thread, there are snow leopard permutations π_1 and π_2 such that $J(b) = \pi_1^o$ and $H(a) = \pi_2^o$. Observe that

$$((1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2)^e = c(J(b)) \oplus 1 \oplus H(a),$$

and the connector in $(1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2$ corresponds with the 1 in $c(J(b)) \oplus 1 \oplus H(a)$. Therefore, by Lemma 3.14, if $c(J(b)) \oplus 1 \oplus H(a)$ is not the leftmost decomposition of H(p) then it begins with its largest entry. This implies that J(b) begins with 1. Now by (6), if $b = a_1^r N b_1 E$ then $J(b) = \emptyset$, so $b_1 = \emptyset$, and b ends with NE. This means b^r begins with NE, and $p = aNb^r E$ has an ascent of length exactly two, contradicting the fact that $p \in ENNE_n$.

Since the right side of (9) is the leftmost decomposition of H(p), by (7) we have

$$F(H(p)) = F(H(a))NG(J(b))^{r}E$$
$$= aNb^{r}E$$
$$= p,$$

by induction.

To show $G \circ J$ is the identity on $NEEN_n$, suppose we have $p \in NEEN_n$, and let $p = a^r NbE$ for Catalan paths a and b. By (6) we have

$$J(p) = (1 \oplus c(H(a)) \oplus 1) \ominus J(b)$$

Since H(a) is an even thread, J(b) is an odd thread, and the decomposition of an odd thread in (2) is unique, by (8) we have

$$G(J(p)) = F(H(a))^r NG(J(b))E.$$

Now the result follows by induction.

5 Entangled Threads

For each $n \ge -1$, there is a bipartite graph whose vertices are the even threads of length n and the odd threads of length n + 1, in which two vertices are adjacent whenever they are entangled. In Section 4 we gave a partial answer to the question of how many vertices this graph has, by giving bijections between the even threads and the Catalan paths with no ascent of length exactly two, and between the odd threads and the Catalan paths with no ascent of length exactly two, and between the odd threads and the Catalan paths with no *NEEN*. Nevertheless, there are numerous other questions one might ask about this graph. In this section we start to answer one of these questions, by characterizing the even threads of length n - 1 which are entangled with the increasing permutation $12 \cdots n$, and by characterizing the odd threads of length n + 1 which are entangled with the decreasing permutation $n \cdots 21$. Throughout we write \Box_n (resp. \Box_n) to denote the increasing (resp. decreasing) permutation $12 \cdots n$ (resp. $n \cdots 21$).

To state and prove our results, we first need some terminology. Recall that an *involution* π is a permutation π with $\pi = \pi^{-1}$. As Guibert (1995) and Egge (2004) have noted, involutions which avoid the classical pattern 3412 have a simple recursive structure: a permutation π is a 3412-avoiding involution if and only if one of the following mutually exclusive conditions holds: (i) $\pi = \emptyset$, (ii) $\pi = 1 \oplus \pi_1$ for a 3412-avoiding involution π_1 , or (iii) $\pi = (1 \oplus \pi_1 \oplus 1) \oplus \pi_2$ for 3412-avoiding involutions π_1 and π_2 . Moreover, when π satisfies (ii) or (iii), the 3412-avoiding involutions π_1 and π_2 are uniquely determined. Using these results, it is routine to show that the number of 3412-avoiding involutions of length n is the Motzkin number M_n , which may be defined by $M_0 = 1$ and $M_n = M_{n-1} + \sum_{k=2}^n M_{k-2}M_{n-k}$ for $n \ge 1$. We will encounter another family of objects counted by the Motzkin numbers in Section 6, and the interested reader can find still more such families of objects in the paper of Donaghey and Shapiro (1977) and Exercise 6.38 of the book of Stanley (1999). As we show next, the even threads entangled with \Box_n and the odd threads entangled with \Box_n are easy to describe in terms of 3412-avoiding involutions.

- **Theorem 5.1.** (i) For any integer $n \ge 1$, the even threads entangled with the odd thread \square_n are the 3412-avoiding involutions of length n 1.
 - (ii) For any integer $n \ge 0$, the odd threads entangled with the even thread \square_n are the complements of the 3412-avoiding involutions of length n + 1.

Proof: The odd thread $\square_1 = 1$ is only entangled with the even thread \emptyset , which is the only 3412-avoiding involution of length 0. Similarly, the even thread $\square_1 = 1$ is entangled with both 12 and 21, which are the complements of the 3412-avoiding involutions of length 2. Therefore, the results holds for n = 1. Now fix $n \ge 2$, and suppose the result holds for all k < n; we argue by induction on n.

Suppose α is an even thread entangled with \Box_n . Then there is a snow leopard permutation π such that $\pi^e = \alpha$ and $\pi^o = \Box_n$. By Theorem 3.1, there are snow leopard permutations π_1 and π_2 such that $\pi = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2$. Since $\pi^o = \Box_n$, we see that $\pi(1) = 1$, which means $\pi_2 = @$ and $\pi = 1 \oplus c(\pi_1) \oplus 1$. Now $\pi_1^e = \Box_{n-2}$, so by induction $c(\pi_1^o)$ is a 3412-avoiding involution. On the other hand, $\alpha = \pi^e = c(\pi_1^o)$, so α is also a 3412-avoiding involution.

For the reverse inclusion, suppose α is a 3412-avoiding involution of length n - 1. By induction $c(\alpha)$ is an odd thread entangled with \Box_{n-2} , so there is a snow leopard permutation π_1 with $\pi_1^o = c(\alpha)$ and $\pi_1^e = \Box_{n-2}$. By Theorem 3.1, the permutation $1 \oplus c(\pi_1) \oplus 1$ is a snow leopard permutation with $\pi^o = \Box_n$ and $\pi^e = \alpha$, so α is an even thread entangled with \Box_n .

Suppose β is an odd thread entangled with \square_n . Then there is a snow leopard permutation π such that $\pi^o = \beta$ and $\pi^e = \square_n$. By Theorem 3.1, there are snow leopard permutations π_1 and π_2 such that $\pi = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2$. In addition,

$$\beta = \pi^o = (1 \oplus c(\pi_1^e) \oplus 1) \ominus \pi_2^o$$

and

$$\square_n = \pi^e = c(\pi_1^o) \ominus 1 \ominus \pi_2^e,$$

which implies $\pi_1^o = \Box_k$ and $\pi_2^e = \Box_{n-k-1}$ for some k with $-1 \le k \le n-1$. By induction and a previous case, π_1^e and $c(\pi_2^o)$ are 3412-avoiding involutions, so $c(\beta) = (1 \ominus \pi_1^e \ominus 1) \oplus c(\pi_2^o)$ is also a 3412-avoiding involution.

For the reverse inclusion, suppose $c(\beta)$ is a 3412-avoiding involution of length n + 1. Then there are 3412-avoiding involutions α_1 of length k and β_1 of length n - k - 1, where $-1 \le k \le n - 1$, such that

$$c(\beta) = (1 \ominus \alpha_1 \ominus 1) \oplus \beta_1$$

By induction and a previous case, α_1 is an even thread entangled with \Box_{k+1} and $c(\beta_1)$ is an odd thread entangled with \Box_{n-k-2} . Therefore, there are snow leopard permutations π_1 and π_2 with $\pi_1^e = \alpha_1$, $\pi_1^o = \Box_{k+1}, \pi_2^o = c(\beta_1)$, and $\pi_2^e = \Box_{n-k-2}$. Now $\pi = (1 \oplus c(\pi_1) \oplus 1) \oplus 1 \oplus \pi_2$ is a snow leopard permutation by Theorem 3.1, and it has

$$\pi^{o} = (1 \oplus c(\pi_{1}^{e}) \oplus 1) \oplus \pi_{2}^{o}$$
$$= (1 \oplus c(\alpha_{1}) \oplus 1) \oplus c(\beta_{1})$$

and

$$\pi^{e} = c(\pi_{1}^{o}) \ominus 1 \ominus \pi_{2}^{e}$$
$$= \boxtimes_{k+1} \ominus 1 \ominus \boxtimes_{n-k-2}$$
$$= \boxtimes_{n}.$$

Snow Leopard Permutations and their Even and Odd Threads

Since $c(\pi^o) = (1 \ominus \alpha_1 \ominus 1) \oplus \beta_1 = c(\beta)$, we see that β is an odd thread entangled with \square_n .

By combining Theorem 5.1 with Proposition 3.11, we can characterize the even threads entangled with a much larger class of odd threads.

Definition 5.2. For $m \ge 1$ and any positive integers l_1, \ldots, l_m , we write \Box_{l_1,\ldots,l_m} to denote the permutation $\Box_{l_1} \ominus \Box_{l_2} \ominus \cdots \ominus \Box_{l_m}$. We sometimes call a permutation of the form \Box_{l_1,\ldots,l_m} up-layered.

Corollary 5.3. For $m \ge 1$ and any positive integers l_1, \ldots, l_m , the permutation $\boxtimes_{l_1,\ldots,l_m}$ is an odd thread, and the even threads entangled with it are the permutations of the form $\pi_1 \ominus 1 \ominus \pi_2 \ominus 1 \ominus \cdots \ominus 1 \ominus \pi_m$, where π_j is a 3412-avoiding involution of length $l_j - 1$ for $1 \le j \le m$.

Proof: This follows from Theorem 5.1(i) and Proposition 3.11 by induction on m.

Corollary 5.4. For any $m \ge 1$ and any positive integers l_1, \ldots, l_m , there are exactly $M_{l_1-1} \cdots M_{l_m-1}$ even threads entangled with the odd thread \Box_{l_1,\ldots,l_m} .

Proof: This is immediate from Corollary 5.3, since the number of 3412-avoiding involutions of length $l_j - 1$ is M_{l_j-1} for $1 \le j \le m$.

It is natural at this point to try to use Theorem 5.1(ii) and Proposition 3.13 to prove a analogues of Corollaries 5.3 and 5.4 for even threads which are layered in the opposite direction. Unfortunately, doing this only allows us to recover Theorem 5.1(ii). Nevertheless, we can use Corollary 5.3 to prove the analogues we seek. We begin by making the phrase "layered in the opposite direction" precise.

Definition 5.5. For $m \ge 1$ and any positive integers l_1, \ldots, l_m , we write \Box_{l_1,\ldots,l_m} to denote the permutation $\Box_{l_1} \oplus \Box_{l_2} \oplus \cdots \oplus \Box_{l_m}$. We sometimes call a permutation of the form \Box_{l_1,\ldots,l_m} down-layered.

We now have a natural analogue of Corollary 5.3 for down-layered even threads.

Theorem 5.6. For $m \ge 1$ and any positive integers l_1, \ldots, l_m , the permutation \Box_{l_1,\ldots,l_m} is an even thread, and the odd threads entangled with it are the permutations of the form $1 \oplus \pi_1 \oplus 1 \oplus \pi_2 \oplus 1 \oplus \cdots \oplus 1 \oplus \pi_m \oplus 1$, where π_j is the complement of a 3412-avoiding involution of length $l_j - 1$ for $1 \le j \le m$.

Proof: First note that the result is Theorem 5.1(ii) when m = 1, so we may assume $m \ge 2$.

Since @ is an even thread and $\square_{l_1,...,l_m}$ is an odd thread, by Theorem 3.8 we see that $\square_{l_1,...,l_m} = c(\square_{l_1,...,l_m} \ominus 1 \ominus @$ is an even thread.

To prove the rest of the result, suppose π is a snow leopard permutation with $\pi^e = \Box_{l_1,...,l_m}$. By Theorem 3.1, there are snow leopard permutations σ_1 and σ_2 such that $\pi = (1 \ominus c(\sigma_1) \ominus 1) \ominus 1 \ominus \sigma_2$. If $\sigma_2 \neq @$ then there is an odd thread β_1 and a nonempty even thread α_1 such that $\pi^e = c(\beta_1) \ominus 1 \ominus \alpha_1$. In $c(\beta_1) \ominus 1 \ominus \alpha_1$, the largest entry appears to the left of the smallest entry, but in $\pi^e = \Box_{l_1,...,l_m}$ the largest entry appears to the right of the smallest entry (since $m \ge 2$). Therefore, $\sigma_2 = @$ and $\pi = 1 \oplus c(\sigma_1) \oplus 1$.

Since $\pi^e = \boxtimes_{l_1,...,l_m}$, we have $\sigma_1^o = c(\boxtimes_{l_1,...,l_m}) = \boxtimes_{l_1,...,l_m}$. Now by Corollary 5.3, the permutation σ_1^e has the form $\pi_1 \ominus 1 \ominus \cdots \ominus 1 \ominus \pi_{l_m}$, where π_j is a 3412-avoiding involution of length $l_j - 1$ for $1 \le j \le m$. This means $\pi^o = 1 \oplus c(\pi_1) \oplus 1 \oplus \cdots \oplus 1 \oplus c(\pi_{l_m}) \oplus 1$, and the result follows. \Box

Corollary 5.7. For $m \ge 1$ and any positive integers l_1, \ldots, l_m , there are exactly $M_{l_1-1} \cdots M_{l_m-1}$ odd threads entangled with the even thread \Box_{l_1,\ldots,l_m} .

Proof: This is immediate from Theorem 5.6, since the number of 3412-avoiding involutions of length $l_j - 1$ is M_{l_j-1} for $1 \le j \le m$.

Inspired by Corollaries 5.4 and 5.7, along with some numerical data, we make the following conjecture.

Conjecture 5.8. For any even (resp. odd) thread α (resp. β), the number of odd (resp. even) threads entangled with α (resp. β) is a product of Motzkin numbers.

We have verified Conjecture 5.8 for all even threads of length eleven or less and all odd threads of length twelve or less.

6 Janus Threads

Having investigated the entanglement relation between even and odd threads, we now turn our attention to those threads which are both even and odd.

Definition 6.1. We say a permutation is a Janus thread whenever it is both an even thread and an odd thread. For convenience, we also regard the antipermutation @ as a Janus thread (of length -1). For each $n \ge -1$, we write JT_n to denote the set of Janus threads of length n.

Remarkably, for small n nearly every odd thread is also a Janus thread: the smallest odd thread that is not also an even thread is 3412. Nevertheless, it's natural to ask how many Janus threads of length n there are. In Table 4 we have the number of Janus threads of length nine or less.

$\frac{n}{ JT_n }$	-1	0	1	2	3	4	5	6	7	8	9
$ JT_n $	1	1	1	2	4	8	17	37	82	185	423

Tab. 4: The number of Janus threads of length nine or less.

As was the case for even and odd threads, Janus threads are related to a certain kind of lattice path. To describe these lattice paths, first recall that a *Motzkin path* of length n is a lattice path from (0,0) to (n,0) consisting of unit Up (1,1), Down (1,-1), and Level (1,0) steps which does not pass below the x-axis. As their name suggests, the Motzkin paths of length n are counted by the Motzkin number M_n , which we met in Section 5. A *peak* in a Motzkin path is a pair of consecutive steps in which the first step is an Up step and the second is a Down step. We say a Motzkin path is *peakless* whenever it has no peaks, and we write UD_n to denote the set of peakless Motzkin paths of length n. For example, UD_5 consists of eight Motzkin paths: *LLLLL*, *LLULD*, *LULDL*, *ULDLL*, *LULLD*, *ULLDL*, *ULLLD*, and *ULLLD*. In Table 5 we have the number of peakless Motzkin paths of length ten or less. These paths are counted by

n	0	1	2	3	4	5	6	7	8	9	10
$ UD_n $	1	1	1	2	4	8	17	37	82	185	423

Tab. 5: The number of peakless Motzkin paths of length ten or less.

a certain sequence of generalized Catalan numbers, which is sequence A004148 in the OEIS; its terms satisfy $a_n = a_{n-1} + \sum_{k=1}^{n-2} a_k a_{n-2-k}$ for $n \ge 0$.

As a comparison of Tables 4 and 5 suggests, Janus threads of length n are in bijection with peakless Motzkin paths of length n + 1. As a first step in constructing a bijection between these two sets, we describe how to construct peakless Motzkin paths recursively. **Theorem 6.2.** Suppose n is a positive integer. For each $p \in UD_n$, exactly one of (i) and (ii) below holds.

- (i) There is a unique Motzkin path $a \in UD_{n-1}$ such that p = La.
- (ii) There are unique integers $k \ge 1$ and $l \ge 0$ and unique Motzkin paths $a \in UD_k$ and $b \in UD_l$ such that n = k + l + 2 and p = UaDb.

Conversely, both of the following hold.

- (iii) For every $a \in UD_{n-1}$, we have $La \in UD_n$.
- (iv) For any integers $k \ge 1$ and $l \ge 0$ with n = k + l + 2, and any $a \in UD_k$ and $b \in UD_l$, we have $UaDb \in UD_n$.

Proof: This is similar to the proof of Theorem 4.2.

We can also use our recursive decompositions of the even and odd threads in Theorem 3.8 to describe how to construct Janus threads recursively.

Theorem 6.3. Suppose γ is a permutation of length at least one which begins with its largest entry. Then γ is a Janus thread if and only if there is a Janus thread γ_1 such that $\gamma = 1 \ominus \gamma_1$. Moreover, when these conditions hold, γ_1 is determined by γ .

Proof: (\Rightarrow) Suppose γ is a Janus thread which begins with its largest entry. Since γ begins with its largest entry, it has the form $1 \ominus \gamma_1$ for a unique permutation γ_1 . In particular, the last statement of the theorem holds.

To show γ_1 is an odd thread, first note that since γ is an odd thread, by (2) there is an even thread α_1 and an odd thread β_1 such that $\gamma = (1 \oplus c(\alpha_1) \oplus 1) \oplus \beta_1$. Since γ begins with its largest entry, we must have $\alpha_1 = @$ and $\beta_1 = \gamma_1$, so γ_1 is an odd thread.

To show γ_1 is an even thread, and therefore a Janus thread, first note that since γ is an even thread, by (1) and Theorem 3.15 there is an even thread α_2 and an odd thread β_2 such that $\gamma = c(\beta_2) \ominus 1 \ominus \alpha_2$, and the 1 corresponds to the leftmost eligible connector in γ . If $c(\beta_2)$ begins with its largest entry, then β_2 begins with 1, and the first entry of γ is an eligible connector. However, this contradicts the fact that the leftmost eligible connector in β_2 . Therefore, $c(\beta_2) \ominus 1 \ominus \alpha_2 = \emptyset$. The first of these contradicts the fact that the leftmost eligible entry, we must have $\beta_2 = \emptyset$ or $\beta_2 = \emptyset$. The first of these contradicts the fact that the leftmost eligible entry of γ does not occur in α_2 , so we must have $\gamma = 1 \ominus \alpha_2$. Therefore, $\gamma_1 = \alpha_2$ is an even thread.

(\Leftarrow) This is immediate from Propositions 3.11 and 3.13, the fact that 1 is an odd thread, and the fact that \emptyset is an even thread.

Theorem 6.4. Suppose γ is a permutation of length at least two which does not begin with its largest entry. Then γ is a Janus thread if and only if there are Janus threads γ_1 and γ_2 such that γ_1 has nonnegative length and $\gamma = (1 \oplus c(\gamma_1) \oplus 1) \oplus 1 \oplus \gamma_2$. Moreover, when these conditions hold, γ_1 and γ_2 are determined by γ .

Proof: (\Rightarrow) Suppose γ is a Janus thread which does not begin with its largest entry. Since γ is an odd thread, by (2) there is an even thread α_1 and an odd thread β_1 such that $\gamma = (1 \oplus c(\alpha_1) \oplus 1) \oplus \beta_1$. The

second 1 in this decomposition corresponds to the largest entry of γ , so this decomposition is unique, and the last statement of the theorem holds. Moreover, since γ does not begin with its largest entry, α_1 has nonnegative length. Since all odd threads have nonnegative length, β_1 also has nonnegative length.

We claim β_1 begins with its largest entry. To prove this, suppose by way of contradiction that β_1 does not begin with its largest entry. Then by (2) there is an even thread α_2 of nonnegative length and an odd thread β_2 such that $\beta_1 = (1 \oplus c(\alpha_2) \oplus 1) \oplus \beta_2$. Repeating this argument as long as β_j does not begin with its largest entry, we find there are even threads $\alpha_1, \ldots, \alpha_k$ of nonnegative length and an odd thread β_{k+1} such that

$$\gamma = (1 \oplus c(\alpha_1) \oplus 1) \oplus (1 \oplus c(\alpha_2) \oplus 1) \oplus \dots \oplus (1 \oplus c(\alpha_k) \oplus 1) \oplus 1 \oplus \beta_{k+1}.$$
(10)

Furthermore, because γ is an even thread, $c(\gamma)$ must have a fixed point which is a left-to-right maximum. This cannot occur in any of the summands $1 \oplus c(\alpha_j) \oplus 1$, so the leftmost eligible connector in γ must correspond to the rightmost 1 in the decomposition in (10). By Theorem 3.15, the permutation $(1 \oplus c(\alpha_1) \oplus 1) \oplus (1 \oplus c(\alpha_2) \oplus 1) \oplus \cdots \oplus (1 \oplus c(\alpha_k) \oplus 1)$ is the complement of an odd thread and β_{k+1} an even thread. Now it follows that $(1 \oplus \alpha_1 \oplus 1) \oplus \cdots \oplus (1 \oplus \alpha_k \oplus 1)$ is an odd thread, and by (2) there is an even thread α and an odd thread β such that $(1 \oplus \alpha_1 \oplus 1) \oplus \cdots \oplus (1 \oplus \alpha_k \oplus 1) = (1 \oplus c(\alpha) \oplus 1) \oplus \beta$. Matching largest entries on each side of this equation, we find $\beta = \alpha_k \oplus 1$. But all of the entries of β in $(1 \oplus c(\alpha) \oplus 1) \oplus \beta$ are less than every entry to the left of the largest entry, while all of the entries of $\alpha_k \oplus 1$ in $(1 \oplus \alpha_1 \oplus 1) \oplus \cdots \oplus (1 \oplus \alpha_k \oplus 1)$ are greater than all of the entries to the left of the largest entry. This contradicts the facts that $k \ge 2$ and $\alpha_1 \neq 0$.

Since β_1 begins with its largest entry, we now know that we have $\gamma = (1 \oplus c(\gamma_1) \oplus 1) \oplus 1 \oplus \gamma_2$, where γ_1 has nonnegative length, γ_1 is an even thread, $1 \oplus \gamma_2$ is an odd thread, and $1 \oplus c(\gamma_1) \oplus 1$ is the complement of an odd thread. Furthermore, the rightmost 1 in this decomposition is the leftmost eligible connector in γ , so by Theorem 3.15 we see that γ_2 is an even thread. On the other hand, since $1 \oplus \gamma_2$ is an odd thread, by Proposition 3.12, we have that γ_2 is also an odd thread. Therefore, γ_2 is a Janus thread.

To see that γ_1 is an odd thread, first note that $1 \oplus c(\gamma_1) \oplus 1$ is the complement of an odd thread means $1 \oplus \gamma_1 \oplus 1$ is an odd thread. Now the result follows from Proposition 3.12.

(\Leftarrow) Suppose γ_1 and γ_2 are Janus threads. The fact that $(1 \oplus c(\gamma_1) \oplus 1) \oplus 1 \oplus \gamma_2$ is an odd thread is immediate from Proposition 3.11, the fact that 1 is an odd thread, and (2). Similarly, since γ_1 is an odd thread by Proposition 3.11. Now the fact that $(1 \oplus c(\gamma_1) \oplus 1) \oplus 1 \oplus \gamma_2$ is an even thread follows from (1), since γ_2 is an even thread.

The fact that our Janus thread decompositions exactly match our peakless Motzkin path decompositions allows us to construct a recursive bijection between these two sets.

Theorem 6.5. For each integer $n \ge -1$, there is a unique bijection

$$K: JT_n \to UD_{n+1}$$

such that $K(@) = \emptyset$, $K(\emptyset) = L$, if $\gamma = 1 \ominus \gamma_1$ for a Janus thread γ then $K(\gamma) = LK(\gamma_1)$ and if $\gamma = (1 \oplus c(\gamma_1) \oplus 1) \ominus 1 \ominus \gamma_2$ for Janus threads γ_1 and γ_2 then $K(\gamma) = UK(\gamma_1)DK(\gamma_2)$.

Proof: This is similar to the proof of Theorems 4.4 and 4.6, using Theorems 6.2, 6.3, and 6.4. \Box

Although our description of K above is recursive, it turns out this map has an elegant direct combinatorial description as well. We close the section with this alternative description of K.

Definition 6.6. For any Janus thread of length $n \ge -1$, we define the lattice path $\mathcal{K}(\gamma)$ as follows. For n = -1 or n = 0 we have $\mathcal{K}(@) = \emptyset$ and $\mathcal{K}(\emptyset) = L$. If $n \ge 1$, then we obtain $\mathcal{K}(\gamma)$ as follows.

- 1. Write n + 1, followed by γ , followed by 0.
- 2. For each pair of consecutive entries in this new sequence, if the entries are consecutive integers (in either order) then write L between them. If the consecutive entries are not consecutive integers, then write U between them if they form an ascent, and D if they form a descent.
- 3. In the resulting sequence of n + 1 Us, Ls, and Ds, number the subsequence of Us and Ds from left to right, beginning with 1. Change every odd-numbered U to a D and every odd-numbered D to a U.

The resulting sequence of Us, Ls, and Ds is the lattice path $\mathcal{K}(\gamma)$.

Example 6.7. When $\gamma = 576894312$ we write 10 576894312 0, and we obtain the sequence DULULDLDLD. Changing the odd-numbered Us to Ds and the odd-numbered Ds to Us, we find that $\mathcal{K}(578694312) = UULDLDLULD$.

In Example 6.7 we obtain a peakless Motzkin path, and in fact it's not difficult to check that $\mathcal{K}(576894312) = K(576894312)$. As we show in our final result, $\mathcal{K}(\gamma)$ is a peakless Motzkin path for every Janus thread γ , and $\mathcal{K}(\gamma) = K(\gamma)$.

Theorem 6.8. For any Janus thread γ , we have $\mathcal{K}(\gamma) = K(\gamma)$.

Proof: The result is easy to check when γ has length less than two, so suppose $|\gamma| \ge 3$; we argue by induction on the length of γ .

If γ begins with its largest entry, then by Theorem 6.3 there is a Janus thread γ_1 for which $\gamma = 1 \ominus \gamma_1$. Since $|\gamma_1| = |\gamma| - 1 \ge 2$, we have $\mathcal{K}(\gamma) = L\mathcal{K}(\gamma_1)$. By induction, $\mathcal{K}(\gamma_1) = K(\gamma_1)$, so $\mathcal{K}(\gamma) = K(\gamma)$ by the definition of K.

If γ does not begin with its largest entry, then by Theorem 6.4 there are Janus threads γ_1 and γ_2 for which $\gamma = (1 \oplus c(\gamma_1) \oplus 1) \oplus 1 \oplus \gamma_2$, where γ_1 has nonnegative length. Examining $\mathcal{K}((1 \oplus c(\gamma_1) \oplus 1) \oplus 1 \oplus \gamma_2)$, we find $\mathcal{K}(\gamma) = U\mathcal{K}(\gamma_1)D\mathcal{K}(\gamma_2)$. Now the result follows by induction and the definition of K. \Box

7 Future Directions

This work originated in a larger effort to give a nonrecursive characterization of the snow leopard permutations. For instance, we have made extensive use of the fact that snow leopard permutations preserve parity, but most permutations which preserve parity are not snow leopard permutations. Similarly, we noted in the Introduction that snow leopard permutations are anti-Baxter permutations, so they avoid the vincular patterns $3\underline{412}$ and $2\underline{143}$. Still, most anti-Baxter permutations which preserve parity are not snow leopard permutations. Caffrey et al. (2015) introduce in their Definition 3.2 a function κ which maps each permutation to a lattice path consisting of North and East steps, and they show that κ is a bijection between the set of snow leopard permutations of length 2n - 1 and the set of Catalan paths from (0,0) to (n,n). However, there are many other permutations π for which $\kappa(\pi)$ is a Catalan path. Indeed, let's call a permutation SLP-like whenever it is an anti-Baxter permutation which preserves parity and maps to a Catalan path under κ . Then in Table 6 we see that most SLP-like permutations are still not snow leopard

length	1	3	5	7	9
number of SLP-like permutations	1	2	7	32	175
number of snow leopard permutations	1	2	5	14	42

Tab. 6: The number of SLP-like permutations compared with the number of snow leopard permutations.

permutations. Giving a nonrecursive characterization of the snow leopard permutations remains an open problem. Similarly, it remains an open problem to prove our Conjecture 5.8. Finally, as far as we know no one has investigated the permutations which are compatible with alternating Baxter permutations, even though Cori et al. (1986) have shown that the number of alternating Baxter permutations of length 2n (resp. 2n + 1) is C_n^2 (resp. $C_n C_{n+1}$). We conjecture that the compatibility relation is a bijection in this case, just as it is in the case of doubly alternating permutations and snow leopard permutations.

References

- A. Asinowski, G. Barequet, M. Bousquet-Mélou, T. Mansour, and R. Y. Pinter. Orders induced by segments in floorplans and (2 14 3, 3 41 2)-avoiding permutations. *Electron. J. Combin.*, 20(2): #R35, 2013.
- G. Baxter. On fixed points of the composite of commuting functions. *Proc. Amer. Math. Soc.*, 15:851–855, 1964.
- D. Bressoud. *Proofs and Confirmations: The Story of the Alternating-Sign Matrix Conjecture.* The Mathematical Association of America, 1999.
- B. Caffrey, E. S. Egge, G. Michel, K. Rubin, and J. V. Steegh. Domino tilings of Aztec diamonds, Baxter permutations, and snow leopard permutations. *Involve*, 8(5):833–858, 2015.
- H. Canary. Aztec diamonds and Baxter permutations. Electron. J. Combin., 17(1):#R105, 2010.
- F. R. K. Chung, R. L. Graham, V. E. H. Jr., and M. Kleiman. The number of Baxter permutations. J. Combin. Theory Ser. A, 24(3):382–394, 1978.
- R. Cori, S. Duluq, and G. Viennot. Shuffle of parenthesis systems and Baxter permutations. J. Combin. Theory Ser. A, 43:1–22, 1986.
- K. Dilks. Involutions on Baxter objects. arXiv:1402.2961 [math.CO], 2014.
- R. Donaghey and L. W. Shapiro. Motzkin numbers. J. Combin. Theory, Series A, 23(3):291-301, 1977.
- E. S. Egge. Restricted 3412-avoiding involutions, continued fractions, and Chebyshev polynomials. *Adv. Appl. Math.*, 33:451–475, 2004.
- N. Elkies, G. Kuperberg, M. Larsen, and J. Propp. Alternating-sign matrices and domino tilings. I. J. *Algebraic Combin.*, 1(2):111–132, 1992.
- O. Guibert. Combinatoire des permutations à motifs exclus en liaison avec mots, cartes planaires et tableaux de Young. PhD thesis, Université Bordeaux I, 1995.

- O. Guibert and S. Linusson. Doubly alternating Baxter permutations are Catalan. *Discrete Math.*, 217 (1–3):157–166, 2000.
- S. Kitaev. Patterns in Permutations and Words. Springer-Verlag, 2011.
- J. Propp. The many faces of alternating-sign matrices. *Discrete Math. Theor. Comput. Sci. Proc., AA*, 2001.
- D. P. Robbins. The story of 1, 2, 7, 42, 429, 7436, Math. Intelligencer, 13(2):12–19, 1991.
- A. Sapounakis, I. Tasoulas, and P. Tsikouras. Counting strings in Dyck paths. *Discrete Math.*, 307(23): 2909–2924, 2007.
- R. P. Stanley. Enumerative Combinatorics, volume 2. Cambridge University Press, 1999.