

Reachability problems for PAMs ^{*}

Oleksiy Kurgansky¹ and Igor Potapov²

¹ Institute of Applied Mathematics and Mechanics, NAS of Ukraine

² Department of Computer Science, University of Liverpool, UK

Abstract. Piecewise affine maps (PAMs) are frequently used as a reference model to show the openness of the reachability questions in other systems. The reachability problem for one-dimensional PAM is still open even if we define it with only two intervals. As the main contribution of this paper we introduce new techniques for solving reachability problems based on p -adic norms and weights as well as showing decidability for two classes of maps. Then we show the connections between topological properties for PAM's orbits, reachability problems and representation of numbers in a rational base system. Finally we show a particular instance where the uniform distribution of the original orbit may not remain uniform or even dense after making regular shifts and taking a fractional part in that sequence.

Keywords: *Reachability problems, piecewise affine maps (PAMs), β -expansion, p -adic analysis*

1 Introduction

The simplification of real programs shows that there is a number of quite basic models/fragments for which we have fundamental difficulties in the design of verification tools. One of them is the model of iterative map that appears in many different contexts, including discrete-event/discrete-time/hybrid systems, qualitative biological models, chaos-based cryptography, etc [1, 8, 16, 23].

The one-dimensional affine piecewise iterative map is a very rich mathematical object and at the same time one of the simplest dynamical system producing very complex and sensitive effects. A function $f: \mathbb{Q} \rightarrow \mathbb{Q}$ is a one-dimensional piecewise-affine map (PAM) if f is of the form $f(x) = a_i x + b_i$ for $x \in X_i$ where all coefficients a_i, b_i and the extremities of a finite number of bounded intervals X_i are rational numbers. Let us consider the sequence of iterations starting from a rational point x : $x, f(x), f^2(x) = f(f(x))$, and so on. The reachability in PAM is a problem to decide for a given f and two rational points x and y whether y is reachable from x . In other words, is there an $n \in \mathbb{N}$ such that $f^n(x) = y$?

The decidability of the reachability problem for one dimensional piecewise-affine map is still an open problem, which is related to other challenging questions in the theory of computation, number theory and linear algebra [7, 14, 15]. This

^{*} This research is supported by EPSRC grant Reachability problems for words, matrices and maps (EP/M00077X/1)

model plays a crucial role in the recent research about verification of hybrid systems [2, 3], timed automata [2] control systems [9, 10], representation of numbers in a rational base (β -expansions) [19, 21], discounted sum automata [11]. In particular PAM is often used as a reference model to show the openness of the reachability questions in other systems. It also has a very natural geometrical interpretation as pseudo-billiard system [17] and Hierarchical Piecewise Constant Derivative (HPCD) system [3]. The reachability problem for one-dimensional PAM is still open even if we define it with only two intervals [2, 3, 5, 12].

The primary goal of this paper is to demonstrate new approaches for solving reachability problem in PAMs, connecting reachability questions with topological properties of maps and widening connections with other important theoretical computer science problems. First, we show new techniques for decidability of the reachability problem in PAMs based on p -adic norms and weights. We illustrate these techniques showing decidability of two classes of PAMs. The algorithm in Theorem 1 solves point to point reachability problem for two-interval injective PAM under the assumption that a PAM has bounded invariant densities. While our numerical experiments shows that the sequence of invariant densities converge to smooth functions it is not yet clear whether it holds for all PAMs or if not whether this property can be algorithmically checked.

Following the proposed approach based on p -adic weights in Theorem 2 we define another fragment of PAMs for which the reachability problem is decidable. In particular we remove the condition on bounded invariant densities and injectivity of piecewise-affine map and consider a PAM f with a constraint on linear coefficients in affine maps. This class of PAMs is also related to encoding of rational numbers in the rational base (β -expansions). The decidability of the point-to-point problem for this class is shown in Theorem 2 and decidability of point-to-set problem for the same class can give someone an answer to the open problem related to β -expansions.

Then we establish the connections of topological properties for PAM's orbits with reachability problems and representation of numbers in a rational base system. We show that the reachability problems for above objects tightly connected to questions about distribution of the fractional parts in the generated sequences and moreover about distribution of the fractional part after regular shifts.

2 Preliminaries and Notations

In what follows we use traditional denotations \mathbb{N} , \mathbb{Z} , $\mathbb{Z}^+ = \{0, 1, 2, \dots\}$, \mathbb{P} , \mathbb{Q} and \mathbb{R} for sets of natural, integers, positive integers, primes, rational and real numbers, respectively. Let us denote by $S^1 = \mathbb{Q}/\mathbb{Z}$ the unit circle which consists only rational numbers. By $\{x\}$, $\lfloor x \rfloor$ and $\lceil x \rceil$ we denote the fractional part ³ of a number, floor and ceiling functions.

Let Y be a set of numbers and x is a single number, then we define their addition and multiplication as follows: $Y + x = x + Y = \{x + y | y \in Y\}$ and

³ It will be clear from the context if brackets are used in other conventional ways, for example, to indicate a set of numbers.

$xY = Yx = \{xy|y \in Y\}$. The application of a function $f : X \rightarrow Y$ to a set $X' \subseteq X$ is defined as $f(X') = \{f(x)|x \in X'\}$. If $f \subseteq X \times Y$ is a nondeterministic map, i.e. $f : X \rightarrow 2^Y$ and $x \in X$, $X' \subseteq X$, we define $f(x) = \{y|(x, y) \in f\}$ and $f(X') = \bigcup_{x \in X'} f(x)$.

p -adic norms and weights: Let us consider an arbitrary finite set of prime numbers $\mathbb{F} = \{p_1, p_2, \dots, p_k\} \subset \mathbb{P}$ in ascending order and define the product of prime numbers from \mathbb{F} by $m = p_1 p_2 \dots p_k$. Let x be a positive rational number that can be represented by primes from a set \mathbb{F} . Then its prime factorization is $x = \prod_{p \in \mathbb{F}} p^{\alpha_p}$, where $\alpha_p \in \mathbb{Z}$, $p \in \mathbb{F}$.

Any nonzero rational number x can be represented by $x = (p^{\alpha_p} r)/s$, where p is a prime number, r and s are integers not divisible by p , and α_p is a unique integer. The p -adic norm of x is then defined by $|x|_p = p^{(-\alpha_p)}$. The p -adic weight of x is defined as $\|x\|_p = \log_p(|x|_p)$, i.e. $\|x\|_p = -\alpha_p$. The following properties of p -adic weights are directly follows from the properties of p -adic norm:

$$\|x\|_p = \|y\|_p \Rightarrow \|x + y\|_p \leq \|x\|_p. \quad (1)$$

$$\|x\|_p < \|y\|_p \Rightarrow \|x + y\|_p = \|y\|_p, \quad (2)$$

$$\|x \cdot y\|_p = \|x\|_p + \|y\|_p, \quad (3)$$

$$\|x^r\|_p = r\|x\|_p, \quad (4)$$

If there is a prime $p \notin \mathbb{F}$ such that $\|x\|_p > 0$, then we define $\|x\|_m = +\infty$, otherwise $\|x\|_m = \max_{p \in \mathbb{F}} \|x\|_p$.

By m -weight and m -vector-weight of x in respect to a set \mathbb{F} we denote $\|x\|_m$ and $(\|x\|)_m = (\|x\|_{p_1}, \|x\|_{p_2}, \dots, \|x\|_{p_k})^T$ respectively. Informally speaking the m -weight of a number x (if $\|x\|_m > 0$) is the number of digits after the decimal point in the representation of x in base m , i.e. x can be written as $x = y \cdot m^{-\|x\|_m}$, where y is an integer which the last digit in the m -ary representation is non zero. If $\|x\|_m \leq 0$, then x is an integer number.

Without loss of generality let us consider from now on only such $x \in X$ for which $\|x\|_m < +\infty$. Alternatively if $\|x\|_m = +\infty$, it is enough to change the set $\mathbb{F} = \{p_1, \dots, p_k\}$ in order to fulfill the requirements of $\|x\|_m < +\infty$.

Lemma 1. *For all rational $x \in [0, 1]$ with an upper bound $a \in \mathbb{Z}^+$ on $\|x\|_m$, i.e. $\|x\|_m < a$, there is a lower bound $b \in \mathbb{Z}$ based on a and \mathbb{F} such that $\|x\|_p \geq b$ for all $p \in \mathbb{P}$.*

Proof. Let us denote two sums of weights: $\alpha = -\sum_{p \in \mathbb{P}, \|x\|_p \leq 0} \|x\|_p$ and $\beta = \sum_{p \in \mathbb{F}, \|x\|_p \geq 1} \|x\|_p$. Assuming that 2 is the smallest possible prime number and p_k is the largest number in \mathbb{F} , we have the following inequality: $\frac{2^\alpha}{p_k^{\beta a}} \leq \frac{2^\alpha}{p_k^\beta} \leq x \leq 1$. Then $-\alpha \geq -k a \log_2 p_k$ and if $b = -\alpha$ we have $\|x\|_p \geq b$ for all $p \in \mathbb{P}$. \square

Corollary 1. *For any $a \in \mathbb{Z}$ there is only a finite number of rational $x \in [0, 1]$ for which $\|x\|_m < a$.*

Reachability problem for PAMs: We say that $f : S^1 \rightarrow S^1$ is a one-dimensional piecewise affine map (PAM) whenever f is of the form $f(x) = a_i x + b_i$, where $a_i, b_i \in \mathbb{Q} \Leftrightarrow x \in X_i$, $S^1 = X_1 \cup X_2 \cup \dots \cup X_l$ and where $\{X_1, \dots, X_l\}$ is a finite family of disjoint (rational) intervals. If the intervals are not disjoint we call it *non-deterministic piecewise affine map* and by default a piecewise-affine mapping is understood to be deterministic. The derivative f' of a PAM f we define as $f'(x) = a_i$ for $x \in X_i$, $1 \leq i \leq l$.

If f is deterministic PAM, an orbit (trajectory) of a point x is denoted by $O_f(x)$ and will be understood either as a set $O_f(x) = \{f^i(x) | i \in \mathbb{Z}^+\}$ or as a sequence, i.e. $O_f(x) : \mathbb{Z}^+ \rightarrow S^1$, $O_f(x)(i) = f^i(x)$. We also define that a point y is reachable from x if $y \in O_f(x)$.

In general the reachability problem for PAM can be defined as follows. Given a PAM f , $x \in S^1$ and $Y \subseteq S^1$, decide whether the intersection $Y \cap O_f(x)$ is empty. If Y is a finite union of intervals, we name the reachability problem as *point-to-set (interval) problem*. If Y is a one element set (i.e. a single point), the reachability problem is known as *point-to-point reachability*.⁴

In this paper we only consider one-dimensional PAMs and by the *reachability problem for PAM* we understand point-to-point reachability and explicitly state the type of the problem when we need to refer to other reachability questions. Note that the point-to-interval reachability can be reduced to point-to-point reachability problem by extending a map with a few intervals in which the current value is just sequentially deleted. It works for all PAMs but may not preserve the properties and the form of the original map.

Generally speaking the piecewise-affine mapping does not need to be defined as $f : X \rightarrow X$, where $X = S^1 = [0, 1)$. However if the set X is a union of any finite number of bounded intervals we always can scale it to S^1 . If $f : X \rightarrow X$ is such that $X \neq [0, 1)$, and $X \subseteq [a, b)$, then by applying conjugation $h(x) = \frac{x-a}{b-a}$ the original reachability problem for f is reduced to the reachability problem for the mapping $g = h \circ f \circ h^{-1}$ from $[0, 1)$ to $[0, 1)$. Moreover the interest to PAMs as $f : S^1 \rightarrow S^1$ is also motivated by their use in the research of chaotic systems.

3 Decidability using p -adic norms

It is well known in dynamical systems research that due to complexity of orbits in iterative maps it is less useful, and perhaps misleading, to compute the orbit of a single point and it is more reasonable to approximate the statistics of the underlying dynamics [13, 20]. This information is encoded in the so-called *invariant measures*, which specify the probability to observe a typical trajectory within a certain region of state space and their corresponding *invariant densities*.

Let us consider a density as an ensemble of initial starting points (i.e. initial conditions). The action of the dynamical system on this ensemble is described by the Perron-Frobenius operator. The ensembles which are fixed under the linear

⁴ Also in a similar way it is possible to define set-to-point and set-to-set reachability problems.

Perron-Frobenius operator is known as invariant densities or in other words, they are eigenfunctions with eigenvalue 1 [13].

Formally under an ensemble A we understand an enumerated set (sequence) of points in phase space. With ensemble we can associate the distribution function and the density function. Let I be a set of points. We denote by $F_I^A(n) = |\{i \in \mathbb{Z}^+ | i \leq n, A(i) \in I\}|$ the number of elements in the sequence A which belong to the set I and which indexes are less or equal n . The distribution function of the ensemble A is defined as $\Phi_A(x) = \lim_{n \rightarrow \infty} \frac{F_{(-\infty, x]}^A(n)}{n}$, if the limits exist. The density function ϕ_A of the ensemble A is defined as $\phi_A(x) = \Phi'_A(x)$.

Suppose given an ensemble A_0 with density ϕ_0 . If we apply PAM f to each point of the ensemble, we get a new ensemble A_1 with some density distribution ϕ_1 . We say that the function ϕ_1 is obtained from ϕ_0 using the Frobenius-Perron or transfer operator, which we denote by L_f . It is known that

$$\phi_1(x) = L_f(\phi_0)(x) = \sum_{y \in f^{-1}(x)} \frac{\phi_0(y)}{|f'(y)|}.$$

If $\phi_1 = \phi_0$ we say that ϕ_0 is a f -invariant density function or an eigenfunction of the transfer operator L_f .

We prove that if for an injective PAM f there exists an invariant bounded density function then the reachability problem for f is decidable.

Lemma 2. *Let f be an injective PAM, and ϕ be a f -invariant density function. If there are $K_{min} > 0$ and $K_{max} < +\infty$ such that for any x from the domain of f the following inequality holds: $K_{min} < \phi(x) < K_{max}$, then for an arbitrary segment of the orbit x_1, x_2, \dots, x_{n+1} , where $x_{i+1} = f(x_i)$, we have $\frac{K_{min}}{K_{max}} \leq |c_1 \cdot c_2 \cdot \dots \cdot c_n| \leq \frac{K_{max}}{K_{min}}$, where $c_i = a_j$ if $x_i \in X_j$.*

Proof. Let ϕ be an eigenfunction of the Perron-Frobenius operator for an injective PAM f . Then injectivity of f and the fact that $y = f(x)$ implies that $\phi(y) = \frac{\phi(x)}{|f'(x)|}$. We denote $f'(x_i)$ by c_i . Then $\phi(x_{n+1}) = \frac{\phi(x_1)}{|c_1 \cdot c_2 \cdot \dots \cdot c_n|}$ and $|c_1 \cdot c_2 \cdot \dots \cdot c_n| = \frac{\phi(x_1)}{\phi(x_{n+1})}$. Now we can bound $|c_1 \cdot c_2 \cdot \dots \cdot c_n|$ by $\frac{K_{min}}{K_{max}}$ and $\frac{K_{max}}{K_{min}}$.

Theorem 1. *Given an injective PAM f with two intervals and the existence of a f -invariant density function ϕ such that there are $K_{min} > 0$ and $K_{max} < +\infty$ and the following inequality holds $K_{min} < \phi(x) < K_{max}$ for all x from the domain of f . Then the reachability problem for f is decidable.*

Proof. A piecewise-affine map f with two intervals X_1 and X_2 is defined as follows: $f(x) = a_i x + b_i$, if $x \in X_i$. Note that we only consider PAMs over rational numbers where a starting point as well as all coefficients and borders of intervals are rational numbers. Let us consider an arbitrary pair of points $x, y \in X$ such that $y \in O_f(x)$ and the sequence of points x_1, x_2, \dots, x_{n+1} from the orbit $O_f(x)$ such that $x_1 = x, x_{n+1} = y, x_{i+1} = f(x_i), 1 \leq i \leq n$.

If it would be possible to find a computable upper bound M on the length n of such reachability sequence from x to y (based on $K_{min}, K_{max}, a_1, a_2, b_1, b_2$,

x, y) we could solve the reachability problem by considering only initial segment of the reachability path of length less or equal to M .

In general the existence of such bound is not obvious, however if we can show that in the path x_1, x_2, \dots, x_{n+1} where $x_1 = x$ and $x_{n+1} = y$ the p -adic weights $\|x_i\|_m$ for all $i \in \{1, \dots, n+1\}$ are bounded by some value M_1 then we can restrict M as follows: $n < M < m^{M_1}$ as $\|x_i\|_m$ is the number of decimal places in the representation of x_i in base m . In particular if there is a orbit's segment of length that is greater than m^{M_1} and which consists of numbers with p -adic weights less than M_1 then it always contains two identical numbers and the orbit loops.

So in order to prove a computable upper bound on $\|x_i\|_m$ it is sufficient to prove that for any $p \in \mathbb{F}$ p -adic weights (i.e. logarithmic norms) $\|x_i\|_p$ are bounded from above by a computable number M_2 for all $i \in \{1, \dots, n+1\}$. Note that by the definition of the logarithmic m -norm, $M_2 = M_1$.

Let us define a number $h = \max\{\|b_1\|_m + 1, \|b_2\|_m + 1, \|x_1\|_m, \|x_{n+1}\|_m\}$. Let us take an arbitrary $p \in \mathbb{F}$ and select a subsequence x_j, x_{j+1}, \dots, x_r in the sequence x_1, x_2, \dots, x_{n+1} such that its elements $x \in \{x_{j+1}, x_{j+2}, \dots, x_{r-1}\}$ have the following property $\|x\|_p > h$, but at the same time $\|x_j\|_p \leq h$ and $\|x_r\|_p \leq h$. For simplicity, but without loss of generality, we assume that $j = 1, r = n+1$ and $\|x_1\|_p = \|x_{n+1}\|_p = h$.

Later we either will find the computable upper bound on n or will show a computable bound M_2 on p -adic weights based on $K_{min}, K_{max}, a_1, a_2$ and h . As stated above, it is enough for proving the theorem.

Let us define $x_{i+1} = f(x_i) = c_i x_i + d_i$, where $c_i \in \{a_1, a_2\}, d_i \in \{b_1, b_2\}$. From the properties (1) and (2) of the p -adic weights and the fact that $\|x_i\|_p > \|b_j\|_p, i \in \{1, 2, \dots, n\}, j \in \{1, 2\}$ follows that $\|x_{i+1}\|_p = \|x_i\|_p + \|c_i\|_p$. This implies $\|x_{n+1}\|_p = \|x_1\|_p + \sum_{i=1}^n \|c_i\|_p = \|x_1\|_p + \sum_{i=1}^n \alpha_i \|a_i\|_p$ for some non-negative numbers α_1 and α_2 , where $\alpha_1 + \alpha_2 = n$. Taking into account that $\|x_1\|_p = \|x_{n+1}\|_p$ we have $\sum_{i=1}^n \alpha_i \|a_i\|_p = 0$.

Let r be the greatest common divisor of α_1 and α_2 . We define $\alpha = \alpha_1/r$ and $\beta = \alpha_2/r$, i.e. α and β are smallest non-negative integers such that $\alpha \|a_1\|_p + \beta \|a_2\|_p = 0$. Thus, we obtain $\sum_{i=1}^n \alpha_i \|a_i\|_p = r(\alpha \|a_1\|_p + \beta \|a_2\|_p)$. By Lemma 2 we have: $\frac{K_{min}}{K_{max}} \leq |c_1 \cdot c_2 \cdot \dots \cdot c_n| \leq \frac{K_{max}}{K_{min}}$, i.e. $\frac{K_{min}}{K_{max}} \leq |a_1^\alpha a_2^\beta|^r \leq \frac{K_{max}}{K_{min}}$,

Now we consider two cases $|a_1^\alpha a_2^\beta| \neq 1$ and $|a_1^\alpha a_2^\beta| = 1$. The first case corresponds to linearly independent columns of the matrix A_f , $\text{rank}(A) = 2$, and the second case to linear dependence, $\text{rank}(A_f) < 2$.

Let $|a_1^\alpha a_2^\beta| \neq 1$, then either $|a_1^\alpha a_2^\beta| > 1$ or $|a_1^\alpha a_2^\beta| < 1$. If $|a_1^\alpha a_2^\beta| > 1$ then from $|a_1^\alpha a_2^\beta|^r \leq \frac{K_{max}}{K_{min}}$ follows that $r \leq \frac{\ln \frac{K_{max}}{K_{min}}}{\ln |a_1^\alpha a_2^\beta|}$. On the other hand if $|a_1^\alpha a_2^\beta| < 1$, then from $|a_1^\alpha a_2^\beta|^r \geq \frac{K_{min}}{K_{max}}$ follows $r \leq \frac{\ln \frac{K_{min}}{K_{max}}}{\ln |a_1^\alpha a_2^\beta|}$.

Now suppose that $|a_1^\alpha a_2^\beta| = 1$. The only interesting case is when $a_1 \neq 1$ and $a_2 \neq 1$ as the cases where $|a_1| = 1, |a_2| = 1$ or both correspond to the trivial case of piecewise-affine mapping (i.e. with no more than one linear factor).

Without loss of generality, we assume that $a_1 > 1$ and $a_2 < 1$. Note also that $\|a_2\|_p = -\frac{\alpha}{\beta}\|a_1\|_p$. Let us now consider an arbitrary subsequence of consecutive points $x_1x_2 \dots x_{j+1}$, $j \leq n$ of the original reachability path. Assuming that α_1, α_2 such that $|c_1c_2 \dots c_j| = |a_1|^{\alpha_1}|a_2|^{\alpha_2}$ we have that

$$\|x_{j+1}\|_p = \|x_1\|_p + \alpha_1\|a_1\|_p + \alpha_2\|a_2\|_p = (\alpha_1 - \alpha_2\frac{\alpha}{\beta})\|a_1\|_p.$$

On the other hand from Lemma 2 we know that: $\frac{K_{min}}{K_{max}} \leq |a_1|^{\alpha_1}|a_2|^{\alpha_2} \leq \frac{K_{max}}{K_{min}}$. Since $|a_1^{\alpha_1}a_2^{\alpha_2}| = 1$, then $|a_2| = |a_1|^{-\frac{\alpha}{\beta}}$ and

$$\frac{K_{min}}{K_{max}} \leq |a_1|^{\alpha_1 - \alpha_2\frac{\alpha}{\beta}} \leq \frac{K_{max}}{K_{min}}.$$

Taking into account assumption that $|a_1| > 1$, we get $\alpha_1 - \alpha_2\frac{\alpha}{\beta} \leq \frac{\ln \frac{K_{max}}{K_{min}}}{\ln |a_1|}$.

Now it follows that:

$$\|x_{j+1}\|_p = \|x_1\|_p + (\alpha_1 - \alpha_2\frac{\alpha}{\beta})\|a_1\|_p \leq h + \left(\frac{\ln \frac{K_{max}}{K_{min}}}{\ln |a_1|} \right) \|a_1\|_m.$$

Thus, we have shown a computable upper bound for p -adic weights of the orbital elements that can reach a point y . Finally, in view of provided reasoning, we shown that the reachability problem for this type of PAMs is decidable. \square

The theorem can be applied for a larger class of PAMs if more information would be known about the convergence of density functions under the action of the Perron-Frobenius operator. Let us call an ensemble A to be statistically fixed with respect to f , if $\phi_A = L_f(\phi_A)$. E.g. if someone can show that in injective PAM all statistically fixed ensembles have identical distribution functions then Theorem 1 can be applied to show decidability in injective PAMs.

Following the proposed approach based on p -adic weights we define another fragment of PAMs for which the reachability problem is decidable. In particularly we remove the condition on eigenfunction of the transfer operator and injectivity of piecewise-affine map and consider a PAM f with only constraints on linear coefficients in affine maps. More specifically we require that the powers of prime numbers from prime factorizations of linear coefficients should have the same signs (i.e. two sets of prime numbers used in nominator and denominator are disjoint). Let us denote for a PAM f a matrix A_f with values (a_{ji}) , where $a_{ji} = \|a_i\|_{p_j}$, $1 \leq i \leq l$, $1 \leq j \leq k$. The rank of A_f is denoted by $\text{rank}(A_f)$.

Theorem 2. *The reachability problem for a PAM f is decidable if every row of a matrix A_f contains values of the same sign, (i.e. $a_{ji} \cdot a_{j'i} \geq 0$, for all i, j such that $1 \leq i \leq l$, $1 \leq j, j' \leq k$).*

Proof. Let us consider a PAM f of the form $f(x) = a_i x + b_i$ for $x \in X_i$ where all coefficients a_i, b_i and the extremities of a finite number of bounded intervals X_i are rational numbers. Let us define $h = \max\{\|b_1\|_m, \|b_2\|_m, \dots, \|b_l\|_m\}$. The

condition of the theorem means that for any prime $p \in \mathbb{F}$ all linear coefficients of the map f have non-zero p -adic weights of the same sign.

In this case, if p -adic weights of linear coefficients of f are non-negative, then for any $x \in X$ from $\|x\|_p > h$ follows that $\|f(x)\|_p \geq \|x\|_p$ and therefore $\|f(x)\|_m \geq \|x\|_m$ (i.e. m -adic weight does not decrease). If p -adic weight of linear coefficients of the mapping are negative, then for any $x \in X$ we have $\|f(x)\|_p \leq \max\{\|x\|_p, h\}$.

Thus, in the sequence of reachable points for an orbit of a map f either all points of the orbit have m -adic weights bounded from above by h , then we have a cyclic orbit, or from some moment when m -adic weight of a reachable point exceeds h it does not decrease and again, either orbit loops or m -adic weight increases indefinitely.

Thus, in order to decide whether y is reachable, i.e. $y \in O_f(x)$, it is sufficient to start generating a sequence of reachable points in the orbit $O_f(x)$ and wait for one of the events, where either 1) a point in the orbit is equal to y (y is reachable), or 2) the orbit will loop and $y \notin O_f(x)$ (y is not reachable), or 3) a point x' is reachable, such that $\|x'\|_m > \max\{h, \|y\|_m\}$, and then $y \notin O_f(x)$ (y is not reachable). \square

Definition 1. A piecewise affine mapping $f : S^1 \rightarrow S^1$ is complete if for a set of disjoint intervals $S^1 = X_1 \cup X_2 \cup \dots \cup X_n$, $f(X_i) = S^1$ for any $i = 1..n$.

Definition 2. Let be $F : \mathbb{R} \rightarrow \mathbb{R}$ is the lifting of a continuous map $f : S^1 \rightarrow S^1$ on \mathbb{R} , i.e. $f(\{x\}) = \{F(x)\}$. Then by the degree $\deg(f)$ of a map f we denote the number $F(x+1) - F(x)$, which is independent from the choice of the point x and the lifting F .

Corollary 2. The reachability problem for complete piecewise affine mappings with two intervals ⁵ is decidable.

Proof. The condition of a piecewise affine map with two intervals $f : S^1 \rightarrow S^1$ to be complete means that $S^1 = X_1 \cup X_2$ and $f(X_1) = f(X_2) = S^1$. Thus, if $X_1 = [0, \frac{m}{n}]$ and $X_2 = [\frac{m}{n}, 1)$, then $f(x) = a_1x + b_1$, where $a_1 = \pm \frac{n}{m}$, when $x \in X_1$, and $f(x) = a_2x + b_2$, where $a_2 = \pm \frac{n}{n-m}$ at $x \in X_2$, $m, n \in \mathbb{N}$, $\gcd(m, n) = 1$. It is clear that $n, m, n - m$ are relatively prime. So the conditions of Theorem 2 are satisfied. \square

4 PAM representation of β -expansions

Given a rational non-integer $\beta > 1$ and the number $x \in [0, 1]$. The target discounted-sum 0-1 problem [22, 11] is defined as follows: *Is there a sequence $w : \mathbb{N} \rightarrow \{0, 1\}$ of zeros and ones such that $x = \sum_{i=1}^{\infty} w(i) \frac{1}{\beta^i}$.*

For any $x \in S^1$, there exists β -expansion $w : \mathbb{N} \rightarrow \{0, 1, \dots, [\beta] - 1\}$ such that $x = \sum_{i=1}^{\infty} w(i) \frac{1}{\beta^i}$. If $w(i) \in \{0, 1\}$ we call it $(0, 1) - \beta$ -**expansion**. Therefore, when $\beta \leq 2$ the answer to the target discounted-sum problem is always

⁵ In particularly the continuous piecewise affine mapping of degree two

positive. Therefore, the only interesting case is when $\beta > 2$. We denote $D = \{0, 1, \dots, \lceil \beta \rceil - 1\}$. Then the minimal and maximal numbers, which are representable in the basis β with digits from the alphabet D , are $min = \sum_{i=1}^{\infty} 0 \frac{1}{\beta^i} = 0$ and $max = \sum_{i=1}^{\infty} (\lceil \beta \rceil - 1) \frac{1}{\beta^i} = \frac{\lceil \beta \rceil - 1}{\beta - 1}$. When $\beta > 2$ then max is always less than two. Let us denote by X_d the interval $[\frac{min+d}{\beta}, \frac{max+d}{\beta})$ for each $d \in D$. If β is not an integer number then two intervals X_d and X_{d+1} intersect. Also taking into account that $max < 2$, then the intervals X_d and X_{d+2} have no common points. Finally from the above construction we get the next lemma:

Lemma 3. *If $\beta > 2$ and β is rational/non-integer number: $X_d \cap X_{d+1} \neq \emptyset$, $d < \lceil \beta \rceil - 1$; $X_d \cap X_{d+2} = \emptyset$, $d < \lceil \beta \rceil - 2$; $[min, max) = \cup_{d \in A} X_d$.*

Proposition 1. *For any β -expansion there is a non-deterministic PAM where a symbolic dynamic of visited intervals (i.e. a sequence of symbols associated with intervals) from an initial point x_0 corresponds to its representation in base β .*

Proof. Let us define the piecewise affine mapping $f \subseteq [min, max) \times [min, max)$ as follows $f = \{(x, \beta x - d) | x \in X_d, d \in D\}$. It directly follows from this definition that $f(X_d) = [min, max)$. Let us consider an orbit $f^i(x) = x(i)$,




Fig. 1. A non-deterministic PAM for $\frac{5}{2}$ -expansion

$i \in \mathbb{Z}^+$. We say that $d_i \in D$ such that $x(i) \in X_{d_i}$. Then for any $n \in \mathbb{N}$ $min < |\beta^n x - d_1 \beta^{n-1} - d_2 \beta^{n-2} - \dots - d_n \beta^0| < max$, and in other form $\frac{min}{\beta^n} < |x - \sum_{i=1}^n (d_i \frac{1}{\beta^i})| < \frac{max}{\beta^n}$. So $|x - \sum_{i=1}^n (d_i \frac{1}{\beta^i})| \rightarrow 0$, $n \rightarrow \infty$, and therefore $x = \sum_{i=1}^{\infty} (d_i \frac{1}{\beta^i})$. Let us consider it in other direction. Let $x = \sum_{i=1}^{\infty} (d_i \frac{1}{\beta^i})$, then the sequence $\mathbf{x}(i)$, where $\mathbf{x}(0) = x$, $\mathbf{x}(i+1) = \beta \mathbf{x}(i) - d_i$, is the orbit of x in PAM f . Let us name the constructed map as the β -**expansion PAM**. \square

The nondeterministic β -expansion can be translated into deterministic maps corresponding to *greedy* and *lazy* expansions as follows:

Definition 3. *A function $f : [min, max) \rightarrow [min, max)$ is the greedy β -expansion PAM if the domain $[min, max)$ is divided on intervals X'_d , $d \in \{0, 1, \dots, \lceil \beta \rceil - 1\}$ such that $X'_{\lceil \beta \rceil - 1} = X_{\lceil \beta \rceil - 1}$, $X'_{d-1} = X_{d-1} - X_d$, $d \in \{1, 2, \dots, \lceil \beta \rceil - 1\}$ and $f(x) = \beta x - d$ iff $x \in X'_d$.*

Since $X_d = [\frac{\min+d}{\beta}, \frac{\max+d}{\beta})$ then $X'_d = [\frac{\min+d}{\beta}, \frac{\min+d+1}{\beta}) = [\frac{d}{\beta}, \frac{d+1}{\beta})$, $d \in \{0, 1, \dots, \lceil \beta \rceil - 2\}$ and the length of the interval X'_d is equal to $\frac{1}{\beta}$, $d < \lceil \beta \rceil - 1$.




Fig. 2. Deterministic greedy (on the left) and lazy (on the right) $\frac{5}{2}$ -expansion PAM

Definition 4. A function $f : [min, max) \rightarrow [min, max)$ is the lazy β -expansion PAM, if the domain $[min, max)$ is divided into intervals X''_d , $d \in \{0, 1, \dots, \lceil \beta \rceil - 1\}$, such that $X''_0 = X_0$, $X''_d = X_d - X_{d-1}$, $d \in \{1, 2, \dots, \lceil \beta \rceil - 1\}$, and $f(x) = \beta x - d$ iff $x \in X''_d$.

Proposition 2. Let f and g are greedy and lazy β -expansion PAM's respectively. f and g are (topologically) conjugate by the homeomorphism $h : h(x) = h^{-1}(x) = \max - x$, i.e. $f = h \circ g \circ h$.

Proof. The statement holds since $X'_d = \max - X''_{\lceil \beta \rceil - 1 - d}$, $d \in \{0, 1, \dots, \lceil \beta \rceil - 1\}$

We would like to highlight that the questions about reachability as well as representation of numbers in rational bases are tightly connected with questions about the density of orbits in PAMs. Moreover if the density of orbits are the same for all non-periodic points then it may be possible to have a wider application of p -adic techniques provided in the beginning of the paper. Let us formulate a hypothesis that goes along with our experimental simulations in PAMs:

Hypothesis 1 The orbit of any rational point in any expanding deterministic PAM is either finite or dense on the whole domain.

Lemma 4. Any $(0, 1) - \beta$ -expansion is greedy.

Proof. Let f be a β -expansion PAM. Assume that there is a point x and the orbit $\mathbf{x}(i)$, where $\mathbf{x}(0) = x$, $\mathbf{x}(i+1) = \beta \mathbf{x}(i) - d_i$ in the map f such that $d_i \in \{0, 1\}$ for all $i \in \mathbb{N}$ and the orbit does not correspond to the β greedy expansion of x .

The intersection of intervals X_0 and X_1 is an interval $X_{01} = [\frac{\min+1}{\beta}, \frac{\max}{\beta})$. Applying a map $y = \beta x$ to X_{01} we see that X_{01} is scaled into $[\min+1, \max) = [1, \frac{\lceil \beta \rceil - 1}{\beta - 1})$. The interval $[1, \frac{\lceil \beta \rceil - 1}{\beta - 1})$ does not have any common points with X_0 as the point 1 lies on the right side of the left border of the interval $X_2 = [\frac{\min+2}{\beta}, \frac{\max+2}{\beta})$ and by Lemma 3 $X_i \cap X_{i+2} = \emptyset$.

Note that when $x > \frac{1}{\beta-1}$ we have $\beta x - 1 > x$. Let us assume that for some i $\mathbf{x}(i) \in X_{01}$ and $\mathbf{x}(i+1) = \beta \mathbf{x}(i) - 0$, i.e. we did not followed a greedy expansion

and therefore $\mathbf{x}(i+1) \in [1, \frac{[\beta]-1}{\beta-1})$. Then $\mathbf{x}(i+2) = \beta\mathbf{x}(i+1) - 1 > \mathbf{x}(i+1)$ and $\mathbf{x}(i+2) \notin X_0$, etc. In this case starting from $\mathbf{x}(i+1)$ there is monotonically increasing sequence of orbital points in the interval X_1 . So points in such orbit should eventually leave the interval X_1 and reach X_d , where $d > 1$. This gives us a contradiction with the original assumption. \square

Corollary 3. *Since the greedy expansion can be expressed by a deterministic map then $(0, 1) - \beta$ -expansion is unique and greedy.*

Theorem 3. *If Hypothesis 1 holds then a non-periodic $(0, 1) - \beta$ -expansion does not exist.*

Proof. Any $(0, 1) - \beta$ -expansion can be constructed by expanding ⁶ deterministic greedy β -expansion PAM. If the orbit of a rational point in greedy β -expansion PAM is non-periodic, then by Hypothesis 1 it should be dense and therefore should intersect all intervals and cannot provide $(0, 1) - \beta$ -expansion. \square

Theorem 4. *If Hypothesis 1 holds then for any rational number its deterministic β -expansion is either eventually periodic or it contains all possible patterns (finite subsequences of digits) from $\{0, 1, \dots, [\beta] - 1\}$.*

Proof. The statement is obvious as Hypothesis 1 implies that the orbit is either periodic or it is dense and the dense orbit is visiting all intervals. \square

It looks that the point-to-interval problem is harder than the point-to-point reachability problem for the expanding PAMs, as for example Theorem 2 gives an algorithm for the point-to-point reachability problem in the β -expansion PAMs, but not for the point-to-interval reachability that is equivalent to the β -expansion problem.

Note that in the β -expansion PAMs all linear coefficients are the same, so the density of the orbit correspond to the density of the following sequence $\mathbf{x}(n) = f^n(x_0)$, where $f(x) = \{\beta x\}$. For example when $\beta = \frac{5}{2}$ and $x_0 = 1$ we get the sequence:

$$\{\frac{5}{2}\}, \{\frac{5}{2}\{\frac{5}{2}\}\}, \{\frac{5}{2}\{\frac{5}{2}\{\frac{5}{2}\}\}\}, \dots$$

The question about the distribution of a similar sequence $\{\frac{3}{2}\}, \{\frac{3^2}{2^2}\}, \{\frac{3^3}{2^3}\}, \dots$, where the integer part is removed once after taking a power of a fraction (for example $3/2$) is known as Mahler's $3/2$ problem, that is a long standing open problem in analytic number theory.

5 Density of orbits and its geometric interpretation

It is well known that $\mathbf{x}(n) = \{\alpha n\}$, where α is an irrational number, has an uniform distribution. Let us give some geometric interpretation of the orbit density. Consider the Cartesian plane with the y-axis x and and the x-axis y (just

⁶ I.e. with linear coefficients that are greater than 1

swapping their places). Now let us divide the set of lines $x = n$, $n \in \mathbb{N}$, by integer points on the segments of the unit length. The set of points (y, x) , where $m \leq y < m + 1$, $x = n$, i.e. the interval $[m, m + 1) \times n$ on the line $x = n$, will be denoted by $S_{m,n}$, $m \in \mathbb{Z}^+$, $n \in \mathbb{N}$. In other words, $S_{m,n} = (m + [0, 1)) \times n$. Let I be an interval such that $I \subseteq [0, 1)$ and by $I_{m,n}$ let us denote the set $(m + I) \times n$.




Fig. 3. Left: An example for two sets $S_{m,n}$ and $I_{m,n}$; Right: A dynamic interval $I(n)$.

Two points of the plane are defined to be equivalent if they belong to a same line passing through the origin. We call α as homogeneous coordinate of a point (y, x) if $y = \alpha x$. By $H(I)$ we denote the set of homogeneous coordinates of all points from $\bigcup_{m \in \mathbb{Z}^+, n \in \mathbb{N}} I_{m,n}$. The sequence $\mathbf{x}(n) = \{\alpha n\}$ is dense in $[0, 1)$ if and only if for any interval $I \subseteq [0, 1)$ there are m and n , such that the line $y = \alpha x$ intersects the set $I_{m,n}$. It is known that $[0, 1) - H(I) \subseteq \mathbb{Q}$ for any interval $I \subseteq [0, 1)$, i.e. for any irrational $\alpha > 0$ the line $y = \alpha x$ intersect the set $\bigcup_{m \in \mathbb{Z}^+, n \in \mathbb{N}} I_{m,n}$. Moreover in the case of irrational factors it is known that the frequency of occurrence of $\mathbf{x}(n) = \{\alpha n\}$ in the interval I is equal to its length.

In some sense the interval I , in the above example, can be named as *static* because it does not change in time n . However in order to study and describe previously mentioned problems such as the target discounted-sum problem, PAMs reachability problems, the Mahler's 3/2 problem we require the notion of “dynamic intervals”.

Let \mathbf{x} be a sequence of numbers from $[0, 1)$. What is the distribution of a sequence $\mathbf{x}'(n) = \{p^{\mathbf{k}(n)} \mathbf{x}(n)\}$, where $\mathbf{k} : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ is a non-decreasing sequence? For example, if $\mathbf{k}(n) = n - 1$ and the number $\mathbf{x}(n)$ has in the base p the following form $\mathbf{x}(n) = 0.a_{n1}a_{n2} \dots a_{nn}a_{n,n+1} \dots$, then $\mathbf{x}'(n) = 0.a_{nn}a_{n,n+1} \dots$.

Let us assume that $I \subseteq S^1$ and $\mathbf{k} : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ is a non-decreasing sequence, $p \in \mathbb{N}$. Now we define “dynamical intervals” as an evolving infinite sequence

$\mathbf{I}(1), \mathbf{I}(2), \mathbf{I}(3), \dots :$

$$\mathbf{I}(1) = I, \mathbf{I}(n) = \bigcup_{j=0}^{p^{\mathbf{k}(n)}-1} \frac{I+j}{p^{\mathbf{k}(n)}}.$$

By $F_{\mathbf{I}}^{\mathbf{x}}(n) = |\{i \in \mathbb{Z}^+ | i \leq n, \mathbf{x}(i) \in \mathbf{I}(i)\}|$ we denote a function representing a frequency of hitting dynamical interval \mathbf{I} by the sequence \mathbf{x} . In contrast to $F_I^{\mathbf{x}}(n)$ which only counts the number of hittings to a fixed interval I , our new function $F_{\mathbf{I}}^{\mathbf{x}}$ counts the number of hittings when both points and intervals are changing in time.

Proposition 3. *The following equation holds: $F_{\mathbf{I}}^{\mathbf{x}'}(n) = F_{\mathbf{I}}^{\mathbf{x}}(n)$.*

The phenomenon that significant digit distribution in real data are not accruing randomly known as Benford's Law. For example the sequence p^1, p^2, p^3, \dots satisfies Benford's Law, under the condition that $\log_{10} p$ is an irrational number, which is a consequence of the *Equidistribution theorem* (proved separately by Weyl, Sierpinski and Bohl). The Equidistribution theorem states that the sequence $\{\alpha\}, \{2\alpha\}, \{3\alpha\}, \dots$ is uniformly distributed on the circle \mathbb{R}/\mathbb{Z} , when α is an irrational number. It gives us the fact that each significant digit of numbers in (p^n) sequence will correspond to the interval \mathbb{R}/\mathbb{Z} and the length of the interval related to the frequency for each appearing digit.

However the question about the distribution of the sequence $\{(3/2)^n\}$ is different in the way that it is not about the distribution of the first digits of 3^n in base 2, i.e. not about the distribution of the sequence $\frac{3^n}{2^{\lceil n \log_2 3 \rceil}}$, but related to the sequence of digits after some shift of the number $\frac{3^n}{2^{\lceil n \log_2 3 \rceil}}$ corresponding to the multiplication by a power of 2.

So in the above notations the distribution of numbers in the sequence $\mathbf{x}'(n) = \{(3/2)^n\}$ corresponds to the $F_{\mathbf{I}}^{\mathbf{x}'}(n)$ for the logarithmic (Benford's law) distributed sequence $\mathbf{x}(n) = \frac{3^n}{2^{\lceil n \log_2 3 \rceil}}$, $p = 2$ and $\mathbf{k}(n) = \lceil n \log_2 3 \rceil - n$.

Now we will show that even if the sequence $\{\alpha\}, \{2\alpha\}, \{3\alpha\}, \dots$ is uniformly distributed on the circle \mathbb{R}/\mathbb{Z} , the irrationality of α is not enough to guarantee uniform distribution or even density of the sequence $\mathbf{x}'(n)$ on the circle corresponding to the linear shifts $\mathbf{k}(n) = n$.

Theorem 5. *Let us define $\alpha = \sum_{i=1}^{\infty} \frac{1}{2^{\Delta_i}}$ where $\Delta_1 = 1, \Delta_{i+1} = 2^{\Delta_i} + \Delta_i, i \geq 1$ (<http://oeis.org/A034797>). Then for all $n \in \mathbb{N} \cup \{0\}$ the sequence $\{2^n n \alpha\}$ is not dense in the interval $[0, 1]$ and $\{2^n n \alpha\} < \frac{1}{2}$.*

Proof. Let us consider a sequence Δ , which initial elements are $\Delta_1 = 1, \Delta_2 = 3, \Delta_3 = 11, \Delta_4 = 2059$ etc.

$$\begin{array}{r}
\Delta_1 \quad \Delta_2 \quad \Delta_3 \\
\mathbf{x}(0) = 0, 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \dots \\
\mathbf{x}(1) = 0, \boxed{1} \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \dots \\
\mathbf{x}(2) = 0, 0 \ \boxed{1} \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 0 \dots \\
\mathbf{x}(3) = 0, 0 \ 1 \ \boxed{1} \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 1 \ 0 \dots \\
\mathbf{x}(4) = 0, 1 \ 0 \ 0 \ \boxed{0} \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 0 \dots \\
\mathbf{x}(5) = 0, 1 \ 0 \ 1 \ 0 \ \boxed{0} \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 0 \dots \\
\mathbf{x}(6) = 0, 1 \ 1 \ 0 \ 0 \ 0 \ \boxed{0} \ 0 \ 0 \ 1 \ 1 \ 0 \ 0 \dots \\
\mathbf{x}(7) = 0, 1 \ 1 \ 1 \ 0 \ 0 \ 0 \ \boxed{0} \ 0 \ 1 \ 1 \ 1 \ 0 \dots \\
\mathbf{x}(8) = 0, 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ \boxed{1} \ 0 \ 0 \ 0 \ 0 \dots \\
\mathbf{x}(9) = 0, 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ \boxed{0} \ 0 \ 1 \ 0 \dots \\
\mathbf{x}(10) = 0, 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ \boxed{1} \ 0 \ 0 \dots \\
\mathbf{x}(11) = 0, 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ \boxed{1} \ 0 \dots \\
\\
\mathbf{x}'(0) = 0, 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \dots \\
\mathbf{x}'(1) = 0, 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \dots \\
\mathbf{x}'(2) = 0, 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 0 \dots \\
\mathbf{x}'(3) = 0, 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 1 \ 0 \dots \\
\mathbf{x}'(4) = 0, 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 0 \dots \\
\mathbf{x}'(5) = 0, 0 \ 0 \ 0 \ 1 \ 0 \ 1 \ 0 \dots \\
\mathbf{x}'(6) = 0, 0 \ 0 \ 1 \ 1 \ 0 \ 0 \dots \\
\mathbf{x}'(7) = 0, 0 \ 1 \ 1 \ 1 \ 0 \dots \\
\mathbf{x}'(8) = 0, 0 \ 0 \ 0 \ 0 \dots \\
\mathbf{x}'(9) = 0, 0 \ 1 \ 0 \dots \\
\mathbf{x}'(10) = 0, 0 \ 0 \dots \\
\mathbf{x}'(11) = 0, 0 \dots
\end{array}$$

Let us prove that when $0 \leq n \leq \Delta_i$ the inequality $\{2^n n \alpha\} < \frac{1}{2}$ if and only if $\{2^n n \sum_{j=1}^i \frac{1}{2^{\Delta_j}}\} < \frac{1}{2}$. The implication from left to right follows from $\{2^n n \sum_{j=1}^i \frac{1}{2^{\Delta_j}}\} < \{2^n n \alpha\}$. Let us show that it also holds in other direction. Assume that $\{2^n n \sum_{j=1}^i \frac{1}{2^{\Delta_j}}\} < \frac{1}{2}$ then $\{2^n n \sum_{j=1}^i \frac{1}{2^{\Delta_j}} + x\} < \frac{1}{2}$ for any $0 \leq x \leq \frac{1}{2^{\Delta_i+1-n}}$. In this case it is enough to show that $2^n n \alpha - 2^n n \sum_{j=1}^i \frac{1}{2^{\Delta_j}} < \frac{1}{2^{\Delta_i+1-n}}$ holds when $n \leq \Delta_i$. In fact we have that

$$\begin{aligned}
2^n n \alpha - 2^n n \sum_{j=1}^i \frac{1}{2^{\Delta_j}} &= 2^n n \sum_{j=i+1}^{\infty} \frac{1}{2^{\Delta_j}} < 2^n n \frac{1}{2^{\Delta_{i+1}-1}} = \\
&= \frac{1}{2^{\Delta_{i+1}-n-\log_2(n)-1}} \leq \frac{1}{2^{\Delta_{i+1}-\Delta_i-\log_2(\Delta_i)-1}} = \frac{1}{2^{2^{\Delta_i}-\log_2(\Delta_i)-1}}.
\end{aligned}$$

Finally we have $\frac{1}{2^{2^{\Delta_i}-\log_2(\Delta_i)-1}} < \frac{1}{2^{\Delta_i+1-n}}$ when $n \leq \Delta_i$ and $i > 1$.

Now for proving the theorem it is enough to show that for $0 \leq n \leq \Delta_i$ the inequality $\{2^n n \sum_{j=1}^i \frac{1}{2^{\Delta_j}}\} < \frac{1}{2}$ holds. When $i = 1$ the statement is trivial. Let

us assume that it holds for $i = k - 1$. We show now that it holds for $i = k$, i.e. we show that for $\Delta_{k-1} \leq n \leq \Delta_k$ inequality $\{2^n n \sum_{j=1}^k \frac{1}{2^{\Delta_j}}\} < \frac{1}{2}$ holds or taking into account $\Delta_{k-1} \leq n$ also we have inequality $\{2^n n \frac{1}{2^{\Delta_k}}\} < \frac{1}{2}$.

For $\Delta_{k-1} \leq n \leq \Delta_k - \Delta_{k-1} - 1$ we get $2^n n \frac{1}{2^{\Delta_k}} \leq 2^{\Delta_k - \Delta_{k-1} - 1} (\Delta_k - \Delta_{k-1} - 1) \frac{1}{2^{\Delta_k}}$. Since $\Delta_k = 2^{\Delta_{k-1}} + \Delta_{k-1}$, then $2^{\Delta_k - \Delta_{k-1} - 1} (\Delta_k - \Delta_{k-1} - 1) \frac{1}{2^{\Delta_k}} = (2^{\Delta_{k-1}} - 1) \frac{1}{2^{\Delta_{k-1} + 1}} = \frac{1}{2} - \frac{1}{2^{\Delta_{k-1} + 1}} < \frac{1}{2}$.

Consider the case when $\Delta_k - \Delta_{k-1} \leq n \leq \Delta_k$, i.e. $2^{\Delta_{k-1}} \leq n \leq 2^{\Delta_{k-1}} + \Delta_{k-1}$ and define $m = n - 2^{\Delta_{k-1}}$. We have that $0 \leq m \leq \Delta_{k-1}$ and $2^n n \frac{1}{2^{\Delta_k}} = 2^{m+2^{\Delta_{k-1}}} (m + 2^{\Delta_{k-1}}) \frac{1}{2^{2^{\Delta_{k-1}} + \Delta_{k-1}}} = 2^m (m + 2^{\Delta_{k-1}}) \frac{1}{2^{\Delta_{k-1}}} = 2^m m \frac{1}{2^{\Delta_{k-1}}} + 2^m$. Therefore $\{2^n n \frac{1}{2^{\Delta_k}}\} = \{2^m m \frac{1}{2^{\Delta_{k-1}}}\}$, where $0 \leq m \leq \Delta_{k-1}$. Now by the induction we have that $\{2^m m \frac{1}{2^{\Delta_{k-1}}}\} < \frac{1}{2}$.

While the question about the distributions for PAM orbits remains open we have unexpectedly shown that in a very similar system, operating with irrational numbers, the uniform distribution of original orbits in maps may not remain uniform or even dense when taking the fractional part after regular shifts. This makes the questions about PAMs even more “mysterious” as it is not clear whether such property may hold for a sequence of points generated by PAMs, β -expansion and Mahler’s problem.

References

1. A. Aswani and C. Tomlin (2007), Reachability algorithm for biological piecewise-affine hybrid systems, HSCC,2007,633-636.
2. E. Asarin, V. Mysore, A. Pnueli, G. Schneider. Low dimensional hybrid systems – decidable, undecidable, don’t know. Inf. Comput. 211: 138-159 (2012)
3. E.Asarin and G.Schneider, Widening the boundary between decidable and undecidable hybrid systems. CONCUR, 2002,193-208 .
4. E.Asarin, O. Maler, A. Pnueli. Reachability analysis of dynamical systems having piecewise-constant derivatives. Theoretical Computer Science, 138, 1995, 35 - 66.
5. Paul C. Bell , Shang Chen, Lisa Jackson: Reachability and Mortality Problems for Restricted Hierarchical Piecewise Constant Derivatives. RP 2014: 32-44
6. P. Bell, I. Potapov. On undecidability bounds for matrix decision problems Theoretical Computer Science 391 (1), 3-13, 2008.
7. Amir M. Ben-Amram: Mortality of iterated piecewise affine functions over the integers: Decidability and complexity. Computability 4(1): 19-56 (2015)
8. M. Blank and L. Bunimovich. Switched flow systems: pseudo billiard dynamics. Dynamical Systems. V.19, No.4 (2004) 359-370
9. V. Blondel, O. Bournez, P. Koiran, C. Papadimitriou and J. Tsitsiklis. Deciding stability and mortality of piecewise affine dynamical systems. Theoretical Computer Sci. 255: (1-2), 687-696, 2001.
10. V. Blondel, O. Bournez, P. Koiran, J. Tsitsiklis: The Stability of Saturated Linear Dynamical Systems Is Undecidable. J. Comput. Syst. Sci. 62(3): 442-462 (2001)
11. Udi Boker and Thomas A. Henzinger and Jan Otop. The Target Discounted-Sum Problem. LICS 2015: 750-761 (2015)

12. Hugo Bazille, Olivier Bournez, Walid Gomaa, Amaury Pouly: On The Complexity of Bounded Time Reachability for Piecewise Affine Systems. RP 2014: 20-31
13. M. Dellnitz, G. Froyland, and S. Sertl, On the isolated spectrum of the Perron-Frobenius operator. *Nonlinearity*, vol. 13, no. 4, pp. 1171-1188, July 2000.
14. P. Koiran, M. Cosnard, and M. Garzon. Computability with low-dimensional dynamical systems. *Theoretical Computer Science*, 132:113-128, 1994
15. P. Koiran: The topological entropy of iterated piecewise affine maps is uncomputable. *Discrete Mathematics and Theor. Computer Science* 4(2): 351-356 (2001)
16. V. Blondel, J. Tsitsiklis. A survey of computational complexity results in systems and control. *Automatica*, 36, 2000, 1249-1274.
17. O. Kurgansky, I. Potapov, F. Sancho-Caparrini: Reachability Problems in Low-Dimensional Iterative Maps. *Int. J. Found. Comput. Sci.* 19(4): 935-951 (2008)
18. O. Kurgansky, I. Potapov. Computation in One-Dimensional Piecewise Maps and Planar Pseudo-Billiard Systems. LNCS, v.3699, 2005, 169-175.
19. A. Renyi. Representations for real numbers and their ergodic properties. *Acta Mathematica Academiae Scientiarum Hungarica*, 8(3-4):477-493, 1957.
20. G. Setti, G. Mazzini, R. Rovatti, and S. Callegari, Statistical modeling of discrete-time chaotic processes: basic finite dimensional tools and applications, *Proceedings of the IEEE*, vol. 90, no. 5, May 2002, pp. 662-690.
21. N. Sidorov. Almost Every Number Has a Continuum of β -Expansions. *The American Mathematical Monthly* 110(9): 838-842 (2003)
22. Mickael Randour, Jean-Francois Raskin, Ocan Sankur. Percentile Queries in Multi-Dimensional Markov Decision Processes. Dagstuhl seminar "Non-zero-sum games and control", 5.02.2015.
23. J. Ouaknine, J. Sousa Pinto, J. Worrell: On Termination of Integer Linear Loops. SODA 2015: 957-969