

OPERADS OF FINITE POSETS

FRÉDÉRIC FAUVET, LOÏC FOISSY, AND DOMINIQUE MANCHON

ABSTRACT. We describe four natural operad structures on the vector space generated by isomorphism classes of finite posets. The three last ones are set-theoretical and can be seen as a simplified version of the first, the same way the NAP operad behaves with respect to the pre-Lie operad. Moreover the two first ones are isomorphic.

Keywords: partial orders, finite posets, Hopf algebras, posets, operads

Math. subject classification: 05E05, 06A11, 16T30.

CONTENTS

1.	Introduction	1
2.	A first operad structure on finite posets	3
2.1.	Reminder on the set operad of sets	3
2.2.	Quotient posets	3
2.3.	The first poset operad	4
3.	The set triple operad of finite connected posets	4
3.1.	Saturation of relations	5
3.2.	The first set operad structure	5
3.3.	Two more set operad structures	6
3.4.	Proof of Theorem 2	8
3.5.	The triple suboperad generated by the connected poset with two elements	9
4.	Compatibilities for the operadic products	10
5.	Algebraic structures associated to these operads	11
5.1.	Products	11
5.2.	Coproducts	13
6.	Suboperads generated in degree 2	14
6.1.	WN Posets	14
6.2.	Compositions \blacktriangle and \blacktriangledown	17
6.3.	Comparison of WN posets and ∇ -compatible posets	19
	References	19

1. INTRODUCTION

A finite poset is a finite set E endowed with a partial order \leq . The Hasse diagram of (E, \leq) is obtained by representing any element of E by a vertex, and by drawing a directed edge from e to e' if and only if e' sits directly above e , i.e. $e < e'$ and, for any $e'' \in E$ such that $e \leq e'' \leq e'$, one has $e'' = e$ or $e'' = e'$. A poset is *connected* if its Hasse diagram is connected. Here are the isomorphism classes of connected posets up to four vertices, represented by their Hasse diagrams:



Let \leq and \leqslant be two partial orders on the same finite set E . We say that \leqslant is finer than \leq if $x \leqslant y \Rightarrow x \leq y$ for any $x, y \in E$. We also say that the poset (E, \leqslant) is finer than the poset (E, \leq) and we write $(E, \leqslant) \preceq (E, \leq)$. The finest partial order on E is the trivial one, for which any $x \in E$ is only comparable with itself.

It is well-known that the linear span \mathcal{H} (over a field \mathbf{k}) of the isomorphism classes of posets is a commutative incidence Hopf algebra: see [10, Paragraph 16], taking for \mathcal{F} the family of all finite posets with the notations therein. The product is given by the disjoint union, and the coproduct is given by:

$$(1) \quad \Delta_* P = \sum_{P_1 \sqcup P_2 = P, P_1 < P_2} P_1 \otimes P_2,$$

where the sum runs over all *admissible cuts* of the poset P , i.e. partitions of P into two (possibly empty) subposets P_1 and P_2 such that $P_1 < P_2$, which means that for any $x \in P_1$ and $y \in P_2$, we have $x \not\leq y$. For algebraic structures on finite posets, see also [8].

The linear species of finite posets \mathbb{P} is defined as follows: to any finite set A one associates the vector space \mathbb{P}_A freely generated by all poset structures on A . The species structure is obviously defined by relabelling. The natural internal involution is defined as follows: for any finite poset $\mathcal{A} = (A, \leq)$ we set $\overline{\mathcal{A}} := (A, \geq)$.

We exhibit in this work several (precisely four) natural operad structures on the linear species \mathbb{P} of posets. Let us briefly describe them by their partial compositions [6, Paragraph 5.3.7]: the first one, denoted by a circle \circ , is obtained by inserting one poset into another one at some vertex summing up over all possibilities. The three others, denoted by a black circle \bullet and two black triangles \blacktriangledown and \blacktriangle , are obtained by retaining three different "optimal" insertions, in a way to be precised. The natural involution on posets preserves the set operad \bullet , and exchanges \blacktriangledown and \blacktriangle .

It happens that the operads (\mathbb{P}, \circ) and (\mathbb{P}, \bullet) are isomorphic: there is a species automorphism Φ of \mathbb{P} such that for any pair $(\mathcal{A}, \mathcal{B})$ of posets and any $a \in \mathcal{A}$, we have:

$$(2) \quad \Phi(\mathcal{A} \bullet_a \mathcal{B}) = \Phi(\mathcal{A}) \circ_a \Phi(\mathcal{B}).$$

The species automorphism Φ is simply defined by:

$$\begin{aligned} \Phi_A : \mathbb{P}_A &\longrightarrow \mathbb{P}_A \\ \mathcal{A} &\longmapsto \sum_{\mathcal{A}' \preceq \mathcal{A}} \mathcal{A}'. \end{aligned}$$

We give in Section 4 some compatibility relations between the three set-theoretic operad structures \bullet , \blacktriangledown and \blacktriangle . Section 5 is devoted to the various binary products on free algebras given by the elements of arity two in the operads under investigation. Three associative products and two non-associative permutative (NAP) products are obtained this way. Coproducts obtained by dualization of the aforementioned associative products are also investigated.

We give in Section 6 a presentation of the suboperad \mathbb{WNP} of (\mathbb{P}, \bullet) generated by the elements of arity two. The posets thus obtained are exactly the WN posets, i.e. the posets which do not contain \mathbb{N} as a subposet [3]. Finally, we give a presentation of the suboperad of ∇ -compatible posets, i.e. the suboperad \mathbb{CP}_{∇} of $(\mathbb{P}, \blacktriangledown)$ generated by the elements of arity two, as well as a set-theoretical species isomorphism $\theta : \mathbb{WNP} \rightarrow \mathbb{CP}_{\nabla}$ from WN posets onto ∇ -compatible posets.

2. A FIRST OPERAD STRUCTURE ON FINITE POSETS

2.1. **Reminder on the set operad of sets.** We refer the reader to [9] for an account of species as well as operads in the species formalism, see also [1]. Recall that a set species (resp. a linear species) is a contravariant¹ functor from the category of finite sets together with bijections, into the category of sets (resp. vector spaces) together with maps (resp. linear maps). The collection of finite sets is endowed with a tautological species structure, given by $A \mapsto A$ for any finite set, and $\varphi \mapsto \varphi^{-1}$ for any bijection $\varphi : A \rightarrow B$.

Let A and B be two finite sets, and let $a \in A$. We introduce the finite set:

$$(3) \quad A \sqcup_a B := A \sqcup B \setminus \{a\}.$$

It is obvious that the partial compositions \sqcup_a above are functorial, and endow the species of finite sets with an operad structure. Indeed, let $a, a' \in A$ with $a \neq a'$, let $b \in B$, and let C be a third finite set. The parallel associativity axiom reads:

$$(4) \quad (A \sqcup_a B) \sqcup_{a'} C = (A \sqcup_{a'} C) \sqcup_a B = A \sqcup B \sqcup C \setminus \{a, a'\}$$

and the nested associativity axiom reads:

$$(5) \quad (A \sqcup_a B) \sqcup_b C = A \sqcup_a (B \sqcup_b C) = A \sqcup B \sqcup C \setminus \{a, b\}.$$

2.2. **Quotient posets.** Let P be a set and $B \subseteq P$ be a subset. We denote by P/B the set $P \setminus B \sqcup \{B\}$. Let $\mathcal{P} = (P, \leq)$ be a poset, and let B a nonempty subset of P . We define a binary relation \leq on P/B as follows: $x \leq y$ if and only if:

- either $x = \{B\}$ and there exists $x' \in B$ such that $x' \leq y$,
- or $y = \{B\}$ and there exists $y' \in B$ such that $x \leq y'$,
- or $x, y \in P \setminus B$ and $x \leq y$,
- or $x, y \in P \setminus B$ and there exist $b, b' \in B$ such that $x \leq b$ and $b' \leq y$.

Recall that the interval $[a, b]$ in a poset $\mathcal{P} = (P, \leq)$ (with $a, b \in P$) is defined by:

$$[a, b] := \{x \in P, a \leq x \leq b\}.$$

We say that B is *convex* if any interval $[x, y]$ of \mathcal{P} with $x, y \in B$ is included in B .

Proposition 1. *Let $\mathcal{P} = (P, \leq)$ be a poset, let B be a subset of P . Then the binary relation \leq defined above is a partial order on P/B if and only if B is convex.*

Proof. If B is not convex, there exist $x, y, z \in P$ with $x, y \in B$, $z \notin B$ and $x \leq z \leq y$. By definition of \leq we have then $\{B\} \leq z \leq \{B\}$, hence \leq is not antisymmetric.

Now suppose that B is convex. Let $x, y, z \in P/B$ with $x \leq z$ and $z \leq y$. If they are not distinct we immediately get $x \leq y$. If $x = \{B\}$, there exists $x' \in B$ with $x' \leq z$. Two subcases can occur: If $z \leq y$ then $x' \leq y$, and then $x = \{B\} \leq y$. If there exist $b, b' \in B$ such that $z \leq b$ and $b' \leq y$, then $x' \leq b$ and $b' \leq y$, hence $x = \{B\} \leq y$ again. The cases $z = \{B\}$ and $y = \{B\}$ are treated similarly. The last case, when the three elements x, z, y are different from $\{B\}$, divides itself into four subcases:

- if $x \leq z$ and $z \leq y$, then $x \leq y$ hence $x \leq y$.
- if there exist $b, b' \in B$ such that $x \leq b$, $b' \leq z$ and $z \leq y$, then $x \leq b$ and $b' \leq y$, hence $x \leq y$.
- if there exist $b'', b''' \in B$ such that $x \leq z$, $z \leq b''$ and $b''' \leq y$, then $x \leq b''$ and $b''' \leq y$, hence $x \leq y$.

¹in order to get right actions of permutation groups. It is purely a matter of convention: [1] and [9] prefer the covariant definition.

- if there exist $b, b', b'', b''' \in B$ such that $x \leq b, b' \leq z, z \leq b''$ and $b''' \leq y$, then $z \in B$ by convexity, which contradicts the hypothesis: this last subcase cannot occur.

This proves the transitivity of the binary relation \leq . Now suppose $x \leq y$ and $y \leq x$. If $x = \{B\}$, and $y \neq x$, there are $b, b' \in B$ such that $b \leq y$ and $y \leq b'$. Hence $y \in B$ by convexity, which is a contradiction. The case $y = \{B\}$ is treated similarly. If both x and y are different from $\{B\}$, four subcases can occur:

- if $x \leq y$ and $y \leq x$, then $x = y$.
- If there exist $b, b' \in B$ with $x \leq b, b' \leq y$ and $y \leq x$, then $b' \leq y \leq x \leq b$, hence $x, y \in B$ by convexity, which is impossible.
- If there exist $b'', b''' \in B$ with $y \leq b'', b''' \leq x$ and $x \leq y$, then $b''' \leq x \leq y \leq b''$, hence $x, y \in B$ by convexity, which is impossible.
- If there exist $b, b', b'', b''' \in B$ such that $x \leq b, b' \leq y$ and $y \leq b'', b''' \leq x$, then $y \in B$ by convexity, which is impossible.

Hence \leq is a partial order. \square

For any poset $\mathcal{P} = (P, \leq)$ and any convex subset of P , we will denote by \mathcal{P}/B the poset $(P/B, \leq)$ thus obtained.

2.3. The first poset operad. Let $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$ be two finite posets, and let $a \in A$. We denote by $\Omega(\mathcal{A}, a, \mathcal{B})$ the set of all partial orders \leq on $A \sqcup_a B$ such that:

- (1) $\leq|_B = \leq_B$,
- (2) B is convex in $(A \sqcup_a B, \leq)$,
- (3) $(A \sqcup_a B, \leq)/B \sim \mathcal{A}$, where the set $(A \sqcup_a B)/B = (A \setminus \{a\}) \sqcup \{B\}$ is naturally identified with A by sending B on a .

The partial compositions of posets, which we denote by \circ_a , are then defined by:

$$(6) \quad \mathcal{A} \circ_a \mathcal{B} := \sum_{\leq \in \Omega(\mathcal{A}, a, \mathcal{B})} (A \sqcup_a B, \leq).$$

For example, we have

$$\mathbf{1}_a^b \circ_a \mathbf{1}_1^2 = {}^b \mathbf{V}_1^2 + \mathbf{1}_1^b$$

and

$$\mathbf{1}_a^c \circ_b \mathbf{1}_1^2 = \mathbf{1}_a^c + {}^c \mathbf{Y}_1^2 + {}^c \mathbf{A}_{1+}^2 + {}^c \mathbf{M}_1^2 + {}^c \mathbf{M}_a^2.$$

Note that a and c are not comparable in the last term, but become comparable (namely $a \leq c$) when the subposet $\mathbf{1}_1^2$ is shrunk.

Theorem 2. *The partial compositions (6) define an operad structure on the species of finite posets.*

We postpone the proof to Paragraph 3.4 in the next section.

3. THE SET TRIPLE OPERAD OF FINITE CONNECTED POSETS

We will define set-theoretical partial compositions by retaining one privileged term in the sum (6) defining the partial compositions \circ_a . It turns out that three different procedures are available leading to three families of partial compositions \bullet_a , \blacktriangledown_a and \blacktriangle_a , providing the species of finite posets with three distinct set operad structures.

For any two finite posets $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$, and $a \in A$, the partial order on $\mathcal{A} \bullet_a \mathcal{B}$ is obtained from \leq_A and \leq_B by saturation, and the operad axioms can be deduced from saturation by stages. We give an account of this general phenomenon in Paragraph 3.1 below

before looking at the partial compositions \bullet_a themselves.

The operad axioms for the family ∇_a , and similarly for \blacktriangle_a , are obtained by a direct check, which can also be used with very few modifications for \bullet_a .

3.1. Saturation of relations. Let E be a set. Let \sim be an equivalence relation on E , let \mathcal{R} be a binary relation on E/\sim and let \mathcal{S} be a binary relation on E . The class of $e \in E$ in E/\sim will be denoted by $[e]$. Let \mathcal{T} be the relation on E defined by $e\mathcal{T}e'$ if and only if, either $[e] \neq [e']$ and $[e]\mathcal{R}[e']$, or $[e] = [e']$ and $e\mathcal{S}e'$.

Proposition 3. *If the two binary relations \mathcal{R} and \mathcal{S} above are partial orders, then \mathcal{T} is also a partial order.*

Proof. Reflexivity is obvious. Now let $e, e', e'' \in E$ such that $e\mathcal{T}e'$ and $e'\mathcal{T}e''$. Several cases can occur:

- (1) If $[e] = [e'] = [e'']$, then $e\mathcal{S}e'$ and $e'\mathcal{S}e''$, hence $e\mathcal{S}e''$, which yields $e\mathcal{T}e''$.
- (2) If $[e] = [e'] \neq [e'']$, then $[e']\mathcal{R}[e'']$. Hence $[e]\mathcal{R}[e'']$, which yields $e\mathcal{T}e''$. Similarly for $[e] \neq [e'] = [e'']$.
- (3) If $[e] \neq [e']$ and $[e'] \neq [e'']$, then $[e]\mathcal{R}[e']$ and $[e']\mathcal{R}[e'']$, hence $[e]\mathcal{R}[e'']$. Moreover, $[e] \neq [e'']$: otherwise, we would have $[e] = [e'] = [e'']$, as \mathcal{R} is a partial order. This yields $e\mathcal{T}e''$.

This proves transitivity of \mathcal{T} . Finally, if $e\mathcal{T}e'$ and $e'\mathcal{T}e$ then:

- (1) If $[e] = [e']$, then $e\mathcal{S}e'$ and $e'\mathcal{S}e$, hence $e = e'$ by antisymmetry of \mathcal{S} .
- (2) If $[e] \neq [e']$, then $[e]\mathcal{R}[e']$ and $[e']\mathcal{R}[e]$, contradiction.

This proves antisymmetry of \mathcal{T} . □

Proposition 4 (Saturation by stages). *Let E be a set. Let \sim be an equivalence relation on E , let \approx be an equivalence relation on E finer than \sim , and let \simeq be the equivalence relation on E/\approx deduced from \sim . The quotient E/\sim is in bijection with $(E/\approx)/\simeq$. Let*

- \mathcal{R} be a binary relation on $E/\sim = (E/\approx)/\simeq$,
- \mathcal{S} be a binary relation on E/\approx ,
- \mathcal{T} be the relation on E/\approx obtained from \mathcal{R} and \mathcal{S} by saturation as explained above,
- \mathcal{S}' be a binary relation on E ,
- \mathcal{S}'' be the binary relation on E obtained from \mathcal{S} and \mathcal{S}' by saturation,
- \mathcal{T}' be the relation on E obtained from \mathcal{T} and \mathcal{S}' by saturation.

Then \mathcal{T}' is also obtained from \mathcal{R} and \mathcal{S}'' by saturation.

Proof. For any $e \in E$, we will denote by $[e]$ the class of e for the relation \approx , and by $[[e]]$ the class of e for \sim , which is the same as the class of $[e]$ for \simeq . We have $e\mathcal{T}'e'$ if and only if either $[e] = [e']$ and $e\mathcal{S}'e'$, or $[e] \neq [e']$ and $[e]\mathcal{T}[e']$. Using the definition of \mathcal{T} we have that $e\mathcal{T}'e'$ if and only if:

- either $[e] = [e']$ and $e\mathcal{S}'e'$,
- or $[e] \neq [e']$, $[[e]] = [[e']]$ and $[e]\mathcal{S}[e']$,
- or $[[e]] \neq [[e']]$ and $[[e]]\mathcal{R}[[e']]$.

Hence $e\mathcal{T}'e'$ if and only if:

- either $[[e]] = [[e']]$ and $e\mathcal{S}''e'$,
- or $[[e]] \neq [[e']]$ and $[[e]]\mathcal{R}[[e']]$,

which proves the claim. □

3.2. The first set operad structure. Let $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$ be two finite posets, and let $a \in A$. Then we define:

$$(7) \quad \mathcal{A} \bullet_a \mathcal{B} := (A \sqcup_a B, \leq),$$

where \leq is obtained from \leq_A on $A = A \sqcup_a B / \sim$ and \leq_B on $A \sqcup B$ by the saturation process described in Paragraph 3.1. Here the equivalence relation \sim is obtained by identifying all elements of B with a , and the partial order \leq_B is trivially extended to $A \sqcup_a B$, i.e. $x \leq y$ if and only if $x = y$ in case x or y does not belong to B . The relation \leq is a partial order by Proposition 3. To sum up, for any $x, y \in A \sqcup_a B$, $x \leq y$ if and only if:

- either $x, y \in B$ and $x \leq_B y$,
- or $x, y \in A \setminus \{a\}$ and $x \leq_A y$,
- or $x \in A \setminus \{a\}$, $y \in B$ and $x \leq_A a$,
- or $x \in B$, $y \in A \setminus \{a\}$ and $a \leq_A y$.

for example, we have:

$$\begin{array}{c} \begin{array}{c} \uparrow^c \\ \uparrow^b \\ \uparrow^a \end{array} \bullet_b \begin{array}{c} \uparrow^2 \\ \uparrow^1 \end{array} = \begin{array}{c} \uparrow^c \\ \uparrow^2 \\ \uparrow^1 \\ \uparrow^a \end{array}, \\ \begin{array}{c} \uparrow^c \\ \uparrow^b \\ \uparrow^a \end{array} \bullet_b \cdot 1 \cdot 2 = 1 \begin{array}{c} \uparrow^c \\ \uparrow^2 \\ \uparrow^1 \\ \uparrow^a \end{array}. \end{array}$$

Remark 5. Note that the partial order \leq defined above is an element of the set $\Omega(A, a, B)$ defined in Paragraph 2.3.

Theorem 6. The partial compositions \bullet_a endow the set species of finite posets with a structure of operad.

Proof. Partial composition of two posets is a single poset, and functoriality with respect to relabelling is obvious. It remains to check the two associativity axioms. The nested associativity axiom is directly derived from Proposition 4, with $E = A \sqcup_b B \sqcup_b C$. The equivalence relation \approx (resp. \sim) is obtained by shrinking C to $\{b\}$ (resp. $B \sqcup_b C$ to $\{a\}$). The set E / \approx is identified with $A \sqcup_a B$, and \simeq is obtained by shrinking B on $\{a\}$. It remains to apply Proposition 4 with the following dictionary:

- \mathcal{R} : partial order \leq_A on $A = E / \sim$.
- \mathcal{S} : partial order \leq_B trivially extended on $E / \approx = A \sqcup_a B$.
- \mathcal{T} : partial order of $\mathcal{A} \bullet_b \mathcal{B}$ on $E / \approx = A \sqcup_a B$.
- \mathcal{S}' : partial order \leq_C trivially extended to E ,
- \mathcal{S}'' : partial order of $\mathcal{B} \bullet_b \mathcal{C}$ trivially extended from $B \sqcup_b C$ to E .
- \mathcal{T}' : partial order of $(\mathcal{A} \bullet_a \mathcal{B}) \bullet_b \mathcal{C}$.

The proof of the parallel associativity is similar and left to the reader. \square

3.3. Two more set operad structures. We define two families of partial compositions \blacktriangle_a and \blacktriangledown_a as follows: for two finite posets $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$ and $a \in A$, the poset $\mathcal{A} \blacktriangle_a \mathcal{B}$ is the set $A \sqcup_a B$ together with the partial order \leq defined by $x \leq y$ if and only if:

- either $x, y \in B$ and $x \leq_B y$,
- or $x, y \in A \setminus \{a\}$ and $x \leq_A y$,
- or $x \in A \setminus \{a\}$, $y \in \max B$ and $x \leq_A a$,
- or $x \in B$, $y \in A \setminus \{a\}$ and $a \leq_A y$.

Similarly, the poset $\mathcal{A} \blacktriangledown_a \mathcal{B}$ is the set $A \sqcup_a B$ together with the partial order \leq defined by $x \leq y$ if and only if:

- either $x, y \in B$ and $x \leq_B y$,
- or $x, y \in A \setminus \{a\}$ and $x \leq_A y$,
- or $x \in A \setminus \{a\}$, $y \in B$ and $x \leq_A a$,
- or $x \in \min B$, $y \in A \setminus \{a\}$ and $a \leq_A y$.

Theorem 7. Both families of partial compositions \blacktriangle_a and \blacktriangledown_a endow the species of finite posets with a set operad structure.

Proof. Let us prove it for the family \blacktriangledown_a . The proof for \blacktriangle_a is entirely similar. Let $\mathcal{A} = (A, \leq_A)$, $\mathcal{B} = (B, \leq_B)$ and $\mathcal{C} = (C, \leq_C)$ be three finite posets, with $a \in A$ and $b \in B$. The poset $(\mathcal{A}\blacktriangledown_a\mathcal{B})\blacktriangledown_b\mathcal{C}$ is the set $A \sqcup_a B \sqcup_b C$ endowed with the partial order \leq defined by: $x \leq y$ if and only if:

- either $x, y \in C$ and $x \leq_C y$,
- or $x, y \in A \sqcup_a B \setminus \{b\}$ and $x \leq_{\mathcal{A}\blacktriangledown_a\mathcal{B}} y$,
- or $x \in A \sqcup_a B \setminus \{b\}$, $y \in C$ and $x \leq_{\mathcal{A}\blacktriangledown_a\mathcal{B}} b$,
- or $x \in \min \mathcal{C}$, $y \in A \sqcup_a B \setminus \{b\}$ and $b \leq_{\mathcal{A}\blacktriangledown_a\mathcal{B}} y$.

This in turn expands into: $x \leq y$ if and only if:

- (1) either $x, y \in C$ and $x \leq_C y$,
- (2) or $x, y \in B \setminus \{b\}$ and $x \leq_B y$,
- (3) or $x, y \in A \setminus \{a\}$ and $x \leq_A y$,
- (4) or $x \in A \setminus \{a\}$, $y \in B \setminus \{b\}$ and $x \leq_A a$,
- (5) or $x \in (\min \mathcal{B}) \setminus \{b\}$, $y \in A \setminus \{a\}$ and $a \leq_A y$,
- (6) or $x \in B \setminus \{b\}$, $y \in C$ and $x \leq_B b$,
- (7) or $x \in A \setminus \{a\}$, $y \in C$ and $x \leq_A a$,
- (8) or $x \in \min \mathcal{C}$, $y \in B \setminus \{b\}$ and $b \leq_B y$,
- (9) or $x \in \min \mathcal{C}$, $y \in A \setminus \{a\}$, $b \in \min \mathcal{B}$ and $a \leq_A y$.

On the other hand, the poset $\mathcal{A}\blacktriangledown_a(\mathcal{B}\blacktriangledown_b\mathcal{C})$ is the set $A \sqcup_a B \sqcup_b C$ endowed with the partial order \leq defined by: $x \leq y$ if and only if:

- either $x, y \in B \sqcup_b C$ and $x \leq_{\mathcal{B}\blacktriangledown_b\mathcal{C}} y$,
- or $x, y \in A \setminus \{a\}$ and $x \leq_A y$,
- or $x \in A \setminus \{a\}$, $y \in B \sqcup_b C$ and $x \leq_A a$,
- or $x \in \min \mathcal{B}\blacktriangledown_b\mathcal{C}$, $y \in A \setminus \{a\}$ and $a \leq_A y$.

This in turn expands into: $x \leq y$ if and only if:

- (1) either $x, y \in C$ and $x \leq_C y$,
- (2) or $x, y \in B \setminus \{b\}$ and $x \leq_B y$,
- (6) or $x \in B \setminus \{b\}$, $y \in C$ and $x \leq_B b$,
- (8) or $x \in \min \mathcal{C}$, $y \in B \setminus \{b\}$ and $b \leq_B y$,
- (3) or $x, y \in A \setminus \{a\}$ and $x \leq_A y$,
- (4) or $x \in A \setminus \{a\}$, $y \in B \setminus \{b\}$ and $x \leq_A a$,
- (7) or $x \in A \setminus \{a\}$, $y \in C$ and $x \leq_A a$,
- (9) or $x \in \min \mathcal{C}$, $b \in \min \mathcal{B}$, $y \in A \setminus \{a\}$ and $a \leq_a y$,
- (5) or $x \in (\min \mathcal{B}) \setminus \{b\}$, $y \in A \setminus \{a\}$ and $a \leq_A y$.

Hence the two partial orders \leq and \leq coincide of $A \sqcup_a B \sqcup_b C$, which proves the nested associativity axiom. Now let $a' \neq a$ be a second element of A . The poset $(\mathcal{A}\blacktriangledown_a\mathcal{B})\blacktriangledown_{a'}\mathcal{C}$ is the set $A \sqcup B \sqcup C \setminus \{a, a'\}$ endowed with the partial order \leq defined by: $x \leq y$ if and only if:

- either $x, y \in C$ and $x \leq_C y$,
- or $x, y \in A \sqcup_a B \setminus \{a'\}$ and $x \leq_{\mathcal{A}\blacktriangledown_a\mathcal{B}} y$,
- or $x \in A \sqcup_a B \setminus \{a'\}$, $y \in C$ and $x \leq_{\mathcal{A}\blacktriangledown_a\mathcal{B}} a'$,
- or $x \in \min \mathcal{C}$, $y \in A \sqcup_a B \setminus \{a'\}$ and $a' \leq_{\mathcal{A}\blacktriangledown_a\mathcal{B}} y$.

This in turn expands into: $x \leq y$ if and only if:

- either $x, y \in C$ and $x \leq_C y$,
- or $x, y \in B$ and $x \leq_B y$,

- or $x, y \in A \setminus \{a, a'\}$ and $x \leq_A y$,
- or $x \in A \setminus \{a, a'\}$, $y \in B$ and $x \leq_A a$,
- or $x \in \min \mathcal{B}$, $y \in A \setminus \{a, a'\}$ and $a \leq_A y$,
- or $x \in \min \mathcal{B}$, $y \in C$ and $a \leq_A a'$,
- or $x \in A \setminus \{a, a'\}$, $y \in C$ and $x \leq_A a'$,
- or $x \in \min \mathcal{C}$, $y \in B$ and $a' \leq_A a$,
- or $x \in \min \mathcal{C}$, $y \in A \setminus \{a, a'\}$ and $a' \leq_A y$.

Exchanging (B, a) with (C, a') leaves the nine conditions above globally unchanged, which proves the parallel associativity axiom. \square

Remark 8. The same proof can be written almost word for word for the operad (\mathbb{P}, \bullet) : it suffices to replace \blacktriangledown_a and \blacktriangledown_b by \bullet_a and \bullet_b respectively, and to suppress "min" everywhere in the proof. Hence we have two proofs for Theorem 6.

We can see the difference between the three structures in this example:

$$\begin{aligned} \begin{array}{c} c \\ \vdots \\ b \\ \vdots \\ a \end{array} \bullet_b \begin{array}{c} 2 \\ \vdots \\ 1 \end{array} &= \begin{array}{c} c \\ \vdots \\ 2 \\ \vdots \\ 1 \\ \vdots \\ a \end{array}, \\ \begin{array}{c} c \\ \vdots \\ b \\ \vdots \\ a \end{array} \blacktriangledown_b \begin{array}{c} 2 \\ \vdots \\ 1 \end{array} &= \begin{array}{c} c \\ \vee \\ 1 \\ \vdots \\ a \end{array}, \\ \begin{array}{c} c \\ \vdots \\ b \\ \vdots \\ a \end{array} \blacktriangle_b \begin{array}{c} 2 \\ \vdots \\ 1 \end{array} &= \begin{array}{c} c \\ \wedge \\ 1 \\ \vdots \\ a \end{array}. \end{aligned}$$

Remark 9. The family of partial compositions \blacktriangledown_a restricts itself to the rooted trees. The suboperad thus defined is nothing but the NAP operad [4]. The pre-Lie operad [2] is however *not* obtained from the family of partial compositions \circ_a , which don't restrict themselves to the rooted trees.

3.4. Proof of Theorem 2. Recall the species automorphism Φ of \mathbb{P} described in the Introduction, defined by:

$$\Phi(\mathcal{A}) = \sum_{\mathcal{A}' \preceq \mathcal{A}} \mathcal{A}'.$$

Let $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$ be two finite posets, and let $a \in A$. Let us introduce the two following sets: $\Omega'(\mathcal{A}, a, \mathcal{B})$ is the set of all partial orders \leq on $A \sqcup_a B$ such that

- B is convex in $(A \sqcup_a B, \leq)$,
- $(B, \leq|_B) \preceq \mathcal{B}$,
- $(A \sqcup_a B, \leq)/B \preceq \mathcal{A}$, where the set $(A \sqcup_a B)/B = (A \setminus \{a\}) \sqcup \{B\}$ is naturally identified with A by sending B on a ,

and $\Omega''(\mathcal{A}, a, \mathcal{B})$ is the set of all partial orders \leq on $A \sqcup_a B$ such that

$$(A \sqcup_a B, \leq) \preceq \mathcal{A} \bullet_a \mathcal{B}.$$

We obviously have:

$$(8) \quad \Phi(\mathcal{A} \bullet_a \mathcal{B}) = \sum_{\leq \in \Omega''(\mathcal{A}, a, \mathcal{B})} (A \sqcup_a B, \leq)$$

and

$$(9) \quad \Phi(\mathcal{A}) \circ_a \Phi(\mathcal{B}) = \sum_{\leq \in \Omega'(\mathcal{A}, a, \mathcal{B})} (A \sqcup_a B, \leq).$$

To prove Theorem 2, it is then enough to prove the equality $\Omega'(\mathcal{A}, a, \mathcal{B}) = \Omega''(\mathcal{A}, a, \mathcal{B})$. Let us first prove the inclusion $\Omega'(\mathcal{A}, a, \mathcal{B}) \subset \Omega''(\mathcal{A}, a, \mathcal{B})$: for any partial order $\leq \in \Omega'(\mathcal{A}, a, \mathcal{B})$ and x, y such that $x \leq y$, we have four cases to look at:

- if $x, y \in B$ then $x \leq|_B y$, hence $x \leq_B y$, hence $x \leq_{\mathcal{A} \bullet_a \mathcal{B}} y$.
- If $x \in A \setminus \{a\}$ and $y \in B$, then $x \leq_A a$, hence $x \leq_{\mathcal{A} \bullet_a \mathcal{B}} y$.
- If $x \in B$ and $y \in A \setminus \{a\}$, then $a \leq_A y$, hence $x \leq_{\mathcal{A} \bullet_a \mathcal{B}} y$.
- If $x, y \in A \setminus \{a\}$, then $x \leq_A y$, hence $x \leq_{\mathcal{A} \bullet_a \mathcal{B}} y$.

To sum up, for any partial order $\leq \in \Omega'(\mathcal{A}, a, \mathcal{B})$ and for any $x, y \in A \sqcup_a B$, $x \leq y \Rightarrow x \leq_{\mathcal{A} \bullet_a \mathcal{B}} y$, which shows the inclusion.

Let us now prove the reverse inclusion $\Omega''(\mathcal{A}, a, \mathcal{B}) \subset \Omega'(\mathcal{A}, a, \mathcal{B})$: first of all, for any partial order $\leq \in \Omega''(\mathcal{A}, a, \mathcal{B})$ and for any $x, z, y \in A \sqcup_a B$ with $x, y \in B$ and $x \leq z \leq y$, we have $x \leq_{\mathcal{A} \bullet_a \mathcal{B}} z \leq_{\mathcal{A} \bullet_a \mathcal{B}} y$, hence $z \in B$ by convexity of B in $\mathcal{A} \bullet_a \mathcal{B}$. Hence B is convex in $(A \sqcup_a B, \leq)$. Now for any $x, y \in B$ we have $x \leq y \Rightarrow x \leq_{\mathcal{A} \bullet_a \mathcal{B}} y \Rightarrow x \leq_B y$. Now let us denote by \leq the partial order on $A \sim (A \sqcup_a B)/B$ corresponding to the poset $(A \sqcup_a B, \leq)/B$. We just have to prove that for any $x, y \in A$, $x \leq y \Rightarrow x \leq_A y$. Four cases can occur:

- (1) if $x, y \in A \setminus \{a\}$, two subcases occur:
 - if $x \leq_A y$ then $x \leq_{\mathcal{A} \bullet_a \mathcal{B}} y$, then $x \leq_A y$.
 - If there exist $b, b' \in B$ such that $x \leq b$ and $b' \leq y$, then $x \leq_{\mathcal{A} \bullet_a \mathcal{B}} b$ and $b' \leq_{\mathcal{A} \bullet_a \mathcal{B}} y$, hence $x \leq_A a$ and $a \leq_A y$, which yields $x \leq_A y$.
- (2) If $x \in A \setminus \{a\}$ and $y = a$, there exists $b \in B$ such that $x \leq b$, hence $x \leq_{\mathcal{A} \bullet_a \mathcal{B}} b$, hence $x \leq_A y$.
- (3) The case $x = a$ and $y \in A \setminus \{a\}$ is treated similarly.
- (4) The case $x = y = a$ is trivial.

This proves the equality $\Omega'(\mathcal{A}, a, \mathcal{B}) = \Omega''(\mathcal{A}, a, \mathcal{B})$, which in turn proves Theorem 2.

Corollary 10. *The partial compositions \circ endow the species \mathbb{P} with a linear operad structure, isomorphic to (\mathbb{P}, \bullet) .*

3.5. The triple suboperad generated by the connected poset with two elements. It is possible to generate all connected posets up to four vertices with the poset \downarrow and the three families of partial compositions \bullet , \blacktriangledown and \blacktriangle . Indeed,

$$\begin{array}{l}
 \begin{array}{ccc}
 \begin{array}{c} c \\ \downarrow \\ b \\ \downarrow \\ a \end{array} = \begin{array}{c} 1 \\ \downarrow \\ a \end{array} \bullet_1 \begin{array}{c} c \\ \downarrow \\ b \end{array}, & \begin{array}{c} b \\ \blacktriangledown \\ a \end{array}^c = \begin{array}{c} b \\ \downarrow \\ 1 \\ \downarrow \\ a \end{array} \blacktriangledown_1 \begin{array}{c} c \\ \downarrow \\ a \end{array}, & \begin{array}{c} a \\ \blacktriangle \\ c \end{array}^b = \begin{array}{c} 1 \\ \downarrow \\ b \end{array} \blacktriangle_1 \begin{array}{c} a \\ \downarrow \\ c \end{array}, \\
 \begin{array}{c} d \\ \downarrow \\ c \\ \downarrow \\ a \end{array} = \begin{array}{c} d \\ \downarrow \\ 1 \\ \downarrow \\ a \end{array} \bullet_1 \begin{array}{c} b \\ \downarrow \\ a \end{array}, & \begin{array}{c} c \\ \blacktriangledown \\ a \end{array}^d = \begin{array}{c} c \\ \downarrow \\ 1 \\ \downarrow \\ a \end{array} \blacktriangledown_1 \begin{array}{c} d \\ \downarrow \\ b \end{array}, & \begin{array}{c} c \\ \blacktriangledown \\ a \end{array}^d = \begin{array}{c} c \\ \downarrow \\ 1 \\ \downarrow \\ b \end{array} \blacktriangledown_1 \begin{array}{c} d \\ \downarrow \\ a \end{array}, \\
 \begin{array}{c} c \\ \downarrow \\ b \\ \downarrow \\ a \end{array}^d = \begin{array}{c} b \\ \downarrow \\ 1 \\ \downarrow \\ a \end{array} \blacktriangledown_1 \begin{array}{c} c \\ \downarrow \\ d \end{array}, & \begin{array}{c} d \\ \blacktriangle \\ b \end{array}^c = \begin{array}{c} d \\ \downarrow \\ 1 \\ \downarrow \\ a \end{array} \blacktriangle_1 \begin{array}{c} c \\ \downarrow \\ b \end{array}, & \begin{array}{c} d \\ \blacktriangle \\ a \end{array}^c = \begin{array}{c} 1 \\ \downarrow \\ a \end{array} \blacktriangle_1 \begin{array}{c} d \\ \downarrow \\ c \end{array}, \\
 \begin{array}{c} d \\ \blacktriangle \\ c \end{array}^b = \begin{array}{c} a \\ \downarrow \\ 1 \\ \downarrow \\ b \end{array} \blacktriangle_1 \begin{array}{c} d \\ \downarrow \\ c \end{array}, & \begin{array}{c} b \\ \blacktriangle \\ d \end{array}^c = \begin{array}{c} b \\ \downarrow \\ a \end{array} \blacktriangle_1 \begin{array}{c} 1 \\ \downarrow \\ d \end{array}, & \begin{array}{c} b \\ \blacktriangle \\ a \end{array}^c = \begin{array}{c} 1 \\ \downarrow \\ 1 \\ \downarrow \\ a \end{array} \blacktriangle_1 \begin{array}{c} c \\ \downarrow \\ d \end{array}, \\
 \begin{array}{c} d \\ \blacktriangle \\ a \end{array}^c = \begin{array}{c} 1 \\ \downarrow \\ a \end{array} \bullet_1 \begin{array}{c} c \\ \downarrow \\ b \end{array} \blacktriangle_1^b \begin{array}{c} d \\ \downarrow \\ a \end{array}.
 \end{array}
 \end{array}$$

One cannot generate all finite posets with the poset \downarrow and the three families of partial compositions \bullet , \blacktriangledown and \blacktriangle . For example, \blacktriangleright cannot be reached this way.

4. COMPATIBILITIES FOR THE OPERADIC PRODUCTS

Proposition 11. *Let $\mathcal{A} = (A, \leq_A)$, $\mathcal{B} = (B, \leq_B)$ and $\mathcal{C} = (C, \leq_C)$ be three finite posets, and let $a, b \in A$, distinct. Then:*

$$\begin{aligned} (\mathcal{A} \blacktriangle_a \mathcal{B}) \bullet_b \mathcal{C} &= (\mathcal{A} \bullet_b \mathcal{C}) \blacktriangle_a \mathcal{B}, \\ (\mathcal{A} \blacktriangledown_a \mathcal{B}) \bullet_b \mathcal{C} &= (\mathcal{A} \bullet_b \mathcal{C}) \blacktriangledown_a \mathcal{B}, \\ (\mathcal{A} \blacktriangledown_a \mathcal{B}) \blacktriangle_b \mathcal{C} &= (\mathcal{A} \blacktriangle_b \mathcal{C}) \blacktriangledown_a \mathcal{B}. \end{aligned}$$

Proof. 1. We put $(\mathcal{A} \blacktriangle_a \mathcal{B}) \bullet_b \mathcal{C} = (A \sqcup_a B \sqcup_b C, \leq_S)$ and $(\mathcal{A} \bullet_b \mathcal{C}) \blacktriangle_a \mathcal{B} = (A \sqcup_b C \sqcup_a B, \leq_T)$. Let $x \in A \sqcup_a B \sqcup_b C$. Then $x \leq_S y$ if:

	$y \in A \setminus \{a, b\}$	$y \in B$	$y \in C$
$x \in A \setminus \{a, b\}$	$x \leq_A y$	$y \in \max(\mathcal{B})$ and $x \leq_A a$	$x \leq_A b$
$x \in B$	$a \leq_A y$	$x \leq_B y$	$x \leq_{\mathcal{A} \blacktriangle_a \mathcal{B}} b \iff a \leq_A b$
$x \in C$	$b \leq_A y$	$b \leq_{\mathcal{A} \blacktriangle_a \mathcal{B}} y \iff$ $y \in \max(\mathcal{B})$ and $b \leq_A a$	$x \leq_C y$

Moreover, $x \leq_T y$ if:

	$y \in A \setminus \{a, b\}$	$y \in B$	$y \in C$
$x \in A \setminus \{a, b\}$	$x \leq_A y$	$y \in \max(\mathcal{B})$ and $x \leq_A a$	$x \leq_A b$
$x \in B$	$a \leq_A y$	$x \leq_B y$	$x \leq_{\mathcal{A} \bullet_b \mathcal{C}} b \iff a \leq_A b$
$x \in C$	$b \leq_A y$	$y \in \max(\mathcal{B})$ and $x \leq_{\mathcal{A} \bullet_b \mathcal{C}} a \iff$ $y \in \max(\mathcal{B})$ and $b \leq_A a$	$x \leq_C y$

So $\leq_S = \leq_T$.

2. Can be deduced from the first point, with the help of the involution on posets.

3. We put $(\mathcal{A} \blacktriangledown_a \mathcal{B}) \blacktriangle_b \mathcal{C} = (A \sqcup_a B \sqcup C, \leq_S)$ and $(\mathcal{A} \blacktriangle_b \mathcal{C}) \blacktriangledown_a \mathcal{B} = (A \sqcup_b C \sqcup_a B, \leq_T)$. Let $x, y \in A \sqcup_a B \sqcup_b C$. Then $x \leq_S y$ if:

	$y \in A \setminus \{a, b\}$	$y \in B$	$y \in C$
$x \in A \setminus \{a, b\}$	$x \leq_A y$	$x \leq_A a$	$y \in \max(\mathcal{C})$ and $x \leq_A a$
$x \in B$	$x \in \min(\mathcal{B})$ and $a \leq_A y$	$x \leq_B y$	$y \in \max(\mathcal{C}), x \in \min(\mathcal{B})$ and $a \leq_A b$
$x \in C$	$b \leq_A y$	$b \leq_{\mathcal{A} \blacktriangledown_a \mathcal{B}} y \iff$ $b \leq_A a$	$x \leq_C y$

Moreover, $x \leq_T y$ if:

	$y \in A \setminus \{a, b\}$	$y \in B$	$y \in C$
$x \in A \setminus \{a, b\}$	$x \leq_A y$	$x \leq_A a$	$y \in \max(\mathcal{C})$ and $x \leq_A a$
$x \in B$	$x \in \min(\mathcal{B})$ and $a \leq_A y$	$x \leq_B y$	$y \in \max(\mathcal{C}), x \in \min(\mathcal{B})$ and $a \leq_{\mathcal{A} \blacktriangle_b \mathcal{C}} y \iff$ $y \in \max(\mathcal{C}), x \in \min(\mathcal{B})$ and $a \leq_A b$
$x \in C$	$b \leq_A y$	$x \leq_{\mathcal{A} \blacktriangle_b \mathcal{C}} a \iff$ $b \leq_A a$	$x \leq_C y$

Hence $\leq_S = \leq_T$. □

Remark 12. *Let $a \in A$ and $b \in B$. In general:*

$$\begin{aligned} (\mathcal{A} \blacktriangle_a \mathcal{B}) \bullet_b \mathcal{C} &\neq \mathcal{A} \blacktriangle_a (\mathcal{B} \bullet_b \mathcal{C}), & (\mathcal{A} \blacktriangledown_a \mathcal{B}) \bullet_b \mathcal{C} &\neq \mathcal{A} \blacktriangledown_a (\mathcal{B} \bullet_b \mathcal{C}), \\ (\mathcal{A} \bullet_a \mathcal{B}) \blacktriangle_b \mathcal{C} &\neq \mathcal{A} \bullet_a (\mathcal{B} \blacktriangle_b \mathcal{C}), & (\mathcal{A} \blacktriangledown_a \mathcal{B}) \blacktriangle_b \mathcal{C} &\neq \mathcal{A} \blacktriangledown_a (\mathcal{B} \blacktriangle_b \mathcal{C}), \\ (\mathcal{A} \bullet_a \mathcal{B}) \blacktriangledown_b \mathcal{C} &\neq \mathcal{A} \bullet_a (\mathcal{B} \blacktriangledown_b \mathcal{C}), & (\mathcal{A} \blacktriangle_a \mathcal{B}) \blacktriangledown_b \mathcal{C} &\neq \mathcal{A} \blacktriangle_a (\mathcal{B} \blacktriangledown_b \mathcal{C}). \end{aligned}$$

For example:

$$\begin{array}{ll}
 (\mathbb{1}^a \blacktriangle_a \mathbb{1}^b) \bullet_b \mathbb{1} = \mathbb{1}^b \bullet_b \mathbb{1} = \mathbb{1}^\wedge, & \mathbb{1}^a \blacktriangle_a (\mathbb{1}^b \bullet_b \mathbb{1}) = \mathbb{1}^a \blacktriangle_a \mathbb{1} = \mathbb{1}^\wedge. \\
 (\mathbb{1}_a \nabla_a \mathbb{1}_b) \bullet_b \mathbb{1} = \mathbb{V}_b \bullet_b \mathbb{1} = \mathbb{Y}, & \mathbb{1}_a \nabla_a (\mathbb{1}_b \bullet_b \mathbb{1}) = \mathbb{1}_a \nabla_a \mathbb{1} = \mathbb{V}. \\
 (\mathbb{1}^a \bullet_a \mathbb{1}^b) \blacktriangle_b \mathbb{1} = \mathbb{1}^b \bullet_b \mathbb{1} = \mathbb{1}^\wedge, & \mathbb{1}^a \bullet_a (\mathbb{1}^b \blacktriangle_b \mathbb{1}) = \mathbb{1}^a \blacktriangle_a \mathbb{1} = \mathbb{1}^\diamond. \\
 (\mathbb{1}_a \nabla_a \mathbb{1}^b) \blacktriangle_b \mathbb{1} = \mathbb{1}^b \mathbb{V} \blacktriangle_b \mathbb{1} = \mathbb{N}, & \mathbb{1}_a \nabla_a (\mathbb{1}^b \blacktriangle_b \mathbb{1}) = \mathbb{1}_a \blacktriangle_a \mathbb{V} = \mathbb{N}. \\
 (\mathbb{1}_a \bullet_a \mathbb{1}_b) \nabla_b \mathbb{1} = \mathbb{1}_b \bullet_b \mathbb{1} = \mathbb{V}, & \mathbb{1}_a \bullet_a (\mathbb{1}_b \nabla_b \mathbb{1}) = \mathbb{1}_a \blacktriangle_a \mathbb{V} = \mathbb{1}^\diamond. \\
 (\mathbb{1}^a \blacktriangle_a \mathbb{1}_b) \nabla_b \mathbb{1} = \mathbb{1}^a \blacktriangle_a \mathbb{V} = \mathbb{N}, & \mathbb{1}^a \blacktriangle_a (\mathbb{1}_b \nabla_b \mathbb{1}) = \mathbb{1}^a \blacktriangle_a \mathbb{V} = \mathbb{N}.
 \end{array}$$

5. ALGEBRAIC STRUCTURES ASSOCIATED TO THESE OPERADS

5.1. Products. Let us introduce some notations: let $\mathcal{A} \in \mathbb{P}_{\{1,2\}}$ and \mathcal{B}, \mathcal{C} be two finite posets. We put:

$$\begin{aligned}
 \mathcal{A} \circ (\mathcal{B}, \mathcal{C}) &= (\mathcal{A} \circ_1 \mathcal{B}) \circ_2 \mathcal{C} = (\mathcal{A} \circ_2 \mathcal{C}) \circ_1 \mathcal{B}, \\
 \mathcal{A} \bullet (\mathcal{B}, \mathcal{C}) &= (\mathcal{A} \bullet_1 \mathcal{B}) \bullet_2 \mathcal{C} = (\mathcal{A} \bullet_2 \mathcal{C}) \bullet_1 \mathcal{B}, \\
 \mathcal{A} \blacktriangle (\mathcal{B}, \mathcal{C}) &= (\mathcal{A} \blacktriangle_1 \mathcal{B}) \blacktriangle_2 \mathcal{C} = (\mathcal{A} \blacktriangle_2 \mathcal{C}) \blacktriangle_1 \mathcal{B}, \\
 \mathcal{A} \nabla (\mathcal{B}, \mathcal{C}) &= (\mathcal{A} \nabla_1 \mathcal{B}) \nabla_2 \mathcal{C} = (\mathcal{A} \nabla_2 \mathcal{C}) \nabla_1 \mathcal{B}.
 \end{aligned}$$

The free algebra on one generator over the different operadic structures on \mathbb{P} is the vector space:

$$F_{\mathbb{P}}(1) = \bigoplus_{n=1}^{\infty} \mathbb{P}(\{1, \dots, n\}) / \mathfrak{S}_n,$$

so it can be identified with the vector space generated by the isomorphism classes (shortly, isoclasses) of finite posets. This space inherits several bilinear products from the operad structures on posets, described below.

Notations. If \mathcal{A} is a finite poset, we denote by $[\mathcal{A}]$ its isomorphism class.

Theorem 13. Let $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$ be two finite posets. Then we have in $F_{\mathbb{P}}(1)$:

$$(1) \ .\mathbb{1} \ .\mathbb{2} \circ ([\mathcal{A}], [\mathcal{B}]) = \ .\mathbb{1} \ .\mathbb{2} \bullet ([\mathcal{A}], [\mathcal{B}]) = \ .\mathbb{1} \ .\mathbb{2} \blacktriangle ([\mathcal{A}], [\mathcal{B}]) = \ .\mathbb{1} \ .\mathbb{2} \nabla ([\mathcal{A}], [\mathcal{B}]) = [\mathcal{A}\mathcal{B}],$$

where $\mathcal{A}\mathcal{B} = (A \sqcup B, \leq)$, with, for all $x, y \in A \sqcup B$; $x \leq y$ if and only if:

- $x, y \in A$ and $x \leq_A y$,
- or $x, y \in B$ and $x \leq_B y$.

This product, defined on isoclasses of posets and also denoted by m , is sometimes called the disjoint union [12]. It is associative and commutative.

$$(2) \ \mathbb{1}_1^2 \bullet ([\mathcal{A}], [\mathcal{B}]) = [\mathcal{A} \downarrow \mathcal{B}], \text{ where } \mathcal{A} \downarrow \mathcal{B} = (A \sqcup B, \leq), \text{ with, for all } x, y \in A \sqcup B; x \leq y \text{ if and only if:}$$

- $x, y \in A$ and $x \leq_A y$,
- or $x, y \in B$ and $x \leq_B y$,
- or $x \in A$ and $y \in B$.

This product, defined on isoclasses of posets and also denoted by \downarrow , is sometimes called the direct sum [12]. It is associative.

$$(3) \ (\ .\mathbb{1} \ .\mathbb{2} + \mathbb{1}_1^2) \circ ([\mathcal{A}], [\mathcal{B}]) \text{ is the sum of isoclasses of posets } \mathcal{C} \text{ on } A \sqcup B \text{ such that:}$$

- If $x, y \in A$, $x \leq_C y$ if, and only if, $x \leq_A y$.
- If $x, y \in B$, $x \leq_C y$ if, and only if, $x \leq_B y$.
- $B < A$ in the sense of the Introduction, that is to say for all $x \in B$, $y \in A$, $x \not\leq_C y$.

This product, denoted by $*$, is associative.

(4) $\mathbf{1}_1^2 \blacktriangle([\mathcal{A}], [\mathcal{B}]) = [\mathcal{A} \blacktriangle \mathcal{B}]$, where $\mathcal{A} \blacktriangle \mathcal{B} = (A \sqcup B, \leq)$, with, for all $x, y \in A \sqcup B$; $x \leq y$ if and only if:

- $x, y \in A$ and $x \leq_A y$,
- or $x, y \in B$ and $x \leq_B y$,
- or $x \in A$ and $y \in \max(\mathcal{B})$.

This product, defined on isoclasses of posets and also denoted by Δ , is non associative permutative: for all $x, y, z \in F_{\mathbb{P}}(1)$,

$$x \Delta (y \Delta z) = (xy) \Delta z = y \Delta (x \Delta z).$$

(5) $\mathbf{1}_1^2 \blacktriangledown([\mathcal{A}], [\mathcal{B}]) = [\mathcal{A} \blacktriangledown \mathcal{B}]$, where $\mathcal{A} \blacktriangledown \mathcal{B} = (A \sqcup B, \leq)$, with, for all $x, y \in A \sqcup B$; $x \leq y$ if:

- $x, y \in A$ and $x \leq_A y$,
- or $x, y \in B$ and $x \leq_B y$,
- or $x \in \min(\mathcal{B})$ and $y \in A$.

This product, defined on isoclasses of posets and also denoted by ∇ , is non associative permutative: for all $x, y, z \in F_{\mathbb{P}}(1)$,

$$x \nabla (y \nabla z) = (xy) \nabla z = y \nabla (x \nabla z).$$

Proof. 1. We prove it for \circ ; the proof is similar in the other cases. First:

$$\begin{aligned} \bullet_1 \bullet_2 \stackrel{(12)}{=} \bullet_1 \bullet_2; \\ \bullet_1 \bullet_2 \circ_1 \bullet_1 \bullet_2 = \bullet_1 \bullet_2 \circ_2 \bullet_1 \bullet_2 = \bullet_1 \bullet_2 \bullet_3, \end{aligned}$$

so the product induced on $F_{\mathbb{P}}(1)$ by $\bullet_1 \bullet_2$ is associative and commutative. Moreover:

$$\bullet_1 \bullet_2 \circ ([\mathcal{A}], [\mathcal{B}]) = [(\bullet_1 \bullet_2 \circ_1 \mathcal{A}) \circ_2 \mathcal{B}].$$

By definition of \circ , $(\bullet_1 \bullet_2 \circ_1 \mathcal{A}) = \mathcal{A} \bullet_2$, and $\mathcal{A} \bullet_2 \circ_2 \mathcal{B} = \mathcal{A} \mathcal{B}$.

2. First:

$$\mathbf{1}_1^2 \bullet_1 \mathbf{1}_1^2 = \mathbf{1}_1^2 \bullet_2 \mathbf{1}_1^2 = \mathbf{1}_1^3,$$

so the product \downarrow is associative. Moreover:

$$\mathbf{1}_1^2 \bullet ([\mathcal{A}], [\mathcal{B}]) = [(\mathbf{1}_1^2 \bullet_1 \mathcal{A}) \bullet_2 \mathcal{B}].$$

The underlying set of the poset $\mathcal{C} = \mathbf{1}_1^2 \bullet_1 \mathcal{A}$ is $A \sqcup \{2\}$, and:

$$\{(x, y) \in (A \sqcup \{2\})^2 \mid x \leq_{\mathcal{C}} y\} = \{(x, y) \in A \mid x \leq_A y\} \sqcup ((A \sqcup \{2\}) \times \{2\}).$$

Hence, the underlying set of the poset $\mathcal{S} = (\mathbf{1}_1^2 \bullet_1 \mathcal{A}) \bullet_2 \mathcal{B}$ is $A \sqcup B$, and:

$$\{(x, y) \in (A \sqcup B)^2 \mid x \leq_{\mathcal{S}} y\} = \{(x, y) \in A \mid x \leq_A y\} \sqcup \{(x, y) \in B \mid x \leq_B y\} \sqcup (A \times B).$$

3. First:

$$\begin{aligned} (\bullet_1 \bullet_2 + \mathbf{1}_1^2) \circ_1 (\bullet_1 \bullet_2 + \mathbf{1}_1^2) &= \bullet_1 \bullet_2 \bullet_3 + \bullet_1 \mathbf{1}_1^3 + \bullet_2 \mathbf{1}_1^3 + \bullet_3 \mathbf{1}_1^2 + {}^2 \mathcal{V}_1^3 + {}_1 \mathcal{K}_2^3 + \mathbf{1}_1^3 \\ &= (\bullet_1 \bullet_2 + \mathbf{1}_1^2) \circ_2 (\bullet_1 \bullet_2 + \mathbf{1}_1^2), \end{aligned}$$

so $*$ is associative. Let us compute $\mathbf{1}_1^2 \circ (\mathcal{A}, \mathcal{B}) = (\mathbf{1}_1^2 \circ_1 \mathcal{A}) \circ_2 \mathcal{B}$. By definition, $\mathbf{1}_1^2 \circ_1 \mathcal{A}$ is the sum of the posets \mathcal{C} on $A \sqcup \{2\}$ such that:

- For all $x, y \in A$, $x \leq_{\mathcal{C}} y$ if, and only if, $x \leq_A y$.
- For all $y \in A$, we do not have $2 \leq_{\mathcal{C}} y$.
- There exists $x \in A$, $x \leq_{\mathcal{C}} 2$.

Hence, $\mathbf{1}_1^2 \circ (\mathcal{A}, \mathcal{B})$ is the sum of all the posets \mathcal{S} on $A \sqcup B$ such that:

- For all $x, y \in A$, $x \leq_{\mathcal{S}} y$ if, and only if, $x \leq_A y$.
- For all $x, y \in B$, $x \leq_{\mathcal{S}} y$ if, and only if, $x \leq_B y$.

- For all $x \in B, y \in A$, we do not have $x \leq_S y$.
- There exists $x \in A, y \in B$, such that $x \leq_S y$.

These conditions are equivalent to:

- For all $x, y \in A, x \leq_S y$ if, and only if, $x \leq_A y$.
- For all $x, y \in B, x \leq_S y$ if, and only if, $x \leq_B y$.
- $A < B$.
- $S \neq \mathcal{A}\mathcal{B}$.

Finally, $\mathfrak{I}_1^2 \circ ([\mathcal{A}], [\mathcal{B}])$ is the sum of isoclasses of posets S on $A \sqcup B$ such that:

- If $x, y \in A, x \leq_S y$ if, and only if, $x \leq_A y$.
- If $x, y \in B, x \leq_S y$ if, and only if, $x \leq_B y$.
- $A <_S B$.
- $S \neq \mathcal{A}\mathcal{B}$.

The first point gives the result for $*$.

4. First:

$$\mathfrak{I}_1^2 \blacktriangle_2 \mathfrak{I}_1^2 = \mathfrak{I}_1^2 \blacktriangle_{1 \cdot 1 \cdot 2} = {}_1 \mathfrak{A}_2^3.$$

So for all $x, y, z \in F_{\mathbb{P}}(1)$, $x \Delta (y \Delta z) = (xy) \Delta z$. The commutativity of the product m gives the permutativity of Δ . Moreover, $[\mathcal{A}] \Delta [\mathcal{B}] = [(\mathfrak{I}_1^2 \blacktriangle_1 \mathcal{A}) \blacktriangle_2 \mathcal{B}]$; the underlying set of the poset $\mathcal{C} = \mathfrak{I}_1^2 \blacktriangle_1 \mathcal{A}$ is $A \sqcup \{2\}$, and:

$$\{(x, y) \in (A \sqcup \{2\})^2 \mid x \leq_{\mathcal{C}} y\} = \{(x, y) \in A^2 \mid x \leq_A y\} \sqcup ((A \sqcup \{2\}) \times \{2\}).$$

so the underlying set of $S = (\mathfrak{I}_1^2 \blacktriangle_1 \mathcal{A}) \blacktriangle_2 \mathcal{B}$ is $A \sqcup B$ and:

$$\{(x, y) \in (A \sqcup B)^2 \mid x \leq_S y\} = \{(x, y) \in A \mid x \leq_A y\} \sqcup \{(x, y) \in B \mid x \leq_B y\} \sqcup (A \times \max(\mathcal{B})).$$

5. Comes from the preceding point, using the involution on posets. \square

The products m, \downarrow and $*$ are extended to $\overline{F}_{\mathbb{P}}(1) = \mathbb{K} \oplus F_{\mathbb{P}}(1)$, by assuming that $1 \in \mathbb{K}$ is the unit for all these products. We now identify 1 with the empty poset.

Remark 14. *We could also work with the free \mathbb{P} -algebra $F_{\mathbb{P}}(D)$ generated by a set D : this is the vector space generated by isoclasses of posets decorated by D , that is to say pairs (\mathcal{A}, d) , where $\mathcal{A} = (A, \leq_A)$ is a poset and $d : A \rightarrow D$ is a map.*

5.2. Coproducts. We identify $\overline{F}_{\mathbb{P}}(1)$ and its graded dual, via the pairing defined on two isoclasses of posets $[\mathcal{A}], [\mathcal{B}]$ by:

$$\langle [\mathcal{A}], [\mathcal{B}] \rangle = s_{[\mathcal{A}]} \delta_{[\mathcal{A}], [\mathcal{B}]},$$

where $s_{[\mathcal{A}]}$ is the number of poset automorphisms of \mathcal{A} . We now define two coproducts on $F_{\mathbb{P}}(1)$. Let $\mathcal{A} \in \mathbb{P}_A$.

- We decompose it as $\mathcal{A} = \mathcal{A}_1 \dots \mathcal{A}_k$, where \mathcal{A}_i are the connected components of \mathcal{A} . Then:

$$\Delta([\mathcal{A}]) = \sum_{I \subseteq \{1, \dots, k\}} \left[\prod_{i \in I} \mathcal{A}_i \right] \otimes \left[\prod_{i \notin I} \mathcal{A}_i \right].$$

This coproduct is dual of the product m .

- We put:

$$\Delta_*([\mathcal{A}]) = \sum_{I \subseteq A, A \setminus I < I} [\mathcal{A}_{|A \setminus I}] \otimes [\mathcal{A}_{|I}].$$

This coproduct is the one of the Introduction, and it is the dual of the product $*$.

Theorem 15. (1) $(\overline{F}_{\mathbb{P}}(1), *, \Delta)$ and $(\overline{F}_{\mathbb{P}}(1), m, \Delta_*)$ are dual bialgebras.
 (2) $(\overline{F}_{\mathbb{P}}(1), \downarrow, \Delta_*)$ is an infinitesimal bialgebra [5].

Proof. Let $\mathcal{A} = (A, \leq_A)$ be a finite poset, and $I \subseteq A$. We shall say that I is an ideal of \mathcal{A} if $A \setminus I < I$.

1. For $(\overline{F}_{\mathbb{P}}(1), m, \Delta_*)$, it remains only to prove the compatibility of m and Δ_* . Let \mathcal{A} and \mathcal{B} be finite posets. The ideals of $\mathcal{A}\mathcal{B}$ are the subposets $I = I_1 I_2$, where I_j is an ideal of \mathcal{A}_j for all j . So:

$$\begin{aligned} \Delta_*([\mathcal{A}\mathcal{B}]) &= \sum_{I \text{ ideal of } \mathcal{A}\mathcal{B}} \sum_{J \text{ ideal of } \mathcal{A}\mathcal{B}} [\mathcal{A}\mathcal{B}]_{I \sqcup B \setminus I \sqcup J} \otimes [\mathcal{A}\mathcal{B}]_{I \sqcup J} \\ &= \sum_{I \text{ ideal of } \mathcal{A}} \sum_{J \text{ ideal of } \mathcal{B}} [\mathcal{A}]_{A \setminus I} [\mathcal{B}]_{B \setminus J} \otimes [\mathcal{A}]_I [\mathcal{B}]_J \\ &= \Delta_*(\mathcal{A}) \Delta_*(\mathcal{B}). \end{aligned}$$

Hence, $(\overline{F}_{\mathbb{P}}(1), m, \Delta_*)$ is a bialgebra. By duality, $(\overline{F}_{\mathbb{P}}(1), *, \Delta)$ also is.

2. It remains to prove the compatibility between \downarrow and Δ_* . Let \mathcal{A} and \mathcal{B} be two finite posets. The ideals of $\mathcal{A} \downarrow \mathcal{B}$ are the ideals of \mathcal{B} and the ideals $I \sqcup B$, where I is an ideal of \mathcal{A} . Note that in this description, B appears two times, as B and $\emptyset \sqcup B$. Hence:

$$\begin{aligned} \Delta_*([\mathcal{A} \downarrow \mathcal{B}]) &= \sum_{I \text{ ideal of } \mathcal{A}} [\mathcal{A} \downarrow \mathcal{B}]_{A \setminus I} \otimes [\mathcal{A} \downarrow \mathcal{B}]_{I \sqcup B} + \sum_{I \text{ ideal of } \mathcal{B}} [\mathcal{A} \downarrow \mathcal{B}]_{A \sqcup B \setminus I} \otimes [\mathcal{A} \downarrow \mathcal{B}]_I \\ &\quad - [\mathcal{A} \downarrow \mathcal{B}]_{\mathcal{A}} \otimes [\mathcal{A} \downarrow \mathcal{B}]_{\mathcal{B}} \\ &= \sum_{I \text{ ideal of } \mathcal{A}} [\mathcal{A}]_{A \setminus I} \otimes [\mathcal{A}]_I \downarrow [\mathcal{B}] + \sum_{I \text{ ideal of } \mathcal{B}} [\mathcal{A}] \downarrow [\mathcal{B}]_{B \setminus I} \otimes [\mathcal{B}]_I - [\mathcal{A}] \otimes [\mathcal{B}] \\ &= \Delta_*([\mathcal{A}]) \downarrow (1 \otimes [\mathcal{B}]) + ([\mathcal{A}] \otimes 1) \downarrow \Delta_*([\mathcal{B}]) - [\mathcal{A}] \otimes [\mathcal{B}]. \end{aligned}$$

So $(\overline{F}_{\mathbb{P}}(1), \downarrow, \Delta_*)$ is indeed an infinitesimal bialgebra. \square

All these objects are graded by the cardinality of posets and are connected. By the rigidity theorem for connected infinitesimal bialgebras [5], $(\overline{F}_{\mathbb{P}}(1), \downarrow, \Delta_*)$ is isomorphic to a tensor algebra, with the concatenation product and the deconcatenation coproduct. Consequently:

Corollary 16. *The commutative bialgebra $(\overline{F}_{\mathbb{P}}(1), m, \Delta_*)$ is cofree; the cocommutative bialgebra $(\overline{F}_{\mathbb{P}}(1), *, \Delta)$ is free.*

6. SUBOPERADS GENERATED IN DEGREE 2

6.1. WN Posets. Let \mathcal{A} be a finite poset. We shall say that it is a *poset without N*, or *WN poset*, if it does not contain any subposet isomorphic to \mathbf{N} [12, 3]. For any finite set A , the space of finite WN poset structures on A is denoted by $\text{WN}\mathbb{P}_A$. We define in this way a linear species $\text{WN}\mathbb{P}$. For example, here are the isoclasses of WN posets of cardinality ≤ 4 :

$$\begin{aligned} &.; \quad .., \mathbf{1}; \quad \dots, .., \mathbf{1}, \mathbf{V}, \mathbf{1}, \mathbf{\Lambda}; \\ &\dots, .., \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{V}, .., \mathbf{1}, \mathbf{\Lambda}, .., \mathbf{V}, \mathbf{1}, \mathbf{V}, \mathbf{Y}, \mathbf{1}, \mathbf{\Lambda}, \mathbf{1}, \mathbf{1}, \mathbf{\Lambda}, \mathbf{\Lambda}, \mathbf{\Lambda}, \mathbf{\Lambda}, \mathbf{\Lambda}. \end{aligned}$$

Lemma 17. (1) *Let \mathcal{A} be a WN poset. It can be written in a unique way as $\mathcal{A} = \mathcal{A}_1 \downarrow \dots \downarrow \mathcal{A}_k$, with for all $1 \leq i \leq k$, $[\mathcal{A}_i] = \cdot$ or \mathcal{A}_i is not connected.*

(2) *Let \mathcal{A} and \mathcal{B} be two finite posets. The following conditions are equivalent:*

- (a) \mathcal{A} and \mathcal{B} are WN.
- (b) $\mathcal{A}\mathcal{B}$ is WN.
- (c) $\mathcal{A} \downarrow \mathcal{B}$ is WN.

Proof. (1) *Existence.* We proceed by induction on $n = |\mathcal{A}|$. If $n = 1$, then $\lfloor \mathcal{A} \rfloor = \cdot$ and the result is obvious. Let us assume the result at all rank $< n$. If \mathcal{A} is not connected, we choose $k = 1$ and $\mathcal{A}_1 = \mathcal{A}$. Let us assume that \mathcal{A} is connected. If \mathcal{A} has a unique maximal element M , then for all $x \in \mathcal{A}$, $x \leq_A M$ so $\mathcal{A} = (\mathcal{A} \setminus \{M\}) \downarrow \{M\}$: we then apply the induction hypothesis to $\mathcal{A} \setminus \{M\}$ and obtain a decomposition of \mathcal{A} . Let us assume that \mathcal{A} has at least two maximal elements. We put:

$$\mathcal{A}' = \{x \in \mathcal{A} \mid \forall M \in \max(\mathcal{A}'), x \leq_A M\}, \quad \mathcal{A}'' = \mathcal{A} \setminus \mathcal{A}'.$$

Let $M \in \max(\mathcal{A})$ and M' be another maximal element of \mathcal{A} . We do not have $M \leq M'$, so $M \in \mathcal{A}''$, hence $\max(\mathcal{A}) \subseteq \mathcal{A}''$. Let $m \in \min(\mathcal{A})$, and let us assume that $m \notin \mathcal{A}'$. So there exists a maximal element M' , such that m and M' are not comparable in \mathcal{A} . Moreover, there exists $m \in \max(\mathcal{A})$, $m \leq_A M$. As \mathcal{A} is connected, there exists elements x_1, \dots, x_{2k-1} such that $x_0 = M \geq_A x_1 \leq_A x_2 \geq_A \dots \geq_A x_{2k-1} \leq_A x_{2k} = M'$. We choose these elements in such a way that k is minimal. As \mathcal{A} is WN, necessarily $k = 1$, and then M, m, x_1, M' is a copy of \mathbf{N} in \mathcal{A} : contradiction. So $\min(\mathcal{A}) \subseteq \mathcal{A}'$: consequently, \mathcal{A}' and \mathcal{A}'' are nonempty.

Let $x \in \mathcal{A}'$ and $y \in \mathcal{A}''$. If $y \in \max(\mathcal{A})$, by definition of \mathcal{A}' , $x \leq_A y$. If not, as $y \in \mathcal{A}''$, there exists $M \in \max(\mathcal{A})$ such that y and M are not comparable in \mathcal{A} ; there exists $M' \in \max(\mathcal{A})$, such that $y \leq_A M'$. As $x \in \mathcal{A}'$, $x \leq_A M, M'$. As $\{x, y, M, M'\}$ cannot be isomorphic to \mathbf{N} , necessarily $x \leq_A y$. Finally, $\mathcal{A} = \mathcal{A}' \downarrow \mathcal{A}''$. Applying the induction hypothesis to \mathcal{A}' and \mathcal{A}'' , we obtain the result.

Unicity. Let us assume that $\mathcal{A} = \mathcal{A}_1 \downarrow \dots \downarrow \mathcal{A}_k = \mathcal{A}'_1 \downarrow \dots \downarrow \mathcal{A}'_l$, with for all i, j , \mathcal{A}_i and \mathcal{A}'_j reduced to a single point or non connected. We proceed by induction on k . First, observe that if $k \geq 2$, \mathcal{A} is connected and not reduced to a single point. So $k = 1$ is equivalent to $l = 1$, which proves the result for $k = 1$. Let us assume the result at rank $k - 1$. We consider $\mathcal{A}' = \{x \in \mathcal{A} \mid \forall M \in \max(\mathcal{A}), x <_A M\}$. Clearly, $\mathcal{A}_1 \downarrow \dots \downarrow \mathcal{A}_{k-1} \subseteq \mathcal{A}'$. If $\mathcal{A}_k = \{M\}$, then M is the unique maximal element of \mathcal{A} , and $M \notin \mathcal{A}'$. If \mathcal{A}_k is not reduced to a single element, then it is not connected. Let $x \in \mathcal{A}_k$ and let M be a maximal element of \mathcal{A}_k which is in a different connected component. Then M is a maximal element of \mathcal{A} and we do not have $x <_A M$, so $x \notin \mathcal{A}'$. Consequently, $\mathcal{A}' = \mathcal{A}_1 \downarrow \dots \downarrow \mathcal{A}_{k-1}$. Similarly, $\mathcal{A}' = \mathcal{A}'_1 \downarrow \dots \downarrow \mathcal{A}'_{l-1}$. Taking the complement, $\mathcal{A}_k = \mathcal{A}'_l$. Using the induction hypothesis, $k = l$, and $\mathcal{A}_i = \mathcal{A}'_i$ for all $1 \leq i \leq k - 1$.

(2) (a) \iff (b). Let \mathcal{N} be a copy of \mathbf{N} in $\mathcal{A}\mathcal{B}$. As \mathbf{N} is connected, it is included in a component of $\mathcal{A}\mathcal{B}$, so is included in A or is included in B . So $\mathcal{A}\mathcal{B}$ is WN if, and only if, \mathcal{A} and \mathcal{B} are.

(a) \iff (c). Let \mathcal{N} be a copy of \mathbf{N} in $\mathcal{A} \downarrow \mathcal{B}$. We put $N_1 = \mathcal{N} \cap A$ and $N_2 = \mathcal{N} \cap B$. Then, for all $x \in N_1$, $y \in N_2$, $x \leq_{\mathcal{A}\downarrow\mathcal{B}} y$, so $\mathcal{N} = \mathcal{N}|_{N_1} \downarrow \mathcal{N}|_{N_2}$. The only possibilities are $(N_1, N_2) = (\mathcal{N}, \emptyset)$ or (\emptyset, \mathcal{N}) , so $\mathcal{N} \subseteq A$ or $\mathcal{N} \subseteq B$. Hence, $\mathcal{A} \downarrow \mathcal{B}$ is WN if, and only if, \mathcal{A} and \mathcal{B} are. \square


The vector space $F_{\text{WN}\mathbb{P}}(1)$ generated by the set of isoclasses of WN posets is an algebra for both products m and \downarrow : this is a 2-As algebra, with the terminology of [7]. As the first product is commutative, we shall say that it is a Com-As algebra.

Theorem 18. $\text{WN}\mathbb{P}$ is a suboperad of (\mathbb{P}, \bullet) . It is generated by $m = \cdot_1 \cdot_2$ and $\downarrow = \mathbf{!}_1^2 \in \mathbb{P}_{\{1,2\}}$, and the relations:

$$m^{(12)} = m, \quad m \bullet_1 m = m \bullet_2 m, \quad \downarrow \bullet_1 \downarrow = \downarrow \bullet_2 \downarrow.$$

Proof. *First step.* Let us first prove that $\text{WN}\mathbb{P}$ is a suboperad of \mathbb{P} for the product \bullet . Let \mathcal{A}, \mathcal{B} be two WN posets, and $a \in \mathcal{A}$. Let us assume that $\mathcal{C} = \mathcal{A} \bullet_a \mathcal{B}$ contains a copy N of \mathbf{N} . The

elements of N are denoted in this way:



As \mathcal{A} and \mathcal{B} are WN, N cannot be included in $\mathcal{A} \setminus \{a\}$, nor in \mathcal{B} . If $N \cap \mathcal{B}$ is a singleton, $\mathcal{C}/\mathcal{B} \twoheadrightarrow \{a\} = \mathcal{A}$ contains a copy of \mathbf{N} , formed by a and the three elements of $N \cap \mathcal{A} \setminus \{a\}$: this is a contradiction. If $N \cap \mathcal{B}$ contains two elements, six cases are possible:

- $N \cap \mathcal{B} = \{x, y\}$: as $y \leq_C t$, we should have $x \leq_C t$: this is a contradiction.
- $N \cap \mathcal{B} = \{x, z\}$: as $y \leq_C z$, we should have $x \leq_C z$: this is a contradiction.
- $N \cap \mathcal{B} = \{x, t\}$: as $x \leq_C z$, we should have $t \leq_C z$: this is a contradiction.
- $N \cap \mathcal{B} = \{y, z\}$: as $x \leq_C z$, we should have $x \leq_C y$: this is a contradiction.
- $N \cap \mathcal{B} = \{y, t\}$: as $y \leq_C z$, we should have $t \leq_C z$: this is a contradiction.
- $N \cap \mathcal{B} = \{z, t\}$: as $x \leq_C z$, we should have $x \leq_C t$: this is a contradiction.

If $N \cap \mathcal{B}$ contains three elements, four cases are possible:

- $N \cap \mathcal{B} = \{x, y, z\}$: as $y \leq_C t$, we should have $x \leq_C t$: this is a contradiction.
- $N \cap \mathcal{B} = \{x, y, t\}$: as $y \leq_C z$, we should have $t \leq_C z$: this is a contradiction.
- $N \cap \mathcal{B} = \{x, z, t\}$: as $y \leq_C t$, we should have $y \leq_C x$: this is a contradiction.
- $N \cap \mathcal{B} = \{y, z, t\}$: as $x \leq_C z$, we should have $x \leq_C y$: this is a contradiction.

In all cases, we obtain a contradiction, so \mathcal{C} is WN.

Second step. We denote by \mathbb{P}' the suboperad of \mathbb{P} generated by $\cdot_1 \cdot_2$ and $\mathbf{!}_1^2$. By the first point, $\mathbb{P}' \subseteq \mathbb{WNP}$. Let us prove the inverse inclusion. Let \mathcal{A} be a WN poset, let us prove that $\mathcal{A} \in \mathbb{P}'$ by induction on $n = |\mathcal{A}|$. This is obvious if $n = 1$ or $n = 2$. Let us assume the result at all rank $< n$, with $n \geq 3$. If \mathcal{A} is not connected, we can write $\mathcal{A} = \mathcal{A}_1 \mathcal{A}_2$, with \mathcal{A}_1 and \mathcal{A}_2 nonempty. By restriction, \mathcal{A}_1 and \mathcal{A}_2 are WN, so belong to \mathbb{P}' by the induction hypothesis. Then:

$$\mathcal{A} = \cdot_1 \cdot_2 \bullet (\mathcal{A}_1, \mathcal{A}_2) \in \mathbb{P}'.$$

If \mathcal{A} is connected, as it is WN we can write it as $\mathcal{A} = \mathcal{A}_1 \downarrow \mathcal{A}_2$, with \mathcal{A}_1 and \mathcal{A}_2 nonempty. By restriction, \mathcal{A}_1 and \mathcal{A}_2 are WN, so belong to \mathbb{P}' by the induction hypothesis. Then:

$$\mathcal{A} = \mathbf{!}_1^2 \bullet (\mathcal{A}_1, \mathcal{A}_2) \in \mathbb{P}'.$$

Last step. In order to give the presentation of \mathbb{WNP} by generators and relations, it is enough to prove that free \mathbb{WNP} -algebras satisfy the required universal property. We restrict ourselves to $F_{\mathbb{WNP}}(1)$, the other cases are proved similarly. More precisely, let (A, m, \downarrow) be a Com-As algebra, that is to say that (A, m) is an associative, commutative algebra, and (A, \downarrow) is an associative algebra, and let $a \in A$. Let us show that there exists a unique morphism ϕ of Com-As algebras from $F_{\mathbb{WNP}}(1)$ to A , sending \cdot to a . For this, we consider an iso-class $[\mathcal{A}]$ of a WN poset. We define $\phi([\mathcal{A}])$ by induction on the cardinality of \mathcal{A} in the following way:

- $\phi(\cdot) = a$.
- If ϕ is not connected, we put $\mathcal{A} = \mathcal{A}_1 \dots \mathcal{A}_k$, where $k \geq 2$ and $\mathcal{A}_1, \dots, \mathcal{A}_k$ are connected. We put:

$$\phi([\mathcal{A}]) = \phi([\mathcal{A}_1]) \dots \phi([\mathcal{A}_k]).$$

As the product m of A is commutative, this does not depend of the chosen order on the connected components of \mathcal{A} , so is well-defined.

- If $[\mathcal{A}] \neq \cdot$ and \mathcal{A} is connected, by the preceding lemma it can be uniquely written as $\mathcal{A} = \mathcal{A}_1 \downarrow \dots \downarrow \mathcal{A}_k$, with $k \geq 2$, with for all i $[\mathcal{A}_i] = \cdot$ or \mathcal{A}_i not connected. We put:

$$\phi([\mathcal{A}]) = \phi([\mathcal{A}_1]) \downarrow \dots \downarrow \phi([\mathcal{A}_k]).$$

It is an easy exercise to prove that ϕ is indeed a Com-As algebra morphism. \square

6.2. Compositions \blacktriangle and \blacktriangledown . We first define the notion of ∇ -compatible poset, inductively on the cardinality. Let $\mathcal{A} = (A, \leq_A)$ be a finite poset of cardinality n . We shall need the following notation: we first put

$$A' = \{y \in A \mid \forall x \in \min(\mathcal{A}), x <_A y\}.$$

We then define $b\mathcal{A} = \mathcal{A}|_{A'}$ and $r\mathcal{A} = \mathcal{A}|_{A \setminus A'}$. Note that the minimal elements of \mathcal{A} do not belong to A' , so $r\mathcal{A}$ is not empty. It may happen that $b\mathcal{A}$ is empty.

Let us now define ∇ -compatible posets by induction on the cardinality. If $n = 1$, then $[\mathcal{A}] = \cdot$ is ∇ -compatible. If $n \geq 2$, we shall say that \mathcal{A} is ∇ -compatible if one of the following conditions holds:

- (1) \mathcal{A} is not connected and all the connected components of \mathcal{A} are ∇ -compatible.
- (2) \mathcal{A} is connected and the following conditions hold:
 - (a) $b\mathcal{A}$ is not empty.
 - (b) $b\mathcal{A}$ and $r\mathcal{A}$ are ∇ -compatible.
 - (c) $\mathcal{A} = b\mathcal{A} \nabla r\mathcal{A}$.

The subspecies of \mathbb{P} of ∇ -compatible posets is denoted by \mathbb{CP}_∇ .

For example, here are the isoclasses of ∇ -compatible posets of cardinality ≤ 4 :

$$\begin{aligned} & \cdot; \quad \dots, \mathbf{I}; \quad \dots, \mathbf{I}, \mathbf{V}, \mathbf{I}, \mathbf{\Lambda}; \\ & \dots, \dots, \mathbf{I}, \mathbf{I}, \mathbf{I}, \mathbf{V}, \mathbf{\cdot}, \mathbf{I}, \mathbf{\cdot}, \mathbf{\Lambda}, \mathbf{\cdot}, \mathbf{V}, \mathbf{V}, \mathbf{Y}, \mathbf{I}, \mathbf{\cdot}, \mathbf{\Lambda}, \mathbf{\Lambda}, \mathbf{N}, \mathbf{M}, \mathbf{\Diamond}. \end{aligned}$$

Lemma 19. *Let $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$ be two finite posets.*

- (1) $b(\mathcal{A} \nabla \mathcal{B}) = Ab\mathcal{B}$ and $r(\mathcal{A} \nabla \mathcal{B}) = r\mathcal{B}$.
- (2) *Let us assume that \mathcal{A} is ∇ -compatible.*
 - (a) \mathcal{A} is connected and different from \cdot if, and only if, $b\mathcal{A}$ is nonempty.
 - (b) If \mathcal{A} is connected, then $[r\mathcal{A}] = \cdot$ or $r\mathcal{A}$ is not connected.

Proof. (1) This comes easily from the observation that $\min(\mathcal{A} \nabla \mathcal{B}) = \min \mathcal{B}$.

(2) (a) \implies . By definition of the ∇ -compatibility. \impliedby . If $[\mathcal{A}] = \cdot$, then obviously $b\mathcal{A}$ is empty. If \mathcal{A} is not connected, let us take $a \in A$. If m is a minimal element of connected component of \mathcal{A} which does not contain a , we do not have $a >_A m$, so $a \notin A'$: $b\mathcal{A}$ is empty.

2. (b) Let us assume that $[r\mathcal{A}] \neq \cdot$ and is connected. By (2) (a), $br\mathcal{A}$ is not empty. By (1), $b\mathcal{A} = b(b\mathcal{A} \nabla r\mathcal{A}) = bAbr\mathcal{A} \neq b\mathcal{A}$, which is a contradiction. \square

Lemma 20. *Let $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$ be two ∇ -compatible posets. Then $\mathcal{A}\mathcal{B}$ and $\mathcal{A} \nabla \mathcal{B}$ are ∇ -compatible.*

Proof. Let $\mathcal{A} = \mathcal{A}_1 \dots \mathcal{A}_k$ and $\mathcal{B} = \mathcal{B}_1 \dots \mathcal{B}_l$ be the decomposition of \mathcal{A} and \mathcal{B} into connected components. By definition, $\mathcal{A}_1, \dots, \mathcal{A}_k, \mathcal{B}_1, \dots, \mathcal{B}_l$ are ∇ -compatible. Then $\mathcal{A}\mathcal{B}$ is not connected and its connected components are $\mathcal{A}_1, \dots, \mathcal{A}_k, \mathcal{B}_1, \dots, \mathcal{B}_l$, so $\mathcal{A}\mathcal{B}$ is ∇ -compatible.

Let us prove that $\mathcal{C} = \mathcal{A} \nabla \mathcal{B}$ is ∇ -compatible. First, observe that $r\mathcal{C} = r\mathcal{B}$ and $b\mathcal{C} = Ab\mathcal{B}$.

- If \mathcal{B} is not connected or reduced to a single element, $r\mathcal{C} = \mathcal{B}$ and $b\mathcal{C} = \mathcal{A}$, so are both ∇ -compatible.
- If not, $r\mathcal{C} = r\mathcal{B}$ and $b\mathcal{C} = Ab\mathcal{B}$ are both ∇ -compatible, as \mathcal{B} is ∇ -compatible.

Moreover:

$$b\mathcal{C}\nabla r\mathcal{C} = (Ab\mathcal{B})\nabla r\mathcal{B} = \mathcal{A}\nabla(b\mathcal{B}\nabla r\mathcal{B}) = \mathcal{A}\nabla\mathcal{B} = \mathcal{C},$$

so \mathcal{C} is ∇ -compatible. \square

Theorem 21. *The species $\mathbb{C}\mathcal{A}_\nabla$ is a suboperad of (\mathbb{P}, ∇) . It is generated by the elements $m = \bullet_1 \bullet_2$ and $\nabla = \mathbf{1}_1^2$ in $\mathbb{C}\mathbb{P}^\nabla(\{1, 2\})$, and the relations:*

$$m^{(12)} = m, \quad m\nabla_1 m = m\nabla_2 m, \quad \nabla\nabla_1 m = \nabla\nabla_2 \nabla.$$

Proof. First step. Let us prove that $\mathbb{C}\mathbb{P}^\nabla$ is a suboperad. Let $\mathcal{A} = (A, \leq_A)$ and $\mathcal{B} = (B, \leq_B)$ be two ∇ -compatible posets, and let $a \in A$; let us show that $\mathcal{C} = \mathcal{A}\nabla_a\mathcal{B}$ is ∇ -compatible. We proceed by induction on the cardinality n of \mathcal{A} . If $n = 1$, then $\mathcal{C} = \mathcal{B}$ and the result is obvious. Let us assume the result at all ranks $< n$. If \mathcal{A} is not connected, we can write $\mathcal{A} = \mathcal{A}_1\mathcal{A}_2$, with \mathcal{A}_1 and \mathcal{A}_2 nonempty ∇ -compatible posets. Then:

$$\begin{aligned} \mathcal{C} &= (\bullet_1 \bullet_2 \nabla(\mathcal{A}_1, \mathcal{A}_2))\nabla_a\mathcal{B} \\ &= \bullet_1 \bullet_2 \nabla(\mathcal{A}_1\nabla_a\mathcal{B}, \mathcal{A}_2) \text{ or } \bullet_1 \bullet_2 \nabla(\mathcal{A}_1, \mathcal{A}_2\nabla_a\mathcal{B}) \\ &= (\mathcal{A}_1\nabla_a\mathcal{B})\mathcal{A}_2 \text{ or } \mathcal{A}_1(\mathcal{A}_2\nabla_a\mathcal{B}). \end{aligned}$$

We conclude with the induction hypothesis applied to \mathcal{A}_1 or \mathcal{A}_2 and with lemma 20. If \mathcal{A} is connected, we can write $\mathcal{A} = \mathcal{A}_1\nabla\mathcal{A}_2$, with $\mathcal{A}_1 = b\mathcal{A}$ and $\mathcal{A}_2 = r\mathcal{A}$ nonempty ∇ -compatible posets. Then:

$$\begin{aligned} \mathcal{C} &= (\mathbf{1}_1^2 \nabla(\mathcal{A}_1, \mathcal{A}_2))\nabla_a\mathcal{B} \\ &= \mathbf{1}_1^2 \nabla(\mathcal{A}_1\nabla_a\mathcal{B}, \mathcal{A}_2) \text{ or } \mathbf{1}_1^2 \nabla(\mathcal{A}_1, \mathcal{A}_2\nabla_a\mathcal{B}) \\ &= (\mathcal{A}_1\nabla_a\mathcal{B})\nabla\mathcal{A}_2 \text{ or } \mathcal{A}_1\nabla(\mathcal{A}_2\nabla_a\mathcal{B}). \end{aligned}$$

We conclude with the induction hypothesis applied to \mathcal{A}_1 or \mathcal{A}_2 and with lemma 20.

Second step. Let \mathbb{P}' be the suboperad of (\mathbb{P}, ∇) generated by $\bullet_1 \bullet_2$ and $\mathbf{1}_1^2$. The first step implies that $\mathbb{P}' \subseteq \mathbb{C}\mathbb{P}^\nabla$. Let us prove the inverse inclusion. Let \mathcal{A} be a ∇ -compatible poset; let us prove that $\mathcal{A} \in \mathbb{P}'$ by induction on its cardinality n . It is obvious if $n = 1$. Let us assume the result at all rank $< n$. If \mathcal{A} is not connected, we can write $\mathcal{A} = \mathcal{A}_1\mathcal{A}_2$, with \mathcal{A}_1 and \mathcal{A}_2 nonempty ∇ -compatible posets. Then $\mathcal{A} = \bullet_1 \bullet_2 \nabla(\mathcal{A}_1, \mathcal{A}_2)$ belongs to \mathbb{P}' by the induction hypothesis. If \mathcal{A} is connected, we can write $\mathcal{A} = \mathcal{A}_1\nabla\mathcal{A}_2$, with $\mathcal{A}_1 = b\mathcal{A}$ and $\mathcal{A}_2 = r\mathcal{A}$ nonempty ∇ -compatible posets. Then $\mathcal{A} = \mathbf{1}_1^2 \nabla(\mathcal{A}_1, \mathcal{A}_2)$ belongs to \mathbb{P}' by the induction hypothesis.

Last step. In order to prove the generation of $\mathbb{C}\mathbb{P}^\nabla$ by generators and relations, it is enough to prove the required universal property for free $\mathbb{C}\mathbb{P}^\nabla$ -algebras. Let us restrict ourselves to the case of one generator; the proof is similar in the other cases. $F_{\mathbb{C}\mathbb{P}^\nabla}(1)$ is the space generated by the isoclasses of ∇ -compatible posets, with the products m and ∇ . Let A be an algebra with an associative and commutative product m and a second product ∇ , such that:

$$\forall x, y, z \in A, \quad x\nabla(y\nabla z) = (xy)\nabla z.$$

Let $a \in A$; let us prove that there exists a unique morphism $\phi : F_{\mathbb{C}\mathbb{P}^\nabla}(1) \rightarrow A$ for the two products m and ∇ , such that $\phi(\bullet) = a$.

We define $\phi(\llbracket \mathcal{A} \rrbracket)$ for any ∇ -compatible poset \mathcal{A} by induction on its cardinality n . If $n = 1$, then $\phi(\llbracket \mathcal{A} \rrbracket) = a$. If $n > 1$ and \mathcal{A} is not connected, let us put $\mathcal{A} = \mathcal{A}_1 \dots \mathcal{A}_k$ be the decomposition of \mathcal{A} into connected components. We then put:

$$\phi(\llbracket \mathcal{A} \rrbracket) = \phi(\llbracket \mathcal{A}_1 \rrbracket) \dots \phi(\llbracket \mathcal{A}_k \rrbracket).$$

As the product m of \mathcal{A} is associative and commutative, this does not depend on the chosen order on the \mathcal{A}_i , so is well-defined. If $n > 1$ and \mathcal{A} is connected, we write $\mathcal{A} = b\mathcal{A}\nabla r\mathcal{A}$ and put:

$$\phi(\lfloor \mathcal{A} \rfloor) = \phi(\lfloor b\mathcal{A} \rfloor)\nabla\phi(\lfloor r\mathcal{A} \rfloor).$$

Let us prove that this is indeed an algebra morphism for both products. It is immediate for m . Let \mathcal{A} and \mathcal{B} be two ∇ -compatible posets. We put $\mathcal{C} = \mathcal{A}\nabla\mathcal{B}$. If \mathcal{A} is not connected, then $b\mathcal{C} = \mathcal{A}$ and $r\mathcal{C} = \mathcal{B}$. By definition of ϕ , $\phi(\lfloor \mathcal{C} \rfloor) = \phi(\lfloor \mathcal{A} \rfloor)\nabla\phi(\lfloor \mathcal{B} \rfloor)$. If \mathcal{A} is not connected, then $b\mathcal{C} = \mathcal{A}b\mathcal{B}$ and $r\mathcal{C} = r\mathcal{A}$. By definition of ϕ :

$$\begin{aligned} \phi(\lfloor \mathcal{C} \rfloor) &= \phi(\lfloor \mathcal{A}b\mathcal{B} \rfloor)\nabla\phi(\lfloor r\mathcal{A} \rfloor) \\ &= (\phi(\lfloor \mathcal{A} \rfloor)\phi(\lfloor b\mathcal{B} \rfloor))\nabla\phi(\lfloor r\mathcal{A} \rfloor) \\ &= \phi(\lfloor \mathcal{A} \rfloor)\nabla(\phi(\lfloor b\mathcal{B} \rfloor)\nabla\phi(\lfloor r\mathcal{A} \rfloor)) \\ &= \phi(\lfloor \mathcal{A} \rfloor)\nabla\phi(\lfloor \mathcal{B} \rfloor). \end{aligned}$$

So the morphism ϕ exists. As $F_{\mathbb{C}\mathbb{P}^\nabla(1)}$ is generated for the products m and ∇ by \cdot , the unicity is immediate. \square

6.3. Comparison of WN posets and ∇ -compatible posets. The preceding observations on WN posets and ∇ -compatible posets gives the following decompositions:

- (1) (a) If \mathcal{A} is a non connected WN poset, it can be written as $\mathcal{A} = \mathcal{A}_1 \dots \mathcal{A}_k$, where $\mathcal{A}_1, \dots, \mathcal{A}_k$ are connected WN posets. This decomposition is unique, up to the order of the factors.
- (b) If \mathcal{A} is connected and different from \cdot , it can be uniquely written as $\mathcal{A} = \mathcal{A}_1 \downarrow \mathcal{A}_2$, where \mathcal{A}_1 is a WN poset, and \mathcal{A}_2 is a WN poset, non connected or equal to \cdot .
- (2) (a) If \mathcal{A} is a non connected ∇ -compatible poset, it can be written as $\mathcal{A} = \mathcal{A}_1 \dots \mathcal{A}_k$, where $\mathcal{A}_1, \dots, \mathcal{A}_k$ are connected ∇ -compatible posets. This decomposition is unique, up to the order of the factors.
- (b) If \mathcal{A} is connected and different from \cdot , it can be uniquely written as $\mathcal{A} = \mathcal{A}_1 \nabla \mathcal{A}_2$, where \mathcal{A}_1 is a ∇ -compatible poset, and \mathcal{A}_2 is a ∇ -compatible poset, non connected or equal to \cdot .

Consequently, the species $\mathbb{W}\mathbb{N}\mathbb{P}$ and $\mathbb{C}\mathbb{P}^\nabla$ are isomorphic (as species, not as operads). An isomorphism is inductively defined by a bijection θ from $\mathbb{W}\mathbb{N}\mathbb{P}_A$ to $\mathbb{C}\mathbb{P}_A^\nabla$ by:

- $\theta(\cdot_a) = \cdot_a$.
- If $\mathcal{A} = \mathcal{A}_1 \dots \mathcal{A}_k$ is not connected, $\theta(\mathcal{A}) = \theta(\mathcal{A}_1) \dots \theta(\mathcal{A}_k)$.
- If $\mathcal{A} = \mathcal{A}_1 \downarrow \mathcal{A}_2$ is connected, with \mathcal{A}_2 non connected or equal to 1, then $\theta(\mathcal{A}) = \theta(\mathcal{A}_1)\nabla\theta(\mathcal{A}_2)$.

For example, if \mathcal{A} is a WN poset of cardinality ≤ 4 , then $\theta(\mathcal{A}) = \mathcal{A}$, except if $\lfloor \mathcal{A} \rfloor = \uparrow^4$. Moreover:

$$\theta\left(\begin{matrix} 2 \\ 1 \end{matrix} \uparrow^4 \begin{matrix} 3 \\ 3 \end{matrix}\right) = \theta(\mathbf{1}_1^2 \cdot_3 \downarrow \cdot_4) = \mathbf{1}_1^2 \cdot_3 \nabla \cdot_4 = {}_1^2 \mathcal{V}_3^4.$$

As a consequence, the sequences $(\dim(\mathbb{W}\mathbb{N}\mathbb{P}_{\{1, \dots, n\}}))_{n \geq 1}$ and $(\dim(\mathbb{C}\mathbb{P}_{\{1, \dots, n\}}^\nabla))_{n \geq 1}$ are both equal to sequence A048172 of the OEIS [11]; considering the isoclasses, the sequences $(\dim(F_{\mathbb{W}\mathbb{N}\mathbb{P}}(1)_n))_{n \geq 1}$ and $(\dim(F_{\mathbb{C}\mathbb{P}^\nabla}(1)_n))_{n \geq 1}$ are both equal to sequence A003430 of the OEIS.

REFERENCES

- [1] M. Aguiar, S. Mahajan, *Monoidal functors, species and Hopf algebras*, CRM Monograph Series **29**, Amer. Math. Soc. (2010).
- [2] F. Chapoton, M. Livernet, *Pre-Lie algebras and the rooted trees operad*, Inten. Math. Res. Notices **8**, 395–408 (2001).
- [3] L. Foissy, *Algebraic structures on double and plane posets*, J. Algebraic Combin. **37**, 39–66 (2013).

- [4] M. Livernet, *A rigidity theorem for pre-Lie algebras*, J. Pure and Appl. Algebra **2017** No1, 1–18 (2006).
- [5] J.-L. Loday, *Scindement d'associativité et algèbres de Hopf*, Sémin. Congr. **9**, 155–172 (2004).
- [6] J.-L. Loday, B. Vallette, *Algebraic operads*, Grundlehren Math. Wiss. **346**, Springer Verlag (2012).
- [7] J.-L. Loday, M. Ronco, *On the structure of cofree Hopf algebras*, J. Reine Angew. Math. **592**, 123–155 (2006).
- [8] C. Malvenuto, C. Reutenauer, *A self-paired Hopf algebra on double posets and a Littlewood-Richardson rule*, J. Combin. Theory Ser. A, **118** No4, 1322–1333 (2011).
- [9] M. A. Mendez, *Set operads in combinatorics and computer science*, Springer Briefs in Mathematics, DOI 10.1007/978-3-319-11713-3, Springer (2015).
- [10] W. Schmitt, *Incidence Hopf algebras*, J. Pure and Applied Algebra **96**, 299–330 (1994).
- [11] N. J. A. Sloane, *On-line Encyclopedia of Integer Sequences*, available at <http://www.research.att.com/~njas/sequences/Seis.html>
- [12] R. Stanley, *Enumerative Combinatorics, Vol. 2*, Cambridge Studies in Advanced Mathematics **62**, Cambridge University Press (1999).

IRMA, 10 RUE DU GÉNÉRAL ZIMMER, 67084 STRASBOURG CEDEX, FRANCE
E-mail address: fauvet@math.unistra.fr

UNIVERSITÉ DU LITTORAL - CÔTE D'OPALE, CALAIS
E-mail address: Loic.Foissy@lmpa.univ-littoral.fr

UNIVERSITÉ BLAISE PASCAL, C.N.R.S.-UMR 6620, 3 PLACE VASARÉLY, CS 60026, 63178 AUBIÈRE, FRANCE
E-mail address: manchon@math.univ-bpclermont.fr
URL: <http://math.univ-bpclermont.fr/~manchon/>