

Counting the number of isosceles triangles in rectangular regular grids

Chai Wah Wu

IBM T. J. Watson Research Center

P. O. Box 218, Yorktown Heights, New York 10598, USA

e-mail: chaiwahwu@member.ams.org

February 6, 2017

Abstract

In general graph theory, the only relationship between vertices are expressed via the edges. When the vertices are embedded in an Euclidean space, the geometric relationships between vertices and edges can be interesting objects of study. We look at the number of isosceles triangles where the vertices are points on a regular grid and show that they satisfy a recurrence relation when the grid is large enough. We also derive recurrence relations for the number of acute, obtuse and right isosceles triangles.

1 Introduction

In general graph theory, the relationship between vertices are expressed via the edges of the graph. In geometric graph theory, the vertices and edges have geometric attributes that are important as well. For instance, a random geometric graph is constructed by embedding the vertices randomly in some metric space and connecting two vertices by an edge if and only if they are close enough to each other. When the vertices lie in an Euclidean space, the edges of vertices can form geometric objects such as polygons. In [Nara et al., 2003], the occurrence of polygons is studied. In [Bautista-Santiago et al., 2013] the number of nontrivial triangles is studied. In this note, we consider this problem when the vertices are arranged on a regular grid. The study of the abundance (or sparsity) of such subgraphs or network motifs [Milo et al., 2002] is important in the characterization of complex networks.

Consider an n by k rectangular regular grid G with $n, k \geq 2$. A physical manifestation of this pattern, called geoboard, is useful in teaching elementary geometric concepts [Carroll, 1992]. Let 3 distinct points be chosen on the grid such that they form the vertices of a triangle with nonzero area (i.e. the points are not collinear). In OEIS sequences A271910, A271911, A271912, A271913, A271915 [OEIS], the number of such triangles that are isosceles are listed for various k and n . Neil Sloane made the conjecture that for a fixed

$n \geq 2$, the number of isosceles triangles in an n by k grid, denoted as $a_n(k)$, satisfies the recurrence relation $a_n(k) = 2a_n(k-1) - 2a_n(k-3) + a_n(k-4)$ for $k > K(n)$ for some number $K(n)$. The purpose of this note is to show that this conjecture is true and give an explicit form of $K(n)$. In particular, we show that $K(n) = (n-1)^2 + 3$ if n is even and $K(n) = (n-1)^2 + 2$ if n is odd and that this is the best possible value for $K(n)$.

2 Counting isosceles triangles

We first start with some simple results:

Lemma 1. *If x, y, u, w are integers such that $0 < x, u \leq n$ and $y > \frac{n^2}{2}$, then $x^2 + y^2 = u^2 + w^2$ implies that $x = u$ and $y = w$.*

Proof. If $y \neq w$, then $|y^2 - w^2| \geq 2y - 1 > n^2 - 1$. On the other hand, $|y^2 - w^2| = |u^2 - x^2| < n^2$, a contradiction. \square

Lemma 2. *If x, y, u, w are integers such that $0 \leq x, u \leq n$ and $y > \frac{n^2+1}{2}$, then $x^2 + y^2 = u^2 + w^2$ implies that $x = u$ and $y = w$.*

Proof. If $y \neq w$, then $|y^2 - w^2| \geq 2y - 1 > n^2$. On the other hand, $|y^2 - w^2| = |u^2 - x^2| \leq n^2$, a contradiction. \square

Lemma 3.

$$2 \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} n - 2m = \left\lfloor \frac{(n-1)^2}{2} \right\rfloor$$

Proof. If n is odd, then

$$2 \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} n - 2m = 2 \sum_{m=1}^{\frac{n-1}{2}} n - 2m = n(n-1) - 2 \frac{n-1}{2} \frac{n+1}{2} = \frac{(n-1)^2}{2}$$

If n is even, then

$$2 \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} n - 2m = 2 \sum_{m=1}^{\frac{n-2}{2}} n - 2m = n(n-2) - 2 \frac{n-2}{2} \frac{n}{2} = \frac{(n-1)^2 - 1}{2} = \left\lfloor \frac{(n-1)^2}{2} \right\rfloor$$

\square

Our main result is the following:

Theorem 1. *Let $a_n(k)$ be the number of isosceles triangles of nonzero area formed by 3 distinct points in an n by k grid. Then $a_n(k) = 2a_n(k-1) - 2a_n(k-3) + a_n(k-4)$ for $k > (n-1)^2 + 3$.*

Proof. for $k > 2$, let the n by k array be decomposed into 3 parts consisting of the first column, the middle part (of size n by $k - 2$) and the last column denoted as p_1 , p_2 and p_3 respectively.

Let $b_n(k)$ be the number of isosceles triangles in the n by k array with vertices in the last column p_3 and let $c_n(k)$ be the number of isosceles triangles in the n by k array with vertices in the first and last columns p_1 and p_3 . It is clear that $b_n(k) = a_n(k) - a_n(k - 1)$. Furthermore, $b_n(k) = b_n(k - 1) + c_n(k)$. Let us partition the isosceles triangles corresponding to $c_n(k)$ into 2 groups, $A(k)$ and $B(k)$ where $A(k)$ are triangles with all 3 vertices in p_1 or p_3 and $B(k)$ are triangles with a vertex in each of p_1 , p_2 and p_3 . Since $k > n$, all triangles in $A(k)$ must be of the form where the two vertices in p_1 (resp. p_3) are an even number of rows apart and the third vertex is in p_3 (resp. p_1) in the middle row between them. Since $k > (n - 1)^2 + 1$, these triangles are all acute (we'll revisit this later). Let us count how many such triangles there are. There are $n - 2m$ pairs of vertices which are $2m$ rows apart, for $1 \leq m \leq \lfloor \frac{n-1}{2} \rfloor$. Thus the total number of triangles in $A(k)$ is

$$2 \sum_{m=1}^{\lfloor \frac{n-1}{2} \rfloor} n - 2m = \left\lfloor \frac{(n-1)^2}{2} \right\rfloor$$

by Lemma 3. Next we consider the isosceles triangles in $B(k)$. Let e_1 be the edge between the vertex in p_1 and the vertex in p_2 and e_2 be the edge between the vertex in p_2 and the vertex in p_3 and e_3 be the edge between the vertices in p_1 and p_3 . There are 2 cases. In case 1, the length of e_1 is equal to the length of e_2 and is expressed as $x^2 + y^2 = u^2 + w^2$ with $0 \leq x, u \leq n - 1$ and $y + w = k - 1$. Without loss of generality, we pick y to be the larger of y and w , i.e., $y \geq \frac{k}{2}$. This is illustrated in Fig. 1.

Since $k > (n - 1)^2 + 1$, Lemma 2 implies that $x = u$ and $y = w$ and this is only possible if k is odd and the vertex in p_1 and in p_3 must be at the same row and p_2 in a different row. Note that such triangles are right triangles for $n = 2$ and obtuse for $n > 2$. In case 2, the length of e_3 is equal to the length of either e_1 or e_2 . Lemma 2 shows that in this case $y = k - 1$ which is not possible. This implies that $c_n(k) = P(n, 2) + \lfloor \frac{(n-1)^2}{2} \rfloor = n(n - 1) + \lfloor \frac{(n-1)^2}{2} \rfloor$ if k is odd and $c_n(k) = \lfloor \frac{(n-1)^2}{2} \rfloor$ if k is even. Thus we have $a_n(k) = a_n(k - 1) + b_n(k) = a_n(k - 1) + b_n(k - 1) + c_n(k) = a_n(k - 1) + a_n(k - 1) - a_n(k - 2) + c_n(k) = 2a_n(k - 1) - a_n(k - 2) + c_n(k)$. Since $k - 2 > n$ and $k - 2 > (n - 1)^2 + 1$, we can apply the same analysis to find that $c_n(k - 2) = c_n(k)$. Since $a_n(k - 2) = 2a_n(k - 3) - a_n(k - 4) + c_n(k - 2)$, we get $a_n(k) = 2a_n(k - 1) - 2a_n(k - 3) + a_n(k - 4) - c_n(k - 2) + c_n(k) = 2a_n(k - 1) - 2a_n(k - 3) + a_n(k - 4)$. \square

It is clear that $a_n(k) = a_k(n)$. The above argument also shows that:

Theorem 2. *Suppose that $k > (n - 1)^2 + 1$. Then $a_n(k) = 2a_n(k - 1) - a_n(k - 2) + n(n - 1) + \lfloor \frac{(n-1)^2}{2} \rfloor$ if k is odd and $a_n(k) = 2a_n(k - 1) - a_n(k - 2) + \lfloor \frac{(n-1)^2}{2} \rfloor$ if k is even.*




Figure 1: Illustrating a triangle in $B(k)$.

3 Obtuse, acute and right isosceles

As noted above, the triangles in $A(k)$ are acute and the triangles in $B(k)$ are right for $n = 2$ and obtuse for $n > 2$. This implies that Theorem 1 is also true when restricted to the set of acute isosceles triangles and restricted obtuse or right isosceles triangles if $n > 2$. As for Theorem 2, we have the following similar results:

Theorem 3. *Let $a_n^a(k)$ be the number of acute isosceles triangles of nonzero area formed by 3 distinct points in an n by k grid. Then $a_n^a(k) = 2a_n^a(k-1) - a_n^a(k-2) + \lfloor \frac{(n-1)^2}{2} \rfloor$ for $k > (n-1)^2 + 1$.*

Theorem 4. *Let $a_n^o(k)$ be the number of obtuse isosceles triangles of nonzero area formed by 3 distinct points in an n by k grid. Suppose $k > \max(3, (n-1)^2 + 1)$. Then $a_n^o(k) = 2a_n^o(k-1) - a_n^o(k-2) + n(n-1)$ if k is odd and $a_n^o(k) = 2a_n^o(k-1) - a_n^o(k-2)$ if k is even.*

Theorem 5. *Let $a_n^r(k)$ be the number of right isosceles triangles of nonzero area formed by 3 distinct points in an n by k grid. Then $a_n^r(k) = 2a_n^r(k-1) - a_n^r(k-2)$ for $k > \max(3, (n-1)^2 + 1)$.*

4 Pythagorean triples and a small improvement

In Theorem 1 we have given an explicit form for the bound $K(n)$ in the conjecture described in Section 1. We next show that for odd n , this bound can be reduced by 1. Let us consider

the case $k = (n - 1)^2 + 1$. Consider the triangles in $B(k)$. Again case 2 is impossible. For case 1, the length of e_1 is equal to the length of e_2 and is expressed as $x^2 + y^2 = u^2 + w^2$ with $0 \leq x, u \leq n - 1$ and $y \geq \frac{k}{2}$ and $y \geq w$. If $x > 0$, this implies that $u > 0$ and we can apply Lemma 1 to show that the only isosceles triangles in $B(k)$ are as in Theorem 1. Suppose $x = 0$. We can eliminate $y = w$ since this results in $u = 0$ and a collinear set of vertices. For $n = 2, k = 2$, it is clear that $w = 0$ and $y = w + 1 = 1$. For $n > 2$, since $u \leq n - 1$, this again means that $y = w + 1$ as otherwise $y^2 - w^2 \geq 4y - 4 \geq 2k - 4 > (n - 1)^2$. This implies that k must necessarily be even such that a, b, c are integers forming a Pythagorean triple satisfying $a^2 = b^2 + c^2$ where $a = \frac{k}{2}$, $b = a - 1$ and $c = \sqrt{a^2 - b^2} = \sqrt{k - 1}$. If n is odd, then $k = (n - 1)^2 + 1$ is odd, and the case of $x = 0$ cannot occur. Thus Theorem 1 can be improved to:

Theorem 6. *Suppose n is odd. Then $a_n(k) = 2a_n(k - 1) - 2a_n(k - 3) + a_n(k - 4)$ for $k > (n - 1)^2 + 2$.*

We can rewrite this in an inhomogeneous form of lower degree:

Theorem 7. *Suppose that n is odd and $k > (n - 1)^2$. Then $a_n(k) = 2a_n(k - 1) - a_n(k - 2) + n(n - 1) + \lfloor \frac{(n-1)^2}{2} \rfloor$ if k is odd and $a_n(k) = 2a_n(k - 1) - a_n(k - 2) + \lfloor \frac{(n-1)^2}{2} \rfloor$ if k is even.*

Again, Theorem 6 is still valid when restricted to acute isosceles triangles. When $n = 3$ and $k = (n - 1)^2 + 1 = 5$, two of the triangles in $B(k)$ are right isosceles and for $n > 3$ and $k = (n - 1)^2 + 1$, all triangles in $B(k)$ are obtuse triangles. Thus we have

Theorem 8. *Suppose n is odd. Then $a_n^a(k) = 2a_n^a(k - 1) - 2a_n^a(k - 3) + a_n^a(k - 4)$ for $k > (n - 1)^2 + 2$. Furthermore, $a_n^o(k) = 2a_n^o(k - 1) - 2a_n^o(k - 3) + a_n^o(k - 4)$ and $a_n^r(k) = 2a_n^r(k - 1) - 2a_n^r(k - 3) + a_n^r(k - 4)$ for $k > \max(7, (n - 1)^2 + 2)$.*

Similiarly we have:

Theorem 9. *Suppose n is odd and $k > \max(5, (n - 1)^2)$. Then $a_n^o(k) = 2a_n^o(k - 1) - a_n^o(k - 2) + n(n - 1)$ for k odd and $a_n^o(k) = 2a_n^o(k - 1) - a_n^o(k - 2)$ for k even.*

When n is even and $k = (n - 1)^2 + 1$, the above analysis shows that there are 4 extra isosceles triangles in $B(k)$ due to the case $x = 0$. These triangles are right triangles for $n = 2$ and obtuse for $n > 2$. This is illustrated in Fig. 2 for the case $k = 10, n = 4$. This means that when restricted to acute isosceles triangles (or right isosceles triangles with $n > 2$), the condition that n is odd is not necessary in Theorem 6. In addition Theorems 3 and 5 can be improved to:

Theorem 10. *Suppose $k > (n - 1)^2$. Then $a_n^a(k) = 2a_n^a(k - 1) - a_n^a(k - 2) + \lfloor \frac{(n-1)^2}{2} \rfloor$.*

This can be rewritten in homogeneous form as:

Corollary 1. *Suppose $k > (n - 1)^2 + 1$. Then $a_n^a(k) = 3a_n^a(k - 1) - 3a_n^a(k - 2) + a_n^a(k - 3)$.*




Figure 2: 4 obtuse isosceles triangles in $B(k)$ corresponding to $x = 0$ for the case $k = 10$, $n = 4$.

Proof. From Theorem 10, we have $a_n^a(k) = 2a_n^a(k-1) - a_n^a(k-2) + \lfloor \frac{(n-1)^2}{2} \rfloor$ and $a_n^a(k-1) = 2a_n^a(k-2) - a_n^a(k-3) + \lfloor \frac{(n-1)^2}{2} \rfloor$. Subtracting these two equations and combining terms we reach the conclusion. \square

A similar homogeneous linear recurrence for right isosceles triangles is:

Theorem 11. *Suppose $n > 2$ and $k > \max(5, (n-1)^2)$. Then $a_n^r(k) = 2a_n^r(k-1) - a_n^r(k-2)$.*

Similarly, we have

Theorem 12. *Suppose that n is even and $k = (n-1)^2 + 1$. Then $a_n(k) = 2a_n(k-1) - a_n(k-2) + \lfloor \frac{(n-1)^2}{2} \rfloor + 4$.*

Theorem 13. *Suppose that $n > 2$ is even and $k = (n-1)^2 + 1$. Then $a_n^o(k) = 2a_n^o(k-1) - a_n^o(k-2) + 4$.*

Because of the recurrence relation in Theorem 1, the generating functions for $a_n(k)$ for $k \geq 1$ will be of the form $\frac{p_n(x)}{(x-1)^3(x+1)}$. The polynomials $p_n(x)$ for the first few values of n are shown in Table 1.

Similarly, the polynomials for obtuse, acute and right isosceles triangles are shown in Tables 2-4. We see that the denominator for the generating function of $a_n^a(k)$ can be reduced to $(x-1)^3$ corresponding to the recurrence relation in Corollary 1. Similarly, we see that the denominator for the generating function of $a_n^r(k)$ can be reduced to $(x-1)^2$ corresponding to the recurrence relation in Theorem 5.

5 Optimal value for $K(n)$

Theorem 12 shows that $(n-1)^2 + 3$ is the best value for $K(n)$ when n is even. Next we show that $(n-1)^2 + 2$ is the best value for $K(n)$ when n is odd.

Consider the case where $n > 2$ is odd and $k = (n-1)^2$. Then k is even and by setting $y = \frac{k}{2}$, $w = \frac{k}{2} - 1$, $x = 1$, $u = n-1$, we get $x^2 + y^2 = u^2 + w^2$ where $x \neq u$ and $y \neq w$ and

n	$p_n(x)$
2	$2x(2x^2 - x - 2)$
3	$2x(2x^4 + 4x^3 + 2x^2 - 8x - 5)$
4	$4x(x^{10} - x^8 + 2x^6 + x^5 + 4x^4 + 4x^3 - 3x^2 - 9x - 4)$
5	$4x(x^{16} - x^{14} + 2x^{10} + 2x^9 - x^8 - x^7 + 5x^6 + 6x^5 + 6x^4 + x^3 - 8x^2 - 15x - 6)$
6	$2x(2x^{26} - 2x^{24} + 2x^{22} - 2x^{20} + 6x^{16} - 4x^{14} + 2x^{13} - 2x^{11} + 6x^{10} + 12x^9 - 6x^7 + 26x^6 + 24x^5 + 6x^4 - 8x^3 - 30x^2 - 43x - 16)$
7	$2x(2x^{36} - 2x^{34} + 2x^{28} + 2x^{26} - 4x^{24} + 4x^{22} - 4x^{20} + 4x^{19} - 2x^{17} + 10x^{16} - 6x^{14} + 4x^{13} + 2x^{12} - 2x^{11} + 10x^{10} + 18x^9 + 2x^8 + 4x^7 + 40x^6 + 22x^5 - 2x^4 - 20x^3 - 44x^2 - 58x - 21)$
8	$4x(x^{50} - x^{48} + x^{46} - x^{44} + 3x^{36} - 2x^{34} - x^{32} + 2x^{28} + 2x^{26} + x^{25} - 3x^{24} - x^{23} + 3x^{22} + x^{21} - 4x^{20} + 4x^{19} + x^{18} - 3x^{17} + 7x^{16} - 3x^{14} + 5x^{13} + 3x^{12} - 2x^{11} + 6x^{10} + 13x^9 + 10x^8 + 7x^7 + 19x^6 + 9x^5 - 7x^4 - 17x^3 - 29x^2 - 37x - 13)$

Table 1: Numerator of generating functions of $a_n(k)$.

n	$p_n^o(x)$
2	$-2x^4$
3	$-2x^4(x^2 + 2)$
4	$2x^3(2x^8 - 2x^6 - x^5 - 2x^3 + 2x^2 - 3x - 2)$
5	$2x(2x^{16} - 2x^{14} + 4x^{10} + x^9 - 4x^8 - 4x^7 - x^5 + 4x^4 - 6x^3 - 3x^2 - 1)$
6	$2x(2x^{26} - 2x^{24} + 2x^{22} - 2x^{20} + 6x^{16} - 6x^{14} + 2x^{13} - 3x^{11} + 6x^{10} + 2x^9 - 6x^8 - 7x^7 + 4x^6 + 2x^4 - 9x^3 - 4x^2 - 2)$
7	$2x(2x^{36} - 2x^{34} + 2x^{28} + 2x^{26} - 4x^{24} + 4x^{22} - 4x^{20} + 2x^{19} + 10x^{16} - 2x^{15} - 10x^{14} + 5x^{13} - 8x^{11} + 8x^{10} + 3x^9 - 8x^8 - 6x^7 + 8x^6 - 3x^5 - 11x^3 - 4x^2 - x - 4)$
8	$2x(2x^{50} - 2x^{48} + 2x^{46} - 2x^{44} + 6x^{36} - 4x^{34} - 2x^{32} + 4x^{28} + 2x^{26} + 2x^{25} - 6x^{24} - 2x^{23} + 8x^{22} + 2x^{21} - 8x^{20} + 2x^{19} + 16x^{16} - 5x^{15} - 16x^{14} + 10x^{13} - 15x^{11} + 10x^{10} + 4x^9 - 4x^8 - 5x^7 + 6x^6 - 6x^5 - 2x^4 - 13x^3 - 4x^2 - 2x - 6)$

Table 2: Numerator of generating functions $\frac{p_n^o(x)}{(x-1)^3(x+1)}$ of $a_n^o(k)$.

n	$p_n^a(x)$
2	0
3	$2x^2(x+1)(3x-4)$
4	$2x^2(x+1)(2x^4-x^3+3x^2+3x-9)$
5	$2x^2(x+1)(2x^7-2x^6+x^5+5x^4+3x^3-x^2+3x-15)$
6	$2x^2(x+1)(2x^{12}-2x^{11}+2x^{10}-2x^9+7x^7-5x^6+x^5+15x^4+2x^3-6x^2+2x-22)$
7	$2x^2(x+1)(2x^{17}-2x^{16}+2x^{13}+2x^{12}-4x^{11}+4x^{10}-x^9-x^8+11x^7-7x^6+10x^5+14x^4+2x^3-13x^2+2x-30)$
8	$2x^2(x+1)(2x^{24}-2x^{23}+2x^{22}-2x^{21}+6x^{17}-4x^{16}-2x^{15}+4x^{13}+4x^{12}-7x^{11}+9x^{10}-3x^9-x^8+16x^7+10x^5+12x^4+x^3-21x^2+3x-39)$

Table 3: Numerator of generating functions $\frac{p_n^a(x)}{(x-1)^3(x+1)}$ of $a_n^a(k)$.

n	$p_n^r(x)$
2	$2x(x-1)(x+1)(x+2)$
3	$2x(x-1)(x+1)(x^3+2x^2+4x+5)$
4	$2x(x-1)(x+1)(x^5+2x^4+4x^3+6x^2+9x+8)$
5	$2x(x-1)(x+1)(x^7+2x^6+4x^5+6x^4+9x^3+12x^2+15x+11)$
6	$2x(x-1)(x+1)(x^9+2x^8+4x^7+6x^6+9x^5+12x^4+16x^3+20x^2+21x+14)$
7	$2x(x-1)(x+1)(x^{11}+2x^{10}+4x^9+6x^8+9x^7+12x^6+16x^5+20x^4+25x^3+29x^2+27x+17)$
8	$2x(x-1)(x+1)(x^{13}+2x^{12}+4x^{11}+6x^{10}+9x^9+12x^8+16x^7+20x^6+25x^5+30x^4+36x^3+38x^2+33x+20)$

Table 4: Numerator of generating functions $\frac{p_n^r(x)}{(x-1)^3(x+1)}$ of $a_n^r(k)$.

$y + w = k - 1$. This corresponds to 4 additional isosceles triangles in $B(k)$. These triangles are right triangles for $n = 3$ (Fig. 3) and obtuse for $n > 3$.




Figure 3: 4 right isosceles triangles in $B(k)$ corresponding to $x = 1$ for the case $k = 4$, $n = 3$.

Theorem 14. *Suppose that n is odd and $k = (n - 1)^2$. Then $a_n(k) = 2a_n(k - 1) - a_n(k - 2) + \lfloor \frac{(n-1)^2}{2} \rfloor + 4$.*

Theorem 15. *Suppose that $n > 3$ is odd and $k = (n - 1)^2$. Then $a_n^o(k) = 2a_n^o(k - 1) - a_n^o(k - 2) + 4$.*

This means that $K(n) = (n - 1)^2 + 3$ for n is even and $K(n) = (n - 1)^2 + 2$ for n is odd are the best possible values for $K(n)$ in the conjecture in Section 1 as expressed in Theorems 1 and 6. They are also optimal when restricted to the set of obtuse isosceles and $n > 3$. In addition, $a_n(k) = 2a_n(k - 1) - 2a_n(k - 3) + a_n(k - 4) - 4$ if $k = (n - 1)^2 + 3$ and n is even or if $k = (n - 1)^2 + 2$ and n is odd. This is due to the fact that for these values of n and k , $c_n(k - 2) = c_n(k) + 4$.

6 Subsets of points on a regular grid containing the vertices of isosceles triangles

Consider an n by k regular grid. Let $S(n, k) = r$ be defined as the smallest number r such that any r points in the grid contains the vertices of an isosceles triangle of nonzero area. This is equivalent to finding the constellation with the largest number of points $T(n, k)$ such that no three points form an isosceles triangle as $S(n, k) = T(n, k) + 1$. The values of $T(n, k)$ for small n and k are listed in OEIS A271914 where it was conjectured that $T(n, k) \leq n + k - 1$. The next Lemma shows that this conjecture is true for $k = 1, 2$.

Lemma 4. *$T(n, k)$ satisfies the following properties:*

1. $T(n, k) = T(k, n)$. If $m \leq n$, then $T(m, k) \leq T(n, k)$.
2. $T(n, k) \geq \max(n, k)$,
3. $T(n, 1) = n$,

4. (subadditivity) $T(n, k + m) \leq T(n, k) + T(n, m)$,

5. $T(n, 2) = n$ for $n > 3$,

6. Suppose $n > 4$. Then $T(n, 3) \geq n + 1$ if n is odd and $T(n, 3) \geq n + 2$ if n is even.

Proof. Property 1 is trivial. Properties 2 and 3 are clear, by considering points all in one row (or column) only. To show property 4, suppose $T(n, k) + T(n, m) + 1$ points are chosen in an n by $k + m$ grid, and consider a decomposition into an n by k grid and an n by m grid. Either the n by k grid has more than $T(n, k)$ points or the n by m grid has more than $T(n, m)$ points. In either case, there is an isosceles triangle. Property 4 implies that $T(n + 2, 2) \leq T(n, 2) + T(2, 2) = T(n, 2) + 2$ and property 2 implies $T(n, 2) \geq n$ for $n \geq 2$. Thus property 5 follows from properties 2 and 4. Suppose $n > 4$. For n odd, the points $\{(1, i) : 2 \leq i \leq n\}$ along with the points $(2, 1)$ and $(3, 1)$ shows that $T(n, 3) \geq n + 1$ (Fig. 4). If n is even, the points $\{(1, i) : 2 \leq i \leq n - 1\}$ along with the points $(2, 1)$, $(3, 1)$, $(2, n)$ and $(3, n)$ shows that $T(n, 3) \geq n + 2$ (Fig. 5). \square




Figure 4: A constellation of $n + 1$ points for which there are no 3 points forming an isosceles triangle for the case when $n > 4$ is odd.




Figure 5: A constellation of $n + 2$ points for which there are no 3 points forming an isosceles triangle for the case when $n > 4$ is even.

Note that for n odd, the constellation in Fig. 5 for $n - 1$ which has $n - 1 + 2 = n + 1$ points will also show that $T(n, 3) \geq n + 1$. We conjecture the following:

Conjecture 1. *If n is even, then $T(n, k) \leq n + k - 2$ for $k \geq 2n$.*

Conjecture 2. *For $n > 4$, $T(n, 3) = n + 1$ if n is odd and $T(n, 3) = n + 2$ if n is even.*

Note that by Lemma 4 Conjecture 1 is true for $n = 2$.

References

- [Carroll, 1992] Carroll, J., “Using the geoboard for teaching primary mathematics”. In Horne, Marj and Supple, Marjorie (eds). *Mathematics : meeting the challenge*. Brunswick, Vic.: Mathematical Association of Victoria. pp. 283-288, 1992.
- [OEIS] The on-line encyclopedia of integer sequences (<http://oeis.org/>), founded in 1964 by N. J. A. Sloane.
- [Nara et al., 2003] C. Nara, T. Sakai and J. Urrutia, “Maximal number of edges in geometric graphs without convex polygons”. in *Discrete and Computational Geometry*, Edited by J. Akiyama and M. Kano, Springer Berlin/Heidelberg, 2003. 215–220.
- [Bautista-Santiago et al., 2013] C. Bautista-Santiago, M. A. Heredia, C. Huemer, A. Ramrez-Vigueras, C. Seara and J. Urrutia, “On the Number of Edges in Geometric Graphs Without Empty Triangles”. in *Graphs and Combinatorics*, vol. 29, no. 6, pp. 1623-1631, 2013.
- [Milo et al., 2002] R. Milo, S. Shen-Orr, S. Itzkovitz, N. Kashtan, D. Chklovskii, and U. Alon, “Network Motifs: Simple Building Blocks of Complex Networks”. in *Science, American Association for the Advancement of Science*, vol. 298, pp. 824-827, 2002.