

# Properties of distance spaces with power triangle inequalities

DANIEL J. GREENHOE

*Abstract:* Metric spaces provide a framework for analysis and have several very useful properties. Many of these properties follow in part from the *triangle inequality*. However, there are several applications in which the triangle inequality does not hold but in which we may still like to perform analysis. This paper investigates what happens if the triangle inequality is removed all together, leaving what is called a *distance space*, and also what happens if the triangle inequality is replaced with a much more general two parameter relation, which is herein called the “*power triangle inequality*”. The power triangle inequality represents an uncountably large class of inequalities, and includes the *triangle inequality*, *relaxed triangle inequality*, and *inframetric inequality* as special cases. The power triangle inequality is defined in terms of a function that is herein called the *power triangle function*. The power triangle function is itself a *power mean*, and as such is *continuous* and *monotone* with respect to its exponential parameter, and also includes the operations of *maximum*, *minimum*, *mean square*, *arithmetic mean*, *geometric mean*, and *harmonic mean* as special cases.

*2010 Mathematics Subject Classification* 54E25 (primary); 54A05,54A20 (secondary)

*Keywords:* metric space, distance space, semimetric space, quasi-metric space, triangle inequality, relaxed triangle inequality, inframetric, arithmetic mean, means square, geometric mean, harmonic mean, maximum, minimum, power mean

## Contents

<b>1</b>	<b>Introduction and summary</b>	<b>2</b>
<b>2</b>	<b>Standard definitions</b>	<b>4</b>
2.1	Standard sets . . . . .	4
2.2	Relations . . . . .	4
2.3	Set functions . . . . .	4
2.4	Order . . . . .	5
<b>3</b>	<b>Background: distance spaces</b>	<b>5</b>
3.1	Fundamental structure of distance spaces . . . . .	6
3.1.1	Definitions . . . . .	6
3.1.2	Properties . . . . .	6
3.2	Open sets in distance spaces . . . . .	6
3.2.1	Definitions . . . . .	6
3.2.2	Properties . . . . .	7
3.3	Sequences in distance spaces . . . . .	9
3.3.1	Definitions . . . . .	9
3.3.2	Properties . . . . .	9
3.4	Examples . . . . .	12



<b>4 Distance spaces with power triangle inequalities</b>	<b>15</b>
4.1 Definitions . . . . .	15
4.2 Properties . . . . .	16
4.2.1 Relationships of the power triangle function . . . . .	16
4.2.2 Properties of power distance spaces . . . . .	17
4.3 Examples . . . . .	24
<b>A Topological Spaces</b>	<b>26</b>
<b>B Finite sums</b>	<b>30</b>
B.1 Convexity . . . . .	30
B.2 Power means . . . . .	31
B.3 Inequalities . . . . .	35
<b>C Metric preserving functions</b>	<b>35</b>
<b>Bibliography</b>	<b>37</b>
<b>Reference Index</b>	<b>44</b>
<b>Subject Index</b>	<b>45</b>

## 1 Introduction and summary

*Metric spaces* provide a framework for analysis and have several very useful properties. Many of these properties follow in part from the *triangle inequality*. However, there are several applications<sup>1</sup> in which the triangle inequality does not hold but in which we would still like to perform analysis. So the questions that naturally follow are:

- Q1. What happens if we remove the *triangle inequality* all together?
- Q2. What happens if we replace the *triangle inequality* with a generalized relation?

A *distance space* is a *metric space* without the *triangle inequality* constraint. Section 3 introduces *distance spaces* and demonstrates that some properties commonly associated with *metric spaces* also hold in any *distance space*:

- |   |                            |
|---|----------------------------|
| D1. $\emptyset$ and $X$ are <i>open</i>                               | (Theorem 3.7 page 7)       |
| D2. the intersection of a finite number of open sets is <i>open</i>   | (Theorem 3.7 page 7)       |
| D3. the union of an arbitrary number of open sets is <i>open</i>      | (Theorem 3.7 page 7)       |
| D4. every Cauchy sequence is <i>bounded</i>                           | (Proposition 3.14 page 9)  |
| D5. any subsequence of a <i>Cauchy</i> sequence is also <i>Cauchy</i> | (Proposition 3.15 page 10) |
| D6. the <i>Cantor Intersection Theorem</i> holds                      | (Theorem 3.18 page 11)     |

<sup>1</sup>references for applications in which the *triangle inequality* may not hold: [Maligranda and Orlicz(1987)] page 54 (“pseudonorm”), [Lin(1998)] (“similarity measures”, Table 6), [Veltkamp and Hagedoorn(2000)] (“shape similarity measures”), [Veltkamp(2001)] (“shape matching”), [Costa et al.(2004)Costa, Castro, Rowstron, and Key] (“network distance estimation”), [Burstein et al.(2005)Burstein, Ulitsky, Tuller, and Chor] page 287 (distance matrices for “genome phylogenies”), [Jiménez and Yukich(2006)] page 224 (“statistical distances”), [Szirmai(2007)] page 388 (“geodesic ball”), [Crammer et al.(2007)Crammer, Kearns, and Wortman] page 326 (“decision-theoretic learning”), [Vitányi(2011)] page 2455 (“information distance”),



The following five properties (M1–M5) *do* hold in any *metric space*. However, the examples from Section 3 listed below demonstrate that the five properties *do not* hold in all *distance spaces*:

M1.	the <i>metric function</i> is <i>continuous</i>	fails to hold in	Example 3.21–Example 3.23
M2.	<i>open balls</i> are <i>open</i>	fails to hold in	Example 3.21 and Example 3.22
M3.	the <i>open balls</i> form a <i>base</i> for a topology	fails to hold in	Example 3.21 and Example 3.22
M4.	the limits of <i>convergent sequences</i> are <i>unique</i>	fails to hold in	Example 3.21
M5.	<i>convergent</i> sequences are <i>Cauchy</i>	fails to hold in	Example 3.22

Hence, Section 3 answers question Q1.

Section 4 begins to answer question Q2 by first introducing a new function, called the *power triangle function* in a *distance space*  $(X, d)$ , as  $\tau(p, \sigma; x, y, z; d) \triangleq 2\sigma \left[ \frac{1}{2}d^p(x, z) + \frac{1}{2}d^p(z, y) \right]^{\frac{1}{p}}$  for some  $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$ .<sup>2</sup> Section 4 then goes on to use this function to define a new relation, called the *power triangle inequality* in  $(X, d)$ , and defined as  $\mathcal{Q}(p, \sigma; d) \triangleq \{(x, y, z) \in X^3 \mid d(x, y) \leq \tau(p, \sigma; x, y, z; d)\}$ .

The *power triangle inequality* is a generalized form of the *triangle inequality* in the sense that the two inequalities coincide at  $(p, \sigma) = (1, 1)$ . Other special values include  $(1, \sigma)$  yielding the *relaxed triangle inequality* (and its associated *near metric space*) and  $(\infty, \sigma)$  yielding the  *$\sigma$ -inframetric inequality* (and its associated  *$\sigma$ -inframetric space*). Collectively, a distance space with a power triangle inequality is herein called a *power distance space* and denoted  $(X, d, p, \sigma)$ .<sup>3</sup>

The *power triangle function*, at  $\sigma = \frac{1}{2}$ , is a special case of the *power mean* with  $N = 2$  and  $\lambda_1 = \lambda_2 = \frac{1}{2}$ . *Power means* have the elegant properties of being *continuous* and *monotone* with respect to a free parameter  $p$ . From this it is easy to show that the *power triangle function* is also *continuous* and *monotone* with respect to both  $p$  and  $\sigma$ . Special values of  $p$  yield operators coinciding with *maximum*, *minimum*, *mean square*, *arithmetic mean*, *geometric mean*, and *harmonic mean*. *Power means* are briefly described in APPENDIX B.2.<sup>4</sup>

Section 4.2 investigates the properties of *power distance spaces*. In particular, it shows for what values of  $(p, \sigma)$  the properties M1–M5 hold. Here is a summary of the results in a *power distance space*  $(X, d, p, \sigma)$ , for all  $x, y, z \in X$ :

(M1)	holds for any $(p, \sigma) \in (\mathbb{R}^* \setminus \{0\}) \times \mathbb{R}^+$ such that $2\sigma = 2^{\frac{1}{p}}$	(Theorem 4.18 page 23)
(M2)	holds for any $(p, \sigma) \in (\mathbb{R}^* \setminus \{0\}) \times \mathbb{R}^+$ such that $2\sigma \leq 2^{\frac{1}{p}}$	(Corollary 4.14 page 21)
(M3)	holds for any $(p, \sigma) \in (\mathbb{R}^* \setminus \{0\}) \times \mathbb{R}^+$ such that $2\sigma \leq 2^{\frac{1}{p}}$	(Corollary 4.12 page 20)
(M4)	holds for any $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$	(Theorem 4.19 page 23)
(M5)	holds for any $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$	(Theorem 4.16 page 21)

APPENDIX A briefly introduces *topological spaces*. The *open balls* of any *metric space* form a *base* for a *topology*. This is largely due to the fact that in a metric space, open balls are *open*. Because of this, in metric spaces it is convenient to use topological structure to define and exploit analytic concepts such as *continuity*, *convergence*, *closed sets*, *closure*, *interior*, and *accumulation point*. For example, in a metric space, the traditional definition of defining continuity using open balls and the topological

<sup>2</sup>where  $\mathbb{R}^*$  is the *set of extended real numbers* and  $\mathbb{R}^+$  is the *set of positive real numbers* (Definition 2.1 page 4)

<sup>3</sup>*power triangle inequality*: Definition 4.4 page 16; *power distance space*: Definition 4.3 page 16; examples of *power distance space*: Definition 4.5 page 16;

<sup>4</sup>*power triangle function*: Definition 4.1 (page 15); *power mean*: Definition B.6 (page 31); power mean is *continuous* and *monotone*: Theorem B.7 (page 31); power triangle function is *continuous* and *monotone*: Corollary 4.6 (page 16); Special values of  $p$ : Corollary 4.7 (page 17), Corollary B.8 (page 34)



definition using open sets, coincide with each other. Again, this is largely because the open balls of a metric space are open.<sup>5</sup>

However, this is not the case for all *distance spaces*. In general, the open balls of a distance space are not open, and they are not a base for a topology. In fact, the open balls of a distance space are a base for a topology if and only if the open balls are open. While the open sets in a distance space do induce a topology, it's open balls may not.<sup>6</sup>

## 2 Standard definitions

### 2.1 Standard sets

**Definition 2.1** Let  $\mathbb{R}$  be the **set of real numbers**. Let  $\mathbb{R}^{\perp} \triangleq \{x \in \mathbb{R} \mid x \geq 0\}$  be the **set of non-negative real numbers**. Let  $\mathbb{R}^+ \triangleq \{x \in \mathbb{R} \mid x > 0\}$  be the **set of positive real numbers**. Let  $\mathbb{R}^* \triangleq \mathbb{R} \cup \{-\infty, \infty\}$  be the set of **extended real numbers**.<sup>7</sup> Let  $\mathbb{Z}$  be the **set of integers**. Let  $\mathbb{N} \triangleq \{n \in \mathbb{Z} \mid n \geq 1\}$  be the **set of natural numbers**. Let  $\mathbb{Z}^* \triangleq \mathbb{Z} \cup \{-\infty, \infty\}$  be the *extended set of integers*.

**Definition 2.2** Let  $X$  be a set. The quantity  $2^X$  is the *power set of  $X$*  such that  

$$2^X \triangleq \{A \subseteq X\} \quad (\text{the set of all subsets of } X).$$

### 2.2 Relations

**Definition 2.3**<sup>8</sup> Let  $X$  and  $Y$  be sets. The **Cartesian product**  $X \times Y$  of  $X$  and  $Y$  is the set  $X \times Y \triangleq \{(x, y) \mid x \in X \text{ and } y \in Y\}$ . An **ordered pair**  $(x, y)$  on  $X$  and  $Y$  is any element in  $X \times Y$ . A **relation**  $\mathcal{R}$  on  $X$  and  $Y$  is any subset of  $X \times Y$  such that  $\mathcal{R} \subseteq X \times Y$ . The set  $2^{X \times Y}$  is the **set of all relations** in  $X \times Y$ . A *relation*  $f \in 2^{X \times Y}$  is a **function** if  $(x, y_1) \in f$  and  $(x, y_2) \in f \implies y_1 = y_2$ . The set  $Y^X$  is the **set of all functions** in  $2^{X \times Y}$ .

### 2.3 Set functions

**Definition 2.4**<sup>9</sup> Let  $2^X$  be the *power set* (Definition 2.2 page 4) of a set  $X$ .

A set  $\mathcal{S}(X)$  is a **set structure** on  $X$  if  $\mathcal{S}(X) \subseteq 2^X$ .

A set structure  $\mathcal{Q}(X)$  is a **paving** on  $X$  if  $\emptyset \in \mathcal{Q}(X)$ .

<sup>5</sup> *open ball*: Definition 3.5 page 6; *metric space*: Definition 4.5 page 16; *base*: Definition A.3 page 27; *topology*: Definition A.1 page 26; *open*: Definition 3.6 page 7; *continuity in topological space*: Definition A.11 page 28; *convergence in distance space*: Definition 3.11 page 9; *convergence in topological space*: Definition A.16 page 28; *closed set*: Definition A.1 page 26; *closure, interior, accumulation point*: Definition A.8 page 27; coincidence in all *metric spaces* and some *power distance spaces*: Theorem 4.15 page 21;

<sup>6</sup> *if and only if* statement: Theorem 3.10 page 8; open sets of a distance space induce a topology: Corollary 3.8 page 8;

<sup>7</sup> [Rana(2002)] pages 385–388 (Appendix A)

<sup>8</sup> [Maddux(2006)] page 4, [Halmos(1960)] pages 26–30, [Suppes(1972)] page 86, [Kelley(1955)] page 10, [Bourbaki(1939)], [Bottazzini(1986)] page 7, [Comtet(1974)] page 4  $\langle |Y^X| \rangle$ ; The notation  $Y^X$  and  $2^{X \times Y}$  is motivated by the fact that for finite  $X$  and  $Y$ ,  $|Y^X| = |Y|^{|X|}$  and  $|2^{X \times Y}| = 2^{|X| \cdot |Y|}$ .

<sup>9</sup> [Molchanov(2005)] page 389, [Pap(1995)] page 7, [Hahn and Rosenthal(1948)] page 254



**Definition 2.5**<sup>10</sup> Let  $\mathcal{Q}(X)$  be a *paving* (Definition 2.4 page 4) on a set  $X$ . Let  $Y$  be a set containing the element 0. A function  $m \in Y^{\mathcal{Q}(X)}$  is a **set function** if  $m(\emptyset) = 0$ .

**Definition 2.6** The *set function* (Definition 2.5 page 5)  $|A| \in \mathbb{Z}^*_{\geq 0}$  is the *cardinality* of  $A$  such that

$$|A| \triangleq \begin{cases} \text{the number of elements in } A & \text{for finite } A \\ \infty & \text{otherwise} \end{cases} \quad \forall A \in \mathcal{2}^X$$

**Definition 2.7** Let  $|X|$  be the *cardinality* (Definition 2.6 page 5) of a set  $X$ . The structure  $\emptyset$  is the **empty set**, and is a set such that  $|\emptyset| = 0$ .

## 2.4 Order

**Definition 2.8**<sup>11</sup> Let  $X$  be a set. A relation  $\leq$  is an **order relation** in  $\mathcal{2}^{X \times X}$  (Definition 2.3 page 4) if

- |  |                         |                  |     |          |
|--|-------------------------|------------------|-----|----------|
| 1. $x \leq x$                                  | $\forall x \in X$       | (reflexive)      | and | preorder |
| 2. $x \leq y$ and $y \leq z \implies x \leq z$ | $\forall x, y, z \in X$ | (transitive)     | and |          |
| 3. $x \leq y$ and $y \leq x \implies x = y$    | $\forall x, y \in X$    | (anti-symmetric) |     |          |

An **ordered set** is the pair  $(X, \leq)$ .

**Definition 2.9**<sup>12</sup> In an *ordered set*  $(X, \leq)$ ,  
 the set  $[x : y] \triangleq \{z \in X \mid x \leq z \leq y\}$  is a **closed interval** and  
 the set  $(x : y] \triangleq \{z \in X \mid x < z \leq y\}$  is a **half-open interval** and  
 the set  $[x : y) \triangleq \{z \in X \mid x \leq z < y\}$  is a **half-open interval** and  
 the set  $(x : y) \triangleq \{z \in X \mid x < z < y\}$  is an **open interval**.

**Definition 2.10** Let  $(\mathbb{R}, \leq)$  be the *ordered set of real numbers* (Definition 2.8 page 5).

The **absolute value**  $|\cdot| \in \mathbb{R}^{\mathbb{R}}$  is defined as<sup>13</sup>  $|x| \triangleq \begin{cases} -x & \text{for } x \leq 0 \\ x & \text{otherwise} \end{cases}$ .

## 3 Background: distance spaces

A *distance space* (Definition 3.1 page 6) can be defined as a *metric space* (Definition 4.5 page 16) without the *triangle inequality* constraint. Much of the material in this section about *distance spaces* is standard in *metric spaces*. However, this paper works through this material again to demonstrate “how far we can go”, and can't go, without the *triangle inequality*.

<sup>10</sup> [Hahn and Rosenthal(1948)], [Choquet(1954)], [Pap(1995)] page 8 (Definition 2.3: extended real-valued set function), [Halmos(1950)] page 30 (§7. MEASURE ON RINGS),

<sup>11</sup> [MacLane and Birkhoff(1999)] page 470, [Beran(1985)] page 1, [Korselt(1894)] page 156 (I, II, (1)), [Dedekind(1900)] page 373 (I–III). An *order relation* is also called a **partial order relation**. An *ordered set* is also called a **partially ordered set** or **poset**.

<sup>12</sup> [Apostol(1975)] page 4, [Ore(1935)] page 409

<sup>13</sup> A more general definition for *absolute value* is available for any *commutative ring*: Let  $R$  be a *commutative ring*. A function  $|\cdot|$  in  $R^R$  is an **absolute value**, or **modulus**, on  $R$  if

- |                             |              |                                     |     |
|-----------------------------|--------------|-------------------------------------|-----|
| 1. $ x  \geq 0$             | $x \in R$    | (non-negative)                      | and |
| 2. $ x  = 0 \iff x = 0$     | $x \in R$    | (nondegenerate)                     | and |
| 3. $ xy  =  x  \cdot  y $   | $x, y \in R$ | (homogeneous / submultiplicative)   | and |
| 4. $ x + y  \leq  x  +  y $ | $x, y \in R$ | (subadditive / triangle inequality) |     |

Reference: [Cohn(2002 December 6)] page 312



### 3.1 Fundamental structure of distance spaces

#### 3.1.1 Definitions

**Definition 3.1**<sup>14</sup> A function  $d$  in the set  $\mathbb{R}^{X \times X}$  (Definition 2.3 page 4) is a **distance** if

1.  $d(x, y) \geq 0 \quad \forall x, y \in X$  (*non-negative*) and
2.  $d(x, y) = 0 \iff x = y \quad \forall x, y \in X$  (*nondegenerate*) and
3.  $d(x, y) = d(y, x) \quad \forall x, y \in X$  (*symmetric*)

The pair  $(X, d)$  is a **distance space** if  $d$  is a *distance* on a set  $X$ .

**Definition 3.2**<sup>15</sup> Let  $(X, d)$  be a *distance space* and  $2^X$  be the *power set* of  $X$  (Definition 2.2 page 4). The **diameter** in  $(X, d)$  of a set  $A \in 2^X$  is  $\text{diam } A \triangleq \begin{cases} 0 & \text{for } A = \emptyset \\ \sup \{d(x, y) \mid x, y \in A\} & \text{otherwise} \end{cases}$

**Definition 3.3**<sup>16</sup> Let  $(X, d)$  be a *distance space*. Let  $2^X$  be the *power set* (Definition 2.2 page 4) of  $X$ . A set  $A$  is **bounded** in  $(X, d)$  if  $A \in 2^X$  and  $\text{diam } A < \infty$ .

#### 3.1.2 Properties

**Remark 3.4** Let  $(x_n)_{n \in \mathbb{Z}}$  be a *sequence* in a *distance space*  $(X, d)$ . The *distance space*  $(X, d)$  does not necessarily have all the nice properties that a *metric space* (Definition 4.5 page 16) has. In particular, note the following:

- |  |        |  |                                       |
|--|--------|--|---------------------------------------|
| 1. $d$ is a <i>distance</i> in $(X, d)$                          | $\iff$ | $d$ is <i>continuous</i> in $(X, d)$                 | (Example 3.23 page 14).               |
| 2. $B$ is an <i>open ball</i> in $(X, d)$                        | $\iff$ | $B$ is <i>open</i> in $(X, d)$                       | (Example 3.22 page 13).               |
| 3. $\mathcal{B}$ is the set of all <i>open balls</i> in $(X, d)$ | $\iff$ | $\mathcal{B}$ is a <i>base</i> for a topology on $X$ | (Example 3.22 page 13). <sup>17</sup> |
| 4. $(x_n)$ is <i>convergent</i> in $(X, d)$                      | $\iff$ | limit is <i>unique</i>                               | (Example 3.21 page 12).               |
| 5. $(x_n)$ is <i>convergent</i> in $(X, d)$                      | $\iff$ | $(x_n)$ is <i>Cauchy</i> in $(X, d)$                 | (Example 3.22 page 13).               |

### 3.2 Open sets in distance spaces

#### 3.2.1 Definitions

**Definition 3.5**<sup>18</sup> Let  $(X, d)$  be a *distance space* (Definition 3.1 page 6). Let  $\mathbb{R}^+$  be the *set of positive real numbers* (Definition 2.1 page 4).

An **open ball** centered at  $x$  with radius  $r$  is the set  $B(x, r) \triangleq \{y \in X \mid d(x, y) < r\}$ .

A **closed ball** centered at  $x$  with radius  $r$  is the set  $\bar{B}(x, r) \triangleq \{y \in X \mid d(x, y) \leq r\}$ .

<sup>14</sup> [Menger(1928)] page 76 (“Abstand  $a$   $b$  definiert ist...” (distance from  $a$  to  $b$  is defined as...)), [Wilson(1931)] page 361 (“§1., “distance”, “semi-metric space”), [Blumenthal(1938)] page 38, [Blumenthal(1953)] page 7 (“DEFINITION 5.1. A distance space is called semimetric provided...”), [Galvin and Shore(1984)] page 67 (“distance function”), [Laos(1998)] page 118 (“distance space”), [Khamisi and Kirk(2001)] page 13 (“semimetric space”), [Bessenyei and Pales(2014)] page 2 (“semimetric space”), [Deza and Deza(2014)] page 3 (“**distance** (or **dissimilarity**)”)

<sup>15</sup> in *metric space*: [Hausdorff(1937)], page 166, [Copson(1968)], page 23, [Michel and Herget(1993)], page 267, [Molchanov(2005)] page 389

<sup>16</sup> in *metric space*: [Thron(1966)], page 154 (definition 19.5), [Bruckner et al.(1997)Bruckner, Bruckner, and Thomson] page 356

<sup>17</sup> [Heath(1961)] page 810 (THEOREM), [Galvin and Shore(1984)] page 71 (2.3 LEMMA)

<sup>18</sup> in *metric space*: [Aliprantis and Burkinshaw(1998)], page 35



**Definition 3.6** Let  $(X, d)$  be a *distance space*. Let  $X \setminus A$  be the *set difference* of  $X$  and a set  $A$ . A set  $U$  is **open** in  $(X, d)$  if  $U \in 2^X$  and for every  $x$  in  $U$  there exists  $r \in \mathbb{R}^+$  such that  $B(x, r) \subseteq U$ . A set  $U$  is an **open set** in  $(X, d)$  if  $U$  is *open* in  $(X, d)$ . A set  $D$  is **closed** in  $(X, d)$  if  $(X \setminus D)$  is *open*. A set  $D$  is a **closed set** in  $(X, d)$  if  $D$  is *closed* in  $(X, d)$ .

### 3.2.2 Properties

**Theorem 3.7**<sup>19</sup> Let  $(X, d)$  be a DISTANCE SPACE. Let  $N$  be any (finite) positive integer. Let  $\Gamma$  be a SET possibly with an uncountable number of elements.

- |    |   |   |          |  |
|----|---|---|----------|--|
| 1. |   | $X$   | is OPEN. |  |
| 2. |   | $\emptyset$                                     | is OPEN. |  |
| 3. | each element in $\{U_n \mid n=1, 2, \dots, N\}$               | $\implies \bigcap_{n=1}^N U_n$                  | is OPEN. |  |
| 4. | each element in $\{U_\gamma \in 2^X \mid \gamma \in \Gamma\}$ | $\implies \bigcup_{\gamma \in \Gamma} U_\gamma$ | is OPEN. |  |

PROOF:

(1) Proof that  $X$  is *open* in  $(X, d)$ :

- (a) By definition of *open set* (Definition 3.6 page 7),  $X$  is *open*  $\iff \forall x \in X \exists r$  such that  $B(x, r) \subseteq X$ .
- (b) By definition of *open ball* (Definition 3.5 page 6), it is always true that  $B(x, r) \subseteq X$  in  $(X, d)$ .
- (c) Therefore,  $X$  is *open* in  $(X, d)$ .

(2) Proof that  $\emptyset$  is *open* in  $(X, d)$ :

- (a) By definition of *open set* (Definition 3.6 page 7),  $\emptyset$  is *open*  $\iff \forall x \in X \exists r$  such that  $B(x, r) \subseteq \emptyset$ .
- (b) By definition of *empty set*  $\emptyset$  (Definition 2.7 page 5), this is always true because no  $x$  is in  $\emptyset$ .
- (c) Therefore,  $\emptyset$  is *open* in  $(X, d)$ .

(3) Proof that  $\bigcup U_\gamma$  is *open* in  $(X, d)$ :

- (a) By definition of *open set* (Definition 3.6 page 7),  $\bigcup U_\gamma$  is *open*  $\iff \forall x \in \bigcup U_\gamma \exists r$  such that  $B(x, r) \subseteq \bigcup U_\gamma$ .
- (b) If  $x \in \bigcup U_\gamma$ , then there is at least one  $U \in \bigcup U_\gamma$  that contains  $x$ .
- (c) By the left hypothesis in (4), that set  $U$  is open and so for that  $x$ ,  $\exists r$  such that  $B(x, r) \subseteq U \subseteq \bigcup U_\gamma$ .
- (d) Therefore,  $\bigcup U_\gamma$  is *open* in  $(X, d)$ .

(4) Proof that  $U_1$  and  $U_2$  are *open*  $\implies U_1 \cap U_2$  is *open*:

- (a) By definition of *open set* (Definition 3.6 page 7),  $U_1 \cap U_2$  is *open*  $\iff \forall x \in U_1 \cap U_2 \exists r$  such that  $B(x, r) \subseteq U_1 \cap U_2$ .
- (b) By the left hypothesis above,  $U_1$  and  $U_2$  are *open*; and by the definition of *open sets* (Definition 3.6 page 7), there exists  $r_1$  and  $r_2$  such that  $B(x, r_1) \subseteq U_1$  and  $B(x, r_2) \subseteq U_2$ .
- (c) Let  $r \triangleq \min\{r_1, r_2\}$ . Then  $B(x, r) \subseteq U_1$  and  $B(x, r) \subseteq U_2$ .
- (d) By definition of *set intersection*  $\cap$  then,  $B(x, r) \subseteq U_1 \cap U_2$ .
- (e) By definition of *open set* (Definition 3.6 page 7),  $U_1 \cap U_2$  is *open*.

(5) Proof that  $\bigcap_{n=1}^N U_n$  is *open* (by induction):

- (a) Proof for  $N = 1$  case:  $\bigcap_{n=1}^N U_n = \bigcap_{n=1}^1 U_n = U_1$  is *open* by hypothesis.

<sup>19</sup> in *metric space*: [Dieudonné(1969)], pages 33–34, [Rosenlicht(1968)] page 39



(b) Proof that  $N$  case  $\implies N + 1$  case:

$$\begin{aligned} \bigcap_{n=1}^{N+1} U_n &= \left( \bigcap_{n=1}^N U_n \right) \cap U_{N+1} && \text{by property of } \bigcap \\ &\implies \text{open} && \text{by “}N\text{ case” hypothesis and (4) lemma page 7} \end{aligned}$$

⇒

**Corollary 3.8** Let  $(X, d)$  be a DISTANCE SPACE. The set  $T \triangleq \{U \in 2^X \mid U \text{ is OPEN in } (X, d)\}$  is a TOPOLOGY on  $X$ , and  $(X, T)$  is a TOPOLOGICAL SPACE.

✎PROOF: This follows directly from the definition of an *open set* (Definition 3.6 page 7), Theorem 3.7 (page 7), and the definition of *topology* (Definition A.1 page 26). ⇒

Of course it is possible to define a very large number of topologies even on a finite set with just a handful of elements;<sup>20</sup> and it is possible to define an infinite number of topologies even on a *linearly ordered* infinite set like the *real line*  $(\mathbb{R}, \leq)$ .<sup>21</sup> Be that as it may, Definition 3.9 (next definition) defines a single but convenient *topological space* in terms of a *distance space*. Note that every *metric space* conveniently and naturally induces a *topological space* because the *open balls* of the metric space form a *base* for the *topology*. This is not the case for all distance spaces. But if the open balls of a *distance space* are all *open*, then those open balls induce a topology (next theorem).<sup>22</sup>

**Definition 3.9** Let  $(X, d)$  be a *distance space*. The set  $T \triangleq \{U \in 2^X \mid U \text{ is open in } (X, d)\}$  is the **topology induced by  $(X, d)$  on  $X$** . The pair  $(X, T)$  is called the **topological space induced by  $(X, d)$** .

For any *distance space*  $(X, d)$ , no matter how strange, there is guaranteed to be at least one *topological space induced by  $(X, d)$* —and that is the *indiscrete topological space* (Example A.2 page 26) because for any distance space  $(X, d)$ ,  $\emptyset$  and  $X$  are *open sets* in  $(X, d)$  (Theorem 3.7 page 7).

**Theorem 3.10** Let  $B$  be the set of all OPEN BALLS in a DISTANCE SPACE  $(X, d)$ .  
 $\{ \text{every OPEN BALL in } B \text{ is OPEN} \} \iff \{ B \text{ is a BASE for a TOPOLOGY} \}$

<sup>20</sup> For a finite set  $X$  with  $n$  elements, there are 29 topologies on  $X$  if  $n = 3$ , 6942 topologies on  $X$  if  $n = 5$ , and 8,977,053,873,043 (almost 9 trillion) topologies on  $X$  if  $n = 10$ . References: ☞ [oei(2014)] (<http://oeis.org/A000798>), ☞ [Brown and Watson(1996)], page 31, ☞ [Comtet(1974)] page 229, ☞ [Comtet(1966)], ☞ [Chatterji(1967)], page 7, ☞ [Evans et al.(1967)Evans, Harary, and Lynn], ☞ [Krishnamurthy(1966)], page 157

<sup>21</sup> For examples of topologies on the real line, see the following: ☞ [Adams and Franzosa(2008)] page 31 (“six topologies on the real line”), ☞ [Salzmann et al.(2007)Salzmann, Grundhöfer, Hähl, and Löwen] pages 64–70 (“Weird topologies on the real line”), ☞ [Murdeswar(1990)] page 53 (“often used topologies on the real line”), ☞ [Joshi(1983)] pages 85–91 (§4.2 Examples of Topological Spaces)

<sup>22</sup> *metric space*: Definition 4.5 page 16; *open ball*: Definition 3.5 page 6; *base*: Definition A.3 page 27; *topology*: Definition A.1 page 26; not all open balls are open in a distance space: Example 3.21 (page 12) and Example 3.22 (page 13);





☞PROOF:

every open ball in  $\mathbf{B}$  is open

$\implies$  for every  $x$  in  $B_y \in \mathbf{B}$  there exists  $r \in \mathbb{R}^+$  such that  $B(x, r) \subseteq B_y$  by definition of *open* (Definition 3.6 page 7)

$\implies$   $\left\{ \begin{array}{l} \text{for every } x \in X \text{ and for every } B_y \in \mathbf{B} \text{ containing } x, \\ \text{there exists } B_x \in \mathbf{B} \text{ such that } x \in B_x \subseteq B_y. \end{array} \right\}$  because  $\forall (x, r) \in X \times \mathbb{R}^+, B(x, r) \subseteq X$

$\implies$   $\mathbf{B}$  is a base for  $T$

by Theorem A.4 page 27

$\implies$   $\left\{ \begin{array}{l} \text{for every } x \in X \text{ and for every } U \subseteq T \text{ containing } x, \\ \text{there exists } B_x \in \mathbf{B} \text{ such that } x \in B_x \subseteq U. \end{array} \right\}$

by Theorem A.4 page 27

$\implies$   $\left\{ \begin{array}{l} \text{for every } x \in X \text{ and for every } B_y \in \mathbf{B} \subseteq T \text{ containing } x, \\ \text{there exists } B_x \in \mathbf{B} \text{ such that } x \in B_x \subseteq B_y. \end{array} \right\}$

by definition of *base* (Definition A.3 page 27)

$\implies$   $\left\{ \begin{array}{l} \text{for every } x \in B_y \in \mathbf{B} \subseteq T, \\ \text{there exists } B_x \in \mathbf{B} \text{ such that } x \in B_x \subseteq B_y. \end{array} \right\}$

$\implies$  every open ball in  $\mathbf{B}$  is open

by definition of *open* (Definition 3.6 page 7)

☞

### 3.3 Sequences in distance spaces

#### 3.3.1 Definitions

**Definition 3.11** <sup>23</sup> Let  $(x_n \in X)_{n \in \mathbb{Z}}$  be a *sequence* in a *distance space*  $(X, d)$ . The sequence  $(x_n)$  **converges** to a **limit**  $x$  if for any  $\varepsilon \in \mathbb{R}^+$ , there exists  $N \in \mathbb{Z}$  such that for all  $n > N$ ,  $d(x_n, x) < \varepsilon$ .

This condition can be expressed in any of the following forms:

1. The **limit** of the sequence  $(x_n)$  is  $x$ .
2. The sequence  $(x_n)$  is **convergent** with limit  $x$ .
3.  $\lim_{n \rightarrow \infty} (x_n) = x$ .
4.  $(x_n) \rightarrow x$ .

A *sequence* that converges is **convergent**.

**Definition 3.12** <sup>24</sup> Let  $(x_n \in X)_{n \in \mathbb{Z}}$  be a *sequence* in a *distance space*  $(X, d)$ . The sequence  $(x_n)$  is a **Cauchy sequence** in  $(X, d)$  if

for every  $\varepsilon \in \mathbb{R}^+$ , there exists  $N \in \mathbb{Z}$  such that  $\forall n, m > N$ ,  $d(x_n, x_m) < \varepsilon$  (*Cauchy condition*).

**Definition 3.13** <sup>25</sup> Let  $(x_n \in X)_{n \in \mathbb{Z}}$  be a *sequence* in a *distance space*  $(X, d)$ .

The sequence  $(x_n \in X)_{n \in \mathbb{Z}}$  is **complete** in  $(X, d)$  if

$(x_n)$  is *Cauchy* in  $(X, d)$   $\implies$   $(x_n)$  is *convergent* in  $(X, d)$ .

#### 3.3.2 Properties

**Proposition 3.14** <sup>26</sup> Let  $(x_n \in X)_{n \in \mathbb{Z}}$  be a *SEQUENCE* in a *DISTANCE SPACE*  $(X, d)$ .

$\{ (x_n) \text{ is CAUCHY in } (X, d) \} \implies \{ (x_n) \text{ is BOUNDED in } (X, d) \}$

<sup>23</sup> in *metric space*: ☞ [Rosenlicht(1968)] page 45, ☞ [Giles(1987)] page 37 (3.2 Definition), ☞ [Khamisi and Kirk(2001)] page 13 (Definition 2.1) “ $\rightarrow$ ” symbol: ☞ [Leathem(1905)] page 13 (section III.11)

<sup>24</sup> in *metric space*: ☞ [Apostol(1975)] page 73 (4.7), ☞ [Rosenlicht(1968)] page 51

<sup>25</sup> in *metric space*: ☞ [Rosenlicht(1968)] page 52

<sup>26</sup> in *metric space*: ☞ [Giles(1987)] page 49 (Theorem 3.30)



PROOF:

$(x_n)$  is *Cauchy*  $\implies$  for every  $\varepsilon \in \mathbb{R}^+$ ,  $\exists N \in \mathbb{Z}$  such that  $\forall n, m > N$ ,  $d(x_n, x_m) < \varepsilon$  (by Definition 3.12 page 9)  
 $\implies \exists N \in \mathbb{Z}$  such that  $\forall n, m > N$ ,  $d(x_n, x_m) < 1$  (arbitrarily choose  $\varepsilon \triangleq 1$ )  
 $\implies \exists N \in \mathbb{Z}$  such that  $\forall n, m \in \mathbb{Z}$ ,  $d(x_n, x_{m+1}) < \max \{ \{1\} \cup \{d(x_p, x_q) \mid p, q \neq N\} \}$   
 $\implies (x_n)$  is *bounded* (by Definition 3.3 page 6)

◻

**Proposition 3.15**<sup>27</sup> Let  $(x_n \in X)_{n \in \mathbb{Z}}$  be a SEQUENCE in a DISTANCE SPACE  $(X, d)$ . Let  $f \in \mathbb{Z}^{\mathbb{Z}}$  (Definition 2.3 page 4) be a STRICTLY MONOTONE function such that  $f(n) < f(n+1)$ .

$\underbrace{(x_n)_{n \in \mathbb{Z}} \text{ is CAUCHY}}_{\text{sequence is CAUCHY}} \implies \underbrace{(x_{f(n)})_{n \in \mathbb{Z}} \text{ is CAUCHY}}_{\text{subsequence is also CAUCHY}}$

PROOF:

$(x_n)_{n \in \mathbb{Z}}$  is *Cauchy*  
 $\implies$  for any given  $\varepsilon > 0$ ,  $\exists N$  such that  $\forall n, m > N$ ,  $d(x_n, x_m) < \varepsilon$  by Definition 3.12 page 9  
 $\implies$  for any given  $\varepsilon > 0$ ,  $\exists N'$  such that  $\forall f(n), f(m) > N'$ ,  $d(x_{f(n)}, x_{f(m)}) < \varepsilon$   
 $\implies (x_{f(n)})_{n \in \mathbb{Z}}$  is *Cauchy* by Definition 3.12 page 9

◻

**Theorem 3.16**<sup>28</sup> Let  $(X, d)$  be a DISTANCE SPACE. Let  $A^-$  be the CLOSURE (Definition A.8 page 27) of a  $A$  in a TOPOLOGICAL SPACE INDUCED BY  $(X, d)$ .

$\left\{ \begin{array}{l} 1. \text{ LIMITS are UNIQUE in } (X, d) \text{ (Definition 3.11 page 9)} \\ 2. (A, d) \text{ is COMPLETE in } (X, d) \text{ (Definition 3.13 page 9)} \end{array} \right\} \text{ and } \implies \underbrace{A \text{ is CLOSED in } (X, d)}_{A = A^-}$

PROOF:

- (1) Proof that  $A \subseteq A^-$ : by Lemma A.10 page 27
- (2) Proof that  $A^- \subseteq A$  (proof that  $x \in A^- \implies x \in A$ ):
  - (a) Let  $x$  be a point in  $A^-$  ( $x \in A^-$ ).
  - (b) Define a *sequence* of open balls  $(B(x, \frac{1}{1}), B(x, \frac{1}{2}), B(x, \frac{1}{3}), \dots)$ .
  - (c) Define a *sequence* of points  $(x_1, x_2, x_3, \dots)$  such that  $x_n \in B(x, \frac{1}{n}) \cap A$ .
  - (d) Then  $(x_n)$  is *convergent* in  $X$  with limit  $x$  by Definition 3.11 page 9
  - (e) and  $(x_n)$  is *Cauchy* in  $A$  by Definition 3.12 page 9.
  - (f) By the hypothesis 2,  $(x_n)$  is therefore also *convergent* in  $A$ .  
Let this limit be  $y$ . Note that  $y \in A$ .
  - (g) By hypothesis 1, limits are *unique*, so  $y = x$ .
  - (h) Because  $y \in A$  (item (2f)) and  $y = x$  (item (2g)), so  $x \in A$ .
  - (i) Therefore,  $x \in A^- \implies x \in A$  and  $A^- \subseteq A$ .

◻

**Proposition 3.17**<sup>29</sup> Let  $(x_n)_{n \in \mathbb{Z}}$  be a sequence in a DISTANCE SPACE  $(X, d)$ . Let  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  be a strictly

<sup>27</sup> in metric space: [Rosenlicht(1968)] page 52

<sup>28</sup> in metric space: [Kubrusly(2001)] page 128 (Theorem 3.40), [Haaser and Sullivan(1991)] page 75 (6.10, 6.11 Propositions), [Bryant(1985)] page 40 (Theorem 3.6, 3.7), [Sutherland(1975)] pages 123–124

<sup>29</sup> in metric space: [Rosenlicht(1968)] page 46



increasing function such that  $f(n) < f(n + 1)$ .

$$\underbrace{(x_n)_{n \in \mathbb{Z}} \rightarrow x}_{\text{sequence converges to limit } x} \implies \underbrace{(x_{f(n)})_{n \in \mathbb{Z}} \rightarrow x}_{\text{subsequence converges to the same limit } x}$$

PROOF:

$$\begin{aligned} (x_n)_{n \in \mathbb{Z}} \rightarrow x &\implies \forall \epsilon > 0, \exists N \text{ such that } \forall n > N, d(x_n, x) < \epsilon && \text{by Theorem 4.15 page 21} \\ &\implies \forall \epsilon > 0, \exists f(N) \text{ such that } \forall f(n) > f(N), d(x_{f(n)}, x) < \epsilon \\ &\implies (x_{f(n)})_{n \in \mathbb{Z}} \rightarrow x && \text{by Theorem 4.15 page 21} \end{aligned}$$

⇒

**Theorem 3.18** (Cantor intersection theorem)<sup>30</sup> Let  $(X, d)$  be a DISTANCE SPACE (Definition 3.1 page 6),  $(A_n)_{n \in \mathbb{Z}}$  a SEQUENCE with each  $A_n \in 2^X$ , and  $|A|$  the number of elements in  $A$ .

$$\left\{ \begin{array}{l} 1. (X, d) \text{ is COMPLETE} \quad (\text{Definition 3.13 page 9}) \text{ and} \\ 2. A_n \text{ is CLOSED} \quad \forall n \in \mathbb{N} \quad (\text{Definition A.1 page 26}) \text{ and} \\ 3. \text{diam } A_n \geq \text{diam } A_{n+1} \quad \forall n \in \mathbb{N} \quad (\text{Definition 3.2 page 6}) \text{ and} \\ 4. \text{diam } (A_n)_{n \in \mathbb{Z}} \rightarrow 0 \quad (\text{Definition 3.11 page 9}) \end{array} \right\} \implies \left\{ \left| \bigcap_{n \in \mathbb{N}} A_n \right| = 1 \right\}$$

PROOF:

- (1) Proof that  $|\bigcap_{n \in \mathbb{Z}} A_n| < 2$ :
  - (a) Let  $A \triangleq \bigcap A_n$ .
  - (b)  $x \neq y$  and  $\{x, y\} \in A \implies d(x, y) > 0$  and  $\{x, y\} \subseteq A_n \forall n$
  - (c)  $\exists n$  such that  $\text{diam } A_n < d(x, y)$  by left hypothesis 4
  - (d)  $\implies \exists n$  such that  $\sup \{d(x, y) \mid x, y \in A_n\} < d(x, y)$
  - (e) This is a contradiction, so  $\{x, y\} \notin A$  and  $|\bigcap A_n| < 2$ .
- (2) Proof that  $|\bigcap A_n| \geq 1$ :
  - (a) Let  $x_n \in A_n$  and  $x_m \in A_m$
  - (b)  $\forall \epsilon, \exists N \in \mathbb{N}$  such that  $\text{diam } A_N < \epsilon$
  - (c)  $\forall m, n > N, x_n \in A_n \subseteq A_N$  and  $x_m \in A_m \subseteq A_N$
  - (d)  $d(x_n, x_m) \leq \text{diam } A_N < \epsilon \implies \{x_n\}$  is a Cauchy sequence
  - (e) Because  $\{x_n\}$  is complete,  $x_n \rightarrow x$ .
  - (f)  $\implies x \in (A_n)^- = A_n$
  - (g)  $\implies |A_n| \geq 1$

⇒

**Definition 3.19**<sup>31</sup> Let  $(X, d)$  be a distance space. Let  $C$  be the set of all convergent sequences in  $(X, d)$ . The distance function  $d$  is **continuous** in  $(X, d)$  if

$$(x_n), (y_n) \in C \implies \lim_{n \rightarrow \infty} (d(x_n, y_n)) = d\left(\lim_{n \rightarrow \infty} (x_n), \lim_{n \rightarrow \infty} (y_n)\right).$$

A distance function is **discontinuous** if it is not continuous.

<sup>30</sup> in metric space: [Davis(2005)], page 28, [Hausdorff(1937)], page 150

<sup>31</sup> [Blumenthal(1953)] page 9 (DEFINITION 6.3)



**Remark 3.20** Rather than defining *continuity* of a *distance function* in terms of the *sequential characterization of continuity* as in Definition 3.19 (previous), we could define continuity using an *inverse image characterization of continuity*<sup>32</sup> (Definition 3.9 page 8). Assuming an equivalent *topological space* is used for both characterizations, the two characterizations are equivalent (Theorem A.20 page 29). In fact, one could construct an equivalence such as the following:

$$\left\{ \begin{array}{l} d \text{ is } \textit{continuous} \text{ in } \mathbb{R}^{X^2} \\ \text{(Definition A.11 page 28)} \\ \text{(inverse image characterization of continuity)} \end{array} \right\} \iff \left\{ \begin{array}{l} ((x_n), (y_n)) \in C \implies \\ \lim_{n \rightarrow \infty} (d(x_n, y_n)) = d\left(\lim_{n \rightarrow \infty} (x_n), \lim_{n \rightarrow \infty} (y_n)\right) \\ \text{(Definition A.16 page 28)} \\ \text{(sequential characterization of continuity)} \end{array} \right\}$$

Note that just as  $(x_n)$  is a sequence in  $X$ , so the ordered pair  $((x_n), (y_n))$  is a sequence in  $X^2$ . The remainder follows from Theorem A.20 (page 29). However, use of the *inverse image characterization* is somewhat troublesome because we would need a topology on  $X^2$ , and we don't immediately have one defined and ready to use. In fact, we don't even immediately have a distance space on  $X^2$  defined or even open balls in such a distance space. The result is, for the scope of this paper, it is arguably not worthwhile constructing the extra structure, but rather instead this paper uses the *sequential characterization* as a definition (as in Definition 3.19).

### 3.4 Examples

Similar distance functions and several of the observations for the examples in this section can be found in  [Blumenthal(1953)] pages 8–13.

In a *metric space*, all *open balls* are *open*, the *open balls* form a *base* for a *topology*, the limits of *convergent* sequences are *unique*, and the *metric function* is *continuous*. In the *distance space* of the next example, none of these properties hold.

**Example 3.21**<sup>32</sup> Let  $(x, y)$  be an *ordered pair* in  $\mathbb{R}^2$ . Let  $(a : b)$  be an *open interval* and  $(a : b]$  a *half-open interval* in  $\mathbb{R}$ . Let  $|x|$  be the *absolute value* of  $x \in \mathbb{R}$ . The function  $d(x, y) \in \mathbb{R}^{\mathbb{R} \times \mathbb{R}}$  such that

$$d(x, y) \triangleq \left\{ \begin{array}{ll} y & \forall (x, y) \in \{4\} \times (0 : 2] \quad \text{(vertical half-open interval)} \\ x & \forall (x, y) \in (0 : 2] \times \{4\} \quad \text{(horizontal half-open interval)} \\ |x - y| & \text{otherwise} \quad \text{(Euclidean)} \end{array} \right\} \text{ is a } \textit{distance} \text{ on } \mathbb{R}.$$

Note some characteristics of the *distance space*  $(\mathbb{R}, d)$ :

- (1)  $(\mathbb{R}, d)$  is not a *metric space* because  $d$  does not satisfy the *triangle inequality*:

$$d(0, 4) \triangleq |0 - 4| = 4 \not\leq 2 = |0 - 1| + 1 \triangleq d(0, 1) + d(1, 4)$$


- (2) Not every *open ball* in  $(\mathbb{R}, d)$  is *open*.

For example, the *open ball*  $B(3, 2)$  is *not open* because  $4 \in B(3, 2)$  but for all  $0 < \varepsilon < 1$

$$B(4, \varepsilon) = (4 - \varepsilon : 4 + \varepsilon) \cup (0 : \varepsilon) \not\subseteq (1 : 5) = B(3, 2)$$

- (3) The *open balls* of  $(\mathbb{R}, d)$  do not form a *base* for a *topology* on  $\mathbb{R}$ .

This follows directly from item (2) and Theorem 3.10 (page 8).

<sup>32</sup> A similar distance function  $d$  and item (4) page 13 can in essence be found in  [Blumenthal(1953)] page 8. Definitions for Example 3.21:  $(x, y)$ : Definition 2.3 (page 4);  $(a : b)$  and  $(a : b]$ : Definition 2.9 (page 5);  $|x|$ : Definition 2.10 (page 5);  $\mathbb{R}^{\mathbb{R} \times \mathbb{R}}$ : Definition 2.3 (page 4); *distance*: Definition 3.1 (page 6); *open ball*: Definition 3.5 (page 6); *open*: Definition 3.6 (page 7); *base*: Definition A.3 (page 27); *topology*: Definition A.1 (page 26); *open set*: Definition 3.6 (page 7); *topological space induced by*  $(\mathbb{R}, d)$ : Definition 3.9 (page 8); *discontinuous*: Definition 3.19 (page 11);



(4) In the *distance space*  $(\mathbb{R}, d)$ , limits are *not unique*;

For example, the sequence  $(\frac{1}{n})_1^\infty$  converges both to the limit 0 and the limit 4 in  $(\mathbb{R}, d)$ :

$$\begin{aligned} \lim_{n \rightarrow \infty} d(x_n, 0) &\triangleq \lim_{n \rightarrow \infty} d(\frac{1}{n}, 0) \triangleq \lim_{n \rightarrow \infty} |\frac{1}{n} - 0| = 0 \implies (\frac{1}{n}) \rightarrow 0 \\ \lim_{n \rightarrow \infty} d(x_n, 4) &\triangleq \lim_{n \rightarrow \infty} d(\frac{1}{n}, 4) \triangleq \lim_{n \rightarrow \infty} (\frac{1}{n}) = 0 \implies (\frac{1}{n}) \rightarrow 4 \end{aligned}$$

(5) The *topological space*  $(X, T)$  induced by  $(\mathbb{R}, d)$  also yields limits of 0 and 4 for the sequence  $(\frac{1}{n})_1^\infty$ , just as it does in item (4). This is largely due to the fact that, for small  $\varepsilon$ , the open balls  $B(0, \varepsilon)$  and  $B(4, \varepsilon)$  are *open*.

$$\begin{aligned} B(0, \varepsilon) \text{ is open} &\implies \text{for each } U \in T \text{ that contains } 0, \exists N \in \mathbb{N} \text{ such that } \frac{1}{n} \in U \quad \forall n > N \\ &\iff (\frac{1}{n}) \rightarrow 0 \quad \text{by definition of convergence (Definition A.16 page 28)} \end{aligned}$$

$$\begin{aligned} B(4, \varepsilon) \text{ is open} &\implies \text{for each } U \in T \text{ that contains } 4, \exists N \in \mathbb{N} \text{ such that } \frac{1}{n} \in U \quad \forall n > N \\ &\iff (\frac{1}{n}) \rightarrow 4 \quad \text{by definition of convergence (Definition A.16 page 28)} \end{aligned}$$

(6) The distance function  $d$  is *discontinuous* (Definition 3.19 page 11):

$$\begin{aligned} \lim_{n \rightarrow \infty} (d(1 - \frac{1}{n}, 4 - \frac{1}{n})) &= \lim_{n \rightarrow \infty} (|(1 - \frac{1}{n}) - (4 - \frac{1}{n})|) = |1 - 4| = 3 \neq 4 = d(0, 4) \\ &= d\left(\lim_{n \rightarrow \infty} (1 - \frac{1}{n}), \lim_{n \rightarrow \infty} (4 - \frac{1}{n})\right) \end{aligned}$$

In a *metric space*, all *convergent* sequences are also *Cauchy*. However, this is not the case for all *distance spaces*, as demonstrated next:

**Example 3.22** <sup>33</sup> The function  $d(x, y) \in \mathbb{R}^{\mathbb{R} \times \mathbb{R}}$  such that

$$d(x, y) \triangleq \left\{ \begin{array}{ll} |x - y| & \text{for } x = 0 \text{ or } y = 0 \text{ or } x = y \quad (\text{Euclidean}) \\ 1 & \text{otherwise} \quad (\text{discrete}) \end{array} \right\} \text{ is a distance on } \mathbb{R}.$$

Note some characteristics of the *distance space*  $(\mathbb{R}, d)$ :

(1)  $(X, d)$  is not a *metric space* because the *triangle inequality* does not hold:

$$d(\frac{1}{4}, \frac{1}{2}) = 1 \not\leq \frac{3}{4} = |\frac{1}{4} - 0| + |0 - \frac{1}{2}| = d(\frac{1}{4}, 0) + d(0, \frac{1}{2})$$

(2) The *open ball*  $B(\frac{1}{4}, \frac{1}{2})$  is *not open* because for any  $\varepsilon \in \mathbb{R}^+$ , no matter how small,

$$B(0, \varepsilon) = (-\varepsilon : +\varepsilon) \not\subseteq \{0, \frac{1}{4}\} = \{x \in X \mid d(\frac{1}{4}, x) < \frac{1}{2}\} \triangleq B(\frac{1}{4}, \frac{1}{2})$$

(3) Even though not all the *open balls* are *open*, it is still possible to have an *open set* in  $(X, d)$ . For example, the set  $U \triangleq \{1, 2\}$  is *open*:

$$B(1, 1) \triangleq \{x \in X \mid d(1, x) < 1\} = \{1\} \subseteq \{1, 2\} \triangleq U$$


$$B(2, 1) \triangleq \{x \in X \mid d(2, x) < 1\} = \{2\} \subseteq \{1, 2\} \triangleq U$$

(4) By item (2) and Theorem 3.10 (page 8), the *open balls* of  $(\mathbb{R}, d)$  do not form a *base* for a *topology* on  $\mathbb{R}$ .

(5) Even though the *open balls* in  $(\mathbb{R}, d)$  do not induce a *topology* on  $X$ , it is still possible to find a set of *open sets* in  $(X, d)$  that is a *topology*. For example, the set  $\{\emptyset, \{1, 2\}, \mathbb{R}\}$  is a *topology* on  $\mathbb{R}$ .

(6) In  $(\mathbb{R}, d)$ , limits of *convergent* sequences are *unique*:

$$(\frac{1}{n}) \rightarrow x \implies \lim_{n \rightarrow \infty} d(x_n, x) = \left\{ \begin{array}{ll} \lim_{n \rightarrow \infty} |x_n - 0| = 0 & \text{for } x = 0 \quad \text{OR} \\ |x - x| = 0 & \text{for constant } (x_n) \text{ for } n > N \quad \text{OR} \\ 1 \neq 0 & \text{otherwise} \end{array} \right\}$$

<sup>33</sup> The distance function  $d$  and item (7) page 14 can in essence be found in  [Blumenthal(1953)] page 9



which says that there are only two ways for a sequence to converge: either  $x = 0$  or the sequence eventually becomes constant (or both). Any other sequence will *diverge*. Therefore we can say the following:

- (a) If  $x = 0$  and the sequence is not constant, then the limit is *unique* and 0.
  - (b) If  $x = 0$  and the sequence is constant, then the limit is *unique* and 0.
  - (c) If  $x \neq 0$  and the sequence is constant, then the limit is *unique* and  $x$ .
  - (d) If  $x \neq 0$  and the sequence is not constant, then the sequence diverges and there is no limit.
- (7) In  $(\mathbb{R}, d)$ , a *convergent* sequence is not necessarily *Cauchy*. For example,
- (a) the sequence  $(\frac{1}{n})_{n \in \mathbb{N}}$  is *convergent* with limit 0:  $\lim_{n \rightarrow \infty} d(\frac{1}{n}, 0) = \lim_{n \rightarrow \infty} \frac{1}{n} = 0$
  - (b) However, even though  $(\frac{1}{n})$  is *convergent*, it is *not Cauchy*:  $\lim_{n, m \rightarrow \infty} d(\frac{1}{n}, \frac{1}{m}) = 1 \neq 0$
- (8) The *distance function*  $d$  is *discontinuous* in  $(X, d)$ :


$$\lim_{n \rightarrow \infty} (d(\frac{1}{n}, 2 - \frac{1}{n})) = 1 \neq 2 = d(0, 2) = d\left(\lim_{n \rightarrow \infty} (\frac{1}{n}), \lim_{n \rightarrow \infty} (2 - \frac{1}{n})\right).$$

**Example 3.23** <sup>34</sup> The function  $d(x, y) \in \mathbb{R}^{\mathbb{R} \times \mathbb{R}}$  such that

$$d(x, y) \triangleq \left\{ \begin{array}{ll} 2|x - y| & \forall (x, y) \in \{(0, 1), (1, 0)\} \quad (\text{dilated Euclidean}) \\ |x - y| & \text{otherwise} \quad (\text{Euclidean}) \end{array} \right\} \text{ is a distance on } \mathbb{R}.$$

Note some characteristics of the *distance space*  $(\mathbb{R}, d)$ :

- (1)  $(\mathbb{R}, d)$  is *not* a *metric space* because  $d$  does *not* satisfy the *triangle inequality*:  
 $d(0, 1) \triangleq 2|0 - 1| = 2 \not\leq 1 = |0 - \frac{1}{2}| + |\frac{1}{2} - 1| \triangleq d(0, \frac{1}{2}) + d(\frac{1}{2}, 1)$
- (2) The function  $d$  is *discontinuous*:  
 $\lim_{n \rightarrow \infty} (d(1 - \frac{1}{n}, \frac{1}{n})) \triangleq \lim_{n \rightarrow \infty} (|1 - \frac{1}{n} - \frac{1}{n}|) = 1 \neq 2 = 2|0 - 1| \triangleq d(0, 1) = d\left(\lim_{n \rightarrow \infty} (1 - \frac{1}{n}), \lim_{n \rightarrow \infty} (\frac{1}{n})\right).$
- (3) In  $(X, d)$ , *open balls* are *open*:
  - (a)  $\rho(x, y) \triangleq |x - y|$  is a *metric* and thus all open balls in that do not contain both 0 and 1 are *open*.
  - (b) By Example C.4 (page 36),  $q(x, y) \triangleq 2|x - y|$  is also a *metric* and thus all open balls containing 0 and 1 only are *open*.
  - (c) The only question remaining is with regards to open balls that contain 0, 1 and some other element(s) in  $\mathbb{R}$ . But even in this case, open balls are still open. For example:  
 $B(-1, 2) = (-1 : 2) = (-1 : 1) \cup (1 : 2)$   
 Note that both  $(-1 : 1)$  and  $(1 : 2)$  are *open*, and thus by Theorem 3.7 (page 7),  $B(-1, 2)$  is *open* as well.
- (4) By item (3) and Theorem 3.10 (page 8), the *open balls* of  $(\mathbb{R}, d)$  *do* form a *base* for a *topology* on  $\mathbb{R}$ .
- (5) In  $(X, d)$ , the limits of *convergent* sequences are *unique*. This is demonstrated in Example 4.22 (page 25) using additional structure developed in Section 4.
- (6) In  $(X, d)$ , *convergent* sequences are *Cauchy*. This is also demonstrated in Example 4.22 (page 25).

<sup>34</sup> The distance function  $d$  and item (2) page 14 can in essence be found in  [Blumenthal(1953)] page 9



The *distance functions* in Example 3.21 (page 12)–Example 3.23 (page 14) were all *discontinuous*. In the absence of the *triangle inequality* and in light of these examples, one might try replacing the *triangle inequality* with the weaker requirement of *continuity*. However, as demonstrated by the next example, this also leads to an arguably disastrous result.

**Example 3.24** <sup>35</sup> The function  $d \in \mathbb{R}^{\mathbb{R} \times \mathbb{R}}$  such that  $d(x, y) \triangleq (x - y)^2$  is a *distance* on  $\mathbb{R}$ . Note some characteristics of the *distance space*  $(\mathbb{R}, d)$ :

- (1)  $(\mathbb{R}, d)$  is *not* a *metric space* because the *triangle inequality* does not hold:  

$$d(0, 2) \triangleq (0 - 2)^2 = 4 \not\leq 2 = (0 - 1)^2 + (1 - 2)^2 \triangleq d(0, 1) + d(1, 2)$$
- (2) The *distance function*  $d$  is *continuous* in  $(X, d)$ . This is demonstrated in the more general setting of Section 4 in Example 4.23 (page 25).
- (3) Calculating the length of curves in  $(X, d)$  leads to a paradox:<sup>36</sup>

- (a) Partition  $[0 : 1]$  into  $2^N$  consecutive line segments connected at the points

$$\left(0, \frac{1}{2^N}, \frac{2}{2^N}, \frac{3}{2^N}, \dots, \frac{2^N - 1}{2^N}, 1\right)$$

- (b) Then the distance, as measured by  $d$ , between any two consecutive points is

$$d(p_n, p_{n+1}) \triangleq (p_n - p_{n+1})^2 = \left(\frac{1}{2^N}\right)^2 = \frac{1}{2^{2N}}$$

- (c) But this leads to the paradox that the total length of  $[0 : 1]$  is 0:

$$\lim_{N \rightarrow \infty} \sum_{n=0}^{2^N - 1} \frac{1}{2^{2N}} = \lim_{N \rightarrow \infty} \frac{2^N}{2^{2N}} = \lim_{N \rightarrow \infty} \frac{1}{2^N} = 0$$

## 4 Distance spaces with power triangle inequalities

### 4.1 Definitions

This paper introduces a new relation called the *power triangle inequality* (Definition 4.3 page 16). It is a generalization of other common relations, including the *triangle inequality* (Definition 4.4 page 16). The *power triangle inequality* is defined in terms of a function herein called the *power triangle function* (next definition). This function is a special case of the *power mean* with  $N = 2$  and  $\lambda_1 = \lambda_2 = \frac{1}{2}$  (Definition B.6 page 31). *Power means* have the attractive properties of being *continuous* and *strictly monotone* with respect to a free parameter  $p \in \mathbb{R}^*$  (Theorem B.7 page 31). This fact is inherited and exploited by the *power triangle inequality* (Corollary 4.6 page 16).

**Definition 4.1** Let  $(X, d)$  be a *distance space* (Definition 3.1 page 6). Let  $\mathbb{R}^+$  be the set of all *positive real numbers* and  $\mathbb{R}^*$  be the set of *extended real numbers* (Definition 2.1 page 4). The **power triangle function**  $\tau$  on  $(X, d)$  is defined as

$$\tau(p, \sigma; x, y, z; d) \triangleq 2\sigma \left[ \frac{1}{2}d^p(x, z) + \frac{1}{2}d^p(z, y) \right]^{\frac{1}{p}} \quad \forall (p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+, \quad x, y, z \in X$$

<sup>35</sup> [Blumenthal(1953)] pages 12–13, [Laos(1998)] pages 118–119

<sup>36</sup> This is the method of “inscribed polygons” for calculating the length of a curve and goes back to Archimedes: [Brunschwig et al.(2003)Brunschwig, Lloyd, and Pellegrin] page 26, [Walmsley(1920)], page 200 (§158),



**Remark 4.2** <sup>37</sup> In the field of *probabilistic metric spaces*, a function called the *triangle function* was introduced by Sherstnev in 1962. However, the *power triangle function* as defined in this present paper is *not* a special case of (is not compatible with) the *triangle function* of Sherstnev. Another definition of *triangle function* has been offered by Bessenyei in 2014 with special cases of  $\Phi(u, v) \triangleq c(u + v)$  and  $\Phi(u, v) \triangleq (u^p + v^p)^{\frac{1}{p}}$ , which are similar to the definition of *power triangle function* offered in this present paper.

**Definition 4.3** Let  $(X, d)$  be a *distance space*. Let  $\mathcal{2}^{X \times X}$  be the set of all trinomial *relations* (Definition 2.3 page 4) on  $X$ . A relation  $\Theta(p, \sigma; d)$  in  $\mathcal{2}^{X \times X}$  is a **power triangle inequality** on  $(X, d)$  if

$$\Theta(p, \sigma; d) \triangleq \{(x, y, z) \in X^3 \mid d(x, y) \leq \tau(p, \sigma; x, y, z; d)\} \quad \text{for some } (p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+.$$

The tuple  $(X, d, p, \sigma)$  is a **power distance space** and  $d$  a **power distance** or **power distance function** if  $(X, d)$  is a *distance space* in which the *triangle relation*  $\Theta(p, \sigma; d)$  holds.

The *power triangle function* can be used to define some standard inequalities (next definition). See Corollary 4.7 (page 17) for some justification of the definitions.

**Definition 4.4** <sup>38</sup> Let  $\Theta(p, \sigma; d)$  be a *power triangle inequality* on a *distance space*  $(X, d)$ .

- |   |  |
|---|--|
| 1. $\Theta(\infty, \sigma/2; d)$ is the <b><math>\sigma</math>-inframetric inequality</b> | 6. $\Theta(1/2, 2; d)$ is the <b>square mean root inequality</b> |
| 2. $\Theta(\infty, 1/2; d)$ is the <b>inframetric inequality</b>                          | 7. $\Theta(0, 1/2; d)$ is the <b>geometric inequality</b>        |
| 3. $\Theta(2, \sqrt{2}\sigma; d)$ is the <b>quadratic inequality</b>                      | 8. $\Theta(-1, 1/4; d)$ is the <b>harmonic inequality</b>        |
| 4. $\Theta(1, \sigma; d)$ is the <b>relaxed triangle inequality</b>                       | 9. $\Theta(-\infty, 1/2; d)$ is the <b>minimal inequality</b>    |
| 5. $\Theta(1, 1; d)$ is the <b>triangle inequality</b>                                    |  |

**Definition 4.5** <sup>39</sup> Let  $(X, d)$  be a *distance space* (Definition 3.1 page 6).

- |   |  |                |
|---|--|----------------|
| 1. $(X, d)$ is a <b>metric space</b>                          | if the <i>triangle inequality</i>                        | holds in $X$ . |
| 2. $(X, d)$ is a <b>near metric space</b>                     | if the <i>relaxed triangle inequality</i>                | holds in $X$ . |
| 3. $(X, d)$ is an <b>inframetric space</b>                    | if the <i>inframetric inequality</i>                     | holds in $X$ . |
| 4. $(X, d)$ is a <b><math>\sigma</math>-inframetric space</b> | if the <i><math>\sigma</math>-inframetric inequality</i> | holds in $X$ . |

## 4.2 Properties

### 4.2.1 Relationships of the power triangle function

**Corollary 4.6** Let  $\tau(p, \sigma; x, y, z; d)$  be the POWER TRIANGLE FUNCTION (Definition 4.1 page 15) in the DISTANCE SPACE (Definition 3.1 page 6)  $(X, d)$ . Let  $(\mathbb{R}, |\cdot|, \leq)$  be the ORDERED METRIC SPACE with the usual ordering relation

<sup>37</sup> [Sherstnev(1962)], page 4, [Schweizer and Sklar(1983)] page 9 (1.6.1)–(1.6.4), [Bessenyei and Pales(2014)] page 2

<sup>38</sup> [Bessenyei and Pales(2014)] page 2, [Czerwik(1993)] page 5 (*b-metric*; (1),(2),(5)), [Fagin et al.(2003a)Fagin, Kumar, and Sivakumar], [Fagin et al.(2003b)Fagin, Kumar, and Sivakumar] (Definition 4.2 (Relaxed metrics)), [Xia(2009)] page 453 (Definition 2.1), [Heinonen(2001)] page 109 (14.1 Quasimetric spaces.), [Kirk and Shahzad(2014)] page 113 (Definition 12.1), [Deza and Deza(2014)] page 7, [Hoehn and Niven(1985)] page 151, [Gibbons et al.(1977)Gibbons, Olkin, and Sobel] page 51 (*square-mean-root (SMR)* (2.4.1)), [Euclid(circa 300BC)] (triangle inequality—Book I Proposition 20)

<sup>39</sup> **metric space:** [Dieudonné(1969)], page 28, [Copson(1968)], page 21, [Hausdorff(1937)] page 109, [Fréchet(1928)], [Fréchet(1906)] page 30 **near metric space:** [Czerwik(1993)] page 5 (*b-metric*; (1),(2),(5)), [Fagin et al.(2003a)Fagin, Kumar, and Sivakumar], [Fagin et al.(2003b)Fagin, Kumar, and Sivakumar] (Definition 4.2 (Relaxed metrics)), [Xia(2009)] page 453 (Definition 2.1), [Heinonen(2001)] page 109 (14.1 Quasimetric spaces.), [Kirk and Shahzad(2014)] page 113 (Definition 12.1), [Deza and Deza(2014)] page 7





$\leq$  and usual metric  $|\cdot|$  on  $\mathbb{R}$ . The function  $\tau(p, \sigma; x, y, z; d)$  is CONTINUOUS and STRICTLY MONOTONE in  $(\mathbb{R}, |\cdot|, \leq)$  with respect to both the variables  $p$  and  $\sigma$ .

✎PROOF:

- (1) Proof that  $\tau(p, \sigma; x, y, z; d)$  is *continuous* and *strictly monotone* with respect to  $p$ : This follows directly from Theorem B.7 (page 31).
- (2) Proof that  $\tau(p, \sigma; x, y, z; d)$  is *continuous* and *strictly monotone* with respect to  $\sigma$ :

$$\begin{aligned} \tau(p, \sigma; x, y, z; d) &\triangleq 2\sigma \underbrace{\left[ \frac{1}{2}d^p(x, z) + \frac{1}{2}d^p(z, y) \right]^{\frac{1}{p}}}_{f(p, x, y, z)} && \text{by definition of } \tau \text{ (Definition 4.1 page 15)} \\ &= 2\sigma f(p, x, y, z) && \text{where } f \text{ is defined as above} \\ &\implies \tau \text{ is } \textit{affine} \text{ with respect to } \sigma \\ &\implies \tau \text{ is } \textit{continuous} \text{ and } \textit{strictly monotone} \text{ with respect to } \sigma: \end{aligned}$$

⇒

**Corollary 4.7** Let  $\tau(p, \sigma; x, y, z; d)$  be the POWER TRIANGLE FUNCTION in the DISTANCE SPACE (Definition 3.1 page 6)  $(X, d)$ .

$$\tau(p, \sigma; x, y, z; d) = \begin{cases} 2\sigma \max \{d(x, z), d(z, y)\} & \text{for } p = \infty, & (\text{MAXIMUM, corresponds to INFRAMETRIC SPACE}) \\ 2\sigma \left[ \frac{1}{2}d^2(x, z) + \frac{1}{2}d^2(z, y) \right]^{\frac{1}{2}} & \text{for } p = 2, & (\text{QUADRATIC MEAN}) \\ \sigma [d(x, z) + d(z, y)] & \text{for } p = 1, & (\text{ARITHMETIC MEAN, corresponds to NEAR METRIC SPACE}) \\ 2\sigma \sqrt{d(x, z)} \sqrt{d(z, y)} & \text{for } p = 0 & (\text{GEOMETRIC MEAN}) \\ 4\sigma \left[ \frac{1}{d(x, z)} + \frac{1}{d(z, y)} \right]^{-1} & \text{for } p = -1 & (\text{HARMONIC MEAN}) \\ 2\sigma \min \{d(x, z), d(z, y)\} & \text{for } p = -\infty, & (\text{MINIMUM}) \end{cases}$$

✎PROOF: These follow directly from Theorem B.7 (page 31).

⇒

**Corollary 4.8** Let  $(X, d)$  be a DISTANCE SPACE.

$$\begin{aligned} 2\sigma \min \{d(x, z), d(z, y)\} &\leq 4\sigma \left[ \frac{1}{d(x, z)} + \frac{1}{d(z, y)} \right]^{-1} \leq 2\sigma \sqrt{d(x, z)} \sqrt{d(z, y)} \\ &\leq \sigma [d(x, z) + d(z, y)] \leq 2\sigma \max \{d(x, z), d(z, y)\} \end{aligned}$$

✎PROOF: These follow directly from Corollary B.8 (page 34).

⇒

## 4.2.2 Properties of power distance spaces

The *power triangle inequality* property of a *power distance space* axiomatically endows a metric with an upper bound. Lemma 4.9 (next) demonstrates that there is a complementary lower bound somewhat similar in form to the *power triangle inequality* upper bound. In the special case where  $2\sigma = 2^{\frac{1}{p}}$ , the lower bound helps provide a simple proof of the *continuity* of a large class of *power distance functions* (Theorem 4.18 page 23). The inequality  $2\sigma \leq 2^{\frac{1}{p}}$  is a special relation in this paper and appears repeatedly in this paper; it appears as an inequality in Lemma 4.13 (page 20), Corollary 4.12 (page 20) and Corollary 4.14 (page 21), and as an equality in Lemma 4.9 (next) and Theorem 4.18 (page 23). It is plotted in Figure 1 (page 18).






Figure 1:  $\sigma = \frac{1}{2}(2^{\frac{1}{p}}) = 2^{\frac{1}{p}-1}$  or  $p = \frac{\ln 2}{\ln(2\sigma)}$  (see Lemma 4.9 page 18, Lemma 4.13 page 20, Corollary 4.14 page 21, Corollary 4.12 page 20, and Theorem 4.18 page 23).

**Lemma 4.9** <sup>40</sup> Let  $(X, d, p, \sigma)$  be a POWER TRIANGLE TRIANGLE SPACE (Definition 4.3 page 16). Let  $|\cdot|$  be the ABSOLUTE VALUE function (Definition 2.10 page 5). Let  $\max\{x, y\}$  be the maximum and  $\min\{x, y\}$  the minimum of any  $x, y \in \mathbb{R}^*$ . Then, for all  $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$ ,

1.  $d^p(x, y) \geq \max\left\{0, \frac{2}{(2\sigma)^p}d^p(x, z) - d^p(z, y), \frac{2}{(2\sigma)^p}d^p(y, z) - d^p(z, x)\right\} \quad \forall x, y, z \in X \quad \text{and}$
2.  $d(x, y) \geq |d(x, z) - d(z, y)| \quad \text{if } p \neq 0 \text{ and } 2\sigma = 2^{\frac{1}{p}} \quad \forall x, y, z \in X.$

PROOF:

(1) lemma:  $\frac{2}{(2\sigma)^p}d^p(x, z) - d^p(z, y) \leq d^p(x, y) \quad \forall (p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$ : Proof:

$$\begin{aligned} \frac{2}{(2\sigma)^p}d^p(x, z) - d^p(z, y) &\leq \frac{2}{(2\sigma)^p} \left[ 2\sigma \left[ \frac{1}{2}d^p(x, y) + \frac{1}{2}d^p(y, z) \right]^{\frac{1}{p}} \right]^p - d^p(z, y) && \text{by power triangle inequality} \\ &= \frac{2(2\sigma)^p}{(2\sigma)^p} \left[ \frac{1}{2}d^p(x, y) + \frac{1}{2}d^p(y, z) \right] - d^p(z, y) \\ &= [d^p(x, y) + d^p(y, z)] - d^p(y, z) && \text{by symmetric property of } d \\ &= d^p(x, y) \end{aligned}$$

(2) Proof for  $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$  case:

$$\begin{aligned} d^p(x, y) &\geq \frac{2}{(2\sigma)^p}d^p(x, z) - d^p(z, y) && \text{by (1) lemma} \\ d^p(x, y) = d^p(y, x) &\geq \frac{2}{(2\sigma)^p}d^p(y, z) - d^p(z, x) && \text{by commutative property of } d \text{ and (1) lemma} \\ d^p(x, y) &\geq 0 && \text{by non-negative property of } d \text{ (Definition 3.1 page 6)} \end{aligned}$$

The rest follows because  $g(x) \triangleq x^{\frac{1}{p}}$  is strictly monotone in  $\mathbb{R}^{\mathbb{R}}$ .

<sup>40</sup> in metric space  $((p, \sigma) = (1, 1))$ : [Dieudonné(1969)] page 28, [Michel and Herget(1993)] page 266, [Berberian(1961)] page 37 (Theorem II.4.1)



(3) Proof for  $2\sigma = 2^{\frac{1}{p}}$  case:

$$\begin{aligned}
 d(x, y) &\geq \max \left\{ 0, \frac{2}{(2\sigma)^p} d^p(x, z) - d^p(z, y), \frac{2}{(2\sigma)^p} d^p(y, z) - d^p(z, x) \right\}^{\frac{1}{p}} && \text{by item (2) (page 18)} \\
 &= \max \{0, d(x, z) - d(z, y), d(y, z) - d(z, x)\} && \text{by } 2\sigma = 2^{\frac{1}{p}} \text{ hypothesis } \iff \frac{2}{(2\sigma)^p} = 1 \\
 &= \max \{0, (d(x, z) - d(z, y)), -(d(x, z) - d(z, y))\} && \text{by symmetric property of } d \\
 &= |(d(x, z) - d(z, y))|
 \end{aligned}$$

⇒

**Theorem 4.10** Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE (Definition 4.3 page 16). Let  $B$  be an OPEN BALL (Definition 3.5 page 6) on  $(X, d)$ . Then for all  $(p, \sigma) \in (\mathbb{R}^* \setminus \{0\}) \times \mathbb{R}^+$ ,

$$\left\{ \begin{array}{l} A. \ 2\sigma \leq 2^{\frac{1}{p}} \\ B. \ q \in B(\theta, r) \end{array} \right\} \text{ and } \left\{ \begin{array}{l} 1. \ \exists r_q \in \mathbb{R}^+ \text{ such that} \\ \quad B(q, r_q) \subseteq B(\theta, r) \end{array} \right\} \implies \left\{ \begin{array}{l} B. \ q \in B(\theta, r) \end{array} \right\}$$

✎PROOF:

(1) lemma:

$$\begin{aligned}
 q \in B(\theta, r) &\iff d(\theta, q) < r && \text{by definition of open ball (Definition 3.5 page 6)} \\
 &\iff 0 < r - d(\theta, q) && \text{by field property of real numbers} \\
 &\iff \exists r_q \in \mathbb{R}^+ \text{ such that } 0 < r_q < r - d(\theta, q) && \text{by The Archimedean Property}^{41}
 \end{aligned}$$

(2) Proof that (A), (B)  $\implies$  (1):

$$\begin{aligned}
 B(q, r_q) &\triangleq \{x \in X \mid d(q, x) < r_q\} && \text{by definition of open ball (Definition 3.5 page 6)} \\
 &= \{x \in X \mid d^p(q, x) < r_q^p \in \mathbb{R}^+\} && \text{because } f(x) \triangleq x^p \text{ is monotone} \\
 &\subseteq \{x \in X \mid d^p(q, x) < r^p - d^p(\theta, q)\} && \text{by hypothesis B and (1) lemma page 19} \\
 &= \{x \in X \mid d^p(\theta, q) + d^p(q, x) < r^p\} && \text{by field property of real numbers} \\
 &= \left\{x \in X \mid [d^p(\theta, q) + d^p(q, x)]^{\frac{1}{p}} < r\right\} && \text{because } f(x) \triangleq x^{\frac{1}{p}} \text{ is monotone} \\
 &\subseteq \left\{x \in X \mid 2^{1-\frac{1}{p}}\sigma [d^p(\theta, q) + d^p(q, x)]^{\frac{1}{p}} < r\right\} && \text{by hypothesis A which implies } 2^{1-\frac{1}{p}}\sigma \leq 1 \\
 &= \left\{x \in X \mid 2\sigma [^{\frac{1}{2}}d(\theta, q) x + ^{\frac{1}{2}}d^p(q, x)]^{\frac{1}{p}} < r\right\} && \text{because } 2^{1-\frac{1}{p}}\sigma = 2\sigma(^{\frac{1}{2}})^{\frac{1}{p}} \\
 &\triangleq \{x \in X \mid \tau(p, \sigma, \theta, x, q) < r\} && \text{by definition of } \tau \text{ (Definition 4.1 page 15)} \\
 &\subseteq \{x \in X \mid d(\theta, x) < r\} && \text{by definition of } (X, d, p, \sigma) \text{ (Definition 4.3 page 16)} \\
 &\triangleq B(\theta, r) && \text{by definition of open ball (Definition 3.5 page 6)}
 \end{aligned}$$

(3) Proof that (B)  $\iff$  (1):

$$\begin{aligned}
 q &\in \{x \in X \mid d(q, x) = 0\} && \text{by nondegenerate property (Definition 3.1 page 6)} \\
 &\subseteq \{x \in X \mid d(q, x) < r_q\} && \text{because } r_q > 0 \\
 &\triangleq B(q, r_q) && \text{by definition of open ball (Definition 3.5 page 6)} \\
 &\subseteq B(\theta, r) && \text{by hypothesis 2}
 \end{aligned}$$

<sup>41</sup> ✎ [Aliprantis and Burkinshaw(1998)] page 17 (Theorem 3.3 (“The Archimedean Property”) and Theorem 3.4), ✎ [Zorich(2004)] page 53 (6° (“The principle of Archimedes”) and 7°)





**Corollary 4.11** Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Then for all  $(p, \sigma) \in (\mathbb{R}^* \setminus \{0\}) \times \mathbb{R}^+$ ,

$$\left\{ 2\sigma \leq 2^{\frac{1}{p}} \right\} \implies \left\{ \text{every OPEN BALL in } (X, d) \text{ is OPEN} \right\}$$

PROOF: This follows from Theorem 4.10 (page 19) and Theorem 3.10 (page 8).

**Corollary 4.12** Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Let  $\mathbf{B}$  be the set of all OPEN BALLS in  $(X, d)$ . Then for all  $(p, \sigma) \in (\mathbb{R}^* \setminus \{0\}) \times \mathbb{R}^+$ ,

$$\left\{ 2\sigma \leq 2^{\frac{1}{p}} \right\} \implies \left\{ \mathbf{B} \text{ is a BASE for } (X, T) \right\}$$

PROOF:

- (1) The set of all *open balls* in  $(X, d)$  is a *base* for  $(X, T)$  by Corollary 4.11 (page 20) and Theorem A.4 (page 27).
- (2)  $T$  is a topology on  $X$  by Definition A.3 (page 27).




Figure 2: *open set* (see Lemma 4.13 page 20)

Lemma 4.13 (next) demonstrates that every point in an open set is contained in an open ball that is contained in the original open set (see also Figure 2 page 20).

**Lemma 4.13** Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Let  $\mathbf{B}$  be an OPEN BALL on  $(X, d)$ . Then for all  $(p, \sigma) \in (\mathbb{R}^* \setminus \{0\}) \times \mathbb{R}^+$ ,

$$\left\{ \begin{array}{l} \text{A. } 2\sigma \leq 2^{\frac{1}{p}} \text{ and} \\ \text{B. } U \text{ is OPEN in } (X, d) \end{array} \right\} \implies \left\{ \begin{array}{l} \text{1. } \forall x \in U, \exists r \in \mathbb{R}^+ \text{ such that} \\ \quad \mathbf{B}(x, r) \subseteq U \end{array} \right\} \implies \left\{ \begin{array}{l} \text{B. } U \text{ is} \\ \quad \text{OPEN in } (X, d) \end{array} \right\}$$

PROOF:

- (1) Proof that for  $((A), (B)) \implies (1)$ :

$$\begin{aligned} U &= \bigcup \{ \mathbf{B}(x_\gamma, r_\gamma) \mid \mathbf{B}(x_\gamma, r_\gamma) \subseteq U \} \\ &\supseteq \mathbf{B}(x, r) \end{aligned}$$

by left hypothesis and Corollary 4.12 page 20

because  $x$  must be in one of those balls in  $U$



(2) Proof that  $((B) \Leftarrow (1))$  case:

$$\begin{aligned}
 U &= \bigcup \{x \in X \mid x \in U\} && \text{by definition of union operation } \bigcup \\
 &= \bigcup \{B(x, r) \mid x \in U \text{ and } B(x, r) \subseteq U\} && \text{by hypothesis (1)} \\
 &\Rightarrow U \text{ is open} && \text{by Corollary 4.12 page 20 and Corollary 3.8 page 8}
 \end{aligned}$$

⇒

**Corollary 4.14** <sup>42</sup> Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Let  $B$  be an OPEN BALL on  $(X, d)$ . Then for all  $(p, \sigma) \in (\mathbb{R}^* \setminus \{0\}) \times \mathbb{R}^+$ ,

$$\left\{ 2\sigma \leq 2^{\frac{1}{p}} \right\} \Rightarrow \left\{ \text{every OPEN BALL } B(x, r) \text{ in } (X, d) \text{ is OPEN} \right\}$$

✎PROOF:

The union of any set of open balls is open by Corollary 4.12 page 20  
 $\Rightarrow$  the union of a set of just one open ball is open  
 $\Rightarrow$  every open ball is open.

⇒

**Theorem 4.15** <sup>43</sup> Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Let  $(X, T)$  be a TOPOLOGICAL SPACE INDUCED BY  $(X, d)$ . Let  $(x_n \in X)_{n \in \mathbb{Z}}$  be a sequence in  $(X, d)$ .

$$\underbrace{(x_n) \text{ converges to a limit } x}_{\text{(Definition A.16 page 28)}} \Leftrightarrow \left\{ \begin{array}{l} \text{for any } \varepsilon \in \mathbb{R}^+, \text{ there exists } N \in \mathbb{Z} \\ \text{such that for all } n > N, \quad d(x_n, x) < \varepsilon \end{array} \right\}$$

✎PROOF:

$$\begin{aligned}
 (x_n) \rightarrow x &\Leftrightarrow x_n \in U \quad \forall U \in \mathcal{N}_x, n > N && \text{by Definition A.16 page 28} \\
 &\Leftrightarrow \exists B(x, \varepsilon) \text{ such that } x_n \in B(x, \varepsilon) \forall n > N && \text{by Lemma 4.13 page 20} \\
 &\Leftrightarrow d(x_n, x) < \varepsilon && \text{by Definition 3.5 page 6}
 \end{aligned}$$

⇒

In *distance spaces* (Definition 3.1 page 6), not all *convergent* sequences are *Cauchy* (Example 3.22 page 13). However in a distance space with any *power triangle inequality* (Definition 4.3 page 16), all *convergent* sequences are *Cauchy* (next theorem).

**Theorem 4.16** <sup>44</sup> Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Let  $B$  be an OPEN BALL on  $(X, d)$ . For any  $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$ ,

$$\left\{ \begin{array}{l} (x_n) \text{ is CONVERGENT} \\ \text{in } (X, d) \end{array} \right\} \Rightarrow \left\{ (x_n) \text{ is CAUCHY} \right\} \Rightarrow \left\{ \begin{array}{l} (x_n) \text{ is BOUNDED} \\ \text{in } (X, d) \end{array} \right\}$$

<sup>42</sup> in *metric space*  $((p, \sigma) = (1, 1))$ : ✎ [Rosenlicht(1968)] pages 40–41, ✎ [Aliprantis and Burkinshaw(1998)] page 35

<sup>43</sup> in *metric space*: ✎ [Rosenlicht(1968)] page 45, ✎ [Giles(1987)] page 37 (3.2 Definition)

<sup>44</sup> in *metric space*: ✎ [Giles(1987)] page 49 (Theorem 3.30), ✎ [Rosenlicht(1968)] page 51, ✎ [Apostol(1975)] pages 72–73 (Theorem 4.6)



PROOF:

(1) Proof that *convergent*  $\implies$  *Cauchy*:

$$\begin{aligned}
 d(x_n, x_m) &\leq \tau(p, \sigma; x_n, x_m, x) && \text{by definition of } \textit{power triangle inequality} \text{ (Definition 4.3 page 16)} \\
 &\triangleq 2\sigma \left[ \frac{1}{2}d^p(x_n, x) + \frac{1}{2}d^p(x_m, x) \right]^{\frac{1}{p}} && \text{by definition of } \textit{power triangle function} \text{ (Definition 4.1 page 15)} \\
 &< 2\sigma \left[ \frac{1}{2}\varepsilon^p + \frac{1}{2}\varepsilon^p \right]^{\frac{1}{p}} && \text{by } \textit{convergence hypothesis} \text{ (Definition A.16 page 28)} \\
 &= 2\sigma\varepsilon && \text{by definition of } \textit{convergence} \text{ (Definition A.16 page 28)} \\
 &\implies \textit{Cauchy} && \text{by definition of } \textit{Cauchy} \text{ (Definition 3.12 page 9)} \\
 \\
 d(x_n, x_m) &\leq \tau(\infty, \sigma; x_n, x_m, x) && \text{by definition of } \textit{power triangle inequality at } p = \infty \\
 &= 2\sigma \max \{d(x_n, x), d(x_m, x)\} && \text{by Corollary 4.7 (page 17)} \\
 &= 2\sigma \max \{\varepsilon, \varepsilon\} && \text{by } \textit{convergent hypothesis} \text{ (Definition A.16 page 28)} \\
 &= 2\sigma\varepsilon && \text{by definition of } \max \\
 \\
 d(x_n, x_m) &\leq \tau(-\infty, \sigma; x_n, x_m, x) && \text{by definition of } \textit{power triangle inequality at } p = -\infty \\
 &= 2\sigma \min \{d(x_n, x), d(x_m, x)\} && \text{by Corollary 4.7 (page 17)} \\
 &= 2\sigma \min \{\varepsilon, \varepsilon\} && \text{by } \textit{convergent hypothesis} \text{ (Definition A.16 page 28)} \\
 &= 2\sigma\varepsilon && \text{by definition of } \min \\
 \\
 d(x_n, x_m) &\leq \tau(0, \sigma; x_n, x_m, x) && \text{by definition of } \textit{power triangle inequality at } p = 0 \\
 &= 2\sigma \sqrt{d(x_n, x)} \sqrt{d(x_m, x)} && \text{by Corollary 4.7 (page 17)} \\
 &= 2\sigma \sqrt{\varepsilon} \sqrt{\varepsilon} && \text{by } \textit{convergent hypothesis} \text{ (Definition A.16 page 28)} \\
 &= 2\sigma\varepsilon && \text{by property of } \mathbb{R}
 \end{aligned}$$

(2) Proof that *Cauchy*  $\implies$  *bounded*: by Proposition 3.14 (page 9).

⇒

**Theorem 4.17** <sup>45</sup> Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Let  $f \in \mathbb{Z}^{\mathbb{Z}}$  be a STRICTLY MONOTONE function such that  $f(n) < f(n+1)$ . For any  $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$

$$\left\{ \begin{array}{l} 1. \ (x_n)_{n \in \mathbb{Z}} \text{ is CAUCHY} \\ 2. \ (x_{f(n)})_{n \in \mathbb{Z}} \text{ is CONVERGENT} \end{array} \right\} \text{ and } \implies \{ (x_n)_{n \in \mathbb{Z}} \text{ is CONVERGENT.} \}$$

PROOF:

$$\begin{aligned}
 d(x_n, x) &= d(x, x_n) && \text{by } \textit{symmetric property of } d \\
 &\leq \tau(p, \sigma; x, x_n, x_{f(n)}) && \text{by definition of } \textit{power triangle inequality} \text{ (Definition 4.3 page 16)} \\
 &\triangleq 2\sigma \left[ \frac{1}{2}d^p(x, x_{f(n)}) + \frac{1}{2}d^p(x_{f(n)}, x_n) \right]^{\frac{1}{p}} && \text{by definition of } \textit{power triangle function} \text{ (Definition 4.1 page 15)} \\
 &= 2\sigma \left[ \frac{1}{2}\varepsilon + \frac{1}{2}d^p(x_{f(n)}, x_n) \right]^{\frac{1}{p}} && \text{by left hypothesis 2} \\
 &= 2\sigma \left[ \frac{1}{2}\varepsilon^p + \frac{1}{2}\varepsilon^p \right]^{\frac{1}{p}} && \text{by left hypothesis 1} \\
 &= 2\sigma\varepsilon && \\
 &\implies \textit{convergent} && \text{by definition of } \textit{convergent} \text{ (Definition A.16 page 28)}
 \end{aligned}$$

<sup>45</sup> in metric space: [Rosenlicht(1968)] page 52





**Theorem 4.18** <sup>46</sup> Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Let  $(\mathbb{R}, q)$  be a metric space of real numbers with the usual metric  $q(x, y) \triangleq |x - y|$ . Then

$$\left\{ 2\sigma = 2^{\frac{1}{p}} \right\} \implies \left\{ d \text{ is CONTINUOUS in } (\mathbb{R}, q) \right\}$$

PROOF:

$$\begin{aligned} |d(x, y) - d(x_n, y_n)| &\leq |d(x, y) - d(x_n, y)| + |d(x_n, y) - d(x_n, y_n)| && \text{by triangle inequality of } (\mathbb{R}, |x - y|) \\ &= |d(x, y) - d(y, x_n)| + |d(y, x_n) - d(x_n, y_n)| && \text{by commutative property of } d \text{ (Definition 3.1 page 6)} \\ &\leq d(x, x_n) + d(y, y_n) && \text{by } 2\sigma = 2^{\frac{1}{p}} \text{ and Lemma 4.9 (page 18)} \\ &= 0 \quad \text{as } n \rightarrow \infty \end{aligned}$$



In *distance spaces* and *topological spaces*, limits of convergent sequences are in general *not unique* (Example 3.21 page 12, Example A.17 page 29). However Theorem 4.19 (next) demonstrates that, in a *power distance space*, limits are unique.

**Theorem 4.19** (Uniqueness of limit) <sup>47</sup> Let  $(X, d, p, \sigma)$  be a POWER DISTANCE SPACE. Let  $x, y \in X$  and let  $(x_n \in X)$  be an  $X$ -valued sequence.

$$\left\{ \begin{array}{l} 1. \left\{ ((x_n), (y_n)) \rightarrow (x, y) \right\} \text{ and} \\ 2. (p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+ \end{array} \right\} \implies \{x = y\}$$

PROOF:

(1) lemma: Proof that for all  $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$  and for any  $\varepsilon \in \mathbb{R}^+$ , there exists  $N$  such that  $d(x, y) < 2\sigma\varepsilon$ :

$$\begin{aligned} d(x, y) &\leq \tau(p, \sigma; x, y, x_n) && \text{by definition of power triangle inequality (Definition 4.3 page 16)} \\ &\triangleq 2\sigma \left[ \frac{1}{2}d^p(x, x_n) + \frac{1}{2}d^p(x_n, y) \right]^{\frac{1}{p}} && \text{by definition of power triangle function (Definition 4.1 page 15)} \\ &< 2\sigma \left[ \frac{1}{2}\varepsilon^p + \frac{1}{2}\varepsilon^p \right]^{\frac{1}{p}} && \text{by left hypothesis and for } p \in \mathbb{R}^* \setminus \{-\infty, 0, \infty\} \\ &= 2\sigma\varepsilon \\ d(x, y) &\leq \tau(\infty, \sigma; x, y, x_n) && \text{by definition of power triangle inequality at } p = \infty \\ &= 2\sigma \max \{d(x, x_n), d(x_n, y)\} && \text{by Corollary 4.7 (page 17)} \\ &< 2\sigma\varepsilon && \text{by left hypothesis} \\ d(x, y) &\leq \tau(-\infty, \sigma; x, y, x_n) && \text{by definition of power triangle inequality at } p = -\infty \\ &= 2\sigma \min \{d(x, x_n), d(x_n, y)\} && \text{by Corollary 4.7 (page 17)} \\ &< 2\sigma\varepsilon && \text{by left hypothesis} \\ d(x, y) &\leq \tau(0, \sigma; x, y, x_n) && \text{by definition of power triangle inequality at } p = 0 \\ &= 2\sigma \sqrt{d(x, x_n)} \sqrt{d(x_n, y)} && \text{by Corollary 4.7 (page 17)} \\ &= 2\sigma \sqrt{\varepsilon} \sqrt{\varepsilon} && \text{by left hypothesis} \\ &< 2\sigma\varepsilon && \text{by property of real numbers} \end{aligned}$$

<sup>46</sup> in metric space  $((p, \sigma) = (1, 1)$  case): [Berberian(1961)] page 37 (Theorem II.4.1)

<sup>47</sup> in metric space: [Rosenlicht(1968)] page 46, [Thomson et al.(2008)Thomson, Bruckner, and Bruckner] page 32 (Theorem 2.8)



(2) Proof that  $x = y$  (proof by contradiction):

$$\begin{aligned}
 x \neq y &\implies d(x, y) \neq 0 && \text{by the } \textit{nondegenerate} \text{ property of } d \text{ (Definition 3.1 page 6)} \\
 &\implies d(x, y) > 0 && \text{by } \textit{non-negative} \text{ property of } d \text{ (Definition 3.1 page 6)} \\
 &\implies \exists \varepsilon \text{ such that } d(x, y) > 2\sigma\varepsilon \\
 &\implies \textit{contradiction} \text{ to (1) lemma page 23} \\
 &\implies d(x, y) = 0 \\
 &\implies x = y
 \end{aligned}$$

⇒

### 4.3 Examples

It is not always possible to find a *triangle relation* (Definition 4.3 page 16)  $\mathcal{O}(p, \sigma; d)$  that holds in every *distance space* (Definition 3.1 page 6), as demonstrated by Example 4.20 and Example 4.21 (next two examples).

**Example 4.20** Let  $d(x, y) \in \mathbb{R}^{\mathbb{R} \times \mathbb{R}}$  be defined such that

$$d(x, y) \triangleq \left\{ \begin{array}{ll} y & \forall (x, y) \in \{4\} \times (0 : 2] \quad (\textit{vertical half-open interval}) \\ x & \forall (x, y) \in (0 : 2] \times \{4\} \quad (\textit{horizontal half-open interval}) \\ |x - y| & \text{otherwise} \quad (\textit{Euclidean}) \end{array} \right\}.$$

Note the following about the pair  $(\mathbb{R}, d)$ :

- (1) By Example 3.21 (page 12),  $(\mathbb{R}, d)$  is a *distance space*, but not a *metric space*—that is, the *triangle relation*  $\mathcal{O}(1, 1; d)$  does not hold in  $(\mathbb{R}, d)$ .
- (2) Observe further that  $(\mathbb{R}, d)$  is *not a power distance space*. In particular, the *triangle relation*  $\mathcal{O}(p, \sigma; d)$  does not hold in  $(\mathbb{R}, d)$  for any finite value of  $\sigma$  (does not hold for any  $\sigma \in \mathbb{R}^+$ ):

$$\begin{aligned}
 d(0, 4) = 4 \not\leq 0 &= \lim_{\varepsilon \rightarrow 0} 2\sigma\varepsilon = \lim_{\varepsilon \rightarrow 0} 2\sigma \left[ \frac{1}{2}|0 - \varepsilon|^p + \frac{1}{2}\varepsilon^p \right]^{\frac{1}{p}} \\
 &\triangleq \lim_{\varepsilon \rightarrow 0} 2\sigma \left[ \frac{1}{2}d^p(0, \varepsilon) + \frac{1}{2}d^p(\varepsilon, 4) \right]^{\frac{1}{p}} \triangleq \lim_{\varepsilon \rightarrow 0} \mathcal{O}(p, \sigma; 0, 4, \varepsilon; d)
 \end{aligned}$$

**Example 4.21** Let  $d(x, y) \in \mathbb{R}^{\mathbb{R} \times \mathbb{R}}$  be defined such that

$$d(x, y) \triangleq \left\{ \begin{array}{ll} |x - y| & \text{for } x = 0 \text{ or } y = 0 \text{ or } x = y \quad (\textit{Euclidean}) \\ 1 & \text{otherwise} \quad (\textit{discrete}) \end{array} \right\}.$$

Note the following about the pair  $(\mathbb{R}, d)$ :

- (1) By Example 3.22 (page 13),  $(\mathbb{R}, d)$  is a *distance space*, but not a *metric space*—that is, the *triangle relation*  $\mathcal{O}(1, 1; d)$  does not hold in  $(\mathbb{R}, d)$ .
- (2) Observe further that  $(\mathbb{R}, d)$  is *not a power distance space*—that is, the *triangle relation*  $\mathcal{O}(p, \sigma; d)$  does not hold in  $(\mathbb{R}, d)$  for any value of  $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$ .

(a) Proof that  $\mathcal{O}(p, \sigma; d)$  does not hold for any  $(p, \sigma) \in \{\infty\} \times \mathbb{R}^+$ :

$$\begin{aligned}
 \lim_{n, m \rightarrow \infty} d(\frac{1}{n}, \frac{1}{m}) &\triangleq 1 \not\leq 0 = 2\sigma \max\{0, 0\} && \text{by definition of } d \\
 &= 2\sigma \lim_{n, m \rightarrow \infty} \max\{d(\frac{1}{n}, 0), d(0, \frac{1}{m})\} && \text{by Corollary 4.7 (page 17)} \\
 &\geq \lim_{n, m \rightarrow \infty} 2\sigma \left[ \frac{1}{2}d^p(\frac{1}{n}, 0) + \frac{1}{2}d^p(0, \frac{1}{m}) \right]^{\frac{1}{p}} && \text{by Corollary 4.6 (page 16)} \\
 &\triangleq \lim_{n, m \rightarrow \infty} \tau(p, \sigma, \frac{1}{n}, \frac{1}{m}, 0) && \text{by definition of } \tau \text{ (Definition 4.1 page 15)}
 \end{aligned}$$





- (b) Proof that  $\mathcal{Q}(p, \sigma; d)$  does not hold for any  $(p, \sigma) \in \mathbb{R}^* \times \mathbb{R}^+$ : By Corollary 4.6 (page 16), the *triangle function* (Definition 4.1 page 15)  $\tau(p, \sigma; x, y, z; d)$  is *continuous* and *strictly monotone* in  $(\mathbb{R}, |\cdot|, \leq)$  with respect to the variable  $p$ . Item 2a demonstrates that  $\mathcal{Q}(p, \sigma; d)$  fails to hold at the best case of  $p = \infty$ , and so by Corollary 4.6, it doesn't hold for any other value of  $p \in \mathbb{R}^*$  either.

**Example 4.22** Let  $d$  be a function in  $\mathbb{R}^{\mathbb{R} \times \mathbb{R}}$  such that

$$d(x, y) \triangleq \left\{ \begin{array}{ll} 2|x - y| & \forall (x, y) \in \{(0, 1), (1, 0)\} \quad (\text{dilated Euclidean}) \\ |x - y| & \text{otherwise} \quad (\text{Euclidean}) \end{array} \right\}.$$

Note the following about the pair  $(\mathbb{R}, d)$ :

- (1) By Example 3.23 (page 14),  $(\mathbb{R}, d)$  is a *distance space*, but not a *metric space*—that is, the *triangle relation*  $\mathcal{Q}(1, 1; d)$  does not hold in  $(\mathbb{R}, d)$ .
- (2) But observe further that  $(\mathbb{R}, d, 1, 2)$  is a *power distance space*:

- (a) Proof that  $\mathcal{Q}(1, 2; d)$  (Definition 4.3 page 16) holds for all  $(x, y) \in \{(0, 1), (1, 0)\}$ :

$$\begin{aligned} d(1, 0) &= d(0, 1) \triangleq 2|0 - 1| = 2 && \text{by definition of } d \\ &\leq 2 \leq 2(|0 - z| + |z - 1|) \quad \forall z \in \mathbb{R} && \text{by definition of } |\cdot| \text{ (Definition 2.10 page 5)} \\ &= 2\sigma(\tfrac{1}{2}|0 - z|^p + \tfrac{1}{2}|z - 1|^p)^{\frac{1}{p}} \quad \forall z \in \mathbb{R} && \text{for } (p, \sigma) = (1, 2) \\ &\triangleq 2\sigma(\tfrac{1}{2}d^p(0, z) + d^p(z, 1))^{\frac{1}{p}} \quad \forall z \in \mathbb{R} && \text{for } (p, \sigma) = (1, 2) \text{ and by definition of } d \\ &\triangleq \tau(1, 2; 0, 1, z) && \text{by definition of } \tau \text{ (Definition 4.1 page 15)} \end{aligned}$$

- (b) Proof that  $\mathcal{Q}(1, 2; d)$  holds for all other  $(x, y) \in \mathbb{R}^* \times \mathbb{R}^+$ :

$$\begin{aligned} d(x, y) &\triangleq 2|x - y| && \text{by definition of } d \\ &\leq (|x - z| + |z - y|) && \text{by property of } \textit{Euclidean metric spaces} \\ &= 2\sigma(\tfrac{1}{2}|0 - z|^p + \tfrac{1}{2}|z - 1|^p)^{\frac{1}{p}} && \text{for } (p, \sigma) = (1, 1) \\ &\triangleq \tau(1, 1; x, y, z) && \text{by definition of } \tau \text{ (Definition 4.1 page 15)} \\ &\leq \tau(1, 2; x, y, z) && \text{by Corollary 4.6 (page 16)} \end{aligned}$$

- (3) In  $(X, d)$ , the limits of *convergent* sequences are *unique*. This follows directly from the fact that  $(\mathbb{R}, d, 1, 2)$  is a *power distance space* (item (2) page 25) and by Theorem 4.19 page 23.
- (4) In  $(X, d)$ , *convergent* sequences are *Cauchy*. This follows directly from the fact that  $(\mathbb{R}, d, 1, 2)$  is a *power distance space* (item (2) page 25) and by Theorem 4.16 page 21.

**Example 4.23** Let  $d$  be a function in  $\mathbb{R}^{\mathbb{R} \times \mathbb{R}}$  such that  $d(x, y) \triangleq (x - y)^2$ . Note the following about the pair  $(\mathbb{R}, d)$ :

- (1) It was demonstrated in Example 3.24 (page 15) that  $(\mathbb{R}, d)$  is a *distance space*, but that it is *not a metric space* because the *triangle inequality* does not hold.
- (2) However, the tuple  $(\mathbb{R}, d, p, \sigma)$  is a *power distance space* (Definition 4.3 page 16) for any  $(p, \sigma) \in \mathbb{R}^* \times [2 : \infty)$ : In particular, for all  $x, y, z \in \mathbb{R}$ , the *power triangle inequality* (Definition 4.3 page 16) must hold. The “worst case” for this is when a third point  $z$  is exactly “halfway between”  $x$  and  $y$  in  $d(x, y)$ ;



that is, when  $z = \frac{x+y}{2}$ :

$$\begin{aligned}
 (x-y)^2 &\triangleq d(x, y) && \text{by definition of } d \\
 &\leq \tau(p, \sigma; x, y, z; d) && \text{by definition } \textit{power triangle inequality} \\
 &\triangleq 2\sigma \left[ \frac{1}{2}d^p(x, z) + \frac{1}{2}d^p(z, y) \right]^{\frac{1}{p}} && \text{by definition } \tau \text{ (Definition 4.1 page 15)} \\
 &\triangleq 2\sigma \left[ \frac{1}{2}(x-z)^{2p} + \frac{1}{2}(z-y)^{2p} \right]^{\frac{1}{p}} && \text{by definition of } d \\
 &= 2\sigma \left[ \frac{1}{2}|x-z|^{2p} + \frac{1}{2}|z-y|^{2p} \right]^{\frac{1}{p}} && \text{because } (x)^2 = |x|^2 \text{ for all } x \in \mathbb{R} \\
 &= 2\sigma \left[ \frac{1}{2} \left| x - \frac{x+y}{2} \right|^{2p} + \frac{1}{2} \left| \frac{x+y}{2} - y \right|^{2p} \right]^{\frac{1}{p}} && \text{because } z = \frac{x+y}{2} \text{ is the "worst case" scenario} \\
 &= 2\sigma \left[ \frac{1}{2} \left| \frac{y-x}{2} \right|^{2p} + \frac{1}{2} \left| \frac{x-y}{2} \right|^{2p} \right]^{\frac{1}{p}} \\
 &= 2\sigma \left[ \left| \frac{x-y}{2} \right|^{2p} \right]^{\frac{1}{p}} = \frac{2\sigma}{4} |x-y|^2 \\
 &\implies (p, \sigma) \in \mathbb{R}^* \times [2 : \infty)
 \end{aligned}$$

- (3) The *power distance function*  $d$  is *continuous* in  $(\mathbb{R}, d, p, \sigma)$  for any  $(p, \sigma)$  such that  $\sigma \geq 2$  and  $2\sigma = p^{\frac{1}{p}}$ . This follows directly from Theorem 4.18 (page 23).

## Appendix A Topological Spaces

**Definition A.1**<sup>48</sup> Let  $\Gamma$  be a set with an arbitrary (possibly uncountable) number of elements. Let  $2^X$  be the *power set* of a set  $X$  (Definition 2.2 page 4). A family of sets  $\mathcal{T} \subseteq 2^X$  is a **topology** on  $X$  if

1.  $\emptyset \in \mathcal{T}$  and
2.  $X \in \mathcal{T}$  and
3.  $U, V \in \mathcal{T} \implies U \cap V \in \mathcal{T}$  and
4.  $\{U_\gamma \mid \gamma \in \Gamma\} \subseteq \mathcal{T} \implies \bigcup_{\gamma \in \Gamma} U_\gamma \in \mathcal{T}$ .

The ordered pair  $(X, \mathcal{T})$  is a *topological space* if  $\mathcal{T}$  is a *topology* on  $X$ . A set  $U$  is **open** in  $(X, \mathcal{T})$  if  $U$  is any element of  $\mathcal{T}$ . A set  $D$  is **closed** in  $(X, \mathcal{T})$  if  $D^c$  is *open* in  $(X, \mathcal{T})$ .

Just as the power set  $2^X$  and the set  $\{\emptyset, X\}$  are algebras of sets on a set  $X$ , so also are these sets topologies on  $X$  (next example):

**Example A.2**<sup>49</sup> Let  $\mathcal{T}(X)$  be the set of topologies on a set  $X$  and  $2^X$  the *power set* (Definition 2.2 page 4) on  $X$ .

- $\{\emptyset, X\}$  is a *topology* in  $\mathcal{T}(X)$  (indiscrete topology or trivial topology)  
 $2^X$  is a *topology* in  $\mathcal{T}(X)$  (discrete topology)

<sup>48</sup> [Munkres(2000)] page 76, [Riesz(1909)], [Hausdorff(1914)], [Tietze(1923)], [Hausdorff(1937)] page 258

<sup>49</sup> [Munkres(2000)], page 77, [Kubrusly(2011)] page 107 (Example 3.J), [Steen and Seebach(1978)] pages 42–43 (II.4), [DiBenedetto(2002)] page 18



**Definition A.3** <sup>50</sup> Let  $(X, T)$  be a *topological space*. A set  $B \subseteq 2^X$  is a **base** for  $T$  if

1.  $B \subseteq T$  and
2.  $\forall U \in T, \exists \{B_\gamma \in B\}$  such that  $U = \bigcup_\gamma B_\gamma$

**Theorem A.4** <sup>51</sup> Let  $(X, T)$  be a TOPOLOGICAL SPACE. Let  $B$  be a subset of  $2^X$  such that  $B \subseteq 2^X$ .  
 $\{B \text{ is a BASE for } T\} \iff \left\{ \begin{array}{l} \text{For every } x \in X \text{ and for every OPEN SET } U \text{ containing } x, \\ \text{there exists } B_x \in B \text{ such that } x \in B_x \subseteq U. \end{array} \right\}$

**Theorem A.5** <sup>52</sup> Let  $(X, T)$  be a TOPOLOGICAL SPACE (Definition A.1 page 26) and  $B \subseteq 2^X$ .

$$B \text{ is a base for } (X, T) \iff \left\{ \begin{array}{l} 1. x \in X \implies \exists B_x \in B \text{ such that } x \in B_x \text{ and} \\ 2. B_1, B_2 \in B \implies B_1 \cap B_2 \in B \end{array} \right.$$

**Example A.6** <sup>53</sup> Let  $(X, d)$  be a *metric space*. The set  $B \triangleq \{B(x, r) \mid x \in X, r \in \mathbb{N}\}$  (the set of all open balls in  $(X, d)$ ) is a *base* for a topology on  $(X, d)$ .

**Example A.7** (the standard topology on the real line) <sup>54</sup> The set  $B \triangleq \{(a, b) \mid a, b \in \mathbb{R}, a < b\}$  is a *base* for the metric space  $(\mathbb{R}, |b - a|)$  (the *usual metric space* on  $\mathbb{R}$ ).

**Definition A.8** <sup>55</sup> Let  $(X, T)$  be a *topological space* (Definition A.1 page 26). Let  $2^X$  be the *power set* of  $X$ .

The set  $A^-$  is the **closure** of  $A \in 2^X$  if  $A^- \triangleq \bigcap \{D \in 2^X \mid A \subseteq D \text{ and } D \text{ is closed}\}$ .

The set  $A^\circ$  is the **interior** of  $A \in 2^X$  if  $A^\circ \triangleq \bigcup \{U \in 2^X \mid U \subseteq A \text{ and } U \text{ is open}\}$ .

A point  $x$  is a **closure point** of  $A$  if  $x \in A^-$ .

A point  $x$  is an **interior point** of  $A$  if  $x \in A^\circ$ .

A point  $x$  is an **accumulation point** of  $A$  if  $x \in (A \setminus \{x\})^-$ .

A point  $x$  in  $A^-$  is a **point of adherence** in  $A$  or is **adherent** to  $A$  if  $x \in A^-$ .

**Proposition A.9** <sup>56</sup> Let  $(X, T)$  be a TOPOLOGICAL SPACE (Definition A.1 page 26). Let  $A^-$  be the CLOSURE,  $A^\circ$  the INTERIOR, and  $\partial A$  the BOUNDARY of a set  $A$ . Let  $2^X$  be the POWER SET of  $X$ .

1.  $A^-$  is CLOSED  $\forall A \in 2^X$ .
2.  $A^\circ$  is OPEN  $\forall A \in 2^X$ .

**Lemma A.10** <sup>57</sup> Let  $A^-$  be the CLOSURE,  $A^\circ$  the INTERIOR, and  $\partial A$  the BOUNDARY of a set  $A$  in a topological space  $(X, T)$ . Let  $2^X$  be the POWER SET of  $X$ .

1.  $A^\circ \subseteq A \subseteq A^-$   $\forall A \in 2^X$ .
2.  $A = A^\circ \iff A \text{ is OPEN}$   $\forall A \in 2^X$ .
3.  $A = A^- \iff A \text{ is CLOSED}$   $\forall A \in 2^X$ .

<sup>50</sup> [Joshi(1983)] page 92 (3.1 Definition), [Davis(2005)] page 46 (Definition 4.15)

<sup>51</sup> [Joshi(1983)] pages 92–93 (3.2 Proposition), [Davis(2005)] page 46

<sup>52</sup> [Bollobás(1999)] page 19

<sup>53</sup> [Davis(2005)] page 46 (Example 4.16)

<sup>54</sup> [Munkres(2000)] page 81, [Davis(2005)] page 46 (Example 4.16)

<sup>55</sup> [Gemignani(1972)] pages 55–56 (Definition 3.5.7), [McCarty(1967)] page 90, [Munkres(2000)] page 95 (Closure and Interior of a Set), [Thron(1966)], pages 21–22 (definition 4.8, definition 4.9), [Kelley(1955)] page 42, [Kubrusly(2001)] pages 115–116

<sup>56</sup> [McCarty(1967)] page 90 (IV.1 THEOREM)

<sup>57</sup> [McCarty(1967)] pages 90–91 (IV.1 THEOREM), [ALIPRANTIS AND BURKINSHAW(1998)] PAGE 59



**Definition A.11** <sup>58</sup> Let  $(X, T_x)$  and  $(Y, T_y)$  be *topological spaces* (Definition A.1 page 26). Let  $f$  be a function in  $Y^X$ . A function  $f \in Y^X$  is **continuous** if

$$\underbrace{U \in T_y}_{\text{open in } (Y, T_y)} \implies \underbrace{f^{-1}(U) \in T_x}_{\text{open in } (X, T_x)}.$$

A function is **discontinuous** in  $(X, T_y)^{(X, T_x)}$  if it is not *continuous* in  $(X, T_y)^{(X, T_x)}$ .




Figure 3: *continuous/discontinuous* functions (Example A.12 page 28)

**Example A.12** Some *continuous/discontinuous* functions are illustrated in Figure 3 (page 28).

Definition A.11 (previous definition) defines continuity using open sets. Continuity can alternatively be defined using closed sets or closure (next theorem).

**Theorem A.13** <sup>59</sup> Let  $(X, T)$  and  $(Y, S)$  be *topological spaces*. Let  $f$  be a function in  $Y^X$ .

The following are equivalent:

1.  $f$  is CONTINUOUS  $\iff$
2.  $B$  is closed in  $(Y, S) \implies f^{-1}(B)$  is closed in  $(X, T) \quad \forall B \in 2^Y$   $\iff$
3.  $f(A^-) \subseteq f(A)^-$   $\forall A \in 2^X \iff$
4.  $f^{-1}(B)^- \subseteq f^{-1}(B^-)$   $\forall B \in 2^Y$

**Remark A.14** A word of warning about defining *continuity* in terms of topological spaces—*continuity* is defined in terms of a pair of *topological spaces*, and whether function is *continuous* or *discontinuous* in general depends very heavily on the selection of these spaces. This is illustrated in Proposition A.15 (next). The ramification of this is that when declaring a function to be *continuous* or *discontinuous*, one must make clear the assumed *topological spaces*.

**Proposition A.15** <sup>60</sup> Let  $(X, T)$  and  $(Y, S)$  be TOPOLOGICAL SPACES. Let  $f$  be a FUNCTION in  $(Y, S)^{(X, T)}$ .

1.  $T$  is the DISCRETE TOPOLOGY  $\implies f$  is CONTINUOUS  $\forall f \in (Y, S)^{(X, T)}$
2.  $S$  is the INDISCRETE TOPOLOGY  $\implies f$  is CONTINUOUS  $\forall f \in (Y, S)^{(X, T)}$

**Definition A.16** <sup>61</sup> Let  $(X, T)$  be a *topological space* (Definition A.1 page 26). A sequence  $(x_n)_{n \in \mathbb{Z}}$  **converges** in  $(X, T)$  to a point  $x$  if for each *open set* (Definition A.1 page 26)  $U \in T$  that contains  $x$  there exists  $N \in \mathbb{N}$  such that

<sup>58</sup> [Davis(2005)] page 34

<sup>59</sup> [McCarty(1967)] pages 91–92 (IV.2 THEOREM), [SEARCÓID(2006)] PAGE 130 (“THEOREM 8.3.1 (CRITERIA FOR CONTINUITY)”, SET IN *metric spaces*)

<sup>60</sup> [Crossley(2006)] page 18 (Proposition 3.9), [Ponnusamy(2002)] page 98 (2.64. Theorem.)

<sup>61</sup> [Joshi(1983)] page 83 ((3.1) Definition), [Leathem(1905)], page 13 (“ $\rightarrow$ ” symbol, section III.11)



$x_n \in U$  for all  $n > N$ .

This condition can be expressed in any of the following forms:

1. The **limit** of the sequence  $(x_n)$  is  $x$ .
2. The sequence  $(x_n)$  is **convergent** with limit  $x$ .
3.  $\lim_{n \rightarrow \infty} (x_n) = x$ .
4.  $(x_n) \rightarrow x$ .

A sequence that converges is **convergent**. A sequence that does not converge is said to **diverge**, or is **divergent**. An element  $x \in A$  is a **limit point** of  $A$  if it is the limit of some  $A$ -valued sequence  $(x_n \in A)$ .

**Example A.17** <sup>62</sup> Let  $(X, T_{31})$  be a *topological space* where  $X \triangleq \{x, y, z\}$  and

$$T_{31} \triangleq \{\emptyset, \{x\}, \{x, y\}, \{x, z\}, \{x, y, z\}\}.$$

In this space, the sequence  $(x, x, x, \dots)$  converges to  $x$ . But this sequence also converges to both  $y$  and  $z$  because  $x$  is in every *open set* (Definition A.1 page 26) that contains  $y$  and  $x$  is in every *open set* that contains  $z$ . So, the *limit* (Definition A.16 page 28) of the sequence is *not unique*.

**Example A.18** In contrast to the low resolution topological space of Example A.17, the limit of the sequence  $(x, x, x, \dots)$  is unique in a *topological space* with sufficiently high resolution with respect to  $y$  and  $z$  such as the following: Define a *topological space*  $(X, T_{56})$  where  $X \triangleq \{x, y, z\}$  and

$$T_{56} \triangleq \{\emptyset, \{y\}, \{z\}, \{x, y\}, \{y, z\}, \{x, y, z\}\}.$$

In this space, the sequence  $(x, x, x, \dots)$  converges to  $x$  only. The sequence does *not* converge to  $y$  or  $z$  because there are *open sets* (Definition A.1 page 26) containing  $y$  or  $z$  that do not contain  $x$  (the open sets  $\{y\}$ ,  $\{z\}$ , and  $\{y, z\}$ ).

**Theorem A.19** (The Closed Set Theorem) <sup>63</sup> Let  $(X, T)$  be a TOPOLOGICAL SPACE. Let  $A$  be a subset of  $X$  ( $A \subseteq X$ ). Let  $A^-$  be the CLOSURE (Definition A.8 page 27) of  $A$  in  $(X, T)$ .

$$\underbrace{A \text{ is CLOSED in } (X, T)}_{(A = A^-)} \iff \left\{ \begin{array}{l} \text{Every } A\text{-valued sequence } (x_n \in A)_{n \in \mathbb{Z}} \\ \text{that CONVERGES in } (X, T) \text{ has its LIMIT in } A \end{array} \right\}$$

**Theorem A.20** <sup>64</sup> Let  $(X, T)$  and  $(Y, S)$  be a TOPOLOGICAL SPACES. Let  $f$  be a function in  $(Y, S)^{(X, T)}$ .

$$\underbrace{\left\{ \begin{array}{l} f \text{ is CONTINUOUS in } (Y, S)^{(X, T)} \\ \text{(Definition A.11 page 28)} \end{array} \right\}}_{\text{INVERSE IMAGE CHARACTERIZATION OF CONTINUITY}} \iff \underbrace{\left\{ \begin{array}{l} (x_n) \rightarrow x \implies f((x_n)) \rightarrow f(x) \\ \text{(Definition A.16 page 28)} \end{array} \right\}}_{\text{SEQUENTIAL CHARACTERIZATION OF CONTINUITY}}$$

 PROOF:


(1) Proof for the  $\implies$  case (proof by contradiction):

- (a) Let  $U$  be an *open set* in  $(Y, T)$  that contains  $f(x)$  but for which there exists no  $N$  such that  $f(x_n) \in U$  for all  $n > N$ .
- (b) Note that the set  $f^{-1}(U)$  is also *open* by the *continuity* hypothesis.

<sup>62</sup>  [Munkres(2000)] page 98 (Hausdorff Spaces)

<sup>63</sup>  [Kubrusly(2001)] page 118 (Theorem 3.30),  [Haaser and Sullivan(1991)] page 75 (6.9 Proposition),

 [Rosenlicht(1968)] pages 47–48

<sup>64</sup>  [Ponnusamy(2002)] pages 94–96 (“2.59. Proposition.”; in the context of *metric spaces*; includes the “*inverse image characterization of continuity*” and “*sequential characterization of continuity*” terminology; this terminology does not seem to be widely used in the literature in general, but has been adopted for use in this text)



(c) If  $(x_n) \rightarrow x$ , then

$f((x_n)) \not\rightarrow f(x) \implies$  there exists no  $N$  such that  $f(x_n) \in U$  for all  $n > N$  by Definition A.16 (page 28)  
 $\implies$  there exists no  $M$  such that  $x_n \in f^{-1}(U)$  for all  $n > M$  by definition of  $f^{-1}$   
 $\implies (x_n) \not\rightarrow x$  by *continuity* hypothesis and def. of *convergence* (Definition A.16 page 28)  
 $\implies$  contradiction of  $(x_n) \rightarrow x$  hypothesis  
 $\implies f((x_n)) \rightarrow f(x)$

(2) Proof for the  $\Leftarrow$  case (proof by contradiction):

(a) Let  $D$  be a *closed* set in  $(Y, S)$ .

(b) Suppose  $f^{-1}(D)$  is *not closed*...

(c) then by the *closed set theorem* (Theorem A.19 page 29), there must exist a *convergent* sequence  $(x_n)$  in  $(X, T)$ , but with limit  $x$  *not* in  $f^{-1}(D)$ .

(d) Note that  $f(x)$  must be in  $D$ . Proof:

(i) by definition of  $D$  and  $f$ ,  $f((x_n))$  is in  $D$

(ii) by left hypothesis, the sequence  $f((x_n))$  is *convergent* with limit  $f(x)$

(iii) by *closed set theorem* (Theorem A.19 page 29),  $f(x)$  must be in  $D$ .

(e) Because  $f(x) \in D$ , it must be true that  $x \in f^{-1}(D)$ .

(f) But this is a contradiction to item (2c) (page 30), and so item (2b) (page 30) must be wrong, and  $f^{-1}(D)$  must be *closed*.

(g) And so by Theorem A.13 (page 28),  $f$  is *continuous*.

$\Rightarrow$

## Appendix B Finite sums

### B.1 Convexity

#### Definition B.1 <sup>65</sup>

A function  $f \in \mathbb{R}^{\mathbb{R}}$  is **convex** if

$$f(\lambda x + [1 - \lambda]y) \leq \lambda f(x) + (1 - \lambda) f(y) \quad \forall x, y \in \mathbb{R} \text{ and } \forall \lambda \in (0 : 1)$$

A function  $g \in \mathbb{R}^{\mathbb{R}}$  is **strictly convex** if

$$g(\lambda x + [1 - \lambda]y) < \lambda g(x) + (1 - \lambda) g(y) \quad \forall x, y \in D, x \neq y, \text{ and } \forall \lambda \in (0 : 1)$$

A function  $f \in \mathbb{R}^{\mathbb{R}}$  is **concave** if  $-f$  is *convex*.

A function  $f \in \mathbb{R}^{\mathbb{R}}$  is **affine** if  $f$  is *convex* and *concave*.

**Theorem B.2** (Jensen's Inequality) <sup>66</sup> Let  $f \in \mathbb{R}^{\mathbb{R}}$  be a function.

$$\left\{ \begin{array}{l} 1. f \text{ is CONVEX (Definition B.1 page 30) and} \\ 2. \sum_{n=1}^N \lambda_n = 1 \quad \text{(WEIGHTS)} \end{array} \right\} \implies \left\{ f\left(\sum_{n=1}^N \lambda_n x_n\right) \leq \sum_{n=1}^N \lambda_n f(x_n) \quad \forall x_n \in D, N \in \mathbb{N} \right\}$$

<sup>65</sup>  [Simon(2011)] page 2,  [Barvinok(2002)] page 2,  [Bollobás(1999)], page 3,  [Jensen(1906)], page 176

<sup>66</sup>  [Mitrinović et al.(2010)Mitrinović, Pečarić, and Fink] page 6,  [Bollobás(1999)] page 3,  [Jensen(1906)] pages 179–180



## B.2 Power means

### Definition B.3 <sup>67</sup>

The  $(\lambda_n)_1^N$  weighted  $\phi$ -mean of a tuple  $(x_n)_1^N$  is defined as

$$M_\phi((x_n)) \triangleq \phi^{-1}\left(\sum_{n=1}^N \lambda_n \phi(x_n)\right)$$

where  $\phi$  is a *continuous* and *strictly monotonic* function in  $\mathbb{R}^{\mathbb{R}^+}$

and  $(\lambda_n)_{n=1}^N$  is a sequence of weights for which  $\sum_{n=1}^N \lambda_n = 1$ .

**Lemma B.4** <sup>68</sup> Let  $M_\phi((x_n))$  be the  $(\lambda_n)_1^N$  weighted  $\phi$ -mean and  $M_\psi((x_n))$  the  $(\lambda_n)_1^N$  weighted  $\psi$ -mean of a tuple  $(x_n)_1^N$ .

$$\begin{aligned} \phi\psi^{-1} \text{ is CONVEX and } \phi \text{ is INCREASING} &\implies M_\phi((x_n)) \geq M_\psi((x_n)) \\ \phi\psi^{-1} \text{ is CONVEX and } \phi \text{ is DECREASING} &\implies M_\phi((x_n)) \leq M_\psi((x_n)) \\ \phi\psi^{-1} \text{ is CONCAVE and } \phi \text{ is INCREASING} &\implies M_\phi((x_n)) \leq M_\psi((x_n)) \\ \phi\psi^{-1} \text{ is CONCAVE and } \phi \text{ is DECREASING} &\implies M_\phi((x_n)) \geq M_\psi((x_n)) \end{aligned}$$

One of the most well known inequalities in mathematics is *Minkowski's Inequality*. In 1946, H.P. Mulholland submitted a result that generalizes Minkowski's Inequality to an equal weighted  $\phi$ -mean.<sup>69</sup> And Milovanović and Milovanović (1979) generalized this even further to a *weighted*  $\phi$ -mean (next).

**Theorem B.5** <sup>70</sup> Let  $\phi$  be a function in  $\mathbb{R}^{\mathbb{R}}$ .

$$\left\{ \begin{array}{l} 1. \phi \text{ is CONVEX and } 2. \phi \text{ is STRICTLY MONOTONE and } \\ 3. \phi(0) = 0 \quad \text{and } 4. \log \circ \phi \circ \exp \text{ is CONVEX} \end{array} \right\} \implies \left\{ \phi^{-1}\left(\sum_{n=1}^N \lambda_n \phi(x_n + y_n)\right) \leq \phi^{-1}\left(\sum_{n=1}^N \lambda_n \phi(x_n)\right) + \phi^{-1}\left(\sum_{n=1}^N \lambda_n \phi(y_n)\right) \right\}$$

**Definition B.6** <sup>71</sup> Let  $M_{\phi(x;p)}((x_n))$  be the  $(\lambda_n)_1^N$  weighted  $\phi$ -mean of a *non-negative* tuple  $(x_n)_1^N$ . A mean  $M_{\phi(x;p)}((x_n))$  is a **power mean** with parameter  $p$  if  $\phi(x) \triangleq x^p$ . That is,

$$M_{\phi(x;p)}((x_n)) = \left(\sum_{n=1}^N \lambda_n (x_n)^p\right)^{\frac{1}{p}}$$

**Theorem B.7** <sup>72</sup> Let  $M_{\phi(x;p)}((x_n))$  be the POWER MEAN with parameter  $p$  of an  $N$ -tuple  $(x_n)_1^N$  in which the elements are NOT all equal.

<sup>67</sup> [Bollobás(1999)] page 5

<sup>68</sup> [Pečarić et al.(1992)Pečarić, Proschan, and Tong] page 107, [Bollobás(1999)] page 5, [Hardy et al.(1952)Hardy, Littlewood, and Pólya] page 75

<sup>69</sup> [Minkowski(1910)] page 115, [Mulholland(1950)], [Hardy et al.(1952)Hardy, Littlewood, and Pólya] <Theorem 24>, [Tolsted(1964)] page 7, [Maligranda(1995)] page 258, [Carothers(2000)], page 44, [Bullen(2003)] page 179

<sup>70</sup> [Milovanović and Milovanović(1979)], [Bullen(2003)] page 306 <Theorem 9>

<sup>71</sup> [Bullen(2003)] page 175, [Bollobás(1999)] page 6

<sup>72</sup> [Bullen(2003)] pages 175–177 <see also page 203>, [Bollobás(1999)] pages 6–8, [Bullen(1990)] page 250, [Besso(1879)], [Bienaymé(1840)] page 68, [Brenner(1985)] page 160



$$M_{\phi(x;p)}(\langle x_n \rangle) \triangleq \left( \sum_{n=1}^N \lambda_n(x_n)^p \right)^{\frac{1}{p}} \text{ is CONTINUOUS and STRICTLY MONOTONE in } \mathbb{R}^*.$$

$$M_{\phi(x;p)}(\langle x_n \rangle) = \begin{cases} \max_{n=1,2,\dots,N} \langle x_n \rangle & \text{for } p = +\infty \\ \prod_{n=1}^N x_n^{\lambda_n} & \text{for } p = 0 \\ \min_{n=1,2,\dots,N} \langle x_n \rangle & \text{for } p = -\infty \end{cases}$$

 PROOF:

(1) Proof that  $M_{\phi(x;p)}$  is strictly monotone in  $p$ :

(a) Let  $p$  and  $s$  be such that  $-\infty < p < s < \infty$ .

(b) Let  $\phi_p \triangleq x^p$  and  $\phi_s \triangleq x^s$ . Then  $\phi_p \phi_s^{-1} = x^{\frac{p}{s}}$ .

(c) The composite function  $\phi_p \phi_s^{-1}$  is convex or concave depending on the values of  $p$  and  $s$ :

	$p < 0$ ( $\phi_p$ decreasing)	$p > 0$ ( $\phi_p$ increasing)
$s < 0$	convex	(not possible)
$s > 0$	convex	concave

(d) Therefore by Lemma B.4 (page 31),

$$-\infty < p < s < \infty \implies M_{\phi(x;p)}(\langle x_n \rangle) < M_{\phi(x;s)}(\langle x_n \rangle).$$

(2) Proof that  $M_{\phi(x;p)}$  is continuous in  $p$  for  $p \in \mathbb{R} \setminus \{0\}$ : The sum of continuous functions is continuous. For the cases of  $p \in \{-\infty, 0, \infty\}$ , see the items that follow.

(3) Lemma:  $M_{\phi(x;-p)}(\langle x_n \rangle) = \{M_{\phi(x;p)}(\langle x_n^{-1} \rangle)\}^{-1}$ . Proof:

$$\begin{aligned} \{M_{\phi(x;p)}(\langle x_n^{-1} \rangle)\}^{-1} &= \left\{ \left( \sum_{n=1}^N \lambda_n(x_n^{-1})^p \right)^{\frac{1}{p}} \right\}^{-1} && \text{by definition of } M_\phi \\ &= \left( \sum_{n=1}^N \lambda_n(x_n)^{-p} \right)^{\frac{1}{-p}} \\ &= M_{\phi(x;-p)}(\langle x_n \rangle) && \text{by definition of } M_\phi \end{aligned}$$

(4) Proof that  $\lim_{p \rightarrow \infty} M_\phi(\langle x_n \rangle) = \max_{n \in \mathbb{Z}} \langle x_n \rangle$ :

(a) Let  $x_m \triangleq \max_{n \in \mathbb{Z}} \langle x_n \rangle$

(b) Note that  $\lim_{p \rightarrow \infty} M_\phi \leq \max_{n \in \mathbb{Z}} \langle x_n \rangle$  because

$$\begin{aligned} \lim_{p \rightarrow \infty} M_\phi(\langle x_n \rangle) &= \lim_{p \rightarrow \infty} \left( \sum_{n=1}^N \lambda_n x_n^p \right)^{\frac{1}{p}} && \text{by definition of } M_\phi \\ &\leq \lim_{p \rightarrow \infty} \left( \sum_{n=1}^N \lambda_n x_m^p \right)^{\frac{1}{p}} && \text{by definition of } x_m \text{ in item (4a) and because } \phi(x) \triangleq x^p \text{ and } \phi^{-1} \text{ are both increasing or both decreasing} \\ &= \lim_{p \rightarrow \infty} \left( x_m^p \underbrace{\sum_{n=1}^N \lambda_n}_1 \right)^{\frac{1}{p}} && \text{because } x_m \text{ is a constant} \end{aligned}$$





$$\begin{aligned}
&= \lim_{p \rightarrow \infty} (x_m^p \cdot 1)^{\frac{1}{p}} \\
&= x_m \\
&= \max_{n \in \mathbb{Z}} \langle x_n \rangle && \text{by definition of } x_m \text{ in item (4a)}
\end{aligned}$$

(c) But also note that  $\lim_{p \rightarrow \infty} M_\phi \geq \max_{n \in \mathbb{Z}} \langle x_n \rangle$  because

$$\begin{aligned}
\lim_{p \rightarrow \infty} M_\phi(\langle x_n \rangle) &= \lim_{p \rightarrow \infty} \left( \sum_{n=1}^N \lambda_n x_n^p \right)^{\frac{1}{p}} && \text{by definition of } M_\phi \\
&\geq \lim_{p \rightarrow \infty} (w_m x_m^p)^{\frac{1}{p}} && \text{by definition of } x_m \text{ in item (4a) and because } \phi(x) \triangleq x^p \text{ and } \phi^{-1} \text{ are both increasing or both decreasing} \\
&= \lim_{p \rightarrow \infty} w_m^{\frac{1}{p}} x_m^{\frac{p}{p}} \\
&= x_m \\
&= \max_{n \in \mathbb{Z}} \langle x_n \rangle && \text{by definition of } x_m \text{ in item (4a)}
\end{aligned}$$

(d) Combining items (b) and (c) we have  $\lim_{p \rightarrow \infty} M_\phi = \max_{n \in \mathbb{Z}} \langle x_n \rangle$ .

(5) Proof that  $\lim_{p \rightarrow -\infty} M_\phi(\langle x_n \rangle) = \min_{n \in \mathbb{Z}} \langle x_n \rangle$ :

$$\begin{aligned}
\lim_{p \rightarrow -\infty} M_{\phi(x;p)}(\langle x_n \rangle) &= \lim_{p \rightarrow \infty} M_{\phi(x;-p)}(\langle x_n \rangle) && \text{by change of variable } p \\
&= \lim_{p \rightarrow \infty} \{M_{\phi(x;p)}(\langle x_n^{-1} \rangle)\}^{-1} && \text{by Lemma in item (3) page 32} \\
&= \lim_{p \rightarrow \infty} \frac{1}{M_{\phi(x;p)}(\langle x_n^{-1} \rangle)} \\
&= \frac{\lim_{p \rightarrow \infty} 1}{\lim_{p \rightarrow \infty} M_{\phi(x;p)}(\langle x_n^{-1} \rangle)} && \text{by property of } \lim^{73} \\
&= \frac{1}{\max_{n \in \mathbb{Z}} \langle x_n^{-1} \rangle} && \text{by item (4)} \\
&= \frac{1}{\left(\min_{n \in \mathbb{Z}} \langle x_n \rangle\right)^{-1}} \\
&= \min_{n \in \mathbb{Z}} \langle x_n \rangle
\end{aligned}$$

(6) Proof that  $\lim_{p \rightarrow 0} M_\phi(\langle x_n \rangle) = \prod_{n=1}^N x_n^{\lambda_n}$ :

$$\begin{aligned}
\lim_{p \rightarrow 0} M_\phi(\langle x_n \rangle) &= \lim_{p \rightarrow 0} \exp \left\{ \ln \left\{ M_\phi(\langle x_n \rangle) \right\} \right\} \\
&= \lim_{p \rightarrow 0} \exp \left\{ \ln \left\{ \left( \sum_{n=1}^N \lambda_n (x_n^p) \right)^{\frac{1}{p}} \right\} \right\} && \text{by definition of } M_\phi
\end{aligned}$$

<sup>73</sup>  [Rudin(1976)] page 85 (4.4 Theorem)



$$\begin{aligned}
&= \exp \left\{ \frac{\frac{\partial}{\partial p} \ln \left( \sum_{n=1}^N \lambda_n (x_n^p) \right)}{\frac{\partial}{\partial p} p} \right\}_{p=0} && \text{by l'Hôpital's rule}^{74} \\
&= \exp \left\{ \frac{\sum_{n=1}^N \lambda_n \frac{\partial}{\partial p} (x_n^p)}{\sum_{n=1}^N \lambda_n (x_n^p)} \right\}_{p=0} && \\
&= \exp \left\{ \frac{\sum_{n=1}^N \lambda_n \frac{\partial}{\partial p} \exp (r \ln (x_n))}{1} \right\}_{p=0} && \\
&= \exp \left\{ \sum_{n=1}^N \lambda_n \exp \{ p \ln x_n \} \ln (x_n) \right\}_{p=0} && \\
&= \exp \left\{ \sum_{n=1}^N \lambda_n \ln (x_n) \right\} && \\
&= \exp \left\{ \ln \prod_{n=1}^N x_n^{\lambda_n} \right\} = \prod_{n=1}^N x_n^{\lambda_n} &&
\end{aligned}$$

⇒

**Corollary B.8** <sup>75</sup> Let  $(x_n)_1^N$  be a tuple. Let  $(\lambda_n)_1^N$  be a tuple of weighting values such that  $\sum_{n=1}^N \lambda_n = 1$ .

$$\min (x_n) \leq \underbrace{\left( \sum_{n=1}^N \lambda_n \frac{1}{x_n} \right)^{-1}}_{\text{harmonic mean}} \leq \underbrace{\prod_{n=1}^N x_n^{\lambda_n}}_{\text{geometric mean}} \leq \underbrace{\sum_{n=1}^N \lambda_n x_n}_{\text{arithmetic mean}} \leq \max (x_n)$$

PROOF:

(1) These five means are all special cases of the *power mean*  $M_{\phi(x;p)}$  (Definition B.6 page 31):

$$\begin{aligned}
p = \infty: & \quad \max (x_n) \\
p = 1: & \quad \text{arithmetic mean} \\
p = 0: & \quad \text{geometric mean} \\
p = -1: & \quad \text{harmonic mean} \\
p = -\infty: & \quad \min (x_n)
\end{aligned}$$

(2) The inequalities follow directly from Theorem B.7 (page 31).

<sup>74</sup> [Rudin(1976)] page 109 (5.13 Theorem)

<sup>75</sup> [Bullen(2003)] page 71, [Bollobás(1999)] page 5, [Cauchy(1821)] pages 457–459 (Note II, theorem 17), [Jensen(1906)] page 183, [Hoehn and Niven(1985)] page 151



- (3) Generalized AM-GM inequality: If one is only concerned with the arithmetic mean and geometric mean, their relationship can be established directly using *Jensen's Inequality*:

$$\begin{aligned} \sum_{n=1}^N \lambda_n x_n &= b^{\log_b(\sum_{n=1}^N \lambda_n x_n)} \\ &\geq b^{(\sum_{n=1}^N \lambda_n \log_b x_n)} && \text{by Jensen's Inequality (Theorem B.2 page 30)} \\ &= \prod_{n=1}^N b^{(\lambda_n \log_b x_n)} = \prod_{n=1}^N b^{(\log_b x_n) \lambda_n} = \prod_{n=1}^N x_n^{\lambda_n} \end{aligned}$$

◻

### B.3 Inequalities

**Lemma B.9** (Young's Inequality) <sup>76</sup>

$$\begin{aligned} xy &< \frac{x^p}{p} + \frac{y^q}{q} \quad \text{with } \frac{1}{p} + \frac{1}{q} = 1 \quad \forall 1 < p < \infty, \quad x, y \geq 0, \quad \text{but } y \neq x^{p-1} \\ xy &= \frac{x^p}{p} + \frac{y^q}{q} \quad \text{with } \frac{1}{p} + \frac{1}{q} = 1 \quad \forall 1 < p < \infty, \quad x, y \geq 0, \quad \text{and } y = x^{p-1} \end{aligned}$$

**Theorem B.10** (Minkowski's Inequality for sequences) <sup>77</sup> Let  $(x_n \in \mathbb{C})_1^N$  and  $(y_n \in \mathbb{C})_1^N$  be complex  $N$ -tuples.

$$\left( \sum_{n=1}^N |x_n + y_n|^p \right)^{\frac{1}{p}} \leq \left( \sum_{n=1}^N |x_n|^p \right)^{\frac{1}{p}} + \left( \sum_{n=1}^N |y_n|^p \right)^{\frac{1}{p}} \quad \forall 1 < p < \infty$$

## Appendix C Metric preserving functions

**Definition C.1** <sup>78</sup> Let  $\mathbb{M}$  be the set of all *metric spaces* (Definition 4.5 page 16) on a set  $X$ .  $\phi \in \mathbb{R}^{+\mathbb{R}^+}$  is a **metric preserving function** if  $d(x, y) \triangleq \phi \circ \rho(x, y)$  is a *metric* on  $X$  for all  $(X, \rho) \in \mathbb{M}$

**Theorem C.2** (necessary conditions) <sup>79</sup> Let  $\mathcal{R}\phi$  be the RANGE of a function  $\phi$ .

$$\left\{ \begin{array}{l} \phi \text{ is a} \\ \text{METRIC PRESERVING FUNCTION} \\ \text{(Definition C.1 page 35)} \end{array} \right\} \implies \left\{ \begin{array}{ll} 1. \phi^{-1}(0) = \{0\} & \text{and} \\ 2. \mathcal{R}\phi \subseteq \mathbb{R}^+ & \text{and} \\ 3. \phi(x + y) \leq \phi(x) + \phi(y) & (\phi \text{ is SUBADDITIVE}) \end{array} \right\}$$

<sup>76</sup> [Young(1912)] page 226, [Hardy et al.(1952)Hardy, Littlewood, and Pólya] (Theorem 24), [Tolsted(1964)] page 5, [Maligranda(1995)] page 257, [Carothers(2000)], page 43

<sup>77</sup> [Minkowski(1910)], page 115, [Hardy et al.(1952)Hardy, Littlewood, and Pólya] (Theorem 24), [Maligranda(1995)] page 258, [Tolsted(1964)], page 7, [Carothers(2000)], page 44, [Bullen(2003)] page 179

<sup>78</sup> [Vallin(1999)], page 849 (Definition 1.1), [Corazza(1999)], page 309, [Deza and Deza(2009)] page 80

<sup>79</sup> [Corazza(1999)], page 310 (Proposition 2.1), [Deza and Deza(2009)] page 80



**Theorem C.3** (sufficient conditions) <sup>80</sup> Let  $\phi$  be a function in  $\mathbb{R}^{\mathbb{R}}$ .

$$\left\{ \begin{array}{l} 1. \ x \geq y \implies \phi(x) \geq \phi(y) \quad \forall x, y \in \mathbb{R}^+ \quad (\text{ISOTONE}) \\ 2. \ \phi(0) = 0 \\ 3. \ \phi(x + y) \leq \phi(x) + \phi(y) \quad \forall x, y \in \mathbb{R}^+ \quad (\text{SUBADDITIVE}) \end{array} \right\} \text{ and } \left\{ \begin{array}{l} \phi \text{ is a METRIC} \\ \text{PRESERVING FUNCTION} \\ (\text{Definition C.1 page 35}). \end{array} \right\}$$




Figure 4: metric preserving functions

**Example C.4** ( $\alpha$ -scaled metric/dilated metric) <sup>81</sup> Let  $(X, d)$  be a *metric space* (Definition 4.5 page 16).  $\phi(x) \triangleq \alpha x, \alpha \in \mathbb{R}^+$  is a *metric preserving function* (Figure 4 page 36 (A))

**PROOF:** The proofs for Example C.4–Example C.9 (page 36) follow from Theorem C.3 (page 36). ⇒

**Example C.5** (power transform metric/snowflake transform metric) <sup>82</sup> Let  $(X, d)$  be a *metric space* (Definition 4.5 page 16).  $\phi(x) \triangleq x^\alpha, \alpha \in (0 : 1]$ , is a *metric preserving function* (see Figure 4 page 36 (B))

**Example C.6** ( $\alpha$ -truncated metric/radar screen metric) <sup>83</sup> Let  $(X, d)$  be a *metric space* (Definition 4.5 page 16).  $\phi(x) \triangleq \min \{ \alpha, x \}, \alpha \in \mathbb{R}^+$  is a *metric preserving function* (see Figure 4 page 36 (C)).

**Example C.7** (bounded metric) <sup>84</sup> Let  $(X, d)$  be a *metric space* (Definition 4.5 page 16).  $\phi(x) \triangleq \frac{x}{1+x}$  is a *metric preserving function* (see Figure 4 page 36 (D)).

**Example C.8** (discrete metric preserving function) <sup>85</sup> Let  $\phi$  be a function in  $\mathbb{R}^{\mathbb{R}}$ .  $\phi(x) \triangleq \begin{cases} 0 & \text{for } x \leq 0 \\ 1 & \text{otherwise} \end{cases}$  is a *metric preserving function* (see Figure 4 page 36 (E)).

**Example C.9** Let  $\phi$  be a function in  $\mathbb{R}^{\mathbb{R}}$ .  $\phi(x) \triangleq \begin{cases} x & \text{for } 0 \leq x < 1, & 1 & \text{for } 1 \leq x \leq 2, \\ x - 1 & \text{for } 2 < x < 3, & 2 & \text{for } x \geq 3 \end{cases}$  is a *metric preserving function* (see Figure 4 page 36 (F)).

<sup>80</sup> [Corazza(1999)] (Proposition 2.3), [Deza and Deza(2009)] page 80, [Kelley(1955)] page 131 (Problem C)

<sup>81</sup> [Deza and Deza(2006)] page 44

<sup>82</sup> [Deza and Deza(2009)] page 81, [Deza and Deza(2006)] page 45

<sup>83</sup> [Giles(1987)], page 33, [Deza and Deza(2006)] pages 242–243

<sup>84</sup> [Vallin(1999)], page 849, [Aliprantis and Burkinshaw(1998)] page 39

<sup>85</sup> [Corazza(1999)], page 311



## References

- [oei(2014)] , 2014. On-line encyclopedia of integer sequences. World Wide Web.  
URL <http://oeis.org/>
- [Adams and Franzosa(2008)] Adams, C. C., Franzosa, R. D., 2008. Introduction to Topology: Pure and Applied. Featured Titles for Topology Series. Pearson Prentice Hall.  
URL <http://books.google.com/books?vid=ISBN0131848690>
- [Aliprantis and Burkinshaw(1998)] Aliprantis, C. D., Burkinshaw, O., 1998. Principles of Real Analysis, 3rd Edition. Academic Press, London.  
URL <http://www.amazon.com/dp/0120502577>
- [Apostol(1975)] Apostol, T. M., 1975. Mathematical Analysis, 2nd Edition. Addison-Wesley series in mathematics. Addison-Wesley, Reading.  
URL <http://books.google.com/books?vid=ISBN0201002884>
- [Barvinok(2002)] Barvinok, A., 2002. A Course in Convexity. Vol. 54 of Graduate studies in mathematics. American Mathematical Society.  
URL <http://books.google.com/books?vid=ISBN0821872311>
- [Beran(1985)] Beran, L., 1985. Orthomodular Lattices: Algebraic Approach. Mathematics and Its Applications (East European Series). D. Reidel Publishing Company, Dordrecht.  
URL <http://books.google.com/books?vid=ISBN902771715X>
- [Berberian(1961)] Berberian, S. K., 1961. Introduction to Hilbert Space. Oxford University Press, New York.  
URL <http://books.google.com/books?vid=ISBN0821819127>
- [Bessenyei and Pales(2014)] Bessenyei, M., Pales, Z., January 8 2014. A contraction principle in semimetric spaces. arXiv.org.  
URL <http://arxiv.org/abs/1401.1709>
- [Besso(1879)] Besso, D., 1879. Teoremi elementari sui massimi i minimi. Annuari Ist. Tech. Roma, 7–24See Bullen(2003) pages 453, 203.
- [Bienaymé(1840)] Bienaymé, M. J., June 13 1840. Société philomatique de paris—extraits des procès-verbaux. Scéance, 67–68See Bullen(2003) pages 453, 203.  
URL <http://www.archive.org/details/extraitsdesproc46183941soci>
- [Blumenthal(1938)] Blumenthal, L. M., 1938. Distance geometries: a study of the development of abstract metrics. The University of Missouri studies. A quarterly of research 13 (2), 145.
- [Blumenthal(1953)] Blumenthal, L. M., 1953. Theory and Applications of Distance Geometry, 1st Edition. Oxford at the Clarendon Press.  
URL <http://books.google.com/books?vid=ISBN0828402426>
- [Bollobás(1999)] Bollobás, B., March 1 1999. Linear Analysis; an introductory course, 2nd Edition. Cambridge mathematical textbooks. Cambridge University Press, Cambridge.  
URL <http://books.google.com/books?vid=ISBN0521655773>
- [Bottazzini(1986)] Bottazzini, U., 1986. The Higher Calculus: A History of Real and Complex Analysis from Euler to Weierstrass. Springer-Verlag, New York.  
URL <http://books.google.com/books?vid=ISBN0387963022>
- [Bourbaki(1939)] Bourbaki, 1939. Éléments de mathématique. Prewmière partie: Les structures fondamentales de l'analyse. Livre I: Théorie des ensembles. Fascicule des résultats. Paris.
- [Brenner(1985)] Brenner, J. L., Fall 1985. Limits of means for large values of the variable. Pi Mu Epsilon Journal 8 (3), 160–163.  
URL <http://www.pme-math.org/journal/issues.html>



- [Brown and Watson(1996)] Brown, J. I., Watson, S., 15 June 1996. The number of complements of a topology on  $n$  points is at least  $2^n$  (except for some special cases). *Discrete Mathematics* 154 (1–3), 27–39.  
URL [http://dx.doi.org/10.1016/0012-365X\(95\)00004-G](http://dx.doi.org/10.1016/0012-365X(95)00004-G)
- [Bruckner et al.(1997)Bruckner, Bruckner, and Thomson] Bruckner, A. M., Bruckner, J. B., Thomson, B. S., 1997. *Real Analysis*. Prentice-Hall, Upper Saddle River, N.J.  
URL <http://books.google.com/books?vid=ISBN013458886X>
- [Brunschwig et al.(2003)Brunschwig, Lloyd, and Pellegrin] Brunschwig, J., Lloyd, G. E. R., Pellegrin, P., 2003. *A Guide to Greek Thought: Major Figures and Trends*. Harvard University Press.  
URL <http://books.google.com/books?vid=ISBN0674021568>
- [Bryant(1985)] Bryant, V., 1985. *Metric Spaces: Iteration and Application*, illustrated, reprint Edition. Cambridge University Press, Cambridge.  
URL <http://books.google.com/books?vid=ISBN0521318971>
- [Bullen(1990)] Bullen, P. S., 1990. Averages still on the move. *Mathematics Magazine* 63 (4), 250–255.  
URL <http://www.maa.org/sites/default/files/Bullen94830627.pdf>
- [Bullen(2003)] Bullen, P. S., 2003. *Handbook of Means and Their Inequalities*, 2nd Edition. Vol. 560 of *Mathematics and Its Applications*. Kluwer Academic Publishers, Dordrecht, Boston.  
URL <http://books.google.com/books?vid=ISBN1402015224>
- [Burstein et al.(2005)Burstein, Ulitsky, Tuller, and Chor] Burstein, D., Ulitsky, I., Tuller, T., Chor, B., May 14–18 2005. Information theoretic approaches to whole genome phylogenies. In: Miyano, S., Mesirov, J., Kasif, S., Istrail, S., Pevzner, P. A., Waterman, M. (Eds.), *Research in Computational Molecular Biology*. 9th Annual International Conference, RECOMB 2005. Vol. 3500 of *Lecture Notes in Computer Science*. Springer Science & Business Media, pp. 283–295.  
URL [http://www.researchgate.net/profile/Tamir\\_Tuller/publication/221530206\\_Information\\_Theoretic\\_Approaches\\_to\\_Whole\\_Genome\\_Phylogenies/links/0912f50b8c7a689215000000.pdf](http://www.researchgate.net/profile/Tamir_Tuller/publication/221530206_Information_Theoretic_Approaches_to_Whole_Genome_Phylogenies/links/0912f50b8c7a689215000000.pdf)
- [Carothers(2000)] Carothers, N., 2000. *Real Analysis*. Cambridge University Press, Cambridge.  
URL <http://books.google.com/books?vid=ISBN0521497566>
- [Cauchy(1821)] Cauchy, A.-L., 1821. Part 1: *Analyse Algèbre*. *Cours D'Analyse de L'école Royale Polytechnique*. Systems design course of the Polytechnic Royal School; 1st Part: Algebraic analysis).  
URL <http://www.archive.org/details/coursdanalysede00caucgoog>
- [Chatterji(1967)] Chatterji, S. D., July 1967. The Number of Topologies on  $n$  Points. No. N67-31144. Techreport.  
URL [http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19670021815\\_1967021815.pdf](http://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19670021815_1967021815.pdf)
- [Choquet(1954)] Choquet, G., 1954. Theory of capacities. *Annales de l'institut Fourier* 5, 131–295.  
URL [http://aif.cedram.org/item?id=AIF\\_1954\\_\\_5\\_\\_131\\_0](http://aif.cedram.org/item?id=AIF_1954__5__131_0)
- [Cohn(2002 December 6)] Cohn, P., 2002 December 6. *Basic Algebra; Groups, Rings and Fields*. Springer.  
URL <http://books.google.com/books?vid=isbn1852335874>
- [Comtet(1966)] Comtet, L., 1966. Recouvrements, bases de filtre et topologies d'un ensemble fini. *Comptes rendus de l'Academie des sciences* 262 (20), A1091–A1094, recoveries, bases and filter topologies of a finite set.
- [Comtet(1974)] Comtet, L., 1974. *Advanced combinatorics: the art of finite and infinite*. D. Reidel Publishing Company, Dordrecht, translated and corrected version of the 1970 French edition.  
URL <http://books.google.com/books?vid=ISBN9027704414>
- [Copson(1968)] Copson, E. T., 1968. *Metric Spaces*. No. 57 in *Cambridge tracts in mathematics and mathematical physics*. Cambridge University Press, London.  
URL <http://books.google.com/books?vid=ISBN0521047226>
- [Corazza(1999)] Corazza, P., April 1999. Introduction to metric-preserving functions. *The American Mathematical Monthly* 104 (4), 309–323.  
URL <http://pcorazza.lisco.com/papers/metric-preserving.pdf>



- [Costa et al.(2004)Costa, Castro, Rowstron, and Key] Costa, M., Castro, M., Rowstron, A., Key, P., 2004. Pic: Practical internet coordinates for distance estimation. In: Distributed Computing Systems, 2004. Proceedings. 24th International Conference on. pp. 178–187.  
URL <http://research.microsoft.com/pubs/67616/tr-2003-53.pdf>
- [Crammer et al.(2007)Crammer, Kearns, and Wortman] Crammer, K., Kearns, M., Wortman, J., 2007. Learning from multiple sources. In: Scholkopf, B., Platt, J., Hofmann, T. (Eds.), Advances in Neural Information Processing Systems 19: Proceedings of the 2006 Conference. Vol. 19 of Neural information processing series. MIT Press, pp. 321–328.  
URL [http://machinelearning.wustl.edu/mlpapers/paper\\_files/NIPS2006\\_46.pdf](http://machinelearning.wustl.edu/mlpapers/paper_files/NIPS2006_46.pdf)
- [Crossley(2006)] Crossley, M. D., 2006. Essential Topology. Springer Undergraduate Mathematics Series. Springer Science & Business Media.  
URL <http://books.google.com/books?vid=ISBN1846281946>
- [Czerwik(1993)] Czerwik, S., 1993. Contraction mappings in b-metric spaces. Acta Mathematica et Informatica Universitatis Ostraviensis 1 (1), 5–11.  
URL <http://dml.cz/dmlcz/120469>
- [Davis(2005)] Davis, S. W., 2005. Topology. McGraw Hill, Boston.  
URL <http://www.worldcat.org/isbn/0071243399>
- [Dedekind(1900)] Dedekind, R., January 8 1900. Ueber die von drei moduln erzeugte dualgruppe. Mathematische Annalen 53, 371–403, regarding the Dual Group Generated by Three Modules.  
URL <http://resolver.sub.uni-goettingen.de/purl/?GDZPPN002257947>
- [Deza and Deza(2006)] Deza, E., Deza, M.-M., 2006. Dictionary of Distances. Elsevier Science, Amsterdam.  
URL <http://books.google.com/books?vid=ISBN0444520872>
- [Deza and Deza(2009)] Deza, M.-M., Deza, E., 2009. Encyclopedia of Distances. Springer.  
URL <http://www.uco.es/users/malfegan/Comunes/asignaturas/vision/Encyclopedia-of-distances-2009.pdf>
- [Deza and Deza(2014)] Deza, M.-M., Deza, E., 2014. Encyclopedia of Distances, 3rd Edition. Springer, Bücher.  
URL <http://books.google.com/books?vid=ISBN3662443422>
- [DiBenedetto(2002)] DiBenedetto, E., 2002. Real Analysis. Birkhäuser Advanced Texts. Birkhäuser, Boston.  
URL <http://books.google.com/books?vid=ISBN0817642315>
- [Dieudonné(1969)] Dieudonné, J. A., 1969. Foundations of Modern Analysis. Academic Press, New York.  
URL <http://books.google.com/books?vid=ISBN1406727911>
- [Euclid(circa 300BC)] Euclid, circa 300BC. Elements.  
URL <http://farside.ph.utexas.edu/euclid.html>
- [Evans et al.(1967)Evans, Harary, and Lynn] Evans, J., Harary, F., Lynn, M., 1967. On the computer enumeration of finite topologies. Communications of the ACM — Association for Computing Machinery 10, 295–297.  
URL <http://portal.acm.org/citation.cfm?id=363282.363311>
- [Fagin et al.(2003a)Fagin, Kumar, and Sivakumar] Fagin, R., Kumar, R., Sivakumar, D., 2003a. Comparing top  $k$  lists. In: In Proceedings of the ACM-SIAM Symposium on Discrete Algorithms. Society for Industrial and Applied Mathematics, pp. 28–36.  
URL <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.119.6597>
- [Fagin et al.(2003b)Fagin, Kumar, and Sivakumar] Fagin, R., Kumar, R., Sivakumar, D., 2003b. Comparing top  $k$  lists. SIAM Journal on Discrete Mathematics 17 (1), 134–160.  
URL <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.86.3234&rep=rep1&type=pdf>
- [Fréchet(1906)] Fréchet, M. R., 1906. Sur quelques points du calcul fonctionnel (on some points of functional calculation). Rendiconti del Circolo Matematico di Palermo 22, 1–74, rendiconti del Circolo Matematico di Palermo (Statements of the Mathematical Circle of Palermo).



- [Fréchet(1928)] Fréchet, M. R., 1928. Les Espaces abstraits et leur théorie considérée comme introduction à l'analyse générale. Borel series. Gauthier-Villars, Paris, abstract spaces and their theory regarded as an introduction to general analysis.  
URL <http://books.google.com/books?id=9czoHQAACAAJ>
- [Galvin and Shore(1984)] Galvin, F., Shore, S. D., March 1984. Completeness in semimetric spaces. Pacific Journal Of Mathematics 113 (1), 67–75.  
URL <http://msp.org/pjm/1984/113-1/pjm-v113-n1-p04-s.pdf>
- [Gemignani(1972)] Gemignani, M. C., 1972. Elementary Topology, 2nd Edition. Addison-Wesley Series in Mathematics. Addison-Wesley Publishing Company, Reading, Massachusetts, a 1990 Dover “unabridged and corrected” edition has been published.  
URL <http://books.google.com/books?vid=ISBN0486665224>
- [Gibbons et al.(1977)] Gibbons, Olkin, and Sobel] Gibbons, J. D., Olkin, I., Sobel, M., 1977. Selecting and Ordering Populations: A New Statistical Methodology. John Wiley & Sons, New York, a 1999 unabridged and corrected republication has been made available as a “SIAM Classics edition”, ISBN 9781611971101.  
URL <http://books.google.com/books?vid=ISBN1611971101>
- [Giles(1987)] Giles, J. R., 1987. Introduction to the Analysis of Metric Spaces. No. 3 in Australian Mathematical Society lecture series. Cambridge University Press, Cambridge.  
URL <http://books.google.com/books?vid=ISBN0521359287>
- [Haaser and Sullivan(1991)] Haaser, N. B., Sullivan, J. A., 1991. Real Analysis. Dover Publications, New York.  
URL <http://books.google.com/books?vid=ISBN0486665097>
- [Hahn and Rosenthal(1948)] Hahn, H., Rosenthal, A., 1948. Set Functions. University of New Mexico Press.  
URL <http://books.google.com/books?vid=ISBN111422295X>
- [Halmos(1950)] Halmos, P. R., 1950. Measure Theory. The University series in higher mathematics. D. Van Nostrand Company, New York, 1976 reprint edition available from Springer with ISBN 9780387900889.  
URL <http://www.amazon.com/dp/0387900888>
- [Halmos(1960)] Halmos, P. R., 1960. Naive Set Theory. The University Series in Undergraduate Mathematics. D. Van Nostrand Company, Inc., Princeton, New Jersey.  
URL <http://books.google.com/books?vid=isbn0387900926>
- [Hardy et al.(1952)] Hardy, Littlewood, and Pólya] Hardy, G. H., Littlewood, J. E., Pólya, G., 1952. Inequalities, 2nd Edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge.  
URL <http://books.google.com/books?vid=ISBN0521358809>
- [Hausdorff(1914)] Hausdorff, F., 1914. Grundzüge der Mengenlehre. Von Veit, Leipzig, properties of Set Theory.  
URL <http://books.google.com/books?id=KTs4AAAAAAAJ>
- [Hausdorff(1937)] Hausdorff, F., 1937. Set Theory, 3rd Edition. Chelsea Publishing Company, New York, 1957 translation of the 1937 German *Grundzüge der Mengenlehre*.  
URL <http://books.google.com/books?vid=ISBN0828401195>
- [Heath(1961)] Heath, R. W., 1961. A regular semi-metric space for which there is no semi-metric under which all spheres are open. Proceedings of the American Mathematical Society 12, 810–811.  
URL <http://www.ams.org/journals/proc/1961-012-05/S0002-9939-1961-0125562-9/>
- [Heinonen(2001)] Heinonen, J., January 1 2001. Lectures on Analysis on Metric Spaces. Universitext Series. Springer Science & Business Media.  
URL <http://books.google.com/books?vid=ISBN0387951040>
- [Hoehn and Niven(1985)] Hoehn, L., Niven, I., 1985 1985. Mathematics Magazine 58 (3), 151–156.  
URL <http://www.jstor.org/stable/2689911>
- [Jensen(1906)] Jensen, J. L. W. V., December 1906. Sur les fonctions convexes et les in'egalite's entre les valeurs moyennes (on the convex functions and the inequalities between the average values). Acta Mathematica 30 (1), 175–193.  
URL <http://www.springerlink.com/content/r55q1411g840j446/>





- [Jiménez and Yukich(2006)] Jiménez, R., Yukich, J. E., 2006. Statistical distances based on Euclidean graphs. pp. 223–240.  
URL <http://books.google.com/books?vid=ISBN0387233946>
- [Joshi(1983)] Joshi, K. D., 1983. Introduction To General Topology, 1st Edition. New Age International.  
URL <http://books.google.com/books?vid=ISBN0852264445>
- [Kelley(1955)] Kelley, J. L., 1955. General Topology. University Series in Higher Mathematics. Van Nostrand, New York, republished by Springer-Verlag, New York, 1975.  
URL <http://books.google.com/books?vid=ISBN0387901256>
- [Khamssi and Kirk(2001)] Khamssi, M. A., Kirk, W., 2001. An Introduction to Metric Spaces and Fixed Point Theory. John Wiley, New York.  
URL <http://books.google.com/books?vid=isbn0471418250>
- [Kirk and Shahzad(2014)] Kirk, W., Shahzad, N., October 23 2014. Fixed Point Theory in Distance Spaces. SpringerLink: Bücher. Springer.  
URL <http://books.google.com/books?vid=isbn3319109278>
- [Korselt(1894)] Korselt, A., March 1894. Bemerkung zur algebra der logik. Mathematische Annalen 44 (1), 156–157, referenced by Birkhoff(1948)p.133.  
URL <http://www.springerlink.com/content/v681m56871273j73/>
- [Krishnamurthy(1966)] Krishnamurthy, V., February 1966. On the number of topologies on a finite set. The American Mathematical Monthly 73 (2), 154–157.  
URL <http://www.jstor.org/stable/2313548>
- [Kubrusly(2001)] Kubrusly, C. S., 2001. The Elements of Operator Theory, 1st Edition. Springer.  
URL <http://books.google.com/books?vid=ISBN0817641742>
- [Kubrusly(2011)] Kubrusly, C. S., 2011. The Elements of Operator Theory, 2nd Edition. Springer.  
URL <http://books.google.com/books?vid=ISBN0817649972>
- [Laos(1998)] Laos, N. K., 1998. Topics in Mathematical Analysis and Differential Geometry. Vol. 24 of Series in pure mathematics. World Scientific.  
URL <http://books.google.com/books?vid=ISBN9810231806>
- [Leathem(1905)] Leathem, J. G., 1905. Volume and surface integrals used in physics, 1st Edition. Vol. 1 of Cambridge Tracts in Mathematics and Mathematical Physics. Cambridge University Press.  
URL <http://archive.org/details/volumesurfaceint01leatuoft>
- [Lin(1998)] Lin, D., 1998. An information-theoretic definition of similarity. In: Proceedings of the International Conference on Machine Learning.  
URL <http://webdocs.cs.ualberta.ca/~lindek/papers/sim.pdf>
- [MacLane and Birkhoff(1999)] MacLane, S., Birkhoff, G., 1999. Algebra, 3rd Edition. AMS Chelsea Publishing, Providence.  
URL <http://books.google.com/books?vid=isbn0821816462>
- [Maddux(2006)] Maddux, R. D., July 27 2006. Relation Algebras, 1st Edition. Elsevier Science.  
URL <http://books.google.com/books?vid=ISBN0444520139>
- [Maligranda(1995)] Maligranda, L., March 1995. A simple proof of the hölder and the minkowski inequality. The American Mathematical Monthly 102 (3), 256–259.  
URL <http://www.jstor.org/stable/2975013>
- [Maligranda and Orlicz(1987)] Maligranda, L., Orlicz, W., March 01 1987. On some properties of functions of generalized variation. Monatshefte für Mathematik 104 (1), 53–65.  
URL [http://gdz.sub.uni-goettingen.de/download/PPN362162050\\_0104/LOG\\_0008.pdf](http://gdz.sub.uni-goettingen.de/download/PPN362162050_0104/LOG_0008.pdf)
- [McCarty(1967)] McCarty, G., 1967. Topology: An Introduction With Application to Topological Groups. International Series in Pure and Applied Mathematics. McGraw-Hill Book Company, New York, 1988 Dover edition available.  
URL <http://www.amazon.com/dp/0486656330>



- [Menger(1928)] Menger, K., 1928. Untersuchungen über allgemeine metrik. *Mathematische Annalen* 100, 75–163, (Investigations on general metric).  
URL <http://link.springer.com/article/10.1007%2FBF01455705>
- [Michel and Herget(1993)] Michel, A. N., Herget, C. J., 1993. *Applied Algebra and Functional Analysis*. Dover Publications, Inc., original version published by Prentice-Hall in 1981.  
URL <http://books.google.com/books?vid=ISBN048667598X>
- [Milovanović and Milovanović(1979)] Milovanović, G. V., Milovanović, I., 1979. On a generalization of certain results of a. ostrowski and a. lupaş. University of Beograd. *Publikacije Elektrotehničkog Fakulteta* (634–677), 62–69, ser. Mat. Fiz.  
URL <http://www.mi.sanu.ac.rs/~gvm/radovi/643.pdf>
- [Minkowski(1910)] Minkowski, H., 1910. *Geometrie der Zahlen*. Druck und Verlag von B.G. Teubner, Leipzig, geometry of Numbers.  
URL <http://www.archive.org/details/geometriederzahl00minkrich>
- [Mitrinović et al.(2010)] Mitrinović, D. S., Pečarić, J. E., Fink, A. M., 2010. *Classical and New Inequalities in Analysis*. Vol. 61 of *Mathematics and its Applications (East European Series)*. Kluwer Academic Publishers, Dordrecht, Boston, London.  
URL <http://www.amazon.com/dp/0792320646>
- [Molchanov(2005)] Molchanov, I. S., 2005. *Theory of Random Sets. Probability and Its Applications*. Springer.  
URL <http://books.google.com/books?vid=ISBN185233892X>
- [Mulholland(1950)] Mulholland, H. P., 1950. On generalizations of minkowski's inequality in the form of a triangle inequality. *Proceedings of the London Mathematical Society* s2-51, 294–307, received 1946 October 10, read 1947 June 19.  
URL <http://plms.oxfordjournals.org/content/s2-51/1/294.extract>
- [Munkres(2000)] Munkres, J. R., 2000. *Topology*, 2nd Edition. Prentice Hall, Upper Saddle River, NJ.  
URL <http://www.amazon.com/dp/0131816292>
- [Murdeswar(1990)] Murdeswar, M. G., 1990. *General Topology*, 2nd Edition. New Age International.  
URL <http://books.google.com/books?vid=isbn8122402461>
- [Ore(1935)] Ore, O., April 1935. On the foundation of abstract algebra. i. *The Annals of Mathematics* 36 (2), 406–437.  
URL <http://www.jstor.org/stable/1968580>
- [Pap(1995)] Pap, E., 1995. *Null-Additive Set Functions*. Vol. 337 of *Mathematics and Its Applications*. Kluwer Academic Publishers.  
URL <http://www.amazon.com/dp/0792336585>
- [Pečarić et al.(1992)] Pečarić, J. E., Proschan, F., Tong, Y. L., 1992. *Convex Functions, Partial Orderings, and Statistical Applications*. Vol. 187 of *Mathematics in Science and Engineering*. Academic Press, San Diego, California.  
URL <http://books.google.com/books?vid=ISBN0125492502>
- [Ponnusamy(2002)] Ponnusamy, S., 2002. *Foundations of Functional Analysis*. CRC Press.  
URL <http://books.google.com/books?vid=ISBN0849317177>
- [Rana(2002)] Rana, I. K., 2002. *An Introduction to Measure and Integration*, 2nd Edition. Vol. 45 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, R.I.  
URL <http://books.google.com/books?vid=ISBN0821829742>
- [Riesz(1909)] Riesz, F., 1909. Stetigkeitsbegriff und abstrakte mengenlehre. In: *Castelnuovo, G. (Ed.), Atti del IV Congresso Internazionale dei Matematici*. Vol. II. Tipografia della R. Accademia dei Lincei, Rome, pp. 18–24, 1908 April 6–11.  
URL <http://www.mathunion.org/ICM/ICM1908.2/Main/icm1908.2.0018.0024.ocr.pdf>



- [Rosenlicht(1968)] Rosenlicht, M., 1968. Introduction to Analysis. Dover Publications, New York.  
URL <http://books.google.com/books?vid=ISBN0486650383>
- [Rudin(1976)] Rudin, W., 1976. Principles of Mathematical Analysis, 3rd Edition. McGraw-Hill, New York, library QA300.R8 1976.  
URL <http://books.google.com/books?vid=ISBN007054235X>
- [Salzmann et al.(2007)Salzmann, Grundhöfer, Hähl, and Löwen] Salzmann, H., Grundhöfer, T., Hähl, H., Löwen, R., 2007. The Classical Fields: Structural Features of the Real and Rational Numbers. Vol. 112 of Encyclopedia of Mathematics and its Applications. Cambridge University Press.  
URL <http://books.google.com/books?vid=ISBN0521865166>
- [Schweizer and Sklar(1983)] Schweizer, B., Sklar, A., 1983. Probabilistic Metric Spaces. Elsevier Science Publishing Co., a 2005 Dover edition (ISBN 9780486143750) has been published which is “an unabridged republication of the work first published by Elsevier Science Publishing Co., Inc., in 1983.”  
URL <http://books.google.com/books?vid=ISBN0486143759>
- [Searcóid(2006)] Searcóid, M. O., 2006. Metric Spaces. Springer Undergraduate Mathematics Series. Springer Science & Business Media.  
URL <http://books.google.com/books?vid=ISBN1846286271>
- [Sherstnev(1962)] Sherstnev, A. N., 1962. Random normed spaces. questions of completeness. Kazan. Gos. Univ. Uchen. Zap. 122 (4), pages 3–20.  
URL <http://mi.mathnet.ru/uzku138>
- [Simon(2011)] Simon, B., May 19 2011. Convexity: An Analytic Viewpoint. Vol. 187 of Cambridge Tracts in Mathematics. Cambridge University Press.  
URL <http://books.google.com/books?vid=ISBN1107007313>
- [Steen and Seebach(1978)] Steen, L. A., Seebach, J. A., 1978. Counterexamples in Topology, 2nd Edition. Springer-Verlag, a 1995 “unabridged and unaltered republication” Dover edition is available.  
URL <http://books.google.com.tw/books?vid=ISBN0486319296>
- [Suppes(1972)] Suppes, P., 1972. Axiomatic Set Theory. Dover Publications, New York.  
URL <http://books.google.com/books?vid=ISBN0486616304>
- [Sutherland(1975)] Sutherland, W. A., 1975. Introduction to Metric and Topological Spaces, illustrated, reprint Edition. Oxford University Press.  
URL <http://books.google.com/books?vid=ISBN0198531613>
- [Szirmai(2007)] Szirmai, J., 2007. The densest geodesic ball packing by a type of nil lattices. Contributions to Algebra and Geometry 48 (2), 383–397.  
URL <http://www.emis.ams.org/journals/BAG/vol.48/no.2/b48h2szi.pdf>
- [Thomson et al.(2008)Thomson, Bruckner, and Bruckner] Thomson, B. S., Bruckner, A. M., Bruckner, J. B., 2008. Elementary Real Analysis, 2nd Edition. www.classicalrealanalysis.com.  
URL <http://classicalrealanalysis.info/com/Elementary-Real-Analysis.php>
- [Thron(1966)] Thron, W. J., 1966. Topological structures. Holt, Rinehart and Winston, New York.  
URL [http://books.google.com/books?id=JRM\\_AAAAIAAJ](http://books.google.com/books?id=JRM_AAAAIAAJ)
- [Tietze(1923)] Tietze, H. F. F., 1923. Beiträge zur allgemeinen topologie i. Mathematische Annalen 88 (3–4), 290–312.  
URL <http://link.springer.com/article/10.1007%2FBF01579182>
- [Tolsted(1964)] Tolsted, E., 1964. An elementary derivation of cauchy, hölder, and minkowski inequalities from young's inequality. Mathematics Magazine 37, 2–12.  
URL <http://mathdl.maa.org/mathDL/22/?pa=content&sa=viewDocument&nodeId=3036>
- [Vallin(1999)] Vallin, R. W., 1999. Continuity and differentiability aspects of metric preserving functions. Real Analysis Exchange 25 (2), 849–868.  
URL [projecteuclid.org/euclid.rae/1230995419](http://projecteuclid.org/euclid.rae/1230995419)



- [Veltkamp(2001)] Veltkamp, R. C., May 2001. Shape matching: Similarity measures and algorithms. Shape Modeling and Applications, SMI 2001 International Conference on, 188–197.  
URL <http://dspace.library.uu.nl/bitstream/handle/1874/2540/2001-03.pdf>
- [Veltkamp and Hagedoorn(2000)] Veltkamp, R. C., Hagedoorn, M., November 2–4 2000. Shape similarity measures, properties and constructions. In: Laurini, R. (Ed.), Advances in Visual Information Systems 4th International Conference, VISUAL 2000 Lyon. Vol. 1929 of Lecture Notes in Computer Science. Springer, France, pp. 467–476.  
URL [http://link.springer.com/chapter/10.1007/3-540-40053-2\\_41](http://link.springer.com/chapter/10.1007/3-540-40053-2_41)
- [Vitányi(2011)] Vitányi, P. M. B., April 2011. Information distance in multiples. IEEE Transactions on Information Theory 57 (4).  
URL <http://arxiv.org/pdf/0905.3347.pdf>
- [Walmsley(1920)] Walmsley, C., 1920. An Introductory Course Of Mathematical Analysis. Cambridge University Press.  
URL <https://archive.org/details/introductorycour032788mbp>
- [Wilson(1931)] Wilson, W. A., April 1931. On semi-metric spaces. American Journal of Mathematics 53 (2), 361–373.  
URL <http://www.jstor.org/stable/2370790>
- [Xia(2009)] Xia, Q., April 2009. The geodesic problem in quasimetric spaces. Journal of Geometric Analysis 19 (2), 452–479.  
URL <http://link.springer.com/article/10.1007/s12220-008-9065-4>
- [Young(1912)] Young, W. H., August 1912. On classes of summable functions and their fourier series. Proceedings of the Royal Society of London 87 (594), 225–229.  
URL <http://www.archive.org/details/philtrans02496252>
- [Zorich(2004)] Zorich, V. A., 2004. Mathematical Analysis I. Universitext Series. Springer Science & Business Media.  
URL <http://books.google.com/books?vid=ISBN3540403868>

## Reference Index

- |  |   |   |
|--|---|---|
| [Aliprantis and Burkinshaw(1998)], 6, 19, 21, 27, 36 | [Bruckner et al.(1997)Bruckner, Bruckner, and Thomson], 6     | [Costa et al.(2004)Costa, Castro, Rowstron, and Key], 2 |
| [Adams and Franzosa(2008)], 8                        | [Brunschwig et al.(2003)Brunschwig, Lloyd, and Pellegrin], 15 | [Crammer et al.(2007)Crammer, Kearns, and Wortman], 2   |
| [Apostol(1975)], 5, 9, 21                            | [Bryant(1985)], 10  | [Crossley(2006)], 28                                    |
| [Barvinok(2002)], 30                                 | [Bullen(1990)], 31  | [Czerwik(1993)], 16                                     |
| [Beran(1985)], 5                                     | [Bullen(2003)], 31, 34, 35                                    | [Davis(2005)], 11, 27, 28                               |
| [Berberian(1961)], 18, 23                            | [Burstein et al.(2005)Burstein, Ulitsky, Tuller, and Chor], 2 | [Dedekind(1900)], 5                                     |
| [Bessenyei and Pales(2014)], 6, 16                   | [Carothers(2000)], 31, 35                                     | [Deza and Deza(2006)], 36                               |
| [Besso(1879)], 31                                    | [Cauchy(1821)], 34  | [Deza and Deza(2009)], 35, 36                           |
| [Bienaymé(1840)], 31                                 | [Chatterji(1967)], 8  | [Deza and Deza(2014)], 6, 16                            |
| [Blumenthal(1938)], 6                                | [Choquet(1954)], 5  | [DiBenedetto(2002)], 26                                 |
| [Blumenthal(1953)], 6, 11–15                         | [Cohn(2002 December 6)], 5                                    | [Dieudonné(1969)], 7, 16, 18                            |
| [Bollobás(1999)], 27, 30, 31, 34                     | [Comtet(1966)], 8   | [Euclid(circa 300BC)], 16                               |
| [Bottazzini(1986)], 4                                | [Comtet(1974)], 4, 8  | [Evans et al.(1967)Evans, Harary, and Lynn], 8          |
| [Bourbaki(1939)], 4                                  | [Copson(1968)], 6, 16   | [Fagin et al.(2003a)Fagin, Kumar, and Sivakumar], 16    |
| [Brenner(1985)], 31                                  | [Corazza(1999)], 35, 36                                       |   |
| [Brown and Watson(1996)], 8                          |   |   |



- [Fagin et al.(2003b)Fagin, Kumar, and Sivakumar], 16  
 [Fréchet(1906)], 16  
 [Fréchet(1928)], 16  
 [Galvin and Shore(1984)], 6  
 [Gemignani(1972)], 27  
 [Gibbons et al.(1977)Gibbons, Olkin, and Sobel], 16  
 [Giles(1987)], 9, 21, 36  
 [Haaser and Sullivan(1991)], 10, 29  
 [Hahn and Rosenthal(1948)], 4, 5  
 [Halmos(1950)], 5  
 [Halmos(1960)], 4  
 [Hardy et al.(1952)Hardy, Littlewood, and Pólya], 31, 35  
 [Hausdorff(1914)], 26  
 [Hausdorff(1937)], 6, 11, 16, 26  
 [Heath(1961)], 6  
 [Heinonen(2001)], 16  
 [Hoehn and Niven(1985)], 16, 34  
 [Jensen(1906)], 30, 34  
 [Jiménez and Yukich(2006)], 2  
 [Joshi(1983)], 8, 27, 28  
 [Kelley(1955)], 4, 27, 36  
 [Khamsi and Kirk(2001)], 6, 9  
 [Kirk and Shahzad(2014)], 16  
 [Korselt(1894)], 5  
 [Krishnamurthy(1966)], 8  
 [Kubrusly(2001)], 10, 27, 29  
 [Kubrusly(2011)], 26  
 [Laos(1998)], 6, 15  
 [Leathem(1905)], 9, 28  
 [Lin(1998)], 2  
 [MacLane and Birkhoff(1999)], 5  
 [Maddux(2006)], 4  
 [Maligranda and Orlicz(1987)], 2  
 [Maligranda(1995)], 31, 35  
 [McCarty(1967)], 27, 28  
 [Menger(1928)], 6  
 [Michel and Herget(1993)], 6, 18  
 [Milovanović and Milovanović(1979)], 31  
 [Minkowski(1910)], 31, 35  
 [Mitrinović et al.(2010)Mitrinović, Pečarić, and Fink], 30  
 [Molchanov(2005)], 4, 6  
 [Mulholland(1950)], 31  
 [Munkres(2000)], 26, 27, 29  
 [Murdeshwar(1990)], 8  
 [Ore(1935)], 5  
 [Pap(1995)], 4, 5  
 [Pečarić et al.(1992)Pečarić, Proschan, and Tong], 31  
 [Ponnusamy(2002)], 28, 29  
 [Rana(2002)], 4  
 [Riesz(1909)], 26  
 [Rosenlicht(1968)], 7, 9, 10, 21–23, 29  
 [Rudin(1976)], 33, 34  
 [Salzmann et al.(2007)Salzmann, Grundhöfer, Hähl, and Löwen], 8  
 [Schweizer and Sklar(1983)], 16  
 [Searcoid(2006)], 28  
 [Sherstnev(1962)], 16  
 [Simon(2011)], 30  
 [Steen and Seebach(1978)], 26  
 [Suppes(1972)], 4  
 [Sutherland(1975)], 10  
 [Szirmai(2007)], 2  
 [Thomson et al.(2008)Thomson, Bruckner, and Bruckner], 23  
 [Thron(1966)], 6, 27  
 [Tietze(1923)], 26  
 [Tolsted(1964)], 31, 35  
 [Vallin(1999)], 35, 36  
 [Veltkamp and Hagedoorn(2000)], 2  
 [Veltkamp(2001)], 2  
 [Vitányi(2011)], 2  
 [Walmsley(1920)], 15  
 [Wilson(1931)], 6  
 [Xia(2009)], 16  
 [Young(1912)], 35  
 [Zorich(2004)], 19

## Subject Index

- $\alpha$ -scaled, 36  
 $\alpha$ -scaled metric, 36  
 $\alpha$ -truncated, 36  
 $\alpha$ -truncated metric, **36**  
 $\phi$ -mean, **31**  
 $\sigma$ -inframetric inequality, 3, 16, **16**  
 $\sigma$ -inframetric space, 3, **16**  
 absolute value, 5, **5**, 12, 18  
 accumulation point, 3, 4, **27**  
 adherent, **27**  
 affine, 17, **30**  
 AM-GM inequality, 34  
 anti-symmetric, 5  
 arithmetic mean, 1, 3, 17  
 arithmetic mean geometric mean inequality, 34  
 b-metric, 16  
 base, 3, 4, 6, 8, 9, 12–14, 20, 27, **27**  
 boundary, 27  
 bounded, 2, **6**, 9, 10, 21, 22, 36  
 bounded metric, 36  
 Cantor Intersection Theorem, 2  
 Cantor intersection theorem, 11, **11**  
 cardinality, 5  
 cardinality of  $A$ , 5  
 Cartesian product, 4  
 Cauchy, 2, 3, 6, 9, 10, 13, 14, 21, 22, 25  
 Cauchy condition, 9  
 Cauchy sequence, **9**  
 Cauchy sequences, 9  
 closed, 7, 7, 10, 11, **26**, 27, 29, 30  
 closed ball, **6**  
 closed interval, 5  
 closed set, 3, 4, 7  
 closed set theorem, 30  
 closure, 3, 4, 10, 27, **27**, 29  
 closure point, **27**  
 commutative, 18, 23  
 commutative ring, 5  
 complete, 9, 10, 11  
 concave, 30, **30**, 31, 32  
 continuity, 3, 4, 12, 15, 17, 28–30  
 continuous, 1, 3, 6, 11, **11**, 12, 15, 17, 23, 25, 26, 28, **28**, 29–32  
 contradiction, 24  
 converge, **9**  
 convergence, 3, 4, 13, 22, 30  
 metric space, 21



- convergent, 3, 6, 9, 10–14, 21, 22, 25, **29**, 30
- convergent sequence, 3
- converges, **28**, 29
- convex, 30, **30**, 31, 32
  - functional, 30
  - strictly, 30
- decreasing, 31
- definitions
  - $\sigma$ -inframetric space, **16**
  - accumulation point, **27**
  - base, **27**
  - Cartesian product, **4**
  - Cauchy sequence, **9**
  - closed ball, **6**
  - closed interval, **5**
  - closed set, **7**
  - closure point, **27**
  - distance space, **6**
  - empty set, **5**
  - extended real numbers, **4**
  - function, **4**
  - half-open interval, **5**
  - inframetric space, **16**
  - interior, **27**
  - interior point, **27**
  - metric space, **16**
  - near metric space, **16**
  - open ball, **6**
  - open interval, **5**
  - open set, **7**
  - ordered pair, **4**
  - ordered set, **5**
  - partially ordered set, **5**
  - paving, **4**
  - point of adherence, **27**
  - poset, **5**
  - power distance space, **16**
  - relation, **4**
  - set function, **5**
  - set of all functions, **4**
  - set of all relations, **4**
  - set of integers, **4**
  - set of natural numbers, **4**
  - set of non-negative real numbers, **4**
  - set of positive real numbers, **4**
  - set of real numbers, **4**
  - set structure, **4**
  - topological space induced by  $(X, d)$ , **8**
  - topology, **26**
  - topology induced by  $(X, d)$  on  $X$ , **8**
  - diameter, **6**
- dilated, 36
- dilated Euclidean, 14, 25
- dilated metric, 36
- discontinuous, **11**, 12–15, 28, **28**
- discrete, 13, 24, 36
- discrete metric preserving function, **36**
- discrete topology, 26, 28
- distance, 6, **6**, 12–15
- distance function, 11, 12, 14, 15
- distance space, 1–6, **6**, 7–17, 21, 23–25
- distance spaces, 3
- diverge, 14, **29**
- divergent, **29**
- empty set, **5**, 7
- Euclidean, 12–14, 24, 25
- Euclidean metric space, 25
- examples
  - $\alpha$ -scaled, 36
  - $\alpha$ -scaled metric, 36
  - $\alpha$ -truncated, 36
  - $\alpha$ -truncated metric, **36**
  - bounded, 36
  - bounded metric, 36
  - dilated, 36
  - dilated metric, 36
  - discrete, 36
  - discrete metric preserving function, **36**
  - power transform, 36
  - power transform metric, 36
  - radar screen, **36**
  - radar screen metric, **36**
  - snowflake, 36
  - snowflake transform metric, 36
  - the standard topology on the real line, **27**
- extended real numbers, **4**, 15
- extended set of integers, 4
- finite, 5
- function, **4**, 28
- functions
  - absolute value, 5, **5**, 12, 18
  - arithmetic mean, 17
  - cardinality of  $A$ , 5
  - distance, 6, **6**, 12–15
  - distance function, 11, 14, 15
  - geometric mean, 17
  - harmonic mean, 17
  - maximum, 17
  - metric, 14
  - metric function, 3, 12
  - metric preserving function, 35, **35**, 36
  - minimum, 17
  - modulus, **5**
  - power distance, **16**
  - power distance function, **16**, 17, 26
  - Power mean, 3, 15
  - power mean, 1, 3, 15, 31, **31**
  - power triangle function, 1, 3, 15, **15**, 16, 17, 22, 23
  - quadratic mean, 17
  - sequence, 6, 9–11
  - set function, 5
  - triangle function, 16, 25
- Generalized AM-GM inequality, 35
- generalized arithmetic mean
- geometric mean inequality, 34
- geometric inequality, **16**, 18
- geometric mean, 1, 3, 17
- half-open interval, **5**, 12
- harmonic inequality, **16**, 18
- harmonic mean, 1, 3, 17
- homogeneous, 5
- horizontal half-open interval, 12, 24
- increasing, 31
- indiscrete topological space, 8
- indiscrete topology, 26, 28
- inequalities
  - AM-GM, 34
  - Jensen's, 30
  - Minkowski (sequences), 35
  - power triangle inequality, 23
  - Young, 35
- inframetric inequality, 1, 16, **16**, 18
- inframetric space, **16**, 17
- interior, 3, 4, 27, **27**
- interior point, **27**
- inverse image characterization, 12
- inverse image characterization of continuity, 12, 29
- isotone, 36
- Jensen's Inequality, **30**, 35
- l'Hôpital's rule, 34



- limit, **9**, 10, 29, **29**
- limit point, **29**
- linearly ordered, 8
- maximum, 1, 3, 17
- mean square, 1, 3
- metric, 14, 35
- metric function, 3, 12
- metric preserving function, 35, **35**, 36
- Metric space, 1, 2
- metric space, 2–16, **16**, 18, 21–25, 27–29, 35, 36
- metrics
  - $\alpha$ -scaled metric, 36
  - $\alpha$ -truncated metric, 36
  - bounded, 36
  - dilated metric, 36
  - power transform metric, 36
  - radar screen, 36
- minimal inequality, **16**, 18
- minimum, 1, 3, 17
- Minkowski's Inequality, 31
- Minkowski's Inequality for sequences, **35**
- modulus, **5**
- monotone, 3
- monotone, 1, 19
- near metric space, 3, 16, **16**, 17
- non-negative, 5, 6, 18, 24, 31
- nondegenerate, 5, 6, 19, 24
- not Cauchy, 14
- not closed, 30
- not open, 12, 13
- not unique, 13, 23, 29
- open, 2–4, 6, 7, 7, 8, 9, 12–14, 20, 21, 26, **26**, 27–29
- open ball, 3, 4, 6, **6**, 7–9, 12–14, 19–21
- open balls, 12, 20
- open interval, **5**, 12
- open set, 7, 7, 8, 12, 13, 20, 27–29
- operations
  - $\phi$ -mean, **31**
  - arithmetic mean, 1, 3
  - closure, **27**
  - geometric mean, 1, 3
  - harmonic mean, 1, 3
  - limit, **9**, **29**
  - maximum, 1, 3
  - mean square, 1, 3
  - minimum, 1, 3
  - open set, 7
  - set difference, 7
  - set intersection, 7
  - SMR, 16
  - square-mean-root, 16
- order relation, **5**, **5**
- ordered metric space, 16
- ordered pair, **4**, 12
- ordered set, **5**, **5**
- ordered set of real numbers, 5
- partial order relation, **5**
- partially ordered set, **5**
- paving, **4**, 5
- point of adherence, **27**
- poset, **5**
- positive real numbers, 15
- power distance, **16**
- power distance function, **16**, 17, 26
- power distance space, 3, 4, **16**, 17, 19–25
- power distance spaces, 3
- Power mean, 3, 15
- power mean, 1, 3, 15, 31, **31**, 34
- power set, 4, 6, 26, 27
- power set of  $X$ , 4
- power transform, 36
- power transform metric, 36
- power triangle function, 1, 3, 15, **15**, 16, 17, 22, 23
- power triangle inequality, 1, 3, 15, 16, **16**, 17, 18, 21–23, 25, 26
- power triangle triangle space, 18
- preorder, **5**
- probabilistic metric spaces, 16
- properties
  - adherent, **27**
  - affine, 17, **30**
  - anti-symmetric, 5
  - bounded, 2, **6**, 9, 10, 21, 22
  - Cauchy, 2, 3, 6, 9, 10, 13, 14, 21, 22, 25
  - Cauchy condition, 9
  - closed, 7, 7, 10, 11, **26**, 27, 29, 30
  - closed set, 4
  - commutative, 18, 23
  - complete, **9**, 10, 11
  - concave, 30, **30**, 31, 32
  - continuity, 3, 4, 12, 15, 17, 28–30
  - continuous, 1, 3, 6, 11, **11**, 12, 15, 17, 23, 25, 26, 28, **28**, 29–32
  - converge, **9**
  - convergence, 3, 4, 13, 22, 30
  - convergent, 3, 6, 9, **9**, 10–14, 21, 22, 25, **29**, 30
  - converges, **28**, 29
  - convex, 30, **30**, 31, 32
  - decreasing, 31
  - dilated Euclidean, 14, 25
  - discontinuous, **11**, 12–15, 28, **28**
  - discrete, 13, 24
  - distance space, 11
  - diverge, 14, **29**
  - divergent, **29**
  - Euclidean, 12–14, 24, 25
  - finite, 5
  - homogeneous, 5
  - increasing, 31
  - isotone, 36
  - limit, **9**
  - linearly ordered, 8
  - monotone, 3
  - monotone, 1, 19
  - non-negative, 5, 6, 18, 24, 31
  - nondegenerate, 5, 6, 19, 24
  - not Cauchy, 14
  - not closed, 30
  - not open, 12, 13
  - not unique, 13, 23, 29
  - open, 2, 3, 6, 7, 7, 8, 9, 12–14, 20, 21, 26, **26**, 27, 28
  - open set, 8
  - power triangle inequality, 17, 18, 26
  - reflexive, 5
  - strictly convex, **30**
  - strictly monotone, 15
  - strictly monotone, 10, 17, 18, 22, 25, 31, 32
  - strictly monotonic, 31
  - subadditive, 5, 35, 36
  - submultiplicative, 5
  - symmetric, 6, 18, 19, 22
  - transitive, 5
  - triangle inequality, 2, 3, 5, 12–15, 23, 25
  - unique, 3, 6, 10, 12–14, 25, 29
- quadratic inequality, **16**
- quadratic mean, 17
- quadratic mean inequality, 18
- radar screen, **36**
- radar screen metric, **36**
- range, 35



- real line, 8
- reflexive, 5
- relation, 4, **4**, 16
- relations
  - $\sigma$ -inframetric inequality, 3, 16, **16**
  - geometric inequality, **16**
  - harmonic inequality, **16**
  - inframetric inequality, 1, 16, **16**
  - minimal inequality, **16**
  - order relation, **5**
  - partial order relation, **5**
  - power triangle inequality, 1, 3, 15, 16, **16**, 21, 22, 25
  - preorder, **5**
  - quadratic inequality, **16**
  - relaxed triangle inequality, 1, 3, 16, **16**
  - square mean root inequality, **16**
  - triangle inequality, 1, 2, 15, 16, **16**
  - triangle relation, 24, 25
- relaxed triangle inequality, 1, 3, 16, **16**
- sequence, 6, 9–11
- sequences
  - Cauchy, 9
- sequential characterization, 12
- sequential characterization of continuity, 12, 29
- set, 5, 7
- set difference, 7
- set function, 5, **5**
- set intersection, 7
- set of all functions, **4**
- set of all relations, **4**
- set of extended real numbers, 3
- set of integers, **4**
- set of natural numbers, **4**
- set of non-negative real numbers, **4**
- set of positive real numbers, 3, 6
- set of positive real numbers, **4**
- set of real numbers, **4**
- set structure, **4**
- sets, 4
- SMR, 16
- snowflake, 36
- snowflake transform metric, 36
- space
  - metric, 7
  - topological, 26
- square mean root (SMR) in-
  - equality, 18
  - square mean root inequality, **16**
  - square-mean-root, 16
  - strictly convex, 30, **30**
  - strictly monotone, 15
  - strictly monotone, 10, 17, 18, 22, 25, 31, 32
  - strictly monotonic, 31
  - structures
    - $\sigma$ -inframetric space, 3, **16**
- 27
  - b-metric, 16
  - base, 3, 4, 6, 8, 9, 12–14, 20, 27, **27**
  - boundary, 27
  - Cartesian product, **4**
  - Cauchy sequence, **9**
  - closed ball, **6**
  - closed interval, **5**
  - closed set, 3, 7
  - closure, 3, 4, 10, 27, 29
  - closure point, **27**
  - commutative ring, 5
  - convergent sequence, 3
  - discrete topology, 26, 28
  - distance function, 12
  - distance space, 1–6, **6**, 7–17, 21, 23–25
  - distance spaces, 3
  - empty set, **5**, 7
  - Euclidean metric space, 25
  - extended real numbers, **4**, 15
  - extended set of integers, 4
  - function, **4**, 28
  - geometric inequality, 18
  - half-open interval, **5**, 12
  - harmonic inequality, 18
  - horizontal half-open interval, 12, 24
  - indiscrete topological space, 8
  - indiscrete topology, 26, 28
  - inframetric inequality, 18
  - inframetric space, **16**, 17
  - interior, 3, 4, 27, **27**
  - interior point, **27**
  - limit, 29
  - metric, 35
  - Metric space, 1, 2
  - metric space, 2–16, **16**, 18, 21–25, 27–29, 35, 36
  - minimal inequality, 18
  - near metric space, 3, 16, **16**, 17
  - open, 4, 29
  - open ball, 3, 4, 6, **6**, 7–9, 12–14, 19–21
  - open balls, 12, 20
  - open interval, **5**, 12
  - open set, 7, **7**, 12, 13, 20, 27–29
  - order relation, 5
  - ordered metric space, 16
  - ordered pair, **4**, 12
  - ordered set, 5, **5**
  - ordered set of real numbers, 5
  - partially ordered set, **5**
  - paving, **4**, 5
  - point of adherence, **27**
  - poset, **5**
  - positive real numbers, 15
  - power distance space, 3, 4, **16**, 17, 19–25
  - power distance spaces, 3
  - power set, 4, 6, 26, 27
  - power set of  $X$ , 4
  - power triangle inequality, 3
  - power triangle triangle space, 18
  - quadratic mean inequality, 18
  - range, 35
  - real line, 8
  - relation, 4, **4**, 16
  - set, 5, 7
  - set function, **5**
  - set of all functions, **4**
  - set of all relations, **4**
  - set of extended real numbers, 3
  - set of integers, **4**
  - set of natural numbers, **4**
  - set of non-negative real numbers, **4**
  - set of positive real numbers, 3, 6
  - set of positive real numbers, **4**
  - set of real numbers, **4**
  - set structure, **4**
  - sets, 4
  - square mean root (SMR) inequality, 18
  - topological space, 3, 4, 8, 12, 23, 26–29
  - topological space  $(X, T)$





- induced by  $(\mathbb{R}, d)$ , 13
  - topological space induced by  $(\mathbb{R}, d)$ , 12
  - topological space induced by  $(X, d)$ , 8, **8**, 10, 21
  - topological space, 8
  - topology, 3, 4, 8, 12–14, 26, **26**
  - topology induced by  $(X, d)$  on  $X$ , **8**
  - triangle inequality, 2, 18
  - triangle relation, 16
  - trivial topology, 26
  - usual metric space, 27
  - vertical half-open interval, 12, 24
    - weights, 30
- subadditive, 5, 35, 36
- submultiplicative, 5
- symmetric, 6, 18, 19, 22
- The Archimedean Property, 19
- The Closed Set Theorem, **29**
- The principle of Archimedes, 19
- the standard topology on the real line, **27**
- theorems
  - Cantor intersection, 11
  - Cantor Intersection Theorem, 2
    - Cantor intersection theorem, **11**
  - closed set theorem, 30
  - Generalized AM-GM inequality, 35
  - inverse image characterization, 12
  - inverse image characterization of continuity, 12, 29
  - Jensen's Inequality, **30**, 35
  - l'Hôpital's rule, 34
  - Minkowski's Inequality, 31
    - Minkowski's Inequality for sequences, **35**
    - sequential characterization, 12
    - sequential characterization of continuity, 12, 29
  - The Archimedean Property, 19
  - The Closed Set Theorem, **29**
  - The principle of Archimedes, 19
  - Uniqueness of limit, **23**
  - Young's Inequality, **35**
- topological space, 3, 4, 8, 12, 23, 26–29
- topological space  $(X, T)$  induced by  $(\mathbb{R}, d)$ , 13
- topological space induced by  $(\mathbb{R}, d)$ , 12
- topological space induced by  $(X, d)$ , 8, **8**, 10, 21
- topologies
  - discrete, 26
  - indiscrete, 26
  - trivial, 26
- topological space, 8
- topology, 3, 4, 8, 12–14, 26, **26**
- topology induced by  $(X, d)$  on  $X$ , **8**
- transitive, 5
- triangle function, 16, 25
- triangle inequality, 1–3, 5, 12–16, **16**, 18, 23, 25
- triangle relation, 16, 24, 25
- trivial topology, 26
- unique, 3, 6, 10, 12–14, 25, 29
- Uniqueness of limit, **23**
- usual metric space, 27
- values
  - cardinality, 5
  - contradiction, 24
  - diameter, **6**
  - limit, 10
  - limit point, **29**
- vertical half-open interval, 12, 24
- weighted, 31
- weights, 30
- Young's Inequality, **35**

*Telecommunications Engineering Department, National Chiao-Tung University, Hsinchu, Taiwan; 國立交通大學 (Gúo Lì Jiāo Tōng Dà Xué) 電信工程學系 (Diàn Xìn Gōng Chéng Xué Xì) 新竹, 台灣 (Xīn Zhú, Tái Wān)*

[dgreenhoe@gmail.com](mailto:dgreenhoe@gmail.com)

Received: aa bb 20YY      Revised: cc dd 20ZZ

