

JORDAN TRIALGEBRAS AND POST-JORDAN ALGEBRAS

FATEMEH BAGHERZADEH, MURRAY BREMNER, AND SARA MADARIAGA

ABSTRACT. We compute minimal sets of generators for the S_n -modules ($n \leq 4$) of multilinear polynomial identities of arity n satisfied by the Jordan product and the Jordan diproduct (resp. pre-Jordan product) in every triassociative (resp. tridendriform) algebra. These identities define Jordan trialgebras and post-Jordan algebras: Jordan analogues of the Lie trialgebras and post-Lie algebras introduced by Dotsenko et al., Pei et al., Vallette & Loday. We include an extensive review of analogous structures existing in the literature, and their interrelations, in order to identify the gaps filled by our two new varieties of algebras. We use computer algebra (linear algebra over finite fields, representation theory of symmetric groups), to verify in both cases that every polynomial identity of arity ≤ 6 is a consequence of those of arity ≤ 4 . We conjecture that in both cases the next independent identities have arity 8, imitating the Glennie identities for Jordan algebras. We formulate our results as a commutative square of operad morphisms, which leads to the conjecture that the squares in a much more general class are also commutative.

CONTENTS

1. Introduction	2
1.1. Operadic generalization of Lie and Jordan algebras	2
1.2. Overview of problems and methods	3
1.2.1. The symmetric operads ΣTriAss , $\Sigma\text{TriDend}$ and BW	4
1.2.2. Computational methods	4
1.2.3. Terminology and notation	4
1.3. Associative, Lie, and Jordan algebras	4
1.4. Results of this paper in context	5
2. Preliminaries on algebraic operads	6
2.1. Free nonsymmetric binary operads	6
2.2. Free symmetric binary operads	7
2.3. Identical relations and operad ideals	7
2.4. Koszul duality for quadratic operads	8
2.5. Diassociative and dendriform algebras	8
2.6. Manin black and white products	10
3. Triassociative and tridendriform algebras	10
4. Jordan trialgebras	13
4.1. Relations of arity 3	14
4.2. The operad BW : skeletons, total order, normal forms	14
4.2.1. Computing the normal form of a multilinear BW monomial	15

2010 *Mathematics Subject Classification.* Primary 17C05. Secondary 05A10, 05C30, 15A69, 15B33, 16W10, 17-04, 17A30, 17A50, 17C50, 18D50, 20C30, 68R15, 68W30.

Key words and phrases. Jordan algebras, di-, and tri-algebras; pre- and post-Jordan algebras; polynomial identities; algebraic operads; triplicators; trisuccessors; Koszul duality; combinatorics of binary trees; computer algebra; linear algebra over prime fields; representation theory of symmetric groups.

The authors thank Vladimir Dotsenko for numerous helpful discussions. The authors were supported by a Discovery Grant from NSERC, the Natural Sciences and Engineering Research Council of Canada.

4.3. Relations of arity 4	16
4.4. Triplicators	18
4.5. Representation theory of the symmetric group	19
4.5.1. All relations: kernel of the expansion map	20
4.5.2. Old relations; symmetries of the skeletons	21
4.6. Relations of arity ≥ 5 : a result, a problem, and two conjectures	22
5. Enumeration of association types and skeletons	22
5.1. One commutative operation	22
5.2. One noncommutative operation	22
5.3. Two operations: one commutative, one noncommutative	23
6. Post-Jordan algebras	23
6.1. Gröbner-Shirshov basis for the tridendriform operad	23
6.2. Relations of arity ≤ 4	24
6.3. Trisuccessors	25
6.4. Relations of arity ≥ 5 : a result and two conjectures	26
7. Concluding remarks	26
7.1. Koszul duality for nonquadratic operads	26
7.2. A commutative diagram: isomorphism of reconfigured operads	27
References	28

1. INTRODUCTION

1.1. Operadic generalization of Lie and Jordan algebras. Lie and Jordan algebras are defined by the polynomial identities of arity $n \leq 3, 4$ satisfied by the (anti)commutator in every associative algebra. Lie dialgebras (Leibniz algebras) were introduced in the early 1990s [37] together with diassociative algebras and the (anti)dicommutator; Jordan dialgebras (quasi-Jordan algebras) appeared 10 years later [52]. Dendriform algebras, governed by the Koszul dual of the diassociative operad, were introduced in the late 1990s [42]; in this case, the (anti)dicommutator produces pre-Lie [26, 53] and pre-Jordan [33] algebras. These constructions stimulated the theory of algebraic operads and were reformulated in terms of Manin white and black products. It was then a short step to triassociative and tridendriform algebras, and their Lie analogues, Lie trialgebras [21] and post-Lie algebras [50].

In this paper, we investigate the Jordan analogues, and define Jordan trialgebras and post-Jordan algebras¹. For $X \in \{\text{Lie}, \text{Jordan}\}$, X -trialgebras combine X -algebras and X -dialgebras in one structure; post- X algebras combine X -algebras and pre- X algebras in one structure. Our methods are primarily computational; we determine the multilinear identities of arity ≤ 6 satisfied by the anticommutator and anticommutator in every triassociative and tridendriform algebra. We use combinatorics of trees, linear algebra over finite fields, and representation theory of symmetric groups. The identities form the nullspace of what we call expansion matrix, which represents (with respect to ordered monomial bases) a morphism from an operad of Jordan type to one of associative type. The defining identities

¹One choice of defining relations for post-Jordan algebras appears in Appendix A.5 of the arXiv version of Bai et al. [3], where these relations are presented as defining the trisuccessor of the Jordan operad. That Appendix does not appear in the published version of the paper, and in any case, our methods to obtain them are entirely different.

have arity ≤ 4 in both cases, and we verify that there are no new identities of arity 5 or 6.

ASSOCIATIVE ALGEBRAS Operation ab Relation $(ab)c \equiv a(bc)$	Self-dual
Lie bracket $[a, b] = ab - ba$ Lie algebras , equation (1)	symmetry $ab \equiv ba$, relation $(ab)c \equiv a(bc)$ Commutative associative algebras
Jordan product $a \circ b = ab + ba$ Jordan algebras , equation (2)	No dual, operad is cubic not quadratic
DIASSOCIATIVE ALGEBRAS Operations $a \dashv b, a \vdash b$, Definition 2.9	DENDRIFORM ALGEBRAS Operations $a < b, a > b$, Definition 2.9
Leibniz bracket $\{a, b\} = a \dashv b - b \vdash a$ Leibniz algebras , Definition 2.12	Zinbiel algebras , Definition 2.12
Perm algebras , Definition 2.16	Pre-Lie product $\{a, b\} = a < b - b > a$ Pre-Lie algebras , Definition 2.16
Jordan diproduct $a \circ b = a \dashv b + b \vdash a$ Jordan dialgebras , Definition 2.18	No dual, operad is cubic not quadratic
No dual, operad is cubic not quadratic	Pre-Jordan product $a \bullet b = a < b + b > a$ Pre-Jordan algebras , Definition 2.18
TRIASSOCIATIVE ALGEBRAS Operations $a \dashv b, a \perp b, a \vdash b$ Definition 3.1	TRIDENDRIFORM ALGEBRAS Operations $a < b, a \wedge b, a > b$ Definition 3.1
Lie bracket $[a, b] = a \perp b - b \perp a$ Leibniz bracket $\{a, b\} = a \dashv b - b \vdash a$ Lie trialgebras , Definition 3.4	Commutative tridendriform algebras , Definition 3.4
Commutative triassociative algebras , Definition 3.8	Lie bracket $[a, b] = a \wedge b - b \wedge a$ Pre-Lie product $\{a, b\} = a < b - b > a$ Post-Lie algebras , Definition 3.8
Jordan product $a \circ b = a \perp b + b \perp a$ Jordan diproduct $a \bullet b = a \dashv b + b \vdash a$ Jordan trialgebras , Section 4	No dual, operad is cubic not quadratic
No dual, operad is cubic not quadratic	Jordan product $a \circ b = a \wedge b + b \wedge a$ Pre-Jordan product $a \bullet b = a < b + b > a$ Post-Jordan algebras , Section 6

TABLE 1. Operadic generalizations of associative, Lie, and Jordan algebras

1.2. Overview of problems and methods. Table 1 displays the generalizations of associativity underlying our results; the number of operations increases from top to bottom. The left column contains operads obtained from associative operations (di- and triassociative); the right column contains their Koszul duals (dendriform and tridendriform). These operads are nonsymmetric (but will be symmetrized); each of them leads to generalizations of Lie and Jordan algebras, defined by the polynomial identities of arity $\leq 3, 4$ satisfied by the analogues of the (anti)commutator and (anti)dicommutator. All the operads we consider are generated by binary operations, with or without symmetry.

1.2.1. *The symmetric operads ΣTriAss , $\Sigma\text{TriDend}$ and BW .* We are primarily concerned with three operads; the first two form a Koszul dual pair:

- The symmetric triassociative operad ΣTriAss (the symmetrization of TriAss) which is generated by three binary operations \dashv, \perp, \vdash with no symmetry, satisfying seven quadratic relations; see Definition 3.1.
- The symmetric tridendriform operad $\Sigma\text{TriDend}$ (the symmetrization of TriDend) which is generated by three binary operations \langle, \wedge, \rangle with no symmetry, satisfying 11 quadratic relations; see Definition 3.1.
- The free symmetric operad BW generated by two binary operations, one commutative \circ and one noncommutative (with no symmetry) \bullet . The operads governing Jordan trialgebras and post-Jordan algebras are quotients of BW .
- The expansion map $E(n): \text{BW}(n) \rightarrow \Sigma\text{TriAss}(n)$ is defined by $a \circ b \mapsto a \perp b + b \perp a$ and $a \bullet b \mapsto a \dashv b + b \vdash a$; its kernel contains the defining identities for Jordan trialgebras.
- The expansion map $E(n): \text{BW}(n) \rightarrow \Sigma\text{TriDend}(n)$ is defined by $a \circ b \mapsto a \wedge b + b \wedge a$ and $a \bullet b \mapsto a \langle b + b \rangle$; its kernel contains the defining identities for post-Jordan algebras.

The standard monograph on the theory of algebraic operads is Loday & Vallette [44].

1.2.2. *Computational methods.* All computations are performed with Maple using arithmetic over \mathbb{Z} or \mathbb{Q} or a finite field. To save time and memory we usually work over a finite prime field \mathbb{F}_p , where p is greater than the arity n of the identities being considered; this guarantees that the group algebra $\mathbb{F}_p S_n$ is semisimple. In arity n , both the domain and the codomain of the expansion map are S_n -modules, and so if $p > n$ then semisimplicity of $\mathbb{F}_p S_n$ guarantees that it is isomorphic to the direct sum of simple two-sided ideals each of them isomorphic to a full matrix algebra. The coefficients in the formulas for the matrix units in $\mathbb{F}_p S_n$ as linear combinations of permutations have denominators which are divisors of $n!$ and hence are well-defined modulo p . We are left with the problem of reconstructing correct rational results from modular calculations, but in this respect we had very good luck: all matrix entries are in $\{0, \pm 1\}$. Using the representation theory of the symmetric group allows us to decompose a large matrix into much smaller pieces.

1.2.3. *Terminology and notation.* The number of arguments in a monomial is often called its *degree*; here we use *arity*: for algebraic operads, *degree* refers to (homological) degree in a differential graded vector space. From the homological point of view, all our vector spaces are graded but concentrated in degree 0 over a base field \mathbb{F} of characteristic 0, unless otherwise stated. We write $\mathbf{Vect}_{\mathbb{F}}$ for the category of vector spaces over \mathbb{F} . If X is a set then $\mathbb{F}X$ is the vector space² with basis X . If \mathcal{O} is a nonsymmetric operad then $\Sigma\mathcal{O}$ denotes its symmetrization.

1.3. **Associative, Lie, and Jordan algebras.** An associative algebra is a vector space A with a bilinear product $m: A \times A \rightarrow A$, $(a, b) \mapsto ab$, satisfying the relation $(ab)c - a(bc) \equiv 0$ which is nonsymmetric (hence multilinear): every term has the identity permutation of the arguments. The symbol \equiv indicates that equality holds for all values of the arguments.

²From a categorical perspective, the functor $X \mapsto \mathbb{F}X$ is left adjoint to the forgetful functor $\mathbf{Vect}_{\mathbb{F}} \rightarrow \mathbf{Sets}$; that is, $\mathbb{F}X$ is the free vector space over \mathbb{F} generated by X .

In every associative algebra, the commutator (Lie bracket) $[a, b] = ab - ba$ is antisymmetric and satisfies the Jacobi identity; these properties define Lie algebras:

$$(1) \quad [a, a] \equiv 0, \quad [[a, b], c] + [[b, c], a] + [[c, a], b] \equiv 0.$$

In every associative algebra, the anticommutator (Jordan product) $a \circ b = ab + ba$ is symmetric and satisfies the Jordan identity; these properties define Jordan algebras:

$$(2) \quad a \circ b \equiv b \circ a, \quad ((a \circ a) \circ b) \circ a \equiv (a \circ a) \circ (b \circ a).$$

Let L be a Lie algebra with bracket $[-, -]$, let J be a Jordan algebra with product $a \circ b$, and let X be either L or J . Then X has a universal associative enveloping algebra $U(X)$ in the following sense: if $f: X \rightarrow A$ is a linear map to an associative algebra A then there is a unique algebra morphism $g: U(X) \rightarrow A$ such that $g \circ h = f$ where $h: X \rightarrow U(X)$ is the natural map arising as follows. If $T(V)$ is the tensor algebra with product $a \cdot b$ of the vector space V , then $U(X) \cong T(X)/I(X)$ for the following (two-sided) ideals:

$$I(L) = \langle a \cdot b - b \cdot a - [a, b] \mid a, b \in L \rangle, \quad I(J) = \langle a \cdot b + b \cdot a - a \circ b \mid a, b \in J \rangle.$$

For every Lie algebra L , h is injective; this follows from the Poincaré-Birkhoff-Witt theorem and implies that every polynomial identity satisfied by the commutator in every associative algebra is a consequence of skewsymmetry and the Jacobi identity. The same does not hold for Jordan algebras. A Jordan algebra J is *special* if it is isomorphic to a subalgebra of A^+ for some associative algebra A , where A^+ is the same vector space with the nonassociative product $a \circ b = ab + ba$. For a Jordan algebra J , the natural map $h: J \rightarrow U(J)$ is injective if and only if J is special. There are polynomial identities satisfied by every special Jordan algebra which are not consequences of symmetry and the Jordan identity [28]; the smallest examples are the so-called Glennie identities and occur in arity 8. Quotients of special Jordan algebras are not necessarily special [20], so special Jordan algebras do not form a variety defined by polynomial identities.




FIGURE 1. Generalizations of the Lie, associative, and Jordan operads

1.4. Results of this paper in context. In later sections, we include concise reviews of analogous structures existing in the literature, and their interrelations, in order to identify the gaps which motivated the writing of this paper. Table 1 shows the algebraic operads with which we are concerned. Our results provide definitions of Jordan trialgebras and post-Jordan algebras which are indicated by boxes. (A somewhat similar diagram appears in the lecture notes [35].) The dotted straight

(resp. curved) up-down arrows indicate the white (resp. black) Manin products with the operads Perm and PreLie (resp. ComTriAss and PostLie); see §2.6. These procedures may also be described as duplicators and disuccessors (resp. triplicators and trisuccessors); see §4.4, §6.3. Reversing an arrow corresponds to realizing a simpler structure as a subalgebra or quotient of a more complex structure. The left-right arrows represent morphisms between operads which replace the (generalized) associative product by one or two (generalized) Lie brackets or Jordan products. We use the symbols $[a, b]$ and/or $\{a, b\}$ for the Lie case, $a \circ b$ and/or $a \bullet b$ for Jordan. Reversing a left-right arrow represents constructing the universal enveloping algebra of the corresponding Lie or Jordan structure. All these operads are binary: the generating operations are bilinear. In columns 1, 2 the operads are quadratic: the terms of the relations contain two operations. In column 3, the operads are cubic: the terms contain three operations. Koszul duality for quadratic operads can be applied to columns 1, 2 but not 3.

2. PRELIMINARIES ON ALGEBRAIC OPERADS

2.1. Free nonsymmetric binary operads. Let $\Omega = \{\circ_1, \dots, \circ_m\}$ be a finite ordered set of binary operations; let O_m be the free nonsymmetric operad in $\mathbf{Vect}_{\mathbb{F}}$ generated by $\mathbb{F}\Omega$. The underlying vector space of O_m is the direct sum over $n \geq 1$ of homogeneous components $O_m(n)$ of arity n . We never add elements of different arities, so we could replace the direct sum by a disjoint union (but then we would leave $\mathbf{Vect}_{\mathbb{F}}$). The following construction of O_m also makes sense in the category \mathbf{Set} of sets. We define sets X of operations of arity n and then consider the vector spaces $\mathbb{F}X$. Disjoint union (resp. direct product) of sets corresponds to direct sum (resp. tensor product) of vector spaces. The operad O_m in \mathbf{Set} may be identified with the free magma on one generator with m binary operations.

Definition 2.1. For $n \geq 1$, let $P_n = \{p_1, \dots, p_{c(n)}\}$ be the **association types** of arity n : the distinct placements of balanced pairs of parentheses for a single binary operation in a sequence of n arguments. If we omit the outermost pair then for $n \geq 2$ there are $n-2$ pairs. The size of P_n is the Catalan number $c(n) = (2n-2)!/(n!(n-1)!)$. We write $P = \bigcup_{n \geq 1} P_n$; the set P is a basis for \mathcal{O}_1 , the free nonsymmetric operad on one binary operation.

Example 2.2. We write x for the arguments and juxtaposition for the operation. For $n = 1$ we have only x , for $n = 2$ only xx , and for $n = 3$ only $(xx)x, x(xx)$. For $n = 4, 5$ we have:

$$\begin{aligned} n = 4 & \quad ((xx)x)x, (x(xx))x, (xx)(xx), x((xx)x), x(x(xx)). \\ n = 5 & \quad (((xx)x)x)x, ((x(xx))x)x, ((xx)(xx))x, (x((xx)x))x, (x(x(xx)))x, \\ & \quad ((xx)x)(xx), (x(xx))(xx), (xx)((xx)x), (xx)(x(xx)), \\ & \quad x(((xx)x)x), x((x(xx))x), x((xx)(xx)), x(x((xx)x)), x(x(x(xx))). \end{aligned}$$

Definition 2.3. The **revdeglex** order $<$ on P is defined as follows. If $f \in P_n, f' \in P_{n'}$, $n < n'$ then $f < f'$. If $f, f' \in P_n$ then we proceed by induction. If $n \leq 2$ then $|P_n| = 1$ and $f = f'$. If $n \geq 3$ then there are unique factorizations $f = gh, f' = g'h'$; we set $f < f'$ if and only if either $h < h'$, or $h = h', g < g'$. (This order was used in Example 2.2.)

If $m \geq 2$ then we must distinguish the m operation symbols in Ω . An element of P_n contains $n-1$ multiplications ordered from left to right, and each may be

replaced by any element of Ω . We identify the m^{n-1} possibilities with Ω^{n-1} whose lex order $<_{\Omega^{n-1}}$ is induced by Ω : if $\alpha, \beta \in \Omega^{n-1}$ then $\alpha <_{\Omega^{n-1}} \beta$ if and only if $\alpha_i <_{\Omega} \beta_i$ where i is minimal for $\alpha_i \neq \beta_i$.

Definition 2.4. Operations are independent of association types, and so we identify a basis of $O_m(n)$ with $P_n^{(m)} = P_n \times \Omega^{n-1}$. The elements of $P_n^{(m)}$ are the **skeletons** of arity n ; each skeleton is an n -tuple $(p_i; \circ_{j_1}, \dots, \circ_{j_{n-1}})$ which represents the basis element of $O_m(n)$ with association type p_i and operations $\circ_{j_1}, \dots, \circ_{j_{n-1}}$ from left to right. The total orders on P_n and Ω^{n-1} extend to $P_n^{(m)}$: for $p, p' \in P_n$ and $\alpha, \alpha' \in \Omega^{n-1}$ we have $(p, \alpha) < (p', \alpha')$ if and only if either $p <_P p'$, or $p = p', \alpha <_{\Omega^{n-1}} \alpha'$.

Definition 2.5. To avoid confusion with the operations in Ω , we denote the **substitution maps** in $O_m(n)$ by an asterisk instead of a circle, $*_i: O_m(n) \otimes O_m(n') \rightarrow O_m(n+n'-1)$ for $1 \leq i \leq n$. For skeletons $f \in P_n^{(m)} \subseteq O_m(n)$ and $f' \in P_{n'}^{(m)} \subseteq O_m(n')$, the skeleton

$$f *_i f' \in P_{n+n'-1}^{(m)} \subseteq O_m(n+n'-1),$$

is obtained by substituting f' for the i -th argument of f . These definitions extend bilinearly to O_m , and the resulting substitutions generate all compositions in O_m .

Definition 2.6. We may now state precisely that the **free nonsymmetric operad** with m binary operations Ω with no symmetry is the direct sum O_m of the finite dimensional vector spaces $O_m(n)$ with the compositions generated by the substitutions $*_i$ of Definition 2.5.

2.2. Free symmetric binary operads. The free nonsymmetric operad O_m has a basis of skeletons containing parentheses and operation symbols; arguments are denoted by x .

Definition 2.7. A **multilinear monomial** of arity n is a skeleton $s \in P_n^{(m)}$ in which the argument symbols have been replaced by a permutation of x_1, \dots, x_n . We identify the set of such monomials with $W_n^{(m)} = P_n^{(m)} \times S_n$ where S_n is the symmetric group, and define $\Sigma O_m(n)$ to be the vector space with basis $W_n^{(m)}$ on which there is a right S_n -action. The direct sum ΣO_m of the homogeneous components $\Sigma O_m(n)$ is the **symmetrization** of O_m . In fact, ΣO_m is the (underlying vector space of the) **free symmetric operad** on m binary operations with no symmetry. Definition 2.5 shows how to compose skeletons, but in ΣO_m we also need to compose monomials: if α, α' are monomials of arities n, n' with arguments x_1, \dots, x_n and $x_1, \dots, x_{n'}$, then $\alpha *_i \alpha'$ ($1 \leq i \leq n$) is a monomial of arity $n+n'-1$ with arguments $x_1, \dots, x_{n+n'-1}$. For further information, see §5.3.2 of [44]. A **regular operad** is a symmetric operad (free or not) which is the symmetrization of a nonsymmetric operad.

2.3. Identical relations and operad ideals. Let P be a free symmetric operad generated by $P(2)$; that is, binary operations with or without symmetry. The symmetries of the operations are determined by the S_2 -module structure of $P(2)$, which is the direct sum of S_2 -modules isomorphic to either the unit module $[+]$ (representing commutativity), the sign module $[-]$ (anticommutativity), or the group algebra $\mathbb{F}S_n$ (no symmetry). (Since $\mathbb{F}S_n \cong [+] \oplus [-]$, an operation with no symmetry can be polarized into two operations, one commutative and one anticommutative, but this will not concern us.) We impose an S_3 -module of quadratic relations $R \subseteq P(3)$ on the operations generating P .

Definition 2.8. The **operad ideal** $I = \langle R \rangle$ generated by the relations R is the smallest sum $\bigoplus_{n \geq 1} I(n)$ of S_n -submodules $I(n) \subseteq P(n)$ which is closed under arbitrary compositions with elements of P . That is, if $f \in I(r)$ and $g \in P(s)$ then $f *_i g \in I(r+s-1)$ for $1 \leq i \leq r$, and $g *_j f \in I(r+s-1)$ for $1 \leq j \leq s$. The **quotient operad** P/I has homogeneous components $P(n)/I(n)$ with compositions defined in the natural way.

For given relations R , we construct a set of S_n -module generators of $I(n)$ for all n . The S_3 -module $R = I(3)$ is generated by a finite set of relations. Assume that we have already constructed a finite set of S_n -module generators of $I(n)$. If f is such a generator then we increment the arity by composing $f(x_1, \dots, x_n)$ with some $g(x_1, x_2) \in P(2)$. By multilinearity, we may assume that $g(x_1, x_2) = x_1 \circ x_2$ where \circ is one of the generating operations. There are n possibilities for $f *_i g$ and two for $g *_j f$; since every relation must be multilinear, we change subscripts of some arguments:

$$(3) \quad \begin{cases} f *_i g = f(x_1, \dots, x_n) *_i g(x_1, x_2) = f(x_1, \dots, x_{i-1}, x_i \circ x_{n+1}, x_{i+1}, \dots, x_n) \\ g *_j f = g(x_1, x_2) *_j f(x_1, \dots, x_n) = \begin{cases} f(x_1, \dots, x_n) \circ x_{n+1} & (j = 1) \\ x_{n+1} \circ f(x_1, \dots, x_n) & (j = 2) \end{cases} \end{cases}$$

As f runs over the set of S_n -module generators of $I(n)$, the elements (3) form a set of S_{n+1} -module generators for $I(n+1)$.

2.4. Koszul duality for quadratic operads. For any category of algebras governed by a quadratic operad, one can construct its Koszul dual operad, which governs another category of algebras. For associative algebras, the nonsymmetric operad is self-dual: $\text{Assoc}^! \cong \text{Assoc}$. The symmetric operads $\text{ComAss}^!$ for commutative associative algebras and Lie for Lie algebras form a Koszul dual pair: $\text{ComAss}^! = \text{Lie}$, $\text{Lie}^! = \text{ComAss}$. (If P is a quadratic operad then $(P^!)^! \cong P$.) If we generalize associativity to more than one binary operation (diassociative, triassociative, ...), the quadratic operads are not self-dual, and define further generalizations of associativity (dendriform, tridendriform, ...).

Loday showed that for an operad (symmetric or nonsymmetric) generated by binary operations with no symmetry, the relations for the Koszul dual can be obtained from the original relations by means of elementary linear algebra; see Theorem 8.5 and Proposition B.3 of [40] and Table 2 of [14]. In this paper we give two examples, one of which extends the algorithm to operations with symmetry; see Lemmas 3.3 and 3.5.

2.5. Diassociative and dendriform algebras.

Definition 2.9. [38], [42], [40]. A vector space with bilinear operations \dashv, \vdash is a **diassociative algebra** if it satisfies these relations:

- left and right associativity: $(a \dashv b) \dashv c \equiv a \dashv (b \dashv c)$, $(a \vdash b) \vdash c \equiv a \vdash (b \vdash c)$,
- inner associativity: $(a \vdash b) \dashv c \equiv a \vdash (b \dashv c)$,
- left and right bar identities: $(a \dashv b) \vdash c \equiv (a \vdash b) \dashv c$, $a \dashv (b \vdash c) \equiv a \vdash (b \dashv c)$.

A vector space with bilinear operations \langle, \rangle is a **dendriform algebra** if it satisfies:

- inner associativity: $(a \rangle b) \langle c \equiv a \rangle (b \langle c)$,
- left-right symmetrization: $(a \langle b) \langle c \equiv a \langle (b \langle c) + a \langle (b \rangle c)$,
- right-left symmetrization: $a \rangle (b \rangle c) \equiv (a \rangle b) \rangle c + (a \langle b) \rangle c$.

Remark 2.10. The diassociative (but not dendriform) relations have the form $m_1 \equiv m_2$ for monomials m_1, m_2 . Dendriform algebras are related to Rota-Baxter operators [1], [24].

Lemma 2.11. [42], [40]. *The operation $a \cdot b = a < b + a > b$ is associative in every dendriform algebra. The operads DiAss (ΣDiAss) and Dend (ΣDend) are Koszul duals.*

Definition 2.12. [6], [37], [39]. A vector space with a bilinear operation $\{a, b\}$ (resp. $a < b$) is a (left) **Leibniz** (resp. **Zinbiel**) algebra if it satisfies the relation

$$\{a, \{b, c\}\} \equiv \{\{a, b\}, c\} + \{b, \{a, c\}\} \quad (\text{resp. } (a < b) < c \equiv a < (b < c) + a < (c < b)).$$

Lemma 2.13. [38, 39, 40]. *The operation $a \dashv b - b \vdash a$ satisfies the left Leibniz identity in every diassociative algebra. In every Zinbiel algebra the anticommutator $a < b + b < a$ is commutative and associative. Every Zinbiel algebra becomes a dendriform algebra if we define $a > b = b < a$; conversely, every dendriform algebra which is commutative ($a < b \equiv b > a$) is a Zinbiel algebra. The operads Leib and Zinb are Koszul duals.*

Remark 2.14. The operation $\{a, b\} = a \dashv b - b \vdash a$ defines a functor between the categories of diassociative and Leibniz algebras; the left adjoint sends a Leibniz algebra to its universal enveloping diassociative algebra; see [2], [7], [29], [40].

Problem 2.15. Determine the polynomial identities satisfied by (1) the dicommutator and anticommutator in Leibniz algebras; (2) the dicommutator in Zinbiel algebras.

Definition 2.16. [16], [18], [26], [53]. A vector space with a bilinear operation $\{a, b\}$ is a (left) **pre-Lie** (or left-symmetric) algebra if it satisfies the relation $(a, b, c) \equiv (b, a, c)$ where the associator is $(a, b, c) = \{\{a, b\}, c\} - \{a, \{b, c\}\}$. A (right) **perm algebra** is associative and **right-commutative**: $abc \equiv acb$.

Lemma 2.17. [5], [17], [38], [40]. *In every dendriform algebra the operation $a < b - b > a$ satisfies the pre-Lie identity. The commutator in a pre-Lie algebra satisfies (anticommutativity and) the Jacobi identity. Every identity for the anticommutator in a pre-Lie algebra is a consequence of commutativity. The operads PreLie and Perm are Koszul duals.*

Definition 2.18. [9], [12], [33], [34], [52]. A (right) **Jordan dialgebra** is a vector space with a bilinear product $a \bullet b$ satisfying the following three relations (one of arity 3 and two of arity 4) where $(a, b, c) = (a \bullet b) \bullet c - a \bullet (b \bullet c)$ is the associator:

$$\begin{array}{ll} \text{right commutativity} & a \bullet (b \bullet c) \equiv a \bullet (c \bullet b) \\ \text{right quasi-Jordan identity} & (b \bullet a) \bullet (a \bullet a) \equiv (b \bullet (a \bullet a)) \bullet a \\ \text{right associator-derivation identity} & (b, a \bullet a, c) \equiv 2(b, a, c) \bullet a \end{array}$$

A (right) **pre-Jordan algebra** is a vector space with a bilinear product $a \bullet b$ satisfying the following two relations of arity 4 where we write for short $a \cdot b = a \bullet b + b \bullet a$:

$$\begin{aligned} & (a \cdot b) \bullet (c \cdot d) + (b \cdot c) \bullet (a \cdot d) + (c \cdot a) \bullet (b \cdot d) \\ & \equiv c \bullet [(a \cdot b) \bullet d] + a \bullet [(b \cdot c) \bullet d] + b \bullet [(c \cdot a) \bullet d], \\ & a \bullet [b \bullet (c \cdot d)] + c \bullet [b \bullet (a \cdot d)] + [(a \cdot c) \cdot b] \bullet d \\ & \equiv c \bullet [(a \cdot b) \bullet d] + a \bullet [(b \cdot c) \bullet d] + b \bullet [(c \cdot a) \bullet d]. \end{aligned}$$

Lemma 2.19. [9], [12], [33]. *In every diassociative algebra the operation $a \vdash b + b \dashv a$ satisfies the identities defining Jordan dialgebras. In every dendriform algebra the operation $a < b + b > a$ satisfies the identities defining pre-Jordan algebras.*

2.6. Manin black and white products. We can interpret the vertical arrows in Table 1 using Manin white and black products of operads; for details see [4], [27], [31], [44], [51].

Starting with ΣAssoc in the center of Table 1, we compute the white product $\Sigma\text{DiAss} \cong \text{Perm} \circ \Sigma\text{Assoc}$ to move up in column 2. Similarly, we obtain the Leibniz operad $\text{Leib} \cong \text{Perm} \circ \text{Lie}$ from the operad Lie , and the di-Jordan operad $\text{DiJor} \cong \text{Perm} \circ \text{Jor}$ from the Jordan operad Jor . Taking the white product with Perm is sometimes called duplication [30], [47]; this process has also been called the KP algorithm [11]. Starting again from ΣAssoc , we move down by computing the black product $\Sigma\text{Dend} \cong \text{PreLie} \bullet \Sigma\text{Assoc}$. Similarly, we obtain $\text{PreLie} \cong \text{PreLie} \bullet \text{Lie}$ and $\text{PreJor} \cong \text{PreLie} \circ \text{Jor}$. Taking the black product with PreLie is sometimes called computing the disuccessor [3].

For any finitely generated binary quadratic operads P and Q , Koszul duality interchanges Manin black and white products: we have $(P \bullet Q)^! \cong P^! \circ Q^!$. Therefore, DiAss and Dend are Koszul dual:

$$\Sigma\text{DiAss}^! \cong (\Sigma\text{Assoc} \circ \text{Perm})^! \cong \Sigma\text{Assoc}^! \bullet \text{Perm}^! \cong \Sigma\text{Assoc} \bullet \text{PreLie} \cong \Sigma\text{Dend}.$$

For the Lie column, the operads Leib and PreLie are not a dual pair, but we have

$$\begin{aligned} \text{Leib} \cong \text{Perm} \circ \text{Lie} &= \text{PreLie}^! \circ \text{ComAss}^! \cong (\text{PreLie} \bullet \text{ComAss})^! = \text{Zinb}^!, \\ \text{PreLie} \cong \text{PreLie} \bullet \text{Lie} &\cong \text{Perm}^! \bullet \text{ComAss}^! \cong (\text{Perm} \circ \text{ComAss})^! \cong \text{Perm}^!. \end{aligned}$$

For the Jordan column, *if the Jordan operad had a Koszul dual*, then we would obtain:

$$\begin{aligned} \text{DiJor} \cong \text{Perm} \circ \text{Jor} &= \text{PreLie}^! \circ (\text{Jor}^!)^! \cong (\text{PreLie} \bullet \text{Jor}^!)^!, \\ \text{PreJor} \cong \text{PreLie} \bullet \text{Jor} &\cong \text{Perm}^! \bullet (\text{Jor}^!)^! \cong (\text{Perm} \circ \text{Jor}^!)^!. \end{aligned}$$

See §7.1 for one way in which a Koszul dual for the Jordan operad could be defined.

Similar considerations apply to the top and bottom rows of Table 1. Going from the middle to the top corresponds to taking the triplicator of the operad [30], [47]; this is the white product with the operad ComTriAss (Definition 3.1):

$$\text{TriLie} \cong \text{ComTriAss} \circ \text{Lie}, \quad \Sigma\text{TriAss} \cong \text{ComTriAss} \circ \Sigma\text{Assoc}, \quad \text{TriJor} \cong \text{ComTriAss} \circ \text{Jor}.$$

Going from the middle to the bottom represents taking the trisuccessor of the operad [3]: the black product with the operad PostLie (Definition 3.8). The operads ΣTriAss and $\Sigma\text{TriDend}$ at top and bottom of the associative column are Koszul dual. For the Lie column,

$$\begin{aligned} \text{TriLie} \cong \text{ComTriAss} \circ \text{Lie} &= \text{PostLie}^! \circ \text{ComAss}^! \cong (\text{PostLie} \bullet \text{ComAss})^! \cong \text{ComTriDend}^!, \\ \text{PostLie} \cong \text{PostLie} \bullet \text{Lie} &\cong \text{ComTriAss}^! \bullet \text{ComAss}^! \cong (\text{ComTriAss} \circ \text{ComAss})^! \cong \text{ComTriAss}^!. \end{aligned}$$

3. TRIASSOCIATIVE AND TRIDENDRIFORM ALGEBRAS

Definition 3.1. [43]. A **triassociative algebra** is a vector space with bilinear operations \dashv, \perp, \vdash satisfying these relations where $*$ $\in \{\dashv, \perp, \vdash\}$:

- left, middle, and right associativity: $(a * b) * c \equiv a * (b * c)$,
- bar identities: $a \dashv (b \dashv c) \equiv a \dashv (b * c)$ and $(a \vdash b) \vdash c \equiv (a * b) \vdash c$,
- inner associativity: if $(*_1, *_2) \in \{(\vdash, \dashv), (\vdash, \perp), (\perp, \dashv)\}$ then $(a *_1 b) *_2 c \equiv a *_1 (b *_2 c)$,
- Loday-Ronco relation: $(a \dashv b) \perp c \equiv a \perp (b \vdash c)$.

A **tridendriform algebra** is a vector space with bilinear operations \langle, \wedge, \rangle satisfying:

- middle associativity: $(a \wedge b) \wedge c \equiv a \wedge (b \wedge c)$,

some permutation of the arguments. The (nonzero) elements of the kernel of $E(n)$ are the (multilinear) polynomial identities of arity n satisfied by the operations \circ, \bullet in every triassociative algebra.

4.1. Relations of arity 3.

Lemma 4.2. *Over a field \mathbb{F} of characteristic 0 or $p > 3$, every element of the kernel of $E(3)$ is a consequence of the commutativity of \circ and the relation $a \bullet (b \circ c) \equiv a \bullet (b \bullet c)$.*

Definition 4.3. The relation of arity 3 in Lemma 4.2 is the **black right bar identity**.

Proof. (of Lemma 4.2). We first find monomial bases for $\mathbb{B}\mathbb{W}(3)$ and $\Sigma\text{TriAss}(3)$ so that we can represent $E(3)$ by a matrix and compute a basis for its nullspace.

Since TriAss is defined by monomial relations, we find a basis for $\text{TriAss}(3)$ as follows. The triassociative relations generate an equivalence relation \sim on the set B of 18 (nonsymmetric) monomials (4): we define $t_1 \sim t_2$ if and only if $t_1 \equiv t_2$ is one of the relations in Definition 3.1. We take the quotient of B by \sim to obtain a set partition B/\sim . In each equivalence class we choose, as the normal form of the elements in the class, the monomial which is minimal with respect to the total order in the proof of Lemma 3.3; we use these representatives as a basis of $\text{TriAss}(3)$. We obtain the following normal forms; the first two classes each contain four monomials, and the last five each contain two:

$$(a \vdash b) \vdash c, (a \vdash b) \vdash c, (a \perp b) \perp c, (a \vdash b) \vdash c, (a \perp b) \vdash c, (a \vdash b) \perp c, (a \vdash b) \perp c.$$

To symmetrize, we apply all 6 permutations of a, b, c to the arguments of the normal forms, and obtain a basis of 42 monomials for $\Sigma\text{TriAss}(3)$.

Since $\mathbb{B}\mathbb{W}$ is a symmetric operad, to find an ordered basis for $\mathbb{B}\mathbb{W}(3)$, we take the 27 monomials (5) from the proof of Lemma 3.5 and replace the symbols \prec, \wedge by \bullet, \circ .

The columns of the 42×27 matrix representing $E(3)$ with these ordered bases are the coefficient vectors of the expansions of the basis monomials of $\mathbb{B}\mathbb{W}(3)$ into $\Sigma\text{TriAss}(3)$; we must replace the triassociative monomials by their normal forms. For example,

$$\begin{aligned} (a \circ b) \bullet c &\mapsto (a \perp b + b \perp a) \vdash c + c \vdash (a \perp b + b \perp a) && \text{expansion map} \\ &= (a \perp b) \vdash c + (b \perp a) \vdash c + c \vdash (a \perp b) + c \vdash (b \perp a) && \text{bilinearity} \\ &= (a \perp b) \vdash c + (b \perp a) \vdash c + (c \vdash a) \perp b + (c \vdash b) \perp a && \text{normal form} \end{aligned}$$

The matrix has rank 21 and nullity 6. We find a basis for the nullspace and write down the corresponding identities, which are the 6 permutations of the black right bar identity. \square

Remark 4.4. The black right bar identity and the commutativity of \circ imply right commutativity for \bullet (Definition 2.18): we have $a \bullet (b \bullet c) \equiv a \bullet (b \circ c) \equiv a \bullet (c \circ b) \equiv a \bullet (c \bullet b)$.

4.2. The operad $\mathbb{B}\mathbb{W}$: skeletons, total order, normal forms. Before studying arities $n \geq 4$, we need to understand skeletons in $\mathbb{B}\mathbb{W}$; as before, a skeleton in arity n is a placement of parentheses and a choice of operation symbols in a sequence of n indistinguishable arguments. For the operad $\mathbb{B}\mathbb{W}$, their enumeration is not trivial due to the commutativity of \circ .

Definition 4.5. It is convenient to consider $\mathbb{B}\mathbb{W}$ not in the monoidal category $\mathbf{Vect}_{\mathbb{F}}$ with tensor product, but in \mathbf{Set} with direct product; we call the latter operad $\mathbf{BW}\text{-Set}$. (Equivalently, we consider a basis of each $\mathbb{B}\mathbb{W}(n)$ rather than the entire vector

space.) Since BW-Set is a symmetric operad, we cannot use the symbol x for every argument. But we achieve the same goal by considering the free algebra BWS over BW-Set generated by x ; this algebra consists of all BW-skeletons . We write $\text{BWS}(n)$ for the subset of BWS consisting of the elements of arity n ; that is, those in which x occurs exactly n times. This device allows us to use the symbol x as a placeholder for unspecified arguments in a symmetric operad.

Definition 4.6. For $n \geq 1$, we inductively define a **total order** \ll_n on $\text{BWS}(n)$:

- (1) For $n = 1$, we have $\text{BWS}(1) = \{x\}$. In what follows we omit the subscript on \ll_n .
- (2) For $n \geq 2$, if $h \in \text{BWS}(n)$ then $h = f \circ g$ or $h = f \bullet g$ with $1 \leq \text{arity}(f), \text{arity}(g) < n$.
 - (a) We set $f_1 \circ g_1 \ll f_2 \bullet g_2$ for all $f_1 \circ g_1, f_2 \bullet g_2 \in \text{BWS}(n)$.
 - (b) Consider $f_1 \bullet g_1$ and $f_2 \bullet g_2$. If $\text{arity}(f_1) > \text{arity}(f_2)$, or $\text{arity}(f_1) = \text{arity}(f_2)$ and $f_1 \ll f_2$, or $f_1 = f_2$ and $g_1 \ll g_2$, then we set $f_1 \bullet g_1 \ll f_2 \bullet g_2$.
 - (c) Consider $f_1 \circ g_1$ and $f_2 \circ g_2$. We may assume $\text{arity}(f_i) \geq \text{arity}(g_i)$ for $i = 1, 2$.
 - (i) If $\text{arity}(f_1) > \text{arity}(f_2)$ then we set $f_1 \circ g_1 \ll f_2 \circ g_2$.
 - (ii) If $\text{arity}(f_1) = \text{arity}(f_2)$ then we also have $\text{arity}(g_1) = \text{arity}(g_2)$.
 - Assume $\text{arity}(f_1) > \text{arity}(g_1)$. If $f_1 \ll f_2$, or $f_1 = f_2$ and $g_1 \ll g_2$, then we set $f_1 \circ g_1 \ll f_2 \circ g_2$.
 - Assume $\text{arity}(f_1) = \text{arity}(g_1)$. In this case, n is even and all four factors have arity $n/2$. By commutativity, $f_i \ll g_i$ (or $f_i = g_i$) for $i = 1, 2$. If $f_1 \ll f_2$, or $f_1 = f_2$ and $g_1 \ll g_2$, then we set $f_1 \circ g_1 \ll f_2 \circ g_2$.

Example 4.7. The 25 skeletons in $\text{BWS}(4)$ in the order of Definition 4.6 are as follows:

$$\begin{aligned} & ((x \circ x) \circ x) \circ x, ((x \bullet x) \circ x) \circ x, ((x \circ x) \bullet x) \circ x, ((x \bullet x) \bullet x) \circ x, (x \bullet (x \circ x)) \circ x, \\ & (x \bullet (x \bullet x)) \circ x, (x \circ x) \circ (x \circ x), (x \circ x) \circ (x \bullet x), (x \bullet x) \circ (x \bullet x), ((x \circ x) \circ x) \bullet x, \\ & ((x \bullet x) \circ x) \bullet x, ((x \circ x) \bullet x) \bullet x, ((x \bullet x) \bullet x) \bullet x, (x \bullet (x \circ x)) \bullet x, (x \bullet (x \bullet x)) \bullet x, \\ & (x \circ x) \bullet (x \circ x), (x \circ x) \bullet (x \bullet x), (x \bullet x) \bullet (x \circ x), (x \bullet x) \bullet (x \bullet x), x \bullet ((x \circ x) \circ x), \\ & x \bullet ((x \bullet x) \circ x), x \bullet ((x \circ x) \bullet x), x \bullet ((x \bullet x) \bullet x), x \bullet (x \bullet (x \circ x)), x \bullet (x \bullet (x \bullet x)). \end{aligned}$$

4.2.1. *Computing the normal form of a multilinear BW monomial.* We apply commutativity of \circ to straighten first the skeleton and then the arguments; for example, $c \circ (b \circ a)$ becomes $(b \circ a) \circ c$ and then $(a \circ b) \circ c$. Straightening the skeleton is equivalent to choosing a unique representative of each equivalence class in $\text{BWS}(n)$ with respect to commutativity: the skeleton $x \circ (x \circ x)$ has representative $(x \circ x) \circ x$. This algorithm is necessary when we compute the consequences in arity $n+1$ of a BW polynomial in arity n : applying the substitution maps of Definition 2.5 to the set of representatives of equivalence classes in $\text{BWS}(n)$ does not necessarily produce representatives of equivalence classes in $\text{BWS}(n+1)$.

Algorithm 4.8. For $n = 1$ there is only one multilinear monomial x_1 which by definition is already in normal form. For $n \geq 2$, every multilinear monomial of arity n has the form $f \circ g$ or $f \bullet g$ (we write $f * g$ to cover both cases) where $1 \leq \text{arity}(f), \text{arity}(g) < n$:

- (1) Recursively compute f' and g' , the normal forms of f and g .
- (2) If $\text{arity}(f') > \text{arity}(g')$ then return $f' * g'$.
- (3) If $\text{arity}(f') < \text{arity}(g')$ then: if $* = \circ$ then return $g' * f'$ else return $f' * g'$.
- (4) If $\text{arity}(f') = \text{arity}(g')$ then:
 - (a) Extract $s(f')$ and $s(g')$, the skeletons of f' and g' .
 - (b) If $s(f') \ll s(g')$ in the total order on $\text{BWS}(n/2)$ then return $f' * g'$.
 - (c) If $s(f') \gg s(g')$ then: if $* = \circ$ then return $g' * f'$ else return $f' * g'$.

- (d) If $s(g') = s(f')$ then:
- (i) Extract $p(f')$ and $p(g')$, the sequences of subscripts of the arguments of f' and g' ; as sets, $p(f') \cap p(g') = \emptyset$ and $p(f') \cup p(g') = \{1, \dots, n\}$.
 - (ii) If $p(f')$ precedes $p(g')$ in lex order then return $f' * g'$, where lex order means to compare the leftmost unequal elements.
 - (iii) If $p(g')$ precedes $p(f')$ in lex order then: if $* = \circ$ then return $g' * f'$ else return $f' * g'$.
 - (iv) If $p(f') = p(g')$ then return $f' * g'$.

4.3. Relations of arity 4.

Theorem 4.9. *Over a field \mathbb{F} of characteristic 0 or $p > 4$, every multilinear polynomial identity of arity ≤ 4 , satisfied by the Jordan product and diproduct in every triassociative algebra, is a consequence of the commutativity of \circ , the black right bar identity, and the (linearizations of the) following identities of arity 4:*

- (6) $((a \circ a) \circ b) \circ a \equiv (a \circ a) \circ (b \circ a),$
- (7) $(a \bullet (b \bullet b)) \bullet b \equiv (a \bullet b) \bullet (b \bullet b),$
- (8) $((a \bullet b) \bullet d) \bullet c + ((a \bullet c) \bullet d) \bullet b + a \bullet ((b \bullet c) \bullet d)$
 $\equiv (a \bullet (b \bullet c)) \bullet d + (a \bullet (b \bullet d)) \bullet b + (a \bullet (c \bullet d)) \bullet b,$
- (9) $(a \circ a) \circ (a \bullet b) \equiv ((a \circ a) \bullet b) \circ a,$
- (10) $((a \bullet d) \circ c) \circ b + ((b \bullet d) \circ c) \circ a + ((a \circ b) \circ c) \bullet d$
 $\equiv ((a \circ b) \bullet d) \circ c + ((a \circ c) \bullet d) \circ b + ((b \circ c) \bullet d) \circ a,$
- (11) $((a \bullet d) \bullet c) \circ b + ((b \bullet d) \bullet c) \circ a + ((a \circ b) \bullet c) \bullet d$
 $\equiv (b \bullet (c \bullet d)) \circ a + ((a \bullet c) \circ b) \bullet d + ((a \bullet d) \circ b) \bullet c,$
- (12) $(a \bullet c) \circ (b \bullet d) + (a \bullet d) \circ (b \bullet c) + (a \circ b) \bullet (c \bullet d)$
 $\equiv (b \bullet (c \bullet d)) \circ a + ((a \bullet c) \circ b) \bullet d + ((a \bullet d) \circ b) \bullet c.$

Proof. Similar to that of Lemma 4.2, but the matrices are larger, and we must deal with the consequences of the black right bar identity of arity 3. We first construct the 135 basis skeletons in $\text{BBB}(4)$, where $*_1, *_2, *_3$ are chosen freely from $\{\vdash, \perp, \dashv\}$:

$$((x*_1x)*_2x)*_3x, (x*_1(x*_2x))*_3x, (x*_1x)*_2(x*_3x), x*_1((x*_2x)*_3x), x*_1(x*_2(x*_3x)).$$

Second, we generate the 165 consequences in arity 4 of the defining relations for TriAss using (3): each relation $f(a, b, c)$ has 15 consequences where $*$ is chosen from $\{\vdash, \perp, \dashv\}$:

$$f(a * d, b, c), \quad f(a, b * d, c), \quad f(a, b, c * d), \quad f(a, b, c) * d, \quad d * f(a, b, c).$$

These consequences generate an equivalence relation on the basis of $\text{BBB}(4)$, which has 15 classes representing a basis of $\text{TriAss}(4)$. We choose the minimal element in each class as the normal form of the (nonsymmetric) monomials in that class:

$$\begin{aligned} &((a \vdash b) \vdash c) \vdash d, & ((a \vdash b) \vdash c) \dashv d, & ((a \vdash b) \vdash c) \perp d, & ((a \vdash b) \dashv c) \dashv d, \\ &((a \vdash b) \dashv c) \perp d, & ((a \vdash b) \perp c) \dashv d, & ((a \vdash b) \perp c) \perp d, & ((a \vdash b) \dashv c) \dashv d, \\ &((a \dashv b) \dashv c) \perp d, & ((a \dashv b) \perp c) \dashv d, & ((a \dashv b) \perp c) \perp d, & ((a \perp b) \dashv c) \dashv d, \\ &((a \perp b) \dashv c) \perp d, & ((a \perp b) \perp c) \dashv d, & ((a \perp b) \perp c) \perp d. & \end{aligned}$$

To construct a monomial basis of the symmetrization, we apply all permutations of a, b, c, d ; thus $\Sigma\text{TriAss}(4)$, the codomain of $E(4)$, has dimension 360.

The domain of $E(4)$ is the space $\text{BW}(4)$. To generate an ordered basis of $\text{BW}(4)$, we start with the 25 skeletons in Example 4.7. In each skeleton, we replace the four occurrences of the symbol x by all permutations of x_1, \dots, x_4 ; for each resulting

multilinear monomial we compute the normal form with respect to commutativity, and save only those monomials which are irreducible (equal to their normal forms). The number of monomials we obtain for each skeleton is $4!/2^s$ where s is the number of commutative symmetries in the skeleton, where by definition a symmetry is a sub-skeleton of the form $f \circ f$. For example, the skeleton $(x \circ x) \circ (x \circ x)$ has three symmetries: one with $f = x \circ x$ and two with $f = x$, so it has only three multilinear monomials, corresponding to permutations 1234, 1324, 1423 of the subscripts. We order these monomials first by skeleton and then by permutation. The total number of multilinear monomials over all skeletons is $\dim \mathbf{BW}(4) = 405$.

With respect to the ordered monomial bases of $\mathbf{BW}(4)$ and $\Sigma\text{TriAss}(4)$, the expansion map $E(4): \mathbf{BW}(4) \rightarrow \Sigma\text{TriAss}(4)$ is represented by the 360×405 matrix E_4 . The (i, j) entry of E_4 is the coefficient of the i -th ΣTriAss monomial in the expansion of the j -th \mathbf{BW} monomial. We must replace each ΣTriAss monomial m in each expansion by the equivalent element in the monomial basis for $\Sigma\text{TriAss}(4)$; we replace the TriAss skeleton of m by the representative of the corresponding equivalence class. For example,

$$\begin{aligned} & E(4)\left((a \bullet b) \circ (c \bullet d)\right) \\ &= (a \dashv b) \perp (c \dashv d) + (c \dashv d) \perp (a \dashv b) + (a \dashv b) \perp (d \dashv c) + (d \dashv c) \perp (a \dashv b) + \\ & \quad (b \dashv a) \perp (c \dashv d) + (c \dashv d) \perp (b \dashv a) + (b \dashv a) \perp (d \dashv c) + (d \dashv c) \perp (b \dashv a) \\ &= ((a \dashv b) \perp c) \dashv d + ((c \dashv d) \perp a) \dashv b + ((a \dashv b) \dashv d) \perp c + ((d \dashv c) \perp a) \dashv b + \\ & \quad ((b \dashv a) \perp c) \dashv d + ((c \dashv d) \dashv b) \perp a + ((b \dashv a) \dashv d) \perp c + ((d \dashv c) \dashv b) \perp a. \end{aligned}$$

In this way, we initialize E_4 , which has entries in $\{0, 1\}$. This matrix is large, so we use modular arithmetic ($p = 101$) to compute its RCF and the corresponding basis for its nullspace.

We find that E_4 has rank 165 and nullity 240. Its nullspace $N(4) \subseteq \mathbf{BW}(4)$ is the S_4 -module which contains the coefficient vectors of the multilinear identities of arity 4 satisfied by the Jordan product and diproduct in every triassociative algebra. The new identities are the (nonzero) elements of the quotient module $N(4)/M(4)$ where $M(4)$ is the S_4 -module generated by the known identities:

- (1) Consequences of the black right bar identity, $f(a, b, c) = a \bullet (b \circ c) - a \bullet (b \bullet c) \equiv 0$, which generate an S_4 -module of dimension 180: $f(a \circ d, b, c)$, $f(a, b \circ d, c)$, $f(a, b, c \circ d)$, $f(a, b, c) \circ d$, $f(a \bullet d, b, c)$, $f(a, b \bullet d, c)$, $f(a, b, c \bullet d)$, $f(a, b, c) \bullet d$, $d \bullet f(a, b, c)$.
- (2) Linearization of the Jordan identity for $a \circ b = a \perp b + b \perp a$. Combining this with the previous generators, we obtain an S_4 -module of dimension 184.
- (3) Remark 4.4 shows that we do not need to consider the consequences of right commutativity for \bullet , since they are also consequences of the black right bar identity.
- (4) Linearizations of the Jordan dialgebra identities for $a \bullet b = a \dashv b + b \dashv a$. Combining these with the other generators, we obtain the S_4 -module $M(4)$ of dimension 200.

We represent $M(4)$ as the row space of a 200×405 matrix M_4 . The quotient module $N(4)/M(4)$ of new identities of arity 4 has dimension 40.

From the RCF of E_4 we extract a ‘‘canonical’’ basis of $N(4)$ as follows: for each standard basis vector v of \mathbb{F}_p^{240} , we set the 240 free variables equal to the components of v and solve for the leading variables. We put the resulting basis vectors into the rows of a 240×405 matrix N_4 whose row space is $N(4)$. (A canonical basis for $N(4)$

would consist of the rows of $\text{RCF}(N_4)$.) Checking the rows of N_4 (all identities) one-by-one against the row space of M_4 (known identities), we obtain a list of 40 rows which increase the rank. Each row contains 6 nonzero entries, representing an identity with 6 terms. The cosets of these row vectors form a basis of the quotient module $N(4)/M(4)$.

From this linear basis of 40 vectors, we extract a minimal set of S_4 -module generators. Checking again the rows ρ of N_4 against the row space of M_4 , but now including all 24 permutations of the identity with coefficient vector ρ , we obtain 5 rows which increase the rank to 204, 216, 222, 232, 240. We find that the third row belongs to the submodule generated by the others. The remaining four generators are identities (9)-(12). \square


Remark 4.10. The operation \circ satisfies the Jordan identity (6); the operation \bullet satisfies the Jordan dialgebra identities (7)-(8); the operations are related by (9)-(12). This pattern of identities is similar to Definition 3.4 of Lie trialgebras: $[a, b]$ defines a Lie algebra, $\{a, b\}$ defines a Leibniz algebra, and other identities relate the two operations.

Definition 4.11. Over a field \mathbb{F} of characteristic 0 or $p > 4$, a vector space J with bilinear operations \circ and \bullet is a **Jordan trialgebra** if:

- (J, \circ) is a Jordan algebra, and (J, \bullet) is a Jordan dialgebra, and
- the two operations are related by identities (9)–(12) of Proposition 4.9.

The corresponding symmetric operad will be denoted TriJor .

4.4. Triplicators. The operad TriJor governing Jordan trialgebras may also be obtained using the techniques of Pei et al. [47]. We start from the linearized Jordan identity, $((ab)d)c + ((ac)d)b + ((bc)d)a - (ab)(cd) - (ac)(bd) - (ad)(bc) \equiv 0$, and its representation in terms of tree monomials, using \cdot for the operation symbol:



Let $L \subseteq \{a, b, c, d\}$ be a nonempty subset of the arguments. For each L we construct a new polynomial identity involving operations $\circ_1, \circ_2, \circ_3$ which we denote simply by 1, 2, 3. We apply the following triplicator algorithm to each tree monomial m in the identity above.

Algorithm 4.12. For each $x \in L$, let $p_m(x)$ be the unique path from the root of m to the leaf x . Let W_m be the set of internal vertices of m , and define $t_m: L \times W_m \rightarrow \{\leftarrow, 0, \rightarrow\}$:


- If the internal vertex $v \in W_m$ does not lie on the path $p_m(x)$ then $t_m(x, v) = 0$.
- If $v \in W_m$ lies on $p_m(x)$ then $t_m(x, v) = \leftarrow$ (resp. \rightarrow) if $p_m(x)$ turns left (resp. right) at v .

There are four possibilities for $T_m(L, v) = \{t_m(x, v) \mid x \in L\} \setminus \{0\}$:


- If $T_m(L, v) = \{\leftarrow\}$ then the internal vertex v receives the new operation symbol 1.
- If $T_m(L, v) = \{\rightarrow\}$ then the internal vertex v receives the new operation symbol 3.
- If $T_m(L, v) = \{\leftarrow, \rightarrow\}$ then the internal vertex v receives the new operation symbol 2.

- If $T_m(L, v) = \emptyset$ then none of the paths $p_m(x)$ pass through v , and in this case v receives the symbol $*$ representing the union of the new operation symbols: $*$ = $\{1, 2, 3\}$.

Since the linearized Jordan identity is symmetric in a, b, c it suffices to consider only these subsets L : $\{a\}, \{d\}, \{a, b\}, \{a, d\}, \{a, b, c\}, \{a, b, d\}, \{a, b, c, d\}$. For example, if $L = \{a, c\}$ then Algorithm 4.12 produces this new identity:



Note that term 5 contains a vertex labelled $*$; it thus produces three new identities after we replace $*$ by 1, then 2, then 3. These new operations satisfy symmetries which follow from the commutativity of the Jordan product: $a \circ_1 b \equiv b \circ_3 a$ and $a \circ_2 b \equiv b \circ_2 a$. We replace $*$ in term 5 by operation 1, and apply the symmetries of the operations to eliminate operation 3 from terms 3 and 6. The final result involves only two new operations 1, 2:



If we replace 2 by \circ , and 1 by \bullet , and rewrite the tree monomials in parenthesized form, then we obtain multilinear identities which we can compare directly to those of Theorem 4.9.

Definition 4.13. The operad defined by the binary operations \circ and \bullet , where \circ is commutative and \bullet has no symmetry, satisfying the multilinear relations resulting from all possible applications of Algorithm 4.12 to the linearized Jordan identity, is denoted TriJor and is called the **triplicator** of the Jordan operad Jor .

Proposition 4.14. Let TriJor be the symmetric operad governing Jordan trialgebras, generated by the operations \circ (commutative) and \bullet (no symmetry) satisfying the identities of Theorem 4.9. Let TripJor be the symmetric operad generated by the same operations but satisfying the identities obtained by applying the triplicator Algorithm 4.12 to the linearized Jordan identity. These two sets of identities generate the same subquotient of the S_4 -module $\text{BW}(4)$, and therefore the two operads are isomorphic: $\text{TriJor} \cong \text{TripJor}$.

4.5. Representation theory of the symmetric group. We can extend these calculations to arities $n \geq 5$, but the matrices become very large, and so we use the representation theory of the symmetric group to decompose the S_n -modules into isotypic components. For the theoretical background see the survey [13], which includes an extensive bibliography. We mention Rutherford’s exposition [48] of Young’s work, Hentzel’s implementation [32] on a computer, and the important contributions by his students Bondari [8] and Clifton [19].

For $n \geq 1$, over a field \mathbb{F} of characteristic 0 or $p > n$, the group algebra $\mathbb{F}S_n$ is semisimple and hence decomposes as the direct sum of simple two-sided ideals isomorphic to full matrix rings. These ideals are in bijection with the partitions of

n : for partition λ we have the matrix ring $M_{d_\lambda}(\mathbb{F})$ where the dimension d_λ is given by the hook formula. If $Y(\lambda)$ is the Young diagram for λ then d_λ is the number of standard tableaux for $Y(\lambda)$. For each λ the projection $R_\lambda: \mathbb{F}S_n \rightarrow M_{d_\lambda}(\mathbb{F})$ defines an irreducible representation of S_n ; we call $R_\lambda(\pi)$ the representation matrix of π for partition λ . If we define the action of S_n on $M_{d_\lambda}(\mathbb{F})$ by $\pi \cdot A = R_\lambda(\pi)A$ for $A \in M_{d_\lambda}(\mathbb{F})$, then $M_{d_\lambda}(\mathbb{F})$ becomes a (left) S_n -module, which is the direct sum of d_λ isomorphic minimal left ideals in $\mathbb{F}S_n$: the column vectors in $M_{d_\lambda}(\mathbb{F})$. We write $[\lambda]$ for the isomorphism class of these minimal left ideals; $[\lambda]$ is the irreducible S_n -module for partition λ . Clifton [19] found an efficient algorithm to compute $R_\lambda(\pi)$.

4.5.1. *All relations: kernel of the expansion map.* We show how to compute the kernel of the expansion map $E(n)$ using representation theory; recall that $E(n)$ is a linear map from $\mathbb{B}\mathbb{W}(n)$ to $\Sigma\text{TriAss}(n)$. Let $\text{BWS}(n) = \{\beta_1, \dots, \beta_{b(n)}\}$ be the ordered set of skeletons for $\mathbb{B}\mathbb{W}(n)$ (Definition 4.5), and let $\text{NTA}(n) = \{\tau_1, \dots, \tau_{t(n)}\}$ be the ordered basis for $\text{TriAss}(n)$ – the representatives of the equivalence classes in $\text{BBB}(n)$ – which are the skeletons for $\Sigma\text{TriAss}(n)$. The advantage of representation theory is that we do not need all multilinear monomials, but only the skeletons: a much smaller set. For each partition λ of n we compute the matrix $E_{n,\lambda}$, which has $b(n)$ rows and $t(n) + b(n)$ columns of $d_\lambda \times d_\lambda$ blocks.

$$(13) \quad E_{n,\lambda} = \left[\begin{array}{cccc|cccc} R_\lambda(X_{1,1}) & R_\lambda(X_{1,2}) & \cdots & R_\lambda(X_{1,t(n)}) & I_{d_\lambda} & 0 & \cdots & 0 \\ R_\lambda(X_{2,1}) & R_\lambda(X_{2,2}) & \cdots & R_\lambda(X_{2,t(n)}) & 0 & I_{d_\lambda} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ R_\lambda(X_{b(n),1}) & R_\lambda(X_{b(n),2}) & \cdots & R_\lambda(X_{b(n),t(n)}) & 0 & 0 & \cdots & I_{d_\lambda} \end{array} \right]$$

For $1 \leq i \leq b(n)$ and $1 \leq j \leq t(n)$, position (i, j) contains the representation matrix $R_\lambda(X_{i,j})$ where $X_{i,j} \in \mathbb{F}S_n$ is determined as follows:

- (i) Start with the i -th skeleton β_i from $\text{BWS}(n)$.
- (ii) Replace the n occurrences of the symbol x in β_i by the identity permutation x_1, \dots, x_n of n indeterminates to obtain the multilinear basis monomial μ_i for $\mathbb{B}\mathbb{W}(n)$.
- (iii) Apply $E(n)$ to μ_i , obtaining $X_i = E(n)(\mu_i) \in \text{TriAss}(n)$. The terms of the expansion $E(n)(\mu_i)$ belong to the symmetrization of $\text{BBB}(n)$, so we replace each BBB skeleton by the representative of its triassociative equivalence class; see the proofs of Lemma 4.2 and Proposition 4.9. After this normalization of the skeletons, $X_i \in \Sigma\text{TriAss}(n)$.
- (iv) Decompose $X_i = X_{i,1} + \cdots + X_{i,t(n)}$ as a sum of $t(n)$ components where $X_{i,j}$ consists of the terms of X_i with triassociative skeleton τ_j .
- (v) In $X_{i,j}$ every term has skeleton τ_j ; the only differences are in the coefficients and the permutations. Thus $X_{i,j}$ is a linear combination of permutations of n , so $X_{i,j} \in \mathbb{F}S_n$.
- (vi) Compute $R_\lambda(X_{i,j})$, the representation matrix of $X_{i,j}$ and store it in block (i, j) of $E_{n,\lambda}$.
- (vii) Finally compute the row canonical form of $E_{n,\lambda}$.

Remark 4.15. The skeletons in $\text{TriAss}(n)$ are in bijection with the equivalence classes of skeletons in $\text{BBB}(n)$: equivalence is determined by the (consequences of the) triassociative relations. Computing this equivalence relation is only practical

in low arities. In general, we use the explicit description of the free trioid on one generator [43].

Definition 4.16. The **toprank** of $\text{RCF}(E_{n,\lambda})$ is the number $\text{top}(n, \lambda)$ of rows with leading 1s to the left of the vertical line in (13). That is, $\text{top}(n, \lambda)$ is the largest i such that the leading 1 of row i is in column $j \leq t(n)d_\lambda$. Every row $i' > i$ has only 0s to the left of the vertical line. The **allrank** of $\text{RCF}(E_{n,\lambda})$ is the number $\text{all}(n, \lambda) = b(n)d_\lambda - \text{top}(n, \lambda)$ of rows with leading 1s to the right of the vertical line.

Lemma 4.17. *The last $\text{all}(n, \lambda)$ rows of $\text{RCF}(E_{n,\lambda})$ are independent S_n -module generators for the isotypic component of partition λ in the kernel of $E(n)$: the sum of all irreducible submodules of $\text{kernel}(E(n))$ isomorphic to $[\lambda]$.*

Definition 4.18. $\text{ALL}(n, \lambda)$, the **matrix of all identities** for arity n , partition λ , is the lower right block of $\text{RCF}(E_{n,\lambda})$, with upper left corner in row $\text{top}(n, \lambda) + 1$ and column $t(n)d_\lambda + 1$.

4.5.2. *Old relations; symmetries of the skeletons.* For each partition λ of n , the nullspace of $E_{n,\lambda}$ contains all polynomial identities in the isotypic component for λ which are satisfied by the Jordan product \circ and diproduct \bullet in every triassociative algebra. In general, many of these are consequences of known identities of lower arity.

We must also consider the symmetries of the skeletons in $\text{BWS}(n)$. If n is small, we can compute a monomial basis and avoid representation theory: to each skeleton, we apply all permutations of the n variables, and retain a monomial only if it equals its normal form. For example, let $n = 3$ and consider $(x \circ x) \bullet x$: commutativity of \circ implies that the six permutations of a, b, c produce only three normal forms: $(a \circ b) \bullet c$, $(a \circ c) \bullet b$, $(b \circ c) \bullet a$. If n is large, so that it is not practical to compute a monomial basis, then we use representation theory, and so we must encode the symmetries in some other way. For a small example, from the skeleton $(x \circ x) \bullet x$ we construct a multilinear identity which can be processed using representation theory: $(x_1 \circ x_2) \bullet x_3 - (x_2 \circ x_1) \bullet x_3 \equiv 0$.

Definition 4.19. We call the identities of this form the **symmetries** of the skeletons. In arity n , each symmetry has the form $m_1 - m_2 \equiv 0$, where m_1, m_2 are monomials of arity n with the same skeleton; m_1 has the identity permutation of the variables, and m_2 has a permutation of order 2; and the commutativity of \circ implies that $m_1 = m_2$.

Lemma 4.20. *The relation between multilinear monomials, the skeletons, and their symmetries, is given by this equation, where $s(\beta)$ is the number of symmetries of skeleton β :*

$$\dim \text{BW}(n) = \sum_{\beta \in \text{BWS}(n)} \frac{n!}{2^{s(\beta)}}$$

Definition 4.21. Recall that $b(n)$ is the size of $\text{BWS}(n)$: the total number of BW skeletons in arity n . Let $\text{sym}(n)$ be the total number of symmetries over all skeletons in arity n . Let $\text{con}(n)$ be the total number of consequences in arity n of the known identities of lower arities. Let $K_{n,\lambda}$ be the matrix of size $(\text{sym}(n) + \text{con}(n))d_\lambda \times b(n)d_\lambda$, consisting of $d_\lambda \times d_\lambda$ blocks; the block in position (i, j) is $R_\lambda(Y_{i,j})$ where $Y_{i,j}$ is the component of symmetry i in skeleton j for $i = 1, \dots, \text{sym}(n)$, and $Y_{\text{sym}(n)+i,j}$ is the component of consequence i in skeleton j for $i = 1, \dots, \text{con}(n)$. The **matrix of old identities** for arity n and partition λ is defined by $\text{OLD}(n, \lambda) = \text{RCF}(K_{n,\lambda})$.

4.6. Relations of arity ≥ 5 : a result, a problem, and two conjectures.

Proposition 4.22. *Every multilinear polynomial identity of arity ≤ 6 relating the Jordan product and diproduct in every triassociative algebra is a consequence of the defining identities for Jordan trialgebras (Definition 4.11).*

Proof. For $n = 5, 6$ and all partitions λ of n , we computed $\text{ALL}(n, \lambda)$ and $\text{OLD}(n, \lambda)$, which are both in row canonical form, and found that they were equal in every case. \square

Problem 4.23. Determine whether or not special identities of arity 8 exist for the Jordan diproduct in the free diassociative algebra; such identities (if they exist) are called special identities for Jordan dialgebras. The existence of such identities for the pre-Jordan product in the free dendriform algebra has been established [12].

Conjecture 4.24. Over a field \mathbb{F} of characteristic 0 or $p > 7$, every multilinear polynomial identity of arity ≤ 7 satisfied by the Jordan product and diproduct in every triassociative algebra is a consequence of the defining identities for Jordan trialgebras.

Conjecture 4.25. There exist identities of arity 8 the Jordan product and diproduct in every triassociative algebra which are not consequences of the identities defining Jordan trialgebras, the (linearizations of) the Glennie identities for Jordan algebras [28], and the multilinear special identities of arity 8 for Jordan dialgebras (if they exist).

5. ENUMERATION OF ASSOCIATION TYPES AND SKELETONS

We consider two integer sequences: the number of skeletons of arity n in the symmetric operad BW generated by two binary operations, one commutative, one noncommutative; the number of multilinear monomials of arity n in BW , which is $\dim \text{BW}(n)$.

5.1. One commutative operation. The number of association types is sequence A001190 in the OEIS (oeis.org), the Wedderburn-Etherington numbers [25, 54]:

1, 1, 1, 2, 3, 6, 11, 23, 46, 98, 207, 451, 983, 2179, 4850, 10905, 24631, 56011, ...

This sequence also enumerates complete rooted binary trees with n leaves, up to abstract graph isomorphism, so that (for example) the two trees with three (unlabelled) leaves are isomorphic. The corresponding number of multilinear monomials is A001147, the double factorial of odd numbers: $(2n-1)!! = 1 \cdot 3 \cdot 5 \cdots (2n-1)$. This is also the number of complete rooted binary trees with leaf labels $1, \dots, n$ up to abstract isomorphism.

5.2. One noncommutative operation. The number of association types is the (shifted) Catalan number $C(n-1)$ where $C(n) = (2n)!/(n!(n+1)!)$. (The shift is required since n in $C(n)$ is the number of operations but for us n is the number of arguments.) This sequence A000108 also enumerates complete rooted binary *plane* trees with n leaves, so that the two trees with three (unlabelled) leaves are *not* isomorphic. In this case, the operation has no symmetry, so the number of multilinear monomials is simply $n!C(n-1) = (2n-2)!/(n-1)!$, which is the quadruple factorial $(4n-2)!!!! = 2 \cdot 6 \cdot 10 \cdots (4n-2)$.

5.3. Two operations: one commutative, one noncommutative. Computational enumeration of the skeletons produced the following sequence, A276277 in the OEIS:

1, 2, 6, 25, 111, 540, 2736, 14396, 77649, 427608, 2392866, 13570386, 77815161, ...

The number of multilinear monomials is the sextuple factorial, sequence A011781:

$$(6n-3)!!!!!! = \prod_{k=1}^{n-1} (6k-3) \implies 1, 3, 27, 405, 8505, 229635, 7577955, 295540245, \dots$$

Problem 5.1. For $p, q \geq 1$ let $\mathbb{B}\bar{W}_{p,q}$ be the free symmetric operad generated by p commutative and q noncommutative binary operations. For $n \geq 1$ determine the number of skeletons and the number of multilinear monomials in $\mathbb{B}\bar{W}_{p,q}(n)$.

6. POST-JORDAN ALGEBRAS

In this section we determine the multilinear polynomial identities of arity ≤ 6 satisfied by the Jordan and pre-Jordan products $a \circ b = a \wedge b + b \wedge a$ and $a \bullet b = a < b + b > a$ in the free tridendriform algebra. In addition to the commutativity of \circ , there are no new identities in arity 3, a nonzero S_4 -module of new identities in arity 4 for which we find a minimal set of generators, and no new identities in arities 5 and 6. The commutativity of \circ together with the new identities in arity 4 define post-Jordan algebras.

6.1. Gröbner-Shirshov basis for the tridendriform operad. The computations for post-Jordan algebras are very similar to those for Jordan trialgebras; in particular, the domain $\mathbb{B}\bar{W}(n)$ of the expansion map $E(n)$ is the same. However, the codomain of the expansion map is no longer ΣTriAss but its Koszul dual $\Sigma\text{TriDend} \cong \Sigma\text{TriAss}^\dagger$. The defining relations for TriAss are monomial relations (Definition 3.1): this allows us to compute normal forms in TriAss very simply in terms of equivalence relations on BBB monomials. On the other hand, not all the defining relations for TriDend are monomial relations (Definition 3.1). Hence computing normal forms in TriDend requires first determining a Gröbner-Shirshov (GS) basis from the defining relations, and this requires fixing a total order on the operations. Checking all six possibilities, we find that the smallest GS basis for TriDend comes from the order $\wedge, <, >$ where the commutative operation comes first. This GS basis for TriDend is almost the same as the original defining relations, except that the relations appear in a different order, as do the terms within each relation.

Lemma 6.1. *Starting with the relations in Definition 3.1, and assuming the order of operations $\wedge, <, >$, we obtain the following Gröbner-Shirshov basis for TriDend :*

$$\begin{aligned} (a < b) < c - a < (b > c) - a < (b < c) - a < (b \wedge c), & (a > b) < c - a > (b < c), \\ (a > b) > c + (a < b) > c + (a \wedge b) > c - a > (b > c), & (a > b) \wedge c - a > (b \wedge c), \\ (a < b) \wedge c - a \wedge (b > c), & (a \wedge b) < c - a \wedge (b < c), & (a \wedge b) \wedge c - a \wedge (b \wedge c). \end{aligned}$$

Proof. Gröbner bases for operads were introduced in [22]; the special case of nonsymmetric operads appears in [23]. Similar computations are explained in detail in [45]. \square

We use this GS basis for TriDend to compute normal forms of nonsymmetric tridendriform polynomials. This also applies in the symmetric case, since $\Sigma\text{TriDend}$ is the symmetrization of TriDend . A monomial $m \in \Sigma\text{TriDend}(n)$ has a skeleton

$s \in \text{TriDend}(n)$ and a permutation $p \in S_n$ of the arguments x_1, \dots, x_n . To find the normal form of m , we compute the normal form of s , and then replace the arguments $x_{p(1)}, \dots, x_{p(n)}$. For details on nonsymmetric operads and their Gröbner bases, see [10].

6.2. Relations of arity ≤ 4 .

Lemma 6.2. *Every multilinear polynomial identity of arity 3 for the Jordan and pre-Jordan products \circ, \bullet in the free tridendriform algebra is a consequence of the commutativity of \circ .*

Proof. The expansion matrix E_3 has size 66×27 : see Table 2, where the matrix is split into top and bottom halves. Following the proof of Lemma 4.2, the columns correspond to the ordered monomial basis of the domain $\text{BW}(3)$ of the expansion map $E(3)$; see the proof of Lemma 3.5 where the symbols $\wedge, <$ are used instead of \circ, \bullet . As for the codomain, since $\dim \text{BBB}(3) = 18$, and the GS basis of Lemma 6.1 has 7 elements, there are 7 leading monomials which are linear combinations of the other 11 monomials, so $\dim \text{TriDend}(3) = 11$. To each nonsymmetric basis monomial we apply all 6 permutations of the arguments; this gives the ordered monomial basis of $\Sigma \text{TriDend}(3)$ corresponding to the 66 rows of E_3 . An easy calculation shows that E_3 has full rank and so its nullspace is 0. \square

TABLE 2. Top and bottom of post-Jordan expansion matrix in arity 3

Theorem 6.3. *Over a field \mathbb{F} of characteristic 0 or $p > 4$, every multilinear polynomial identity of arity ≤ 4 satisfied by the Jordan and pre-Jordan products in the free tridendriform algebra is a consequence of the commutativity of \circ and the (linearizations of the) following 7 identities of arity 4 (there are no new identities of arity 3):*

$$\begin{aligned}
& ((c \bullet a) \circ b) \circ d + ((d \bullet a) \circ b) \circ c - ((b \circ c) \bullet a) \circ d - ((b \circ d) \bullet a) \circ c \\
& \quad - ((c \circ d) \bullet a) \circ b + ((c \circ d) \circ b) \bullet a \equiv 0, \\
& ((b \circ c) \bullet a) \circ d + ((b \circ d) \bullet a) \circ c + ((c \circ d) \bullet a) \circ b - (b \circ c) \circ (d \bullet a) \\
& \quad - (b \circ d) \circ (c \bullet a) - (c \circ d) \circ (b \bullet a) \equiv 0, \\
& ((a \circ b) \circ d) \circ c + ((a \circ c) \circ d) \circ b + ((b \circ c) \circ d) \circ a - (a \circ b) \circ (c \circ d) \\
& \quad - (a \circ c) \circ (b \circ d) - (a \circ d) \circ (b \circ c) \equiv 0,
\end{aligned}$$

$$\begin{aligned}
 & ((c \bullet a) \bullet b) \circ d + ((d \bullet a) \bullet b) \circ c - (d \bullet (a \circ b)) \circ c - (d \bullet (a \bullet b)) \circ c \\
 & \quad - (d \bullet (b \bullet a)) \circ c - ((c \bullet a) \circ d) \bullet b - ((c \bullet b) \circ d) \bullet a + ((c \circ d) \bullet b) \bullet a \equiv 0, \\
 & (d \bullet (a \circ b)) \circ c + (d \bullet (a \bullet b)) \circ c + (d \bullet (b \bullet a)) \circ c - (c \bullet a) \circ (d \bullet b) \\
 & \quad - (c \bullet b) \circ (d \bullet a) + ((c \bullet a) \circ d) \bullet b + ((c \bullet b) \circ d) \bullet a - (c \circ d) \bullet (a \circ b) \\
 & \quad - (c \circ d) \bullet (a \bullet b) - (c \circ d) \bullet (b \bullet a) \equiv 0, \\
 & (d \bullet (a \circ b)) \bullet c + (d \bullet (a \circ c)) \bullet b + (d \bullet (b \circ c)) \bullet a + (d \bullet (a \bullet b)) \bullet c \\
 & \quad + (d \bullet (a \bullet c)) \bullet b + (d \bullet (b \bullet a)) \bullet c + (d \bullet (b \bullet c)) \bullet a + (d \bullet (c \bullet a)) \bullet b \\
 & \quad + (d \bullet (c \bullet b)) \bullet a - (d \bullet a) \bullet (b \circ c) - (d \bullet b) \bullet (a \circ c) - (d \bullet c) \bullet (a \circ b) \\
 & \quad - (d \bullet a) \bullet (b \bullet c) - (d \bullet a) \bullet (c \bullet b) - (d \bullet b) \bullet (a \bullet c) - (d \bullet b) \bullet (c \bullet a) \\
 & \quad - (d \bullet c) \bullet (a \bullet b) - (d \bullet c) \bullet (b \bullet a) \equiv 0, \\
 & ((d \bullet a) \bullet c) \bullet b + ((d \bullet b) \bullet c) \bullet a - (d \bullet (a \circ b)) \bullet c - (d \bullet (a \circ c)) \bullet b \\
 & \quad - (d \bullet (b \circ c)) \bullet a - (d \bullet (a \bullet b)) \bullet c - (d \bullet (a \bullet c)) \bullet b - (d \bullet (b \bullet a)) \bullet c \\
 & \quad - (d \bullet (b \bullet c)) \bullet a - (d \bullet (c \bullet a)) \bullet b - (d \bullet (c \bullet b)) \bullet a + d \bullet ((a \circ b) \circ c) \\
 & \quad + d \bullet ((a \bullet b) \circ c) + d \bullet ((b \bullet a) \circ c) + d \bullet ((a \circ b) \bullet c) + d \bullet ((a \bullet b) \bullet c) \\
 & \quad + d \bullet ((b \bullet a) \bullet c) + d \bullet (c \bullet (a \circ b)) + d \bullet (c \bullet (a \bullet b)) + d \bullet (c \bullet (b \bullet a)) \equiv 0.
 \end{aligned}$$


Proof. All the techniques have already been discussed, so we will be very brief. The expansion matrix E_4 has size 1080×405 and rank 345; its nullity is 60, and every nonzero vector in the nullspace is a new identity, since there are no consequences from arity 3. We extract the canonical basis vectors for the nullspace and sort the identities by increasing number of terms. We find a subset of seven identities which generates the nullspace as an S_4 -module; none belongs to the S_4 -module generated by the others. These identities have 6, 6, 6, 8, 10, 18, 20 terms. The first is the linearized Jordan identity which contains only the operation \circ . Every other identity contains both operations. If we remove every term containing \circ from these seven identities, then the first five identities become 0, and the last two identities become the defining identities in arity 4 for pre-Jordan algebras. \square

Remark 6.4. In a Jordan trialgebra, the Jordan product \circ is commutative and satisfies the Jordan identity; the Jordan diproduct \bullet satisfies the defining relations for Jordan dialgebras. Thus a Jordan trialgebra is a sum or split extension (roughly speaking) of a Jordan algebra by a Jordan dialgebra. In a post-Jordan algebra, the Jordan product \circ is commutative and satisfies the Jordan identity; however, the pre-Jordan product \bullet *does not* satisfy the defining identities for pre-Jordan algebras. In order to obtain the pre-Jordan identities, we must set the Jordan product to zero: remove every term containing \circ . Thus a post-Jordan algebra is a *non-split* extension of a Jordan algebra by a pre-Jordan algebra.

Definition 6.5. Over a field \mathbb{F} of characteristic 0 or $p > 4$, a vector space J with bilinear operations \circ and \bullet is a **post-Jordan algebra** if \circ is commutative and \circ, \bullet together satisfy the multilinear polynomial identities of Theorem 6.3. In particular, (J, \circ) is a Jordan algebra. The corresponding symmetric operad is denoted PostJor .

6.3. Trisuccessors. The operad PostJor governing post-Jordan algebras may be obtained using the techniques of [47]. As in §4.4, we start from the linearized Jordan identity represented in terms of tree monomials. We apply Algorithm 4.12 but with the difference that $*$ no longer represents the union $\{\circ_1, \circ_2, \circ_3\}$ but the sum $\circ_1 + \circ_2 + \circ_3$. As before, we consider the example $L = \{a, c\}$. In the resulting tree polynomial, term 5 has a vertex labelled $*$, which is now replaced by the sum of three terms $5', 5'', 5'''$ obtained by substituting (operations) 1, 2, 3 for $*$; the result

is a tree polynomial with eight terms. As in §4.4, the three new operations satisfy symmetries: $a \circ_1 b \equiv b \circ_3 a$ and $a \circ_2 b \equiv b \circ_2 a$. These allow us to rewrite the tree polynomial using only two operations:



We replace operation 2 by \circ , and 1 by \bullet , and convert the tree monomials to parentheses and permutations. We perform this calculation for all nonempty $L \subseteq \{a, b, c, d\}$ and obtain multilinear identities which can be compared directly to those of Theorem 6.3.

Definition 6.6. The operad defined by the binary operations \circ and \bullet , where \circ is commutative and \bullet has no symmetry, satisfying the multilinear relations obtained from all possible applications of the trisuccessor algorithm to the linearized Jordan identity, is denoted TriSucJor and is called the **trisuccessor** of the Jordan operad Jor .

Proposition 6.7. Let PostJor be the symmetric operad governing post-Jordan algebras, generated by the binary operations \circ (commutative) and \bullet (no symmetry) satisfying the multilinear identities of Theorem 6.3. Let TriSucJor be the symmetric operad generated by the same operations but satisfying the multilinear identities obtained by applying the trisuccessor algorithm to the linearized Jordan identity. These two sets of identities generate the same subquotient S_4 -module of $\text{BW}(4)$, and hence the two operads are isomorphic.

6.4. Relations of arity ≥ 5 : a result and two conjectures.

Proposition 6.8. Every multilinear polynomial identity of arity ≤ 6 relating the Jordan and pre-Jordan products in every tridendriform algebra is a consequence of the defining identities for post-Jordan algebras.

Proof. Similar to the proof of Proposition 4.22. \square

Conjecture 6.9. Over a field \mathbb{F} of characteristic 0 or $p > 7$, every multilinear polynomial identity of arity ≤ 7 satisfied by the Jordan and pre-Jordan products in every tridendriform algebra is a consequence of the defining identities for post-Jordan algebras.

Conjecture 6.10. There exist identities of arity 8 which are not consequences of the defining identities for post-Jordan algebras, the Glennie identities for Jordan algebras [28], and the special identities of arity 8 for pre-Jordan algebras [12].

7. CONCLUDING REMARKS

7.1. Koszul duality for nonquadratic operads. Koszul duality for associative algebras has been extended [23] from the quadratic case ($n = 2$) to the n -homogeneous case ($n > 2$). If a similar extension exists of Koszul duality for operads, it could be

applied to the operads governing the Jordan structures studied in this paper. There is another approach: introduce a new ternary operation to lower the weight of the relations. Consider an operad generated by a commutative binary operation ab . If we define a new ternary operation by $(a, b, c) = (ab)c$ then every binary monomial of weight 2 equals a ternary monomial of weight 1, and every binary monomial of weight 3 equals a binary-ternary monomial of weight 2. Any cubic relation in the binary operation can be rewritten as a quadratic relation in both operations. For example, consider the linearized Jordan identity stated at the start of §4.4. We replace monomials of the forms $((wx)y)z$ and $(wx)(yz)$ by (wx, y, z) and (w, x, yz) respectively:

$$(ab, d, c) + (ac, d, b) + (bc, d, a) - (a, b, cd) - (a, c, bd) - (b, c, ad) \equiv 0.$$

This relation is quadratic: each term involves one binary operation and one ternary operation. However, we must also include the relation $(a, b, c) - (ab)c \equiv 0$, which is homogeneous in the arity, but inhomogeneous in the weight. To go further, we need the theory of inhomogeneous Koszul duality for operads [44]. The Koszul dual cooperad will be a differential graded cooperad with a nonzero differential.

7.2. A commutative diagram: isomorphism of reconfigured operads. A conjecture relating the polynomial identities produced by the KP algorithm (duplicators) to those satisfied by the operations produced by the BSO algorithm (diproduts) was stated in [11]; a year later the conjecture was reformulated and proved by [36]. Those papers deal exclusively with dialgebras, duplicators, and diproduts; there should be a generalization to trialgebras, and then to pre-algebras and post-algebras. The generalized conjecture would state, roughly speaking, that Table 1 is a commutative diagram of morphisms between operads. We attempt to state the generalization as precisely as possible.

Conjecture 7.1. The following diagram of morphisms between operads commutes:

$$(14) \quad \begin{array}{ccccc} \mathcal{P} & \xrightarrow{\Omega} & \mathcal{P}_\Omega & \xrightarrow{\cong} & \mathcal{P}'_{\Omega'} \\ \downarrow \frown & \Downarrow \frown & \downarrow \frown & & \downarrow \frown \\ \widehat{\mathcal{P}} & \xrightarrow{\widehat{\Omega}} & \widehat{\mathcal{P}}_{\widehat{\Omega}} & \xrightarrow{\cong} & \widehat{\mathcal{P}}'_{\widehat{\Omega}'} \stackrel{?}{=} \widehat{\mathcal{P}}'_{\Omega'} \end{array}$$

The symbols in this diagram are defined as follows:

- (1) We start by considering operads defined in terms of operations in an operad \mathcal{P} .
 - (a) Let \mathcal{P} be symmetric, not necessarily binary or quadratic, and in the category \mathbf{Vect}_F .
 - (b) Let \mathbf{A} be the category of \mathcal{P} -algebras.
 - (c) Let $\omega_i \in \mathcal{P}(n_i)$ for $1 \leq i \leq m$ be (multilinear) operations in \mathcal{P} , which we regard as new operations defined on the underlying vector spaces of the \mathcal{P} -algebras in \mathbf{A} .
 - (d) For example, if $m = 1, n_1 = 2$ then ω_1 could be the Lie bracket $ab - ba$ or the Jordan product $ab + ba$, both of which have (skew-)symmetry.
 - (e) Let $\Omega = \{\omega_1, \dots, \omega_m\}$ and let \mathcal{P}_Ω be the suboperad of \mathcal{P} generated by Ω .
 - (f) Let R be the set of all relations satisfied by the operations Ω in \mathcal{P} . That is, R consists of the polynomial identities satisfied by the operations Ω in every \mathcal{P} -algebra.

- (g) To separate \mathcal{P}_Ω from its embedding into \mathcal{P} , we make a copy, $\Omega' = \{\omega'_1, \dots, \omega'_m\}$: symbols representing abstract multilinear operations of arities n_1, \dots, n_m .
 - (h) By definition, the operad $\mathcal{Q} = \mathcal{P}'_{\Omega'}$ has generators Ω' with the same symmetries as those of Ω , satisfying relations R' which are copies of the relations R .
 - (i) For example, if $m = 1, n_1 = 2, \omega_1 = ab - ba$ then $\mathcal{Q} = \mathcal{P}'_{\Omega'}$ is the Lie operad, since every identity satisfied by the Lie bracket in every associative algebra is a consequence of anticommutativity and the Jacobi identity.
 - (j) On the other hand, if $m = 1, n_1 = 2, \omega_1 = ab + ba$ then $\mathcal{Q} = \mathcal{P}'_{\Omega'}$ is not the Jordan operad: the Glennie identities are satisfied by the Jordan product in every associative algebra but are not consequences of commutativity and the Jordan identity.
 - (k) For an integer $d \geq 1$, the operad $\mathcal{Q}_d = \mathcal{P}'_{\Omega',d}$ is defined as $\mathcal{Q} = \mathcal{P}'_{\Omega'}$ except that R' includes only (copies of) the relations in R of arity $\leq d$.
 - (l) If $m = 1, n_1 = 2, \omega_1 = ab + ba, 4 \leq d \leq 7$ then $\mathcal{Q}_d = \mathcal{P}'_{\Omega',d}$ is the Jordan operad.
- (2) Next, we assume that we have an algorithm which takes as input an operad \mathcal{P} , and produces as output a “reconfigured” operad $\widehat{\mathcal{P}}$. For example, the KP algorithm produces, from a given category of algebras, the corresponding category of dialgebras. In terms of operads, this is the Manin white product with Perm .
- (3) Finally, we assume that we have a corresponding algorithm for operations, also denoted by hat, which constructs $k(i)$ “reconfigured” operations $\widehat{\omega_{i,j_1}}, \dots, \widehat{\omega_{i,j_k(i)}}$ from each ω_i . For simplicity, we assume that $k(i)$ equals the arity of ω_i . For example, if ω_i is the Lie bracket then the BSO algorithm produces two “reconfigured” operations, the left and right Leibniz products, $\omega_{i,1} = a \dashv b - b \dashv a$ and $\omega_{i,2} = a \dashv b - b \dashv a$.

The conjecture amounts briefly to the statement that the “hat” and the “prime” commute.

REFERENCES

- [1] M. AGUIAR: Pre-Poisson algebras. *Lett. Math. Phys.* 54 (2000), no. 4, 263–277.
- [2] M. AYMÓN, P.-P. GRIVEL: Un théorème de Poincaré-Birkhoff-Witt pour les algèbres de Leibniz. *Comm. Algebra* 31 (2003), no. 2, 527–544.
- [3] C. BAI, O. BELLIER, L. GUO, X. NI: Splitting of operations, Manin products, and Rota-Baxter operators. arxiv.org/abs/1106.6080 (submitted 29 June 2011). *Int. Math. Res. Not. IMRN* 2013, no. 3, 485–524.
- [4] R. BANDIERA: A trick to compute certain Manin products of operads. arxiv.org/abs/1602.08051 (submitted 25 February 2016).
- [5] N. BERGERON, J.-L. LODAY: The symmetric operation in a free pre-Lie algebra is magmatic. *Proc. Amer. Math. Soc.* 139 (2011), no. 5, 1585–1597.
- [6] A. BLOH: On a generalization of the concept of Lie algebra. *Dokl. Akad. Nauk SSSR* 165 (1965) 471–473.
- [7] L. A. BOKUT, Y. CHEN, C. LIU: Gröbner-Shirshov bases for dialgebras. *Internat. J. Algebra Comput.* 20 (2010), no. 3, 391–415.
- [8] S. BONDARI: Constructing the polynomial identities and central identities of degree < 9 of 3×3 matrices. *Linear Algebra Appl.* 258 (1997) 233–249.
- [9] M. R. BREMNER: On the definition of quasi-Jordan algebra. *Comm. Algebra* 38 (2010), no. 12, 4695–4704.
- [10] M. R. BREMNER, V. DOTSENKO: *Algebraic Operads: An Algorithmic Companion*. CRC Press, Boca Raton, 2016.

- [11] M. R. BREMNER, R. FELIPE, J. SÁNCHEZ-ORTEGA: Jordan triple disystems. *Comput. Math. Appl.* 63 (2012), no. 6, 1039–1055.
- [12] M. R. BREMNER, S. MADARIAGA: Special identities for the pre-Jordan product in the free dendriform algebra. *Linear Algebra Appl.* 439 (2013), no. 2, 435–454.
- [13] M. R. BREMNER, S. MADARIAGA, L. A. PERESI: Structure theory for the group algebra of the symmetric group, with applications to polynomial identities for the octonions. arXiv: 1407.3810 (submitted 14 July 2014)
- [14] M. R. BREMNER, J. SÁNCHEZ-ORTEGA: Self-dual nonsymmetric operads with two binary operations. arxiv.org/abs/1606.01982 (submitted 7 June 2016).
- [15] D. BURDE: Left-symmetric algebras, or pre-Lie algebras in geometry and physics. *Cent. Eur. J. Math.* 4 (2006), no. 3, 323–357.
- [16] F. CHAPOTON: Un endofoncteur de la catégorie des opérades. *Dialgebras and Related Operads*, pages 105–110. Lecture Notes in Math., 1763. Springer, Berlin, 2001.
- [17] F. CHAPOTON: On some anticyclic operads. *Algebr. Geom. Topol.* 5 (2005) 53–69.
- [18] F. CHAPOTON, M. LIVERNET: Pre-Lie algebras and the rooted trees operad. *Int. Math. Res. Not. IMRN* (2001), no. 8, 395–408.
- [19] J. M. CLIFTON: A simplification of the computation of the natural representation of the symmetric group S_n . *Proc. Amer. Math. Soc.* 83 (1981), no. 2, 248–250.
- [20] P. M. COHN: On homomorphic images of special Jordan algebras. *Canadian J. Math.* 6 (1954) 253–264.
- [21] V. DOTSENKO, J. GRIFFIN: Cacti and filtered distributive laws. arXiv:1109.5345 (submitted on 25 September 2011). *Algebr. Geom. Topol.* 14 (2014), no. 6, 3185–3225.
- [22] V. DOTSENKO, A. KHOROSHKIN: Gröbner bases for operads. *Duke Math. J.* 153 (2010), no. 2, 363–396.
- [23] V. DOTSENKO, B. VALLETTE: Higher Koszul duality for associative algebras. *Glasg. Math. J.* 55 (2013), no. A, 55–74.
- [24] K. EBRAHIMI-FARD: Loday-type algebras and the Rota-Baxter relation. *Lett. Math. Phys.* 61 (2002), no. 2, 139–147.
- [25] I. M. H. ETHERINGTON: Non-associate powers and a functional equation. *Math. Gaz.*, 21, 242 (Feb. 1937) 36–39.
- [26] M. GERSTENHABER: The cohomology structure of an associative ring. *Ann. of Math.* (2) 78 (1963) 267–288.
- [27] V. GINZBURG, M. KAPRANOV: Koszul duality for operads. *Duke Math. J.* 76 (1994), no. 1, 203–272. Erratum: *Duke Math. J.* 80 (1995), no. 1, 293.
- [28] C. M. GLENNIE: Some identities valid in special Jordan algebras but not valid in all Jordan algebras. *Pacific J. Math.* 16 (1966) 47–59.
- [29] F. GOICHOT: Un théorème de Milnor-Moore pour les algèbres de Leibniz. *Dialgebras and Related Operads*, pages 111–133. Lecture Notes in Math., 1763. Springer, Berlin, 2001.
- [30] V. GUBAREV, P. S. KOLESNIKOV: Embedding of dendriform algebras into Rota-Baxter algebras. *Cent. Eur. J. Math.* 11 (2013), no. 2, 226–245.
- [31] V. GUBAREV, P. S. KOLESNIKOV: On the computation of Manin products for operads. arxiv.org/abs/1204.0894 (submitted 4 Apr 2012).
- [32] I. R. HENTZEL: Processing identities by group representation. *Computers in Nonassociative Rings and Algebras*, (Special Session, 82nd Annual Meeting, Amer. Math. Soc., San Antonio, Texas, 1976), pages 13–40. Academic Press, New York, 1977.
- [33] D. HOU, X. NI, C. BAI: Pre-Jordan algebras. *Math. Scand.* 112 (2013), no. 1, 19–48.
- [34] P. S. KOLESNIKOV: Varieties of dialgebras, and conformal algebras. *Sibirsk. Mat. Zh.* 49 (2008), no. 2, 322–339; translation in *Sib. Math. J.* 49 (2008), no. 2, 257–272.
- [35] P. S. KOLESNIKOV: Algebras with operators, Koszul duality, and conformal algebras. *Third Mile High Conference on Nonassociative Mathematics*, 11-17 August 2013, University of Denver, Colorado, USA. Slides: www.cs.du.edu/~petr/milehigh/2013/Kolesnikov.pdf
- [36] P. S. KOLESNIKOV, V. Y. VORONIN: On special identities for dialgebras. *Linear Multilinear Algebra* 61 (2013), no. 3, 377–391.
- [37] J.-L. LODAY: Une version non commutative des algèbres de Lie: les algèbres de Leibniz. *Enseign. Math.* (2) 39 (1993), no. 3-4, 269–293.
- [38] J.-L. LODAY: Algèbres ayant deux opérations associatives (digèbres). *C. R. Acad. Sci. Paris Sér. I Math.* 321 (1995), no. 2, 141–146.
- [39] J.-L. LODAY: Cup-product for Leibniz cohomology and dual Leibniz algebras. *Math. Scand.* 77 (1995), no. 2, 189–196.

- [40] J.-L. LODAY: Dialgebras. *Dialgebras and Related Operads*, pages 7–66. Lecture Notes in Math., 1763. Springer, Berlin, 2001.
- [41] J.-L. LODAY: Generalized bialgebras and triples of operads. *Astérisque* 320 (2008). [arXiv:math/0611885](https://arxiv.org/abs/math/0611885) (submitted 28 November 2006).
- [42] J.-L. LODAY, M. O. RONCO: Hopf algebra of the planar binary trees. *Adv. Math.* 139 (1998), no. 2, 293–309.
- [43] J.-L. LODAY, M. O. RONCO: Trialgebras and families of polytopes. *Homotopy Theory: Relations with Algebraic Geometry, Group Cohomology, and Algebraic K-Theory*, pages 369–398. Contemp. Math., 346. Amer. Math. Soc., Providence, 2004.
- [44] J.-L. LODAY, B. VALLETTE: *Algebraic Operads*. Grundlehren Math. Wiss., 346. Springer, Heidelberg, 2012.
- [45] S. MADARIAGA: Gröbner-Shirshov bases for the non-symmetric operads of dendriform algebras and quadri-algebras. *J. Symbolic Comput.* 60 (2014) 1–14.
- [46] X. NI: *Rota-Baxter Algebras, Renormalization on Kausz Compactifications, and Replicating of Binary Operads*. Ph.D. Thesis, California Institute of Technology, Pasadena, California, USA. Defended 19 May 2016.
- [47] J. PEI, C. BAI, L. GUO, X. NI: Replicating of binary operads, Koszul duality, Manin products and average operators. [arXiv:1212.0177](https://arxiv.org/abs/1212.0177) (submitted 2 December 2012).
- [48] D. E. RUTHERFORD: *Substitutional Analysis*. Edinburgh, at the University Press, 1948.
- [49] K. UCHINO: Derived bracket construction and Manin products. *Lett. Math. Phys.* 93 (2010), no. 1, 37–53.
- [50] B. VALLETTE: Homology of generalized partition posets. *J. Pure Appl. Algebra* 208 (2007), no. 2, 699–725.
- [51] B. VALLETTE: Manin products, Koszul duality, Loday algebras and Deligne conjecture. *J. Reine Angew. Math.* 620 (2008), 105–164.
- [52] R. VELÁSQUEZ, R. FELIPE: Quasi-Jordan algebras. *Comm. Algebra* 36 (2008), no. 4, 1580–1602.
- [53] E. B. VINBERG: Convex homogeneous domains. *Dokl. Akad. Nauk SSSR* 141 (1961) 521–524.
- [54] J. H. M. WEDDERBURN: The functional equation $g(x^2) = 2ax + [g(x)]^2$. *Ann. of Math. (2)* 24, 2 (Dec. 1922), 121–140.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, CANADA
E-mail address: bagherzadeh@math.usask.ca

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, CANADA
E-mail address: bremner@math.usask.ca

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF LA RIOJA, SPAIN
E-mail address: sara.madariaga@unirioja.es