Lattice paths with catastrophes

Cyril Banderier^{1*}

Michael Wallner^{2†}

July 10, 2017

¹: LIPN UMR-CNRS 7030, Université de Paris Nord, France. ²: TU Wien, Austria.

Abstract

In queuing theory, it is usual to have some models with a "reset" of the queue. In terms of lattice paths, it is like having the possibility of jumping from any altitude to zero. These objects have the interesting feature that they do not have the same intuitive probabilistic behaviour like classical Dyck paths (the typical properties of which are strongly related to Brownian motion theory), and this article quantifies some relations between these two types of paths. We give a bijection with some other lattice paths and a link with a continued fraction expansion, and prove several formulae for related combinatorial structures conjectured in the On-line Encyclopedia of Integer Sequences. Thanks to the kernel method and via analytic combinatorics, we derive the enumeration and limit laws of these "lattice paths with catastrophes", for any finite set of jumps. We end with an algorithm to generate such lattice paths uniformly at random.

Keywords: Lattice path, generating function, algebraic function, kernel method, context-free grammar, random generation

^{*}http://lipn.fr/~banderier

[†]http://dmg.tuwien.ac.at/mwallner/

1 Introduction

Lattice paths are a natural model in queuing theory: indeed, the evolution of a queue can be seen as a sum of jumps, a subject e.g. considered in Feller (1968). In this article we consider jumps restricted to a given finite set of integers J, where each jump $j \in J$ is associated with a weight (or probability) p_j . The evolution of a queue naturally corresponds to lattice paths constrained to be non-negative. For example, if $J = \{-1, +1\}$, this corresponds to the so-called Dyck paths. Moreover, we also consider the model where "catastrophes" are allowed.

Definition 1.1. A *catastrophe* is a jump from an altitude j > 0 ($j \notin J$) to altitude 0, see Figure 1.

Such a jump corresponds to a "reset" of the queue. The model of queues with catastrophes was e.g. considered in Krinik, Rubino, Marcus, Swift, Kasfy, and Lam (2005) or Hunter, Krinik, Nguyen, Switkes, and Von Bremen (2008). In financial mathematics, this also gives a natural, simple model of the evolution of stock markets, allowing bankruptcies at any time with a small probability q.




Figure 1: Decomposition of a Dyck path with 3 catastrophes into 5 arches. A_{cat} stands for an "arch ending with a catastrophe" (a walk for which the first return to altitude 0 is a catastrophe), while A_{nocat} stands for an "arch with no catastrophe".

Link with a continued fraction expansion. We first start with the observation that the generating function of these lattice paths has the following continued fraction expansion:

$$H(z) = \sum_{n \ge 0} h_n z^n = \frac{1}{1 - \frac{z^2}{1 - z - \frac{z^2}{1 - \frac{$$

We give two proofs of this phenomenon in Theorem 3.1. In this article, we also tackle the question of what happens for more general jumps than Dyck paths, and we give the enumeration and asymptotics of the corresponding lattice path models.

Link with generating trees. In combinatorics, such lattice paths are related to generating trees, which are a convenient tool to enumerate and generate many combinatorial structures in some incremental way (like e.g. permutations avoiding some pattern), see e.g. West (1996). In such trees, the distribution of the children of each node follows exactly the same dynamics as lattice paths with some "extended" jumps, as was intensively investigated by the Florentine school of combinatorics, e.g. in Barcucci, Del Lungo, Pergola, and Pinzani (1999), Duchi, Fédou, and Rinaldi (2004), Ferrari, Pergola, Pinzani, and Rinaldi (2011). For example, these "extended" jumps can be a continuous set of jumps: from altitude k, one can jump to any altitude between 0 and k, possibly with some weights, plus a finite set of bounded jumps. This model can be seen as an intermediate model between Dyck paths and our lattice paths with catastrophes; we investigated it in our series of articles Banderier, Bousquet-Mélou, Denise, Flajolet, Gardy, and Gouyou-Beauchamps (2002), Banderier (2002), Banderier and Merlini (2003), Banderier, Fédou, Garcia, and Merlini (2003). In this article, we will however see that several statistics of lattice paths with catastrophes behave in a rather different way than these walks with a continuous set of jumps, even if they share the following unusual property: both of them correspond to random walks with an "infinite negative drift" (in fact, a space-dependent drift tending to $-\infty$ when the altitude increases). At the same time, they are constrained to remain at non-negative altitudes; this leads to some counter-intuitive behaviour: unlike classical directed lattice paths, the limiting object is no more directly related to Brownian motion theory.

Enumeration and asymptotics: why context-free grammars would be a wrong idea here. One way to analyse our lattice paths could be to use a context-free grammar approach, see Labelle and Yeh (1990): this leads to a system of algebraic equations, and therefore we already know "for free" that the corresponding generating functions are algebraic. However, this system involves nearly $(c+d)^2$ equations (where c is the largest negative jump and d the largest positive jump), so solving it (with resultants or Gröbner bases) leads to computations taking a lot of time and memory (exponential in $(c+d)^2$): even for c = d = 10, the needed memory to compute the algebraic equation with this method would be more than the expected number of particles in the universe! Another drawback of this method is that it would be a "case-by-case" analysis: for each new set of jumps, one would have to do new computations from scratch. Hence, with this method, there is no way to access "universal" asymptotic results: while it is well known that the coefficients of algebraic functions exhibit an asymptotic behaviour of the type $f_n \sim C.A^n n^{\alpha}$, only the "critical exponent" α can be proven to belong to a specific set (see Banderier and Drmota (2015)), and there is no hope to have easy access to C and A with this context-free grammar approach, in a way which is independent of a case-by-case computation (which, what is more, would be impossible for c + d > 20).

The solution: kernel method and analytic combinatorics. In this article, we offer an alternative to context-free grammars. Our approach uses methods of analytic combinatorics for directed lattice paths: the kernel method and singularity analysis, as presented in Banderier and Flajolet (2002), Flajolet and Sedgewick (2009). It allows to get exact enumeration, the typical behaviour of lattice paths with catastrophes, and has the advantage of offering universal results for the asymptotics and generic closed forms, whatever the set of jumps is.

Plan of this article. First, in Section 2, we present the model of walks with catastrophes and derive their generating functions. In Section 3, we establish a bijection between two generalizations of Dyck paths. In Section 4, we analyse our model in more detail and first derive the asymptotic number of excursions and meanders. Then we use these results to obtain limit laws for the number of catastrophes, the number of returns to zero, the final altitude, the total amplitude of catastrophes, the average height of a catastrophe, and the waiting time for the first catastrophe (see Figures 2 and 3). In Section 5, we discuss the uniform random generation of such lattice paths. In Section 6, we state a summary of our results.




Figure 2: In this article, we analyse the number of Dyck paths with catastrophes, the waiting time for the first catastrophe, its amplitude (here, the -6 jump, in red), the number of returns to 0 (here, 5), the number of catastrophes (here, 3, in red), the total sum of their amplitudes (here, 6 + 7 + 4), and therefore the average amplitude of a catastrophe.




Figure 3: A Dyck path with catastrophes drawn uniformly at random (amongst excursions of length n = 400). For this example, the walk has 35 returns to zero, of which 22 are catastrophes. The waiting time for the first catastrophe is 8, its amplitude is 5, the sum of the amplitude of all catastrophes is 124 (therefore an average catastrophe has amplitude 5.63). For all these parameters, this article shows how these quantities evolve when n gets larger. More generally, we give the corresponding limit laws for walks with catastrophes allowing any finite set of jumps.

2 Generating functions

In this section, we give some explicit formulae for the generating functions of non-negative lattice paths with catastrophes, for which the set of jumps is encoded by

$$P(u) = \sum_{i=-c}^{d} p_i u^i.$$

Every catastrophe is also assigned a weight q > 0.

The weights p_j and q are real and non-negative: in fact, even if they would be taken in \mathbb{C} , all our enumerative formulae would remain valid (also if $d = +\infty$). The non-negativity only plays a role for establishing the universal asymptotic phenomena presented in Section 4.

The generating functions of directed lattice paths can be expressed in terms of the small branches $u_i(z)$, i = 1, ..., c, which are solutions of the *kernel equation*

$$1 - zP(u_i(z)) = 0. (1)$$

In general this equation has c + d solutions. The *small branches* are the c solutions with the property $u_i(z) \sim 0$ for $z \sim 0$. The remaining d solutions are called *large branches* as they satisfy $|v_i(z)| \sim +\infty$ for $z \sim 0$. The generating functions of four classical types of directed lattice paths are shown in Table 1.




Table 1: The four types of paths: walks, bridges, meanders and excursions, and the corresponding generating functions for directed lattice paths, from Banderier and Flajolet (2002, Fig. 1).

These results follow from the expression for the bivariate generating function M(z, u) of meanders: Let $m_{n,k}$ be the number of meanders of length n going from altitude 0 to altitude k, then

$$M(z,u) = \sum_{n,k\geq 0} m_{n,k} z^n u^k = \sum_{k\geq 0} M_k(z) u^k = \frac{\prod_{i=1}^c (u - u_i(z))}{u^c (1 - zP(u))}.$$
(2)

This formula is obtained by the kernel method, which mainly consists in setting $u = u_i(z)$ in the functional equation which mimics the recursive definition of a meander. This results in new and simpler equations which lead to the closed form (2). The generating function of excursions is E(z) := M(z, 0).

Let us now investigate which perturbation is introduced by allowing catastrophes in this model. First, we partition the jump set $J = J_+ \cup J_- \cup J_0$ into the set of positive jumps ($j \in J_+$ iff j > 0), the set of negative jumps ($j \in J_-$ iff j < 0), and the possible zero jump ($j \in J_0$ iff j = 0).

Theorem 2.1 (Generating functions for lattice paths with catastrophes). Let $f_{n,k}$ be the number of meanders with catastrophes of length n from altitude 0 to altitude k. Then the generating function $F(z, u) = \sum_{k\geq 0} F_k(z)u^k = \sum_{n,k\geq 0} f_{n,k}u^k z^n$ is algebraic and satisfies

$$F(z,u) = D(z)M(z,u) = D(z)\frac{\prod_{i=1}^{c}(u-u_i(z))}{u^c(1-zP(u))},$$
(3)

$$F_k(z) = D(z)M_k(z) = D(z)\frac{1}{p_d z} \sum_{\ell=1}^d v_\ell^{-k-1} \prod_{j \neq \ell} \frac{1}{v_j - v_\ell}, \quad \text{for } k \ge 0,$$
(4)

where $D(z) = \frac{1}{1-Q(z)}$ is the generating function of excursions ending with a catastrophe, $Q(z) = zq \left(M(z) - E(z) - \sum_{-j \in J_{-}} M_j(z)\right)$, and where, for any set of jumps encoded by P(u), the u_i 's and the v_i 's are the small roots and the large roots of the kernel equation (1).

Proof. Take an arbitrary non-negative path of length n. Let ω_0 be the last time it returns to the *x*-axis with a catastrophe (or $\omega_0 \coloneqq 0$ if the path contains no catastrophe). This point gives a unique decomposition into an initial excursion which ends with a catastrophe (this might be empty), and a meander without any catastrophes. This directly gives (3).

What remains is to describe the initial part D(z). Consider an arbitrary excursion ending with a catastrophe. We decompose it with respect to its catastrophes, into a sequence of minimal excursions with catastrophes which we count by Q(z). Such paths have only one catastrophe at their very ends and none before. Thus,

$$D(z) = \frac{1}{1 - Q(z)}.$$
(5)

Because of Definition 1.1 of a catastrophe, Q(z) is given by the generating function of meanders that are neither excursions nor meanders ending at altitudes |j| $(j \in J_{-})$ followed by a final catastrophe. This implies the shape of $Q(z) = zq \left(M(z) - E(z) - \sum_{-j \in J_{-}} M_{j}(z)\right)$.

Technically, D(z), the generating function of excursions ending with a catastrophe, acts like a prefix on directed lattice paths.

Remark 2.2. Our results obviously depend on the choice of Definition 1.1 of catastrophes. Some slightly different definitions could be used without changing their structure. For example, one could consider allowing catastrophes from any altitude (in this case, Q(z) = zqM(z)). In order to ensure an easy adaptation to different models, we will state all our subsequent results in terms of a generic Q(z).

Let us now consider a famous class of lattice paths (see e.g. Stanley (2011)).

Definition 2.3. A Dyck meander is a path constructed from the possible jumps +1 and -1, each with weight 1, and being constrained to stay weakly above the x-axis. A (classical) Dyck path or also called a Dyck excursion is additional constrained to return to the x-axis. Accordingly, the polynomial encoding the allowed jumps is $P(u) = u^{-1} + u$.

For these paths, when one also allows catastrophes of weight q = 1, one gets the following generating functions.

Corollary 2.4 (Generating functions for Dyck paths with catastrophes). The generating function of Dyck meanders with catastrophes, $F(z, 1) = \sum_{n>0} m_n z^n$, satisfies

$$F(z,1) = \frac{z(u_1(z)-1)}{z^2 + (z^2 + z - 1)u_1(z)} = 1 + z + 2z^2 + 4z^3 + 8z^4 + 17z^5 + 35z^6 + O(z^7),$$

where the root $u_1(z)$ of the kernel is in fact the generating functions of Catalan numbers: $u_1(z) = \frac{1-\sqrt{1-4z^2}}{2}$. Moreover, m_n is also the number of equivalence classes of Dyck excursions of length 2n + 2 for the pattern duu, see OEIS A274115¹.

The generating function of Dyck excursions with catastrophes, $F_0(z) = \sum_{n>0} e_n z^n$, is

$$F_0(z) = \frac{(2z-1)u_1(z)}{z^2 + (z^2 + z - 1)u_1(z)} = 1 + z^2 + z^3 + 3z^4 + 5z^5 + 12z^6 + 23z^7 + O(z^8).$$

This sequence e_n corresponds to OEIS A224747. Moreover, e_{2n} is also the number of Dumont permutations of the first kind of length 2n avoiding the patterns 1423 and 4132, see OEIS A125187.

Proof. The formulae for F(z,1) and $F_0(z)$ are a direct application of Theorem 2.1. Then, one notes that $(F_0(z) + F_0(-z))/2$ equals the generating function of Dumont permutations of the first kind of length 2n avoiding the patterns 1423 and 4132, see Burstein (2005) for the definition of such permutations, and the derivation of their generating function. In Manes, Sapounakis, Tasoulas, and Tsikouras (2016), two Dyck excursions are said equivalent if the occurrences of the pattern duu are at the same place. They derived the generating function for the number of equivalence classes, which appears to be equal to F(z, 1).


In the next section we will analyse Dyck paths with catastrophes in more detail. On the way we solve some conjectures of the On-Line-Encyclopedia of Integer Sequences.

¹Such references are links to the web-page dedicated to the corresponding sequence in the On-Line Encyclopedia of Integer Sequences, http://oeis.org.

3 Bijection for Dyck paths with catastrophes

The goal of this section is to establish a bijection between two classes of extensions of Dyck paths². We consider two extensions of classical Dyck paths (see Figure 4 for an illustration):

- 1. Dyck paths with catastrophes are Dyck paths with the additional option of jumping to the x-axis from any altitude h > 1; and
- 2. 1-horizontal Dyck paths are Dyck paths with the additional allowed horizontal step (1,0) at altitude 1.



(a) Dyck arch ending with a catastrophe



Figure 4: The bijection of Theorem 3.1 transforming Dyck arches ending with catastrophes into 1-horizontal Dyck arches, and vice versa.

Theorem 3.1 (Bijection for Dyck paths with catastrophes). The number e_n of Dyck paths with catastrophes of length n is equal to the number h_n of 1-horizontal Dyck paths of length n: $e_n = h_n$.

Proof. A first proof that $e_n = h_n$ consists of using the continued fraction approach (each level k + 1 of the continued fraction encodes the jumps allowed at altitude k):

$$H(z) = \sum_{n \ge 0} h_n z^n = \frac{1}{1 - \frac{z^2}{1 - z - z^2 C(z)}},$$

where C(z) is the generating function of classical Dyck paths, $C(z) = 1/(1 - z^2C(z))$. One then gets that H(z) equals the closed form of $F_0(z)$ given in Corollary 2.4. We now give a bijective procedure which transforms every Dyck path with catastrophes into a 1-horizontal Dyck path, and vice versa.

Every Dyck path with catastrophes can be decomposed into a sequence of "minimal" excursions, in the sense that their only contact with the x-axis is at the starting and end point. Such paths are called *arches*, see Figure 1. There are two types of arches: arches ending with

 $^{^2} We$ thus prove several conjectures by Alois P. Heinz, R. J. Mathar, and other contributors in the On-Line-Encyclopedia of Integer Sequences, see sequences A224747 and A125187 therein.

catastrophes $A_{\text{cat}}(z)$ and arches ending with a jump $j \in J$ given by $A_{\text{nocat}}(z)$. This gives the alternative decomposition to (4) of the kind (compare also Figure 1)

$$F_0(z) = \frac{1}{1 - (A_{\text{cat}}(z) + A_{\text{nocat}}(z))}.$$

Thus, without loss of generality, we continue our discussion only for arches. The following procedure is visualized in Figure 4.

Let us start with an arbitrary arch of Dyck paths with catastrophes. It is either a classical Dyck path, and therefore also a 1-horizontal Dyck path, or it ends with a catastrophe of height h. First, we associated the catastrophe with h up-steps (1,1). Specifically, we draw horizontal lines to the left, until we hit an up-step. All but the first one are replaced by horizontal steps. Finally, we replace the catastrophe by a down step (1,-1). All parts in between stay the same. Note, that we replaced h-1 up-steps, and therefore lost a height of h-1, but we also replaced the catastrophe of height h by a down-step, which represents a gain of height by h-1. Thus, we again return to the x-axis. Furthermore, all horizontal steps are at altitude 1. Thus, we always stay weakly above the x-axis, and we got an arch of a 1-horizontal Dyck path. The inverse mapping is analogous.

The most important building blocks in the previous bijection were arches ending with a catastrophe. These can be made even more explicit.

Proposition 3.2 (Dyck arches ending with a catastrophe). Let $A(z) = \sum_{n\geq 0} a_n z^n$ be the generating function of arches ending with a catastrophe. Then, one has the following closed forms

$$a_n = \binom{n-2}{\lfloor \frac{n-3}{2} \rfloor},$$

$$A_{\text{cat}}(z) = z \frac{M(z) - E(z) - M_1(z)}{E(z)} = \frac{1}{2} \frac{2z^2 + z - 1 + \sqrt{(1-2z)(1+2z)(1-z)^2}}{1-2z}$$

$$= z^3 + z^4 + 3z^5 + 4z^6 + 10z^7 + 15z^8 + 35z^9 + O(z^{10}),$$

where M(z), E(z) and $M_1(z)$ are the generating functions of classical Dyck paths for meanders, excursions, and meanders ending at 1, respectively, see OEIS A037952.

Proof. Every excursion ending with a catastrophe can be uniquely decomposed into an initial excursion and a final arch with a catastrophe. By Theorem 2.1 we get the generating function of $A_{\text{cat}}(z) = \frac{Q(z)}{E(z)}$.

In order to compute a_n , additionally drop the initial +1-jump which is necessary for all such arches of positive length. The remaining part is a Dyck meander (always staying weakly above the *x*-axis) that does not end on the *x*-axis. Thus,

$$a_{n+2} = \underbrace{\binom{n}{\left\lfloor \frac{n}{2} \right\rfloor}}_{\text{meanders}} - \underbrace{\frac{1}{n/2 + 1} \binom{n}{\frac{n}{2}}}_{\text{excursions}} \llbracket n \text{ even} \rrbracket,$$

where [[P]] denotes the lverson bracket, which is 1 if condition P is true, and 0 otherwise. \Box

4 Asymptotics and limit laws

The natural model in which all paths of length n have the same weight creates a probabilistic model in which the drift of the walk is then space-dependent: it converges to minus infinity when the altitude of the paths is increasing. So, unlike the easier classical Dyck paths (and their generalization via directed lattice paths, having a finite set of given jumps), we are losing the intuition offered by Brownian motion theory. This leads to the natural question of what the asymptotics of the fundamental parameters of our "lattice paths with catastrophes" are. This is the question we are going to answer now.

To this aim, some useful results of Banderier and Flajolet (2002) are the asymptotic enumeration formulae for the four types of paths shown in Table 1. A key result is the fact that the principal small branch $u_1(z)$ and the principal large branch $v_1(z)$ are conjugated to each other at their dominant singularity $\rho = \frac{1}{P(\tau)}$ where $\tau > 0$ is the minimal positive real solution of $P'(\tau) = 0$. In particular it holds that

$$u_1(z) = \tau - C\sqrt{1 - z/\rho} + O(1 - z/\rho),$$

$$v_1(z) = \tau + C\sqrt{1 - z/\rho} + O(1 - z/\rho),$$

with the constant $C \coloneqq \sqrt{2\frac{P(\tau)}{P''(\tau)}}$. This singularity ρ of $u_1(z)$ and $v_1(z)$ turns out to be the dominant singularity of the generating functions of directed lattice paths.

Definition 4.1. We say that a function F(z) has periodic support p or (for short) F(z) is *p*-periodic if there exists a function H(z) and an integer b such that $F(z) = z^b H(z^p)$. If this holds only for p = 1, the function is said to be *aperiodic*.

In Banderier and Wallner (2016, Lemma 8.7 and Theorem 8.8) we show how to deduce the asymptotics of walks having periodic jump polynomials from the results on aperiodic ones. Therefore, without loss of generality we consider only the aperiodic jump polynomials in this article.

4.1 Asymptotics of lattice paths

We start by analysing the function $D(z) = \frac{1}{1-Q(z)}$. In particular, we need to find its singularities, which are given by the behaviour of

$$Q(z) = zq\left(M(z) - E(z) - \sum_{-j \in J_{-}} M_j(z)\right).$$

Caveat: Even if we already know that the radii of convergence $\rho_M, \rho_E, \rho_{M_j}$ of M(z), E(z), $M_j(z)$, it is a priori not granted that Q(z) does not have a larger radius of convergence (some cancellations could occur). In fact, Banderier and Flajolet (2002) allows to prove via the asymptotics of the coefficients of M(z), E(z), and $M_j(z)$ that no such cancellations occur here. Therefore the radius of convergence of Q(z) is $\rho_Q = \min(\rho_M, \rho_E, \rho_{M_j})$.

We now determine the radius of convergence ρ_D of D(z).

Lemma 4.2. Let \mathcal{Z} be the set of zeros of 1 - Q(z) of minimal modulus with $|z| \leq \rho$. The set \mathcal{Z} is either empty or has exactly one real positive element which we call ρ_0 . The sign of the drift $\delta := P'(1)$ of the walk dictates the location of the radius of convergence ρ_D . If $\delta \geq 0$, we have $\rho_D = \rho_0 < \frac{1}{P(1)} \leq \rho$. While, if $\delta < 0$, it also depends on the value $Q(\rho)$:

$$\begin{cases} Q(\rho) > 1 & \Longleftrightarrow & \rho_D = \rho_0 < \rho, \\ Q(\rho) = 1 & \Longleftrightarrow & \rho_D = \rho_0 = \rho, \\ Q(\rho) < 1 & \Longleftrightarrow & \rho_D = \rho \text{ and } \mathcal{Z} \text{ is empty.} \end{cases}$$

Remark: In the remainder of this article we use the notational shorthand $\nexists \rho_0$ to denote the case \mathcal{Z} is empty.

Proof. As D(z) = 1/(1-Q(z)) is a generating function with positive coefficients, Pringsheim's Theorem implies that it has a dominant singularity on the real positive axis, which we call r. This singularity is either $r = \rho_Q$, the singularity of Q(z), or it is the smallest real positive zero of 1 - Q(z) (if it exists, it is denoted by ρ_0 and it is therefore such that $\rho_0 \le \rho_Q$).

What is more, D(z) has no other dominant singularity: When $r = \rho_Q$ this follows from the aperiodicity of E(z), M(z), and $M_j(z)$ proven in Banderier and Flajolet (2002). When $r = \rho_0$ this follows from the strong triangle inequality. Indeed, as Q(z) has non-negative coefficients and is aperiodic, one has 1 = |Q(z)| < Q(|z|) for any |z| = r, $z \neq r$.

Next, note that Q(z) consists of 3 different parts: the generating function of meanders M(z), the generating function of excursions E(z), and the sum of meanders ending at fixed altitudes $\sum_{-j \in J_-} M_j(z)$. The functions E(z) and $M_j(z)$ are analytic for $|z| < \rho$, but the behaviour of M(z) depends on the drift $\delta = P'(1)$. For $\delta \ge 0$ it possesses a simple pole at $\rho_1 \coloneqq \frac{1}{P(1)} \le \rho$, i.e. $\rho_Q = \rho_1$. Thus, $\lim_{z \to \rho_1^-} Q(z) = +\infty$, and together with Q(0) = 0 this implies that there is a solution $0 < \rho_0 < \rho_1$.

For $\delta < 0$ we have that |Q(z)| is bounded for $|z| < \rho$, and we have $\rho_Q = \rho$. Thus, for a fixed jump polynomial P(u) any case can be attained by a variation of q. As Q(z) is monotonically increasing on the real axis, it suffices to compare its value at its maximum $Q(\rho)$.

It remains to consider the lower bound in the case $Q(\rho) > 1$. Because of $u_1(\rho_1) = 1$ the singularity in the denominator is cancelled by the factor $1 - u_1(z)$. Due to the domination property $|u_i(z)| < |u_1(z)| < 1$ for $z \in (0, \rho)$ the remaining factors are strictly smaller than 1. A detailed discussion of this behaviour can be found in the proof of Banderier and Flajolet (2002, Theorem 4).

Note that Q(z) strongly depends on the weight of the catastrophes q > 0. Therefore, for a fixed step set P(u) with negative drift one can model any of the three possible cases by a proper choice of q.

Theorem 4.3. The asymptotics of excursions ending with a catastrophe d_n depends on the structural radius ρ and the possible singularity ρ_0 :

$$d_{n} = \begin{cases} \frac{\rho_{0}^{-n}}{\rho_{0}Q'(\rho_{0})} + o(K^{n}), & \rho_{0} < \rho \quad \text{(for some } K \text{ such } K < 1/\rho_{0}), \\ \frac{\rho^{-n}}{\eta\sqrt{\pi n}} \left(1 + O\left(\frac{1}{n}\right)\right), & \rho_{0} = \rho, \\ \frac{D(\rho)^{2}\eta\rho^{-n}}{2\sqrt{\pi n^{3}}} \left(1 + O\left(\frac{1}{n}\right)\right), & \nexists \rho_{0}, \end{cases}$$

where η is given by the Puiseux expansion of $Q(z) = Q(\rho) - \eta \sqrt{1 - z/\rho} + O(1 - z/\rho)$ for $z \to \rho$. The last two cases occur only when $\delta < 0$.

Proof. The critical exponent α in the Puiseux expansion of D(z) in $(1 - z/r)^{\alpha}$ for $r = \rho_0$ or $r = \rho$, respectively, satisfies

- $\alpha = -1$ for $\rho_0 < \rho$,
- $\alpha = -1/2$ for $\rho_0 = \rho$,
- $\alpha = 1/2$ for $\nexists \rho_0$.

Indeed, the singularity of D(z) arises at the minimum of ρ and ρ_0 , as derived in Lemma 4.2. In the first case $\rho_0 < \rho$, the singularity is a simple pole as the first derivative of the denominator at ρ_0 is strictly positive. We get for $z \rightarrow \rho_0$

$$1 - Q(z) = \underbrace{(1 - Q(\rho_0))}_{=0} + \rho_0 Q'(\rho_0) (1 - z/\rho_0) + O\left((1 - z/\rho_0)^2\right).$$
(6)

This yields a simple pole at ρ_0 for $\frac{1}{1-Q(z)}$ and singularity analysis then gives the asymptotics of d_n .

For ρ_0 = ρ or $\nexists \, \rho_0$ we get a square root behaviour for $z \to \rho$

$$1 - Q(z) = (1 - Q(\rho)) + \eta \sqrt{1 - z/\rho} + O(1 - z/\rho).$$
(7)

For $\rho_0 = \rho$ the constant term is 0, and we get for $z \rightarrow \rho$

$$D(z) = \frac{1}{\eta \sqrt{1 - z/\rho}} \left(1 + O(\sqrt{1 - z/\rho}) \right).$$
(8)

Yet, for $\nexists \rho_0$ the constant term does not vanish. This gives for $z \rightarrow \rho$

$$D(z) = D(\rho) - \eta D(\rho)^2 \sqrt{1 - z/\rho} + O(1 - z/\rho).$$
(9)

Applying singularity analysis yields the result.

With the help of the last result we are able to derive the asymptotics of lattice paths with catastrophes. Let us state the result for excursions next.

Theorem 4.4. The number of excursions with catastrophes e_n is asymptotically equal to

$$e_{n} = \begin{cases} \frac{E(\rho_{0})}{\rho_{0}Q'(\rho_{0})}\rho_{0}^{-n} + o(K^{n}) & \text{(for some } K \text{ such } K < 1/\rho_{0}) & \text{if } \rho_{0} < \rho_{1} \\ \frac{E(\rho)}{\eta}\frac{\rho^{-n}}{\sqrt{\pi n}}\left(1 + O\left(\frac{1}{n}\right)\right) & \text{if } \rho_{0} = \rho_{1} \\ \frac{F_{0}(\rho)}{2}\left(\sqrt{2\frac{P(\tau)}{P''(\tau)}}\frac{1}{\tau} + \eta D(\rho)\right)\frac{\rho^{-n}}{\sqrt{\pi n^{3}}}\left(1 + O\left(\frac{1}{n}\right)\right) & \text{if } \nexists \rho_{0}. \end{cases}$$

Proof. Since $F_0(z) = D(z)E(z)$, the singularity is either at ρ_0 or $\rho = \frac{1}{P(\tau)}$. Combining the results from Theorem 4.3 and Banderier and Flajolet (2002, Theorem 3) gives the result. Note that the cases $\rho_0 = \rho$ and $\nexists \rho_0$ are only possible for $\delta < 0$.

Next we also state the asymptotics for the number of meanders. The only difference is the appearance of M(z) instead of E(z), and a factor $\frac{1}{\tau-1}$ instead of $\frac{1}{\tau}$ in the first term when $\nexists \rho_0$.

Theorem 4.5. The number of meanders with catastrophes m_n is asymptotically equal to

$$m_{n} = \begin{cases} \frac{M(\rho_{0})}{\rho_{0}Q'(\rho_{0})}\rho_{0}^{-n} + o(K^{n}) & \text{(for some } K \text{ such } K < 1/\rho_{0}) & \text{if } \rho_{0} < \rho, \\ \frac{M(\rho)}{\eta}\frac{\rho^{-n}}{\sqrt{\pi n}}\left(1 + O\left(\frac{1}{n}\right)\right) & \text{if } \rho_{0} = \rho, \\ \frac{F(\rho, 1)}{2}\left(\sqrt{2\frac{P(\tau)}{P''(\tau)}}\frac{1}{\tau - 1} + \eta D(\rho)\right)\frac{\rho^{-n}}{\sqrt{\pi n^{3}}}\left(1 + O\left(\frac{1}{n}\right)\right) & \text{if } \not\equiv \rho_{0}. \end{cases}$$

Proof. Analogous to the proof of Theorem 4.4 the result follows after some tiresome computations from the fact that F(z,1) = D(z)M(z). Combining the results from Theorem 4.3 and Banderier and Flajolet (2002, Theorem 4) gives the result.

Remark 4.6. In the previous proofs we needed that P(u) is an aperiodic jump set. Otherwise, the generating function Q(z) does not have a unique singularity on its radius of convergence, but several. In such cases one needs to consider all singularities and sum their contributions; however this can lead to cancellations, thus extra care is necessary. A systematic approach of the periodic cases is treated in Banderier and Wallner (2016). These considerations about periodicity are only necessary when the dominant asymptotics come from the singularity ρ , while when $\rho_0 < \rho$, we have a unique dominant simple pole (the possibly periodic functions E(z) and $M_j(z)$ do not contribute to the asymptotics). This polar behaviour occurs e.g. for Dyck paths.

Corollary 4.7. The number of Dyck paths with catastrophes e_n , and Dyck meanders with catastrophes m_n is asymptotically equal to

$$e_n = C_e \rho_0^{-n} \left(1 + O\left(\frac{1}{n}\right) \right) \quad (C_e \approx 0.10381 \text{ is the unique positive root of } 31C_e^3 - 62C_e^2 + 35C_e - 3),$$

$$m_n = C_m \rho_0^{-n} \left(1 + O\left(\frac{1}{n}\right) \right) \quad (C_m \approx 0.32679 \text{ is the unique positive root of } 31C_m^3 - 31C_m^2 + 16C_m - 3),$$

where $\rho_0 \approx 0.46557$ is the unique positive root of $\rho_0^3 + 2\rho_0^2 + \rho_0 - 1$. Accordingly, $\mathbb{P}(\text{meander of length } n \text{ is an excursion}) = e_n/m_n \approx 0.31767$. *Proof.* We apply the results of Theorem 4.4. This directly gives

$$\rho_0 = \frac{1}{6} \left(116 + 12\sqrt{93} \right)^{1/3} + \frac{2}{3} \left(116 + 12\sqrt{93} \right)^{-1/3} - \frac{2}{3} \approx 0.46557,$$

which is strictly smaller than $\rho_0 = 1/2$. The defining equations for these algebraic numbers are computed by resultants. Finally, the asymptotics of meanders is computed in the same way, where $C_m = \frac{M(\rho_0)}{\rho_0 Q'(\rho_0)}$.

Remark 4.8. Contrary to classical models of lattice paths the number of excursions is of the same order of magnitude as the one of meanders. In probabilistic terms this means that the set of excursions is not a null set with respect to the set of meanders.

Remark 4.9. It is one of the surprising behaviours of Dyck paths with catastrophes: they involve algebraic quantities of degree 3; this was quite counter-intuitive to predict a priori, as Dyck path statistics usually involve by design algebraic quantities of degree 2.

In the next sections, we will need the following variant of the supercritical composition scheme from Flajolet and Sedgewick (2009, Proposition IX.6), in which we add a perturbation function q(z). In the following, we denote by ρ_f the radius of convergence of a function f(z).

Proposition 4.10 (Perturbed supercritical composition). Consider a combinatorial structure \mathcal{F} constructed from \mathcal{H} components according to the bivariate composition schema F(z, u) = q(z)g(uh(z)). Assume that g(z) and h(z) satisfy the supercriticality condition $h(\rho_h) > \rho_g$, that g is analytic in |z| < R for some $R > \rho_g$, with a unique dominant singularity at ρ_g , which is a simple pole, and that h is aperiodic. Furthermore, let q(z) be analytic for $|z| < \rho_h$. Then the number χ of \mathcal{H} -components in a random \mathcal{F} -structure of size n, corresponding to the probability distribution $[u^k z^n]F(z, u)/[z^n]F(z, 1)$ has a mean and variance that are asymptotically proportional to n; after standardization, the parameter χ satisfies a limiting Gaussian distribution, with speed of convergence $O(1/\sqrt{n})$.

Proof. As q(z) is analytic at the dominant singularity, it contributes only a constant factor to the asymptotics. Then, Hwang's Quasi-power theorem, see Flajolet and Sedgewick (2009, Theorem IX.8), gives the claim.

A simple (and useful) application of this result in the context of sequences leads to:

Proposition 4.11 (Perturbed supercritical sequences). Consider a sequence schema $\mathcal{F} = \mathcal{Q} \times SEQ(u\mathcal{H})$ that is supercritical, i.e., the value of h at its dominant positive singularity satisfies $h(\rho_h) > 1$. Assume that h is aperiodic, h(0) = 0, and q(z) is analytic for $|z| < \rho$, where ρ is the positive root of $h(\rho) = 1$. Then, the number X_n of \mathcal{H} -components in a random \mathcal{F} -structure of size n is, after standardization, asymptotically Gaussian with³

$$\mathbb{E}(X_n) \sim \frac{n}{\rho h'(\rho)}, \qquad \qquad \mathbb{V}(X_n) \sim n \frac{\rho h''(\rho) + h'(\rho) - \rho h'(\rho)^2}{\rho^2 h'(\rho)^3}.$$

³The formula for the asymptotics of $\mathbb{V}(X_n)$ in Flajolet and Sedgewick (2009, Proposition IX.7) contains some typos and misses the ρ -factors in the numerator and one in the denominator.

The number $X_n^{(m)}$ of components of some fixed size m is asymptotically Gaussian with asymptotic mean ~ $\theta_m n$, where $\theta_m = h_m \rho^m / (\rho h'(\rho))$.

Proof. The proof follows exactly the same lines as Flajolet and Sedgewick (2009, Proposition IX.7). We state it for completeness. The first part is a direct consequence of Proposition 4.10 with $g(z) = (1 - z)^{-1}$ and ρ_g replaced by 1. The second part results from the bivariate generating function

$$F(z, u) = \frac{q(z)}{1 - (u - 1)h_m z^m - h(z)}$$

and from the fact, that u close to 1 induces a smooth perturbation of the pole of F(z,1) at ρ , corresponding to u = 1.

4.2 Average number of catastrophes

In Theorem 2.1 we had seen that excursions consist of two parts: a prefix containing all catastrophes followed by the type of path one is interested in. If we want to count the number of catastrophes, it suffices therefore to analyse this prefix given by D(z). What is more, by (5) we know already how to count catastrophes: by counting occurrences of Q(z). Thus, let $d_{n,k}$ be the number of excursions ending with a catastrophe of length n with k catastrophes, then

$$D(z,v) := \sum_{n,k \ge 0} d_{n,k} z^n v^k = \frac{1}{1 - vQ(z)}.$$

Let $c_{n,k}$ be the number of excursions with k catastrophes, we get

$$C(z,v) \coloneqq \sum_{n,k\geq 0} c_{n,k} z^n v^k = D(z,v) E(z).$$
(10)

Theorem 4.12. The number of catastrophes of a random excursion with catastrophes of length n admits a limit distribution, with the limit law being dictated by the relation between the singularities ρ_0 and ρ .

1. In case $\rho_0 < \rho$, the standardized random variable

$$\frac{X_n - \mu n}{\sigma \sqrt{n}}, \qquad \mu = \frac{1}{\rho_0 Q'(\rho_0)}, \qquad \sigma^2 = \frac{\rho_0 Q''(\rho_0) + Q'(\rho_0) - \rho_0 Q'(\rho_0)^2}{\rho_0^2 Q'(\rho_0)^3},$$

converges in law to a standard Gaussian variable $\mathcal{N}(0,1)$:

$$\lim_{n \to \infty} \mathbb{P}\left(\frac{X_n - \mu n}{\sigma \sqrt{n}} \le x\right) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-y^2/2} \, dy.$$

2. In case $\rho_0 = \rho$, the normalized random variable

$$\frac{X_n}{\vartheta \sqrt{n}}, \qquad \qquad \vartheta = \frac{\sqrt{2}}{\eta},$$

converges in law to a Rayleigh distributed random variable with density $xe^{-x^2/2}$:

$$\lim_{n \to \infty} \mathbb{P}\left(\frac{X_n}{\vartheta \sqrt{n}} \le x\right) = 1 - e^{-x^2/2}$$

3. In case ρ_0 does not exist in $[0, \rho]$, the limit distribution is a discrete one:

$$\mathbb{P}(X_n = k) = \frac{(k\eta/\lambda + C/\tau)\lambda^k}{\eta D(\rho)^2 + C/\tau D(\rho)} \left(1 + O\left(\frac{1}{n}\right)\right),$$

where η is defined as in Theorem 4.3, $\lambda = Q(\rho)$, $C = \sqrt{2\frac{P(\tau)}{P''(\tau)}}$, and $\tau > 0$ the unique positive real root of P'(u) = 0. In particular, X_n converges to the random variable given by the law $\eta \operatorname{NegBinom}(2, \lambda) + \frac{C}{\tau} \operatorname{NegBinom}(1, \lambda)$.

Proof. First, for $\rho_0 < \rho$ we see from (10) that we are in the case of a perturbed supercritical composition scheme from Proposition 4.11. It is supercritical because Q(z) is singular at ρ_0 and $\lim_{z\to\rho_0} Q(z) = \infty$. The perturbation E(z) is analytic for $|z| < \rho$, and the other conditions are also satisfied. Hence, we get convergence to a normal distribution.

Second, for $\rho_0 = \rho$, we start with the asymptotic expansion of E(z) at $z \sim \rho$. Due to Banderier and Flajolet (2002, Theorem 3) we have

$$E(z) = E(\rho) \left(1 - \frac{C}{\tau} \sqrt{1 - z/\rho} \right) + O(1 - z/\rho),$$
(11)

for $z \sim \rho$. This implies by (7) the asymptotic expansion

$$\frac{1}{C(z,v)} = \frac{1}{E(\rho)} \left((1-v) + \eta \sqrt{1-z/\rho} \right) + O((1-z/\rho) + O\left((1-v)\sqrt{1-z/\rho} \right) + O$$

for $z \sim \rho$ and $v \sim 1$. The shape above is the one necessary for the Drmota–Soria limit scheme in Drmota and Soria (1997, Theorem 1) which implies a Rayleigh distribution. By a variant of the implicit function theorem applied to the small branches, the function satisfies the analytic continuation properties required to apply this theorem.

Third, we know by Theorem 4.3 that D(z) possesses a square-root singularity. Thus, combining the expansions (7), (9), and (11) we get the asymptotic expansion of C(z, v), which is of the same type of a square root as the one from Theorem 4.4. Extracting coefficients with the help of singularity analysis and normalizing by the result of Theorem 4.4 shows the claim.

In the last case of a discrete limit law, the probability generating function is asymptotically equal to

$$\frac{\eta D(\rho, v)^2 + (C/\tau)D(\rho, v)}{\eta D(\rho)^2 + (C/\tau)D(\rho)} \left(1 + O\left(\frac{1}{n}\right)\right).$$

Let us end this discussion with an application to Dyck paths.

Corollary 4.13. The number of catastrophes of a random Dyck path with catastrophes of length n is normally distributed. The standardized version of X_n ,

 $\frac{X_n - \mu n}{\sigma \sqrt{n}}, \qquad \mu \approx 0.0708358118, \qquad \sigma^2 \approx 0.05078979113,$

where μ is the unique positive real root of $31\mu^3 + 31\mu^2 + 40\mu - 3$, and σ is the unique positive real root of $29791\sigma^6 - 59582\sigma^4 + 60579\sigma^2 - 2927$, converges in law to a Gaussian variable $\mathcal{N}(0, 1)$.

4.3 Average number of returns to zero

In order to count the number of returns to zero, we decompose $F_0(z)$ into a sequence of arches. Let A(z) be the corresponding generating function. (Caveat: this is not the same generating function as A(z) in Proposition 3.2.) Then,

$$A(z) = 1 - \frac{1}{F_0(z)}.$$

Let $g_{n,k}$ be the number of excursions with catastrophes of length n and k returns to zero. Then,

$$G(z,v) \coloneqq \sum_{n,k\geq 0} g_{n,k} z^n v^k = \frac{1}{1 - vA(z)}.$$

From now on, let X_n be the random variable giving the number of returns to zero in excursions with catastrophes of length n drawn uniformly at random:

$$\mathbb{P}(X_n = k) = \frac{[z^n v^k]G(z, v)}{[z^n]G(z, 1)}$$

Applying the same ideas and techniques we used in the proof of Theorem 4.12, we get the following result.

Theorem 4.14. The number of returns to zero of a random excursion with catastrophes of length n admits a limit distribution, with the limit law being dictated by the relation between the singularities ρ_0 and ρ .

1. In case $\rho_0 < \rho$, the standardized random variable

$$\frac{X_n - \mu n}{\sigma \sqrt{n}}, \qquad \mu = \frac{1}{\rho_0 A'(\rho_0)}, \qquad \sigma^2 = \frac{\rho_0 A''(\rho_0) + A'(\rho_0) - \rho_0 A'(\rho_0)^2}{\rho_0^2 A'(\rho_0)^3},$$

converges in law to a standard Gaussian variable $\mathcal{N}(0,1)$.

2. In case $\rho_0 = \rho$, the normalized random variable

$$\frac{X_n}{\vartheta\sqrt{n}},\qquad\qquad\qquad\vartheta=\sqrt{2}\frac{E(\rho)}{\eta},$$

converges in law to a Rayleigh distributed random variable with density $xe^{-x^2/2}$.

3. In case ρ_0 does not exist, the limit distribution is NegBinom $(2, \lambda)$:

$$\mathbb{P}\left(X_n = k\right) = \frac{n\lambda^n}{F_0(\rho)^2} \left(1 + O\left(\frac{1}{n}\right)\right), \qquad \lambda = A(\rho) = 1 - \frac{1}{F_0(\rho)}.$$

Again, we give the concrete statement for Dyck paths with catastrophes.

Corollary 4.15. The number of returns to zero of a random Dyck path with catastrophes of length n is normally distributed. The standardized version of X_n ,

$$\frac{X_n - \mu n}{\sigma \sqrt{n}}, \qquad \mu \approx 0.1038149281, \qquad \sigma^2 \approx 0.1198688826,$$

where μ is the unique positive real root of $31\mu^3 - 62\mu^2 + 35\mu - 3$, and σ is the unique positive real root of $29791\sigma^6 + 231\sigma^2 - 79$, converges in law to a Gaussian variable $\mathcal{N}(0, 1)$.

It is interesting to compare the results of Corollaries 4.13 and 4.15 for Dyck paths: more than 10% of all steps are returns to zero, and more than 7% are catastrophes. This implies that among all returns to zero approximately 70% are catastrophes and 30% are -1-jumps.

4.4 Average final altitude

In this section we want to analyse the final altitude of a path after a certain number of steps. The *final altitude* of a path is defined as the ordinate of its endpoint. Theorem 2.1 already encodes this parameter using u:

$$F(z,u) = D(z)M(z,u), \qquad M(z,u) = \frac{\prod_{i=1}^{c} (u - u_i(z))}{u^c(1 - zP(u))},$$

where M(z, u) is the bivariate generating function of meanders.

Let X_n be the random variable giving the final altitude of paths with catastrophes of length n drawn uniformly at random:

$$\mathbb{P}(X_n = k) = \frac{[z^n u^k] F(z, u)}{[z^n] F(z, 1)}$$

This random variable exhibits an interesting fractal-like behaviour, as can be observed in Figure 5, and is more formally stated in the following theorem.

Theorem 4.16. The final altitude of a random lattice path with catastrophes of length n admits a discrete limit distribution:

$$\lim_{n \to \infty} \mathbb{P}(X_n = k) = [u^k] \omega(u), \quad \text{where } \omega(u) = \begin{cases} \prod_{\ell=1}^d \frac{1 - v_\ell(\rho_0)}{u - v_\ell(\rho_0)}, & \text{for } \rho_0 \le \rho, \\ \frac{\eta D(\rho) + \frac{C}{\tau - u}}{\eta D(\rho) + \frac{C}{\tau - 1}} \prod_{\ell=1}^d \frac{1 - v_\ell(\rho)}{u - v_\ell(\rho)}, & \text{for } \nexists \rho_0. \end{cases}$$

Proof. Let us distinguish three cases. First, in the case of $\rho_0 < \rho$ the function D(z) is responsible for the singularity of F(z, u). Thus, by Pólya and Szegő (1925, Problem 178) (see also Flajolet and Sedgewick (2009, Theorem VI.12)) we get the asymptotic expansion

$$\lim_{n \to \infty} \frac{[z^n]F(z,u)}{[z^n]F(z,1)} = \frac{M(\rho_0,u)}{M(\rho_0,1)} = \prod_{\ell=1}^d \frac{1 - v_\ell(\rho_0)}{u - v_\ell(\rho_0)}.$$

For the other cases by Lemma 4.2 we require $\delta < 0$. Yet, in this case we know from Banderier and Flajolet (2002, Theorem 6) that M(z, u) admits a discrete limit distribution. It holds that M(z, u) admits the expansion

$$M(z,u) = M(\rho,u) \left(1 + \frac{C}{u-\tau} \sqrt{1-z/\rho} \right) + O\left(1-z/\rho\right), \quad \text{for } z \to \rho.$$

In the second case and third case for $\rho_0 = \rho$ and $\nexists \rho_0$, we get by multiplying this expansion with the one of D(z) from (8) and (9), respectively, the expansion of F(z, u). Normalizing with the results of Theorem 4.5 yields the result.

Corollary 4.17. The final altitude of a random Dyck path with catastrophes of length n admits a geometric limit distribution with parameter $\lambda = v_1(\rho_0)^{-1} \approx 0.6823278$:

$$\mathbb{P}(X_n = k) \sim (1 - \lambda) \lambda^k$$

The parameter is the unique positive real root of $\lambda^3 + \lambda - 1$ and is given by

$$\lambda = \frac{1}{6} \left(108 + 12\sqrt{93} \right)^{1/3} - 2 \left(108 + 12\sqrt{93} \right)^{-1/3}$$

The situation for Dyck path with catastrophes hides a larger variety of behaviour, as seen in Figure 5.




Figure 5: The limit law of the final altitude for different jump polynomials P(u).

The main periodicity observed in these pictures is due to the fact that the limit law is a sum of "geometric limit laws" of *complex* parameter (as given in Theorem 4.16). The pictures show a combination of a "macroscopic" and a "microscopic" behaviour. On the macroscopic level we see a period of the size of the largest positive jump d. On the microscopic level, we see smaller fluctuations related to the small amplitude jumps (with some additional periodic behaviour if the support of these small amplitude jumps is periodic).

It is noteworthy that the way these pictures approach 0 has some links with the Skolem– Pisot problem (i.e., deciding if a rational function $R(u) \in \mathbb{Z}[[u]]$ has a 0 term in its Taylor expansion): the respective location of the poles of R(u) (and their residues) dictates how near from zero one can get.

4.5 Total amplitude of catastrophes

Another interesting parameter is the total amplitude of catastrophes of excursions of length n. Let $a_{n,k}$ be the number of excursions with catastrophes of length n and total amplitude k of all catastrophes contained in the path. Then its bivariate generating function $A_{tot}(z, u) = \sum_{n,k\geq 0} a_{n,k} z^n u^k$ is given by

$$A_{tot}(z, u) = D(z, u)E(z), \quad \text{where}$$
$$D(z, u) = \frac{1}{1 - Q(z, u)}, \quad \text{and}$$
$$Q(z, u) = zq\left(M(z, u) - E(z) - \sum_{-j \in J_{-}} u^{j}M_{j}(z)\right)$$

The generating function Q(z, u) keeps track of the altitudes of used catastrophes. The new parameter u does not influence the singular expansion of Q(z) analysed in Theorem 4.3. We get for $z \rightarrow \rho$ - and $0 \le u \le 1$ the following expansion

$$Q(z,u) = Q(\rho,u) - \eta(u)\sqrt{1 - z/\rho} + O(1 - z/\rho),$$
(12)

where $\eta(u)$ is a nonzero function, and in terms of the previous expansion of Q(z) we have that $\eta(1) = \eta$.

Let X_n be the random variable giving the total amplitude of catastrophes in lattice paths with catastrophes of length n drawn uniformly at random:

$$\mathbb{P}(X_n = k) = \frac{[z^n u^k] A_{\mathsf{tot}}(z, u)}{[z^n] A_{\mathsf{tot}}(z, 1)}.$$

Theorem 4.18. The total amplitude of catastrophes of a random excursion with catastrophes of length n admits a limit distribution, with the limit law being dictated by the relation between the singularities ρ_0 and ρ .

1. In case $\rho_0 < \rho$, the standardized random variable

$$\frac{X_n - \mu n}{\sigma \sqrt{n}}, \qquad \mu = \frac{Q_u(\rho_0, 1)}{\rho_0 Q_z(\rho_0, 1)},$$
$$\sigma^2 = \left(1 + \frac{\rho_0 Q_{uu}(\rho_0, 1)}{Q_z(\rho_0, 1)}\right) \mu^2 + \left(1 - \frac{2Q_{zu}(\rho_0, 1)}{Q_z(\rho_0, 1)} + \frac{Q_{zz}(\rho_0, 1)}{Q_z(\rho_0, 1)}\right) \mu,$$

converges for $\sigma^2 > 0$ in law to a standard Gaussian variable $\mathcal{N}(0,1)$.

2. In case $\rho_0 = \rho$, the normalized random variable

$$\frac{X_n}{\vartheta\sqrt{n}},\qquad\qquad\qquad \vartheta=\sqrt{2}\frac{Q_u(\rho,1)}{\eta},$$

converges in law to a Rayleigh distributed random variable with density $xe^{-x^2/2}$.

3. In case ρ_0 does not exist, the limit distribution is discrete and given by:

$$\lim_{n \to \infty} \mathbb{P}(X_n = k) = [u^k] \frac{\eta(u)D(\rho, u)^2 + \frac{C}{\tau}D(\rho, u)}{\eta D(\rho)^2 + \frac{C}{\tau}D(\rho)}.$$

Proof. In the first case $\rho_0 < \rho$ we will use the meromorphic schema from Flajolet and Sedgewick (2009, Theorem IX.9), which is a generalization of Hwang's quasi-power theorem. In order to apply it we need to check three conditions. First, the *meromorphic perturbation condition*: We know already from the proof of Theorem 4.3 that ρ_0 is a simple pole. What remains is to show that in a domain $\mathcal{D} = \{(z, u) : |z| < r, |u - 1| < \varepsilon\}$ the function admits the following representation

$$A_{\mathsf{tot}}(z,u) = \frac{B(z,u)}{C(z,u)},$$

where B(z, u) and C(z, u) are analytic for $(z, u) \in \mathcal{D}$. There exists a $\delta > 0$ such that $r := \rho_0 + \delta < \rho$. For this value the representation holds, as $B(z, u) = u^c(1 - zP(u))E(z)$ and $C(z, u) = u^c(1 - zP(u)) - zq \prod_{i=1}^c (1 - u_i(z))$ are only singular for $z = \rho$ or u = 0.

Next, the non-degeneracy $Q_u(\rho, 1)Q_z(\rho, 1) \neq 0$ is easily checked. It ensures the existence of a non-constant $\rho(u)$ analytic at u = 1, such that $1 - Q(\rho(u), u) = 0$.

Finally, the variability condition $r''(1) + r'(1) - r'(1)^2 \neq 0$ for $r(u) = \frac{\rho(1)}{\rho(u)}$ is also satisfied due to

$$\rho(1) = \rho_0, \qquad \rho'(1) = -\frac{Q_u(\rho, 1)}{Q_z(\rho, 1)},$$

$$\rho''(1) = -\frac{1}{Q_z(\rho, 1)} \left(Q_{zz}(\rho, 1)\rho'(1) + 2Q_{z,u}(\rho, 1)\rho'(1) + Q_{uu}(\rho, 1) \right).$$

This implies the claimed normal distribution.

In the second case $\rho_0 = \rho$ we apply again the Drmota–Soria limit theorem Drmota and Soria (1997, Theorem 1) which leads to a Rayleigh distribution. As $Q(\rho_0, 1) = 1$, like in Equation (6), we have a cancellation of the constant term in the Puiseux expansion (for $z \sim \rho, u \sim 1$). Thus, using the asymptotic expansions (11) and (12) leads to

$$\frac{1}{A_{\text{tot}}(z,u)} = \frac{Q_u(\rho,1)}{E(\rho)}(1-u) + \frac{\eta}{E(\rho)}\sqrt{1-z/\rho} + O((1-z/\rho)) + O((u-1)(1-z/\rho)) + O((u-1)^2)$$

Note that the analyticity and the other technical conditions required to apply this theorem follow from the respective properties of the generating functions M(z, u), E(z), and $M_j(z)$. This implies the claimed Rayleigh distribution with the normalizing constant $\vartheta = \sqrt{2} \frac{Q_u(\rho, 1)}{n}$.

In the third case $\nexists \rho_0$, the singularity arises at $z = \rho$. In particular, there arises no zero in the denominator. Thus, after combining the known expansions (11) and (12), singularity analysis yields the given discrete form. This implies the claimed discrete distribution.

Corollary 4.19. The total amplitude of catastrophes of a random Dyck path with catastrophes of length n is normally distributed. The standardized version of X_n ,

 $\frac{X_n - \mu n}{\sigma \sqrt{n}}, \qquad \mu \approx 0.2938197987, \qquad \sigma^2 \approx 0.5809693987,$

where μ is the unique positive real root of $31\mu^3 + 62\mu^2 + 71\mu - 27$, and σ is the unique positive real root of $29791\sigma^6 - 59582\sigma^4 + 298411\sigma^2 - 159099$, converges in law to a Gaussian variable $\mathcal{N}(0,1)$.

4.6 Amplitude of an average catastrophe

As one of the last parameters of our lattice paths with catastrophes, we want to determine the law behind the altitude of a random catastrophe among all lattice paths of length n. In other words, one draws uniformly at random a catastrophe among all possible catastrophes of all lattice paths of length n. Note that this is also the law behind the altitude of the first (or last) catastrophe, as cyclic shifts of the excursions ending with a catastrophe transform any catastrophe into the first (or last) one.

We can construct it from the generating function counting the number of catastrophes. It is given in (10) where each catastrophe is marked by a variable v.

Lemma 4.20. The bivariate generating function $A_{avg}(z, u)$ counting the altitude of a random catastrophe among all excursions with catastrophes is given by

$$A_{\mathsf{avg}}(z,u) = E(z) + Q(z,u)D(z)^2E(z).$$

Proof. A random excursion with catastrophes either contains no catastrophes and is counted by E(z), or it contains at least one catastrophe. In the latter we choose one of its catastrophes and its associated excursion ending with this catastrophe. Then we replace it with an excursion ending with a catastrophe whose altitude has been marked. This corresponds to

$$A_{\text{avg}}(z,u) = E(z) + \frac{Q(z,u)}{Q(z)} \left. \frac{\partial}{\partial v} C(z,v) \right|_{v=1}$$

Computing this expression proves the claim.

As before we define a random variable X_n for our parameter as

$$\mathbb{P}(X_n = k) = \frac{[z^n u^k] A_{\mathsf{avg}}(z, u)}{[z^n] A_{\mathsf{avg}}(z, 1)}.$$

Due to the factor Q(z, u) the situation is similar to final altitude in Section 4.4.

Theorem 4.21. The altitude of a random catastrophe of a lattice path of length n admits a discrete limit distribution:

$$\lim_{n \to \infty} \mathbb{P} \left(X_n = k \right) = \begin{bmatrix} u^k \end{bmatrix} \omega(u), \quad \text{where} \\ \omega(u) = \begin{cases} Q(\rho_0, u), & \text{for } \rho_0 \le \rho, \\ \frac{\frac{C}{\tau} + \left(\frac{C}{\tau} D(\rho)^2 + 2\eta D(\rho)^3 \right) Q(\rho, u) + \eta(u) D(\rho)^2}{\frac{C}{\tau} + \left(\frac{C}{\tau} D(\rho)^2 + 2\eta D(\rho)^3 \right) Q(\rho, 1) + \eta D(\rho)^2}, & \text{for } \nexists \rho_0. \end{cases}$$

Proof. The proof is similar to the one of Theorem 4.16.

First, for $\rho_0 < \rho$ it holds that only $D(z)^2$ is singular at ρ_0 , where all other terms are analytic. Thus, by Pólya and Szegő (1925, Problem 178) the claim holds.

Second, in case $\rho_0 = \rho$ we combine the singular expansions (8), (11), and (12) to get

$$A_{\text{avg}}(z,u) = \frac{E(\rho)Q(\rho,u)}{\eta^2(1-z/\rho)} + O\left((1-z/\rho)^{-1/2}\right).$$

In other words, the polar singularity of $D(z)^2$ dominates, and the situation is similar to the one before.

In the final case, $\nexists \rho_0$, we again combine the singular expansions. Yet this time the expansion of D(z) is given by (9). This implies a contribution of all terms, as all of them are singular at once and all of them have the same type of singularity.

Corollary 4.22. The amplitude of a random catastrophe among all Dyck paths with catastrophes of length n admits a (shifted) geometric limit distribution with parameter $\lambda \approx 0.6823278$:

$$\lim_{n \to \infty} \mathbb{P} \left(X_n = k \right) = \begin{cases} (1 - \lambda) \,\lambda^{k-2}, & \text{for } k \ge 2, \\ 0, & \text{for } k = 0, 1 \end{cases}$$

The parameter is the unique positive real root of $\lambda^3 + \lambda - 1$ and given by

$$\lambda = \frac{1}{6} \left(108 + 12\sqrt{93} \right)^{1/3} - 2 \left(108 + 12\sqrt{93} \right)^{-1/3}.$$

Comparing this result to the one for the final altitude of meanders in Corollary 4.17, we see that the type of the law is of the same nature (yet shifted for the amplitude of catastrophes), and that the parameter λ is the same. The following lemma explains this connection.

Lemma 4.23. Let d = 1, i.e., $P(u) = p_{-c}u^{-c} + \cdots + p_1u^1$ be the jump polynomial. Then, the generating function of excursions of length n (marked by z) ending with a catastrophe of altitude k (marked by u) admits the following decomposition

$$Q(z,u) = qp_1 z^2 u^{c+1} M(z,u) M_c(z) + qz \sum_{\substack{-j \notin J_- \\ -c < j < 0}} u^{-j} M_j(z).$$

Proof. The idea is a last passage decomposition with respect to reaching level c + 1. First, assume that $p_{-c}, \ldots, p_{-1} \neq 0$. Then, the smallest catastrophe is of altitude c+1. We decompose the excursion with respect to the last jump from altitude c to altitude c+1, see Figure 6. Left of it, there is a meander ending at altitude c, and right of it there is a meander starting at altitude c+1 and always staying above altitude c+1. The altitude of the ending catastrophe is then given by the final altitude plus c+1, this gives the factor $zu^{c+1}M(z,u)$. This proves the first part.

For the second part, note that if one of the p_{-c}, \ldots, p_{-1} is equal to 0, then catastrophes of the respective height are allowed. These are given by meanders ending at this altitude and a jump to 0.




Figure 6: Decomposition of an excursion ending with a catastrophe for d = 1 from Lemma 4.23.

This lemma shows that the probability generating function of the final altitude and of the amplitude of an average catastrophe are connected. In particular for d = 1 and $\rho_0 < \rho$ we have

$$\lim_{n \to \infty} \frac{[z^n]M(z,u)}{[z^n]M(z,1)} = \frac{1}{u^{c+1}} \frac{Q(\rho_0, u) - \sum_{j \notin J_-, -c < j < 0} q\rho_0 u^{-j} M_j(\rho_0)}{1 - \sum_{j \notin J_-, -c < j < 0} q\rho_0 M_j(\rho_0)}.$$

We see the shift by u^{-c-1} of the probability generating function. It is now obvious how these laws are related: the parameters are the same, there is just a shift in the parameter and a subtraction of certain, initial values.

The results of Lemma 4.23 can be generalized to $d \ge 2$, but the result is more complicated. For example, for d = 2 there are 4 different cases in the last passage decomposition: a +1-jump from c to c + 1, a +2-jump from c to c + 2, a +2-jump from altitude c - 1 to c + 1, all followed by a meander, and a +2-jump from c to c + 2 followed by a path always staying above c + 1.

However, in all cases there is a factor M(z, u) in Q(z, u) if $p_{-c}, \ldots, p_{-1} \neq 0$.

4.7 Waiting time for the first catastrophe

We end the discussion on limit laws with a parameter that might be of the biggest interest in applications: the waiting time for the first catastrophe. Let $w_{n,k}$ be the number of excursions with catastrophes of length n such that the first catastrophe appears at the k-th steps for k > 0. Let $w_{n,0}$ be the number of such paths without a catastrophe. Then its bivariate generating function $W(z, u) = \sum_{n,k\geq 0} w_{n,k} z^n u^k$ is given by

$$W(z, u) = E(z) + Q(zu)D(z)E(z).$$

This is easily derived from Theorem 2.1 as the prefix D(z) is a sequence of excursions with only one catastrophe at the very end. Thus, marking the length of the first of such excursions marks the position of the first catastrophe.

As done repeatedly we define a random variable X_n for our parameter as

$$\mathbb{P}(X_n = k) = \frac{[z^n u^k]W(z, u)}{[z^n]W(z, 1)}.$$

Theorem 4.24. The waiting time for the first catastrophe in a lattice path with catastrophes of length n admits a discrete limit distribution:

$$\lim_{n \to \infty} \mathbb{P}(X_n = k) = [u^k] \omega(u), \quad \text{where} \quad \omega(u) = \begin{cases} Q(\rho_0 u), & \text{for } \rho_0 \le \rho, \\ 1 - Q(\rho) + Q(\rho u), & \text{for } \nexists \rho_0. \end{cases}$$

Proof. The proof follows again the same lines as the one of Theorem 4.16. In this particular case, we combine the asymptotic expansion of D(z)E(z) from Theorem 4.4 with the asymptotic expansion of E(z) from (Banderier and Flajolet, 2002, Theorem 3).

In the case of Dyck paths we have

$$Q(z) = \frac{1}{2z^2} \left(z^2 - z - 1 - (z^2 + z - 1) \sqrt{\frac{1+2z}{1-2z}} \right).$$

The corresponding limit law which consists of the sum of two discrete distributions for the odd and even case, is shown in Figure 7. We see a periodic behaviour with a distribution for the even and odd steps. This arises from fact that not only at altitude 0 but also at altitude 1 catastrophes are not allowed. Starting from the origin this effects only the odd numbered steps. The probabilities for catastrophes at an odd step are lower than the ones at the following even step, because we can reach altitude 1 from below and from above, whereas the only restriction for the even steps is at altitude 0 which can only be reached from above. In particular, it is reasonable that the later the first catastrophe appears the more unlikely it gets. Yet, it is still interesting to discover that the probability of the first catastrophe being at step 4 is lower than at step 6.




Figure 7: The red dots represent the discrete limit distribution of the waiting time for the first catastrophe in the case of Dyck paths. It is the sum of two discrete distributions for the even (brown and below) case and odd (black and above).

5 Uniform random generation

In order to generate our lattice paths with catastrophes, we can build on some key methods from the last 30 years.

For sure, it is possible to generate lattice paths with catastrophes via a dynamic programming approach, but this would require $O(n^3)$ bits in memory, our next theorem shows we can do much better:

Theorem 5.1 (Uniform random generation). Dyck paths with catastrophes can be generated in linear time. Lattice paths with catastrophes of length n can be generated uniformly at random in time $O(n \ln n)$ with O(n) memory, or in time $O(n^{3/2})$ with O(1) memory (if outpout is given as a stream).

Proof. First, via the bijection of Theorem 3.1, the linear-time approach of Bacher, Bodini, and Jacquot (2013) for Motzkin trees can be applied to Dyck paths with catastrophes. The other cases can be tackled in 2 ways. A first solution is to see that classical Dyck paths (and generalized Dyck paths) can be generated by pushdown-automata, or equivalently, by a context-free grammar. The same holds trivially for generalized lattice paths with catastrophes. Then, using the recursive method of Flajolet, Zimmerman, and Van Cutsem (1994) (which can be seen as a wide generalization to combinatorial structures of what Hickey and Cohen (1983) did for context-free grammars), such paths of length n can be generated in $O(n \ln n)$ average-time. Goldwurm (1995) proved that this can be done with the same time-complexity, with only O(n) memory. The Boltzmann method introduced by Duchon, Flajolet, Louchard, and Schaeffer (2004) is also a way to get a linear average-time random generator for paths of length within $[(1 - \epsilon)n, (1 + \epsilon)n]$.

A second solution relies on a generating tree approach Banderier, Bousquet-Mélou, Denise, Flajolet, Gardy, and Gouyou-Beauchamps (2002), where each transition is computed via

$$\mathbb{P}\left(\begin{cases} \text{jump } j \text{ when at altitude } k, \text{ and length } m, \\ \text{ending at } 0 \text{ at length } n \end{cases}\right) = \frac{f_{m,k}^0 f_{n-(m+1),0}^{k+j}}{f_{n,0}^0},$$

where $f_{m,k}^i$ is the number of paths with catastrophes of length m, starting at altitude i and ending at altitude k. Then, for each pair (i, k), the theory of D-finite functions applied to the algebraic functions derived similarly to Theorem 2.1 allows us to get the recurrence for the corresponding f_m (see e.g. Banderier and Drmota (2015)). In order to get the m-th term f_m of a P-recursive sequence, there is a $O(\sqrt{m})$ algorithm due to Chudnovsky and Chudnovsky (1986). It is possible to win space complexity and bit complexity by computing the f_m 's in floating point arithmetic, instead of rational numbers (although all the f_m are integers, it is often the case that the leading term of the P-recursive recurrence is not 1, and thus it then implies rational number computations, and time loss in gcd computations). All of this leads to a cost $\sum_{m=1}^{n} (O(\sqrt{m}) + O(\sqrt{n-m})) = O(n^{3/2})$, moreover, a O(1) memory is enough to output the n jumps of the lattice path, step after step, as a stream.

6 Conclusion

In this article, motivated by a natural model in queuing theory where one allows a "reset" of the queue, we analysed the corresponding combinatorial model: lattice paths with catastrophes. We showed how to enumerate them, how to get closed forms for their generating functions.

En passant, we gave a bijection (Theorem 3.1) which extends directly to lattice paths with a -1 jump and an arbitrary set of positive jumps (they are sometimes called Łukasiewiecz paths). Łukasiewiecz paths with catastrophes could be considered as a kind of Galton–Watson process with catastrophes, in which some pandemic suddenly kills the full population. Our results quantify the probability of such a pandemic over long periods.

It is known that the limiting objects associated to classical Dyck paths behave like Brownian excursions, Brownian meanders (see Marchal (2003)). It was therefore interesting to see what type of different behaviour lattice paths with catastrophes exhibit. These are illustrated by our results on the asymptotics and on the limit laws of several parameters. It is unusual to see that this leads to some apparently fractal like limit laws (see Figure 5), that we in fact explained via our analytic combinatorics approach.

In conclusion, it is pleasant that the kernel method is once more allowing to solve a variant of lattice paths, giving the exact and asymptotic enumeration, and also leading to uniform random generation.

Remark on this version

This article is the long extended version of the article with the same title which appeared in the volume dedicated to the GASCom'2016 conference, published by Electronic Notes in Discrete Mathematics Banderier and Wallner (2017). In this long version, we included more details, we gave the proofs for the asymptotics results, and we also added the analysis of three new parameters: subsections 4.5 (total sums of catastrophes), 4.6 (average amplitude of a catastrophe), and 4.7 (waiting time for the first catastrophe). We also added the section 5 dedicated to uniform random generation issues.

Acknowledgments. The authors thank Alan Krinik and Gerardo Rubino who suggested considering this model of walks with catastrophes during the Lattice Paths Conference'15, which held in August 2015 in California in Pomona. We also thank Jean-Marc Fédou and the organizers of the GASCom 2016 Conference, which was held in Corsica in June 2016, where we presented the first short version of this work. The second author was supported by the Austrian Science Fund (FWF) grant SFB F50-03.

References

- Axel Bacher, Olivier Bodini, and Alice Jacquot. Exact-size sampling for Motzkin trees in linear time via Boltzmann samplers and holonomic specification. In SIAM Workshop on Analytic Algorithmics and Combinatorics (ANALCO'13), 2013.
- Cyril Banderier. Limit laws for basic parameters of lattice paths with unbounded jumps. In *Mathematics and computer science, II*, Trends Math., pages 33–47. Birkhäuser, 2002.
- Cyril Banderier and Michael Drmota. Formulae and asymptotics for coefficients of algebraic functions. *Combin. Probab. Comput.*, 24(1):1–53, 2015. URL http://dx.doi.org/10. 1017/S0963548314000728.
- Cyril Banderier and Philippe Flajolet. Basic analytic combinatorics of directed lattice paths. *Theoret. Comput. Sci.*, 281(1-2):37–80, 2002. URL http://dx.doi.org/10.1016/ S0304-3975(02)00007-5.
- Cyril Banderier and Donatela Merlini. Algebraic succession rules and lattice paths with an infinite set of jumps. *Proceedings of FPSAC'02*, pages 1–10, 2003.
- Cyril Banderier and Michael Wallner. The kernel method for lattice paths below a rational slope. In *Lattice paths combinatorics and applications*, Developments in Mathematics Series. Springer, 2016. To appear.
- Cyril Banderier and Michael Wallner. Lattice paths with catastrophes. *Electronic Notes in Discrete Mathematics*, 59:131–146, 2017. URL https://doi.org/10.1016/j.endm.2017. 05.010. Random Generation of Combinatorial Structures (GASCom 2016).
- Cyril Banderier, Mireille Bousquet-Mélou, Alain Denise, Philippe Flajolet, Danièle Gardy, and Dominique Gouyou-Beauchamps. Generating functions for generating trees. *Discrete Mathematics*, 246(1-3):29–55, 2002. Formal power series and algebraic combinatorics (Barcelona, 1999).
- Cyril Banderier, Jean-Marc Fédou, Christine Garcia, and Donatela Merlini. Algebraic succession rules and lattice paths with an infinite set of jumps. *Preprint*, pages 1–32, 2003.
- Elena Barcucci, Alberto Del Lungo, Elisa Pergola, and Renzo Pinzani. ECO: a methodology for the enumeration of combinatorial objects. *Journal of Difference Equations and Applications*, 5:435–490, 1999.
- Alexander Burstein. Restricted Dumont permutations. Ann. Comb., 9(3):269–280, 2005. URL http://dx.doi.org/10.1007/s00026-005-0256-4.
- David V. Chudnovsky and Gregory V. Chudnovsky. On expansion of algebraic functions in power and Puiseux series. I. J. Complexity, 2(4):271–294, 1986. URL http://dx.doi.org/10. 1016/0885-064X(86)90006-3.

- Michael Drmota and Michèle Soria. Images and preimages in random mappings. SIAM Journal on Discrete Mathematics, 10(2):246–269, 1997.
- Enrica Duchi, Jean-Marc Fédou, and Simone Rinaldi. From object grammars to ECO systems. *Theoret. Comput. Sci.*, 314(1-2):57–95, 2004. URL http://dx.doi.org/10.1016/j.tcs. 2003.10.037.
- Philippe Duchon, Philippe Flajolet, Guy Louchard, and Gilles Schaeffer. Boltzmann samplers for the random generation of combinatorial structures. *Combin. Probab. Comput.*, 13(4-5): 577–625, 2004. URL http://dx.doi.org/10.1017/S0963548304006315.
- William Feller. An introduction to probability theory and its applications. Vol. I. Third edition. John Wiley & Sons, 1968.
- Luca Ferrari, Elisa Pergola, Renzo Pinzani, and Simone Rinaldi. Some applications arising from the interactions between the theory of Catalan-like numbers and the ECO method. *Ars Combin.*, 99:109–128, 2011.
- Philippe Flajolet and Robert Sedgewick. *Analytic Combinatorics*. Cambridge University Press, 2009.
- Philippe Flajolet, Paul Zimmerman, and Bernard Van Cutsem. A calculus for the random generation of labelled combinatorial structures. *Theoret. Comput. Sci.*, 132(1-2):1–35, 1994. URL http://dx.doi.org/10.1016/0304-3975(94)90226-7.
- Massimiliano Goldwurm. Random generation of words in an algebraic language in linear binary space. Inform. Process. Lett., 54(4):229–233, 1995. URL http://dx.doi.org/10.1016/ 0020-0190(95)00025-8.
- Timothy Hickey and Jacques Cohen. Uniform random generation of strings in a context-free language. SIAM J. Comput., 12(4):645-655, 1983. URL http://dx.doi.org/10.1137/ 0212044.
- Blake Hunter, Allan Krinik, Cynthia Nguyen, Jennifer M. Switkes, and Hubertus F. Von Bremen. Gambler's ruin with catastrophes and windfalls. *Journal of Statistical Theory and Practice*, 2008.
- Alan Krinik, Gerardo Rubino, Daniel Marcus, Randall J. Swift, Hassan Kasfy, and Holly Lam. Dual processes to solve single server systems. J. Statist. Plann. Infer., 135(1):121–147, 2005. URL http://dx.doi.org/10.1016/j.jspi.2005.02.010.
- Jacques Labelle and Yeong Nan Yeh. Generalized Dyck paths. *Discrete Math.*, 82(1):1–6, 1990. URL http://dx.doi.org/10.1016/0012-365X(90)90039-K.
- K. Manes, A. Sapounakis, I. Tasoulas, and P. Tsikouras. Equivalence classes of ballot paths modulo strings of length 2 and 3. *Discrete Math.*, 339(10):2557–2572, 2016. URL http: //dx.doi.org/10.1016/j.disc.2016.05.001.

- Philippe Marchal. Constructing a sequence of random walks strongly converging to Brownian motion. In Cyril Banderier and Christrian Krattenthaler, editors, *Discrete random walks* (*Paris, 2003*), Discrete Math. Theor. Comput. Sci. Proc., AC, pages 181–190. 2003.
- George Pólya and Gábor Szegő. *Aufgaben und Lehrsätze aus der Analysis. Erster Band: Reihen, Integralrechnung Funktionentheorie.* Springer, 1st edition, 1925. ISBN 978-3-662-37599-0,978-3-662-38381-0.
- Richard P. Stanley. *Enumerative combinatorics. Volume 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, second edition, 2011. ISBN 978-1-107-60262-5.
- Julian West. Generating trees and forbidden subsequences. *Discrete Mathematics*, 157:363–374, 1996.