# ENUMERATIONS RELATING BRAID AND COMMUTATION CLASSES

SUSANNA FISHEL, ELIZABETH MILIĆEVIĆ, REBECCA PATRIAS, AND BRIDGET EILEEN TENNER

ABSTRACT. We obtain an upper and lower bound for the number of reduced words for a permutation in terms of the number of braid classes and the number of commutation classes of the permutation. We classify the permutations that achieve each of these bounds, and enumerate both cases.

#### 1. INTRODUCTION

The symmetric group  $S_n$  is the Coxeter group generated by the adjacent transpositions  $\{s_1, \ldots, s_{n-1}\}$ . Every  $w \in S_n$  can be expressed as a minimal-length product of these generators, and a given permutation may have more than one such representation. These representations are *reduced expressions* for w, each of which can be encoded as a *reduced word* for w. The set of reduced words R(w) possesses a rich combinatorial structure that has been studied from many different perspectives. For example, Stanley showed that the number |R(w)| of reduced words for w can be calculated in terms of Young tableaux of particular shapes [Sta84], but this is not an easy value to calculate outside of special cases. Another common technique explores various quotients of R(w) under the relations governing the adjacent transpositions. For example, the reduced words  $\mathbf{u}$  and  $\mathbf{v}$  for  $w \in S_n$  are in the same *commutation class* if we can obtain  $\mathbf{u}$  from  $\mathbf{v}$  by applying a sequence of commutation relations of the braid relation  $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$ . In this paper, we give upper and lower bounds for |R(w)|, in terms of the number of braid and commutation classes of the permutation w.

Quotients of R(w) under these Coxeter relations have been studied before, but almost all of that work has focused on the commutation classes, with very little attention paid to the braid classes. In [Eln97], Elnitsky presented a bijection from the commutation classes of w to rhombic tilings of a particular polygon that depends on w, which Tenner utilized extensively in [Ten06]. Jonsson and Welker [JW09] considered certain cell complexes where the commutation classes appear. Bergeron, Ceballos, and Labbé [BCL15] studied a similar scenario, but their work does consider quotienting out by more general Coxeter relations. Meng [Men10] studied both the number of commutation classes and their network relationships, and Bédard developed recursive formulas for the number of reduced words in each commutation class in [Béd99], as well as more detailed statistics for the case of the longest permutation  $w_0$ . Such focus on certain fixed elements has also been a productive approach to studying the set of reduced words. For example, the higher Bruhat order B(n, 2), studied

<sup>2010</sup> Mathematics Subject Classification. Primary: 05A05; Secondary: 05E15.

Key words and phrases. reduced word, braid class, commutation class, permutation.

SF was partially supported by grant 359602 from the Simons Foundation, as well as by funds from the Mathematical Sciences Research Institute. EM was partially supported by NSF grant DMS-1600982. BT was partially supported by Simons Foundation Collaboration Grant for Mathematicians 277603.

by Manin and Schechtman [MS89] and Ziegler [Zie93] is a partial order on the commutation classes of  $w_0$ .

Another important family of permutations for which the study of the commutation classes has been especially fruitful are the fully commutative elements, which are those having a single commutation class. In their study of Schubert polynomials, Billey, Jockusch, and Stanley [BJS93] showed that fully commutative permutations are 321-avoiding. Stembridge investigated and enumerated the fully commutative elements in all Coxeter groups in [Ste96, Ste97, Ste98], and he provided further examples of their appearance in algebra. More recently, Green and Losonczy [GL02] and Hanusa and Jones [HJ10] have worked on adaptations of full commutativity to root systems and affine permutations, respectively.

The equivalence classes of R(w) under the braid relations, on the other hand, have been noticeably less well studied, both in terms of how they partition the collection of reduced words, and as objects of structure themselves. Zollinger [Zol94] used an encoding of the reduced words to provide formulas for the size of the braid classes (cf. Proposition 3.5, below), and the previously cited work in [BCL15] showed that the graph on braid classes, with edges indicating when an element of one class can be transformed into an element of the other class by a commutation move, is bipartite.

As with commutation classes, focusing on a particular element in  $S_n$ , such as the longest word  $w_0$ , has been a fruitful approach to understanding some of the relevance and influence of braid moves in reduced words. Reiner and Roichman [RR13] defined a graph whose vertices are the elements of  $R(w_0)$  with edges indicating braid relations, and they calculate the diameter of this graph. Building on Reiner's proof [Rei05] that the expected number of braid moves in a random element of  $R(w_0)$  equals one, Schilling, Thiéry, White, and Williams [STWW17] extend this probabilistic result to the case for one particular reduced word for the longest element.

Although braid classes and commutation classes are highly related, recognition of this fact has thus far been under-utilized in the literature. In this paper, we seek to remedy that by studying the relationship between the number of reduced words of a permutation and the numbers of commutation classes and braid classes that it has. Our driving principle is to utilize the fact that commutation classes and braid classes are partitions of the same set. This allows us to leverage one against the other when studying them in tandem. Theorem 4.6 illustrates this approach by providing upper and lower bounds on |R(w)| in terms of the number of braid and commutation classes. More precisely,

$$|B(w)| + |C(w)| - 1 \le |R(w)| \le |B(w)| \cdot |C(w)|$$

where we define this notation in the next section. We demonstrate that these bounds are sharp by characterizing in Proposition 4.8 (and Corollary 4.10) and Proposition 4.15 those permutations that achieve the upper and lower bounds, respectively. Those collections of permutations are enumerated in Corollaries 4.11 and 4.16. In Conjecture 5.2, we suggest an alternate approach to studying these relationships in terms of intervals in the weak order, rather than working directly with the Coxeter relations on reduced words.

We focus on reduced words for elements of the symmetric group, and Section 2 provides the required background information about Coxeter groups, as well as the relevant terminology and notation used throughout the paper. In Section 3, we recall facts about the graph with vertex set R(w) and edges indicating a single commutation or braid move, and obtain initial results about the quantity and structure of braid classes. This sets the stage for Section 4,

which contains the majority of our results, including those discussed above, and the paper concludes in Section 5 with a proposed connection to aspects of intervals in the weak order.

Acknowledgements. This project began at the "Algebraic Combinatorixx 2" workshop at the Banff International Research Station in May 2017. We thank the organizers and staff for putting together such a wonderful conference, and thank Kyoungsuk Park for early discussions. The computations in this paper were performed by using Maple<sup>TM</sup>, a trademark of Waterloo Maple Inc. John Stembridge's posets and coxeter packages for Maple were essential.

### 2. Coxeter groups

We now discuss relevant background from the theory of Coxeter groups, and we refer the reader to [BB05] for more information.

A finitely generated Coxeter group is a group W with a presentation of the form

$$\langle s \in S \mid s^2 = 1 \text{ for all } s \in S \text{ and } (st)^{m(s,t)} = 1 \text{ for all } s, t \in S \rangle,$$

where S is a finite set and  $m(s,t) = m(t,s) \in \{2,3,\ldots\} \cup \{+\infty\}$ . The elements of S are the simple reflections. Because the simple reflections generate the group, any element of Wmay be written as a product  $w = s_{i_1} s_{i_2} \cdots s_{i_k}$ , where each  $s_{i_j} \in S$ . When this k is minimal, it is the *length* of w, and this minimal value of k is denoted  $\ell(w)$ . An expression of w as  $s_{i_1}s_{i_2}\cdots s_{i_{\ell(w)}}$  is a reduced expression for w, and the word formed by the indices  $i_1i_2\cdots i_{\ell(w)}$ is a reduced word for w. The set of reduced words for  $w \in W$  is denoted R(w), and a word formed by taking an ordered subset of the indices in a reduced word  $i_1 i_2 \cdots i_{\ell(w)}$  for w, namely  $i_{j_1}i_{j_2}\cdots i_{j_m}$  where  $1 \leq j_1 < \cdots < j_m \leq \ell(w)$ , is a subword. A factor is a subword in which those indices are consecutive.

In this work, we focus on finite Coxeter groups of type A. We should note, however, that the objects in this paper (such as length, reduced expressions, and the commutation and braid classes defined below) have analogues in other types. Indeed some of the results cited below have been proved in that broader context. The Coxeter group of type  $A_n$  is generated by the simple reflections  $S = \{s_1, \ldots, s_n\}$  subject to the Coxeter relations:

- R1.  $s_i^2 = 1$  for all  $i \in [n]$ , R2.  $(s_i s_{i+1})^3 = 1$  for all  $i \in [n-1]$ , and
- R3.  $(s_i s_j)^2 = 1$  for |i j| > 1.

where [n] denotes the set  $\{1, 2, \ldots, n\}$ . This is, of course, a presentation of the symmetric group  $S_{n+1}$ , where the simple reflection  $s_i$  is the adjacent transposition swapping i and i + 1. Most often, we will express elements of this Coxeter group as reduced words, but occasionally we also use the *window* or *one-line* notation, which records the action of the permutation  $w \in S_n$  on the elements of [n] as follows  $w = [w(1) \ w(2) \ \cdots \ w(n)]$ . The length of a permutation w also has an interpretation in terms of the window notation. A pair (i, j)with i < j is an *inversion* of w if w(i) > w(j).

Relation R2 is the *braid relation* and can be reformulated as

$$s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}.$$

Relations of the form R3 are commutation relations:  $s_i s_j = s_j s_j$  when |i-j| > 1. We refer to an application of a braid relation as a *braid move*, an application of a commutation relation as a *commutation move*, and an application of either relation as a *Coxeter move*.

For a fixed permutation w, the set R(w) can be partitioned by either of these Coxeter moves, forming the classes

$$C(w) := R(w)/(ij \sim ji)$$
 when  $|i - j| > 1$ , and  
 $B(w) := R(w)/(i(i+1)i \sim (i+1)i(i+1)).$ 

The first of these, C(w), denotes the *commutation classes* of R(w), and the second, B(w), denotes the *braid classes*. That is, two elements of some  $C \in C(w)$  can be obtained from each other by a sequence of commutation moves, whereas elements of some  $B \in B(w)$  can be obtained from each other by a sequence of braid moves.

**Example 2.1.** Consider  $[25314] \in S_5$ . Then

$$R([25314]) = \{12432, 14232, 41232, 14323, 41323, 43123\}$$
$$C([25314]) = \{\{12432, 14232, 41232\}, \{14323, 41323, 43123\}\}$$
$$B([25314]) = \{\{12432\}, \{14232, 14323\}, \{41232, 41323\}, \{43123\}\}.$$

Consider a reduced word  $\mathbf{u} = u_1 \cdots u_k$ . Let  $\mathbf{b}_i$  denote the braid relation that affects the subword  $u_{i-1}u_iu_{i+1}$  in the case when  $u_{i-1} = u_{i+1}$  and  $|u_{i\pm 1} - u_i| = 1$ , and acts as the identity otherwise. Let  $\mathbf{c}_i$  denote the commutation relation that swaps  $u_i$  with  $u_{i+1}$  if  $|u_i - u_{i+1}| > 1$ , and acts as the identity otherwise.

**Definition 2.2.** If  $\mathfrak{b}_i(\mathbf{u}) \neq \mathbf{u}$  then  $\mathbf{u}$  supports a braid move in position i; equivalently, the subword  $u_{i-1}u_iu_{i+1}$  supports a braid move. If  $\mathfrak{c}_i(\mathbf{u}) \neq \mathbf{u}$  then  $\mathbf{u}$  supports a commutation move in position i; equivalently, the subword  $u_iu_{i+1}$  supports a commutation move.

**Example 2.3.** We have  $\mathfrak{b}_4(14232) = 14323$  and  $\mathfrak{c}_2(14232) = 12432$ , while  $\mathfrak{b}_2(14232) = 14232$  because the factor 142 does not support a braid move, and  $\mathfrak{c}_4(14232) = 14232$  because the factor 32 does not support a commutation move.

It is sometimes relevant to know whether a pair of Coxeter moves can act on intersecting factors in a reduced word. Note that it is impossible for a reduced word to support both  $\mathfrak{b}_i$  and  $\mathfrak{b}_{i+1}$ .

**Definition 2.4.** If a reduced word **u** supports both  $\mathfrak{b}_i$  and  $\mathfrak{b}_j$  for  $i \neq j$ , then  $\mathfrak{b}_i$  and  $\mathfrak{b}_j$  act as an overlapping pair if |i - j| = 2, and act independently otherwise. If a reduced word **u** supports both  $\mathfrak{c}_i$  and  $\mathfrak{c}_j$  for  $i \neq j$ , then  $\mathfrak{c}_i$  and  $\mathfrak{c}_j$  act as an overlapping pair if |i - j| = 1, and act independently otherwise.

**Example 2.5.** In the reduced word  $1216343 \in R([3254176])$ , the braid moves  $\mathfrak{b}_2$  and  $\mathfrak{b}_6$  act independently, and the commutation moves  $\mathfrak{c}_3$  and  $\mathfrak{c}_4$  act as an overlapping pair.

## 3. Observations about braid and commutation classes

In this section, we highlight certain known or straightforward results about braid classes, commutation classes, and the relationships between them.

The reduced words of a permutation w can be drawn as the vertices of a graph G(w), where two words are connected by a *commutation edge*, or *c-edge*, if they differ by a commutation move, and by a *braid edge*, or *b-edge*, if they differ by a braid move. **Example 3.1.** In the following graph of reduced words for the permutation [25314], c-edges are indicated by solid lines and b-edges are indicated by dashed lines.



The graph G(w) has nice properties, many of which were studied by Reiner and Roichman in [RR13]. We will use the fact that this graph is connected in a critical way, and so we record this property separately in the following theorem.

**Theorem 3.2** (See, for example, [BB05, Theorem 3.3.1]). The graph G(w) is connected.

Let  $G_c(w)$  be the graph that results from contracting the c-edges of G(w), and let  $G_b(w)$ be the graph that results from contracting the b-edges of G(w). The vertices of these two new graphs correspond to the commutation classes C(w) and the braid classes B(w), respectively. Two commutation (respectively, braid) classes are adjacent if an element of one can be obtained from an element of the other by a braid (respectively, commutation) move.

# Example 3.3.



In general, the graphs  $G_c(w)$  and  $G_b(w)$ , along with their analogues in other types, share a certain structural feature.

**Theorem 3.4** ([BCL15, Theorem 3.1]). The graphs  $G_c(w)$  and  $G_b(w)$  are bipartite.

For any permutation w, the braid moves supported by a reduced word  $\mathbf{u} \in R(w)$  must either be entirely disjoint, or may overlap in a single letter. Because commutations are not allowed within a braid class  $B \in B(w)$ , this forces a certain structure to the elements of such a set B.

**Proposition 3.5.** For any permutation w and any  $B \in B(w)$ , we have

$$|B| = 2^x 3^y$$

for some positive integers x and y such that  $3x + 5y \leq \ell(w)$ . Moreover, for any  $B \in B(w)$ , considered as a subgraph  $B \subseteq G(w)$ , the collection B is the following product of path graphs.



*Proof.* Suppose that a reduced word **u** supports distinct braid moves  $\mathfrak{b}_i$  and  $\mathfrak{b}_j$  for 0 < j - i < 3. Because **u** is reduced, we cannot have j - i equal to 1. Thus j - i = 2 and so the factor  $u_{i-1}u_iu_{i+1}u_{i+2}u_{i+3}$  of **u** is of the form  $a(a \pm 1)a(a \mp 1)a$  for some value a. In order for **u** to be reduced, it is impossible for **u** to support additional braid moves  $\mathfrak{b}_k$  for any  $k \in \{i-2, i-1, i+3, i+4\}$ . Therefore braid moves within the class  $B \in B(w)$ , by which all elements of B are related, either act independently, or in overlapping pairs. The number of letters,  $\ell(w)$ , in a reduced word for w, restricts how many independent braid moves (say, x) and overlapping pairs of braid moves (say, y) are possible, because each independent braid move suitilizes three letters in the word, and each overlapping pair utilizes five letters in the word. Therefore

$$3x + 5y \le \ell(w)$$

The braid moves supported by a non-overlapped factor  $a(a \pm 1)a$  can each appear in two presentations:

$$a(a \pm 1)a$$
 and  $(a \pm 1)a(a \pm 1)$ ,

and is represented in the graph G(w) by a path with two nodes. Each overlapped factor  $a(a \pm 1)a(a \mp 1)a$  has exactly three presentations:

$$a(a \pm 1)a(a \mp 1)a, (a \pm 1)a(a \pm 1)(a \mp 1)a, \text{ and } a(a \pm 1)(a \mp 1)a(a \pm 1)$$

and is represented in the graph G(w) by a path with three nodes. Therefore  $|B| = 2^x 3^y$ , and the independence of these factors means that the structure of B is modeled by the product of all such path graphs.

**Example 3.6.** The braid class containing the reduced word  $12143465676 \in R([32547861])$  has  $2^2 \cdot 3^1$  elements, as depicted below.



Recall that the goal of this paper is to leverage the fact that braid classes and commutation classes are partitions of the same set, so that we can study them simultaneously in relation to the collection of reduced words of a permutation. The first step in this process is to recognize that braid moves and commutation moves are, in a sense, orthogonal to each other. Proposition 3.7 recovers the result [Zol94, Corollary 10], proved here in a different manner.

**Proposition 3.7.** Fix a permutation w. For all  $B \in B(w)$  and  $C \in C(w)$ ,

$$|B \cap C| \le 1.$$

*Proof.* Suppose that  $\mathbf{u} \neq \mathbf{v}$  are two reduced words for w and that  $\mathbf{u}$  and  $\mathbf{v}$  are in the same braid class. Since these two words are distinct, we must have done at least one braid move  $i(i+1)i \longleftrightarrow (i+1)i(i+1)$  to get between tehm. It follows that the subword obtained from  $\mathbf{u}$  by restricting to letters "i" and "i + 1" is not the same as the subword obtained from  $\mathbf{v}$  using this same restriction. Since commutation relations cannot change the subword formed by reading only the occurrences of i and (i + 1), the reduced words  $\mathbf{u}$  and  $\mathbf{v}$  are not in the same commutation class.

We can classify all permutations that fall into the extreme cases when either |C(w)| = 1or |B(w)| = 1, and these classifications will be relevant in the next section. We list these classifications as propositions, noting that first is essentially a definition (see [Ste96]).

Characterizing |C(w)| = 1 references a pattern avoidance property, which we pause to define. A permutation  $w \in S_n$  is 321-avoiding if there is no triple of indices, i < j < k, for which w(i) > w(j) > w(k).

**Proposition 3.8** ([BJS93, Theorem 2.1]). |C(w)| = 1 if and only if w is fully commutative; equivalently, if and only if w is 321-avoiding.

Characterizing |B(w)| = 1 is also straightforward because this strict condition requires that any two reduced words be related by braid moves and that no commutation moves are ever possible.

**Proposition 3.9.** |B(w)| = 1 if and only if w has a reduced word of the form i(i+1)i, or of the form  $i(i+\varepsilon)(i+2\varepsilon)\cdots(i+k\varepsilon)$  for some  $\varepsilon \in \{\pm 1\}$  and  $k \ge 0$ ; equivalently, if and only if any two inversions in w share a letter.

*Proof.* By Proposition 3.7, reduced words for w cannot support commutation moves, meaning that all letters must be consecutive in every reduced word for w. The only possibilities, then, are reduced words of the form

$$i(i+\varepsilon)(i+2\varepsilon)\cdots(i+k\varepsilon)$$

for  $\varepsilon \in \{\pm 1\}$  and some  $k \ge 0$ , or

$$i(i+1)i$$
.

This is equivalent to requiring that any two inversions in w share a letter.

## Example 3.10.

(a) The permutation  $[241563] \in S_6$  is 321-avoiding. Thus it is fully commutative and |C([241563])| = 1. More precisely,

 $C([241563]) = \{\{13245, 31245, 13425, 13452, 31425, 31452, 34125, 34152\}\}.$ 

(b) In the permutation  $[124563] \in S_6$ , any two inversions share at least one letter. Thus |B([124563])| = 1. More precisely,

$$B([124563]) = \{\{345\}\}.$$

## 4. Charting reduced words and bounding sizes

Throughout this section, fix a permutation w. The primary tool for our investigation is an organizational scheme for the reduced words in R(w). This relies on the fact that the braid classes B(w) and the commutation classes C(w) give two different partitions of the collection R(w) of reduced words of a permutation.

4.1. Bounds on the number of reduced words. In this subsection, we develop bounds and relationships among the sizes of R(w), B(w), and C(w). These bounds are sharp, and the subsequent subsections will be devoted to characterizing when the upper and lower bounds are each achieved.

**Definition 4.1.** Fix a permutation w, and let  $\mathcal{T}(w)$  be a table whose rows are indexed by B(w), and whose columns are indexed by C(w). For each  $\mathbf{u} \in R(w)$ , if  $\mathbf{u} \in B \in B(w)$  and  $\mathbf{u} \in C \in C(w)$ , then place  $\mathbf{u}$  in row B and column C of the table  $\mathcal{T}(w)$ .

Of course, the choice of order for the rows and columns of  $\mathcal{T}(w)$  has an impact on the table itself, but it has no effect on the properties we will discuss (and care about) below.

**Example 4.2.** Let w = [25314]. The braid classes and commutation classes of this element were discussed in previous examples, and we now label them as shown.

$$B_{1} = \{12432\}$$

$$B_{2} = \{14232, 14323\}$$

$$B_{3} = \{41232, 41323\}$$

$$B_{4} = \{43123\}$$

$$C_{1} = \{12432, 14232, 41232\}$$

$$C_{2} = \{14323, 41323, 43123\}$$

The corresponding table,  $\mathcal{T}(w)$ , for elements of R(w) is as follows.

|       | $C_1$ | $C_2$ |  |  |
|-------|-------|-------|--|--|
| $B_1$ | 12432 |       |  |  |
| $B_2$ | 14232 | 14323 |  |  |
| $B_3$ | 41232 | 41323 |  |  |
| $B_4$ |       | 43123 |  |  |

As is obvious in Example 4.2, some cells of  $\mathcal{T}(w)$  may be empty. On the other hand, there is a limit to how many reduced words each cell might contain.

**Corollary 4.3.** Each cell in  $\mathcal{T}(w)$  contains at most one element.

*Proof.* This is equivalent to Proposition 3.7.

Certain features of these charts dictate how many cells might be empty (put another way, how many cells must be nonempty). Since these features do not care about the type of objects (in our case, reduced words) filling the cells of the charts, we state and prove this result in greater generality.

**Proposition 4.4.** Let T be an  $r \times c$  rectangular array of data, some of whose cells may be empty. Suppose that:

- each column of T contains at least one nonempty cell,
- each row of T contains at least one nonempty cell, and
- one can get from any nonempty cell in T to any other nonempty cell by a sequence of in-row or in-column jumps, always from nonempty cells to nonempty cells.

Then T contains at least c + r - 1 nonempty cells.

*Proof.* Without loss of generality, assume that  $c \ge r$ . Then, by the first requirement, the array T must have at least c nonempty cells.

If r > 1, then the remaining requirements for T imply that there must be a column of T with multiple nonempty cells. In fact, in order to be able to reach nonempty cells in all r rows, there must be x columns of the array in which one may jump, and these x columns must contain at least r + x - 1 nonempty cells.

Because we initially accounted for one nonempty cell in each column, we now have that T must contain at least

$$c + (r + x - 1) - x = c + r - 1$$

nonempty cells.

**Example 4.5.** Proposition 4.4 is demonstrated in Figure 1, which gives a qualifying  $7 \times 9$  array with 15 nonempty cells. Note that a violation of at least one of the three requirements

|   |   |   |   |   |   | • |   |   |
|---|---|---|---|---|---|---|---|---|
| ٠ |   |   |   |   |   |   |   |   |
|   |   |   | • |   |   | • |   |   |
| • |   | • |   | • |   |   |   |   |
|   |   | • |   |   | • |   | • |   |
|   |   | • | ٠ |   |   |   |   |   |
|   | ٠ |   |   |   |   |   | • | ٠ |

FIGURE 1. A  $7 \times 9$  array satisfying the requirements of Proposition 4.4 and containing 15 = 7 + 9 - 1 nonempty cells.

of Proposition 4.4 would occur if any of the nonempty cells in this array were to be made empty. For example, if the cell in the fifth row and eighth column were to be made empty, then there would be no way to get from the bottom right nonempty cell to any nonempty cell outside of its row by a legal sequence of in-row and in-column jumps.

Having organized the elements of R(w) into a chart  $\mathcal{T}(w)$  satisfying the requirements of Proposition 4.4, we immediately obtain the following bounds.

**Theorem 4.6.** For any permutation w,

$$|B(w)| + |C(w)| - 1 \le |R(w)| \le |B(w)| \cdot |C(w)|.$$

*Proof.* The upper bound in this result is an immediate consequence of the definition of  $\mathcal{T}(w)$  and of Corollary 4.3.

To prove the lower bound, first note that by definition of  $\mathcal{T}(w)$ , each row contains at least one nonempty cell, as does each column, and Theorem 3.2 says that one can travel from any nonempty cell of the chart to any other nonempty cell of the chart by traveling only within rows and within columns, and only jumping from nonempty cells to nonempty cells. Thus the chart  $\mathcal{T}(w)$  satisfies the requirements of Proposition 4.4. Corollary 4.3 implies that the number of nonempty cells in  $\mathcal{T}(w)$  is exactly |R(w)|, and hence Proposition 4.4 gives

$$|R(w)| \ge |B(w)| + |C(w)| - 1.$$

It is easy to see that these bounds are sharp, because they are simultaneously achieved by any fully commutative w. More precisely, if w is fully commutative, then |C(w)| = 1 and |B(w)| = |R(w)| by Proposition 3.7, and both bounds of Theorem 4.6 are achieved. The bounds are similarly realized by any w for which |B(w)| = 1.

We devote the next two subsections to characterizing when the upper and lower bounds of Theorem 4.6 are achieved.

4.2. Characterizing the upper bound. We first turn our attention to the upper bound of Theorem 4.6. As discussed at the end of the last subsection, fully commutative permutations (for which |C(w)| = 1), achieve this bound. The next lemma shows that if either |C(w)| or |B(w)| is equal to 1, then the other cardinality must be equal to |R(w)|.

**Lemma 4.7.** Fix a permutation w and suppose that |C(w)| = 1 (respectively, |B(w)| = 1). Then |B(w)| = |R(w)| (respectively, |C(w)| = |R(w)|).

*Proof.* Suppose that |C(w)| = 1. Then by Theorem 4.6,

$$|B(w)| + 1 - 1 \le |R(w)| \le |B(w)| \cdot 1.$$

Hence |B(w)| = |R(w)|. The hypothesis |B(w)| = 1 proceeds analogously.

Therefore, by Lemma 4.7, if either |C(w)| or |B(w)| is equal to 1, then the upper bound of Theorem 4.6 is automatically achieved. In fact, as we prove below, these are the only permutations that will achieve that upper bound.

**Proposition 4.8.**  $|R(w)| = |B(w)| \cdot |C(w)|$  if and only if |B(w)| = 1 or |C(w)| = 1.

*Proof.* If |B(w)| = 1 or |C(w)| = 1, then Lemma 4.7 immediately implies  $|R(w)| = |B(w)| \cdot |C(w)|$ . This proves one direction of the proposition.

We prove the other direction of the proposition by contrapositive. Suppose that |C(w)| > 1 and |B(w)| > 1. Since w has more than one commutation class, this w is not fully commutative, meaning that there is some reduced word for w that contains a factor of the form i(i + 1)i. Choose such a reduced word, **u**, so that the factor i(i + 1)i is as far to the left in **u** as possible. Since w has more than one braid class, **u** must contain at least one additional letter. In particular, we have, without loss of generality, that

(1) 
$$\mathbf{u} = \mathbf{d}ji(i+1)i\mathbf{e},$$

with j > i + 1 and where **d** and **e** are possibly empty subwords.

Let  $B \in B(w)$  and  $C \in C(w)$  be the braid and commutation classes, respectively, containing  $\mathbf{u} \in R(w)$ . From Equation (1), we see that

$$\mathbf{v} = \mathbf{b}_k \mathbf{u} = \mathbf{d}j(i+1)i(i+1)\mathbf{e}$$

is also an element of B, where k is the position of the indicated i + 1 in **u**. Additionally,

$$\mathbf{t} = \mathbf{c}_{k-2}\mathbf{u} = \mathbf{d}ij(i+1)i\mathbf{e}$$

is also an element of C.

By Proposition 3.7, in order to satisfy  $|R(w)| = |B(w)| \cdot |C(w)|$ , the intersection of each braid class with each commutation class must be exactly one reduced word. Let  $\mathbf{v} \in C' \in C(w)$  and  $\mathbf{t} \in B' \in B(w)$ . Thus there must be exactly one reduced word  $\mathbf{a} \in B' \cap C'$ , in the same commutation class as  $\mathbf{v}$  and the same braid class as  $\mathbf{t}$ . In other words,  $\mathbf{a} = \mathfrak{c}_{i_1} \cdots \mathfrak{c}_{i_m} \mathfrak{b}_k \mathbf{u} = \mathfrak{b}_{i_1} \cdots \mathfrak{b}_{i_p} \mathfrak{c}_{k-2} \mathbf{u}$ , for some m and p.

As in the proof of Proposition 3.7, consider the subword of **a** obtained by restricting to the entries i and i + 1. Because **a** and **v** differ by commutation moves, this subword for **a** must be equal to the analogous subword for **v**. By choice of **u**, no factor of the form i(i+1)ican be produced in the prefix dij(i+1) of **t** by Coxeter moves. Therefore any braid moves applied to **t** will not change the prefix of the  $\{i, i+1\}$  subword formed by using only the letters i and i + 1, meaning that this subword can never be identical to the corresponding subword for **v**. Therefore there is no such **a**, and  $B' \cap C' = \emptyset$ . Hence  $|R(w)| \neq |B(w)| \cdot |C(w)|$ , concluding the proof.

The narrowity of Proposition 4.8 actually implies that those permutations achieving the upper bound of Theorem 4.6 are also among those that achieve the lower bound of Theorem 4.6.

## **Corollary 4.9.** If |B(w)| = 1 or |C(w)| = 1, then

$$|B(w)| + |C(w)| - 1 = |R(w)| = |B(w)| \cdot |C(w)|.$$

Note that almost all of the permutations described in Proposition 3.9 are fully commutative. In fact, the only elements with a single braid class that are not also fully commutative are those for which  $R(w) = \{i(i+1)i, (i+1)i(i+1)\}$  for some *i*. Thus we can reformulate Proposition 4.8 as follows.

**Corollary 4.10.**  $|R(w)| = |B(w)| \cdot |C(w)|$  if and only if

- w is fully commutative, or
- $R(w) = \{i(i+1)i, (i+1)i(i+1)\}$  for some *i*; equivalently, w(i) = i+2 and w(i+2) = i for some *i*, and all other values are fixed by *w*.

The characterization in the previous corollary enables us to enumerate the permutations in  $S_n$  achieving the upper bound of Theorem 4.6, for any n.

**Corollary 4.11.** The number of permutations in  $S_n$  for which  $|R(w)| = |B(w)| \cdot |C(w)|$  is equal to

$$\begin{cases} 1 & \text{if } n = 1, \text{ and} \\ C_n + n - 2 & \text{if } n > 1, \end{cases}$$

where  $C_n = \binom{2n}{n}/(n+1)$  is the *n*th Catalan number.

*Proof.* For n = 1, one of the categories given in Corollary 4.10 is not feasible, and the enumeration is quick to do by hand.

Now assume that n > 1. First observe that the two possibilities of Corollary 4.10 are mutually exclusive. Proposition 3.8 says that fully commutative permutations are exactly those that avoid the pattern 321, and it is well-known that the number of 321-avoiding permutations in  $S_n$  is the *n*th Catalan number  $C_n$ ; see [BJS93], for example. It remains to count the permutations  $w \in S_n$  for which  $R(w) = \{i(i+1)i, (i+1)i(i+1)\}$ , and these can be described by choosing the letter  $i \in [1, n-2]$ .

This is sequence A290953 in [Slo96].

4.3. Characterizing the lower bound. Characterizing which permutations achieve the lower bound of Theorem 4.6 has a noticeably different flavor from that of the upper bound characterization. Instead of two classes of permutations, one of which is broad and one of which is quite specific, the lower bound achievers fall into more classes, all of which are highly specified.

**Definition 4.12.** The collection of reduced words R(w) has a *circuit* if there exists a sequence of elements in R(w), say

 $\mathbf{u_0}, \mathbf{u_1}, \mathbf{u_2}, \dots, \mathbf{u_{2t}} = \mathbf{u_0}$ 

for some integer t such that for all s, either

$$\begin{cases} \mathbf{u_{2s}}, \mathbf{u_{2s+1}} \in B_s \text{ and} \\ \mathbf{u_{2s+1}}, \mathbf{u_{2s+2}} \in C_s \end{cases}$$
$$\begin{cases} \mathbf{u_{2s}}, \mathbf{u_{2s+1}} \in C_s \text{ and} \\ \mathbf{u_{2s+1}}, \mathbf{u_{2s+2}} \in B_s \end{cases}$$

or

where  $B_s \in B(w)$  and  $C_s \in C(w)$ . Equivalently, define a graph  $\Gamma(w)$  whose vertices are  $B(w) \cup C(w)$ , with an edge between  $B_i$  and  $C_j$  if and only if  $B_i \cap C_j \neq \emptyset$ ; then R(w) has a circuit if the graph  $\Gamma(w)$  contains a circuit, i.e., if there is some vertex v in  $\Gamma(w)$  such that one can travel from v to itself following a nontrivial sequence of edges and vertices. Equivalently, R(w) has a circuit if and only if  $\Gamma(w)$  is not a tree.

**Definition 4.13.** A permutation w is *circuit-free* if its collection R(w) of reduced words does not have a circuit.

Avoidance of circuits in the set of reduced words completely characterizes those permutations that achieve the lower bound of Theorem 4.6, because having such a circuit would indicate that there are more reduced words than are minimally required.

**Proposition 4.14.** A permutation w is circuit-free if and only if

$$|R(w)| = |B(w)| + |C(w)| - 1.$$

The main idea in the proof of Proposition 4.14 is that R(w) has a circuit if and only if removing a reduced word from a chart  $\mathcal{T}(w)$  leaves a chart that still satisfies the requirements of Proposition 4.4. In other words, R(w) contains a circuit if and only if one can detour around a given reduced word in a chart  $\mathcal{T}(w)$  without violating the rules of Proposition 4.4. Proof of Proposition 4.14. Recall that  $\Gamma(w)$  is the graph whose vertices are  $B(w) \cup C(w)$ , with an edge between  $B_i$  and  $C_j$  if and only if  $B_i \cap C_j \neq \emptyset$ . By Theorem 3.2, this graph is connected. Recall also that because R(w) has a circuit if and only if the graph  $\Gamma(w)$  has a circuit, the permutation w is circuit-free if and only if  $\Gamma(w)$  is a tree.

Recall that a finite connected graph is a tree if and only if it has one fewer edge than vertices. Due to Proposition 3.7, each edge in this graph corresponds to exactly one reduced word for w, and each reduced word for w corresponds to exactly one edge in this graph. Therefore the number of edges is |R(w)|. The number of vertices in the graph  $\Gamma(w)$  is |B(w)| + |C(w)| by definition.

Thus we have that w is circuit-free if and only if  $\Gamma(w)$  is a tree, and this graph is a tree if and only if

$$|R(w)| = |B(w)| + |C(w)| - 1,$$

which means precisely that w achieves the lower bound of Theorem 4.6.

We can now exploit the idea of circuit-free permutations to explicitly describe the permutations achieving the lower bound of Theorem 4.6.

**Proposition 4.15.** A permutation w is circuit-free (that is, it achieves the lower bound of Theorem 4.6) if and only if

- |B(w)| = 1, or
- |C(w)| = 1, or
- $\mathbf{u}i(i+1)i\mathbf{v} \in R(w)$  such that, up to symmetry, one of the following conditions holds:

\* 
$$\mathbf{u} = \emptyset$$
 and  $\mathbf{v} \in \{(i-1)i, (i-1)(i-2)\cdots(i-1-t)\}$  for some  $t \ge 0$ , or

$$\star \mathbf{u} = \mathbf{v} = i - 1$$
, or

\*  $\mathbf{u} = (i-1-t)\cdots(i-2)(i-1)$  and  $\mathbf{v} = (i+2)(i+3)\cdots(i+2+t')$  for some  $t, t' \ge 0$ ,

where the allowable symmetries are those that arise from the standard operations of reversal, complementation, and reverse complementation.

*Proof.* First note that if |B(w)| = 1 or |C(w)| = 1, then w is circuit-free by Corollary 4.9 and Proposition 4.14.

Now assume that w is circuit-free, and that |B(w)| > 1 and |C(w)| > 1. Then w has a reduced word of the form

$$\mathbf{u}i(i+1)i\mathbf{v}$$

where  $\mathbf{u}$  and  $\mathbf{v}$  are could each (but not both) be empty subwords.

If **u** supports a commutation move, say  $c_k \mathbf{u} = \mathbf{u}'$ , then the four words

$$\mathbf{u}i(i+1)i\mathbf{v}, \ \mathbf{u}(i+1)i(i+1)\mathbf{v}, \ \mathbf{u}'(i+1)i(i+1)\mathbf{v}, \ \mathbf{u}'i(i+1)i\mathbf{v}$$

form a circuit in R(w). Thus **u** (and, analogously, **v**) cannot support any commutation moves, meaning that adjacent letters are consecutive.

If **u** has a reduced word of the form  $\mathbf{u} = \overline{\mathbf{u}}x$ , where x is a letter that commutes with both i and i + 1 (that is,  $x \notin \{i - 1, i, i + 1, i + 2\}$ ), then this would also produce a circuit, consisting of the four elements

$$\overline{\mathbf{u}}xi(i+1)i\mathbf{v}\,,\ \overline{\mathbf{u}}i(i+1)ix\mathbf{v}\,,\ \overline{\mathbf{u}}(i+1)i(i+1)x\mathbf{v}\,,\ \overline{\mathbf{u}}x(i+1)i(i+1)\mathbf{v}$$

in R(w). Thus if **u** is nonempty, then its rightmost letter is either i - 1 or i + 2. Similarly, if **v** is nonempty, then its leftmost letter is also either i - 1 or i + 2.

To summarize these findings so far, if **u** (respectively, **v**) is nonempty, then its letters must be sequentially consecutive with rightmost (resp., leftmost) letter equal to i-1 or i+2. Call this **Rule 1**.

If **u** is non-monotonic, then it supports a braid move, say  $\mathbf{b}_k \mathbf{u} = \mathbf{u}'$ . Then  $\mathbf{u}'i(i+1)i\mathbf{v} \in R(w)$  and the nonempty word  $\mathbf{u}'$  does not satisfy **Rule 1**: either it is not sequentially consecutive, or its rightmost letter is not equal to i-1 or i+2. Thus **u** and **v** must each be monotonic. Call this **Rule 2**.

If  $\mathbf{u} = (i+t)\cdots i(i-1)$ , then we claim that  $\mathbf{u}$  must contain at most two letters. Suppose not; that is, suppose that  $t \ge 1$ , and let  $\overline{\mathbf{u}} = \mathbf{b}_{t+2}\mathbf{u}$ . Then R(w) would have a circuit because

 $(\mathfrak{c}_{t+3}\mathfrak{c}_t)(\mathfrak{b}_{t+2}\mathfrak{b}_{t+4})(\mathfrak{c}_{t+2})(\mathfrak{b}_{t+1}\mathfrak{b}_{t+3})(\mathfrak{c}_{t+4}\mathfrak{c}_{t+1})(\mathfrak{b}_{t+3}\mathfrak{b}_{t+1})(\mathfrak{c}_{t+2})(\mathfrak{b}_{t+4}\mathfrak{b}_{t+2})\overline{\mathbf{u}}=\overline{\mathbf{u}}.$ 

Therefore, by this and symmetric arguments, we conclude that if the second rightmost (respectively, leftmost) letter of  $\mathbf{u}$  (resp.,  $\mathbf{v}$ ) is i or i + 1, then  $\mathbf{u}$  (resp.,  $\mathbf{v}$ ) has no additional letters. Call this **Rule 3**.

We now use **Rules 1–3** to show that a reduced word for w must have one of the forms given in the statement of the proposition. For each of those, we check using Proposition 4.14 that the resulting permutation w is indeed circuit-free, which will conclude the proof. To this end, consider the following three cases, which are exhaustive, up to symmetry.

- (i) **u** and **v** each have at most one letter. The possible reduced words for w are as follows.
  - (a) i(i+1)i(i-1) and symmetric cases

|R(w)| = 3, |B(w)| = 2, |C(w)| = 2, and 3 = 2 + 2 - 1.

(b) (i-1)i(i+1)i(i-1) and symmetric cases

|R(w)| = 6, |B(w)| = 4, |C(w)| = 3, and 6 = 4 + 3 - 1.

- (c) (i+2)i(i+1)i(i-1) and symmetric cases
- |R(w)| = 4, |B(w)| = 3, |C(w)| = 2, and 4 = 3 + 2 1.
- (ii) Without loss of generality,  $\mathbf{u} = \emptyset$  and  $\mathbf{v}$  has at least two letters. The possible reduced words for w are as follows.
  - (a) i(i+1)i(i-1)i and symmetric cases
    - |R(w)| = 5, |B(w)| = 3, |C(w)| = 3, and 5 = 3 + 3 1.
  - (b)  $i(i+1)i(i-1)\cdots(i-1-t)$  and symmetric cases

|R(w)| = t + 3, |B(w)| = t + 2, |C(w)| = 2, and t + 3 = t + 2 + 2 - 1.

- (iii)  $\mathbf{u}$  and  $\mathbf{v}$  are both nonempty and, without loss of generality,  $\mathbf{u}$  has at least two letters. The possible reduced words for w are as follows.
  - (a)  $i(i-1)i(i+1)ix\overline{\mathbf{v}}$ , where  $x \in \{i-1, i+2\}$ , and symmetric cases After a braid move, we have  $(i-1)i(i-1)(i+1)ix\overline{\mathbf{v}} \in R(w)$ . By **Rule 1** applied to the bracketed subword, the letters of  $\mathbf{v}' = (i+1)ix\overline{\mathbf{v}}$  must be sequentially consecutive, which means that x = i - 1. On the other hand, the subword  $\mathbf{v}' = (i+1)i(i-1)\overline{\mathbf{v}}$  now violates **Rule 3**, since  $\mathbf{v}'$  has more than two letters. This contradiction implies that all permutations of this form are *not* circuit-free.
  - (b)  $\overline{\mathbf{u}}(i-2)(i-1)i(i+1)i(i-1)\overline{\mathbf{v}}$  and symmetric cases After a braid move and two commutation moves, we can write  $\overline{\mathbf{u}}(i-2)(i+1)(i-1)i(i-1)(i+1)\overline{\mathbf{v}} \in R(w)$ , breaking **Rule 1** since the subword  $\mathbf{u}' =$

 $\overline{\mathbf{u}}(i-2)(i+1)$  does not have sequentially consecutive letters. Therefore, these permutations are *not* circuit-free.

(c)  $(i-1-t)\cdots(i-1)i(i+1)i(i+2)\cdots(i+2+t')$  for  $t \ge 1$  and  $t' \ge 0$ , and symmetric cases |R(w)| = t+t'+4, |B(w)| = t+t'+3, |C(w)| = 2, and t+t'+4 = t+t'+3+2-1.

Having characterized the permutations that achieve the lower bound of Theorem 4.6, we now enumerate this class. That is, we count the permutations characterized by Proposition 4.15. This will provide an analogue to the enumeration of upper bound achievers given in Corollary 4.11.

**Corollary 4.16.** The number of permutations in  $S_n$  for which the lower bound of Theorem 4.6 is achieved is

$$\begin{cases} n & \text{if } n \le 2, \text{ and} \\ C_n + \frac{n^3 - 3n^2 + 8n - 21}{3} & \text{if } n > 2, \end{cases}$$

where  $C_n = \binom{2n}{n}/(n+1)$  is the *n*th Catalan number.

*Proof.* For  $n \in \{1, 2\}$ , many of the categories given in Proposition 4.15 are not feasible, and the enumeration is quick to do by hand.

Now assume that n > 2.

The first two options described in bullet points in the characterization given in Proposition 4.15 are exactly those permutations that (also) achieve the upper bound of Theorem 4.6, and these were counted in Corollary 4.11 as  $C_n + n - 2$ .

Note that there is no overlap between the permutations w for which |B(w)| = 1 or |C(w)| = 1 and those permutations satisfying the third bullet point in the statement of Proposition 4.15. Thus if we can enumerate the permutations satisfying that third bullet point and add  $C_n + n - 2$ , then we will have the desired total. In other words, it remains only to enumerate permutations w with  $\mathbf{u}i(i+1)i\mathbf{v} \in R(w)$  such that, up to symmetry, one of the following conditions holds:

Option 1: 
$$\mathbf{u} = \emptyset$$
 and  $\mathbf{v} \in \{(i-1)i, (i-1)(i-2)\cdots(i-1-t)\}$  for some  $t \ge 0$ , or

Option 2: u = v = i - 1, or

Option 3: 
$$\mathbf{u} = (i-1-t)\cdots(i-2)(i-1)$$
 and  $\mathbf{v} = (i+2)(i+3)\cdots(i+2+t')$  for  
some  $t, t' \ge 0$ .

It is straightforward to check that these three options are disjoint (indeed, as designed). Thus the enumeration we want can be obtained by adding the number of permutations in each of the three categories.

**Option 1:** Including all symmetries, the permutations in this category have reduced words of one of the following forms, where the required braid move has been bracketed for clarity.

- i(i+1)i(i-1)i, which can be rewritten in the form j(j-1)j(j+1)j
- $\overline{i(i+1)i(i+2)(i+1)}$ , which can be rewritten in the form (j+1)(j+2)j(j+1)j

• 
$$\overline{i(i+1)i(i-1)(i-2)\cdots(i-1-t)}$$

•  $\overline{i(i+1)i(i+2)(i+3)\cdots(i+2+t)}$ 

- $(i-1-t)\cdots(i-2)(i-1)\frac{i(i+1)i}{i(i+2)i}$   $(i+2+t)\cdots(i+3)(i+2)\frac{i(i+1)i}{i(i+1)i}$

Other than the equivalences noted in the first two bullet points above, none of these words can be transformed into one of the other configurations by a sequence of braid and commutation moves. There are n-3 choices of i for each of the first two categories, all of which yield different permutations, contributing 2(n-3) permutations to the total. Now consider the remaining four categories. For each  $i \in [2, n-2]$ , we can choose any  $t \in [0, i-2]$ , which gives

$$\sum_{i=2}^{n-2} (i-1) = \frac{(n-2)(n-3)}{2}$$

choices for (i, t). Each choice yields a distinct permutation, meaning that these four categories contirbute 2(n-2)(n-3) permutations to the total. Therefore, the number of permutations satisfying Option 1 is

$$2(n-3) + 2(n-2)(n-3) = 2(n-3)(n-1).$$

**Option 2:** Including all symmetries, the permutations in this category have reduced words of the following form, where the required braid move has been bracketed for clarity.

• (i-1)i(i+1)i(i-1), which can be rewritten in the form (j+2)j(j+1)j(j+2)

Since there are n-3 choices for i (or j), the number of permutations satisfying Option 2 is

n - 3.

**Option 3:** Including all symmetries, the permutations in this category have reduced words of one of the following forms, where the required braid move has been bracketed for clarity.

- $(i-1-t)\cdots(i-2)(i-1)\frac{i(i+1)i(i+2)(i+3)\cdots(i+2+t')}{i(i+2)i(i-1)i(i-2)\cdots(i-1-t)}$

Neither of these words can be transformed into the other configuration by a sequence of braid and commutation moves, and any choice of (i, t, t') identifies a unique permutation in this category. For each  $i \in [2, n-3]$ , we need  $t \in [0, i-2]$  and  $t' \in [0, n-3-i]$ . Thus, for a fixed i, there are (i-1)(n-2-i) distinct permutations. It follows that, in total, the number of permutations satisfying Option 3 is

$$2\sum_{i=2}^{n-3} (i-1)(n-2-i) = 2\left[(n-2)\sum_{i=2}^{n-3} (i-1) - \sum_{i=2}^{n-3} i^2 + \sum_{i=2}^{n-3} i\right]$$
$$= (n-2)(n-3)(n-4) - \frac{(n-3)(n-2)(2n-5)}{3} + (n-3)(n-2)$$
$$= \frac{(n-2)(n-3)(n-4)}{3}.$$

Therefore, the number of permutations achieving the lower bound of Theorem 4.6 is

$$C_n + n - 2 + 2(n - 3)(n - 1) + n - 3 + \frac{(n - 2)(n - 3)(n - 4)}{3},$$

which simplifies to the desired result.

This is sequence A290954 in [Slo96].

### 5. A proposed connection to weak order

Proposition 4.15 gives a characterization of permutations  $w \in S_n$  that achieve the lower bound of Theorem 4.6 by describing their possible reduced words. Based on experimentation in Maple, it also appears that the shape of the interval  $[e, w] = \{u \in S_n \mid e \leq u \leq w\}$  in weak order provides an alternative characterization. Roughly speaking, permutations whose intervals have a narrow shape achieve the lower bound in Theorem 4.6, whereas permutations with a wider interval structure have too many reduced words to achieve this lower bound. We formalize this observation in Conjecture 5.2 below.

We require several additional definitions in order to state our conjecture.

**Definition 5.1.** The weak order on Coxeter groups also appears naturally in the study of reduced words. Let u and v be elements of a finitely generated Coxeter group W. Then  $u \leq v$  in (right) weak order if there is a sequence of simple reflections  $s_{i_1}, \ldots, s_{i_k}$  such that  $us_{i_1}s_{i_2}\cdots s_{i_k} = v$  and  $\ell(us_{i_1}\cdots s_{i_j}) + 1 = \ell(us_{i_1}\cdots s_{i_j+1})$  for all  $j = 1, \ldots, k-1$ . Each reduced expression for w then corresponds to a maximal chain from e to w in the weak order on W, and vice versa.

The set of indices appearing in any, equivalently every, reduced word for w is the *support* of w, and we denote the *size* of this support by  $\sup(w)$ . The poset [e, w] is ranked, and we define  $r_i$  to be the number of elements at rank i, where  $i \in \{0, \ldots, \ell(w)\}$ . The *width* of w is the maximum of  $\{r_0, r_1, \ldots, r_{\ell(w)}\}$ , which we denote by wid(w). Figure 2 provides several examples that calculate the width and size of the support of three different permutations.



FIGURE 2. This figure shows the interval [e, w] for three permutations w, where the top element represents one choice from R(w). For the leftmost figure, wid(w) = 3 and sup(w) = 4. In the middle figure, wid $(w) = \sup(w) = 3$ . On the right, wid(w) = 2 while sup(w) = 3.

We now provide a conjectural reformulation of our proposition characterizing the permutations that achieve the lower bound in Theorem 4.6. In particular, we rephrase the conditions on the reduced words articulated in Proposition 4.15 in terms of the width and support of the permutation.

Conjecture 5.2. The permutation w is circuit-free, and thus achieves the lower bound in Theorem 4.6, if and only if at least one of the following conditions is satisfied:

(1) |C(w)| = 1

(2) |B(w)| = 1(3) wid(w) = 2 (4) wid(w) = sup(w) = 3

**Example 5.3.** Consider again the three permutations referenced in Figure 2. The permutation [152463], with reduced word 34532, has width three and support four, so Conjecture 5.2 predicts that it should not achieve the lower bound and indeed we have |R([152463])| = 6, while |B([152463])| + |C([152463])| - 1 = 5. The permutation [3421], with reduced word 12312, satisfies Condition (4) with width and support three, and it achieves the lower bound with |R([3421])| = |B([3421])| + |C([3421])| - 1 = 5. Condition (3) holds for the permutation [4132], with reduced word 2321, and |R([4132])| = |B([4132])| + |C([4132])| - 1 = 3.

Of course, since the characterization in Proposition 4.15 provides an exhaustive and concrete list of possible reduced expressions for permutations that achieve the desired lower bound, it is straightforward to check case by case that each of those permutations satisfies at least one of the width or support criteria in Conjecture 5.2. The more difficult direction seems to be proving that these conditions on the width and support are sufficient in order for the permutation to achieve the lower bound. More generally, it would thus be interesting to develop a conceptual understanding of precisely how the width of the permutation affects the number of reduced words in terms of the number of braid and commutation classes. Having such an interpretation in terms of intervals in the weak order might also offer a natural pathway for generalizing some of the results in this paper to other Coxeter groups.

#### References

- [BB05] Anders Björner and Francesco Brenti. Combinatorics of Coxeter groups, volume 231 of Graduate Texts in Mathematics. Springer, New York, 2005.
- [BCL15] Nantel Bergeron, Cesar Ceballos, and Jean-Philippe Labbé. Fan realizations of type A subword complexes and multi-associahedra of rank 3. *Discrete Comput. Geom.*, 54(1):195–231, 2015.
- [Béd99] Robert Bédard. On commutation classes of reduced words in Weyl groups. *European J. Combin.*, 20(6):483–505, 1999.
- [BJS93] Sara C. Billey, William Jockusch, and Richard P. Stanley. Some combinatorial properties of Schubert polynomials. J. Algebraic Combin., 2(4):345–374, 1993.
- [Eln97] Serge Elnitsky. Rhombic tilings of polygons and classes of reduced words in Coxeter groups. J. Combin. Theory Ser. A, 77(2):193–221, 1997.
- [GL02] R. M. Green and J. Losonczy. Freely braided elements of Coxeter groups. Ann. Comb., 6(3-4):337–348, 2002.
- [HJ10] Christopher R. H. Hanusa and Brant C. Jones. The enumeration of fully commutative affine permutations. *European J. Combin.*, 31(5):1342–1359, 2010.
- [JW09] Jakob Jonsson and Volkmar Welker. Complexes of injective words and their commutation classes. *Pacific J. Math.*, 243(2):313–329, 2009.
- [Men10] D. Meng. Reduced decompositions and commutation classes. arXiv.org/1009.0886, 2010.
- [MS89] Yu. I. Manin and V. V. Schechtman. Arrangements of hyperplanes, higher braid groups and higher Bruhat orders. In Algebraic number theory, volume 17 of Adv. Stud. Pure Math., pages 289–308. Academic Press, Boston, MA, 1989.
- [Rei05] Victor Reiner. Note on the expected number of Yang-Baxter moves applicable to reduced decompositions. European J. Combin., 26(6):1019–1021, 2005.
- [RR13] Victor Reiner and Yuval Roichman. Diameter of graphs of reduced words and galleries. Trans. Amer. Math. Soc., 365(5):2779–2802, 2013.
- [Slo96] Neil James Alexander Sloane. The On-line Encyclopedia of Integer Sequences (OEIS). OEIS Foundation, Incorporated, 1996.

- [Sta84] Richard P. Stanley. On the number of reduced decompositions of elements of Coxeter groups. *European J. Combin.*, 5(4):359–372, 1984.
- [Ste96] John R. Stembridge. On the fully commutative elements of Coxeter groups. J. Algebraic Combin., 5(4):353–385, 1996.
- [Ste97] John R. Stembridge. Some combinatorial aspects of reduced words in finite Coxeter groups. Trans. Amer. Math. Soc., 349(4):1285–1332, 1997.
- [Ste98] John R. Stembridge. The enumeration of fully commutative elements of Coxeter groups. J. Algebraic Combin., 7(3):291–320, 1998.
- [STWW17] Anne Schilling, Nicolas M. Thiéry, Graham White, and Nathan Williams. Braid moves in commutation classes of the symmetric group. European J. Combin., 62:15–34, 2017.
- [Ten06] Bridget Eileen Tenner. Reduced decompositions and permutation patterns. J. Algebraic Combin., 24(3):263–284, 2006.
- [Zie93] Günter M. Ziegler. Higher Bruhat orders and cyclic hyperplane arrangements. *Topology*, 32(2):259–279, 1993.
- [Zol94] Debbie McIntyre Zollinger. Equivalence classes of reduced words. Master's thesis, University of Minnesota, 1994.

SUSANNA FISHEL, SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNI-VERSITY, TEMPE, AZ, USA

*E-mail address*: sfishel1@asu.edu

ELIZABETH MILIĆEVIĆ, DEPARTMENT OF MATHEMATICS & STATISTICS, HAVERFORD COLLEGE, HAVER-FORD, PA, USA

 $E\text{-}mail\ address:\ \texttt{emilicevic@haverford.edu}$ 

REBECCA PATRIAS, LACIM, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, MONTRÉAL (QUÉBEC), CANADA *E-mail address*: patriasr@lacim.ca

BRIDGET EILEEN TENNER, DEPARTMENT OF MATHEMATICAL SCIENCES, DEPAUL UNIVERSITY, CHICAGO, IL, USA

E-mail address: bridget@math.depaul.edu