

From subKautz digraphs to cyclic Kautz digraphs *

C. Dalfo

Departament de Matemàtiques
Universitat Politècnica de Catalunya
Barcelona, Catalonia
cristina.dalfo@upc.edu

Abstract

The Kautz digraphs $K(d, \ell)$ are a well-known family of dense digraphs, widely studied as a good model for interconnection networks. Closely related to these, the cyclic Kautz digraphs $CK(d, \ell)$ were recently introduced by Böhmová, Huemer and the author, and some of its distance-related parameters were fixed. In this paper we propose a new approach to the cyclic Kautz digraphs by introducing the family of the subKautz digraphs $sK(d, \ell)$, from where the cyclic Kautz digraphs can be obtained as line digraphs. This allows us to give exact formulas for the distance between any two vertices of both $sK(d, \ell)$ and $CK(d, \ell)$. Moreover, we compute the diameter and the semigirth of both families, also providing efficient routing algorithms to find the shortest path between any pair of vertices. Using these parameters, we also prove that $sK(d, \ell)$ and $CK(d, \ell)$ are maximally vertex-connected and super-edge-connected. Whereas $K(d, \ell)$ are optimal with respect to the diameter, we show that $sK(d, \ell)$ and $CK(d, \ell)$ are optimal with respect to the mean distance, whose exact values are given for both families when $\ell = 3$. Finally, we provide a lower bound on the girth of $CK(d, \ell)$ and $sK(d, \ell)$.

Mathematics Subject Classifications: 05C20, 05C50.

Keywords: Digraph, distance, diameter, mean distance, routing, Kautz digraph, line digraph, (vertex-)connectivity, edge-connectivity, superconnectivity, semigirth, girth.

*This research is supported by *MINECO* under project MTM2014-60127-P, and the *Catalan Research Council* under project 2014SGR1147. This research has also received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 734922.

1 Introduction

Originally, the Kautz digraphs were introduced by Kautz [9] in 1968. They have many applications, for example, they are useful as network topologies for connecting processors. The Kautz digraphs have the smallest diameter among all digraphs with their number of vertices and degree.

The cyclic Kautz digraphs $CK(d, \ell)$ were recently introduced by Böhmová, Huemer and the author [2, 3], as subdigraphs with special symmetries of the Kautz digraphs $K(d, \ell)$, see for example Fiol, Yebra and Alegre [7]. In contrast with these, the set of vertices of the cyclic Kautz digraphs is invariant under cyclic permutations of the sequences representing them. Thus, apart from their possible applications in interconnection networks, the cyclic Kautz digraphs $CK(d, \ell)$ could be relevant in coding theory, because they are related to cyclic codes. A linear code C of length ℓ is called cyclic if, for every codeword $c = (c_1, \dots, c_\ell)$, the codeword $(c_\ell, c_1, \dots, c_{\ell-1})$ is also in C . This cyclic permutation allows to identify codewords with polynomials. For more information about cyclic codes and coding theory, see Van Lint [10] (Chapter 6). With respect to other properties of the cyclic Kautz digraphs $CK(d, \ell)$, their number of vertices follows sequences that have several interpretations. For example, for $d = 2$ (that is, 3 different symbols) and $\ell = 2, 3, \dots$, the number of vertices follows the sequence 6, 6, 18, 30, 66, ... According to the On-Line Encyclopedia of Integer Sequences [12], this is the sequence A092297. For $d = 3$ (4 different symbols) and $\ell = 2, 3, \dots$, we get the sequence 12, 24, 84, 240, 732, ... corresponding to A226493 and A218034 in [12].

In this paper we give an alternative definition of $CK(d, \ell)$, by introducing the family of the subKautz digraphs $sK(d, \ell)$, from where the cyclic Kautz digraphs can be obtained as line digraphs. We present the exact formula of the distance between any two vertices of $sK(d, \ell)$ and $CK(d, \ell)$. This allows us to compute the diameter and the semigirth of both families, also providing an efficient routing algorithm to find the shortest path between any pair of vertices. Using these parameters, we also prove that $sK(d, \ell)$ and $CK(d, \ell)$ are maximally vertex-connected and super-edge-connected. Whereas $K(d, \ell)$ are optimal with respect to the diameter, we show that $sK(d, \ell)$ and $CK(d, \ell)$ are optimal with respect to the mean distance, whose exact values are given for both families when $\ell = 3$. Finally, we provide a lower bound on the girth of $sK(d, \ell)$ and $CK(d, \ell)$.

1.1 Notation

We consider simple digraphs (or directed graphs) without loops or multiple arcs, and we follow the usual notation for them. That is, a *digraph* $G = (V, E)$ consists of a (finite) set $V = V(G)$ of vertices and a set $E = E(G)$ of arcs (directed edges) between vertices of G . If $a = (u, v)$ is an arc between vertices u and v , then the vertex u is *adjacent to* the vertex v , and the vertex v is *adjacent from* u . Let $\Gamma^+(v)$ and $\Gamma^-(v)$ denote the set of vertices adjacent from and to the vertex v , respectively. Their cardinalities are the *out-degree*

$\delta^+(v) = |\Gamma^+(v)|$ of the vertex v , and the *in-degree* $\delta^-(v) = |\Gamma^-(v)|$ of the vertex v . A digraph G is called *d-out-regular* if $\delta^+(v) = d$, *d-in-regular* if $\delta^-(v) = d$, and *d-regular* if $\delta^+(v) = \delta^-(v) = d$, for all $v \in V$. The minimum degree $\delta = \delta(G)$ of G is the minimum over all the in-degrees and out-degrees of the vertices of G . A *digon* is a directed cycle on 2 vertices. For other notation, and unless otherwise stated, we follow the book by Bang-Jensen and Gutin [1].

In the *line digraph* $L(G)$ of a digraph G , each vertex represents an arc of G , $V(L(G)) = \{uv : (u, v) \in E(G)\}$, and a vertex uv is adjacent to a vertex wz when $v = w$, that is, when in G the arc (u, v) is adjacent to the arc (w, z) : $u \rightarrow v(=w) \rightarrow z$. Fiol and Lladó defined in [6] the *partial line digraph* $PL(G)$ of a digraph G , where some (but not necessarily all, as in the line digraph $L(G)$) of the arcs in G become vertices in $PL(G)$. Let $E' \subseteq E$ be a subset of arcs which are incident to all vertices of G , that is, $\{v : (u, v) \in E'\} = V$. A digraph $PL(G)$ is said to be a *partial line digraph* of G if its vertices represent the arcs of E' , that is, $V(PL(G)) = \{uv : (u, v) \in E'\}$, and a vertex uv is adjacent to the vertices $v'w$, for each $w \in \Gamma_G^+(v)$, where

$$v' = \begin{cases} v & \text{if } vw \in V(PL(G)), \\ \text{any other vertex of } \Gamma_G^-(w) \text{ such that } v'w \in V(PL(G)) & \text{otherwise.} \end{cases}$$

A digraph G is *strongly connected* when, for any pair of vertices $x, y \in V$, there always exists an $x \rightarrow y$ path, that is, a path from the vertex x to the vertex y . The *strong connectivity* $\kappa = \kappa(G)$ (or strong vertex-connectivity) of G is the smallest number of vertices whose deletion results in a digraph that is either not strongly connected or trivial. Analogously, the *strong arc-connectivity* $\lambda = \lambda(G)$ of G is the smallest number of arcs whose deletion results in a not strongly connected digraph. Since we only deal with strong connectivities, from now on we are going to refer to them simply as connectivities. Now we only consider connected digraphs, so $\delta \geq 1$. It is known that $\kappa \leq \lambda \leq \delta$, see Geller and Harary [8]. A digraph G is *maximally connected* when $\kappa = \lambda = \delta$.

If G is a *maximally arc-connected* digraph ($\lambda = \delta$), then any set of arcs adjacent from [to] a vertex x with out-degree [in-degree] δ is a minimum order arc-disconnecting set. Similarly, if G is a *maximally vertex-connected* digraph ($\kappa = \delta$), the set of vertices adjacent from [to] x is a minimum order vertex-disconnecting set. In this context, these arc or vertex sets are called *trivial*. Note that the deletion of any trivial set isolates a vertex of in-degree or out-degree δ . A digraph G is *super- κ* if every minimum vertex-disconnecting set is trivial. Analogously, G is *super- λ* if all its minimum arc-disconnecting sets are trivial. If G is super- κ , then $\kappa = \delta$, and if G is super- λ , then $\lambda = \delta$. In general, the converses are not true.

We say that a digraph is *weakly antipodal* when every vertex u has exactly one vertex v at maximum distance (the diameter), and it is *antipodal* when simultaneously u and v are at maximum distance from each other. For instance, the directed cycle C_n is weakly antipodal, whereas the symmetric directed cycle C_n^* with even n is antipodal.

1.2 The semigirth

We recall the definition of the *semigirth*: For a given digraph G , let $\gamma = \gamma(G)$, for $1 \leq \gamma \leq D$, where D is the diameter, be the greatest integer such that for any two (not necessarily different) vertices $x, y \in V$,

- (a) if $\text{dist}(x, y) < \gamma$, then the shortest $x \rightarrow y$ path is unique, and there is no an $x \rightarrow y$ path of length $\text{dist}(x, y) + 1$;
- (b) if $\text{dist}(x, y) = \gamma$, then there is only one shortest $x \rightarrow y$ path.

Note that γ is well defined when G has no loops. In [5], Fàbrega and Fiol proved that, if a digraph G (different from a directed cycle) has semigirth γ , then its line digraph $L(G)$ has semigirth $\gamma + 1$. The diameter also has the same behaviour, that is, if the diameter of G is D , then its line digraph $L(G)$ has diameter $D + 1$.

We also recall two results from Fàbrega and Fiol [5] on the connectivities and super-connectivities.

Theorem 1 ([5]). *Let $G = (V, E)$ be a loopless digraph with minimum degree $\delta > 1$, semigirth γ , diameter D and connectivities λ and κ .*

- (a) *If $D \leq 2\gamma$, then $\lambda = \delta$.*
- (b) *If $D \leq 2\gamma - 1$, then $\kappa = \delta$.*

Theorem 2 ([5]). *Let $G = (V, E)$ be a loopless digraph with minimum degree $\delta \geq 3$, semigirth γ , and diameter D .*

- (a) *If $D \leq 2\gamma$, then G is super- λ .*
- (b) *If $D \leq 2\gamma - 2$, then G is super- κ .*

1.3 Moore digraphs with respect to the diameter and the mean distance

The Moore bound on the number of vertices for digraphs with diameter D and maximum degree Δ is $N(\Delta, D) = \frac{\Delta^{D+1}-1}{\Delta-1}$ for $\Delta > 1$ and $N(1, D) = D + 1$. Notice that $N \sim O(\Delta^D)$.

The digraphs that attain the Moore bound $N(\Delta, D)$ are called Moore digraphs. The only Moore digraphs are the directed cycles on $D + 1$ vertices and the complete digraphs on $\Delta + 1$ vertices. For $D > 1$ and $\Delta > 1$, there are no Moore digraphs. For more information, see the survey by Miller and Širáň [11].

The mean distance corresponding to a digraph attaining the Moore bound is given in the following result. As the only Moore digraphs are the directed cycles and the complete digraphs, this bound gives an idea of how close is a digraph (with diameter D and maximum degree Δ) of being a Moore digraph.

Lemma 1. *The mean distance $\bar{\delta}(\Delta, D)$ of a digraph with diameter D and maximum degree Δ attaining the Moore bound would be*

$$\bar{\delta}(\Delta, D) = \frac{D\Delta^{D+2} - (1 + D)\Delta^{D+1} + \Delta}{\Delta^{D+2} - \Delta^{D+1} - \Delta + 1}.$$




Figure 1: Some examples of the Kautz and the subKautz digraphs.

Proof. We compute $\bar{\partial}(\Delta, D)$ taking into account that the maximum number of vertices at distance k is Δ^k .

$$\begin{aligned} \bar{\partial}(\Delta, D) &= \frac{1}{N(\Delta, D)} \sum_{k=0}^D k \Delta^k = \frac{\Delta}{N(\Delta, D)} \sum_{k=0}^D k \Delta^{k-1} = \frac{\Delta}{N(\Delta, D)} \left(\sum_{k=0}^D \Delta^k \right)' \\ &= \frac{\Delta}{N(\Delta, D)} \left(\frac{\Delta^{D+1} - 1}{\Delta - 1} \right)' = \frac{D\Delta^{D+2} - (1+D)\Delta^{D+1} + \Delta}{\Delta^{D+2} - \Delta^{D+1} - \Delta + 1}. \end{aligned}$$

□

We can define a digraph as optimal with respect to the diameter (the maximum delay in a message transmission), but also with respect to the mean distance (the average delay in a message transmission). So, we can say that a digraph is optimal when, if N is of the order of Δ^k , then its mean distance is of the order of k , that is, when $\bar{\partial} \sim O(\log_{\Delta} N)$.

2 Kautz-like digraphs

The Kautz $K(d, \ell)$, the subKautz $sK(d, \ell)$, the cyclic Kautz $CK(d, \ell)$, and the modified cyclic Kautz $MCK(d, \ell)$ digraphs have vertices represented by words on an alphabet, and adjacencies between vertices correspond to shifts of the words. In these Kautz-like digraphs a path $\mathbf{x} \rightarrow \mathbf{y}$ corresponds to a sequence beginning with $\mathbf{x} = x_1 x_2 \dots x_{\ell}$ and finishing with $\mathbf{y} = y_1 y_2 \dots y_{\ell}$, where every subsequence of length ℓ corresponds to a vertex of the corresponding digraph.




Figure 2: An example of a cyclic Kautz digraph and a modified cyclic Kautz digraph.

2.1 Kautz and subKautz digraphs

Next, we recall the definitions of the Kautz $K(d, \ell)$, and we define a new family of Kautz-like digraphs called the subKautz digraphs $sK(d, \ell)$. See examples of both in Figure 1.

A *Kautz digraph* $K(d, \ell)$ has the vertices $x_1x_2 \dots x_\ell$, where $x_i \in \mathbb{Z}_{d+1}$, with $x_i \neq x_{i+1}$ for $i = 1, \dots, \ell - 1$, and adjacencies

$$x_1x_2 \dots x_\ell \rightarrow x_2x_3 \dots x_\ell y, \quad y \neq x_\ell.$$

Given integers d and ℓ , with $d, \ell \geq 2$, a *subKautz digraph* $sK(d, \ell)$ has set of vertices $V = \{x_1x_2 \dots x_\ell : x_i \neq x_{i+1}, i = 1, \dots, \ell - 1\} \subset \mathbb{Z}_{d+1}^\ell$, and adjacencies

$$x_1x_2 \dots x_\ell \rightarrow x_2 \dots x_\ell x_{\ell+1}, \quad x_{\ell+1} \neq x_1, x_\ell. \quad (1)$$

Hence, the subKautz digraph $sK(d, \ell)$ has $d^\ell + d^{\ell-1}$ vertices, as the Kautz digraph $K(d, \ell)$. Besides, the out-degree of a vertex $x_1x_2 \dots x_\ell$ is d if $x_1 = x_\ell$, and $d - 1$ otherwise. In particular, the subKautz digraph $sK(d, 2)$ is $(d - 1)$ -regular and can be obtained from the Kautz digraph $K(d, 2)$ by removing all its arcs forming a digon.

Note that the subKautz digraph $sK(d, \ell)$ is a subdigraph of the Kautz digraph $K(d, \ell)$.

2.2 Cyclic Kautz and modified cyclic Kautz digraphs

Next, we recall the definitions of the cyclic Kautz digraphs $CK(d, \ell)$ and the modified cyclic Kautz digraphs $MCK(d, \ell)$. See an example of both in Figure 2.

A *cyclic Kautz digraph* $CK(d, \ell)$ has the vertices $x_1x_2 \dots x_\ell$, where $x_i \in \mathbb{Z}_{d+1}$, with $x_i \neq x_{i+1}$ for $i = 1, \dots, \ell - 1$, and $x_\ell \neq x_1$, and adjacencies

$$x_1x_2 \dots x_\ell \rightarrow x_2x_3 \dots x_\ell y, \quad y \neq x_2, x_\ell.$$

Note that the cyclic Kautz digraphs $CK(d, \ell)$ are subdigraphs of the Kautz digraph $K(d, \ell)$. It was proved in [3] that when $d = 2$ the cyclic Kautz digraphs $CK(2, \ell)$ are not connected (except for the case $\ell = 4$), and when $\ell = 2$ the cyclic Kautz digraphs $CK(d, 2)$ coincide with the Kautz digraphs $K(d, 2)$.

Recall that the diameter of the Kautz digraphs is optimal, that is, for a fixed out-degree d and number of vertices $(d+1)d^{\ell-1}$, the Kautz digraph $K(d, \ell)$ has the smallest diameter ($D = \ell$) among all digraphs with $(d+1)d^{\ell-1}$ vertices and degree d (see, for example, Miller and Širáň [11]). Since the diameter of the cyclic Kautz digraphs $CK(d, \ell)$ is greater than the diameter of the Kautz digraphs $K(d, \ell)$, in [4] we constructed the *modified cyclic Kautz digraphs* $MCK(d, \ell)$ by adding some arcs to $CK(d, \ell)$, in order to obtain the same diameter as $K(d, \ell)$, without increasing the maximum degree. In a cyclic Kautz digraph $CK(d, \ell)$, a vertex labeled with $a_2 \dots a_{\ell+1}$ is forbidden if $a_2 = a_{\ell+1}$. For each label, we replace the first symbol a_2 by one of the possible symbols a'_2 such that now $a'_2 \neq a_3, a_{\ell+1}$ (so $a'_2 \dots a_{\ell+1}$ represents a vertex). Then, we add arcs from the vertex $a_1 \dots a_\ell$ to the vertex $a'_2 \dots a_{\ell+1}$, with $a_1 \neq a_\ell$ and $a'_2 \neq a_3, a_{\ell+1}$. Note that $CK(d, \ell)$ and $MCK(d, \ell)$ have the same vertices, because we only add arcs to $CK(d, \ell)$ to obtain $MCK(d, \ell)$.

Lemma 2. (a) *The cyclic Kautz digraph $CK(d, \ell)$ is the line digraph of the subKautz digraph $sK(d, \ell - 1)$, that is, $CK(d, \ell) = L(sK(d, \ell - 1))$.*

(b) *The modified cyclic Kautz digraph $MCK(d, \ell)$ is the partial line digraph of the Kautz digraph $K(d, \ell - 1)$, that is, $MCK(d, \ell) = PL(K(d, \ell - 1))$.*

Proof. (a) From (1) we can write the arcs $(x_1x_2 \dots x_{\ell-1}, x_2 \dots x_{\ell-1}x_\ell)$ of $sK(d, \ell - 1)$ as $x_1x_2 \dots x_{\ell-1}x_\ell$ with $x_i \neq x_{i+1}$ and $x_1 \neq x_\ell$, which corresponds to the vertices of $CK(d, \ell)$. Moreover, two arcs are adjacent in $sK(d, \ell - 1)$ if

$$x_1x_2 \dots x_\ell \rightarrow x_2 \dots x_\ell x_{\ell+1},$$

where $x_1 \neq x_\ell$, as required for the vertices of $CK(d, \ell)$.

(b) This was proved in [4]. In taking the partial line digraph, it suffices to consider only the arcs in $K(d, \ell - 1)$ that are also in $sK(d, \ell - 1)$. \square

By using spectral techniques, the order $n_{d, \ell}$ of a cyclic Kautz digraph $CK(d, \ell)$ was given in [2, 3]. Here we use a combinatorial proof of this result.

Proposition 1. *The order $n_{d, \ell}$ of a cyclic Kautz digraph $CK(d, \ell)$ (that coincide with the size of the subKautz digraph $sK(d, \ell - 1)$) is $n_{d, 1} = d + 1$ and*

$$n_{d, \ell} = d^\ell + (-1)^\ell d \quad \text{for } \ell \geq 2. \quad (2)$$

Proof. The number $N_{d,\ell}$ of sequences $x_1x_2\dots x_\ell$ with $x_i \neq x_{i+1}$ for $i = 1, \dots, \ell-1$ (vertices of $K(d, \ell)$) is $d^\ell + d^{\ell-1}$. Then, to compute $n_{d,\ell}$, we must subtract from $N_{d,\ell}$ the number $n'_{d,\ell}$ of sequences $x_1x_2\dots x_\ell$ such that $x_1 = x_\ell$. But this is the same as the number of sequences $x_2\dots x_\ell$ with $x_2 \neq x_\ell$ and $x_i \neq x_{i+1}$ for $i = 2, \dots, \ell-1$, which is $n_{d,\ell-1}$. Consequently, we get the recurrence

$$n_{d,\ell} = d^\ell + d^{\ell-1} - n_{d,\ell-1} \quad \text{for } \ell \geq 3. \quad (3)$$

Thus, (2) follows by applying recursively (3) and using that $n_{d,2} = d^2 + d$. \square

In the following result we prove a way of finding an $sK(d, \ell)$ a from the Kautz digraphs $K(d, \ell)$. We use the cyclic Kautz digraphs $CK(d, \ell)$ in the proof.

Lemma 3. *The subKautz digraphs $sK(d, \ell)$ can be obtained from the Kautz digraphs $K(d, \ell)$ by removing all the arcs of the closed walks of length ℓ in the complete symmetric digraph K_{d+1}^* .*

Proof. From their definition, the subKautz digraphs $sK(d, \ell)$ are obtained from $K(d, \ell)$ by removing the arcs of the form $x_1x_2\dots x_\ell \rightarrow x_2\dots x_\ell x_1$, which correspond to the vertices $x_1x_2\dots x_\ell x_1$ of $K(d, \ell+1)$, which in turn correspond to the closed walks of length ℓ in the complete symmetric digraph K_{d+1}^* . \square

A simple property of symmetry shared by all the Kautz-like digraphs is the following. The converse digraph is obtained by changing the direction of all the arcs in the original digraph.

Lemma 4. *The Kautz digraphs $K(d, \ell)$, the subKautz digraphs $sK(d, \ell)$, and the cyclic Kautz digraphs $CK(d, \ell)$ are isomorphic to their converses.*

Proof. Since the mapping $\Psi(x_1x_2\dots x_\ell) = x_\ell\dots x_2x_1$ satisfies

$$\begin{aligned} \Psi(\Gamma^+(\{x_1x_2\dots x_\ell\})) &= \Psi(\{x_2x_3\dots x_\ell y : y \in \mathbb{Z}_{d+1}, y \neq x_\ell\}) \\ &= \{yx_\ell\dots x_3x_2 : y \in \mathbb{Z}_{d+1}, y \neq x_\ell\} \\ &= \Gamma^-(\{x_\ell\dots x_2x_1\}) = \Gamma^-(\Psi(\{x_1x_2\dots x_\ell\})), \end{aligned}$$

where in the case of $CK(d, \ell)$ also $y \neq x_2$, it is an isomorphism between every of such digraphs and its converse. \square

3 Routing, distances and girth in $CK(d, \ell)$

In this section, we only need to consider the cases with $d \geq 3$ and $\ell \geq 3$ because, as said in the Introduction, when $d = 2$ the cyclic Kautz digraphs $CK(2, \ell)$ are not connected (except for the case $\ell = 4$), and when $\ell = 2$, the cyclic Kautz digraphs $CK(d, 2)$ coincide with the Kautz digraphs $K(d, 2)$.




Figure 3: (a) The subKautz digraph $sK(3,3)$ whose line digraph is $CK(3,4)$ (the lines without direction represent two arcs with opposite directions). (b) The cyclic Kautz digraph $CK(3,3)$ with 24 vertices and diameter 5 (the vertices at maximum distance from 012 are 210 and 213).

3.2 Routing and distances when $d = 3$ or $\ell = 3$

Looking at the case (c3) above, if $d = 3$ and all the elements z_2, x_4, y_6, y_1 are different, then z_3 has no possible value. Analogously, if $\ell = 3$, there must exist two vertices $\mathbf{x} = x_1x_2x_3$ and $\mathbf{y} = y_1y_2y_3$, such that $|\tilde{\mathbf{x}} \sqcap \mathbf{y}| = 2$ (not smaller than $\ell - 1$), and with $y_1 = x_3$. Thus, neither of the strategies in the above cases (c) and (b) can be applied. However, the following reasoning shows that we always can find a path of length $2\ell - 1$. First, we deal with the case $d = 3$, where for simplicity we assume that $\ell = 5$.

(d) We reason as if $|\tilde{\mathbf{x}} \sqcap \mathbf{y}| = 0$:

$$\begin{aligned}
 & x_1 \quad x_2 \quad x_3 \quad x_4 \quad x_5 \quad \overline{x_2} \quad \overline{x_3} \quad \overline{x_4} \quad \overline{x_5} \\
 & \overline{y_1} \quad \overline{y_2} \quad \overline{y_3} \quad \overline{y_4} \quad y_1 \quad y_2 \quad y_3 \quad y_4 \quad y_5 \\
 & = z_1 \quad z_2 \quad z_3 \quad z_4 \quad y_1 \quad y_2 \quad y_3 \quad y_4 \quad y_5
 \end{aligned}$$

where we would need the following conditions:

- (d1) $z_1 \neq x_2, x_5, y_1, 1$,
- (d2) $z_2 \neq z_1, x_3, y_2$,
- (d3) $z_3 \neq z_2, x_4, y_3$,
- (d4) $z_4 \neq z_3, x_5, y_4, y - 1$.

If $d \geq 4$ (for $\ell = 3$), this conditions can always be fulfilled, and the required path is guaranteed.

If $d = 3$, and either $y_1 = x_2$, or $y_2 = x_3$, or $y_3 = x_4$, or $y_4 = x_5$, or $y_1 = x_5$, then there is always a possible choice of z_1, z_2, z_3 and z_4 in \mathbb{Z}_4 . Consequently, $\text{dist}(\mathbf{x}, \mathbf{y}) \leq 9$. Otherwise, if $y_i \neq x_{i+1}$ for $i = 1, \dots, 4$ and $y_1 \neq x_5$, we can reason as if $|\tilde{\mathbf{x}} \cap \mathbf{y}| = 4 (= \ell - 1)$. In this case, the path from \mathbf{x} to \mathbf{y} is:

$$\mathbf{x} = x_1x_2x_3x_4x_5 \rightarrow x_2x_3x_4x_5y_1 \rightarrow x_3x_4x_5y_1y_2 \rightarrow \cdots \rightarrow y_1y_2y_3y_4y_5 = \mathbf{y},$$

which implies that $\text{dist}(\mathbf{x}, \mathbf{y}) \leq 5$.

Thus, in any case,

$$\text{dist}(\mathbf{x}, \mathbf{y}) \leq 2\ell - 1.$$

This leads to the following result.

Proposition 2. (i) *The diameter of the cyclic Kautz digraphs $CK(3, \ell)$ with $\ell \neq 4$ and that of $CK(d, \ell)$ with $\ell = 3$ is $2\ell - 1$.*

(ii) *The diameter of the cyclic Kautz digraph $CK(3, 4)$ is $2\ell - 2 = 6$.*

Proof. (i) We only need to exhibit two vertices at distance $2\ell - 1$. For $CK(3, \ell)$ with $\ell \geq 5$, when ℓ is odd, we can take the vertices $\mathbf{x} = 0101\dots012$ and $\mathbf{y} = 21010\dots10$. When ℓ is even, two vertices at maximum distance are $\mathbf{x} = 102020\dots2012$ and $\mathbf{y} = 2130202\dots02010$. In both cases, it was proved that these vertices are at maximum distance in [3]. The case of the cyclic Kautz digraph $CK(3, 3)$, shown in Figure 3 (b), can be easily checked to have diameter $2\ell - 1 = 5$, for instance, the vertices at maximum distance from 012 are 210 and 213. In general, for $CK(d, 3)$, we show that two vertices at maximum distance 5 are $\mathbf{x} = x_1x_2x_3$ and $\mathbf{y} = x_3x_2y_3$ as follows. If this distance were 2, then we would get the sequence $x_1x_2x_3x_2y_3$, but $x_2x_3x_2$ is not a vertex of $CK(d, 3)$. If this distance were 3, then we would get the sequence $x_1x_2x_3x_3x_2y_3$, but $x_2x_3x_3$ is not a vertex of $CK(d, 3)$. If this distance were 4, then we would get the sequence $x_1x_2x_3y_1x_3x_2y_3$, but $x_3y_1x_3$ is not a vertex of $CK(d, 3)$. Then, the distance is 5, with the sequence $x_1x_2x_3y_1y_2x_3x_2y_3$.

(ii) The cyclic Kautz digraph $CK(3, 4)$ on 84 vertices with labels $x_1x_2x_3x_4$, $x_i \in \mathbb{Z}_4$, is the line digraph of the subKautz digraph $sK(3, 3)$ shown in Figure 3 (a). Then, since $sK(3, 3)$ has diameter 5, we conclude that $CK(3, 4)$ has diameter 6, as claimed. \square

Corollary 2. (i) *The diameter of the subKautz digraphs $sK(d, \ell)$ with either $d = 3$ and $\ell \geq 4$ or $d \geq 3$ and $\ell = 2$ is 2ℓ .*

(ii) *The diameter of the subKautz digraph $sK(3, 3)$ is $2\ell - 1 = 5$.*

See Figure 4 for a summary of the diameters of $sK(d, \ell)$ and $CK(d, \ell)$.

3.3 The girth

Now we give a lower bound on the girth of a cyclic Kautz digraph $CK(d, \ell)$.

$d \backslash \ell$	2	3	≥ 4
3	2ℓ	$2\ell-1$	2ℓ
≥ 4			

$d \backslash \ell$	3	4	≥ 5
3	$2\ell-1$	$2\ell-2$	$2\ell-1$
≥ 4			

Figure 4: Summary of the diameters of $sK(d, \ell)$ and $CK(d, \ell)$, depending on the values of d and ℓ .

Lemma 5. *The girth g of the cyclic Kautz digraph $CK(d, \ell)$ is at least the minimum positive integer k such that ℓ is not congruent with $1 \pmod{k}$.*

Proof. A cycle of minimum length g , rooted to a vertex \mathbf{x} , corresponds to a path from \mathbf{x} to \mathbf{x} of the same length. This means that the maximum length of the (nontrivial) intersection $\mathbf{x} \sqcap \mathbf{x}$ is $\ell - g$. For instance, with $\ell = 7$ and $g = 4$ we would have the intersection pattern

$$\begin{array}{cccccccccccc} x_1 & x_2 & x_3 & x_4 & x_5 & x_6 & x_7 & \overline{x_2} & \overline{x_3} & \overline{x_4} & \overline{x_5} & \overline{x_6} & \overline{x_7} \\ & & & & & & & x_1 & x_2 & x_3 & x_4 & x_5 & x_6 & x_7. \end{array}$$

Then, in general, this means that the sequence representing \mathbf{x} is periodic: $x_i = x_{i+g}$ for every $i = 1, 2, \dots, \ell - g$. Now, if $\ell \equiv r \pmod{g}$, then $x_\ell = x_r$, which is possible if $r \neq 1$, and in this case the cycle would be

$$\begin{aligned} \mathbf{x} &= x_1x_2 \dots x_g \dots x_1x_2 \dots x_g x_1x_2 \dots x_r \\ &\rightarrow x_2 \dots x_g \dots x_1x_2 \dots x_g x_1x_2 \dots x_r x_{r+1} \\ &\rightarrow \dots \rightarrow x_{g-r+1} \dots x_g \dots x_1x_2 \dots x_g x_1x_2 \dots x_r x_{r+1} \dots x_g \\ &\rightarrow x_{g-r+2} \dots x_g \dots x_1x_2 \dots x_g x_1x_2 \dots x_r x_{r+1} \dots x_g x_1 \\ &\rightarrow \dots \rightarrow x_1x_2 \dots x_g \dots x_1x_2 \dots x_r x_{r+1} \dots x_g x_1 \dots x_r = \mathbf{x}. \end{aligned}$$

This completes the proof. □

Note that the girth reaches the bound when there exists a vertex \mathbf{x} that satisfies the cases (a), (b), (c) or (d) (given at the beginning of this section) for the existence of a path of length g from \mathbf{x} to $\mathbf{y} = \mathbf{x}$. In particular, this is fulfilled if d is large enough. As an example, if $\ell = 13$ Lemma 5 gives $g \geq 5$. However, a possible vertex \mathbf{x} only exists for $d \geq 4$. Indeed, assume that $\mathbf{x} = x_1x_2x_3x_4x_5x_1x_2x_3x_4x_5x_1x_2x_3$, where $x_i \in \mathbb{Z}_4$ for $i = 1, \dots, 5$. Since $x_2 \neq x_1$ and $x_3 \neq x_2, x_1$, we can take, without loss of generality $\mathbf{x} = 012x_4x_5012x_4x_5012$. Then, a path of length $g = 5$ from \mathbf{x} to \mathbf{x} should be

$$\begin{aligned} \mathbf{x} &= 012x_4x_5012x_4x_5012 \rightarrow 12x_4x_5012x_4x_5012x_4 \rightarrow 2x_4x_5012x_4x_5012x_4x_5 \\ &x_4x_5012x_4x_5012x_4x_50 \rightarrow x_5012x_4x_5012x_4x_501 \rightarrow 012x_4x_5012x_4x_5012 = \mathbf{x}. \end{aligned}$$

$d \backslash \ell$	2	3	≥ 4
3	super- λ	super- λ	super- λ
≥ 4		max v-c	

$d \backslash \ell$	3	4	≥ 5
3	super- λ max v-c	super- λ	super- λ max v-c
≥ 4		super- κ	

Figure 5: Summary of the connectivities of $sK(d, \ell)$ and $CK(d, \ell)$, depending on the values of d and ℓ .

Therefore, since $x_4 \neq 2, 1$ and $0 \neq x_4$, then $x_4 = 3$. Moreover, since $x_5 \neq x_4, 2$, $0 \neq x_5$, and $1 \neq x_5$, then $x_5 \notin \{0, 1, 2, 3\}$, which is a contradiction. In fact, when $d = 3$, it turns out that $CK(3, 13)$ has girth $g = 7$, for example, with the vertex $x = 0120123012012$.

A direct consequence of this result is that there exist cyclic Kautz digraphs with arbitrarily large girth. Indeed, if $\ell = \text{lcm}(2, 3, \dots, n) + 1$, we have that $\ell = 1 \pmod{i}$ for every $i = 2, 3, \dots, n$. Then, according to Lemma 5, $CK(d, \ell)$ must have girth $g > n$.

It is known that if a digraph G has girth g , then its line digraph $L(G)$ also has girth g , see Fàbrega and Fiol [5]. Since $L(sK(d, \ell)) = CK(d, \ell + 1)$, both digraphs have the same girth.

4 Connectivity and superconnectivity

It is well-known that the Kautz digraphs $K(d, \ell)$ have maximal (edge- and vertex-) connectivities (see Fàbrega and Fiol [5]). The following result shows that this is also the case for the other Kautz-like digraph studied here, see Figure 5 for a summary.

Proposition 3. (i) *The subKautz digraph $sK(d, \ell)$ with $d \geq 3$ and $\ell \geq 2$ is super- λ .*

(ii) *The subKautz digraph $sK(d, \ell)$ with either $d = \ell = 3$, or $d \geq 4$ and $\ell \geq 3$, is maximally vertex-connected.*

(iii) *The cyclic Kautz digraph $CK(d, \ell)$ with $d \geq 3$ and $\ell \geq 3$ is super- λ .*

(iv) *The cyclic Kautz digraph $CK(d, \ell)$ with either $d = 3$ and $\ell = 4$, or $d, \ell \geq 4$, is super- κ .*

(v) *The cyclic Kautz digraph $CK(d, \ell)$ with either $d = 3$ and $\ell \neq 4$, or $d \geq 4$ and $\ell = 3$, is maximally vertex-connected.*

Proof. Since both $sK(d, \ell)$ and $CK(d, \ell)$ are subdigraphs of $K(d, \ell)$, with semigirth ℓ (see Fàbrega and Fiol [5]), then the semigirths of these digraphs are at least ℓ . Hence, by using that the diameters of $sK(d, \ell)$ and $CK(d, \ell)$ are given in Theorem 3, Proposition 2, and Corollaries 1 and 2, the result follows from Theorems 1 and 2. \square

5 Cyclic Kautz digraphs $CK(d, 3)$ with $d \geq 3$

The cyclic Kautz digraphs $CK(d, 3)$ with $d \geq 3$ have some special properties that, in general, are not shared with $CK(d, \ell)$ with $\ell > 3$. These properties are listed in the following result.

Lemma 6. *The cyclic Kautz digraphs $CK(d, 3)$ with $d \geq 3$ satisfy the following properties:*

- (a) $(d - 1)$ -regular.
- (b) Number of vertices: $N = d^3 - d$, number of arcs: $m = (d + 1)d(d - 1)^2$.
- (c) Diameter: $2\ell - 1 = 5$.
- (d) $CK(d, 3)$ are the line digraphs of the subKautz digraphs $sK(d, 2)$, which are obtained from the Kautz digraphs $K(d, 2)$ by removing the arcs of the digons.
- (e) Vertex-transitive.
- (f) Eulerian and Hamiltonian.

Proof. (a), (b), (c) and (d) come from the properties of general $CK(d, \ell)$. (e) Since $sK(d, 2)$ (with $d \geq 3$) are vertex-transitive and arc-transitive, their line digraphs $CK(d, 3)$ are vertex-transitive. (f) $sK(d, 2)$ and $CK(d, 3)$ with $d \geq 3$ are Eulerian, because they are $(d - 1)$ -regular. Since $sK(d, 2)$ (with $d \geq 3$) are Eulerian, their line digraphs $CK(d, 3)$ are Hamiltonian. \square

5.1 Mean distance

As said before, $CK(d, \ell)$ are asymptotically optimal with respect to the mean distance. Now, we give the exact formulas for the mean distance of $sK(d, 2)$ and $CK(d, 3)$ with $d \geq 3$. Let n and N be the numbers of vertices of $sK(d, 2)$ and $CK(d, 3)$, respectively.

Lemma 7. (a) *The mean distance of the antipodal subKautz digraph $sK(d, 2)$ with $d \geq 3$ is*

$$\bar{\partial}^* = \frac{2d^2 + 3d - 1}{d^2 + d}. \quad (4)$$

(b) *The mean distance of the cyclic Kautz digraph $CK(d, 3)$ with $d \geq 3$ is*

$$\bar{\partial} = \frac{3d^3 + d^2 - 5d - 2}{d^3 - d}. \quad (5)$$

Proof. Since $CK(d, 3)$ (and also $sK(2, 2)$) with $d \geq 3$ is vertex-transitive, we can compute the number of vertices from any given vertex. First, we fix the distance layers in $sK(2, 2)$. Thus, in Table 1, we give the numbers $n_k(u, v)$ of vertices at distance $k = 0, 1, \dots, 4$ from vertex $u = 01$ to vertex $v \in \{01, 1x, \dots, 10\}$.

Then, the total numbers $n_i = n_i(u)$ of vertices at distance $i = 0, 1, \dots, 4$ from u turn out to be

$$n_0 = 1, \quad n_1 = d - 1, \quad n_2 = (d - 1)^2, \quad n_3 = 2(d - 1), \quad n_4 = 1,$$

u	v	$k = \text{dist}(u, v)$	$n_k(u, v)$
01	01	0	1
01	$1x$	1	$d - 1$
01	$x0$	2	$d - 1$
01	xy	2	$(d - 1)(d - 2)$
01	$x1$	3	$d - 1$
01	$0x$	3	$d - 1$
01	10	4	1

Table 1: Numbers of vertices v at distance k from $u = 01$.

with $n = n_0 + n_1 + \dots + n_4 = d^2 + d$, and showing that $sK(2, 2)$ is antipodal.

Now we use again that $CK(d, 3)$ is the line digraph of $sK(d, 2)$ to conclude that, in the former, the numbers N_i of vertices at distance $i = 0, 1, \dots, 5$ from a given vertex, say 201, are

$$N_0 = n_0 = 1, \quad N_1 = n_1 = d - 1, \quad N_2 = (d - 1)n_1 = (d - 1)^2, \quad N_3 = (d - 1)n_2 - 1 \\ = (d - 1)^3 - 1, \quad N_4 = (d - 1)n_3 = 2(d - 1)^2, \quad N_5 = (d - 1)n_4 = d - 1,$$

satisfying $N = N_0 + N_1 + \dots + N_5 = d^3 - d$, as requested.

Note that in $N_3 = (d - 1)n_2 - 1$ we subtract one unit due to the presence in $sK(d, 2)$ of the cycle of length 3: $20 \rightarrow 01 \rightarrow 12 \rightarrow 20$. Then, the mean distances of $CK(d, 3)$ with $d \geq 3$ are, respectively, $\bar{\delta}^* = \frac{1}{n} \sum_{k=0}^4 kn_k$, and $\bar{\delta} = \frac{1}{N} \sum_{k=0}^5 kN_k$, which gives the results. \square

Observe that, since $CK(d, 3)$ is the line digraph of $sK(d, 2)$, the respective mean distance satisfies the inequality $\bar{\delta} < \bar{\delta}^*$, in concordance with the results by Fiol, Yebra, and Alegre [7]. Also, note that the mean distances of $sK(d, 2)$ and $CK(d, 3)$, with $d \geq 3$, tend, respectively, to 2 and 3 for large degree $d - 1$, that is, they are asymptotically optimal.

References

- [1] J. Bang-Jensen, G. Gutin, *Digraphs: Theory, Algorithms and Applications*, Springer-Verlag, London, 2007.
- [2] K. Böhmová, C. Dalfó, C. Huemer, The diameter of cyclic Kautz digraphs, *Electron. Notes Discrete Math.* **49** (2015) 323–330.

- [3] K. Böhmová, C. Dalfó, C. Huemer, On the diameter of cyclic Kautz digraphs, *Filomat* (2017), to appear.
- [4] K. Böhmová, C. Dalfó, C. Huemer, New cyclic Kautz digraphs with optimal diameter, submitted (2016).
- [5] J. Fàbrega, M. A. Fiol, Maximally connected digraphs, *J. Graph Theory* **13** (1989) 657–668.
- [6] M. A. Fiol, A. S. Lladó, The partial line digraph technique in the design of large interconnection networks, *IEEE Trans. Comput.* **41** (1992) 848–857.
- [7] M. A. Fiol, J. L. A. Yebra, I. Alegre, Line digraph iterations and the (d, k) digraph problem, *IEEE Trans. Comput.* **C-33** (1984) 400–403.
- [8] D. Geller, F. Harary, Connectivity in digraphs, *Lect. Notes Math.* **186** (1970) 105–114.
- [9] W. H. Kautz, Bounds on directed (d, k) graphs, in *Theory of Cellular Logic Networks and Machines*, AFCRL-68-0668 Final Rep., 1968, 20–28.
- [10] J. H. van Lint, *An Introduction to Coding Theory*, 3rd edition, Springer-Verlag, New York, 1999.
- [11] M. Miller, J. Širáň, Moore graphs and beyond: A survey of the degree/diameter problem, *Electron. J. Combin.* **20(2)**, #DS14v2, 2013.
- [12] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, <http://oeis.org>.