

Some computations
with the
 \mathcal{F} -homotopy limit spectral
sequence

DISSERTATION ZUR ERLANGUNG DES DOKTORGRADES
DER NATURWISSENSCHAFTEN (DR. RER. NAT.)
DER FAKULTÄT FÜR MATHEMATIK
DER UNIVERSITÄT REGENSBURG

vorgelegt von
Koenraad van Woerden
aus Groningen, die Niederlande
im Jahr 2017

Promotionsgesuch eingereicht am: 27. April 2017

Die Arbeit wurde angeleitet von: Dr. Justin Noel

Prüfungsausschuss:

Vorsitzender:

Prof. Dr. Harald Garcke

1. Gutachter:

Dr. Justin Noel

2. Gutachter:

Prof. Dr. Niko Naumann

weiterer Prüfer:

Prof. Dr. Ulrich Bunke

Contents

1	Introduction	1
1.1	Nilpotence in Borel equivariant cohomology	1
1.2	Acknowledgements	4
1.3	Notations and conventions	4
2	The \mathcal{F}-homotopy limit spectral sequence	5
2.1	Introduction	5
2.2	Families of subgroups	5
2.3	Orbit categories	6
2.4	The \mathcal{F} -homotopy limit spectral sequence	6
2.5	\mathcal{F} -nilpotence	7
2.6	Comparison with the Lyndon-Hochschild-Serre spectral sequence	10
2.7	A decomposition of the E_2 -page.	11
	2.7.1 Homotopy cofinality	11
	2.7.2 Homotopy cofinality in orbit categories.	12
	2.7.3 A short exact sequence of coefficient systems	12
3	Exponent lemmas	16
3.1	Introduction	16
3.2	Lemmas for \mathcal{F} -exponents	16
3.3	From representations to exponents via the projective bundle theorem	20
	3.3.1 The projective bundle theorem	21
	3.3.2 Free R -modules	22
	3.3.3 From representations to exponents	24
3.4	A bound on the exponent for the family of proper subgroups	25
4	Mod-2 group cohomology computations of 2-groups using the \mathcal{F}-homotopy limit spectral sequence	27
4.1	Introduction	27
4.2	The 2-groups of order at most 16	28
	4.2.1 Abelian 2-groups	28
	4.2.2 The dihedral groups	28
	4.2.3 The quaternion groups	29
	4.2.4 The groups of order 8	29
	4.2.5 The groups of order 16	29
4.3	Abelian 2-groups	30
	4.3.1 Cyclic 2-groups	31

4.3.2	Abelian 2-groups	31
4.4	Dihedral groups	33
4.4.1	Introduction	33
4.4.2	The exponent	33
	The upper bound	34
4.4.3	The spectral sequence	35
	Summary of the computation	35
	The orbit category	36
	The LHSSS of $H_i \rightarrow D_8 \rightarrow C_2$	36
	The LHSSS of $Z \rightarrow D_8 \rightarrow C_2 \times C_2$	38
	The map j	39
	The E_2 -page	40
	Multiplicative extensions	40
	The lower bound illustrated	41
4.4.4	The dihedral groups D_{2^n} with $n \geq 4$	43
	The decomposition	43
	The LHSSS of $D_{2^{n-1}} \rightarrow D_{2^n} \rightarrow C_2$	43
	The LHSSS of $C_{2^{n-2}} \rightarrow D_{2^n} \rightarrow C_2 \times C_2$	44
	The map j	44
	The kernel of j	45
	The image of j	45
	The E_2 -page	45
	Multiplicative structure	46
4.5	Quaternion groups	46
4.5.1	An upper bound on the exponent	46
4.5.2	The $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence	46
4.5.3	The $\mathcal{E}_{(2)}$ -exponent	47
4.6	SD_{16}	47
4.6.1	The E_2 -page	48
4.6.2	Differentials and $E_4 = E_\infty$	48
4.6.3	Poincaré Series	49
4.6.4	An upper bound on the $\mathcal{E}_{(2)}$ -exponent	50
4.6.5	The exponent	51
4.7	M_{16}	52
4.7.1	Introduction	52
4.7.2	An upper bound on the exponent	52
4.7.3	The orbit category	53
4.7.4	The E_2 -page	53
4.7.5	Differentials on the E_2 -page	56
	Generating differentials	56
	Differentials in the even rows	56
	Differentials in the odd rows	57
4.7.6	The E_3 -page	59
4.7.7	Poincaré Series	60
4.7.8	The exponent	61
4.8	$D_8 * C_4$	61
4.8.1	Introduction	61
4.8.2	Summary of the computation	61
4.8.3	The orbit category	62
4.8.4	An upper bound on the \mathcal{A} -exponent	64

4.8.5	The decomposition	65
4.8.6	Automorphisms of $D_8 * C_4$	65
4.8.7	Low dimensional Betti numbers	66
4.8.8	The LHSSS for $C \rightarrow D_8 * C_4 \rightarrow C_2 \times C_2$	66
	Differentials on E_2 and E_3	67
	The $E_4 = E_\infty$ -page	68
4.8.9	The LHSSS for $A_i \rightarrow D_8 * C_4 \rightarrow C_2$	69
	The local coefficient system	69
	The E_2 -page	70
4.8.10	Summary of LHSSS computations	72
4.8.11	The map j^*	73
	The B -summand	75
	The A -summand	77
	The E_2 -page	77
4.8.12	Differentials	78
4.8.13	Poincaré series	82
4.8.14	Differentials originating from the $s = 0$ -line	82
4.8.15	Exponents for various families	85
4.9	$C_4 \rtimes C_4$	87
4.9.1	An upper bound on the exponent	87
4.9.2	The spectral sequence	89
4.9.3	Poincaré series	92
4.9.4	The exponent	92
4.10	$(C_4 \times C_2) \overset{\psi_5}{\rtimes} C_2$	93
4.10.1	Differentials	94
4.10.2	Poincaré series	95
4.10.3	The exponent	95
4.11	$Q_8 \times C_2$	96
4.11.1	The exponent	97
4.12	$D_8 \times C_2$	97
4.12.1	Introduction	97
4.12.2	The decomposition	98
4.12.3	The Serre spectral sequence for the subgroups F_i	98
4.12.4	The Serre spectral sequence for the subgroup Z	99
4.12.5	The map j^*	99
	Inclusion of the center Z in F_1	100
	Inclusion of the center Z in F_2	100
	The quotient of $D_8 \times C_2/F_1$ by C_2	100
	The quotient of $D_8 \times C_2/F_2$ by C_2	101
	Summary	101
	The kernel of j^*	101
	The image of j^*	102
4.12.6	The E_2 -page	102
4.12.7	Multiplicative extensions	102
4.12.8	The exponent	102

5	Properties of the \mathcal{F}-homotopy limit spectral sequence for equivariant K-theory	104
5.1	Elementary abelian 2-groups	104
5.1.1	Introduction	104
5.1.2	The exponent of the cokernel of the edge map	105
	Splitting of the augmentation ideal	105
	A basis for $I(C_2^{\times n})$	105
	A basis for $\lim_{\vartheta(C_2^{\times n})_{\mathcal{E}}^{\text{op}}} I(-)$	106
	The edge map	107
5.1.3	A lower bound on $\exp_{\mathcal{E}} KU$	108
5.1.4	Odd primes	110
5.1.5	The real case for $p = 2$	111
A	Sage computations	112
A.1	Introduction	112
A.2	Summary of computations	112
A.2.1	$p = 2$	112
A.2.2	$p = 3$	113
A.2.3	$p = 5$	113
A.2.4	$p = 7$	114
A.2.5	$p = 11$	114
A.3	Sage code	114

Chapter 1

Introduction

1.1 Nilpotence in Borel equivariant cohomology

For a finite group G , consider a cohomology class $u \in H^*(BG; \mathbf{F}_p)$ that restricts to zero on all elementary abelian p -subgroups (groups of the form $(\mathbf{Z}/p)^{\times l}$). It is a theorem of Quillen that u is nilpotent. In fact, Quillen showed

1.1.1 Theorem ([Qui71c, Thm. 7.1]) *The map*

$$(1.1) \quad \widetilde{\text{res}}: H^*(BG; \mathbf{F}_p) \rightarrow \lim_{E \subset G \text{ el. ab. } p\text{-gp.}} H^*(BE; \mathbf{F}_p)$$

where the maps in the indexing category for the limit are given by restricting along subgroups and conjugation, is a uniform \mathcal{F}_p -isomorphism ([Qui71c, Thm. 7.1], [Qui71c, p. 556]), which means that there is an n such that

1. Every $u \in \ker(\widetilde{\text{res}})$ satisfies $u^n = 0$.
2. Every $v \in \lim_E H^*(BE; \mathbf{F}_p) \setminus \text{Im}(\widetilde{\text{res}})$ satisfies $v^{p^n} \in \text{Im}(\widetilde{\text{res}})$.

This result in group cohomology is a consequence of the following landmark theorem by Quillen in Borel equivariant cohomology, which is the main theorem of the same paper. For a G -space X , we write $H_G^*(X; \mathbf{F}_p)$ for the mod- p Borel equivariant cohomology of X .

1.1.2 Theorem ([Qui71c, Thm. 6.2]) *For X any paracompact G -space of finite cohomological dimension, the map*

$$\widetilde{\text{res}}: H_G^*(X; \mathbf{F}_p) \rightarrow \lim_{E \subset G \text{ el. ab. } p\text{-gp.}} H_E^*(X; \mathbf{F}_p)$$

is a uniform \mathcal{F}_p -isomorphism.

These results led to a lot of structural results in group cohomology. Quillen himself immediately deduced

1.1.3 Corollary ([Qui71c, Cor. 7.8]) *The Krull dimension of $H^*(BG; \mathbf{F}_p)$ equals the rank of the maximal elementary abelian p -subgroup of G .*

Results directly building on Theorem 1.1.2 include a theorem of Duflo on the depth of $H^*(BG; \mathbf{F}_p)$ ([Duf81, Thm. 1]), a theorem on the complexity of kG -modules by Alperin-Evens ([AE81]), Benson's description of the image of the transfer map ([Ben93, Thm. 1.1]), and a theorem on the depth of group cohomology rings by Carlson ([Car95, Thm. 2.3]). These results, and Quillen's original result all indicate the importance of the elementary abelian p -subgroups in Borel equivariant \mathbf{F}_p -cohomology in general, and group cohomology with \mathbf{F}_p -coefficients in particular.

It is natural to ask what one can say about the n in Theorem 1.1.1. One approach to this question is in [MNN15], where the map (1.1) is realized as the edge homomorphism of a homotopy limit spectral sequence, called the \mathcal{F} -homotopy limit spectral sequence,

$$(1.2) \quad E_2^{s,t} = \lim_{\mathcal{O}(G)_{\mathcal{F}}^{\text{op}}}^s H_H^t(X; \mathbf{F}_p) \Rightarrow H_G^*(X; \mathbf{F}_p)$$

converging strongly to the target ([MNN15, Prop. 2.24]), where X can be any G -space, and \mathcal{F} any family of subgroups of G which contains at least the family $\mathcal{E}_{(p)}$ of elementary abelian p -subgroups. The indexing category is the subcategory of the orbit category $\mathcal{O}(G)$ spanned by the orbits G/H with H in the family \mathcal{F} . The key property of the \mathcal{F} -homotopy limit spectral sequence is that it collapses at a finite page with a horizontal vanishing line ([MNN15, Thm. 2.25]).

This implies that every computation with the \mathcal{F} -homotopy limit spectral sequence is a finite one. Moreover, in many concrete situations we can establish a bound on the height of the horizontal vanishing line, and a bound on which page it will appear. Besides implying when the \mathcal{F} -homotopy limit spectral sequence will have collapsed, it can also be used to deduce differentials. This was illustrated in the computation of the cohomology of the quaternion group of order 8 in [MNN15, Ex. 5.18]. The first principal goal of this thesis is to demonstrate the computational utility of the \mathcal{F} -homotopy limit spectral sequence and the use of the vanishing line in determining differentials. This will be done by using it to compute the cohomology of all 2-groups up to order 16.

Varying X over all G -spectra, this horizontal vanishing line turns out to have a uniform bound in height ([MNN15, Prop. 2.26]). The minimal upper bound of this height is one of the equivalent definitions of the $\mathcal{E}_{(p)}$ -exponent $\exp_{\mathcal{E}_{(p)}} \underline{H}\mathbf{F}_{pG}$. In practice one can often determine this $\mathcal{E}_{(p)}$ -exponent, and this leads to a quantified version of Theorem 1.1.1, because one has $n \leq \exp_{\mathcal{E}_{(p)}} \underline{H}\mathbf{F}_{pG}$ ([MNN15, Thm. 3.24, Rem. 3.26]). The identification of $\mathcal{E}_{(2)}$ -exponents for 2-groups is the second principal goal of this thesis, and leads to the main theorem:

1.1.4 Theorem *Let G be a finite group with a 2-Sylow subgroup of order ≤ 16 , let X be any G -space, and let I be the kernel of*

$$\widetilde{\text{res}}: H_G^*(X; \mathbf{F}_2) \rightarrow \lim_{E \subset G \text{ cl. ab. 2-gp.}} H_E^*(X; \mathbf{F}_2).$$

Then $I^4 = 0$. Moreover, if u is not in the image of $\widetilde{\text{res}}$, then u^8 is.

This follows from combining [MNN15, Thm. 3.24] and Lemma 3.2.14 with the upper bounds on the exponents obtained in this thesis and summarized in Table 1.1. For a description of the groups appearing in the table we refer to

Section 4.2. The first column lists the groups of order ≤ 16 , the second column the exponent or an interval in which the exponent lies, and the last column gives a forward reference for the claim.

G	$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2G}$	Reference
e	1	Proposition 4.3.1
C_2	1	Proposition 4.3.1
$C_2 \times C_2$	1	Proposition 4.3.1
C_4	2	Proposition 4.3.1
$C_2^{\times 3}$	1	Proposition 4.3.1
$C_2 \times C_4$	2	Proposition 4.3.1
C_8	2	Proposition 4.3.1
D_8	2	Proposition 4.4.1
Q_8	4	[MNN15, Ex. 5.18]
$C_2^{\times 4}$	1	Proposition 4.3.1
$C_2^{\times 2} \times C_4$	2	Proposition 4.3.1
$C_4 \times C_4$	3	Proposition 4.3.1
$C_8 \times C_2$	2	Proposition 4.3.1
C_{16}	2	Proposition 4.3.1
D_{16}	2	Proposition 4.4.1
Q_{16}	4	Proposition 4.5.4
$SD_{16} = C_8 \rtimes^3 C_2$	[3, 4]	Proposition 4.6.2
$M_{16} = C_8 \rtimes^5 C_2$	[3, 4]	Proposition 4.7.4
$D_8 * C_4$	4	Proposition 4.8.22
$C_4 \rtimes C_4$	[3, 4]	Proposition 4.9.6
$(C_4 \times C_2) \rtimes^{\psi_5} C_2$	2	Proposition 4.10.1
$Q_8 \times C_2$	4	Proposition 4.11.1
$D_8 \times C_2$	2	Proposition 4.12.2

Table 1.1: The $\mathcal{E}_{(2)}$ -exponents of the groups of order ≤ 16 .

For specific 2-Sylow subgroups of order ≤ 16 it is possible to improve the previous theorem by using the upper bound on the relevant exponent.

The upper bounds in Table 1.1 have been obtained by either using the projective bundle theorem as in [MNN15, Ex. 5.18] or by using Euler classes and group cohomology computation with (1.2) as in [MNN15, Ex. 5.22]. We also refer to Proposition 3.3.3 for the former and Corollary 3.2.10 for the latter method.

The lower bounds on the exponents have been obtained from the computations of group cohomology with (1.2). These then serve a threefold purpose: firstly, they illustrate the computational power of the \mathcal{F} -homotopy limit spectral sequence, secondly they give us lower bounds on $\mathcal{E}_{(2)}$ -exponents, and lastly they can give us upper bounds on exponents. Not all our computations of group cohomology with (1.2) are done with the family $\mathcal{E}_{(2)}$, sometimes we chose a family which was strictly larger, if it lead us to a more manageable computation.

The structure of this thesis is as follows. In Chapter 2 we collect some results from [MNN15] about the \mathcal{F} -homotopy limit spectral sequence and \mathcal{F} -exponents. Some lemmas that can aid in the computation of exponents will be proven in Chapter 3. In Chapter 4, which is the main chapter, we determine

upper and lower bounds on $\exp_{\mathcal{E}_{(2)}} \underline{HF}_2$ for various small 2-groups, and use the \mathcal{F} -homotopy limit spectral sequence to compute group cohomology.

In Chapter 5, which is mostly unrelated to the previous chapters, we consider complex equivariant K -theory KU , which also admits \mathcal{F} -homotopy limit spectral sequences, but one needs to replace the family $\mathcal{E}_{(2)}$ by the family \mathcal{C} to get a horizontal vanishing line on a finite page ([MNN15, Prop. 5.6]). We prove lower bounds on $\exp_{\mathcal{C}} KU$ for E -equivariant K theory, where E is an elementary abelian 2-group. For primes $p \neq 2$ we conjecture a lower bound, which we verify for several small cases using Sage in Appendix A.

1.2 Acknowledgements

I would like to thank my supervisor Justin Noel in general for teaching me practically everything that I know about stable equivariant homotopy theory, and in particular for the help with the statement and proofs of the exponent lemmas in Section 3.2.

I would like to thank Akhil Mathew for a detailed explanation of the use of the projective bundle theorem in [MNN15, Ex. 5.18].

This thesis was written at SFB 1085 Higher Invariants, and I would like to thank all my colleagues for the great working atmosphere.

I would like to thank Kim and Christoph for all the nice discussions we had in their office, whose door was always open, for mathematics or otherwise.

I would like to thank Johannes for making more coffee than anyone else in the SFB, inventing analog divisibility testers, and helping me with the formalities of handing in a thesis.

I would like to thank Oriol for patiently answering all my questions about stable homotopy theory, and discovering and taking me to all the hidden cultural events in Regensburg.

I would like to thank Florent for initiating a shared effort to learn German, giving history lessons and bringing pastry to the office.

Lastly I would like to thank my parents and my brother, for their interest, support, and advice.

1.3 Notations and conventions

Throughout, G will denote a finite group.

Let G be a finite group, and $g_1, \dots, g_n \in G$ such that $\overline{g_1}, \dots, \overline{g_n} \in G^{\text{ab}}$ are a basis for $G^{\text{ab}} \otimes \mathbf{F}_p$ as an \mathbf{F}_p -vector space. We then denote the cohomology classes in $H^1(BG; \mathbf{F}_p)$ dual to $\overline{g_i} \in G^{\text{ab}}$ by $\delta_{\overline{g_i}}$.

For G a finite group, we denote by S_G the ∞ -category obtained from the simplicial model category of topological G -spaces with the level model structure. We denote by Sp_G the ∞ -category obtained from the symmetric monoidal simplicial model category of orthogonal G -spectra with the positive stable model structure (see [MM02, Def. 5.2]).

The cyclic group of order n we denote by C_n .

Chapter 2

The \mathcal{F} -homotopy limit spectral sequence

2.1 Introduction

In this chapter we recall the \mathcal{F} -homotopy limit spectral sequence as constructed in [MNN15], as well as some of the properties of this spectral sequence. For details of the constructions and for proofs of the statements, we refer to [MNN15] and [MNN17].

This chapter is organized as follows. We first recall the notion of a family of subgroups and the orbit category, which we will then use to state the \mathcal{F} -homotopy limit spectral sequence. One of its key properties is the fact that it has a horizontal vanishing line at a finite page. The height of this vanishing line and the page on which it appears can be bounded by the \mathcal{F} -exponent, which we then describe next. Finally we show that for specific families \mathcal{F} , the \mathcal{F} -homotopy spectral sequence (for $X = \text{pt}$) is isomorphic to a Lyndon-Hochschild-Serre spectral sequence in group cohomology.

2.2 Families of subgroups

The \mathcal{F} -homotopy limit spectral sequence depends on a family of subgroups \mathcal{F} . We recall this notion and describe some examples that will feature in this thesis.

2.2.1 Definition A family \mathcal{F} of subgroups of G is a non-empty set of subgroups closed under taking subgroups and conjugation.

2.2.2 Examples The following are examples:

1. The elementary abelian p -subgroups (groups isomorphic to $C_p^{\times n}$) with p a prime. This family is denoted $\mathcal{E}_{(p)}$.
2. The family \mathcal{C} of cyclic subgroups.
3. The family \mathcal{A} of abelian subgroups.
4. The family \mathcal{All} of all subgroups.

5. For a family \mathcal{F} of subgroups of G , and N any normal subgroup of G , let \mathcal{F}_N be the subset of groups in \mathcal{F} contained in N . Then \mathcal{F}_N is a family of subgroups of N , and also of G .

2.3 Orbit categories

The \mathcal{F} -homotopy limit spectral sequence has an E_2 -term the derived limit out of the opposite of a subcategory of the orbit category, of which we now recall the definition.

2.3.1 Definition For G a finite group, the orbit category $\mathcal{O}(G)$ is the subcategory of the category of G -sets spanned by the transitive G -sets G/H , where H ranges over the subgroups of G . For \mathcal{F} a family of subgroups of G , the category $\mathcal{O}(G)_{\mathcal{F}}$ is defined to be the full subcategory of $\mathcal{O}(G)$ spanned by the orbits G/H with the H ranging over the subgroups in \mathcal{F} .

2.4 The \mathcal{F} -homotopy limit spectral sequence

We now state the spectral sequence which is the main object of study of this thesis. We will use the following notation.

2.4.1 Notation We denote by $H_G^*(X; \mathbf{F}_2)$ the mod-2 Borel equivariant cohomology of a G -space X , which is by definition the ordinary cohomology $H^*(X \times_G EG; \mathbf{F}_2)$ of the homotopy orbits of X . The G -spectrum representing this cohomology theory we denote by $\underline{H}\mathbf{F}_{2G}$, or by $\underline{H}\mathbf{F}_2$ if the group is clear from the context.

2.4.2 Theorem ([MNN15, Def. 2.23, Thm. 2.25, Cor. 3.6, Prop. 5.16]) *Let G be a finite group, let X be a G -spectrum, and let \mathcal{F} be family of subgroups containing $\mathcal{E}_{(2)}$. Then there is a spectral sequence, called the \mathcal{F} -homotopy limit spectral sequence, given by*

$$E_2^{s,t}(X) = \lim_{G/H \in \mathcal{O}(G)_{\mathcal{F}}^{\text{op}}}^s H_H^t(X; \mathbf{F}_2) \Rightarrow H_G^{s+t}(X; \mathbf{F}_2).$$

This spectral sequence converges strongly to its target. Moreover, it collapses at a finite page with a horizontal vanishing line: there are $N, l \geq 1$ such that $E_{N+1}^{s,}(X) = 0$ for all $s \geq l$.*

In Chapter 4, we will illustrate the computational use of the \mathcal{F} -homotopy limit spectral sequence by using it when $X = \text{pt}$, is a point, to compute the cohomology of all 2-groups up to order 16.

2.4.3 Remark We abbreviate $E_2^{s,t} = E_2^{s,t}(\text{pt})$ for the \mathcal{F} -homotopy limit spectral sequence for $X = \text{pt}$.

The \mathcal{F} -homotopy limit spectral sequence can be constructed for any generalized equivariant cohomology theory E ([MNN15, Def. 2.23]). The case considered in Theorem 2.4.2 is $E = \underline{H}\mathbf{F}_{2G}$. For different choices of E a statement about a vanishing line analogous to the one in Theorem 2.4.2 also holds, but one needs to change the family $\mathcal{E}_{(2)}$ to the derived defect base of E ([MNN15, remark after def. 1.3, Thm. 2.25]). We refer to [MNN15] for the details.

The vanishing line mentioned in Theorem 2.4.2 and the page on which it appears turn out to have an upper bound uniform in the spectrum X . To describe it, we introduce the notion of \mathcal{F} -nilpotence in the next section.

2.5 \mathcal{F} -nilpotence

We recall the notion of \mathcal{F} -nilpotence from [MNN15], for which we first need to recall the following space.

2.5.1 Definition Let \mathcal{F} be a family of subgroups of G . Then the **universal \mathcal{F} -space** $E\mathcal{F}$ is the G -space given by the homotopy colimit

$$E\mathcal{F} = \operatorname{hocolim}_{\mathcal{O}(G)} G/H.$$

The space $E\mathcal{F}$ is, up to G -equivalence, characterized by ([LMSM86, Def. II.2.10])

$$E\mathcal{F}^H \simeq \begin{cases} \emptyset & \text{if } H \notin \mathcal{F}, \\ \text{pt} & \text{if } H \in \mathcal{F}. \end{cases}$$

Using the space $E\mathcal{F}$, we can give one of the equivalent definitions of \mathcal{F} -nilpotence.

2.5.2 Definition (cf. [MNN15, Def. 1.3]) Let M be a G -spectrum. Then M is said to be \mathcal{F} -nilpotent if there is an n such that M is a retract of $F(\operatorname{sk}_{n-1} E\mathcal{F}_+, M)$. The minimal $n \geq 0$ for which this holds is called the \mathcal{F} -exponent of M , and denoted $\exp_{\mathcal{F}} M$.

Being \mathcal{F} -nilpotent is a strong condition, it implies for example the following.

2.5.3 Proposition ([MNN15, Prop. 2.8]) *Let M be \mathcal{F} -nilpotent. Then M is \mathcal{F} -complete and \mathcal{F} -colocal, that is, the \mathcal{F} -completion map $M \rightarrow M(E\mathcal{F}_+, M)$ and the colocalization map $M \wedge E\mathcal{F}_+ \rightarrow M$ are weak equivalences.*

The primary case of interest for us is $M = \underline{HF}_{2G}$.

2.5.4 Proposition ([MNN15, Prop. 5.16]) *For G any group, the G -spectrum \underline{HF}_{2G} is $\mathcal{E}_{(2)}$ -nilpotent.*

2.5.5 Remark In fact, [MNN15, Prop. 5.16] shows that $\mathcal{E}_{(2)}$ is the minimal family \mathcal{F} such that \underline{HF}_{2G} is \mathcal{F} -nilpotent. In the terminology of [MNN15] we have that $\mathcal{E}_{(2)}$ is the derived defect base of \underline{HF}_{2G} .

A trivial example is the following.

2.5.6 Proposition *Every G -spectrum M is $\mathcal{A}ll$ -nilpotent with $\exp_{\mathcal{A}ll} M \leq 1$, and $\exp_{\mathcal{A}ll} M = 0$ if and only if M is contractible.*

PROOF We have $E\mathcal{A}ll \simeq \text{pt}$, and every spectrum M is a retract of $F(S, M) \simeq M$.

A spectrum M has $\mathcal{A}ll$ -exponent 0 if and only if it is a retract of $F(\text{pt}, M) \simeq \text{pt}$, which happens if and only if M is contractible. \square

The following two propositions are immediate with some alternative definitions of \mathcal{F} -nilpotence, but not with the one we have given.

2.5.7 Proposition ([MNN17, Prop. 6.39]) *A G -spectrum M is \mathcal{F} -nilpotent and \mathcal{G} -nilpotent if and only if M is $\mathcal{F} \cap \mathcal{G}$ -nilpotent.*

2.5.8 Proposition ([MNN15, Def. 1.3]) *If M is \mathcal{F} -nilpotent and $\mathcal{G} \supset \mathcal{F}$, then M is \mathcal{G} -nilpotent.*

Combining Proposition 2.5.6, Proposition 2.5.7 and Proposition 2.5.8 shows that every G -spectrum M has a minimal family \mathcal{F} such that M is \mathcal{F} -nilpotent (cf. the remark after [MNN15, Def. 1.3]).

2.5.9 Definition ([MNN15, remark after Def. 1.3]) This minimal family \mathcal{F} such that a G -spectrum M is \mathcal{F} -nilpotent is called the **derived defect base** of M .

The following is the main case of interest for us.

2.5.10 Proposition ([MNN15, Prop. 5.16]) *For G any finite group, the derived defect base of $H\mathbf{F}_{2G}$ is $\mathcal{E}_{(2)}$.*

We can now state the uniform upper bound in the height of the horizontal vanishing of the \mathcal{F} -homotopy limit spectral sequence and the page on which it appears.

2.5.11 Proposition ([MNN15, Prop. 2.26, Rem. 2.27]) *Let G be a finite group, and $\mathcal{F} \supset \mathcal{E}_{(2)}$ a family of subgroups, M an \mathcal{F} -nilpotent G -spectrum. Then the following integers equal:*

- (1) *The \mathcal{F} -exponent of M .*
- (2) *The minimal N such that for all G -spectra X , the \mathcal{F} -homotopy limit spectral sequence $E_*^{*,*}(X)$ admits a vanishing line of height N on the $N+1$ -page: $E_{N+1}^{s,*} = E_\infty^{s,*} = 0$ for all $s \geq N$.*
- (3) *The minimal n such that the canonical map $F(E_{\mathcal{F}_+}, M) \simeq M \rightarrow F(\mathrm{sk}_{n-1} E_{\mathcal{F}_+}, M)$ admits a section.*
- (4) *The minimal n' such that there is an $(n-1)$ -dimensional CW-complex X with isotropy in \mathcal{F} such that M is a retract of $F(X_+, M)$.*
- (5) *The minimal m such that the canonical map $\mathrm{sk}_{m-1} E_{\mathcal{F}} \wedge M \rightarrow M$ admits a retraction.*
- (6) *The minimal m' such that there is an $(m'-1)$ -dimensional CW-complex X with isotropy in \mathcal{F} such that M is a retract of $X_+ \wedge M$.*

Moreover, if M' is any G -spectrum, then the existence of an integer for M' as in any one of the items from (2) to (6) implies that M' is \mathcal{F} -nilpotent.

An immediate consequence is that exponents are invariant under suspension:

2.5.12 Corollary *Let M be a G -spectrum, \mathcal{F} a family and s an integer. Then M is \mathcal{F} -nilpotent with \mathcal{F} -exponent n if and only if $\Sigma^s M$ is \mathcal{F} -nilpotent with \mathcal{F} -exponent n .*

PROOF This follows from suspending the retraction sequences from Proposition 2.5.11 (5). \square

We end this section by recalling from [MNN15] some properties of exponents that will be used in the next chapter to prove lemmas about exponents.

2.5.13 Proposition *Let $H \in \mathcal{F}$. Then G/H_+ is \mathcal{F} -nilpotent with $\exp_{\mathcal{F}} G/H_+ = 1$.*

PROOF Because $F(G/H_+, G/H_+) \simeq G/H_+ \wedge G/H_+$ by self duality of G -sets in Sp_G ([LMSM86, Cor. 6.3]), the result follows the suspending the following composite of G -spaces, which is the composite of the diagonal followed by the projection

$$G/H \rightarrow G/H \times G/H \rightarrow G/H$$

to get a retraction \square

$$G/H_+ \rightarrow G/H_+ \wedge G/H_+ \rightarrow G/H_+$$

2.5.14 Proposition ([MNN17, Cor. 4.15]) *If M is an \mathcal{F} -nilpotent spectrum and X is any G -spectrum, then $F(X, M)$ is \mathcal{F} -nilpotent with $\exp_{\mathcal{F}} F(X, M) \leq \exp_{\mathcal{F}} M$.*

2.5.15 Proposition *If N is an \mathcal{F} -nilpotent G -spectrum and M is any G -spectrum then $M \wedge N$ is \mathcal{F} -nilpotent with $\exp_{\mathcal{F}} M \wedge N \leq \exp_{\mathcal{F}} N = n$.*

PROOF Because N is \mathcal{F} -nilpotent there is, by Proposition 2.5.11 (6), an $(n-1)$ -dimensional CW-complex with isotropy in \mathcal{F} and a retraction

$$N \rightarrow X_+ \wedge N \rightarrow N,$$

Smashing this with M gives a retraction

$$M \wedge N \rightarrow X_+ \wedge M \wedge N \rightarrow M \wedge N,$$

and another application of Proposition 2.5.11 (6) gives the result. \square

2.5.16 Proposition ([MNN17, Prop. 4.9]) *1. If M is a retract of an \mathcal{F} -nilpotent spectrum N , then M is \mathcal{F} -nilpotent and $\exp_{\mathcal{F}} M \leq \exp_{\mathcal{F}} N$.*

2. If M' and M'' are \mathcal{F} -nilpotent and $M' \rightarrow M \rightarrow M''$ is a cofiber sequence then M is \mathcal{F} -nilpotent and $\exp_{\mathcal{F}} M \leq \exp_{\mathcal{F}} M' + \exp_{\mathcal{F}} M''$.

2.5.17 Proposition *Let M_α be a set of \mathcal{F} -nilpotent spectra with \mathcal{F} -exponents bounded uniformly by n . Then $\bigvee_{\alpha} M_\alpha$ is \mathcal{F} -nilpotent with \mathcal{F} -exponent $\leq n$.*

PROOF By Proposition 2.5.11 (5) there are retractions

$$M_\alpha \rightarrow \mathrm{sk}_{n-1} E\mathcal{F}_+ \wedge M_\alpha \rightarrow M_\alpha.$$

Wedge them together and use that smashing commutes with arbitrary colimits to obtain a retraction

$$\bigvee_{\alpha} M_{\alpha} \rightarrow \mathrm{sk}_{n-1} E_{\mathcal{F}} \wedge \bigvee_{\alpha} M_{\alpha} \rightarrow \bigvee_{\alpha} M_{\alpha}$$

and apply Proposition 2.5.11 (5) again. □

2.5.18 Proposition *Let X be an $(n - 1)$ -dimensional G -CW-complex with isotropy in \mathcal{F} . Then X_+ is \mathcal{F} -nilpotent and $\exp_{\mathcal{F}} X_+ \leq n$.*

PROOF By induction on n . For $n = 1$, X is of the form

$$\bigvee_{H \in \mathcal{F}} \left(G/H \wedge \left(\bigvee_{\alpha} S^0 \right) \right),$$

which is \mathcal{F} -nilpotent of exponent 1 by Proposition 2.5.15, Proposition 2.5.17 and Proposition 2.5.13.

Assume we have the result for $n - 1$, and let X be an n -dimensional CW-complex. Let $X^{(n-1)}$ be the $(n - 1)$ -skeleton of X . Then X sits in a cofiber sequence

$$X^{(n-1)} \rightarrow X \rightarrow \bigvee_{H \in \mathcal{F}} \left(G/H \wedge \left(\bigvee_{\alpha} S^n \right) \right).$$

Now $\exp_{\mathcal{F}} X^{(n-1)} \leq n$ by induction, and the right hand side has \mathcal{F} -exponent ≤ 1 by Proposition 2.5.15, Proposition 2.5.17 and Proposition 2.5.13. □

2.5.19 Proposition *Let X be a finite G -CW complex with isotropy in \mathcal{F} . Then the equivariant Spanier-Whitehead dual $\mathbb{D}(X_+)$ of X_+ is \mathcal{F} -nilpotent, and $\exp_{\mathcal{F}} \mathbb{D}(X_+) = \exp_{\mathcal{F}} X_+$.*

PROOF Write $n = \exp_{\mathcal{F}} X_+$. Let Y_+ be an $(n - 1)$ finite-dimensional G -CW complex with isotropy in \mathcal{F} such that there is a retraction

$$X_+ \rightarrow F(Y_+, X_+) \simeq \mathbb{D}(Y_+) \wedge X_+ \rightarrow X_+.$$

Applying $\mathbb{D}(-)$ to this retraction exhibits $\mathbb{D}(X_+)$ as a retraction of $Y_+ \wedge \mathbb{D}(X_+)$, which has exponent $\leq n$ by Proposition 2.5.19 and Proposition 2.5.15. Therefore $\exp_{\mathcal{F}} \mathbb{D}(X_+) \leq \exp_{\mathcal{F}} X_+$, and replacing X_+ by $\mathbb{D}(X_+)$ in this inequality shows equality. □

2.6 Comparison with the Lyndon-Hochschild-Serre spectral sequence

Let N be a normal subgroup of G , and let $\mathcal{A}ll_N$ be the family of subgroups of N . We then have the following identification of spectral sequences.

2.6.1 Proposition ([MNN15, Lem. A.3]) *The $\mathcal{A}ll_N$ -homotopy limit spectral sequence*

$$\lim_{\mathcal{O}(G)_{\mathcal{A}ll_N}^{\text{op}}}^s H^t(BH; \mathbf{F}_2) \Rightarrow H^{s+t}(BG; \mathbf{F}_2)$$

is isomorphic to the Lyndon-Hochschild-Serre spectral sequence

$$H^s(BG/N; H^t(BN; \mathbf{F}_2)) \Rightarrow H^{s+t}(BG; \mathbf{F}_2).$$

We will make use of this identification in all our computations. In some cases we extend the family $\mathcal{E}_{(2)}$ to a family which is of the form $\mathcal{A}ll_N$, but this is not always possible because there is not always a proper normal subgroup containing all the elementary abelian subgroups.

2.7 A decomposition of the E_2 -page.

In Section 2.6, we saw that the \mathcal{F} -homotopy limit spectral sequence reduces to the LHSSS in case the family \mathcal{F} is of the form $\mathcal{F} = \mathcal{A}ll_N$, for some normal subgroup N of G . In case the family \mathcal{F} is not of this form, we will compute the E_2 -term of the \mathcal{F} -homotopy limit spectral sequence following the strategy of [MNN15, App. B]. In this section we describe this strategy in the abstract.

2.7.1 Homotopy cofinality

To compute the ordinary categorical limit of a diagram $I \rightarrow \mathbf{C}$ indexed by a category I , one can restrict along a left cofinal functor $I' \rightarrow I$. The corresponding notion for homotopy limits is homotopy left finality, which can be characterized as follows for homotopy limits indexed by ordinary categories. We have taken the dual of the cited Proposition. We denote by $N(-)$ the nerve of an ordinary category.

2.7.1 Proposition ([Lur09, Prop. 4.1.1.8]) *Let $\iota: I' \rightarrow I$ be a functor of ordinary categories. The following are equivalent.*

- (1) *The functor ι is homotopy left cofinal.*
- (2) *For \mathbf{C} any ∞ -category and $I \rightarrow \mathbf{C}$ any diagram, the induced map of homotopy limits by ι is a weak equivalence. That is, taking right Kan extensions of*

$$\begin{array}{ccc} N(I) & \longrightarrow & \mathbf{C} \\ \downarrow & \nearrow \text{dashed} & \\ \text{pt} & & \end{array}$$

and of

$$\begin{array}{ccc} N(I') & \longrightarrow & \mathbf{C} \\ \downarrow & \nearrow \text{dashed} & \\ \text{pt} & & \end{array}$$

yields parallel dashed arrows

$$\begin{array}{ccccc} N(I') & \longrightarrow & N(I) & \longrightarrow & C \\ & \searrow & \downarrow & \dashrightarrow & \uparrow \\ & & \text{pt} & & \end{array}$$

with a natural transformation between them. This natural transformation is an equivalence.

2.7.2 Homotopy cofinality in orbit categories.

Let \mathcal{F} be a family of subgroups of G , and let $\{H_i\}$ be the maximal subgroups in \mathcal{F} . Assume there is a subgroup K of G such that K equals any intersection of at least two of the H_i :

$$\bigcap_{i \in I} H_i = K,$$

for all I with $|I| \geq 2$. In particular, the set $\{H_i\} \cup \{K\}$ is closed under intersections. Because the set $\{H_i\}$ is closed under conjugations, so is $\{H_i\} \cup \{K\}$. Denote by $\mathcal{O}_{\mathcal{F}}$ the subcategory of $\mathcal{O}(G)_{\mathcal{F}}$ spanned by the G/J with J in the set $\{H_i\} \cup \{K\}$. Now [MNN17, Prop. 6.31] implies the following:

2.7.2 Proposition *The subcategory $\mathcal{O}_{\mathcal{F}}^{\text{op}} \rightarrow \mathcal{O}(G)_{\mathcal{F}}^{\text{op}}$ is homotopy left cofinal.*

The construction of the homotopy limit spectral sequence by Bousfield-Kan ([BK72, Ch. XI]), applied to the diagram

$$\begin{array}{c} \mathcal{O}(G)_{\mathcal{F}}^{\text{op}} \rightarrow \text{Sp}, \\ G/H_+ \mapsto F(G/H_+, \underline{HF}_2)^G, \end{array}$$

yields the \mathcal{F} -homotopy limit spectral sequence ([MNN15, Prop. 2.24]).

2.7.3 Proposition *The inclusion functor $\iota: \mathcal{O}_{\mathcal{F}} \rightarrow \mathcal{O}(G)_{\mathcal{F}}$ induces a morphism of homotopy limit spectral sequences which is an isomorphism from E_2 onward.*

PROOF The functor ι induces a map of the cosimplicial spectra used in the construction of the homotopy limit spectral sequence, hence a map of homotopy limit spectral sequences. This map is an isomorphism on E_2 by the identification of E_2 in [BK72, §7.1]. \square

In our computations with the \mathcal{F} -homotopy limit spectral sequence we will make use of the previous two propositions by restricting to homotopy left cofinal subcategories of the orbit category when computing the E_2 -page of the \mathcal{F} -homotopy limit spectral sequence.

2.7.3 A short exact sequence of coefficient systems

We assume in addition that all the H_i and the K are normal subgroups of G .

Let $\underline{\mathbf{Z}}$ be the constant \mathbf{Z} -coefficient system on $\mathcal{O}(G)_{\mathcal{F}}$. For a subgroup H of G , we denote by $\mathbf{Z}[H]$ the coefficient system obtained from restricting $\underline{\mathbf{Z}}$ to

$\mathcal{O}(G)_{\mathcal{A}ll_H}$ and left Kan extending back up to $\mathcal{O}(G)_{\mathcal{F}}$. Summing the counit maps $\mathbf{Z}[H_i] \rightarrow \mathbf{Z}$ yields a short exact sequence of coefficient systems

$$(2.1) \quad 0 \rightarrow \bigoplus_{|I|-1} \mathbf{Z}[K] \xrightarrow{j} \bigoplus_{i \in I} \mathbf{Z}[H_i] \rightarrow \mathbf{Z} \rightarrow 0$$

where the map j is given by the matrix

$$(2.2) \quad \begin{pmatrix} 1 & 0 & \cdots & \ddots & \\ -1 & 1 & 0 & \ddots & \\ 0 & -1 & 1 & \ddots & \\ \vdots & 0 & -1 & \ddots & \vdots \\ & \vdots & 0 & \ddots & 0 \\ & & \vdots & \ddots & 1 \\ & & & \ddots & -1 \end{pmatrix}$$

with $|I|$ rows and $|I| - 1$ columns.

2.7.4 Lemma *The short exact sequence (2.1) lifts to a cofiber sequence of spectra*

$$(2.3) \quad \bigvee_{|I|-1} E\mathcal{A}ll_{K_+} \rightarrow \bigvee_{i \in I} E\mathcal{A}ll_{H_{i+}} \rightarrow E\mathcal{F}_+.$$

PROOF The diagram of categories

$$(2.4) \quad \begin{array}{ccc} \mathcal{O}(G)_{\mathcal{A}ll_K} & \hookrightarrow & \mathcal{O}(G)_{\mathcal{A}ll_{H_i}} \\ \downarrow & & \\ \mathcal{O}(G)_{\mathcal{A}ll_{H_{i+1}}} & & \end{array}$$

assemble together to a coequalizer diagram

$$(2.5) \quad \coprod_{|I|-1} \mathcal{O}(G)_{\mathcal{A}ll_K} \rightrightarrows \coprod_{i \in I} \mathcal{O}(G)_{\mathcal{A}ll_{H_i}} \dashrightarrow \mathcal{O}(G)_{\mathcal{F}}$$

with coequalizer the category $\mathcal{O}(G)_{\mathcal{F}}$. For any family \mathcal{G} , denote by $\iota_{\mathcal{G}}: \mathcal{O}(G)_{\mathcal{G}} \rightarrow S_G$ the inclusion into G -spaces. Then $E\mathcal{G} \simeq \text{hocolim}_{\mathcal{O}(G)_{\mathcal{G}}} \iota_{\mathcal{G}}$ ([MNN15, p. 10]). Applying this to (2.5) and the fact that homotopy colimits commute with coproducts yields a cofiber sequence of G -spaces

$$(2.6) \quad \bigvee_{|I|-1} E\mathcal{A}ll_K \rightarrow \bigvee_{i \in I} E\mathcal{A}ll_{H_i} \rightarrow E\mathcal{F},$$

and, after taking suspension spectra, a cofiber sequence of G -spectra. After applying $H_*(-; \mathbf{Z})$ to this cofiber sequence of G -spectra, we get a long exact

sequence that reduces to a short exact sequence because all the homology is concentrated in degree 0. The left hand map is given by the matrix (2.1) but with all -1 entries replaced by 1. The bottom entry in every column comes from the vertical maps in the (2.4), the top entry in every column entries comes from the horizontal maps in the (2.4), for all i . To change the bottom entries into -1 , which is necessary to make a lift of (2.1), we do the following: since suspension preserves homotopy colimits, we can replace at the level of G -spectra the vertical maps by the vertical maps smashed with $S \xrightarrow{-1} S$, which preserves cofibrations. We then get a strict map of homotopy colimit diagrams on objects given by either identity maps or the identity smashed with $S \xrightarrow{-1} S$, and therefore an equivalence of the homotopy colimits. We now have that

$$(2.7) \quad 0 \rightarrow \bigoplus_{|I|-1} H_0(E\mathcal{A}\ell(-); \mathbf{Z}) \rightarrow \bigoplus_{i \in I} H_0(E\mathcal{A}\ell_{H_i}(-); \mathbf{Z}) \rightarrow H_0(E\mathcal{F}(-); \mathbf{Z}) \rightarrow 0.$$

coincides with (2.1) on the level of objects and the left map. For any family \mathcal{G} we have $\mathbf{Z}[\mathcal{G}](-) = H_0(E\mathcal{G}(-); \mathbf{Z})$ ([MNN15, sec. 3]), so the objects of the short exact sequences (2.1) and (2.7) are equal. By construction, the left maps of (2.1) and (2.7) are equal. The right hand maps possibly differ up to a sign, but postcomposing with the weak equivalence $E\mathcal{F}_+ \xrightarrow{-1} E\mathcal{F}_+$ makes sure that the right hand maps are equal on the level of $H_0(-)$. \square

For a normal subgroup H of G , we denote by $E_*^{*,*}(H)$ the LHSSS associated to the group extension $H \rightarrow G \rightarrow G/H$. By Proposition 2.6.1, the $\mathcal{A}\ell_H$ -homotopy limit spectral sequence is isomorphic to $E_*^{*,*}(H)$. The \mathcal{F} -homotopy limit spectral sequence we denote by $E_*^{*,*}$.

2.7.5 Lemma *Mapping (2.3) into \underline{HF}_2 and taking Atiyah-Hirzebruch spectral sequences yields a long exact sequence of spectral sequences, which on the rows $E_2^{s,*}$ of the E_2 -pages is given by*

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \bigoplus_{|I|-1} E_2^{2,*}(K) & \xleftarrow{j^2} & \bigoplus_{i \in I} E_2^{2,*}(H_i) & \xleftarrow{\partial^1} & E_2^{2,*} \\ & & & & & \searrow & \\ & & \bigoplus_{|I|-1} E_2^{1,*}(K) & \xleftarrow{j^1} & \bigoplus_{i \in I} E_2^{1,*}(H_i) & \xleftarrow{\partial^0} & E_2^{1,*} \\ & & & & & \searrow & \\ & & \bigoplus_{|I|-1} E_2^{0,*}(K) & \xleftarrow{j^0} & \bigoplus_{i \in I} E_2^{0,*}(H_i) & \xleftarrow{\quad} & E_2^{0,*} \xleftarrow{\quad} 0 \end{array}$$

Figure 2.1: A long exact sequence of rows $E_2^{s,*}$ of E_2 -pages.

where we have written j^s for the map in degree s in (2.1) induced by j , and ∂^s for the degree s -boundary morphism in (2.1). The maps j^s are given by the transpose of the matrix (2.2). Summing the columns in (2.1) gives the long exact sequence of spectral sequences.

PROOF We use the following standard construction of the AHSS: take a Postnikov tower on \underline{HF}_2 , and take homotopy fibers at each stage (i.e. the Postnikov

sections of $\underline{H}\mathbf{F}_2$). Apply $F(W, -)$ to this Postnikov tower for W the three objects in the cofiber sequence (2.3). Taking homotopy groups gives three exact couples, with associated spectral sequences the AHSS's $H^*(W; \pi_*^{(-)} \underline{H}\mathbf{F}_2)$. Moreover, since the objects W sit in a cofiber sequence, we get a long exact sequence of exact couples, whence of spectral sequences. The identification of the AHSS's with \mathcal{F} -homotopy limit spectral sequence from the E_2 -page onward is [MNN15, Prop. 2.24]), the identification of $\mathcal{A}ll_K$ - and $\mathcal{A}ll_{H_i}$ -homotopy limit spectral sequences with LHSS's is Proposition 2.6.1.

The identification of the maps j^s follows from the identifications

$$H^s(E\mathcal{G}; \pi_*^{(-)} \underline{H}\mathbf{F}_2) \cong \text{Ext}_{\mathbf{Z}\ell(G)}^{s,t}(\mathbf{Z}[\mathcal{G}], \pi_*^{(-)} \underline{H}\mathbf{F}_2)$$

from [MNN15, Cor. 3.6], for every family \mathcal{G} , and the fact that (2.6) lifts (2.1). \square

We write

$$(2.8) \quad \begin{aligned} A^* &= \text{Im } \partial^{*-1} = \text{coker } j^{*-1}, \\ B^* &= \text{coker } \partial^{*-1} = \text{ker } j^*, \end{aligned}$$

(where we have suppressed t -degrees) and obtain the following proposition:

2.7.6 Proposition *The E_2 -page of the homotopy limit spectral sequence fits in a short exact sequence*

$$(2.9) \quad 0 \rightarrow A^* \rightarrow E_2^* \rightarrow B^* \rightarrow 0$$

of spectral sequences (where we suppressed t -degrees). This short exact sequence splits as a short exact sequence of bigraded \mathbf{F}_2 -modules.

PROOF The short exact sequence is obtained from summing (2.8) over the $s = *$ -degree. Any short exact sequence splits as \mathbf{F}_2 -modules, and necessarily preserves the bigrading because the maps in the short exact sequence preserve the bigrading. \square

Chapter 3

Exponent lemmas

3.1 Introduction

We discuss some lemmas that will be of use in determining (bounds on) exponents. Some of these statements appear as exercises in [MNN17, sec. 4].

3.2 Lemmas for \mathcal{F} -exponents

We now give various lemmas which describe how \mathcal{F} -exponents can change as the family \mathcal{F} varies.

3.2.1 Lemma *Let $\mathcal{F}_1, \mathcal{F}_2$ be two families of subgroups. Then $E(\mathcal{F}_1 \cap \mathcal{F}_2)_+ \simeq E\mathcal{F}_{1+} \wedge E\mathcal{F}_{2+}$.*

PROOF This follows from comparing fixed points. \square

3.2.2 Notation For \mathcal{F} a family of subgroups of G , and H a normal subgroup of G , denote by \mathcal{F}_H the subgroups of \mathcal{F} that are contained in H . If \mathcal{F} is a family that make sense for all groups G , such as the family of all subgroups, the family of elementary abelian p -groups, etc., we write $\mathcal{F}(G)$ for this family of subgroups of G . For instance, we write $\mathcal{E}_{(2)}(D_8)$ for the elementary abelian subgroups of the dihedral group of order 8.

3.2.3 Lemma *If \mathcal{F} is a family of subgroups of G , and H is a normal subgroup of G , then $\text{Res}_H^G E\mathcal{F} \simeq E\mathcal{F}_H$.*

PROOF This follows from comparing fixed points. \square

3.2.4 Lemma *Let M be an \mathcal{F} -nilpotent G -spectrum, $H \subset G$ a subgroup. Then $\text{Res}_H^G M$ is \mathcal{F}_H -nilpotent, and $\exp_{\mathcal{F}_H} \text{Res}_H^G M \leq \exp_{\mathcal{F}} M$.*

PROOF This follows from [MNN17, Cor. 4.13] and Lemma 3.2.3. \square

Recall that for a group G , we denote the spectrum representing Borel G -equivariant \mathbf{F}_p -cohomology by $\underline{H\mathbf{F}}_p_G$ (Notation 2.4.1).

3.2.5 Corollary *Let G be a group and $H \subset G$ a subgroup. Then*

$$\exp_{\mathcal{E}(p)(H)} \underline{HF}_{p_H} \leq \exp_{\mathcal{E}(p)(G)} \underline{HF}_{p_G}.$$

3.2.6 Lemma *Let $\mathcal{F}_1, \mathcal{F}_2$ be two families of subgroups, and let M be a G -spectrum which is both \mathcal{F}_1 - and \mathcal{F}_2 -nilpotent, with exponents m, n respectively. Then M is $\mathcal{F}_1 \cap \mathcal{F}_2$ -nilpotent, and*

$$\exp_{\mathcal{F}_1 \cap \mathcal{F}_2} M \leq m + n - 1.$$

PROOF The fact that M is $\mathcal{F}_1 \cap \mathcal{F}_2$ -nilpotent is part of [MNN17, Prop. 6.39]. The assumption on the exponents implies that both maps in

$$(3.1) \quad \begin{aligned} \text{sk}_{m-1} E\mathcal{F}_{1+} \wedge \text{sk}_{n-1} E\mathcal{F}_{2+} \wedge M &\rightarrow \text{sk}_{m-1} E\mathcal{F}_{1+} \wedge E\mathcal{F}_{2+} \wedge M \\ &\rightarrow E\mathcal{F}_{1+} \wedge E\mathcal{F}_{2+} \wedge M \end{aligned}$$

have a retraction, hence the composite has a retraction. The composite (3.1) factors as

$$\begin{array}{ccc} \text{sk}_{m-1} E\mathcal{F}_{1+} \wedge \text{sk}_{n-1} E\mathcal{F}_{2+} & \xrightarrow{(*)} & E\mathcal{F}_{1+} \wedge E\mathcal{F}_{2+} \wedge M \\ \downarrow (**) & \nearrow (***) & \\ \text{sk}_{m+n-2} (E\mathcal{F}_{1+} \wedge E\mathcal{F}_{2+}) \wedge M & & \end{array}$$

Composing the retraction of $(*)$ with $(**)$ gives a retraction of $(***)$. Since $E\mathcal{F}_{1+} \wedge E\mathcal{F}_{2+} \simeq E(\mathcal{F}_1 \cap \mathcal{F}_2)_+$ by Lemma 3.2.1, the desired bound follows from Proposition 2.5.11 (5). \square

3.2.7 Notation For a G -space X , we denote by $\mathcal{I}(X)$ the minimal family containing the isotropy groups of X .

3.2.8 Example For an orthogonal G -representation V , the unit sphere $S(V)$ in V inherits a G -action. Then $\mathcal{I}(S(V))$ the smallest family containing the isotropy groups of $S(V)$.

3.2.9 Lemma (cf. Proof of [MNN15, Thm. 2.3]) *Let R be a ring G -spectrum with multiplicative Thom classes (see [MNN15, Def. 5.1]). Let V be a G -representation with corresponding oriented Euler class*

$$(\chi(V): S^{-|V|} \rightarrow R) \in R^*$$

Suppose $\chi(V)$ is nilpotent with $\chi(V)^n = 0$. Then R is $\mathcal{I}(S(V))$ -nilpotent and

$$\exp_{\mathcal{I}(S(V))} R \leq n \dim_{\mathbf{R}} V.$$

PROOF The fact that $\chi(V)^n = 0$ is equivalent to the oriented Euler class

$$R \xrightarrow{e_n V} S^{nV} \wedge R$$

being nullhomotopic (see [MNN15, Lem. 5.3] and [MNN15, Rem. 2.4]). Hence the left map in the fiber sequence

$$S(nV)_+ \wedge R \rightarrow R \rightarrow S^{nV} \wedge R$$

has a section: R is a retract of $S(nV)_+ \wedge R$. But $S(nV)$ is a $(n \dim_{\mathbf{R}} V - 1)$ -dimensional G -CW complex, hence by Proposition 2.5.19, $\exp_{\mathcal{F}} S(nV)_+ \leq n \dim_{\mathbf{R}} V$, and hence the same bound holds for R by Proposition 2.5.16. \square

3.2.10 Corollary *Let V be a d -dimensional G -representation with corresponding oriented Euler class (see [MNN15, Def. 5.1])*

$$(\chi(V): S^{-d} \rightarrow \underline{H\mathbf{F}}_2) \in H^{|\mathbf{V}|}(BG; \mathbf{F}_2).$$

Suppose $\chi(V)$ is nilpotent with $\chi(V)^n = 0$. Then $\underline{H\mathbf{F}}_2$ is $\mathcal{S}(S(V))$ -nilpotent with $\exp_{\mathcal{S}(S(V))} \underline{H\mathbf{F}}_2 \leq nd$.

3.2.11 Lemma *Let $f: G \rightarrow C_2 \cong O(1)$ be a 1-dimensional real representation with oriented Euler class $e \in H^1(BG; \mathbf{F}_2)$. Suppose e is nilpotent with n the minimal integer ≥ 0 such that $e^n = 0$. Then $\underline{H\mathbf{F}}_2$ is nilpotent for the family $\mathcal{A}\ell_{\ker f}$ of subgroups of $\ker f$ with $\exp_{\mathcal{A}\ell_{\ker f}} \underline{H\mathbf{F}}_2 = n$.*

PROOF The upper bound for the exponent follows immediately from Corollary 3.2.10.

For the lower bound on the exponent, denote a class representing e on the E_{∞} -page of the $\mathcal{A}\ell_{\ker f}$ -limit spectral sequence converging to $H^*(BG; \mathbf{F}_2)$ by \tilde{e} . Since e restricts to 0 on $\ker f$, hence on all subgroups in $\mathcal{A}\ell_{\ker f}$, we know that \tilde{e} lives in filtration degree ≥ 1 on E_{∞} . By assumption, $e^{n-1} \neq 0$. This does not imply $\tilde{e}^{n-1} \neq 0$, however, if $\tilde{e}^{n-1} = 0$, then there must be a class above it in higher filtration which detects e^{n-1} on E_{∞} , since $e^{n-1} \neq 0$. We see that on E_{∞} there are elements in filtration degree $\geq n-1$, hence $\exp_{\mathcal{A}\ell_{\ker f}} R \geq n$. \square

3.2.12 Lemma *For a G -space X , $H \subset G$ a subgroup, the composite of maps of Borel cohomology rings*

$$H_G^*(X; \mathbf{F}_p) \xrightarrow{\text{Res}_H^G} H_H^*(X; \mathbf{F}_p) \xrightarrow{\text{Ind}_H^G} H_G^*(X; \mathbf{F}_p)$$

is multiplication by $[G : H]$.

PROOF This follows from [Bor60, Prop. III.2.4] (note that the induction morphism also goes by the name of transfer morphism in this context, and that the morphism constructed in the proof of [Bor60, Prop. III.2.4] is the transfer morphism).

The result also follows by considering directly what happens on the cochain level. \square

3.2.13 Corollary *Let G be a group and $P \subset G$ a p -Sylow. Then the maps of G -spectra*

$$(3.2) \quad \underline{H\mathbf{F}}_{p_G} \rightarrow G/P_+ \wedge \underline{H\mathbf{F}}_{p_G} \rightarrow \underline{H\mathbf{F}}_{p_G}$$

representing the natural transformations Res_P^G and Ind_P^G exhibit $\underline{H\mathbf{F}}_{p_G}$ as a retract of $G/P_+ \wedge \underline{H\mathbf{F}}_{p_G}$.

PROOF We need to show that for all subgroups $G' \subset G$, applying $\pi_*^{G'}$ to (3.2) yields an isomorphism. This composite is given precisely by Lemma 3.2.12 for $X = G/G'$ and $H = P$. But multiplication by $[G : P]$ is an isomorphism because $p \nmid [G : P]$, whence $[G : P] \in \mathbf{F}_p^\times$. \square

3.2.14 Lemma *Let G be a group, P a p -Sylow subgroup of G . Then*

$$\exp_{\mathcal{E}(p)(G)} \underline{HF}_{p_G} = \exp_{\mathcal{E}(p)(P)} \underline{HF}_{p_P}.$$

PROOF First,

$$(3.3) \quad \text{Res}_P^G \underline{HF}_{p_G} = \underline{HF}_{p_P},$$

$$(3.4) \quad \text{Res}_P^G E_{\mathcal{E}(p)}(G) = E_{\mathcal{E}(p)}(P).$$

Hence by Corollary 3.2.5 (cf. [MNN17, Cor. 4.13]),

$$\exp_{\mathcal{E}(p)(P)} \underline{HF}_{p_P} \leq \exp_{\mathcal{E}(p)(G)} \underline{HF}_{p_G}.$$

For the upper bound, write $n = \exp_{\mathcal{E}(p)(P)} \underline{HF}_{p_P}$. We then have that

$$(3.5) \quad \text{sk}_{n-1} E_{\mathcal{E}(p)}(P) \wedge \underline{HF}_{p_P} \rightarrow \underline{HF}_{p_P}$$

admits a section. Note that for every $X \in \text{Sp}_G$, we have $\text{Ind}_H^G \text{Res}_H^G X = G/H_+ \wedge X$. Furthermore we have that, if we use the model for $E_{\mathcal{E}(p)}(P)$ from (3.4),

$$\text{sk}_{n-1} E_{\mathcal{E}(p)}(P) = \text{Res}_P^G \text{sk}_{n-1} E_{\mathcal{E}(p)}(G),$$

Applying this, the fact that Res_P^G is monoidal, and (3.3) to (3.5) yields a section of

$$G/P_+ \wedge \text{sk}_{n-1} E_{\mathcal{E}(p)}(G) \wedge \underline{HF}_{p_G} \rightarrow G/P_+ \wedge \underline{HF}_{p_G}.$$

Hence

$$\exp_{\mathcal{E}(p)(G)} G/P_+ \wedge \underline{HF}_{p_G} \leq n.$$

Applying Corollary 3.2.13 and Proposition 2.5.16 yields \square

$$\exp_{\mathcal{E}(p)} \underline{HF}_{p_G} \leq n.$$

3.2.15 Lemma *Let \mathcal{F} and \mathcal{G} be families of subgroups, and let M be both \mathcal{F} - and \mathcal{G} -nilpotent. Then for every $K \in \mathcal{F}$, $\text{Res}_K^G M$ is \mathcal{G}_K -nilpotent by Lemma 3.2.4. Write $n = \exp_{\mathcal{F}} M$, $m_K = \exp_{\mathcal{G}_K} \text{Res}_K^G M$ for all $K \in \mathcal{F}$, and $m = \max_K m_K$. Then*

$$\exp_{\mathcal{G}} M \leq mn.$$

PROOF The \mathcal{G}_K -nilpotence of $\text{Res}_K^G M$ implies that, for all $K \in \mathcal{F}$,

$$\text{sk}_{m_K-1} E\mathcal{G}_K \wedge \text{Res}_K^G M \rightarrow \text{Res}_K^G M$$

admits a section. By inducing up to G , we see that $\exp_{\mathcal{G}}(G/K_+ \wedge M) \leq m_K$. By taking the coproduct over all $K \in \mathcal{F}$, we get that $\exp_{\mathcal{G}} \text{sk}_0 E\mathcal{F} \wedge M \leq m$. We now argue by induction that

$$(3.6) \quad \exp_{\mathcal{G}} \text{sk}_{d-1} E\mathcal{F} \wedge M \leq md,$$

for all $d \geq 1$, of which the base case $d = 1$ we have just established.

Hence assume (3.6) has been established for some $d \geq 1$, and consider the cofiber sequence

$$\text{sk}_{d-1} E\mathcal{F} \wedge M \rightarrow \text{sk}_d E\mathcal{F} \wedge M \rightarrow \bigvee (S^d \wedge G/K_+) \wedge M$$

that ends in a wedge of spheres with isotropy in \mathcal{F} smashed with M . By induction, the \mathcal{G} -exponent of the left term is $\leq md$, and the \mathcal{G} -exponent of the right hand side is $\leq m$ because we already saw that $\exp_{\mathcal{G}}(G/K_+ \wedge M) \leq m$ for all $K \in \mathcal{F}$. Hence by [MNN17, Prop. 4.9.2] the \mathcal{G} -exponent of the middle term is $\leq m(d+1)$, which completes the induction.

We have by \mathcal{F} -nilpotency of M that M is a retract of $\text{sk}_{n-1} E\mathcal{F} \wedge M$, hence taking $d = n$ in (3.6) yields the result. \square

3.3 From representations to exponents via the projective bundle theorem

In [MNN15, Ex. 5.16] an upper bound on the $\mathcal{E}_{(2)}$ -exponent of $\underline{H}\mathbf{F}_2_{Q_8}$ is determined using the projective bundle theorem. We will also repeatedly use this technique to give upper bounds on the \mathcal{F} -exponent of $\underline{H}\mathbf{F}_2_G$ for various 2-groups G and families $\mathcal{F} \supset \mathcal{E}_{(2)}$. Therefore in this section we will describe this argument in some detail. We will need the following classical notion:

3.3.1 Definition Let V be a real or complex vector space. Then the *projectivation* $P(V)$ of V is the space of all lines in V .

3.3.2 Remark Note that if V comes equipped with a linear G -action (i.e., is a G -representation), then $P(V)$ inherits a natural G -action.

The goal of this section is to prove:

3.3.3 Proposition *Let G be a finite group, $n \geq 0$ a natural number and suppose G has a real n -dimensional representation V such that the projectivation $P(V)$ has isotropy groups contained in some family \mathcal{F} . That is, for every real line $L \subset V$ we assume that the isotropy group $G_L \leq G$ of elements of G fixing L satisfies $G_L \in \mathcal{F}$. Then $\underline{H}\mathbf{F}_2$ is \mathcal{F} -nilpotent and the exponent satisfies $\exp_{\mathcal{F}} \underline{H}\mathbf{F}_2 \leq n$.*

3.3.4 Proposition *Let G be a finite group, $n \geq 0$ a natural number and suppose G has a complex n -dimensional representation V such that the projectivation $P(V)$ has isotropy groups contained in some family \mathcal{F} . That is, for every complex line $L \subset V$, we assume that the isotropy group $G_L \leq G$ of elements of G fixing L satisfies $G_L \in \mathcal{F}$. Then $\underline{H}\mathbf{Z}$ is \mathcal{F} -nilpotent, and the exponent satisfies $\exp_{\mathcal{F}} \underline{H}\mathbf{Z} \leq 2n - 1$.*

3.3.1 The projective bundle theorem

To prove Proposition 3.3.3, we will use the projective bundle theorem, which can be used to develop the theory of Stiefel-Whitney classes and Chern classes. This is carried out for instance in [Hus94, Ch. 17]. We will follow the treatment and notation of [Hus94, §17.2], but only discussing what we need. For an overview of all bundles and their relations that will appear we refer to diagram (3.7) below, where the dashed arrows indicate pullbacks.

As in [Hus94, Ch. 17], we consider real and complex vector bundles at the same time. For the case of real vector bundles, we write $c = 1$, we consider cohomology with coefficients in $K_c = \mathbf{Z}/2$, and we let F be the field \mathbf{R} of real numbers. For the case of complex vector bundles, we write $c = 2$, we consider cohomology with coefficients in $K_c = \mathbf{Z}$, and we let F be the field \mathbf{C} of complex numbers.

We will write $E(\eta)$ resp. $B(\eta)$ for the total, resp. base space, of a fiber (not necessarily vector) bundle η . Let $\xi: E \xrightarrow{p} B$ be an n -dimensional vector bundle. Let E_0 be the non-zero vectors in E . Let E' be the quotient of E_0 where we identify non-zero vectors in the same line. This yields a factorization

$$\begin{array}{ccc} E_0 & \xrightarrow{\quad} & B \\ & \searrow & \nearrow q \\ & E' & \end{array},$$

and $E' \xrightarrow{q} B$ is a fibre bundle with fibre FP^{n-1} , called the projectivation of ξ and denoted $P(\xi)$.

For every point $b \in B$ in the base, the inclusion of the fibre $F^n \rightarrow p^{-1}(b) \subset E$ defines a natural inclusion $j_b: FP^{n-1} \rightarrow q^{-1}(b) \subset E(P(\xi))$. Pulling back the bundle ξ along q yields a bundle $q^*(\xi): E(q^*(\xi)) \rightarrow E(P(\xi))$, which admits the canonical line bundle λ_ξ as a subbundle, where a point in the total space $E(\lambda_\xi)$ is a pair (L, x) where L and x lie over the same base point and $x \in L$.

3.3.5 Proposition (see, e.g., [Hus94, Prop. 17.2.]) *Let*

$$j_b: FP^{n-1} \rightarrow E(P(\xi))$$

be the inclusion of the fibre. Then $j_b^(\lambda_\xi)$ is isomorphic to the tautological line bundle on FP^{n-1} .*

All the bundles that appear and their relations are as follows, with dashed lines indicating pullbacks:

$$(3.7) \quad \begin{array}{ccccc} E(j_b^*(\lambda_\xi)) & \dashrightarrow & E(\lambda_\xi) & & \\ & & \downarrow & & \\ & & E(q^*(\xi)) & \dashrightarrow & E \\ & & \downarrow & \nearrow E_0 & \downarrow p \\ & & & & B \\ FP^{n-1} & \xrightarrow{j_b} & E(P(\xi)) & \xrightarrow{q} & B \end{array}$$

Let $f: E(\mathbb{P}(\xi)) \rightarrow FP^\infty$ be a classifying map for the line bundle λ_ξ , i.e. $f^*(\gamma) \cong \lambda_\xi$, where γ is the tautological line bundle on FP^∞ . Let $z \in H^c(FP^\infty; K_c)$ be the polynomial generator, and let $a_\xi = f^*(z)$. Since f is unique up to homotopy, a_ξ is well defined. We then have the following Theorem:

3.3.6 Theorem (Projective bundle theorem, see [Hus94, Thm. 17.2.5])
 For an n -dimensional vector bundle ξ , the classes $1, a_\xi, \dots, a_\xi^{n-1}$ form a basis of the free $H^*(B(\xi); K_c)$ -module $H^*(E(\mathbb{P}(\xi)); K_c)$. Moreover,

$$q^*: H^*(B(\xi); K_c) \rightarrow H^*(E(\mathbb{P}(\xi)); K_c)$$

is a monomorphism.

3.3.2 Free R -modules

In this subsection we discuss some properties of free R -modules. The statements and notions are not new, but collected here for convenience.

3.3.7 Definition Let R_* be a graded ring, M_* a graded R_* -module. We say that M_* is a **free** R_* -module if there are integers n_i and an isomorphism of R_* -modules

$$g: \bigoplus_i \Sigma^{n_i} R_* \xrightarrow{\cong} M_*.$$

Here we write $(\Sigma N)_* = N_{*+1}$ for the suspension of graded abelian groups.

Likewise, we have:

3.3.8 Definition Let R be a (possibly equivariant) ring spectrum, and let M be an R -module. Then M is a **free** R -module if there are integers n_i and a weak equivalence of R -modules

$$g: \bigvee_I \Sigma^{n_i} R \xrightarrow{\simeq} M.$$

This weak equivalence is understood to be an equivariant weak equivalence in the equivariant case.

A map $f: R \rightarrow T$ of homotopy ring spectra makes T into an R -module. The next proposition says that one can check freeness of T as an R -module on the level of homotopy groups. The next proposition considers the non-equivariant case, the equivariant case is treated later.

3.3.9 Proposition Let $f: R \rightarrow T$ be a map of non-equivariant homotopy ring spectra. Then the following are equivalent:

- (1) The map f makes T into a free R -module.
- (2) The map $\pi_* f$ makes $\pi_* T$ into a free $\pi_* R$ -module.
- (3) There are integers n_i and homotopy classes of maps $\alpha_i: S^{n_i} \rightarrow T$ such that the α_i form a basis of $\pi_* T$ as a $\pi_* R$ -module.

PROOF (1) \Rightarrow (2) This follows from applying π_* to the equivalence $\bigvee_i \Sigma^{n_i} R \simeq T$ of R -modules.

(2) \Rightarrow (3) This follows from applying $\pi_* f$ to the canonical basis of $\bigoplus_i \Sigma^{n_i} \pi_* R$.

(3) \Rightarrow (1) The α_i define an isomorphism

$$\bigoplus_i \pi_* \Sigma^{n_i} R \rightarrow \pi_* T.$$

The result follows by lifting this isomorphism to an equivalence of spectra, which one can do as follows. For every i , the composite

$$R \wedge S^{n_i} \xrightarrow{f \wedge \alpha_i} T \wedge T \xrightarrow{\mu_T} T$$

is a map of R -modules. Taking the coproduct over i yields a map of R -modules

$$\bigvee_i \Sigma^{n_i} R \rightarrow T$$

which is the desired lift. \square

Combining this Proposition with Theorem 3.3.6 yields the following Corollary:

3.3.10 Corollary *Let the notation be as in Theorem 3.3.6, and let $H\mathbf{F}_2$ be the Eilenberg-MacLane spectrum representing cohomology with \mathbf{F}_2 -coefficients. Then the map $F(B(\xi)_+, H\mathbf{F}_2) \rightarrow F(E(P(\xi))_+, H\mathbf{F}_2)$ induced by q makes the mapping spectrum $F(E(P(\xi))_+, H\mathbf{F}_2)$ a free $F(B(\xi)_+, H\mathbf{F}_2)$ -module.*

PROOF According to Theorem 3.3.6, q_* makes $H^*(E(P(\xi)), \mathbf{F}_2)$ into a free $H^*(B(\xi), \mathbf{F}_2)$ -module, which precisely means that the map

$$\pi_* F(B(\xi)_+, H\mathbf{F}_2) \rightarrow \pi_* F(E(P(\xi))_+, H\mathbf{F}_2)$$

induced by q makes $\pi_* F(E(P(\xi))_+, H\mathbf{F}_2)$ into a free $\pi_* F(B(\xi)_+, H\mathbf{F}_2)$ -module. Applying Proposition 3.3.9 yields the desired result. \square

3.3.11 Corollary *Some suspension of the mapping spectrum $F(B(\xi), H\mathbf{F}_2)$ is a retract of the mapping spectrum $F(E(P(\xi)), H\mathbf{F}_2)$.*

PROOF Every free R -module admits some suspension of R as a retract. \square

Similarly, equivariantly we have

3.3.12 Proposition *Let $R \rightarrow T$ be a map of homotopy G -ring spectra. Then the following are equivalent:*

- (1) *The map f makes T into a free equivariant R -module,*
- (2) *There are homotopy classes of maps $\alpha_i: S^{n_i} \rightarrow T$ such that for every subgroup $H \leq G$, the canonical map $\bigoplus_i \pi_*^H \Sigma^{n_i} R \rightarrow \pi_*^H T, x \mapsto x \cdot \text{Res}_H^G \alpha_i$ is an isomorphism of $\pi_*^H R$ -modules.*

PROOF (1) \Rightarrow (2) follows from moving the canonical basis of $\bigoplus_i \pi_*^G \Sigma^{n_i} R$ through the equivalence of R^G -modules $\bigvee_i \Sigma^{n_i} R^G \simeq T^G$.

(2) \Rightarrow (1) follows from taking the coproduct over i of the composite

$$R \wedge S^{n_i} \rightarrow T \wedge T \rightarrow T$$

to get a map

$$\bigvee_i \Sigma^{n_i} R \rightarrow T.$$

The assumption on the restrictions says precisely that this is an equivariant equivalence. \square

3.3.3 From representations to exponents

In the proof of Proposition 3.3.3, we will take the projectivation of the Borel construction of a G -representation V . The proof will make use of the fact that the projectivation and the Borel construction commute, which is the content of the next lemma.

3.3.13 Lemma *Let V be a real or complex G -representation, and let H be a subgroup of G . Consider the Borel construction*

$$V \rightarrow V \times_H EG \rightarrow BH.$$

Then the projectivation of the Borel construction equals the Borel construction of the projectivation, that is,

$$P(V \times_H EG) = P(V) \times_H EG.$$

PROOF Write V_0 for the non-zero vectors in V , write P for the product $V \times EG$, and write $P_0 = V_0 \times EG$. Then F^* acts on the left coordinate of P_0 by multiplication, and trivially on the right coordinate, and H acts coordinatewise on P_0 . Moreover, these actions commute, by linearity of the action of G (hence of H) on V . Therefore P_0 has in fact an action of $F^* \times H$. Dividing out the action of F^* first and then dividing out the action of H gives the Borel construction of the projectivation, dividing out the action of H first and then the action of F^* gives the projectivation of the Borel construction. Both equal the space obtained by dividing out $F^* \times H$, and they are therefore equal. \square

We are now ready to prove Proposition 3.3.3 and Proposition 3.3.4.

PROOF (OF PROPOSITION 3.3.3 AND PROPOSITION 3.3.4) Consider the Borel construction on V for an arbitrary subgroup $H \leq G$, and call the resulting bundle ξ_H :

$$\xi_H : V \rightarrow V \times_H EG \rightarrow BH.$$

Note that this is natural with respect to inclusions of subgroups. The associated projective bundle is

$$P(\xi_H) : P(V) \rightarrow P(V \times_H EG) \rightarrow BH.$$

Observe that $P(V \times_H EG) = P(V) \times_H EG$ by Lemma 3.3.13. Hence by Theorem 3.3.6,

$$F((P(V) \times_H EG)_+, HK_c) \cong F(P(V)_+, \underline{HK}_c)^H$$

is a free $F(BH, K_c) = \underline{HK}_c^H$ -module. Recall that the *real* dimension of V is cn . A basis is given by the proof of [Hus94, Prop. 17.3.3], which shows that there are classes $1, a_{\xi_H}, \dots, a_{\xi_H}^{n-1}$ that form a basis of $\pi_*^H F(P(V), \underline{HK}_c)$ as a $\pi_*^H \underline{HK}_c$ -module. Moreover, the element a_{ξ_H} is natural with respect to restriction to subgroups. It follows that all basis elements, being powers of a_{ξ_H} , are natural with respect to restriction to subgroups. In particular,

$$\text{Res}_H^G a_{\xi_G} = a_{\xi_H}$$

for all $H \leq G$. Hence by Proposition 3.3.12, $F(P(V), \underline{HK}_c)$ is a free \underline{HK}_c -module. Hence a suspension of \underline{HK}_c is a retract of $F(P(V), \underline{HK}_c)$. Therefore by [MNN17, Prop. 4.9] and Corollary 2.5.12,

$$\exp_{\mathcal{F}} \underline{HK}_c \leq \exp_{\mathcal{F}} F(P(V), \underline{HK}_c),$$

but V is cn -dimensional, therefore $P(V)$ admits the structure of an $(cn - c)$ -dimensional G -CW-complex ([Ill83, Cor. 7.2]) with isotropy (by assumption) contained in \mathcal{F} , therefore by [MNN15, Prop. 2.26] the exponent of the right hand side is

$$\leq cn - c + 1.$$

In the case of a real representation we have $c = 1$ and this equals n , in the case of a complex representation we have $c = 2$ and this equals $2n - 1$. \square

3.4 A bound on the exponent for the family of proper subgroups

Let G be a finite non-abelian 2-group of size 2^k . The goal of this section is to prove

$$\exp_{\mathcal{F}} \underline{HF}_{2G} \leq 2 \lfloor \sqrt{|G| - k} \rfloor - 1,$$

see Corollary 3.4.2 below. The argument is an adaption of the ones found in [PY03, Lem. 4.3] and [Sym91].

3.4.1 Lemma *Let G be a finite non-abelian p -group of size p^k . Then G has an irreducible complex representation V with $\dim_{\mathbf{C}} V \geq 2$, and moreover all such V satisfy*

$$\dim_{\mathbf{C}} V \leq \lfloor \sqrt{|G| - 1} \rfloor.$$

PROOF Denote by n_1, \dots, n_h the \mathbf{C} -dimensions of the irreducible \mathbf{C} -representations of G . We have

$$n_1^2 + n_2^2 + \dots + n_h^2 = |G|,$$

(see, e.g., [Ser77, Cor. 2.4.2]). Since G is non-abelian, some $n_i \geq 2$ ([Ser77, Thm. 3.1.9]). Assume without loss of generality that it is n_1 . Then

$$\begin{aligned} |G| &= n_1^2 + n_2^2 + \cdots + n_h^2 \\ &\geq n_1^2 + 1, \end{aligned}$$

because there is also a trivial representation, hence

$$n_1 \leq \sqrt{|G| - 1}.$$

Flooring this preserves the inequality and does not change the integer on the left hand side. This yields the result. \square

3.4.2 Corollary *Let G be a finite non-abelian 2-group of size 2^k , and let \mathcal{P} be the family of proper subgroups of G . Then*

$$\exp_{\mathcal{P}} \underline{HF}_{2G} \leq 2 \lfloor \sqrt{|G| - 1} \rfloor - 1.$$

PROOF Let V be an irreducible complex representation of G satisfying

$$2 \leq \dim_{\mathbf{C}} V \leq \lfloor \sqrt{|G| - 1} \rfloor,$$

as furnished by Lemma 3.4.1. Then the complex projectivation $P(V)$ has isotropy in \mathcal{P} the family of proper subgroups of G , for if $L \in P(V)$ were fixed by all of G , V would not be irreducible, since $\dim_{\mathbf{C}} V \geq 2$. An application of Proposition 3.3.4 yields the result. \square

Chapter 4

Mod-2 group cohomology computations of 2-groups using the \mathcal{F} -homotopy limit spectral sequence

4.1 Introduction

In this chapter we apply the \mathcal{F} -homotopy limit spectral sequence to the case $X = \text{pt}$, $\mathcal{F} \supset \mathcal{E}_{(2)}$ and G all finite 2-groups of order ≤ 16 , to compute group cohomology. For the indexing of the \mathcal{F} -homotopy limit spectral sequence, we use the following non-standard convention:

4.1.1 Convention For a 2-group G and family $\mathcal{F} \supset \mathcal{E}_{(2)}$, the \mathcal{F} -homotopy limit spectral sequence is given by

$$E_2^{s,t} = \lim_{(-) \in \mathcal{O}(G)_{\mathcal{F}}^{\text{op}}}^s H^t(B(-); \mathbf{F}_2) \Rightarrow H^{s+t}(BG; \mathbf{F}_2).$$

We use Adams' grading convention for displaying the spectral sequences, which means that we put $s + t$ on the horizontal axis and s on the vertical axis. This yields a spectral sequence drawn in the first quadrant, with d_n going 1 over to the right and n up.

The group cohomology rings that we study in this chapter are well known. A classical technique for computing group cohomology is to use the Lyndon-Hochschild-Serre spectral sequence, as was for example done by Quillen to compute the group cohomology of the extra-special 2-groups ([Qui71b]). Rusin computed the cohomology of 2-groups up to order 32 in [Rus89] using the Eilenberg-Moore spectral sequence. Using large initial segments of projective resolutions, and a completion criterion of Benson ([Ben04]), Carlson computed the group cohomology of all 2-groups up to order 64 ([CTVEZ03]) using a computer. Green and King improved these methods by deriving a variant of Benson's completion criterion that allows for earlier detection of completion ([GK11, Thm. 3.3]) and

using this computed the cohomology of all 2-groups up to order 64 ([GK11] and [GK15]).

The \mathcal{F} -homotopy limit spectral sequence provides a fourth method for computing group cohomology. The advantage of this spectral sequence is that, if one takes $\mathcal{F} \supset \mathcal{E}_{(2)}$, there will be a horizontal vanishing line at a finite page. Hence the computation is a finite one, and moreover the knowledge that a vanishing line must eventually appear can also help determine differentials. However, these two advantages come with a trade-off compared to using the Lyndon-Hochschild-Serre or Eilenberg-Moore spectral sequence: compared to the latter two, it is in general rather difficult to compute the E_2 -page of the \mathcal{F} -homotopy limit spectral sequence. Moreover, the product structure on the E_2 -page of the \mathcal{F} -homotopy limit spectral sequence is rather mysterious, which makes propagating differentials difficult. But the trade-off is at least good enough to still be able to compute the group cohomology of all the 2-groups up to order 16, at least additively, as shown in this chapter.

4.2 The 2-groups of order at most 16

The classification of 2-groups up to order 16 has been long known (see, e.g., [Bur55, §118]). In this section we recall this classification, and provide forward references to the bounds on the exponents of \underline{HF}_{2G} for all these groups.

Each classification subsection is preceded by one or two subsections defining families of 2-groups needed in the classification.

4.2.1 Abelian 2-groups

For all finite abelian 2-groups A , we compute the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BA; \mathbf{F}_2)$ in Section 4.3. Additionally, we obtain the following exponent:

Proposition (see Proposition 4.3.1) *Let A be a finite abelian 2-group, and let n be the minimal number of generators of the subgroup of 2-divisible elements of A , i.e., of the image of the multiplication-by-2 map $A \xrightarrow{\cdot 2} A$. (We consider the trivial group to be generated by 0 elements.) Then*

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2A} = n + 1.$$

Because all 2-groups up to order 4 are abelian, Proposition 4.3.1 in particular determines all the $\mathcal{E}_{(2)}$ -exponents of all 2-groups up to order 4.

4.2.2 The dihedral groups

Denote by $R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ the rotation by θ -matrix, and by $S = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ the reflection-in-the- x -axis matrix. Then $R_{2\pi/2^{n-1}}$ and S generate a subgroup of order 2^n of $O(2)$, called the dihedral group of order 2^n , and denoted D_{2^n} . It is the automorphism group of the regular 2^{n-1} -gon centered at the origin and with two of its sides bisected by the x -axis.

The map given by $R_{2\pi/2^{n-1}} \mapsto \rho$ and $S \mapsto \sigma$ gives an isomorphism of D_{2^n} with the finitely presented group

$$\langle \sigma, \rho \mid \sigma^2 = \rho^{2^{n-1}} = e, \sigma\rho\sigma^{-1} = \rho^{-1} \rangle.$$

4.2.3 The quaternion groups

The generalized quaternion group Q_{2^n} of order 2^n is the finite subgroup of quaternionic space \mathbf{H} generated multiplicatively by the elements of unit length $\{e^{2\pi i/2^{n-1}}, j\}$ (see, e.g., [CE56, XII.§7]). Denoting these generators by r and s , respectively, one gets the presentation

$$Q_{2^n} = \langle r, s \mid r^{2^{n-2}} = s^2, rsr = s \rangle.$$

4.2.4 The groups of order 8

Now that we have defined the families of dihedral and quaternion groups, we are able to state the classification of the groups of order 8.

4.2.1 Theorem (see, e.g., [Arm88, Thm. 13.3]) *There are, up to isomorphism, exactly 5 groups of order 8. They are given by $C_2^{\times 3}$, $C_2 \times C_4$, C_8 , D_8 and Q_8 .*

The $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BD_8; \mathbf{F}_2)$ is computed in Section 4.4, and $\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2D_{2^n}}$ is determined.

The $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BQ_{2^n}; \mathbf{F}_2)$ was already computed for $n = 3$ in [MNN15, Ex. 5.18], as well as $\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2Q_{2^n}}$ in that case. We give a straightforward generalization for both computations to the case $n > 3$ in Section 4.5. In summary, the exponents of the groups of order 8 are then given by

G	$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2G}$	Reference
$C_2^{\times 3}$	1	Proposition 4.3.1
$C_2 \times C_4$	2	Proposition 4.3.1
C_8	2	Proposition 4.3.1
D_8	2	Proposition 4.4.1
Q_8	4	[MNN15, Ex. 5.18]

Table 4.1: The $\mathcal{E}_{(2)}$ -exponents for the groups of order 8.

4.2.5 The groups of order 16

We state the classification of groups of order 16, following the treatment in [Wil05]. We first need to introduce some notation as used in [Wil05].

4.2.2 Notation Given groups N and H and an homomorphism $\phi: H \rightarrow \text{Aut}(N)$, we denote the semidirect product of N by H with respect to ϕ by $N \rtimes_{\phi} H$.

In the special case $H = C_2$, $N = C_8 = \langle x \rangle$, the possible actions of C_2 on C_8 are given by $x \mapsto x^n$ with n odd, and for all such actions we denote the induced semidirect product by $C_8 \rtimes^n C_2$. In particular, $C_8 \rtimes^1 C_2 \cong C_8 \times C_2$ and

$C_8 \rtimes^7 C_2 \cong D_{16}$. These semidirect products for $n = 3, 5$ also have names: $C_8 \rtimes^3 C_2$ is also known as the semidihedral group of order 16, denoted SD_{16} . The group $C_8 \rtimes^5 C_2$ is also known as the modular group of order 16, denoted M_{16} .

4.2.3 Notation If N and H are groups such that there is an up to isomorphism unique non-trivial semi-direct product of N by H , we denote it by $N \rtimes H$.

4.2.4 Definition Let $C_4 = \langle x \rangle$, and $C_2 = \langle y \rangle$. Following [Wil05, Fact 4], we denote by ψ_5, ψ_6 the elements of $\text{Aut}(C_4 \times C_2)$ given by $\psi_5(x) = xy, \psi_5(y) = y, \psi_6(x) = x^3$ and $\psi_6(y) = x^2y$.

We are now able to list all groups of order 16, and the bounds on the exponents that we obtained.

4.2.5 Theorem *The following table lists in its first column all groups of order 16, see, e.g., [Wil05, Thm. 2]. The second column either gives the $\mathcal{E}_{(2)}$ -exponent or an interval in which the exponent lies. The third column gives a forward reference for the proof of the contents of the second column. The fourth column gives a forward reference to the computation of an \mathcal{F} -homotopy limit spectral sequence with $\mathcal{F} \supset \mathcal{E}_{(2)}$ converging to the group cohomology of the first column.*

G	$\exp_{\mathcal{E}_{(2)}} \underline{HF}_2 G$	Reference	holim SS
$C_2^{\times 4}$	1	Proposition 4.3.1	Section 4.3
$C_2^{\times 2} \times C_4$	2	Proposition 4.3.1	Section 4.3
$C_4 \times C_4$	3	Proposition 4.3.1	Section 4.3
$C_8 \times C_2$	2	Proposition 4.3.1	Section 4.3
C_{16}	2	Proposition 4.3.1	Section 4.3
D_{16}	2	Proposition 4.4.1	Section 4.4
Q_{16}	4	Proposition 4.5.4	Section 4.5
$SD_{16} = C_8 \rtimes^3 C_2$	[3, 4]	Proposition 4.6.2	Section 4.6
$M_{16} = C_8 \rtimes^5 C_2$	[3, 4]	Proposition 4.7.4	Section 4.7
$D_8 * C_4 = (C_4 \times C_2) \rtimes^{\psi_6} C_2$	4	Proposition 4.8.22	Section 4.8
$C_4 \rtimes C_4$	[3, 4]	Proposition 4.9.6	Section 4.9
$(C_4 \times C_2) \rtimes^{\psi_5} C_2$	2	Proposition 4.10.1	Section 4.10
$Q_8 \times C_2$	4	Proposition 4.11.1	Section 4.11
$D_8 \times C_2$	2	Proposition 4.12.2	Section 4.12

Table 4.2: Summary of results for groups of order 16.

4.3 Abelian 2-groups

In this section we compute the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence for finite 2-groups. We first recall some basic results on the cohomology rings of cyclic 2-groups and the LHSSS's computing these. We will omit the \mathbf{F}_2 -coefficients from the notation for cohomology groups in this section.

4.3.1 Cyclic 2-groups

Let C_{2^n} the cyclic group of order 2^n , generated by t . Denote by $\delta_t \in H^1(BC_{2^n})$ the cohomology class dual to t , where C_{2^n} necessarily acts trivially on \mathbf{F}_2 since \mathbf{F}_2 has only the identity as automorphism. Denote by $\beta_n(-)$ the n -th order Bockstein. Then $\beta_n \delta_t \in H^2(BC_{2^n})$ is the cohomology class corresponding to the unique non-trivial group extension of C_{2^n} by C_2 , and furthermore (see, e.g., [Eve91, sec. 3.2]),

$$H^*(BC_{2^n}) \cong \mathbf{F}_2[\delta_t]/(\delta_t^2) \otimes \mathbf{F}_2[\beta_n \delta_t],$$

with, for $n = 1$, the multiplicative extension $\beta_1 \delta_t = \text{Sq}^1 \delta_t = \delta_t^2$.

The group C_{2^n} has a unique maximal elementary abelian subgroup, which is generated by $t^{2^{n-1}}$. By Proposition 2.6.1, the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence for the group C_{2^n} reduces to the LHSSS of the group extension

$$C_2 \rightarrow C_{2^n} \rightarrow C_{2^{n-1}}.$$

For $n = 1$, the right hand side group is the trivial group, and therefore the LHSSS collapses to the $s = 0$ -line at E_2 .

For $n > 1$, we write $a = \delta_{\bar{t}}$, $\beta_{n-1} a = \beta_{n-1} \delta_{\bar{t}}$ the $(n-1)$ -th order Bockstein on a , and $x = \delta_{t^{2^{n-1}}}$. Because the the group C_{2^n} is abelian, the local coefficient system is trivial, and the E_2 -page is isomorphic to

$$E_2^{*,*} \cong \mathbf{F}_2[a, \beta_{n-1} a, x]/(a^2),$$

with, for $n = 2$, the multiplicative extension $\beta_1 a = a^2$, and with (s, t) -degrees given by $|a| = (1, 0)$, $|\beta_{n-1} a| = (2, 0)$ and $|x| = (0, 1)$.

Because the \mathbf{F}_2 -dimension of $\text{Hom}(C_{2^n}, \mathbf{F}_2)$ is 1, the dimension of $H^1(BC_{2^n})$ is 1. But E_2 has 2 classes in the $s+t = 1$ -stem, so one of them has to die, which can only be x for degree reasons, which can only possibly hit $\beta_{n-1} a$ with a d_2 . The class a is a permanent cycle for degree reasons, and hence

$$E_3 \cong \mathbf{F}_2[a, [x^2]]/(a^2).$$

This has a vanishing line of height 2, and therefore the spectral sequence collapses at this stage.

4.3.2 Abelian 2-groups

Let A be a finite abelian 2-group. Then A is isomorphic to

$$\prod_{j \in J} C_{2^{n_j}} \times \prod_{k \in K} C_2$$

with $n_j \geq 2$, for some indexing sets J and K . We compute the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BA)$. By the Künneth isomorphism, this ring is isomorphic to $\mathbf{F}_2[x_j, \beta_{n_j} x_j, x_k]/(x_j^2)$, with degrees $|x_j| = |x_k| = 1$, and $|\beta_{n_j-1}(x_j)| = 2$.

Since A has a unique maximal elementary abelian 2-subgroup $\prod_{j \in J} C_2 \times \prod_{k \in K} C_2$, this subgroup is normal and hence by Proposition 2.6.1 this reduces

the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence to the LHSSS of the extension

$$(4.1) \quad \prod_{j \in J} C_2 \times \prod_{k \in K} C_2 \rightarrow A \rightarrow \prod_{j \in J} C_{2^{n_j-1}}.$$

Write t_j for the generator of $C_{2^{n_j}}$, and t_k for the generator of C_2 indexed by k .

For all $j \in J$, the LHSSS corresponding to the extension

$$C_2 \rightarrow C_{2^{n_j}} \rightarrow C_{2^{n_j-1}}$$

has as $E_2 = \mathbf{F}_2[a_j, \beta_{n_j-1}a_j, x_j]/(a_j^2)$ with $a_j = \delta_{t_j}$, $\beta_{n_j-1}a_j = \beta_{n_j-1}\delta_{t_j}$, $x_i = \delta_{t_i^{2^{n_i-1}}}$, and for $n_j = 2$ with multiplicative extension $\beta_1 a_j = a_j^2$. The (s, t) -degrees are given by $|a_j| = (1, 0)$, $|\beta_{n_j-1}a_j| = (2, 0)$ and $|x_j| = (0, 1)$. Furthermore, $d_2 x_j = \beta_{n_j-1}a_j$ and

$$E_3 = E_\infty = \mathbf{F}_2[a_j, [x_j^2]]/(a_j^2).$$

Using the Künneth isomorphism, the E_2 -page of the LHSSS of (4.1) is given by

$$E_2^{s,t} = \mathbf{F}_2[a_j, \beta_{n_j-1}a_j, x_j, x_k]/(a_j^2 \mid j \in J)$$

with (s, t) -degrees as above, and for $n_j = 2$ with the multiplicative extensions $\beta_1 a_j = a_j^2$.

By naturality, the generators on E_2 supporting a d_2 are the x_j , and

$$d_2 x_j = \beta_{n_j-1}a_j,$$

for all j . Hence

$$E_3 = \mathbf{F}_2[a_j, [x_j^2], x_k]/(a_j^2 \mid j \in J).$$

By the Künneth theorem, the stems of E_3 have Betti numbers equal to the Betti numbers of the target, hence $E_3 = E_\infty$.

In particular, $\prod_{j \in J} a_j \in E_\infty^{\#J, 0}$ is a non-zero element in filtration degree $\#J$, which shows that

$$(4.2) \quad \exp_{\mathcal{E}_{(2)}} \underline{H\mathbf{F}}_2 \geq \#J + 1.$$

Conversely, for $j \in J$, consider the projection of A onto the C_2 in the j -th factor:

$$p_j: A \rightarrow C_2 \cong O(1).$$

The corresponding Euler class is $a_j \in H^1(BA; C_2)$. Since $a_j^2 = 0$, we get $\exp_{\mathcal{A}ll_{\ker p_j}} \underline{H\mathbf{F}}_2 = 2$. Now $\bigcap_{j \in J} \ker p_j = \mathcal{E}_{(2)}$, and hence by Lemma 3.2.6, $\exp_{\mathcal{E}_{(2)}} \underline{H\mathbf{F}}_2 \leq \#J + 1$. Combined with Equation (4.2) this shows that

4.3.1 Proposition *Let A be a group isomorphic to*

$$\prod_{j \in J} C_{2^{n_j}} \times \prod_{k \in K} C_2$$

with $n_j \geq 2$. Then

$$(4.3) \quad \exp_{\mathcal{E}(2)} \underline{H\mathbf{F}}_{2A} = \#J + 1.$$

4.3.2 Corollary *Let G be a group with subgroup isomorphic to $\prod_{j \in J} C_{2^{n_j}}$ with all $n_j \geq 2$. Then*

$$\exp_{\mathcal{E}(2)} \underline{H\mathbf{F}}_2 \geq \#J + 1.$$

PROOF This follows from Corollary 3.2.5 and Equation (4.3). \square

4.4 Dihedral groups

4.4.1 Introduction

Denote by D_{2^n} the dihedral group of order 2^n . We apply the $\mathcal{E}(2)$ -homotopy limit spectral sequence to compute the group cohomology of D_{2^n} . We first do computation for $D_{2^n} = D_8$, and in the last section we do the computation for $n \geq 4$.

The cohomology rings $H^*(BD_{2^n}; \mathbf{F}_2)$ are all isomorphic to

$$\mathbf{F}_2[x, y, w]/(xy)$$

with degrees $|x| = |y| = 1$, $|w| = 2$, see, e.g., [AM04, Thm. 2.7]. Since $\underline{H\mathbf{F}}_2$ is $\mathcal{E}(2)$ -nilpotent ([MNN15, Prop. 5.14]), the spectral sequence will admit a vanishing line, which is bounded by the exponent $\exp_{\mathcal{E}(2)} \underline{H\mathbf{F}}_2$. In fact, the computation will show that the spectral sequence has a vanishing line of height 1, which is equivalent to the fact that D_{2^n} has its cohomology detected on elementary abelian 2-subgroups, which is [Qui71a, Lem. 4.6].

Before proceeding with the calculation, we first identify the exponent of D_{2^n} -Borel equivariant \mathbf{F}_2 -cohomology for all n .

4.4.2 The exponent

In this section we will show the following.

4.4.1 Proposition

$$\exp_{\mathcal{E}(2)} \underline{H\mathbf{F}}_{2D_{2^n}} = \begin{cases} 2 & \text{for } n \equiv 0 \pmod{4}, \\ 1 & \text{otherwise.} \end{cases}$$

PROOF Write $n = 2^v m$ with $2 \nmid m$. Then $D_{2^{v+1}} \subset D_{2^n}$ is a 2-Sylow subgroup and by Lemma 3.2.14 we are reduced to showing that

$$\exp_{\mathcal{E}(2)} \underline{HF}_2 D_{2^k} \begin{cases} = 2 & \text{for } k \geq 3, \\ = 1 & \text{for } k = 2. \end{cases}$$

For the case $k \geq 3$, we have $C_{2^{k-1}} \subset D_{2^k}$, and hence by Corollary 4.3.2, $\exp_{\mathcal{E}(2)} \underline{HF}_2 \geq 2$. Combining this with the upper bound furnished by Corollary 4.4.3 below gives the claimed exponent.

For the case $k = 2$, we have $D_4 \cong C_2 \times C_2$, and hence by Proposition 2.5.6 we get the claimed exponent. \square

The upper bound

In this section we will show using Proposition 3.3.3 that for D_{2^n} the dihedral group of order 2^n we have

$$\exp_{\mathcal{E}(2)} \underline{HF}_2 D_{2^n} \leq 2,$$

see Corollary 4.4.3 below. To this end, consider the presentation

$$D_{2^n} = \langle \sigma, \rho \mid \sigma^2 = \rho^{2^{n-1}} = e, \sigma \rho \sigma^{-1} = \rho^{-1} \rangle.$$

For an angle θ , denote the matrix representing rotation in \mathbf{R}^2 by θ by

$$R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

Denote by T the matrix representing reflection in the x -axis:

$$T = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The group D_{2^n} is the subgroup of $O(2)$ fixing the regular 2^{n-1} -gon centered at the origin in \mathbf{R}^2 with one of its sides bisected in the right half-plane by the x -axis. The inclusion $D_{2^n} \rightarrow O(2)$ is on elements given by $\sigma \rightarrow T$, $\rho \rightarrow R_{2\pi/2^{n-1}}$. This yields a linear 2-dimension real representation of D_{2^n} , which we call V . We apply Proposition 3.3.3 to compute an upper bound on the exponent with respect to the minimal family \mathcal{F} containing the isotropy of $P(V)$. This minimal family is determined in the next lemma.

4.4.2 Lemma *The minimal family containing the isotropy groups of the projectivization $P(V)$ is $\mathcal{E}(2)$, the family of elementary abelian 2-groups in D_{2^n} .*

PROOF The proof is by elementary linear algebra. Since V is an orthogonal representation, a linear subspace L spanned by a vector v is fixed by an element g of D_{2^n} if and only if g has eigenvalue 1 or -1 (we say that g has eigenvalue λ if the matrix by which g acts on V has eigenvalue λ).

First consider the rotations in D_{2^n} , that is the elements of the form ρ^j . We note that R_θ has characteristic polynomial

$$(\cos(\theta) - \lambda)^2 + \sin^2 \theta = 0,$$

which for $\lambda = 1$ is 0 if and only if $\theta \equiv 0 \pmod{2\pi}$, and for $\lambda = -1$ is 0 if and only if $\theta = \pi \pmod{2\pi}$. Hence ρ^j fixes a line if and only if $2\pi j/2^{n-2} \equiv \pi \pmod{2\pi}$, that is if and only if $j \equiv 2^{n-2} \pmod{2^{n-1}}$. Hence $\rho^{2^{n-2}}$ is the only non-trivial rotation that fixes a line. Since $\rho^{2^{n-2}}$ acts via R_π , the elements $\rho^{2^{n-2}}$ fixes every line in V .

For the lines fixed by a reflection $\sigma\rho^j$, we first note that, writing $e_1 = (1 \ 0)^t$, TR_θ has eigenvectors $R_{-\theta/2}e_1$ and $R_{-\theta/2-\pi}e_1$, with eigenvalues 1 and -1 , respectively. Therefore these are, up to scalar, all eigenvectors of TR_θ , since V is 2-dimensional. Assume $\sigma\rho^j$ and $\sigma\rho^k$ fix a common line, spanned by a vector v . If v had the same eigenvalue for $\sigma\rho^j$ and $\sigma\rho^k$, we would have

$$\frac{-2\pi j}{2^{n-1}} \equiv \frac{-2\pi k}{2^{n-1}} \pmod{2\pi},$$

which is equivalent to $k \equiv j \pmod{2^{n-1}}$, in other words $\sigma\rho^j = \sigma\rho^k$.

Alternatively, if v has different eigenvalues for $\sigma\rho^j$ and $\sigma\rho^k$, then we have

$$\frac{-2\pi j}{2^{n-2}} \equiv \frac{-2\pi k}{2^{n-2}} - \pi \pmod{2\pi},$$

which is equivalent to $j - k \equiv 2^{n-2} \pmod{2^{n-1}}$. Hence if a line is fixed by a reflection $\sigma\rho^j$, it is fixed by exactly one more reflection, namely $\sigma\rho^{j+2^{n-2}}$. Note that

$$\sigma\rho^j\sigma\rho^{j+2^{n-2}} = \rho^{2^{n-2}} = \sigma\rho^{j+2^{n-2}}\sigma\rho^j. \quad \square$$

Therefore $\langle \sigma\rho^j, \sigma\rho^{j+2^{n-2}} \rangle$ is isomorphic to $C_2 \times C_2$.

We conclude that a line in V is fixed either by $\langle \rho^{2^{n-2}} \rangle$, or by $\langle \sigma\rho^j, \sigma\rho^{j+2^{n-2}} \rangle$ for exactly one $j \pmod{2^{n-2}}$. All these subgroups are elementary abelian, proving the result.

4.4.3 Corollary *The $\mathcal{E}_{(2)}$ -exponent of $\underline{H}\mathbf{F}_{2D_{2^n}}$ satisfies*

$$\exp_{\mathcal{E}_{(2)}} \underline{H}\mathbf{F}_{2D_{2^n}} \leq 2.$$

PROOF Immediate from Proposition 3.3.3 and Lemma 4.4.2 and the fact that V is 2-dimensional. \square

4.4.3 The spectral sequence

Summary of the computation

We will now compute the group cohomology ring $H^*(BD_8; \mathbf{F}_2)$ using the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BD_8; \mathbf{F}_2)$. Since $\mathcal{E}_{(2)}$ is the derived defect base of $\underline{H}\mathbf{F}_2$ ([MNN15, Prop. 5.16]), the spectral sequence will have a horizontal vanishing line at a finite page. In fact, by Proposition 4.4.1, the spectral sequence will have a horizontal vanishing line of height 2 on the E_3 -page. In particular, $E_3 = E_\infty$.

The computation, which follows the strategy of [MNN15, App. B], can be summarized as follows. The most work goes into computing the E_2 -term. We

find a homotopy final subcategory \mathcal{O}_f of $\mathcal{O}(D_8)_{\mathcal{E}(2)}$ (using Proposition 2.7.2). We then write a short exact sequence of coefficient systems ending in the constant coefficient system \underline{Z} on \mathcal{O}_f , which lifts to a long exact sequence of spectral sequences, constituting of two Lyndon-Hochschild-Serre spectral sequences and the $\mathcal{E}(2)$ -homotopy limit spectral sequence. We compute the E_2 -pages of the LHSSS's, which allows us to determine the E_2 -page of the $\mathcal{E}(2)$ -homotopy limit spectral sequence.

The vanishing line of height 2 on E_3 implies that the only non-trivial differential is a d_2 , which implies enough about the possible patterns of differentials on the E_2 -page to determine $E_3 = E_\infty$ (see Figure 4.4).

The orbit category

A generating set of morphism for the orbit category $\mathcal{O}(D_8)_{\mathcal{E}(2)}$ is given by




Figure 4.1: The orbit category $\mathcal{O}(D_8)_{\mathcal{E}(2)}$. The highlighted subcategory is homotopy final.

Write

$$\begin{aligned} Z &= Z(D_8) = \langle \rho^2 \rangle, \\ H_1 &= \langle \sigma, \sigma \rho^2 \rangle, \\ H_2 &= \langle \sigma \rho, \sigma \rho^3 \rangle. \end{aligned}$$

With these H_i and Z , we apply the decomposition of Section 2.7. Therefore we next compute the LHSSS's obtained from the extensions $H_i \rightarrow D_8 \rightarrow C_2$ and $Z \rightarrow D_8 \rightarrow C_2 \times C_2$.

The LHSSS of $H_i \rightarrow D_8 \rightarrow C_2$

We compute the E_2 -page of the LHSSS of the extension $H_1 \rightarrow D_8 \rightarrow C_2$, the case for H_2 follows from the automorphism $\sigma \leftrightarrow \sigma \rho$ of D_8 .

The group $C_2 = \langle \bar{\rho} \rangle$ acts on H_1 by

$$\begin{aligned} \rho(\sigma)\rho^{-1} &= \sigma\rho^2, \\ \rho(\sigma\rho)\rho^{-1} &= \sigma. \end{aligned}$$

Hence it acts on $H^*(H_1)$ by interchanging δ_σ and $\delta_{\sigma\rho^2}$. Write $u_{1,1} = \delta_\sigma + \delta_{\sigma\rho^2}$ and $u_{1,2} = \delta_\sigma\delta_{\sigma\rho^2}$ for the first and second elementary symmetric polynomial. It is well known that the invariants of a polynomial algebra of the action of the symmetric group on the generators is given by the polynomial algebra on the symmetric polynomials.

Taking the minimal periodic C_2 -resolution of \mathbf{Z} , applying the hom-functor of C_2 -modules $\text{Hom}_{C_2}(-, H^*(H_1))$, and taking cohomology, we see that the $s = 0$ -row of the LHSSS is given by the C_2 -invariants

$$\mathbf{F}_2[u_{1,1}, u_{2,1}].$$

Write $a_1 = \delta_{\bar{\rho}}$. The image of the differential in the long exact sequence computing $E_2^{s,*}$ maps a polynomial to its C_2 -symmetrization:

$$P(\delta_\sigma, \delta_{\sigma\rho^2}) \mapsto P(\delta_\sigma, \delta_{\sigma\rho^2}) + P(\delta_{\sigma\rho^2}, \delta_\sigma).$$

The image are the symmetrized polynomials, which is $\mathbf{F}_2[u_{1,1}, u_{2,1}]\{u_{1,1}\}$. The rows $E_2^{s,*}$ for $s > 1$ are then given by the C_2 -invariants modulo the symmetrized polynomials, which are

$$\mathbf{F}_2[u_{2,1}]\{a_1^s\}.$$

Assembling the rows together gives the E_2 -page

$$E_2 = \mathbf{F}_2[a_1, u_{1,1}, u_{2,1}]/(a_1u_{1,1}),$$

with (s, t) -degrees $|a_1| = (1, 0)$, $|u_{1,1}| = (0, 1)$ and $|u_{2,1}| = (0, 2)$.

For the extension $H_2 \rightarrow D_8 \rightarrow C_2$, we write $u_{1,2} = \delta_{\sigma\rho} + \delta_{\sigma\rho^3}$, $u_{2,2} = \delta_{\sigma\rho}\delta_{\sigma\rho^3}$, $a_2 = \delta_{\bar{\rho}}$, and obtain as an E_2 -page

$$E_2 = \mathbf{F}_2[a_2, u_{1,2}, u_{2,2}]/(a_2u_{1,2}).$$

Because the number of classes in the 1-stem and 2-stem equals the Betti numbers of $H^1(BD_8)$ respectively $H^2(BD_8)$ ([AM04, Thm. 2.7]), the LHSSS collapses at E_2 .




Figure 4.2: The $E_2 = E_\infty$ -page of the LHSSS obtained from the extension $H_i \rightarrow D_8 \rightarrow C_2$.

The LHSSS of $Z \rightarrow D_8 \rightarrow C_2 \times C_2$

The extension

$$Z \xrightarrow{\rho^2} D_8 \rightarrow \overline{C}_2 \times \overline{C}_2$$

is central, and therefore the associated LHSSS has trivial coefficient system. Writing $b = \delta_{\overline{\sigma}}$, $c = \delta_{\overline{\sigma\rho}}$ and $x = \delta_{\rho^2}$, the E_2 -page then is

$$E_2 = \mathbf{F}_2[b, c, x],$$

with (s, t) -degrees $|b| = |c| = (1, 0)$, $|x| = (0, 1)$. The class x supports a d_2 with $d_2(x) = bc$ ([AM04, §IV.2]), and the LHSSS collapses at E_3 .




Figure 4.3: The E_2 -page of the LHSSS obtained from the extension $Z \rightarrow D_8 \rightarrow C_2 \times C_2$.

The map j

In this subsection we prove the following proposition, which describes the map j from (2.1) from Section 2.7.

4.4.4 Proposition *The map j from (2.1) is given on cohomology classes by*

$$\begin{aligned} u_{1,i} &\mapsto 0, \\ u_{2,i} &\mapsto x^2, \\ a_1 &\mapsto c, \\ a_2 &\mapsto b. \end{aligned}$$

PROOF The map j is in cohomology determined by the inclusions $Z \rightarrow H_i$ and the corresponding quotient maps $D_8/Z \rightarrow D_8/H$.

The map $Z \rightarrow H_1$ is on elements given by $\rho^2 \mapsto \sigma \cdot \sigma\rho^2$, therefore on cohomology classes by

$$\begin{aligned} \delta_\sigma &\mapsto \delta_{\rho^2} = x, \\ \delta_{\sigma\rho^2} &\mapsto \delta_{\rho^2} = x, \end{aligned}$$

and therefore

$$\begin{aligned} u_{1,1} &= \delta_\sigma + \delta_{\sigma\rho^2} \mapsto 0, \\ u_{2,1} &\mapsto x^2. \end{aligned}$$

The map $D_8/Z \cong C_2 \times C_2 \rightarrow D_8/H_1 \cong C_2$ is on elements given by $\bar{\sigma} \mapsto \bar{e}$ and $\bar{\sigma}\bar{\rho} \mapsto \bar{\rho}$, and therefore on cohomology by

$$a_1 = \delta_{\bar{\rho}} \mapsto \delta_{\bar{\sigma}\bar{\rho}} = c.$$

This completes the formulas for $i = 1$, the formulas for $i = 2$ are obtained from the automorphism $\sigma \leftrightarrow \sigma\rho$ of D_8 . \square

The E_2 -page

We write $u_{1,1} = (u_{1,1}, 0)$, $u_{1,2} = (0, u_{1,2})$ and $u = (u_{2,1}, u_{2,2})$. Proposition 4.4.4 shows that the kernel of j is concentrated in the $s = 0$ -line, and equals

$$\ker(j) = \mathbf{F}_2[u_{1,1}, u_{1,2}, u]/(u_{1,1}u_{1,2}),$$

with bidegrees given by $|u_{1,1}| = |u_{1,2}| = (0, 1)$, and $|u| = (0, 2)$. Moreover, it shows that the image of j is given by

$$\mathrm{Im}(j) = \mathbf{F}_2[c, x^2]\{c\} \oplus \mathbf{F}_2[b, x^2]\{b\} \oplus \mathbf{F}_2[x^2].$$

The split exact sequence (Proposition 2.7.6) then implies that the E_2 -page of the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence is given by




Figure 4.4: The E_2 -page. Red classes come from the B -summand, blue classes from the A -summand. The labeled red classes generated the red summand. The entire page is not finitely generated as a ring, but we labeled two blue classes for ease of comparison with Figure 4.3.

Because the $\mathcal{E}_{(2)}$ -exponent is 2, there can only be non-zero differentials on E_2 , and on E_3 there must be a vanishing line of height 2. The differentials are coming from the LHSSS of the extension $Z \rightarrow D_8 \rightarrow C_2$, by naturality, see Figure 4.3. So there is in fact a horizontal vanishing line of height 1 on the $E_3 = E_\infty$ -page.

Multiplicative extensions

Since the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence is concentrated in the $s = 0$ -line on E_∞ , there are no multiplicative extension problems to solve, and

$$\begin{aligned} H^*(BD_8) &\cong E_\infty^{0,*} \\ &= \mathbf{F}_2[u_{1,1}, u_{1,2}, u]/(u_{1,1}u_{1,2}). \end{aligned}$$

The lower bound illustrated

For $G = D_{2n}$ with $n \equiv 0 \pmod{4}$ and $n \geq 4$ we established in Proposition 4.4.1 that $\exp_{\mathcal{E}(2)} \overline{H\mathbf{F}}_2 = 2$. We illustrate this exponent with a $\mathcal{E}(2)$ -homotopy limit spectral sequence which has a vanishing line of height 2 on the E_∞ -page.

Let $C_n = \langle \rho \rangle \subset D_{2n}$ act on $z \in S^1 \subset \mathbf{C}$ via

$$\rho \cdot z = e^{2\pi i/n} z.$$

Let X be the D_{2n} -space given by inducing the C_n -space S^1 up to D_{2n} :

$$(4.4) \quad X := D_{2n} \times_{C_n} S^1$$

The space X looks like




Figure 4.5: The free D_{2n} -space X .

The D_{2n} -action on X is evidently free. As coset representatives of D_{2n}/C_n we pick $\{e, \sigma\}$. The underlying non-equivariant space is the coproduct $S_e^1 \sqcup S_\sigma^1$, where the subscripts are decorations for the corresponding coset. Denoting the elements of X by (e, z_1) and (σ, z_2) (i.e., by choosing the representatives of the elements of the quotient in Equation (4.4) which have as their first coordinate our chosen coset representatives), the D_{2n} -action on X is explicitly given by

$$\begin{aligned} \rho \cdot (e, z) &= (e, \rho z), \\ \rho \cdot (\sigma, z) &= (\sigma, \rho^{-1} z), \\ \sigma \cdot (e, z) &= (\sigma, z), \\ \sigma \cdot (\sigma, z) &= (e, z). \end{aligned}$$

In particular, $\sigma\rho(\sigma, z) = (e, \rho^{-1}z)$, whereas $\rho\sigma(\sigma, z) = (e, \rho z)$, hence the actions of σ and ρ do not commute.

Since X is a free D_{2n} -space, its homotopy orbits coincide with the actual orbits:

$$X // D_{2n} \simeq \frac{X \times ED_{2n}}{D_{2n}} \simeq X/D_{2n} \simeq S^1/C_n \simeq S^1.$$


Hence the Borel equivariant \mathbf{F}_2 -cohomology of X is:

$$(4.5) \quad H_{D_{2n}}^*(X; \mathbf{F}_2) \cong H^*(S^1; \mathbf{F}_2) \cong \mathbf{F}_2[x]/(x^2).$$

Consider the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence of X :

$$E_2^{s,t} = \lim_{\mathcal{O}(G)_{\mathcal{E}_{(2)}}^{\text{op}}}^s H_{(-)}^t(X, \mathbf{F}_2) \Rightarrow H_{D_{2n}}^{s+t}(X; \mathbf{F}_2).$$

We will show that its E_∞ -page has an element in filtration degree 1, which implies the desired lower bound on the exponent. Note that, given Equation (4.5) and the fact that the functors $H_{(-)}^t(X; \mathbf{F}_2)$ are 0 for $t < 0$, only two E_∞ -pages are possible:



and




Figure 4.6: First possibility for E_∞ .

Figure 4.7: Second possibility for E_∞ .

We will show that the E_∞ -page is the second one, by showing that $x \in H_{D_{2n}}^1(X; \mathbf{F}_2)$ is in filtration ≥ 1 , which is equivalent to showing that x is in the kernel of the edge map

$$H_{D_{2n}}^1(X; \mathbf{F}_2) \rightarrow \lim_{\mathcal{O}(G)_{\mathcal{E}_{(2)}}^{\text{op}}}^0 H_{(-)}^1(X; \mathbf{F}_2).$$

In other words, we must show that for all elementary abelian 2-groups $A \subset D_{2n}$ we have that the map induced by the quotient

$$q: X // A \rightarrow X // D_{2n}$$

in cohomology

$$q^*: H_{D_{2n}}^1(X; \mathbf{F}_2) \rightarrow H_A^1(X; \mathbf{F}_2)$$

satisfies $q^*(x) = 0$. For this, we may restrict to the maximal elementary abelian 2-groups in D_{2n} , because all elementary abelian subgroups are a subgroup of a maximal elementary abelian subgroup. Hence if an element restricts to 0 on all maximal elementary abelian subgroup, it also restricts further down to 0 on all elementary abelian subgroups.

The elements of order 2 of D_{2n} are the reflections $\sigma\rho^i$, and the generator of the center $\rho^{n/2}$. Two reflections $\sigma\rho^i$ and $\sigma\rho^j$ commuting is equivalent to $\rho^{j-i} = \rho^{i-j}$, which is equivalent to $i \equiv j \pmod{n/2}$. Since $\sigma\rho^i\sigma\rho^{i+n/2} = \rho^{n/2}$, we see that all maximal elementary abelian subgroups of D_{2n} are given by $\langle \sigma\rho^i, \sigma\rho^{i+n/2} \rangle$ with $0 \leq i < n/2$.

First dividing out $\sigma\rho^i$ identifies

$$(e, z) \leftrightarrow (\sigma, \rho^i z).$$

Hence every element of $X/\langle\sigma\rho^i\rangle$ is represented uniquely by a (e, z) , and this space is isomorphic to S^1 . We suppress the coset coordinate from now on.

Dividing this out by the central element $\rho^{n/2}$ identifies

$$z \leftrightarrow \rho^{n/2}z,$$

and every element becomes represented uniquely by a $e^{\theta i}$ with $0 \leq \theta < \pi$.

Finally, dividing out by the entire group D_{2n} identifies

$$z \leftrightarrow \rho z,$$

hence every element of X/D_{2n} is uniquely represented by a $e^{\theta i}$ with $0 \leq \theta < 2\pi/n$.

The map $S^1 \cong X/\langle\sigma\rho^i, \sigma\rho^{i+n/2}\rangle \rightarrow X/D_{2n} \cong S^1$ is given by

$$\theta \pmod{\pi} \mapsto \theta \pmod{2\pi/n},$$

which is a $(\frac{n}{2} : 1)$ -cover. Hence the map q^* is given by multiplication by $n/2$.

We assumed $n \equiv 0 \pmod{4}$, hence that $n/2 \equiv 0 \pmod{2}$, and therefore $q^*(x) = 0$, as desired.

4.4.4 The dihedral groups D_{2n} with $n \geq 4$

For the dihedral groups D_{2n} with $n \geq 4$ we will not compute the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence, but instead we will compute the \mathcal{F} -homotopy limit spectral sequence for some family \mathcal{F} which strictly contains $\mathcal{E}_{(2)}$, to be described next.

For $n \geq 4$, the dihedral group has exactly two subgroups isomorphic to D_{2n-1} , which we denote by $H_1 = \langle\sigma, \sigma\rho^2\rangle$ and $H_2 = \langle\sigma\rho, \sigma\rho^3\rangle$. Additionally we write $J = H_1 \cap H_2 = \langle\rho^2\rangle \cong C_{2n-2}$. Write \mathcal{F} for the family $\mathcal{F} = \mathcal{A}ll_{H_1} \cap \mathcal{A}ll_{H_2}$. Then $\mathcal{F} \supset \mathcal{E}_{(2)}$, and therefore $\underline{HF}_2 D_{2n}$ is \mathcal{F} -nilpotent. In this subsection we will compute the \mathcal{F} -homotopy limit spectral sequence for D_{2n} . This spectral sequence will turn out to be additively isomorphic to the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence for D_8 . The computation will also be almost identical.

The decomposition

We apply the decomposition of Section 2.7 with the H_i as in the previous section and $K = J$. Therefore we proceed by computing the LHSS's for the extensions $H_i \rightarrow D_{2n} \rightarrow C_2$ and $J \rightarrow D_{2n} \rightarrow C_2 \times C_2$.

The LHSS of $D_{2n-1} \rightarrow D_{2n} \rightarrow C_2$

We consider the case $D_{2n-1} = H_1$, the other case being obtained by the automorphism $\sigma \leftrightarrow \sigma\rho$. Write $x_1 = \delta_{\overline{\sigma}}$ and $y_1 = \delta_{\overline{\sigma\rho^2}}$ for the classes in $H^1(BD_{2n-1}) = \text{Hom}(D_{2n-1}, \mathbf{F}_2)$. Let w_1 be the unique class in $H^2(BD_{2n-1})$ that classifies $C_2 \rightarrow D_{2n} \rightarrow D_{2n-1}$ and that restricts to zero on both $\langle\sigma\rangle$ and $\langle\sigma\rho^2\rangle$ (see, e.g., [AM04, Thm. 2.7]). Then $C_2 = \langle\overline{\rho}\rangle$ acts on $H_1^{\text{ab}} = H_1/\langle\rho^4\rangle \cong C_2 \times C_2$ by

$$\begin{aligned} \rho\sigma\rho^{-1} &\equiv \sigma\rho^{-2} \equiv \sigma\rho^2 \pmod{\langle\rho^4\rangle}, \\ \rho\sigma\rho^2\rho^{-1} &\equiv \sigma \pmod{\langle\rho^4\rangle}, \end{aligned}$$

hence it interchanges x_1 and y_1 . The class w_1 is uniquely determined by conditions which are preserved by the action of C_2 , hence is fixed.

Write $u_{1,1} = x_1 + y_1$ and $a_1 = \delta_{\bar{\rho}}$. Taking a projective C_2 -resolution of \mathbf{Z} , we see that the $s = 0$ row of the E_2 -page of the LHSSS is given by the C_2 -invariants

$$E_2^{0,*} \cong \mathbf{F}_2[u_{1,1}, w],$$

and the rows for $s \geq 1$ are given by

$$E_2^{s,*} \cong \mathbf{F}_2[w]\{a^s\}.$$

Assembling all rows together gives the E_2 -page

$$E_2^{*,*}(H_1) = \mathbf{F}_2[a, u_{1,1}, w_1]/(a_1 u_{1,1}),$$

with (s, t) -degrees $|a| = (1, 0)$, $|u_{1,1}| = (0, 1)$, and $|w_1| = (0, 2)$.

Consider now the extension $H_2 \rightarrow D_{2^n} \rightarrow C_2$. Write $x_2 = \delta_{\bar{\sigma}\rho}$, $y_2 = \delta_{\bar{\sigma}\rho^3}$, w_2 the unique class in $H^2(BH_2)$ classifying $C_2 \rightarrow D_{2^n} \rightarrow D_{2^{n-1}}$ and restricting to zero on $\langle \sigma\rho \rangle$ and $\langle \sigma\rho^3 \rangle$, $a_2 = \delta_{\bar{\sigma}}$ and $u_{1,2} = x_2 + y_2$. Then by applying the automorphism of D_{2^n} determined by $\sigma \leftrightarrow \sigma\rho$ we get that the E_2 -page of the LHSSS of the extension $H_2 \rightarrow D_{2^n} \rightarrow C_2$ is given by

$$E_2^{*,*}(H_2) = \mathbf{F}_2[a_2, u_{1,2}, w_1](a_2 u_{1,2}).$$

The LHSSS of $C_{2^{n-2}} \rightarrow D_{2^n} \rightarrow C_2 \times C_2$

The extension

$$J \cong C_{2^{n-2}} \rightarrow D_{2^n} \rightarrow C_2 \times C_2$$

gives a LHSSS with trivial coefficient system, for $H^*(BC_{2^{n-2}})$ has only trivial automorphisms. Writing $b = \delta_{\bar{\sigma}}$, $c = \delta_{\bar{\sigma}\rho}$, $z = \delta_{\rho^2}$ and $\beta_{n-2}(z) = \beta_{n-2}(\delta_{\rho^2})$ (the $(n-2)$ -th order Bockstein) we get

$$E_2^{*,*}(J) = \mathbf{F}_2[b, c, z, \beta_{n-2}(z)]/(z^2).$$

The map j

We now determine the effect of the map j from (2.1) from Section 2.7.

4.4.5 Proposition *The map j in the long exact sequence (2.1) is given by*

$$\begin{aligned} u_{1,i} &\mapsto 0, \\ w_i &\mapsto \beta_{n-2}(z), \\ a_1 &\mapsto c, \\ a_2 &\mapsto b. \end{aligned}$$

PROOF The map j is determined by the inclusions $J \rightarrow H_i$ and the corresponding quotient maps $D_{2^n}/J \rightarrow D_{2^n}/H_i$.

For $i = 1$, the inclusion $J = \langle \rho^2 \rangle \hookrightarrow H_1 = \langle \sigma, \sigma\rho^2 \rangle$ is on elements given by $\rho^2 \mapsto \sigma \cdot \sigma\rho^2$, hence on abelianizations by $\overline{\rho^2} \mapsto \overline{\sigma} \cdot \overline{\sigma\rho^2}$, hence on degree-1 cohomology classes by

$$\begin{aligned} x_1 &= \delta_{\overline{\sigma}} \mapsto \delta_{\rho^2} = z, \\ y_1 &= \delta_{\overline{\sigma\rho^2}} \mapsto \delta_{\rho^2} = z, \end{aligned}$$

and hence

$$u_{1,1} = x_1 + y_1 \mapsto 0.$$

To see how the class w_1 restricts, we recall that it classifies the extension $C_2 \rightarrow D_{2^n} \rightarrow D_{2^{n-1}}$, which pulls back along $C_{2^{n-1}} \rightarrow D_{2^{n-1}}$ as

$$\begin{array}{ccccc} C_2 & \longrightarrow & C_{2^{n-1}} & \longrightarrow & C_{2^{n-2}} \\ \downarrow & & \downarrow & & \downarrow \\ C_2 & \longrightarrow & D_{2^n} & \longrightarrow & D_{2^{n-1}} \end{array},$$

which is non split, and therefore the class w_1 restricts to a non zero class, the only possibility being

$$w_1 \mapsto \beta_{n-2}(z).$$

The map $D_{2^n}/J \rightarrow D_{2^n}/H_1$ is on elements given by $\overline{\sigma\rho} \mapsto \overline{\rho}$ and $\overline{\sigma} \mapsto e$, and therefore on cohomology classes by

$$a_1 = \delta_{\overline{\rho}} \mapsto \delta_{\overline{\sigma\rho}} = c.$$

The formulas for $i = 2$ are obtained by applying the automorphism $\sigma \leftrightarrow \sigma\rho$. Observe that in particular this automorphism interchanges H_1 and H_2 , hence b and c . \square

The kernel of j

We write $u_{1,1} = (u_{1,1}, 0)$, $u_{1,2} = (0, u_{1,2})$, $w = (w_1, w_2)$. Then Proposition 4.4.5 shows that the kernel of j is concentrated in the 0-line and equals

$$\ker(j) = \mathbf{F}_2[u_{1,1}, u_{1,2}, w]/(u_{1,1}u_{1,2}).$$

The image of j

Proposition 4.4.5 shows that the image of j is given by

$$\mathbf{F}_2[b, c, \beta_{n-2}(z)].$$

The E_2 -page

This gives an E_2 -page which additively is isomorphic to the one in Figure 4.4. But also all the differentials in Figure 4.4 need to happen in this case, because it is the only possibility for a vanishing line of height 2 on E_3 , which must occur by Corollary 4.4.3.

Multiplicative structure

Because the $E_3 = E_\infty$ -page is concentrated in the $s = 0$ -line, there are no multiplicative extension problems, and we get

$$H^*(BD_{2^n}; \mathbf{F}_2) \cong \mathbf{F}_2[u_{1,1}, u_{1,2}, w]/(u_{1,1}u_{1,2}).$$

4.5 Quaternion groups

The generalized quaternion group Q_{2^n} of order 2^n is the finite subgroup of quaternionic space \mathbf{H} generated multiplicatively by the elements of unit length $\{e^{2\pi i/2^{n-1}}, j\}$ (see, e.g., [CE56, XII.§7]). Denoting these generators by r and s , respectively, one gets the presentation

$$Q_{2^n} = \langle r, s \mid r^{2^{n-2}} = s^2, rsr = s \rangle.$$

for all $n \geq 3$. The subgroup generated by $\langle s^2 \rangle$ is central and of order 2, which makes Q_{2^n} into a central extension ([AM04, IV.2]).

$$(4.6) \quad C_2 \rightarrow Q_{2^n} \rightarrow D_{2^{n-1}}.$$

See Proposition 4.5.2 and Proposition 4.5.3 below for a recollection on this cohomology ring and the LHSSS computing it.

4.5.1 An upper bound on the exponent

In this subsection we will prove the following upper bound on the $\mathcal{E}_{(2)}$ -exponent.

4.5.1 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2, Q_{2^n}} \leq 4.$$

PROOF The proof is a straightforward adaption of the argument in [MNN15, Ex. 5.18].

Let $\mathbf{H} \cong \mathbf{R}^4$ be the 4-dimensional real representation coming from the embedding $Q_{2^n} \hookrightarrow \mathbf{H}$. This is a free action, and restricts to a free action on S^3 . The subgroup $\langle \pm 1 \rangle$ is central, and therefore $Q_{2^n}/\langle \pm 1 \rangle$ acts on $S^3/\langle \pm 1 \rangle = \mathbf{P}(\mathbf{R}^4)$ with isotropy in $\langle \pm 1 \rangle$. The result now follows from Proposition 3.3.3. \square

4.5.2 The $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence

The group Q_{2^n} has a unique maximal elementary abelian 2-subgroup, given by $\langle \pm 1 \rangle$. Therefore, by Proposition 2.6.1, the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence reduces to the LHSSS corresponding to the central extension

$$\langle -1 \rangle = Z(Q_{2^n}) \rightarrow Q_{2^n} \rightarrow D_{2^{n-1}}.$$

This spectral sequence is well known, we recall it here for convenience.

4.5.2 Proposition (see, e.g., [AM04, Lem. 2.10]) *For $n = 3$, the E_2 -page of the LHSSS associated to the extension (4.6) is*

$$E_2^{*,*} = H^*(BC_2^{\times 2}; \mathbf{F}_2) \otimes_{\mathbf{F}_2} H^*(BC_2; \mathbf{F}_2) \cong \mathbf{F}_2[x, y, e],$$

with all the generators in degree 1. The differentials are generated under the Leibniz rule by

$$\begin{aligned} d_2(e) &= x^2 + xy + y^2, \\ d_3(e^2) &= \text{Sq}^1(x^2 + xy + y^2) = x^2y + xy^2. \end{aligned}$$

The spectral sequence collapses at E_4 with a vanishing line of height 4, with resulting cohomology ring

$$H^*(BQ_8; \mathbf{F}_2) \cong \mathbf{F}_2[x, y, e_4]/(x^2 + xy + y^2, x^2y + xy^2).$$

with degrees given by $|x| = |y| = 1$, and $|e_4| = 4$.

4.5.3 Proposition (see, e.g., [AM04, Lem. 2.11]) *Denote the classes $x = \delta_{\bar{\sigma}}$ and $y = \delta_{\bar{\sigma}\bar{\rho}}$.*

For $n \geq 4$, the E_2 -page of the LHSSS associated to the extension (4.6) is given by

$$E_2^{*,*} = H^*(BD_{2^{n-1}}; \mathbf{F}_2) \otimes_{\mathbf{F}_2} H^*(BC_2; \mathbf{F}_2) \cong \mathbf{F}_2[x, y, w, e]/(xy).$$

The differentials are generated under the Leibniz rule by

$$\begin{aligned} d_2(e) &= x^2 + y^2 + w, \\ d_3(e^2) &= \text{Sq}^1(x^2 + y^2 + w) = (x + y)w. \end{aligned}$$

The spectral sequence collapses at E_4 with a vanishing line of height 4, and the resulting cohomology ring is

$$H^*(BQ_{2^n}; \mathbf{F}_2) \cong \mathbf{F}_2[x, y, e_4]/(xy, x^3 + y^3).$$

with $|x| = |y| = 1$, and $|e_4| = 4$.

4.5.3 The $\mathcal{E}_{(2)}$ -exponent

Combining Proposition 4.5.1 with the vanishing lines of height 4 in Proposition 4.5.2 and Proposition 4.5.3 gives the following.

4.5.4 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2Q_{2^n}} = 4.$$

4.6 SD_{16}

Let

$$SD_{16} = \langle s, r \mid s^2 = r^8 = e, srs^{-1} = r^3 \rangle$$

be the semidihedral group of order 16. This group has no unique maximal elementary abelian 2-subgroup, but the subgroup $\langle s, r^2 \rangle \cong D_8$ does contain all the elementary abelian 2-subgroups. We will compute the $\mathcal{A}ll_{D_8}$ -homotopy limit spectral sequence converging to $H^*(BSD_{16}; \mathbf{F}_2)$, which is isomorphic to ([CTVEZ03, App. C, #13(16)])

$$\mathbf{F}_2[z, y, x, w]/(zy, y^3, yx, z^2w + x^2),$$

with degrees $|z| = |y| = 1$, $|x| = 3$ and $|w| = 4$.

By Proposition 2.6.1, the $\mathcal{A}ll_{D_8}$ -homotopy limit spectral sequence is isomorphic to the LHSSS corresponding to the group extension

$$D_8 \xrightarrow{s, r^2} SD_{16} \rightarrow C_2^{\bar{r}}$$

4.6.1 The E_2 -page

Since $rsr^{-1} = sr^2$, C_2 acts on D_8 by interchanging the generators s and sr^2 of D_8 . Write $H^*(BD_8; \mathbf{F}_2) = \mathbf{F}_2[x, y, w]/(xy)$, with $x = \delta_s$, $y = \delta_{sr^2}$, and where w is the unique class in degree 2 that restricts to 0 on $\langle s \rangle$ and $\langle sr^2 \rangle$ and classifies an extension of D_8 isomorphic to D_{16} (see, e.g., [AM04, Thm. 2.7]). Then $C_2 = \langle \bar{r} \rangle$ acts by interchanging x and y . Of course it sends an extension isomomorphic to D_{16} to another such extension, and since s and sr^2 get interchanged, we see that the C_2 -action leaves w fixed, by the restriction property mentioned before.

To determine $E_2^{s,t} = H^s(BC_2; H^t(BD_8; \mathbf{F}_2))$, let

$$\dots \xrightarrow{1-\bar{r}} \mathbf{Z}C_2 \xrightarrow{1+\bar{r}} \mathbf{Z}C_2 \xrightarrow{1-\bar{r}} \mathbf{Z}C_2 \xrightarrow{\epsilon} \mathbf{Z}$$

be the minimal free C_2 -resolution of \mathbf{Z} . Applying $\text{Hom}_{C_2}(-, H^*(BD_8; \mathbf{F}_2))$, we see that the differentials are on monomials given by

$$\begin{aligned} (x^i + y^i)w^j &\leftarrow x^i w^j, \\ (x^k + y^k)w^l &\leftarrow y^k w^l. \end{aligned}$$


Hence the kernel has as an \mathbf{F}_2 -basis the elements of the form $w^j P(x, y)$, where $P(x, y)$ is a symmetric polynomial in x and y . Taking the cokernel divides out the symmetrized polynomials in x and y . Hence the result has as an \mathbf{F}_2 -basis the polynomials of the form $x^i y^i w^j$, which is 0 unless $i = 0$. Writing $a = \delta_{\bar{r}}$, and $\sigma_1 = x + y$, we get

$$E_2^{s,t} = \mathbf{F}_2[a, \sigma_1, w]/(a\sigma_1),$$

with (s, t) -degrees given by $|a| = (1, 0)$, $|\sigma_1| = (0, 1)$ and $|w| = (0, 2)$.

4.6.2 Differentials and $E_4 = E_\infty$

Since $SD_{16}^{\text{ab}} = SD_{16}/\langle r^4 \rangle \cong C_2 \times C_4$, we have $\dim_{\mathbf{F}_2} H^1(BSD_{16}; \mathbf{F}_2) = 2$. Hence $x + y$ is a permanent cycle. Since there is no vanishing line on E_2 , w will have to support a differential, which can only be $d_3 w = a^3$. Hence the E_3 -page is

Figure 4.8: The E_3 -page.

Therefore the E_4 -page is

$$\mathbf{F}_2[a, \sigma_1, [\sigma_1 w], [w^2]] / (a\sigma_1, a[\sigma_1 w], \sigma_1^2[w^2] - [\sigma_1 w]^2, a^3).$$

Hence the E_4 -page has a vanishing line of height 3, and therefore $E_4 = E_\infty$. There are several multiplicative extension problems, which we leave unresolved.

4.6.3 Poincaré Series

The E_∞ -page has as $s = 0$ -line

$$\begin{aligned} & \mathbf{F}_2[\sigma_1, [\sigma_1 w], [w^2]] / (\sigma_1^2[w^2] - [\sigma_1 w]^2) \\ & \cong \mathbf{F}_2[\sigma_1, [w^2]] \{1, [\sigma_1 w]\}, \end{aligned}$$

which has Poincaré series

$$\frac{1 + t^3}{(1 - t)(1 - t^4)}.$$

The $s = 1$ -line of the E_∞ -page is given by

$$\mathbf{F}_2[[w^2]] \{a\},$$

which has Poincaré series

$$\frac{t}{1 - t^4}.$$

The $s = 2$ -line of the E_∞ -page is given by

$$\mathbf{F}_2[[w^2]] \{a^2\},$$

which has Poincaré series

$$\frac{t^2}{1-t^4}.$$

This shows that the Poincaré series of $H^*(BSD_{16}; \mathbf{F}_2)$ is given by

$$\begin{aligned} \frac{1+t^3}{(1-t)(1-t^4)} + \frac{t+t^2}{1-t^4} &= \frac{(1+t^3) + t(1-t) + t^2(1-t)}{(1-t)(1-t^4)} \\ &= \frac{1+t}{(1-t)(1-t^4)} \\ &= \frac{1}{(1-t)^2(1+t^2)}, \end{aligned}$$

cf. [CTVEZ03, App. B and App. G].

4.6.4 An upper bound on the $\mathcal{E}_{(2)}$ -exponent

In this subsection we will prove the following upper bound on the $\mathcal{E}_{(2)}$ -exponent.

4.6.1 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2SD_{16}} \leq 4.$$

PROOF We will use the notation R_θ and T from the proof of Proposition 4.7.1, and also use the fact proven there that R_θ fixes a line only if $\theta \in \mathbf{Z}\{\pi\}$. Write $\alpha = 2\pi/8$. Let V be the real SD_{16} -representation given by

$$\begin{aligned} s &\mapsto T, \\ r &\mapsto R_\alpha \oplus R_\alpha \end{aligned}$$

from [Tot14, Lem. 13.4]. The same argument as in the proof of Proposition 4.7.1 shows that the only powers of r fixing a line are e and r^4 , and that the only elements of the form sr^k that can possibly fix a line are s, sr^2, sr^4, sr^6 . The elements

$$e, r^4, s, sr^2, sr^4, sr^6.$$

do not generate an elementary abelian 2-group as in the proof of Proposition 4.7.1, but all the isotropy groups of lines that occur consist of proper subsets of the above elements that do form elementary abelian 2-groups, as we will now show.

First, r^4 fixes every line, so is in every isotropy group.

Write $[x_1 : y_1 : x_2 : y_2]$ for the real homogenous coordinates in \mathbf{RP}^3 . Then

$$s \cdot [x_1 : y_1 : x_2 : y_2] = [x_2 : y_2 : x_1 : y_1],$$

which is fixed if and only if $(x_1, y_1) = (x_2, y_2)$ or $(x_1, y_1) = -(x_2, y_2)$. Computation of $sr^4 \cdot [x_1 : y_1 : x_2 : y_2]$ shows that this element fixes exactly the same lines.

Computing

$$sr^2 \cdot [x_1 : y_1 : x_2 : y_2] = [-y_2 : x_2 : y_1 : -x_1]$$

shows that sr^2 fixes precisely those lines with $(x_1, y_1) = (-y_2, x_2)$ or $(x_1, y_1) = (y_2, -x_2)$. Computing $sr^6 \cdot [x_1 : y_1 : x_2 : y_2]$ shows that this element fixes precisely the same lines.

The sets of lines $\{x_1 = x_2, y_1 = y_2\} \cup \{x_1 = -x_2, y_1 = -y_2\}$ and $\{x_1 = -y_2, y_1 = x_2\} \cup \{x_1 = y_2, y_1 = -x_2\}$ have empty intersection, because a line in the intersection $\{x_1 = x_2, y_1 = y_2\} \cap \{x_1 = -y_2, y_1 = x_2\}$ would satisfy

$$y_2 = y_1 = x_2 = x_1 = -y_2,$$

a line in the intersection $\{x_1 = x_2, y_1 = y_2\} \cap \{x_1 = y_2, y_1 = -x_2\}$ would satisfy

$$y_1 = y_2 = x_1 = x_2 = -y_1,$$

a line in the intersection $\{x_1 = -x_2, y_1 = -y_2\} \cap \{x_1 = -y_2, y_1 = x_2\}$ would satisfy

$$y_2 = -y_1 = -x_2 = x_1 = -y_2,$$

and finally a line in the intersection $\{x_1 = -x_2, y_1 = -y_2\} \cap \{x_1 = y_2, y_1 = -x_2\}$ would satisfy

$$y_2 = x_1 = -x_2 = y_1 = -y_2.$$

Hence the isotropy groups that occur are

$$\langle r^4 \rangle, \langle s, sr^4 \rangle, \langle sr^2, sr^6 \rangle,$$

all of which are elementary abelian 2-groups. Applying Proposition 3.3.3 gives the desired result. \square

4.6.5 The exponent

The 1-dimensional real representation

$$SD_{16} \rightarrow SD_{16}/D_8 \cong C_2$$

has Euler class a . This has isotropy in $\mathcal{A}ll_{D_8}$, and the vanishing line shows that a is of nilpotence degree 3, hence

$$\exp_{\mathcal{A}ll_{D_8}} \underline{HF}_2 = 3.$$

Since increasing the family only can make the exponent go down, we have as a consequence

4.6.2 Proposition *The exponent $\mathcal{E}_{(2)}$ -exponent satisfies*

$$3 \leq \exp_{\mathcal{E}_{(2)}} \underline{HF}_{2SD_{16}} \leq 4.$$

4.7 M_{16}

4.7.1 Introduction

Let

$$M_{16} = \langle r, f \mid r^8 = f^2 = e, frf^{-1} = r^5 \rangle$$

be the modular group of order 16. The group has this name because its lattice of subgroups is modular (see, e.g., [Bir67, I.§7]).

In this section we will compute the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BM_{16}; \mathbf{F}_2)$, which is isomorphic to ([CTVEZ03, App. C, #11(16)])

$$\mathbf{F}_2[z, y, x, w]/(z^2, zy^2, zx, x^2)$$

with degrees $|z| = |y| = 1$, $|x| = 3$, $|w| = 4$.

4.7.2 An upper bound on the exponent

In this subsection we prove the following upper bound on the $\mathcal{E}_{(2)}$ -exponent.

4.7.1 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2M_{16}} \leq 4.$$

PROOF For an angle θ , let R_θ be the rotation-by- θ matrix $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$.

Let T be the matrix

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

which interchanges the summands of $\mathbf{R}^2 \oplus \mathbf{R}^2$. Write $\alpha = 2\pi/8$. Let V be the 4-dimensional real M_{16} -representation given by

$$\begin{aligned} f &\mapsto T, \\ r &\mapsto R_\alpha \oplus R_{5\alpha}. \end{aligned}$$

This is the representation from [Tot14, Lem. 13.3].

We will now determine the isotropy of the projectivation $\mathbf{P}(V)$.

A power of r fixes a line in $\mathbf{R}^2 \oplus \mathbf{R}^2$ if and only if it fixes a line in one of the summands. The matrix R_θ has characteristic polynomial $P(\lambda) = \lambda^2 - 2\lambda \cos \theta + 1$, which has discriminant $\Delta = -4\sin^2 \theta$. Hence R_θ has real eigenvalues only if $\theta \in \mathbf{Z}\{\pi\}$. Since $r \mapsto R_{2\pi/8}$, r^k fixes a line only if $\alpha k \in \mathbf{Z}\{\pi\}$ or $5\alpha k \in \mathbf{Z}\{\pi\}$, that is only if $k \in \mathbf{Z}\{4\}$. Therefore e and r^4 are the only elements powers of r that can possibly fix a line. The element r^4 acts by $-\text{Id}$, and therefore fixes all lines. The element e of course also fixes all lines.

We now consider the remaining elements of M_{16} , which are of the form fr^k , and which act by $T(R_{\alpha k} \oplus R_{5\alpha k})$. This matrix certainly only fixes a line if its square does, which is $R_{6\alpha k} \oplus R_{6\alpha k}$. These summands only fix a line if $6\alpha k \in \mathbf{Z}\{\pi\}$, which implies that k is even. Hence the only possible elements of the form fr^k fixing a line are f, fr^2, fr^4 , and fr^6 . One verifies directly that fr^2 acts by the matrix

$$\begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

which fixes no lines. Consequently its inverse fr^6 neither fixes lines. One also easily checks that fr^4 acts by

$$\begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

which fixes the lines $[x_1 : y_1 : x_2 : y_2]$ in $\{x_1 = -x_2, y_1 = -y_2\}$. Hence the only elements of M_{16} fixing a line are e, f, fr^4, r^4 , which together form a Klein four group. Applying Proposition 3.3.3 gives the desired result. \square

4.7.3 The orbit category

The group M_{16} has a unique maximal elementary abelian 2-subgroup, given by

$$C_2 \times C_2 \cong \langle f, fr^4 \rangle.$$

By Proposition 2.6.1, this identifies the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence with the LHSSS associated to the group extension

$$C_2 \times C_2 \rightarrow M_{16} \rightarrow \langle \bar{r} \rangle \cong C_4.$$

4.7.4 The E_2 -page

The action of $C_4 = \langle \bar{r} \rangle$ on $C_2 \times C_2 = \langle f, fr^4 \rangle$ is given by $rfr^{-1} = fr^4$, i.e. \bar{r} interchanges the generators f and fr^4 . Denote the duals by $x := \delta_f, y := \delta_{fr^4} \in H^1(BC_2 \times C_2; \mathbf{F}_2)$. Then $H^*(BC_2 \times C_2; \mathbf{F}_2) = \mathbf{F}_2[x, y]$, and \bar{r} acts by interchanging x and y .

A 2-periodic projective C_4 -resolution of \mathbf{Z} is given by

$$\cdots \rightarrow \mathbf{Z}C_4 \xrightarrow{1-\bar{r}} \mathbf{Z}C_4 \xrightarrow{1+\bar{r}+\bar{r}^2+\bar{r}^3} \mathbf{Z}C_4 \xrightarrow{1-\bar{r}} \mathbf{Z}C_4 \xrightarrow{\epsilon} \mathbf{Z}.$$

Applying $\text{Hom}_{\mathbf{Z}C_4}(-, \mathbf{F}_2[x, y])$ (where $\mathbf{F}_2[x, y]$ carries the C_4 -action described above) yields

$$\cdots \xleftarrow{0} \mathbf{F}_2[x, y] \xleftarrow{0} \mathbf{F}_2[x, y] \xleftarrow{0} \mathbf{F}_2[x, y] \xleftarrow{0} \mathbf{F}_2[x, y],$$

where the maps going from odd to even degrees are 0, and the maps from even to odd degrees are on monomials given by

$$x^i y^j + x^j y^i \leftarrow x^i y^j.$$

This shows that the even rows of the E_2 -page are given by the C_4 -invariants of $\mathbf{F}_2[x, y]$, whereas the odd rows are given by the coinvariants.

We denote $c_0 := \delta_{\bar{r}} \in H^1(BC_4; \mathbf{F}_2)$, and let $\beta_2(c_0) \in H^2(BC_4; \mathbf{F}_2)$ be the second order Bockstein of c_0 . Then $H^*(BC_4; \mathbf{F}_2) = \mathbf{F}_2[c_0, \beta_2(c_0)]/(c_0^2)$. In addition we write $\sigma_1 = x + y$ and $\sigma_2 = xy$ for the first and second elementary symmetric polynomials in x and y .

The above discussion shows that $E_2^{0,*} = \mathbf{F}_2[\sigma_1, \sigma_2]$, and more generally that the the even rows of the E_2 -page equals

$$(4.7) \quad E_2^{0 \pmod{2},*} \cong \mathbf{F}_2[\sigma_1, \sigma_2, \beta_2(c_0)].$$

The $s \equiv 1 \pmod{2}$ -rows of E_2 are isomorphic to the coinvariants, and are as such modules over the $s = 0$ -row of E_2 , which is the invariants. In the following lemma we describe this module structure, because we will use it to organize the calculation.

4.7.2 Lemma *The $\mathbf{F}_2[x, y]^{C_2}$ -module $\mathbf{F}_2[x, y]_{C_2}$ is (non-freely) generated by $\{[1], [x]\}$. (Recall that we denote by $[P]$ the coinvariance class of $P \in \mathbf{F}_2[x, y]$.)*

PROOF Writing $\sigma_1 = x + y$, $\sigma_2 = xy$ for the first and second elementary symmetric polynomials, we prove that the degree n -part of $\mathbf{F}_2[x, y]_{C_2}$ is generated by $\{[1], [x]\}$ by induction on n . Since the degree 0 and 1 part of $\mathbf{F}_2[x, y]_{C_2}$ are given by $\mathbf{F}_2\{[1]\}$ and $\mathbf{F}\{[x]\}$ respectively, the lemma holds for $n = 0, 1$.

Assume now $n \geq 2$. The degree n -part of $\mathbf{F}_2[x, y]_{C_2}$ is given by

$$\mathbf{F}_2\{[x^n], [x^{n-1}y], \dots, [x^{\lfloor n/2 \rfloor} y^{\lfloor n/2 \rfloor}]\}.$$

All elements $[x^i y^j] \neq [x^n]$ in this basis have strictly positive x - and y -exponent, and therefore

$$[x^i y^j] = xy[x^{i-1}y^{j-1}],$$

and xy is an invariant and $[x^{i-1}y^{j-1}]$ is a coinvariant of degree strictly smaller than n , hence by induction in the $\mathbf{F}_2[x, y]^{C_2}$ -module generated by $\{[1], [x]\}$.

The element $[x^n]$, finally, equals

$$[x^n] = (x + y)[x^{n-1}] + xy[x^{n-2}],$$

where the coinvariance classes on the right hand side are also of degree strictly smaller than n .

Notice that we used the induction hypothesis for $n - 1$ and $n - 2$, but not smaller, so that we do not use more than the base case provided. \square

Let $\mathbf{F}_2[\sigma_1, \sigma_2]$ be the free graded ring with $|\sigma_1| = 1$, $|\sigma_2| = 2$, and let

$$\mathbf{F}_2[\sigma_1, \sigma_2]\{c_0, c_1\}$$

be the free graded module with $|c_0| = 0$ and $|c_1| = 1$. By Lemma 4.7.2, the morphism of $\mathbf{F}_2[\sigma_1, \sigma_2]$ -modules given by

$$(4.8) \quad \begin{aligned} \mathbf{F}_2[\sigma_1, \sigma_2]\{c_0, c_1\} &\rightarrow \mathbf{F}_2[x, y]_{C_2} \\ c_0 &\mapsto [1] \\ c_1 &\mapsto [x] \end{aligned}$$

is surjective.

4.7.3 Lemma *The kernel of the map (4.8) is $\mathbf{F}_2[\sigma_1, \sigma_2]\{\sigma_1 c_0\}$.*

PROOF This follows from a Poincaré series calculation. The Poincaré series of the right hand side of (4.8) is

$$\frac{1+t}{(1-t^2)^2} = \frac{1}{(1-t)(1-t^2)}.$$

The Poincaré series of the left hand side of (4.8) is

$$\frac{1+t}{(1-t)(1-t^2)}.$$

Certainly, $\mathbf{F}_2[\sigma_1, \sigma_2]\{\sigma_1 c_0\}$ is contained in the kernel, for $\sigma_1 c_0 \mapsto (x+y)[1] = 0$. The module $\mathbf{F}_2[\sigma_1, \sigma_2]\{\sigma_1 c_0\}$ has Poincaré series

$$\frac{t}{(1-t)(1-t^2)}.$$

Because

$$\frac{1+t}{(1-t)(1-t^2)} - \frac{t}{(1-t)(1-t^2)} = \frac{1}{(1-t)(1-t^2)},$$

we see that in fact this is all of the kernel. \square

Returning to our description of the E_2 -page, Lemma 4.7.3 shows that the odd rows are given by

$$(4.9) \quad \begin{aligned} E_2^{1 \pmod{2},*} &\cong \frac{\mathbf{F}_2[\sigma_1, \sigma_2]\{c_0, c_1\}}{\mathbf{F}_2[\sigma_1, \sigma_2]\{\sigma_1 c_0\}} \otimes \mathbf{F}_2[\beta_2(c_0)] \\ &\cong \frac{E_2^{0,*}\{c_0, c_1\}}{E_2^{0,*}\{\sigma_1 c_0\}} \otimes \mathbf{F}_2[\beta_2(c_0)]. \end{aligned}$$

Combining (4.9) with (4.7) gives the E_2 -page:

$$E_2^{*,*} \cong \mathbf{F}_2[\sigma_1, \sigma_2, c_0, \beta_2(c_0), c_1] / (c_0^2, c_1^2, c_0 c_1, \sigma_1 c_0)$$

with (s, t) -degrees given by $|\sigma_1| = (0, 1)$, $|\sigma_2| = (0, 2)$, $|c_0| = (1, 0)$, $|\beta_2(c_0)| = (2, 0)$, $|c_1| = (1, 1)$. The E_2 -page is depicted (without differentials) in the following figure:




Figure 4.9: The E_2 -page, without differentials. The even rows, which are the invariants, are colored green, the odd rows, which are the coinvariants, are colored orange.

4.7.5 Differentials on the E_2 -page

Generating differentials

Denote the dimension of $H^i(BM_{16}; \mathbf{F}_2)$, i.e. the i -th Betti number, by h^i . We assume known that $h^1 = 2$ and $h^2 = 2$ in determining the differentials. The first follows from $M_{16}^{\text{ab}} \cong C_2 \times C_2$, the second can be verified either by hand, or using a computer algebra system such as GAP.

The classes c_0 and $\beta_2(c_0)$ are permanent cycles for degree reasons, and σ_1 is a permanent cycle because $h^1 = 1$ and there are 2 classes in the 1-stem. There are 4 classes in the 2-stem, but $h^2 = 2$, so two will have to support a differential, and by the previous discussion, these will have to be σ_2 and c_1 . The only possibility for c_1 to support a differential is

$$(4.10) \quad d_2(c_1) = c_0\beta_2(c_0).$$

Given this, the only remaining possibility for σ_2 to support a differential is

$$(4.11) \quad d_2(\sigma_2) = \sigma_1\beta_2(c_0).$$

We will study the propagation of these differentials using the Leibniz rule for the even and odd rows separately.

Differentials in the even rows

The even rows of the E_2 -page are given by

$$\begin{aligned} E_2^{0 \pmod{2},*} &\cong \mathbf{F}_2[\sigma_1, \sigma_2, \beta_2(c_0)] \\ &\cong \mathbf{F}_2[\sigma_1, [\sigma_2^2], \beta_2(c_0)] \oplus \mathbf{F}_2[\sigma_1, [\sigma_2^2], \beta_2(c_0)]\{\sigma_2\}. \end{aligned}$$

Propagating the differential (4.11) gives the following pattern of differentials:




Figure 4.10: The differentials in the even rows on the E_2 -page.

This also shows that there is no possibility for more d_2 's in the even rows on the E_2 -page, and that the even rows of the E_3 -page are given by

$$E_3^{0 \pmod{2},*} \cong \mathbf{F}_2[\sigma_1, [\sigma_2^2], \beta_2(c_0)]/(\sigma_1\beta_2(c_0)).$$

Differentials in the odd rows

In the odd rows, propagating the differentials (4.10) and (4.11) using the Leibniz rule, we get

$$\begin{aligned} d_2(\sigma_2 c_1) &= \sigma_1 c_1 \beta_2(c_0) + \sigma_2 c_0 \beta_2(c_0), \\ d_2(c_1) &= c_0 \beta_2(c_0), \\ d_2(\sigma_1 \sigma_2 c_1) &= \sigma_1^2 c_1 \beta_2(c_0). \end{aligned}$$


This implies that the classes in the odd rows supporting a differential generate the submodule

$$\begin{aligned} & \left(\mathbf{F}_2[\sigma_2^2]\{\sigma_2 c_1\} \oplus \mathbf{F}_2[\sigma_2^2]\{c_1\} \right. \\ & \left. \oplus \mathbf{F}_2[\sigma_1, \sigma_2^2]\{\sigma_1 \sigma_2 c_1\} \right) \otimes \mathbf{F}_2[\beta_2(c_0)], \end{aligned}$$

and that the classes that get hit are given by

$$\begin{aligned} & \left(\mathbf{F}_2[\sigma_2^2]\{\sigma_1 c_1 + \sigma_2 c_0\} \oplus \mathbf{F}_2[\sigma_2^2]\{c_0\} \right. \\ & \left. \oplus \mathbf{F}_2[\sigma_1, \sigma_2^2]\{\sigma_1^2 c_1\} \right) \otimes \mathbf{F}_2[\beta_2(c_0)]\{\beta_2(c_0)\}. \end{aligned}$$

This pattern of differentials is depicted in the following figure.

Figure 4.11: The differentials in the odd rows of E_2 .


This implies that the classes in the $s = 1$ -line surviving to E_3 are

$$E_3^{1,*} \cong \mathbf{F}_2[\sigma_1, [\sigma_2^2]]\{\sigma_1 c_1\} \oplus \mathbf{F}_2[[\sigma_2^2]]\{c_0, [\sigma_2 c_0]\}$$

and that the odd s -rows with $s \geq 3$ surviving to E_3 are

$$\begin{aligned} E_3^{s,*} &\cong \frac{\mathbf{F}_2[\sigma_1, [\sigma_2^2]]\{[\sigma_1 c_1]\}}{\mathbf{F}_2[\sigma_1, [\sigma_2^2]]\{\sigma_1[\sigma_1 c_1]\}} \\ &\cong \mathbf{F}_2[[\sigma_2^2]]\{[\sigma_1 c_1]\}. \end{aligned}$$

Putting the differentials in the even and odd rows together gives the following E_2 -page:

Figure 4.12: The E_2 -page with differentials.

4.7.6 The E_3 -page

For a horizontal vanishing line of height 4 on E_5 there must be a non-zero

$$d_3: E_3^{1,2} \rightarrow E_3^{4,0}.$$

We have $E_3^{1,2} = \mathbf{F}_2\{\sigma_1 c_1, \sigma_2 c_0\}$, hence at least one of these two classes has to support a differential, the only possible target for both is to hit $\beta_2(c_0)^2$ with a d_3 . Because $\beta_2(c_0)^2$ is the only class in $E_3^{4,0}$, we have that $d_3([\sigma_1 c_1]) \neq 0$ if and only if $d_3([\sigma_1 c_1]) = \beta_2(c_0)^2$. Similarly, we have that $d_3([\sigma_1 c_1]\beta_2(c_0)) \neq 0$ if and only if $d_3([\sigma_1 c_1]\beta_2(c_0)) = \beta_2(c_0)^3$. By the Leibniz rule, $d_3([\sigma_1 c_1]) = \beta_2(c_0)^2$ implies that $d_3([\sigma_1 c_1]\beta_2(c_0)) = \beta_2(c_0)^3$. Furthermore, if $d_3([\sigma_1 c_1]) = 0$ then, also by the Leibniz rule, we have $d_3([\sigma_1 c_1]\beta_2(c_0)) = 0$. Therefore $d_3([\sigma_1 c_1]) = \beta_2(c_0)^2$ if and only if $d_3([\sigma_1 c_1]\beta_2(c_0)) = \beta_2(c_0)^3$.

Similarly, $d_3(\sigma_2 c_0) = \beta_2(c_0)^2$ if and only if $d_3(\sigma_2 c_0 \beta_2(c_0)) = \beta_2(c_0)^3$. But since

$$[\sigma_1 c_1 \beta_2(c_0)] = [\sigma_2 c_0 \beta_2(c_0)],$$


as a consequence of

$$d_2(\sigma_2 c_1) = (\sigma_1 c_1 + \sigma_2 c_0)\beta_2(c_0)$$

we have:

$$d_3(\sigma_1 c_1) = d_3(\sigma_2 c_0) = \beta_2(c_0)^2.$$

This is depicted in

Figure 4.13: The E_3 -page with differentials.

This shows that E_4 has a horizontal vanishing line of height 3, and is therefore the E_∞ -page.

4.7.7 Poincaré Series

The 0-line $E_\infty^{0,*}$ is given by $\mathbf{F}_2[\sigma_1, [\sigma_2^2]]$, which has Poincaré series $1/((1-t)(1-t^4))$. The 2-line $E_\infty^{2,*}$ is given by $\mathbf{F}_2[[\sigma_2^2]]\{\beta_2(c_0)\}$, which has Poincaré series $t^2/(1-t^4)$.

The 1-line of E_3 is given by

$$E_3^{1,*} \cong \mathbf{F}_2[\sigma_1, [\sigma_2^2]]\{\sigma_1 c_1\} \oplus \mathbf{F}_2[[\sigma_2^2]]\{c_0, [\sigma_2 c_0]\},$$

which has Poincaré series

$$\frac{t^3}{(1-t)(1-t^4)} + \frac{t}{1-t^2}.$$

The d_3 -differentials originating from the $s = 1$ -line kill 1 class in each $s+t$ -stem with $s+t \equiv 3 \pmod{4}$, hence $E_4^{1,*} = E_\infty^{1,*}$ has Poincaré series

$$\frac{t^3}{(1-t)(1-t^4)} + \frac{t}{1-t^2} - \frac{t^3}{1-t^4}.$$

Taking everything together, and using

$$1 + (1-t)t^2 + t^3 + t(1+t^2)(1-t) - t^3(1-t) = 1+t,$$

we get that $H^*(BM_{16}; \mathbf{F}_2)$ has Poincaré series

$$\frac{1+t}{(1-t)(1-t^4)} = \frac{1}{(1-t)^2(1+t^2)},$$

4.7.8 The exponent

The above computation shows that $\exp_{\mathcal{E}(2)} \underline{HF}_2 \geq 3$, which together with Proposition 4.7.1 shows that

4.7.4 Proposition *The $\mathcal{E}(2)$ -exponent satisfies*

$$3 \leq \exp_{\mathcal{E}(2)} \underline{HF}_2 \leq 4.$$

4.8 $D_8 * C_4$

4.8.1 Introduction

Let $D_8 = \langle \sigma, \rho \rangle$ be the dihedral group of order 8 and let $C_4 = \langle \gamma \rangle$ be the cyclic group of order 4. Both these groups have central cyclic subgroups of order 2, for D_8 this is $\langle \rho^2 \rangle$ and for C_4 this is $\langle \gamma^2 \rangle$. The *central product* of D_8 and C_4 is defined to be the direct product with these central subgroups identified:

$$D_8 * C_4 := D_8 \times C_4 / \langle \rho^2 \gamma^{-2} \rangle.$$

In this section we will evaluate the \mathcal{F} -homotopy limit spectral sequence converging to $H^*(BD_8 * C_4; \mathbf{F}_2)$ for the family $\mathcal{F} = \mathcal{A}$ of abelian subgroups of $D_8 * C_4$. This ring is isomorphic to ([CTVEZ03, App. C, #8(16)])

$$\mathbf{F}_2[z, y, x, w] / (zx + y^2 + x^2, z^2x + zx^2).$$

with degrees $|z| = |y| = |x| = 1$, and $|w| = 4$.

4.8.2 Summary of the computation

Because the computation of the \mathcal{A} -homotopy limit spectral sequence of $D_8 * C_4$ is rather long, we summarize it here.

The cohomology of $D_8 * C_4$ is detected on abelian subgroups (see Lemma 4.8.1 below). This lemma is part of the input of the calculation, and the proof of this lemma uses the depth of the cohomology ring of $D_8 * C_4$ as input. In particular, the computation of the \mathcal{A} -homotopy limit spectral sequence carried out here does not provide a new computation of the cohomology of $D_8 * C_4$.

The strategy is the same as the one followed for the calculation of the $\mathcal{E}(2)$ -homotopy limit spectral sequence of D_8 (Section 4.4), that is, we apply the decomposition from Section 2.7.

Therefore we compute the LHSS's needed for this decomposition. We need 4 LHSS's, but 3 of them we can identify using automorphisms of $D_8 * C_4$, which we construct first.

After the computation of the LHSS's we compute the map j from the LES in (2.1). This then allows us to calculate the A - and B - summand from (2.8), and hence the E_2 -page.

Next we compute an upper bound of 3 on the \mathcal{A} -exponent of $\underline{HF}_2_{D_8 * C_4}$, which implies that the \mathcal{A} -homotopy limit spectral sequence will collapse at E_4 with a vanishing line of height 4. However, because $D_8 * C_4$ has its cohomology detected on abelian subgroups, there will be even a vanishing line of height 1, i.e., the spectral sequence will collapse to the $s = 0$ -line.

Using naturality and the fact that the spectral sequence will collapse to the $s = 0$ -line at E_4 , we can compute the rank of each differential in the \mathcal{A} -homotopy limit spectral sequence. We are not able to determine precisely which class hits which, but we are able to determine which classes in the $s = 0$ -line support a differential.

4.8.3 The orbit category

The family $\mathcal{F} = \mathcal{A}$ of abelian subgroups gives the following subcategory of the orbit category:

The Weyl groups are not displayed in Figure 4.14. They all turn out to be elementary abelian.

We introduce the following notation for the maximal abelian subgroups of $D_8 * C_4$, all of which are isomorphic to $C_4 \times C_2$

$$\begin{aligned} A_1 &= \langle \gamma, \rho\gamma \rangle, \\ A_2 &= \langle \gamma, \sigma \rangle, \\ A_3 &= \langle \gamma, \sigma\rho \rangle. \end{aligned}$$

In addition, we denote the cyclic group of order 4 generated by γ by

$$C = \langle \gamma \rangle.$$

The group C equals any non-redundant intersection of the A_i 's:

$$C = A_i \cap A_j = A_1 \cap A_2 \cap A_3$$

for all $i \neq j$. We apply the decomposition of Section 2.7 with the $H_i = A_i$, and with $K = C$. Therefore we will compute the LHSSS's obtained from the extensions $C \rightarrow D_8 * C_4 \rightarrow C_2 \times C_2$ and $A_i \rightarrow D_8 * C_4 \rightarrow C_2$. Before doing so, we determine an upper bound on the \mathcal{A} -exponent of $\underline{HF}_{2D_8 * C_4}$.

4.8.4 An upper bound on the \mathcal{A} -exponent

To give an upper bound on $\exp_{\mathcal{A}} \underline{HF}_{2D_8 * C_4}$, we will use the following lemma.

4.8.1 Lemma *The \mathbf{F}_2 -cohomology of $D_8 * C_4$ is detected on abelian subgroups.*

PROOF The centralizers of the elementary abelian 2-subgroups of rank 2 of $D_8 * C_4$ are precisely the A_i . If there would be a cohomology class which is not detected on the A_i , then this would imply by [Car95, Thm. 2.3] that the depth of $H^*(BD_8 * C_4; \mathbf{F}_2)$ is < 2 . But the ring $H^*(BD_8 * C_4; \mathbf{F}_2)$ has depth 2 (see, e.g., [CTVEZ03, App. C, #8(16)]), which gives the desired result. \square

4.8.2 Corollary *The \mathcal{A} -homotopy limit spectral sequence converging to $H^*(BD_8 * C_4)$ collapses at a finite page to the $s = 0$ -line.*

In addition, Lemma 4.8.1 allows us to prove the following.

4.8.3 Proposition *The \mathcal{A} -exponent of $\underline{HF}_{2D_8 * C_4}$ is ≤ 3 .*

PROOF For each i , let τ_i denote the pullback of the sign representation along the quotient map $D_8 * C_4 \rightarrow D_8 * C_4 / A_i \cong C_2$. Then $T = \bigoplus_i \tau_i$ is a real representation of dimension 3 with isotropy in \mathcal{A} . Moreover, T has a trivial summand when restricted to any of the A_i , hence the Euler class e_T is 0 by Lemma 4.8.1. By Corollary 3.2.10 the result follows.

4.8.5 The decomposition

We apply the decomposition of Section 2.7 with $K = C$ and the $H_i = A_i$.

We therefore proceed by computing the LHSS's $E_*^{*,*}(A_i)$ and $E_*^{*,*}(C)$, and subsequently A^* and B^* , which will then give us the E_2 -page of the \mathcal{A} -homotopy limit spectral sequence. But first we discuss some automorphisms of the group of $D_8 * C_4$ and some low-dimensional Betti numbers that will go into these computations.

4.8.6 Automorphisms of $D_8 * C_4$

The elements $\rho\gamma$, σ and $\sigma\rho$ form a generating set of $D_8 * C_4$. In this section we construct an automorphism of $D_8 * C_4$ that allows us to identify the LHSS's of $A_i \rightarrow D_8 * C_4 \rightarrow C_2$ for all i .

4.8.4 Proposition *Every permutation of the generators $\rho\gamma$, σ and $\sigma\rho$ extends to an automorphism of $D_8 * C_4$.*

PROOF Number the 16 elements of $D_8 * C_4$ as follows:

1:	e	9:	γ
2:	ρ	10:	$\rho\gamma$
3:	ρ^2	11:	$\rho^2\gamma$
4:	ρ^3	12:	$\rho^3\gamma$
5:	σ	13:	$\sigma\gamma$
6:	$\sigma\rho$	14:	$\sigma\rho\gamma$
7:	$\sigma\rho^2$	15:	$\sigma\rho^2\gamma$
8:	$\sigma\rho^3$	16:	$\sigma\rho^3\gamma$

This numbering induces an inclusion $i: D_8 * C_4 \rightarrow \Sigma_{16}$ into the symmetric group on 16 elements, which on the generators given above is given by

$$\begin{aligned} \rho\gamma &\mapsto (1, 10)(2, 11)(3, 12)(4, 9)(5, 16)(6, 13)(7, 14)(8, 15), \\ \sigma &\mapsto (1, 5)(2, 6)(3, 7)(4, 8)(9, 13)(10, 14)(11, 15)(12, 16), \\ \sigma\rho &\mapsto (1, 6)(2, 7)(3, 8)(4, 5)(9, 14)(10, 15)(11, 16)(12, 13). \end{aligned}$$

One easily checks that the element of Σ_{16} given by

$$(5, 6)(2, 4)(8, 7)(14, 15)(11, 9)(13, 16)$$

induces an inner automorphism ϕ of Σ_{16} which interchanges $i(\sigma)$ and $i(\sigma\rho)$ and leaves $i(\rho\gamma)$ fixed. Identifying $D_8 * C_4$ with its image in Σ_{16} and restricting $\phi|_{\text{Im}(i)}$ yields an automorphism of $D_8 * C_4$ which interchanges σ and $\sigma\rho$, and leaves $\rho\gamma$ fixed.

Likewise, the element of Σ_{16} given by

$$(5, 10)(14, 16)(12, 7)(4, 15)(9, 11)(13, 2)$$

yields an automorphism of $D_8 * C_4$ that interchanges σ and $\rho\gamma$, and leaves $\sigma\rho$ fixed.

The two automorphisms we constructed restrict to two generating transpositions on the set $\{\rho\gamma, \sigma, \sigma\rho\}$, and hence we are done. \square

Using the automorphisms constructed in the proof of Proposition 4.8.4, we can show the following.

4.8.5 Proposition *There is an automorphism of $D_8 * C_4$ that cyclically permutes the elements of the set $\{\rho\gamma, \sigma, \sigma\rho\}$ and fixes γ . In particular this automorphism permutes the subgroups in $\{A_1, A_2, A_3\}$ cyclically.*

PROOF The existence of an automorphism that cyclically permutes the elements of $\{\rho\gamma, \sigma, \sigma\rho\}$ immediately follows from Proposition 4.8.4. For the statement about γ , we note that $\gamma = (\sigma \cdot \sigma\rho)^3 \cdot \rho\gamma$, and a direct computation shows that the effect of any transposition on γ is given by

$$\begin{aligned} (\sigma\rho \cdot \sigma)^3 \cdot \rho\gamma &= \gamma^{-1}, \\ (\rho\gamma \cdot \sigma\rho)^3 \cdot \sigma &= \gamma^{-1}, \\ (\sigma \cdot \rho\gamma)^3 \cdot \sigma\rho &= \gamma^{-1}. \end{aligned}$$

Hence the cyclic permutation of 3 elements, being the composite of two transpositions, fixes γ . \square

4.8.7 Low dimensional Betti numbers

We will denote the dimension of $H^i(BD_8 * C_4; \mathbf{F}_2)$, i.e. the i -th Betti number, by h^i . In the subsequent computations, we will freely use the following.

4.8.6 Proposition *The Betti numbers h^i for $0 \leq i \leq 4$ are given by the following table:*

i	0	1	2	3	4
h^i	1	3	5	6	7

Table 4.3: The low dimensional Betti numbers h^i .

PROOF For $i = 0$ this is clear, and for $i = 1$ we observe that $(D_8 * C_4)^{\text{ab}} = D_8 * C_4 / \langle \rho^2 \rangle \cong C_2^{\times 3}$.

The statements for $i \geq 2$ can, for instance, be verified using computer algebra software such as GAP.

4.8.8 The LHSSS for $C \rightarrow D_8 * C_4 \rightarrow C_2 \times C_2$

We now compute the LHSSS for the central extension

$$(4.12) \quad C_4 \cong \overset{\gamma}{C} \rightarrow D_8 * C_4 \rightarrow \overset{\bar{\sigma}}{C}_2 \times \overset{\bar{\sigma}\rho}{C}_2,$$

where the elements above the groups are generators. We introduce the following notation for the elements of $H^*(BC_2 \times C_2)$:

$$\begin{aligned} a &= \delta_{\bar{\sigma}}, \\ b &= \delta_{\bar{\sigma}\rho}, \end{aligned}$$

and of $H^*(BC)$:

$$\begin{aligned} x &= \delta_\gamma, \\ \beta_2 x &= \beta_2 \delta_\gamma, \end{aligned}$$

where β_2 denotes the second order Bockstein. Since the extension (4.12) is central, the local coefficient system of the associated LHSSS is trivial, and we get

$$E_2^{*,*} = \mathbf{F}_2[a, b, x, \beta_2 x]/(x^2),$$

with (s, t) -degrees given by $|a| = |b| = (1, 0)$, $|x| = (0, 1)$ and $|\beta_2 x| = (0, 2)$. See Figure 4.15 for a depiction of the E_3 -page, which up to differentials will turn out to be isomorphic to the E_2 -page, because there are no non-zero differentials on E_2 , as we will see in the next section.

Differentials on E_2 and E_3

The classes a and b are permanent cycles for degree reasons, and x is a permanent cycle because $h^1 = 3$. Since the 2-stem of the E_2 -page has 6 classes, but $h^2 = 5$, $\beta_2 x$ will have to support a differential, which for degree reasons will be either a d_2 or a d_3 . There are finitely many possibilities, namely:

$$\begin{aligned} d_2(\beta_2 x) &\in \mathbf{F}_2\{a^2 x, abx, b^2 x\}, \\ d_3(\beta_2 x) &\in \mathbf{F}_2\{a^3, a^2 b, ab^2, b^3\}. \end{aligned}$$

We first apply naturality to the map of central extensions

$$\begin{array}{ccccc} \tilde{C} & \longrightarrow & \tilde{C}_4 \times \tilde{C}_2 & \longrightarrow & \bar{\tilde{C}}_2 \\ \parallel & & \downarrow & & \downarrow \\ \tilde{C} & \longrightarrow & D_8 * C_4 & \longrightarrow & \bar{\tilde{C}}_2 \times \bar{\tilde{C}}_2 \end{array}$$

The rightmost vertical map is given by $\bar{\rho} \mapsto \bar{\sigma} \cdot \bar{\sigma}\bar{\rho}$, and therefore on cohomology by $a = \delta_{\bar{\sigma}} \mapsto \delta_{\bar{\rho}}$ and $b = \delta_{\bar{\sigma}\bar{\rho}} \mapsto \delta_{\bar{\rho}}$. The LHSSS of the top row collapses at E_2 by the Künneth theorem. Therefore $d_2(\beta_2 x)$ and $d_3(\beta_2 x)$, whatever they are, will have to map to 0 in the LHSSS of the top row. This cuts down the possibilities to

$$(4.13) \quad \begin{aligned} d_2(\beta_2 x) &\in \mathbf{F}_2\{(a^2 + b^2)x\}, \\ d_3(\beta_2 x) &\in \mathbf{F}_2\{a^3 + b^3, a^2 b + ab^2\}. \end{aligned}$$

Applying naturality again, this time to the map of central extensions

$$\begin{array}{ccccc} \tilde{C} & \longrightarrow & \tilde{C}_4 \times \tilde{C}_2 & \longrightarrow & \bar{\tilde{C}}_2 \\ \parallel & & \downarrow & & \downarrow \\ \tilde{C} & \longrightarrow & D_8 * C_4 & \longrightarrow & \bar{\tilde{C}}_2 \times \bar{\tilde{C}}_2 \end{array},$$

for which the right hand vertical map is given by $\bar{\sigma} \mapsto \bar{\sigma}$, hence on cohomology classes by $a = \delta_{\bar{\sigma}} \mapsto \delta_{\bar{\sigma}}$ and $b = \delta_{\bar{\sigma}^2} \mapsto 0$. Again, the LHSSS of the top row collapses at E_2 . Combined with (4.13) this shows that $d_2(\beta_2 x) = 0$, and that

$$d_3(\beta_2 x) \in \mathbf{F}_2\{a^2 b + ab^2\}.$$

Since $\beta_2 x$ has to support some differential, and we have determined the only possible non-zero differential, we arrive at

$$(4.14) \quad d_3(\beta_2 x) = a^2 b + ab^2.$$

The $E_4 = E_\infty$ -page

The fact that $d_2 = 0$ implies $E_3 \cong E_2$, and the differential (4.14) shows that the E_3 -page is given by




Figure 4.15: The E_3 -page.

Therefore the E_4 -page is given by

$$E_4 \cong \mathbf{F}_2[a, b, x, [(\beta_2 x)^2]] / (x^2, a^2 b + ab^2).$$

The E_4 -page has 7 classes in the 4-stem, and since $h^4 = 7$ (Proposition 4.8.6), the class $[(\beta_2 x)^2]$ is a permanent cycle. Alternatively, one can apply Kudo's transgression theorem: the class $\beta_2 x$ transgresses with $d_3(\beta_2 x) = a^2 b + ab^2$. This implies that the class $[(\beta_2 x)^2] = [\text{Sq}^2(\beta_2 x)]$ also transgresses, i.e., $d_3([(\beta_2 x)^2]) = 0$ and $d_4([(\beta_2 x)^2]) = 0$, and $d_5([(\beta_2 x)^2]) = [\text{Sq}^2(a^2 b + ab^2)]$. Applying the Cartan formula several times shows

$$\begin{aligned} \text{Sq}^2(a^2 b) &= \text{Sq}^2(a^2) b + \text{Sq}^1(a^2) \text{Sq}^1(b) + a \text{Sq}^2(b) \\ &= a^4 b, \end{aligned}$$

and hence

$$\text{Sq}^2(a^2b + ab^2) = a^4b + ab^4.$$

But since a and b are permanent cycles, we have

$$\begin{aligned} d_3((a^2 + ab + b^2)\beta_2x) &= (a^2 + ab + b^2)(a^2b + ab^2) \\ &= a^4b + ab^4, \end{aligned}$$

so $d_5([\beta_2x]^2) = 0$ also. In any case, $[(\beta_2x)^2]$ is a permanent cycle and the spectral sequence collapses at E_4 .

4.8.9 The LHSSS for $A_i \rightarrow D_8 * C_4 \rightarrow C_2$

We now compute the LHSSS for the extensions

$$A_i \rightarrow D_8 * C_4 \rightarrow C_2.$$

By Proposition 4.8.5, we can restrict to the case $i = 1$, which is the LHSSS of the extension

$$\overset{\gamma}{C}_4 \times \overset{\rho\gamma}{C}_2 \rightarrow D_8 * C_4 \rightarrow \overset{\bar{\sigma}}{C}_2.$$

We introduce the following notation for the following element of $H^*(BC_2)$:

$$c_1 = \delta_{\bar{\sigma}},$$

and for the elements of $H^*(BC_4 \times C_2)$:

$$\begin{aligned} y_1 &= \delta_{\gamma}, \\ \beta_2 y_1 &= \beta_2 \delta_{\gamma}, \\ z_1 &= \delta_{\rho\gamma}, \end{aligned}$$

where β_2 denotes the second order Bockstein.

The local coefficient system

Since

$$\begin{aligned} \sigma\gamma\sigma^{-1} &= \gamma, \\ \sigma(\rho\gamma)\sigma^{-1} &= \gamma^2 \cdot \rho\gamma, \end{aligned}$$

(4.15)

we have that the action of C_2 on $H^1(BC_4 \times C_4; \mathbf{F}_2)$ is determined by

$$\begin{aligned} y_1 &\mapsto y_1, \\ z_1 &\mapsto z_1. \end{aligned}$$

To calculate the action on $\beta_2 y_1$, we note that the second order Bockstein is natural, but only well-defined up to the image of the first order Bockstein, which in degree 2 is given by the squares of the elements in degree 1, i.e.,

$\mathbf{F}_2\{z_1^2\}$. Therefore $\sigma \cdot \beta_2 y_1 \in \{\beta_2 y_1, \beta_2 y_1 + z_1^2\}$. To determine which of the two possibilities we have, we apply naturality to the inclusion of groups

$$(4.16) \quad C_2 \times C_2 \xrightarrow{\rho\gamma} C_4 \times C_2$$

which on cohomology is given by

$$\begin{aligned} y_1 &= \delta_\gamma \mapsto 0, \\ z_1 &= \delta_{\rho\gamma} \mapsto \delta_{\rho\gamma}, \\ \beta_2 y_1 &\mapsto \delta_\gamma^2. \end{aligned}$$

The automorphism (4.15) on $C_4 \times C_2$ restricts along the inclusion (4.16) to the automorphism of $C_2 \times C_2$ which on elements is given by

$$\begin{aligned} \gamma &\mapsto \gamma, \\ \rho\gamma &\mapsto \gamma^2 \cdot \rho\gamma, \end{aligned}$$

and therefore on the cohomology classes by

$$(4.17) \quad \begin{aligned} \delta_{\gamma^2}^2 &\mapsto \delta_{\gamma^2}^2 + \delta_{\rho\gamma}^2, \\ \delta_{\rho\gamma} &\mapsto \delta_{\rho\gamma}. \end{aligned}$$

This implies, by naturality, that $\sigma \cdot \beta_2 y_1$ must be class of $H^*(BC_4 \times C_2; \mathbf{F}_2)$ that restricts via (4.16) to a class with a non-zero $\delta_{\rho\gamma}^2$ -coefficient in $H^*(BC_2 \times C_2; \mathbf{F}_2)$. But there is only one such class in $\{\beta_2 y_1, \beta_2 y_1 + z_1^2\}$ by (4.17), namely $\beta_2 y_1 + z_1^2$ and therefore

$$\sigma \cdot \beta_2 y_1 = \beta_2 y_1 + z_1^2.$$

The E_2 -page

We have $H^*(BC_4 \times C_2) \cong \mathbf{F}_2[y_1, z_1, \beta_2 y_1]/(y_1^2)$, which has a $C_2 \cong \langle \bar{\sigma} \rangle$ -action which we determined in the previous subsection: y_1 and z_1 are fixed and $\beta_2 y_1 \mapsto \beta_2 y_1 + z_1^2$. The E_2 -page of the LHSSS is given by the group cohomology of C_2 with these coefficients.

The C_2 -module $H^*(BC_4 \times C_2)$ splits as

$$\mathbf{F}_2[y_1, z_1, \beta_2 y_1]/(y_1^2) = \mathbf{F}_2[z_1, \beta_2 y_1] \oplus \mathbf{F}_2[z_1, \beta_2 y_1]\{y_1\}$$

(as C_2 -modules). The C_2 -module $\mathbf{F}_2[z_1, \beta_2 y_1]$ splits further as

$$(4.18) \quad \mathbf{F}_2[z_1^2, \beta_2 y_1] \oplus \mathbf{F}_2[z_1^2, \beta_2 y_1]\{z_1\}$$

(as C_2 -modules). This splitting enables the computation of the C_2 -cohomology with coefficients in $\mathbf{F}_2[y_1, z_1, \beta_2 y_1]/(y_1^2)$ by computing the C_2 -cohomology with coefficients in the C_2 -module (4.18).

The C_2 -cohomology $H^*(BC_2; M)$ with coefficients in any C_2 -module M is in degree $* = 0$ given by the C_2 -invariants of M , and in degrees $* \geq 1$ by the C_2 -invariants of M modulo the C_2 -coinvariants of M , as can be seen from mapping the standard minimal free $\mathbf{F}_2[C_2]$ -resolution of \mathbf{F}_2 into M .

The C_2 -module $\mathbf{F}_2[z_1^2, \beta_2 y_1]$, equals $\mathbf{F}_2[z_1^2 + \beta_2 y_1, \beta_2 y_1]$. On this latter polynomial algebra, C_2 acts by interchanging the polynomial generators. The invariants of this action are well-known and are given by

$$\mathbf{F}_2[z_1^2, (z_1^2 + \beta_2 y_1)\beta_2 y_1].$$

We introduce the notation

$$q_1 = (z_1^2 + \beta_2 y_1)\beta_2 y_1,$$

so that the invariants are given by $\mathbf{F}_2[z_1^2, q_1]$, and the invariants modulo the coinvariants are given by $\mathbf{F}_2[q_1]$.

Assembling back the splitting gives the invariants

$$\mathbf{F}_2[y_1, z_1, q_1]/(y_1^2),$$

and the invariants modulo coinvariants

$$\mathbf{F}_2[y_1, z_1, q_1]/(y_1^2, z_1^2).$$

Assembling all the lines together then gives as the E_2 -page (recall that $c_1 = \delta_{\bar{\sigma}}$):

$$E_2 = \mathbf{F}_2[c_1, y_1, z_1, q_1]/(c_1 z_1^2, y_1^2)$$

with (s, t) -degrees given by $|c_1| = (1, 0)$, $|y_1| = |z_1| = (0, 1)$ and $|q_1| = (0, 4)$. A depiction of the E_2 -page is




Figure 4.16: The $E_2 = E_\infty$ -page.

The class c_1 is a permanent cycle for degree reasons, and y_1 , z_1 and q_1 are permanent cycles because h^1 and h^4 equal to the number of classes in the 1-stem and 4-stem respectively (Proposition 4.8.6). Hence the spectral sequence collapses at E_2 .

4.8.7 Remark To get a better understanding of the multiplicative structure depicted in Figure 4.16 it can be helpful to observe that $E_2 \cong \mathbf{F}_2[y_1]/(y_1^2) \otimes \mathbf{F}_2[c_1, z_1, q_1]/(c_1 z_1^2)$ and to display only the second tensor factor:




Figure 4.17: The tensor factor $\mathbf{F}_2[c_1, z_1, q_1]/(c_1 z_1^2)$ of the E_2 -page.

4.8.10 Summary of LHSSS computations

We briefly summarize the LHSSS computations that we have carried out. The $E_2 = E_\infty$ -page of the LHSSS of the extension $A_1 \rightarrow D_8 * C_4 \rightarrow C_2$ is given by

$$E_2 = E_\infty \cong \mathbf{F}_2[c_1, y_1, z_1, q_1]/(c_1 z_1^2, y_1^2).$$

For the subgroup A_2 , we introduce the notation

$$c_2 = \delta_{\bar{\rho}}$$

for the element of $H^*(BD_8 * C_4/A_2)$ and

$$y_2 = \delta_\gamma,$$

$$z_2 = \delta_\sigma,$$

$$q_2 = (\delta_\sigma^2 + \beta_2 \delta_\gamma) \beta \delta_\gamma$$

for the elements of $H^*(BA_2)$.

For the subgroup A_3 we introduce the notation

$$c_3 = \delta_{\bar{\sigma}},$$

$$y_3 = \delta_\gamma,$$

$$z_3 = \delta_{\sigma\rho},$$

$$q_3 = (\delta_{\sigma\rho}^2 + \beta_2 \delta_\gamma) \beta_2 \delta_\gamma.$$

Then, by the automorphism of Proposition 4.8.5 the $E_2 = E_\infty$ -page of the LHSSS of the extensions $A_i \rightarrow D_8 * C_4 \rightarrow C_2$ is given by

$$E_2 = E_\infty \cong \mathbf{F}_2[c_i, y_i, z_i, q_i]/(c_i z_i^2, y_i^2)$$

with (s, t) -degrees given by $|c_i| = (1, 0)$, $|y_i| = |z_i| = (0, 1)$ and $|q_i| = (0, 4)$.

The E_2 -page of the LHSSS of the central extension $C \rightarrow D_8 * C_4 \rightarrow C_2 \times C_2$ is given by

$$E_2 \cong \mathbf{F}_2[a, b, x, \beta_2 x]/(x^2).$$

The differential $d_2 = 0$, and d_3 is generated by $d_3(\beta_2 x) = a^2 b + ab^2$. We introduce the notation $r = [(\beta_2 x)^2]$. Then the $E_4 = E_\infty$ -page is given by

$$E_4 = E_\infty \cong \mathbf{F}_2[a, b, x, r]/(x^2, a^2 b + ab^2)$$

with (s, t) -degrees given by $|a| = |b| = (1, 0)$, $|x| = (0, 1)$ and $|r| = (0, 4)$.

4.8.11 The map j^*

In this subsection we determine the effect of the map j^* in the long exact sequence (2.1), which in terms of group cohomology is given by

$$(4.19) \quad \bigoplus_{i=1}^3 H^*(BC_2; H^*(BA_i; \mathbf{F}_2)) \xrightarrow{j^*} \bigoplus_{k=1}^2 H^*(BC_2 \times C_2; H^*(BC; \mathbf{F}_2)),$$

given by the matrix

$$(4.20) \quad \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix},$$

and the inclusions $C \rightarrow A_i$ and the quotient maps $D_8 * C_4/C \rightarrow D_8 * C_4/A_i$. In (4.19) we have used that finite products and coproducts coincide in graded \mathbf{F}_2 -algebras.

4.8.8 Notation For the elements of $\bigoplus_{i=1}^3 H^*(BC_2; H^*(BA_i; \mathbf{F}_2))$ we will write $c_1 = (c_1, 0, 0)$, $y_1 = (y_1, 0, 0)$, $c_2 = (0, c_2, 0)$, etc.

For the elements of $\bigoplus_{k=1}^2 H^*(BC_2 \times C_2; H^*(BC; \mathbf{F}_2))$ we write $a_1 = (a, 0)$, $b_1 = (b, 0)$, $a_2 = (0, a)$, etc. For example, we have $a_1 + a_2 = (a, a)$.

4.8.9 Proposition *The map j from the long exact sequence (2.1) in terms of*

(4.19) is given by

$$\begin{aligned}
c_1 &\mapsto a_1 + b_1 = (a + b, 0), \\
c_2 &\mapsto b_1 + b_2 = (b, b), \\
c_3 &\mapsto a_2 = (0, a), \\
y_1 &\mapsto x_1 = (x, 0), \\
y_2 &\mapsto x_1 + x_2 = (x, x), \\
y_3 &\mapsto x_2 = (0, x), \\
z_i &\mapsto 0, \\
\beta_2 y_1 &\mapsto \beta_2 x_1 = (\beta_2 x, 0), \\
\beta_2 y_2 &\mapsto \beta_2 x_1 + \beta_2 x_2 = (\beta_2 x, \beta_2 x), \\
\beta_2 y_3 &\mapsto \beta_2 x_2 = (0, \beta_2 x), \\
q_1 &\mapsto (\beta_2 x_1)^2 = ((\beta_2 x)^2, 0), \\
q_2 &\mapsto (\beta_2 x_1)^2 + (\beta_2 x_2)^2 = ((\beta_2 x)^2, (\beta_2 x)^2), \\
(4.21) \quad q_3 &\mapsto (\beta_2 x_2)^2 = (0, (\beta_2 x)^2).
\end{aligned}$$

PROOF The map j^* in the long exact sequence (2.1) is determined by the inclusions $C \rightarrow A_i$, which is given by $\gamma \mapsto \gamma$, and the induced maps on Weyl groups $D_8 * C_4/C \rightarrow D_8 * C_4/A_i$, which for $i = 1$ is given by

$$\begin{aligned}
\bar{\sigma} &\mapsto \bar{\sigma}, \\
\bar{\sigma}\bar{\rho} &\mapsto \bar{\sigma},
\end{aligned}$$

for $i = 2$ is given by

$$\begin{aligned}
\bar{\sigma} &\mapsto \bar{e}, \\
\bar{\sigma}\bar{\rho} &\mapsto \bar{\rho},
\end{aligned}$$

and for $i = 3$ is given by

$$\begin{aligned}
\bar{\sigma} &\mapsto \bar{\sigma}, \\
\bar{\sigma}\bar{\rho} &\mapsto \bar{e}.
\end{aligned}$$

This and the matrix (4.20) shows that on cohomology classes we have

$$\begin{aligned}
c_1 &= \delta_{\bar{\sigma}} \mapsto \delta_{\bar{\sigma}_1} + \delta_{\bar{\sigma}\bar{\rho}_1} = a_1 + b_1, \\
c_2 &= \delta_{\bar{\rho}} \mapsto \delta_{\bar{\sigma}\bar{\rho}_1} + \delta_{\bar{\sigma}\bar{\rho}_2} = b_1 + b_2, \\
c_3 &= \delta_{\bar{\sigma}} \mapsto \delta_{\bar{\sigma}_2} = a_2,
\end{aligned}$$

which takes care of the first three assignments of (4.21).

For the remaining four assignments of (4.21), we observe that the restrictions

$$H^*(BA_i; \mathbf{F}_2) \rightarrow H^*(BC; \mathbf{F}_2)$$

are for all i on low dimensional cohomology classes given by

$$\begin{aligned}
y_i &= \delta_{\gamma} \mapsto \delta_{\gamma} = x, \\
z_i &\mapsto 0, \\
\beta_2 y_i &= \beta_2 \delta_{\gamma} \mapsto \beta_2 x, \\
q_i &= (z_i^2 + \beta_2 y_i) \beta_2 y_i \mapsto (\beta_2 x)^2.
\end{aligned}$$

This together with the matrix (4.20) completes the proof of the remaining assignments. \square

The B -summand

Having determined j^* , we now compute the $B = \ker(j^*)$ -summand of the short exact sequence (2.9). The degree ≥ 1 part of the domain of j^* is given by

$$\bigoplus_{i=1}^3 \mathbf{F}_2[c_i, y_i, z_i, q_i]/(c_i z_i^2, y_i^2)\{c_i\}.$$

We have that the map j^* maps

$$\begin{aligned} c_1 &\mapsto a_1 + b_1, \\ c_2 &\mapsto b_1 + b_2, \\ c_3 &\mapsto a_2, \end{aligned}$$

in particular, the c_i map to algebraically independent polynomial generators of the codomain. Therefore the kernel of j^* in degree ≥ 1 is given by the direct sum of the kernels of the 3 summands.

The classes q_i map to a polynomial generator in the codomain. The classes y_i map to a non-zero element that squares to 0, but the classes y_i themselves also square to 0. Finally, the classes z_i map to 0. Therefore the kernel of each summand is given by

$$\mathbf{F}_2[c_i, y_i, z_i, q_i]/(y_i^2, c_i z_i^2)\{c_i z_i\} = \mathbf{F}_2[c_i, y_i, q_i]/(y_i^2)\{c_i z_i\},$$

and hence the kernel of j^* is in degrees ≥ 1 given by

$$\ker(j^*)^{s \geq 1} = \bigoplus_{i=1}^3 \mathbf{F}_2[c_i, y_i, q_i]/(y_i^2)\{c_i z_i\}.$$

For the computation of the degree = 0 part of $\ker(j^*)$, we introduce the notation

$$\begin{aligned} y &= (y_1, y_2, y_3), \\ q &= (q_1, q_2, q_3), \end{aligned}$$

and recall the notation

$$\begin{aligned} z_1 &= (z_1, 0, 0), \\ z_2 &= (0, z_2, 0), \\ z_3 &= (0, 0, z_3). \end{aligned}$$

The domain of j^* in degree = 0 is given by

$$\bigoplus_{i=1}^3 \mathbf{F}_2[y_i, z_i, q_i]/(y_i^2).$$

The classes y_i map to a non-zero class that squares to 0, but the y_i also square to 0. The classes q_i map to polynomial generators, and the classes z_i map to 0. The images of the y_i satisfy the single relation

$$\sum_{i=1}^3 j^*(y_i) = 0,$$

and the images of the q_i satisfy also the single relation

$$\sum_{i=1}^3 j^*(q_i) = 0.$$

Therefore, for a class to be in the kernel of j^* , it needs to be a multiple of at least one of the following:

$$z_1, z_2, z_3, y, q,$$

which shows that the kernel in degree = 0 is given by

$$\ker(j^*)^{s=0} = \mathbf{F}_2[z_1, z_2, z_3, y, q]/(z_1z_2, z_1z_3, z_2z_3, y^2).$$

The entire kernel is depicted in the following diagram:




Figure 4.18: The B -summand, which equals $\ker(j^*)$.

The A -summand

The determination of j^* allows us to write down $\text{Im}(j^*)$. For $s \geq 1$ the image splits because the map j^* itself splits:

$$\begin{aligned} \text{Im}(j^*)^{s \geq 1} = & \mathbf{F}_2[(a + b, 0), (x, 0), ((\beta_2 x)^2, 0)] / ((x, 0)^2) \\ & \oplus \mathbf{F}_2[(b, b), (x, x), ((\beta_2 x)^2, (\beta_2 x)^2)] / ((x, x)^2) \\ & \oplus \mathbf{F}_2[(0, a), (0, x)(0, (\beta_2 x)^2)] / ((0, x)^2). \end{aligned}$$

For $s = 0$ the image is

$$\text{Im}(j^*)^{s=0} = \mathbf{F}_2[(x, 0), ((\beta_2 x)^2, 0), (0, x), (0, (\beta_2 x)^2)].$$

The cokernel $\text{coker}(j^*)$ is then given by




Figure 4.19: The cokernel of j^* , with the differentials induced from the map of spectral sequences.

The E_2 -page

Since $E_2 \cong A \oplus B$ as bigraded \mathbf{F}_2 -modules, the E_2 -page, without differentials, is given by




Figure 4.20: The E_2 -page of the \mathcal{A} -homotopy limit spectral sequence converging to the group cohomology of $D_8 * C_4$, without differentials.


4.8.12 Differentials

By Proposition 4.8.3, the \mathcal{A} -exponent of $HF_2 D_8 * C_4$ is ≤ 3 , and hence the \mathcal{A} -homotopy limit spectral sequence will collapse at E_4 with a vanishing line of height 3. Furthermore, by Corollary 4.8.2, the spectral sequence will even collapse to the $s = 0$ -line.

We will see that the horizontal vanishing line implies that the d_2 -differentials supported at the $E_2^{s,t} \cap B$ with $s > 0$ and $t \equiv 1, 2 \pmod{4}$ map injectively to $E_2^{s+t, t-1} \cap A$ (cf. Figure 4.22). In particular, all these classes in the B -summand die at E_2 . We are however not able to determine precisely which class in the B -summand hits which class in the A -summand.

We also show that the vanishing line implies that the d_2 -differentials supported at $E_2^{0,t}$ with $t \equiv 1, 2 \pmod{4}$ map surjectively to $E_2^{2, t-1} \cap A$. Using this statement about surjectivity, we are after the fact able to determine which class gets in $E_2^{0,t}$ supports a differential. We postpone this to Section 4.8.14.

The long exact sequence (2.1) respects differentials. This implies that no (red) B -summand class will hit another B -summand class, and that we have the d_3 coming from the LHSSS:

Figure 4.21: A d_3 coming from naturality.

Note however that, a priori, some of the classes involved supporting or getting hit by the d_3 in Figure 4.21 could already have died on the E_2 -page. We will see however that this is not the case: all the differentials in Figure 4.21 actually occur. To see this, observe that the the (blue) A-summand classes in the $t \equiv 0, 1 \pmod{4}$ diagonals that do not get hit by the d_3 from Figure 4.21 need to die (since the spectral sequence will collapse to the $s = 0$ -line), and the only possibility for them to do so is to get hit by (red) B-summand classes (by naturality they cannot get hit by (blue) A-summand classes).

The classes in the A-summand with degree (s, t) with $s + t \geq 5$ and $t \equiv 1 \pmod{4}$ that do not get hit by a d_3 from the A-summand can only get hit by classes from the B-summand with degrees $(s - 2, t + 1)$ or $(s - 3, t + 2)$. But there are no classes of degree $(s - 3, t + 2)$ in the B-summand satisfying this inequality and congruence, therefore these classes in the A-summand will have to get hit by a d_2 originating from the B-summand.

Now the classes in the B-summand with degree (s, t) with $s = 2$ and $t \equiv 1 \pmod{4}$ also will die. By naturality, they will not get hit by other classes from the B-summand, hence they will hit a class from the A-summand, which for degree reasons can only be a class in degree $(4, t - 1)$ with a d_2 . If such a class in the B-summand would hit the target of a d_3 in the A-summand, then the pre-image of this target in the A-summand, which lives in degree $(1, t + 1)$, which would then be a permanent cycle, contrary to the fact that the spectral sequence will collapse to the $s = 0$ -line.

For the same reason, classes in the B-summand with degree (s, t) with $s = 1$ and $t \equiv 2 \pmod{4}$ cannot hit a class of the A-summand of degree $(4, t - 2)$, for there would remain permanent cycles in the $s = 1$ -line. The only remaining possibility for these classes in the B-summand to die is to hit the three classes of the A-summand with degree $(3, t - 1)$ that do not get hit by a d_3 by a class in the A-summand.

Finally, the only remaining possibility for the classes in the B-summand with

degree (s, t) with $s = 1$ and $t \equiv 1 \pmod{4}$ to die is to hit the three classes in the A -summand with degree $(3, t - 1)$.

This pattern of differentials is expanded in Figure 4.22.




Figure 4.22: The only possible pattern of d_2 's and d_3 's originating from the classes in the B -summand with s -degree ≥ 1 . The pattern only involves non-zero d_2 -differentials.

With these d_2 's determined, we also see that all the d_3 's from Figure 4.21 need to happen.

Putting the d_2 and d_3 differentials that we've deduced so far together gives




Figure 4.23: The d_2 and d_3 differentials deduced so far.

Of course the (blue) A-summand classes with $s = 2$ also need to die, and they can only be killed by (red) B-summand classes on the $s = 0$ -line. It is an interesting question which classes in the $s = 0$ -line precisely support a differential, for this information is required if we want to have any hope of deducing the ring structure on $H^*(BD_8 * C_4)$ from this spectral sequence. We will leave this question for the next section, and for now only depict the \mathcal{A} -homotopy limit spectral sequence with all the differentials drawn:




Figure 4.24: The \mathcal{A} -homotopy limit spectral sequence converging to the \mathbf{F}_2 -group cohomology of $D_8 * C_4$.

Since there are non-zero d_3 -differentials, we have $\exp_{\mathcal{A}} \underline{H}\mathbf{F}_2_{D_8 * C_4} \geq 3$. Combined with Proposition 4.8.3, this gives

4.8.10 Proposition *The \mathcal{A} -exponent satisfies*

$$\exp_{\mathcal{A}} \underline{H}\mathbf{F}_2_{D_8 * C_4} = 3.$$

4.8.13 Poincaré series

To compute the Poincaré series, we first recall that the $s = 0$ -line of the E_2 -page was given by

$$E_2^{0,*} = \mathbf{F}_2[z_1, z_2, z_3, y, q]/(z_1 z_2, z_1 z_3, z_2 z_3, y^2).$$

Write $R = \mathbf{F}_2[y, q]/(y^2)$ for the subring of this 0-line. Then, as $\mathbf{F}_2[q]$ -modules, we have

$$R \cong \mathbf{F}_2[q]\{1, y\},$$

and therefore R has Poincaré series $(1+t)/(1-t^4)$. Furthermore, the $s = 0$ -line is, as an R -module, isomorphic to

$$E_2^{0,*} \cong R \oplus \bigoplus_{i=1}^3 R[z_i]\{z_i\}.$$

Therefore $E_2^{0,*}$ has Poincaré series

$$\begin{aligned} \frac{1+t}{1-t^4} + 3 \cdot \frac{t(1+t)}{(1-t)(1-t^4)} &= \frac{(1+t)(1-t) + 3t(1+t)}{(1-t)(1-t^4)} \\ &= \frac{1+3t+2t^2}{(1-t)(1-t^4)}. \end{aligned}$$

There is exactly one differential originating from $E_*^{0,k}$ for every $k \equiv 1, 2 \pmod{4}$, and therefore $E_\infty^{0,*}$ has Poincaré series

$$\begin{aligned} \frac{1+3t+2t^2}{(1-t)(1-t^4)} - \frac{t+t^2}{1-t^4} &= \frac{1+2t+2t^2+t^3}{(1-t)(1-t^4)} \\ &= \frac{(1+t)(1+t+t^2)}{(1-t)^2(1+t)(1+t^2)} \\ &= \frac{1+t+t^2}{(1-t)^2(1+t^2)}, \end{aligned}$$

cf. [CTVEZ03, App. C, #8(16)].

4.8.14 Differentials originating from the $s = 0$ -line

We now determine the d_2 -differentials originating from the $s = 0$ -line. These classes kill the (blue) A-summand classes with $s = 2$, which are given by $[(r^k x^\epsilon a, 0)]$ with $k \geq 0$ and $\epsilon = 0, 1$.

4.8.11 Proposition *The equations*

$$d_2(y) = d_2(z_1) = d_2(z_2) = d_2(z_3) = [(a, 0)]$$

completely describe the differential $d_2: E_2^{0,1} \rightarrow E_2^{2,0}$.

PROOF We are looking for the classes that hit $[(a, 0)] \in E^{2,0}$. If the element $y = (y_1, y_2, y_3)$ were to survive to E_∞ , this would imply that $H^*(BD_8 * C_4)$ had an element that restricted to y_i on all abelian subgroups. However, the classes that do restrict to y_i for the various i are given by:

$$\begin{aligned} A_1: & \quad \{\delta_{\bar{\sigma}}, \delta_{\bar{\sigma}\bar{\rho}}\} \\ A_2: & \quad \{\delta_{\bar{\sigma}\bar{\rho}}, \delta_{\bar{\rho}\bar{\gamma}}\} \\ A_3: & \quad \{\delta_{\bar{\sigma}}, \delta_{\bar{\rho}\bar{\gamma}}\} \end{aligned}$$

Since the intersection of these sets is empty, such a class does not exist, and y will have to die, hence $d_2(y) = [(a, 0)]$.

Suppose $(z_1, 0, 0)$ did *not* support a d_2 . Then $H^*(BD_8 * C_4)$ would have a class that restricted to $\delta_{\bar{\rho}\bar{\gamma}}$ on A_1 and to 0 on A_2 and A_3 . But such a class does not exist. Therefore $d_2((z_1, 0, 0)) = [(a, 0)]$, and by symmetry (Proposition 4.8.5 and the fact that there is a unique (blue) A-summand class in $E^{2,0}$) we also get $d_2((0, z_2, 0)) = d_2((0, 0, z_3)) = [(a, 0)]$. This completes the proof. \square

4.8.12 Proposition *The equations*

$$d_2(z_1y) = d_2(z_2y) = d_2(z_3y) = [(xa, 0)]$$

completely describe differential $d_2: E_2^{0,2} \rightarrow E^{2,1}$.

PROOF We know that the class $[(xa, 0)]$ dies. Squares cannot support d_2 -differentials, hence some z_iy , and therefore all z_iy by Proposition 4.8.5, must hit $[(xa, 0)]$. \square

4.8.13 Remark The Leibniz rule together with the proof of Proposition 4.8.12 shows that we have at E_2 the relation

$$y[(a, 0)] + z_i[(a, 0)] = [(xa, 0)]$$

for all i .

4.8.14 Lemma *For all $k \geq 0$, $l \geq 3$, $\epsilon = 0, 1$ and $i = 1, 2, 3$ we have*

$$d_2(q^k z_i^l y^\epsilon) = 0$$

PROOF The class q is a permanent cycle, since there are no differentials originating from $E_*^{0,4}$, by our computations in Section 4.8.12. First observe that

$$d_2(z_i^3) = z_i^2 dz_i = 0,$$

because there is no differential originating from $E_*^{0,3}$, by our computations in Section 4.8.12. This then implies for $l \geq 3$ that

$$d_2(q^k z_i^l) = q^k z_i^{l-1} dz = 0.$$

Likewise, because there are no differentials originating from $E_2^{0,3}$, by our computations in Section 4.8.12, we have

$$d_2(z_i^2 y) = z_i^2 dy = 0,$$

and because there are no differentials originating from $E_2^{0,4}$ either, we have

$$d_2(z_i^3 y) = y z_i^2 dz_i + z_i^3 dy = 0.$$

This then shows for $k \geq 0, l \geq 3$ that

$$d_2(q^k z_i^l y) = 0. \quad \square$$

4.8.15 Proposition *The equations*

$$d_2(q^k z_i) = d_2(q^k y) = [(r^k a, 0)]$$

completely describe the differentials $d_2: E_2^{0,4k+1} \rightarrow E_2^{2,4k}$.

PROOF By Lemma 4.8.14, the only classes that possibly can support a non-zero differential are the $q^k z_i$ and $q^k y$. We also know that there must be *some* non-zero differential. By Proposition 4.8.11, we have

$$\begin{aligned} d_2(q^k z_i) &= q^k [(a, 0)], \\ d_2(q^k y) &= q^k [(a, 0)]. \end{aligned}$$

Since at least one of these differentials must be non-zero, they both are (since they hit the same class). \square

4.8.16 Remark The proof of Proposition 4.8.15 shows that we have at E_2 the relation

$$q^k [(a, 0)] = [(r^k a, 0)].$$

In a completely similar manner we have

4.8.17 Proposition *The equations*

$$d_2(q^k z_i y) = [(r^k x a, 0)]$$

completely describe the differentials $d_2: E_2^{0,4k+2} \rightarrow E_2^{2,4k+1}$.

PROOF There is at least some class in $E_2^{0,4k+2}$ that kills $[(r^k x a, 0)]$. By Lemma 4.8.14, the only candidates are the $q^k z_i y$. Since for all i we have

$$d_2(q^k z_i y) = q^k [(x a, 0)],$$

we must have that $q^k [(x a, 0)] = [(r^k x a, 0)]$, since that is the only possibility for the latter class to die. \square

4.8.18 Remark The proof of Proposition 4.8.17 shows that we have at E_2 the relation

$$q^k[(xa, 0)] = [(r^k xa, 0)].$$

4.8.15 Exponents for various families

In this subsection we will determine upper bounds on the \mathcal{F} -exponent of $\underline{HF}_2_{D_8 * C_4}$ for two different families, one of which is $\mathcal{E}_{(2)}$.

The group $D_8 * C_4$ has three subgroups isomorphic to D_8 : $\langle \sigma, \sigma\rho \rangle$, $\langle \sigma\rho, \rho\gamma \rangle$, and $\langle \rho\gamma, \sigma \rangle$, which are all normal. Denote them by H_1 , H_2 and H_3 . Denote by τ_i the real 1-dimensional representation of $D_8 * C_4$ obtained from pulling back the sign representation along the quotient map $D_8 * C_4 \rightarrow D_8 * C_4 / H_i \cong C_2$, and let e_{τ_i} be the corresponding Euler classes. Let e be the Euler class of $\bigoplus_{i=1}^3 \tau_i$.

4.8.19 Lemma *The $\bigcup_{i=1}^3 \mathcal{A}ll_{H_i}$ -exponent satisfies*

$$\exp_{\bigcup_{i=1}^3 \mathcal{A}ll_{H_i}} \underline{HF}_2 \leq 3.$$

PROOF The Euler classes e_{τ_i} are all distinct, because $\text{res}_{H_i}^{D_8 * C_4} e_{\tau_j} = 0$ if and only if $i = j$. They are one dimensional cohomology classes, hence detected on the E_∞ -page of the \mathcal{A} -homotopy limit spectral sequence considered in the previous sections. Since the Euler classes e_{τ_i} are of cohomological degree 1, and since the \mathcal{A} -homotopy limit spectral sequence collapsed to the 0-line, they are detected in

$$(4.22) \quad E_\infty^{0,1} = \mathbf{F}\{[z_1 + y], [z_2 + y], [z_3 + y]\}.$$

The automorphism from Proposition 4.8.5 that cyclically permutes the generators $\{\sigma, \sigma\rho, \rho\gamma\}$ also cyclically permutes the groups $\{H_1, H_2, H_3\}$, since it cyclically permutes the generators of these dihedral groups. This automorphism also cyclically permutes the basis elements of (4.22).

We now consider three cases. The first case is that e_{τ_1} is detected by $[z_i + y]$ for some i . Then e_{τ_2} is detected by $[z_{i+1} + y]$ and e_{τ_3} is detected $[z_{i+2} + y]$, where the indices of the z 's are understood to be classes modulo 3. Because

$$\prod_{i=1}^3 [z_i + y] = [z_1 y + z_2 y][z_3 + y] = 0,$$

we have that e is 0 up to higher filtration, of which there is none. Hence in this case $e = 0$.

The second case is that e_{τ_1} is detected by some $[z_i + y] + [z_j + y]$ with $i \neq j$. Again the automorphism from Proposition 4.8.5 then shows that e_{τ_2} is detected by $[z_{i+1} + y] + [z_{j+1} + y]$ and e_{τ_3} is detected by $[z_{i+2} + y] + [z_{j+2} + y]$, where the indices of the z 's are understood to be classes modulo 3. Because

$$\begin{aligned} \prod_{i=1}^3 ([z_i + y] + [z_{i+1} + y]) &= \prod_{i=1}^3 [z_i + z_{i+1}] \\ &= [z_2^2][z_3 + z_1] \\ &= 0, \end{aligned}$$

we have again that e is 0 up to higher filtration, of which there is none. Hence also in this case $e = 0$.

The third and final case is that e_{τ_1} is detected by $\sum_{i=1}^3 [z_i + y]$. But the latter is fixed by the automorphism of Proposition 4.8.5, hence this implies that e_{τ_1} is fixed up to higher filtration, of which there is none. This contradicts the fact that the automorphism maps $e_{\tau_1} \mapsto e_{\tau_2}$ and that the e_{τ_i} are distinct, so in fact this case does not occur.

We conclude that in any case, $e = 0$, and by Corollary 3.2.10, the result follows. \square

4.8.20 Corollary *The $\mathcal{E}_{(2)}$ -exponent satisfies $\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2D_8 * C_4} \leq 5$.*

PROOF This follows from Proposition 4.8.3, Lemma 4.8.19, $\mathcal{A} \cap \left(\bigcup_{i=1}^3 \mathcal{A} \ell_{H_i} \right) = \mathcal{E}_{(2)}$, Lemma 3.2.6, and $3 + 3 - 1 = 5$. \square

We can slightly improve the bound in Corollary 4.8.20 as follows.

4.8.21 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies $\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2D_8 * C_4} \leq 4$.*

PROOF Let

$$\sigma \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \rho \mapsto \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \gamma \mapsto \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

be the underlying 4-dimensional real representation from [Tot14, Lem. 13.5]. The following table gives, up to inverses, all elements of $D_8 * C_4$, the effect of this representation on a line $[x_1 : y_1 : x_2 : y_2]$, and the set of lines each element fixes.

g	$g[x_1 : y_1 : x_2 : y_2]$	fixes the lines in the set(s)
ρ	$[-x_2 : -y_2 : x_1 : y_1]$	\emptyset
ρ^2	$[-x_1 : -y_1 : -x_2 : -y_2]$	\emptyset
γ	$[-y_2 : x_1 : -y_2 : x_2]$	\emptyset
$\sigma\rho\gamma$	$[y_2 : -x_2 : y_1 : -x_1]$	\emptyset
$\sigma\gamma$	$[-y_1 : x_1 : y_2 : -x_2]$	\emptyset
σ	$[x_1 : y_1 : -x_2 : -y_2]$	$\{x_1 = y_1 = 0\}, \{x_2 = y_2 = 0\}$
$\sigma\rho^2$	$[-x_1 : -y_1 : x_2 : y_2]$	$\{x_1 = y_1 = 0\}, \{x_2 = y_2 = 0\}$
$\sigma\rho$	$[-x_2 : -y_2 : -x_1 : -y_1]$	$\{x_1 = x_2, y_1 = y_2\}, \{x_1 = -x_2, y_1 = -y_2\}$
$\sigma\rho^3$	$[x_2 : y_2 : x_1 : y_1]$	$\{x_1 = x_2, y_1 = y_2\}, \{x_1 = -x_2, y_1 = -y_2\}$
$\rho\gamma$	$[y_2 : -x_2 : -y_1 : x_1]$	$\{x_1 = y_2, y_1 = -x_2\}, \{x_1 = -y_2, y_1 = x_2\}$
$\rho^3\gamma$	$[-y_2 : x_1 : y_2 : -x_2]$	$\{x_1 = y_2, y_1 = -x_2\}, \{x_1 = -y_2, y_1 = x_2\}$

Table 4.4: The elements of $D_8 * C_4$, their effect on a line in the above representation, and the lines fixed by each element.

Write

$$\begin{aligned} L_1 &= \{x_1 = y_1 = 0\} \cup \{x_2 = y_2 = 0\}, \\ L_2 &= \{x_1 = -x_2, y_2 = -y_1\} \cup \{x_1 = x_2, y_1 = y_2\}, \\ L_3 &= \{x_1 = y_2 = y_1 = -x_2\} \cup \{x_1 = -y_2 = y_1 = x_2\}. \end{aligned}$$

Then the L_i , together with the set of all lines, are the sets of lines that occur as the lines fixed by an element of $D_8 * C_4$. We immediately see from Table 4.4 that the lines in L_1 are fixed by $\langle \sigma, \sigma\rho^2 \rangle$, the lines in L_2 are fixed by $\langle \sigma\rho, \sigma\rho^3 \rangle$, and the lines in L_3 are fixed by $\langle \rho\gamma, \rho\gamma^3 \rangle$. Moreover, the L_i are pairwise disjoint. To see this, note that a line in $L_1 \cap L_2$ would be fixed by both σ and $\sigma\rho$, hence by their product $\sigma \cdot \sigma\rho = \rho$. But ρ does not fix any line, by Table 4.4. Similarly, a line in $L_2 \cap L_3$ would be fixed by both $\sigma\rho$ and $\rho^3\gamma$, but $\sigma\rho \cdot \rho^3\gamma = \sigma\gamma$ does not fix any line. Lastly, a line in $L_3 \cap L_1$ would be fixed by $\sigma \cdot \rho\gamma$, which does not fix any line. Therefore, no line is fixed by a strict supergroup of $\langle \sigma, \sigma\rho^2 \rangle$, $\langle \sigma\rho, \sigma\rho^3 \rangle$, $\langle \rho\gamma, \rho^3\gamma \rangle$, which are therefore the maximal isotropy groups of a line. All these three groups are elementary abelian, so the result follows from Proposition 3.3.3. \square

We can use the previous proposition to determine the $\mathcal{E}_{(2)}$ -exponent.

4.8.22 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_2 = 4.$$

PROOF The upper bound is Proposition 4.8.21, the lower bound follows from the fact that $D_8 * C_4$ has a subgroup isomorphic to the quaternion group Q_8 of order 8, generated by $\langle \rho, \sigma\gamma \rangle$. Now Proposition 4.5.4 and Lemma 3.2.4 imply the desired lower bound. \square

4.9 $C_4 \rtimes C_4$

Let r and s generate two copies of C_4 : $\langle r \mid r^4 = e \rangle$, $\langle s \mid s^4 = e \rangle \cong C_4$. Let $\langle s \rangle$ act on $\langle r \rangle$ by $s \cdot r = r^{-1}$. Then $C_4 \rtimes C_4$ is defined to be the semi-direct product $\langle r \rangle \rtimes \langle s \rangle$. A presentation is $C_4 \rtimes C_4 = \langle r, s \mid r^4 = s^4 = e, srs^{-1} = r^3 \rangle$. We will compute the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BC_4 \rtimes C_4; \mathbf{F}_2)$ which is isomorphic to ([CTVEZ03, App. C, #10(16)])

$$\mathbf{F}_2[z, y, x, w] / (z^2 + y^2, zy).$$

with degrees $|z| = |y| = 1$, $|x| = |w| = 2$.

4.9.1 An upper bound on the exponent

In this section we show the following upper bound on the $\mathcal{E}_{(2)}$ -exponent:

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2C_4 \rtimes C_4} \leq 8.$$

4.9.1 Lemma *For \mathcal{P} the family of proper subgroups of $C_4 \rtimes C_4$, we have the upper bound:*

$$\exp_{\mathcal{P}} \underline{HF}_{2C_4 \rtimes C_4} \leq 4.$$

PROOF Consider the linear action of $C_4 \rtimes C_4$ on \mathbf{R}^4 from [Tot14, Lem. 13.7] given by

$$s \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad r \mapsto \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We compute the lines fixed by this action. A group element g fixes a line L if and only if g^{-1} fixes L . Furthermore, the central element $r^2 s^2$ fixes all lines. Hence the table below suffices to determine the isotropy groups of all lines:

g	$g[x_1 : y_1 : x_2 : y_2]$	fixes the lines in the set(s)
r	$[-y_1 : x_1 : x_2 : y_2]$	$\{x_1 = y_1 = 0\}$
r^2	$[-x_1 : -y_1 : x_2 : y_2]$	$\{x_1 = y_1 = 0\}, \{x_2 = y_2 = 0\}$
s	$[x_1 : -y_1 : -y_2 : x_2]$	$\{x_2 = y_2 = x_1 = 0\}, \{x_2 = y_2 = y_1 = 0\}$
rs	$[y_1 : x_1 : -y_2 : x_2]$	$\{x_2 = y_2 = x_1 - y_1 = 0\}, \{x_2 = y_2 = x_1 + y_1 = 0\}$
$r^2 s$	$[-x_1 : y_1 : -y_2 : x_2]$	$\{x_2 = y_2 = x_1 = 0\}, \{x_2 = y_2 = y_1 = 0\}$
$r^3 s$	$[-y_1 : -x_1 : -y_2 : x_2]$	$\{x_2 = y_2 = x_1 - y_1 = 0\}, \{x_2 = y_2 = x_1 + y_1 = 0\}$
s^2	$[x_1 : y_1 : -x_2 : -y_2]$	$\{x_1 = y_1 = 0\}, \{x_2 = y_2 = 0\}$
rs^2	$[-y_1 : x_1 : -x_2 : -y_2]$	$\{x_1 = y_1 = 0\}$

From this table we read of that the maximal isotropy groups are:

1. The group $\langle r, s^2 \rangle \cong C_4 \times C_2$, fixing the lines in the set $\{x_1 = y_1 = 0\}$.
2. The group $\langle r^2, s \rangle \cong C_2 \times C_4$ fixing the lines in the sets $\{x_2 = y_2 = x_1 = 0\}$ and $\{x_2 = y_2 = y_1 = 0\}$.
3. The group $\langle r^2, rs \rangle \cong C_2 \times C_4$, fixing the lines in the sets $\{x_2 = y_2 = x_1 - y_1 = 0\}$ and $\{x_2 = y_2 = x_1 + y_1 = 0\}$.

These are precisely the maximal proper subgroups of $C_4 \rtimes C_4$, hence the minimal family containing all the isotropy of $\mathbf{P}(\mathbf{R}^4)$ is \mathcal{P} . An application of Proposition 3.3.3 yields the desired result. \square

4.9.2 Proposition For $\mathcal{E}_{(2)}$ the family elementary abelian subgroups, we have

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2C_4 \rtimes C_4} \leq 8.$$

PROOF We apply Lemma 3.2.15 with $\mathcal{F} = \mathcal{P}$ and $\mathcal{G} = \mathcal{E}_{(2)}$. All maximal subgroups K in \mathcal{P} are isomorphic to $C_4 \times C_2$, and for those we know (Proposition 4.3.1) that $\exp_{\mathcal{E}_{(2)}} \text{Res}_K^G \underline{HF}_2 = 2$. Hence, in the notation of Lemma 3.2.15, $m := \max_{K \in \mathcal{P}} m_K = 2$. Combined with $n = \exp_{\mathcal{P}} \underline{HF}_2 = 4$ this yields by Lemma 3.2.15

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_2 \leq 4 \cdot 2 = 8. \quad \square$$

4.9.2 The spectral sequence

We will compute the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence of $H\mathbf{F}_2$.

The elements of order precisely 2 in G are r^2, s^2, r^2s^2 . Hence all elementary abelian 2-subgroups are contained in $\langle r^2, s^2 \rangle =: V_4 \cong C_2 \times C_2$, a copy of the Klein four-group, which is also the center of G . Hence, by Proposition 2.6.1, the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence reduces to the LHSS associated to the central extension

$$V_4 \rightarrow G \rightarrow \langle \bar{r}, \bar{s} \rangle,$$

We observe that G/V_4 is also isomorphic to the Klein four-group, and call this group V'_4 . Hence the spectral sequence takes the form.

$$E_2^{s,t} = H^s(BV'_4; H^t(BV_4; \mathbf{F}_2)) \Rightarrow H^{s+t}(BG; \mathbf{F}_2).$$

Since the action of V'_4 on the center is trivial the local coefficient system is also trivial. Hence the E_2 -page is isomorphic to

$$\mathbf{F}_2[\delta_{\bar{r}}, \delta_{\bar{s}}] \otimes \mathbf{F}_2[\delta_{r^2}, \delta_{s^2}],$$

with (s, t) -degrees $|\delta_{\bar{r}}| = |\delta_{\bar{s}}| = (1, 0)$, and $|\delta_{r^2}| = |\delta_{s^2}| = (0, 1)$.




Figure 4.25: E_2 -page for $C_4 \rtimes C_4$, with no differentials drawn.

To compute d_2 we consider the map of spectral sequences given by restriction to various subgroups.

First consider the restriction $\langle r \rangle \cong C_4 \hookrightarrow G$. This has one elementary 2-group, $\langle r^2 \rangle \cong C_2$, and we get

$$\begin{array}{ccccccc} 1 & \rightarrow & \underbrace{C_2}_{=\langle r^2 \rangle} & \longrightarrow & \underbrace{C_4}_{=\langle r \rangle} & \longrightarrow & \underbrace{C_2}_{=\langle \bar{r} \rangle} \rightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \rightarrow & \underbrace{V_4}_{=\langle r^2, s^2 \rangle} & \longrightarrow & G & \longrightarrow & \underbrace{C_2 \times C_2}_{=\langle \bar{r}, \bar{s} \rangle} \rightarrow 1 \end{array}$$

We already saw that the LHSS has a d_2 given by $d_2(\delta_{r^2}) = \delta_{\bar{r}}^2$ (Section 4.3.1),

where we use the same notation for dual elements in the cohomology of the subgroups. This gives the following two conclusions for d_2 of the LHSSS G :

1. $d_2\delta_{r^2}$ has a non-zero δ_r^2 -coefficient;
2. $d_2\delta_{s^2}$ has a zero δ_r^2 -coefficient.

Second we consider the restriction $\langle s \rangle \cong C_4 \hookrightarrow G$. This gives a similar diagram

$$\begin{array}{ccccccc} 1 & \rightarrow & \underbrace{C_2}_{=\langle s^2 \rangle} & \longrightarrow & \underbrace{C_4}_{=\langle s \rangle} & \longrightarrow & \underbrace{C_2}_{=\langle \bar{s} \rangle} \rightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \rightarrow & \underbrace{V_4}_{=\langle r^2, s^2 \rangle} & \longrightarrow & G & \longrightarrow & \underbrace{C_2 \times C_2}_{=\langle \bar{r}, \bar{s} \rangle} \rightarrow 1 \end{array}$$

Again by dimension and degree reasons we infer that

3. $d_2\delta_{s^2}$ has a non-zero δ_s^2 -coefficient;
4. $d_2\delta_{r^2}$ has a zero δ_s^2 -coefficient.

Finally, we consider the restriction to $\langle rs \rangle \cong C_4 \hookrightarrow G$. Then $(rs)^2 = s^2$, and we get the diagram

$$\begin{array}{ccccccc} 1 & \rightarrow & \underbrace{C_2}_{=\langle s^2 \rangle} & \longrightarrow & \underbrace{C_4}_{=\langle rs \rangle} & \longrightarrow & \underbrace{C_2}_{=\langle \bar{r}\bar{s} \rangle} \rightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \rightarrow & \underbrace{V_4}_{=\langle r^2, s^2 \rangle} & \longrightarrow & G & \longrightarrow & \underbrace{C_2 \times C_2}_{=\langle \bar{r}, \bar{s} \rangle} \rightarrow 1 \end{array}$$

Now the inclusion $C_2 \hookrightarrow C_2 \times C_2: \bar{r}\bar{s} \mapsto \bar{r}\bar{s}$ gives the restriction map in cohomology, which is given by

$$\begin{aligned} H^*(BC_2 \times C_2; \mathbf{F}_2) &\rightarrow H^*(BC_2; \mathbf{F}_2) \\ \delta_{\bar{r}} &\mapsto \delta_{\bar{r}\bar{s}} \\ \delta_{\bar{s}} &\mapsto \delta_{\bar{r}\bar{s}}. \end{aligned}$$

Again, the LHSSS supports a differential at δ_{s^2} with image $\delta_{\bar{r}\bar{s}}^2$, which implies that

5. $d_2\delta_{s^2}$ has an *odd* number of non-zero $\{\delta_r^2, \delta_{\bar{r}}\delta_{\bar{s}}, \delta_{\bar{s}}^2\}$ -coefficients.
6. $d_2\delta_{r^2}$ has an *even* number of non-zero $\{\delta_r^2, \delta_{\bar{r}}\delta_{\bar{s}}, \delta_{\bar{s}}^2\}$ -coefficients.

Taking all these things together we obtain

$$(4.23) \quad \begin{aligned} d_2\delta_{r^2} &= \delta_r^2 + \delta_{\bar{r}}\delta_{\bar{s}}, \\ d_2\delta_{s^2} &= \delta_{\bar{s}}^2. \end{aligned}$$




Figure 4.26: E_2 -page for $C_4 \rtimes C_4$, with the generating differentials drawn.

Therefore we get that the E_3 -page is

$$E_3 \cong \mathbf{F}_2[\delta_{\bar{r}}, \delta_{\bar{s}}]/(\delta_{\bar{r}}^2, \delta_{\bar{r}}\delta_{\bar{s}} + \delta_{\bar{r}}^2) \otimes_{\mathbf{F}_2} \mathbf{F}_2[[\delta_{r^2}^2], [\delta_{s^2}^2]].$$




Figure 4.27: E_3 -page for $C_4 \rtimes C_4$. There is a vanishing line of height 3.

This page has a vanishing line of height 3, and therefore

$$\begin{aligned} E_\infty &= E_3 \\ &= \mathbf{F}_2[\delta_{\bar{r}}, \delta_{\bar{s}}]/(\delta_{\bar{r}}^2, \delta_{\bar{r}}\delta_{\bar{s}} + \delta_{\bar{r}}^2) \otimes_{\mathbf{F}_2} \mathbf{F}_2[[\delta_{r^2}^2], [\delta_{s^2}^2]]. \end{aligned}$$

4.9.3 Poincaré series

The resulting Poincaré series of $H^*(BC_4 \rtimes C_4; \mathbf{F}_2)$ is

$$\frac{1 + 2t + t^2}{(1 - t^2)^2} = \frac{1}{(1 - t)^2},$$

(cf. [CTVEZ03, App. B and App. G, #10(16)]).

4.9.4 The exponent

Using the computation of the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence, we improve the upper bound on the $\mathcal{E}_{(2)}$ -exponent from Proposition 4.9.2 by showing that

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_{2C_4 \rtimes C_4} \leq 4.$$

4.9.3 Lemma *The Euler class e of the 1-dimensional real representation given by pulling back the sign representation along the quotient map*

$$C_4 \rtimes C_4 \rightarrow C_4 \rtimes C_4 / \langle r, s^2 \rangle \cong C_2 :$$

satisfies $e^2 = 0$.

PROOF We have the composite of quotient maps

$$\overset{r}{C_4} \rtimes \overset{s}{C_4} \rightarrow \overset{\bar{s}}{C_2} \times \overset{\bar{r}}{C_2} \rightarrow \overset{\bar{s}}{C_2},$$

the composite being the quotient-by- $\langle r, s^2 \rangle$ -map. The Euler class of the sign representation of C_2 is $\delta_{\bar{s}} \in H^1(BC_2)$. Euler classes are natural, so e is the pullback of $\delta_{\bar{s}} \in H^1(BC_2 \times C_2)$. In other words, the edge map $E_2^{1,0} \rightarrow H^1(BC_4 \rtimes C_4)$ maps $\delta_{\bar{s}} \rightarrow e$. Hence $\delta_{\bar{s}}$ detects e in the LHSSS. Since $\delta_{\bar{s}}^2 = 0$ on E_3 on account of the differentials in (4.23), e^2 is zero up to higher filtration, of which there is none. \square

4.9.4 Lemma *The Euler class e of the 1-dimensional real representation given by pulling back the sign representation along the quotient map*

$$C_4 \rtimes C_4 \rightarrow C_4 \rtimes C_4 / \langle r^2, s \rangle \cong C_2$$

satisfies $e^3 = 0$.

PROOF The proof is the same as the proof of Lemma 4.9.3, but now e is detected by $\delta_{\bar{r}}$ instead of $\delta_{\bar{s}}$. The class $\delta_{\bar{r}}^2$ is non-zero, but $\delta_{\bar{r}}^3 = 0$, hence e^3 is zero up to higher filtration, of which there is none. \square

4.9.5 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$\underline{HF}_{2C_4 \rtimes C_4} \leq 4.$$

PROOF We just have to remark that $\langle r^2, s^2 \rangle$, which is the unique maximal elementary abelian 2-subgroup of $C_4 \rtimes C_4$, is the intersection of $\langle r^2, s \rangle$ and $\langle r, s^2 \rangle$, the groups we divided out by in Lemma 4.9.3 and Lemma 4.9.4. Hence by Lemma 3.2.11 and Lemma 3.2.6, $\exp_{\mathcal{E}_{(2)}} \underline{HF}_2 \leq 2 + 3 - 1 = 4$. \square

Proposition 4.9.5 together with the vanishing line of height 3 on E_∞ implies that

4.9.6 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$3 \leq \exp_{\mathcal{E}_{(2)}} \underline{H\mathbf{F}}_2 \leq 4.$$

4.10 $(C_4 \times C_2) \overset{\psi_5}{\rtimes} C_2$

Let $C_4 = \langle x \rangle$, $C_2 = \langle y \rangle$, and recall from Definition 4.2.4 that ψ_5 was defined to be the element of $\text{Aut}(C_4 \times C_2)$ defined by $x \mapsto xy$, $y \mapsto y$. In this section we will compute the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence for the group $G = (C_4 \times C_2) \overset{\psi_5}{\rtimes} C_2$ converging to its cohomology ring, which is isomorphic to ([CTVEZ03, App. C, #9(16)])

$$\mathbf{F}_2[u, v, w, r, t]/(u^2, uv, uw, v^2r + w^2).$$

with degrees $|u| = |v| = 1$, $|w| = |r| = |t| = 2$.

Under the correspondence $a \leftrightarrow x$, $b \leftrightarrow y$ and $c \leftrightarrow z$, this group admits a presentation given by

$$G = \langle a, b, c \mid a^4 = b^2 = c^2 = e, ab = ba, bc = cb, cac^{-1} = ab \rangle.$$

Let A be the normal subgroup $\langle a^2, bc, c \rangle$ of G . The group A is the unique maximal elementary abelian 2-subgroup of G , hence by Proposition 2.6.1 the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence reduces to the LHSS obtained from the extension $A \rightarrow G \rightarrow C_2$.

The Weyl group $W_G(A) = \langle \bar{a} \rangle$ acts on the generators of A by

$$\begin{aligned} a(a^2)a^{-1} &= a^2, \\ a(bc)a^{-1} &= a(bc)a^3 = (ab)^4c = c, \\ a(c)a^{-1} &= bc. \end{aligned}$$

Writing $x = \delta_{a^2}$, $y = \delta_{bc}$, and $z = \delta_c$, we get that C_2 interchanges y and z and fixes x .

Write $\sigma_1 = y + z$, $\sigma_2 = yz$, and $b = \delta_{\bar{a}}$. Taking a C_2 -projective resolution of \mathbf{Z} and applying $\text{Hom}(-, H^*(BA))$, we get that the $s = 0$ -line of the E_2 -page is given by the invariants

$$E_2^{0,*} = \mathbf{F}[\sigma_1, \sigma_2, x],$$

and the s -lines for $s \geq 1$ are given by the invariants modulo the symmetrized classed, which is

$$E_2^{s,*} = \mathbf{F}_2[\sigma_2, x]\{b^s\},$$

which assembles to

$$E_2 = \mathbf{F}_2[b, \sigma_1, \sigma_2, x]/(b\sigma_1)$$

with (s, t) -degrees given by $|b| = (1, 0)$, $|\sigma_1| = (0, 1)$, $|\sigma_2| = (0, 2)$, $|x| = (0, 1)$.

4.10.1 Differentials

The subgroup $C_4 = \langle a \rangle$ is normal because b commutes with a and $ca^2c^{-1} = (ab)^2 = a^2$.

The map of extensions

$$(4.24) \quad \begin{array}{ccccc} \overset{a^2}{C_2} & \longrightarrow & \overset{a}{C_4} & \longrightarrow & \overset{\bar{a}}{C_2} \\ \downarrow & & \downarrow & & \parallel \\ A & \longrightarrow & G & \longrightarrow & \overset{\bar{a}}{C_2} \end{array}$$

gives a map of LHSS's. This is induced by the maps in cohomology of the left and right vertical maps, which are the inclusion $C_2 \rightarrow A$ and the corresponding quotient $C_2 \rightarrow C_2$ (which is the identity). The inclusion $C_2 \hookrightarrow A$ is on elements given by $a^2 \mapsto a^2$, hence on cohomology by

$$\begin{aligned} x = \delta_{a^2} &\mapsto \delta_{a^2}, \\ y = \delta_{bc} &\mapsto 0, \\ z = \delta_c &\mapsto 0, \end{aligned}$$

and therefore

$$\begin{aligned} \sigma_1 = y + z &\mapsto 0, \\ \sigma_2 = yz &\mapsto 0. \end{aligned}$$

The right hand vertical map $C_2 \rightarrow C_2$ of (4.24) is the identity, and hence in cohomology also given by the identity map $b = \delta_{\bar{a}} \mapsto \delta_{\bar{a}}$.

Because the differentials of the LHSS of the top row are given by $d_2(\delta_{a^2}) = \delta_{\bar{a}}^2$ (Section 4.3.1), we must have

$$\begin{aligned} d_2(x) &= b^2, \\ d_2(\sigma_1) &= 0, \\ d_2(\sigma_2) &= 0, \end{aligned}$$

Observe that $\sigma_1 x$ survives to E_3 because

$$d_2(\sigma_1 x) = \sigma_1 b^2 = 0.$$

The E_2 -page with differentials is depicted in the next picture.




Figure 4.28: The E_2 -page with differentials.

The E_3 -page is given by

$$E_3 = \mathbf{F}_2[b, \sigma_1, \sigma_2, [x^2], [\sigma_1 x]] / (b\sigma_1, b^2, [\sigma_1 x]^2 - \sigma_1^2[x^2], b[\sigma_1 x]),$$

which has a vanishing line of height 2, and therefore the spectral sequence collapses as E_3 .

4.10.2 Poincaré series

The E_2 -page of the LHSSS is $E_2 = \mathbf{F}_2[b, \sigma_1, \sigma_2, x] / (b\sigma_1)$. The polynomial ring $\mathbf{F}_2[b, \sigma_1, \sigma_2, x]$ has Poincaré series $1/((1-t)^3(1-t^2))$, the ideal $(b\sigma_1)$ in this ring, which is a free module over the ring, has therefore Poincaré series $t^2/((1-t)^3(1-t^2))$. Again because the ideal is a free module, the quotient has Poincaré series $(1-t^2)/((1-t)^3(1-t^2))$.

The classes supporting the d_2 -differential generate the sub- \mathbf{F}_2 -module of E_2 given by

$$\mathbf{F}_2[b, \sigma_2, [x^2]]\{x\},$$

and their image is

$$\mathbf{F}_2[b, \sigma_2, [x^2]]\{b^2\}.$$

The first has Poincaré series $t/((1-t)(1-t^2)^2)$, the second has Poincaré series $t^2/((1-t)(1-t^2)^2)$, and therefore $H^*(BG; \mathbf{F}_2)$ has Poincaré series

4.10.3 The exponent

Consider the quotient map in the group extension

$$A \rightarrow G \rightarrow C_2 \cong \langle \bar{a} \rangle$$

of which we just computed the LHSSS. This gives a 1-dimensional real representation of G with oriented Euler class e . Because e restricts to 0 on A , we see that e lives in filtration > 0 on E_∞ , in other words $e = \delta_{\bar{a}}$. The computation showed that $\delta_{\bar{a}}^2 = 0$, since there is nothing in higher filtration. Hence by Lemma 3.2.11, $\exp_{\mathcal{E}_{(2)}} \underline{HF}_2 \leq 2$, which together with the vanishing line of height 2 on the E_∞ -page implies

4.10.1 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_2 = 2.$$

4.11 $Q_8 \times C_2$

In this section we will compute the \mathcal{F} -homotopy limit spectral sequence converging to the group cohomology of $Q_8 \times C_2$ and using the family $\mathcal{F} = \mathcal{E}_{(2)}$ of elementary abelian subgroups. The computation suggests that there might be a better way to do this using a Künneth-type theorem for the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence for \underline{HF}_2 , but we have not been able to prove such a theorem. The computation presented here is just a straightforward adaption of the computation of the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence for Q_8 .

Let i, j be the usual elements of Q_8 , and let σ be the generator of C_2 . The group $Q_8 \times C_2$ has a maximal elementary abelian subgroup, $C_2 \times C_2 \cong \langle -1, \sigma \rangle$, which is furthermore central. Hence, by Proposition 2.6.1, the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence reduces to the LHSSS for the central extension

$$(4.25) \quad \langle -1, \sigma \rangle \cong C_2 \times C_2 \rightarrow Q_8 \times C_2 \rightarrow C_2 \times C_2 \cong \langle \bar{i}, \bar{j} \rangle.$$

Since $\langle -1, \sigma \rangle$ is central, the coefficient system of $B(-)$ applied to (4.25) is trivial. Writing $a = \delta_{\bar{i}}$, $b = \delta_{\bar{j}}$, $x = \delta_{-1}$ and $y = \delta_\sigma$, the E_2 -page of the $\mathcal{E}_{(2)}$ homotopy limit spectral sequence is therefore

$$\begin{aligned} E_2^{s,t} &\cong H^s(BC_2 \times C_2; H^t(BC_2 \times C_2)) \\ &\cong \mathbf{F}_2[a, b, x, y] \end{aligned}$$

with (s, t) -degrees given by $|a| = |b| = (1, 0)$, $|x| = |y| = (0, 1)$.

Applying naturality to the map of extensions

$$\begin{array}{ccccc} \langle -1, \sigma \rangle & \longrightarrow & Q_8 \times C_2 & \longrightarrow & \langle \bar{i}, \bar{j} \rangle \\ \downarrow & & \downarrow & & \downarrow \\ \langle \sigma \rangle & \longrightarrow & \langle \sigma \rangle & \longrightarrow & \{e\} \end{array}.$$

shows that $\delta_\sigma = y$ is a permanent cycle.

The differentials of the LHSSS for the extension

$$\langle -1 \rangle \cong C_2 \rightarrow Q_8 \rightarrow C_2 \times C_2 \cong \langle \bar{i}, \bar{j} \rangle$$

are well known (see Proposition 4.5.2). Using this, naturality, and the map of

extensions

$$\begin{array}{ccccc} \langle -1, \sigma \rangle & \longrightarrow & Q_8 \times C_2 & \longrightarrow & \langle \bar{i}, \bar{j} \rangle \\ \downarrow & & \downarrow & & \parallel \\ \langle -1 \rangle & \longrightarrow & Q_8 & \longrightarrow & \langle \bar{i}, \bar{j} \rangle \end{array}$$

shows that

$$\begin{aligned} d_2(x) &= a^2 + ab + b^2 \\ d_3([x^2]) &= a^2b + ab^2. \end{aligned}$$

Hence

$$(4.26) \quad E_4 \cong \mathbf{F}_2[a, b, [x^4], y]/(a^2 + ab + b^2, a^2b + ab^2)$$

which has a vanishing line of height 4. Therefore the spectral sequence collapses and $E_4 = E_\infty$.

4.11.1 The exponent

The above computation allows us to prove

4.11.1 Proposition *The $\mathcal{E}_{(2)}$ -exponent is*

$$\exp_{\mathcal{E}_{(2)}} \underline{H}\mathbf{F}_2_{Q_8 \times C_2} = 4.$$

PROOF Elementary abelian subgroups of Q_8 pull back to elementary abelian subgroups of $Q_8 \times C_2$ along the projection map $Q_8 \times C_2 \rightarrow Q_8$. Therefore, pulling back the 4-dimensional real representation of Q_8 whose projectivization has isotropy in $\mathcal{E}_{(2)}$ considered in [MNN15, Ex. 5.18] along the projection map gives a 4-dimensional real representation of $Q_8 \times C_2$ with projectivization with isotropy in $\mathcal{E}_{(2)}(Q_8 \times C_2)$. Hence, by Proposition 3.3.3, the exponent satisfies $\exp_{\mathcal{E}_{(2)}} \underline{H}\mathbf{F}_2_{Q_8 \times C_2} \leq 4$.

Conversely, the vanishing line of height 4 in (4.26) shows the reverse inclusion, giving the desired result. \square

4.12 $D_8 \times C_2$

4.12.1 Introduction

In this section we analyze the behaviour of the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BD_8 \times C_2; \mathbf{F}_2)$. The computation suggest, as the computation for $Q_8 \times C_2$ did, that there might be a better way to do this computation using a Künneth-type theorem for the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence for $\underline{H}\mathbf{F}_2$, but we have not been able to prove such a theorem. The computation presented here is just a straightforward adaption of the computation for D_8 .

The elementary abelian subgroups of D_8 pull back to elementary abelian subgroups of $D_8 \times C_2$ along the projection

$$D_8 \times C_2 \rightarrow D_8,$$

and since D_8 has a 2-dimensional real representation V such that the projectivization $P(V)$ has istropy in $\mathcal{E}_{(2)}$, the same is true for $D_8 \times C_2$. Hence

$$\exp_{\mathcal{E}_{(2)}} \underline{H\mathbf{F}}_2 \leq 2.$$

Therefore the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence will collapse on the E_3 -page with a vanishing line of height ≤ 2 . In fact, since the cohomology of both D_8 and C_2 is detected on elementary abelian subgroups (see, e.g., [Qui71a, Lem. 4.6]), the same is true for $D_8 \times C_2$, by the Künneth Theorem. Therefore the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence will be concentrated in the $s = 0$ -line on $E_3 = E_\infty$.

4.12.2 The decomposition

Let $D_8 = \langle \sigma, \rho \mid \sigma^2 = \rho^4 = e, \sigma\rho\sigma^{-1} = \rho^{-1} \rangle$, $C_2 = \langle u \mid u^2 = e \rangle$. The maximal elementary abelian subgroups of $D_8 \times C_2$ are $F_1 := \langle \sigma, \sigma\rho^2, u \rangle \cong C_2^{\times 3}$ and $F_2 := \langle \sigma\rho, \sigma\rho^3, u \rangle \cong C_2^{\times 3}$. Closing this set of subgroups under intersections adds the additional group

$$F_1 \cap F_2 = \langle \rho^2, u \rangle =: Z \cong C_2 \times C_2$$

We apply the decomposition of Section 2.7 with $K = Z$, and $H_i = F_i$. We therefore proceed by computing the LHSSS's $E_*^{*,*}(F_i)$ and $E_*^{*,*}(Z)$.

4.12.3 The Serre spectral sequence for the subgroups F_i

We do the case $i = 1$, the case $i = 2$ being completely similar. We consider the extension

$$(4.27) \quad \langle \sigma, \sigma\rho^2, u \rangle \cong F_1 \rightarrow D_8 \times C_2 \rightarrow C_2 \cong \langle \bar{\rho} \rangle.$$

Write $x_1 = \delta_\sigma$, $y_1 = \delta_{\sigma\rho^2}$, $z_1 = \delta_u$, and $a_1 = \delta_{\bar{\rho}}$. The extension (4.27) induces an action of C_2 on F_1 given by interchanging σ and $\sigma\rho^2$ and fixing u . Hence the $s = 0$ -row of the Serre spectral sequence is isomorphic to the invariants in x_1 and y_1 tensored with $\mathbf{F}_2[z_1]$. Writing $\sigma_{1,1} = x_1 + y_1$ and $\sigma_{2,1} = x_1 y_1$ for the first and second elementary symmetric polynomial in x_1 and y_1 , we have

$$E_2^{0,*} = \mathbf{F}_2[\sigma_{1,1}, \sigma_{2,1}] \otimes_{\mathbf{F}_2} \mathbf{F}_2[z_1].$$

The rows with $s > 0$ are given by the symmetric polynomials in x_1 and y_1 divided out by the symmetrized polynomials, i.e., for $s > 0$,

$$E_2^{s,*} = a_1^s \mathbf{F}_2[\sigma_{2,1}] \otimes_{\mathbf{F}_2} \mathbf{F}_2[z_1].$$

Combining this gives

$$E_2^{s,*} = \mathbf{F}_2[a_1, \sigma_{1,1}, \sigma_{2,1}] / (a_1 \sigma_{1,1}) \otimes_{\mathbf{F}_2} \mathbf{F}_2[z_1].$$

with (s, t) degrees given by $|a_1| = (1, 0)$, $|\sigma_{1,1}| = |z_1| = (0, 1)$, $|\sigma_{2,1}| = (0, 2)$.

For every $k \geq 0$, the k -column $\bigoplus_s E_2^{s, k-s}$ of the E_2 -page has the dimension of $H^k(BD_8 \times C_2; \mathbf{F}_2)$, as can be seen from the Künneth-formula, hence the spectral sequence collapses at E_2 .

We don't repeat the calculation for the case $i = 2$, but we do define for later use for $i = 2$ the classes $x_2 = \delta_{\sigma\rho}$, $y_2 = \delta_{\sigma\rho^3}$, $z_2 = \delta_u$, $a_2 = \delta_{\bar{\rho}}$, $\sigma_{1,2} = x_2 + y_2$ and $\sigma_{2,2} = x_2 y_2$.

4.12.4 The Serre spectral sequence for the subgroup Z

We consider the central extension

$$\langle \rho^2, u \rangle = Z \rightarrow D_8 \times C_2 \rightarrow C_2 \times C_2 = \langle \bar{\rho} \rangle.$$

Writing $a_3 = \delta_{\bar{\sigma}}$, $b_3 = \delta_{\bar{\sigma}\bar{\rho}}$, $x_3 = \delta_{\rho^2}$, $z_3 = \delta_u$, we have as the E_2 -page:

$$E_2 = \mathbf{F}_2[a_3, b_3, x_3, z_3]$$

with (s, t) -degrees given by $|a_3| = |b_3| = (1, 0)$, $|x_3| = |z_3| = (0, 1)$.

Applying naturality to the map of extensions

$$\begin{array}{ccccc} \langle \rho^2 \rangle & \longrightarrow & D_8 & \longrightarrow & C_2 \times C_2 \\ \downarrow & & \downarrow & & \downarrow \\ \langle \rho^2, u \rangle & \longrightarrow & D_8 \times C_2 & \longrightarrow & C_2 \times C_2 \end{array}$$

shows that

$$d_2(x_3) = a_3 b_3.$$

Applying naturality to the map of extensions

$$\begin{array}{ccccc} \langle \rho^2, u \rangle & \longrightarrow & D_8 \times C_2 & \longrightarrow & C_2 \times C_2 \\ \downarrow & & \downarrow & & \downarrow \\ \langle u \rangle & \longrightarrow & \langle u \rangle & \longrightarrow & \{e\} \end{array}$$

shows that z_3 is a permanent cycle. The classes a_3 and b_3 are permanent cycles for degree reasons. Hence

$$E_3 = \mathbf{F}_2[a_3, b_3, [x_3^2], z_3]/(a_3 b_3).$$

Again by naturality, we see that $[x_3^2]$ is a permanent cycle, hence $E_3 = E_\infty$.

4.12.5 The map j^*

We now compute the map j^* in the long exact sequence (2.1) as a map of spectral sequences coming from the maps of extensions

$$\begin{array}{ccccc} Z & \longrightarrow & D_8 \times C_2 & \longrightarrow & C_2 \times C_2 \\ \downarrow & & \downarrow & & \downarrow \\ F_i & \longrightarrow & D_8 \times C_2 & \longrightarrow & C_2. \end{array}$$

Hence we determine the maps $Z \rightarrow F_i$ and $C_2 \times C_2 \rightarrow C_2$ and their effect on cohomology classes.

Inclusion of the center Z in F_1

The inclusion

$$\langle \rho^2, u \rangle \rightarrow \langle \sigma, \sigma \rho^2, u \rangle$$

is on elements given by

$$\begin{aligned} \rho^2 &\mapsto \sigma \cdot \sigma \rho^2, \\ u &\mapsto u \end{aligned}$$

and therefore is in cohomology given by

$$\begin{aligned} x_3 &= \delta_{\rho^2} \leftarrow \delta_{\sigma} = x_1 \\ x_3 &= \delta_{\rho^2} \leftarrow \delta_{\sigma \rho^2} = y_1 \\ z_3 &= \delta_u \leftarrow \delta_u = z_1. \end{aligned}$$

Inclusion of the center Z in F_2

The inclusion

$$\langle \rho^2, u \rangle \rightarrow \langle \sigma \rho, \sigma \rho^3, u \rangle$$

is on elements given by

$$\begin{aligned} \rho^2 &\mapsto \sigma \rho \cdot \sigma \rho^3, \\ u &\mapsto u \end{aligned}$$

and therefore is in cohomology given by

$$\begin{aligned} x_3 &= \delta_{\rho^2} \leftarrow \delta_{\sigma \rho} = x_2 \\ x_3 &= \delta_{\rho^2} \leftarrow \delta_{\sigma \rho^3} = y_2 \\ z_3 &= \delta_u \leftarrow \delta_u = z_1. \end{aligned}$$

The quotient of $D_8 \times C_2/F_1$ by C_2

The quotient

$$D_8 \times C_2 / \langle \rho^2, u \rangle \rightarrow D_8 \times C_2 / \langle \sigma, \sigma \rho^2, u \rangle$$

is on elements given by

$$\begin{aligned} \bar{\sigma} &\mapsto \bar{e} \\ \bar{\sigma} \bar{\rho} &\mapsto \bar{\rho} \end{aligned}$$

and therefore is in cohomology given by

$$b_3 = \delta_{\bar{\sigma} \bar{\rho}} \leftarrow \delta_{\bar{\rho}} = a_1.$$

The quotient of $D_8 \times C_2/F_2$ by C_2

The quotient

$$D_8 \times C_2 / \langle \rho^2, u \rangle \rightarrow D_8 \times C_2 / \langle \sigma\rho, \sigma\rho^3, u \rangle$$

is on elements given by

$$\begin{aligned} \bar{\sigma} &\mapsto \bar{\rho} \\ \bar{\sigma}\bar{\rho} &\mapsto \bar{e} \end{aligned}$$

and therefore is in cohomology given by

$$a_3 = \delta_{\bar{\sigma}} \leftarrow \delta_{\bar{\rho}} = a_2.$$

Summary

Summarizing, in the long exact sequence (2.1), we have

$$\begin{aligned} E_2(Z) &= \mathbf{F}_2[a_3, b_3, x_3, z_3], \\ E_2(F_1) &= \mathbf{F}_2[a_1, \sigma_{1,1}, \sigma_{2,1}, z_1] / (a_1\sigma_{1,1}), \\ E_2(F_2) &= \mathbf{F}_2[a_2, \sigma_{1,2}, \sigma_{2,2}, z_2] / (a_2\sigma_{1,2}). \end{aligned}$$

The map j^* is given by

$$(4.28) \quad \begin{aligned} a_1 &\mapsto b_3, \\ \sigma_{1,2} = x_1 + y_1 &\mapsto x_3 + x_3 = 0, \\ \sigma_{2,1} &\mapsto x_3^2, \\ z_1 &\mapsto z_3, \\ a_2 &\mapsto a_3, \\ \sigma_{1,2} &\mapsto 0, \\ \sigma_{2,2} &\mapsto x_3^2, \\ z_2 &\mapsto z_3. \end{aligned}$$

The kernel of j^*

The effect (4.28) of j^* on cohomology classes shows that in s -degree $s > 0$, j^* is injective because a_1 and a_2 map to different non-nilpotent elements of $E_2(Z)$. Hence $\ker j^*$ is concentrated in s -degree equal to 0, and from (4.28) we deduce

$$(4.29) \quad \ker j^* = \mathbf{F}_2[(\sigma_{1,1}, 0), (0, \sigma_{1,2}), (\sigma_{2,1}, \sigma_{2,2}), (z_1, z_2)] / ((\sigma_{1,1}, 0)(0, \sigma_{1,2})).$$

4.12.1 Remark As already mentioned, $D_8 \times C_2$ has its \mathbf{F}_2 -cohomology detected on the elementary abelian 2-subgroups. Hence the edge map to the $s = 0$ -line of the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence, which is precisely (4.29), is injective. Moreover, the Poincaré series of (4.29) is precisely equal to the one of $H^*(BD_8 \times C_2; \mathbf{F}_2)$, hence the edge morphism is also surjective. In particular, there will be no differentials emitting from the $s = 0$ -line. We argued before that $\exp_{\mathcal{E}_{(2)}} \underline{H}\mathbf{F}_2 \leq 2$, hence all classes above the $s = 0$ -line will either support a differential or be hit by one on E_2 .

The image of j^*

The effect (4.28) of j^* on cohomology classes shows that

$$\mathrm{Im} j^* = \mathbf{F}_2\{a_3^{i_1} x_3^{2l_1} z_3^{k_1}\} \cup \{b_3^{i_2} x_3^{2l_2} z_3^{k_2}\}.$$

In other words, a class $a_3^i b_3^{i'} x_3^l z_3^k$ *not* being in the image of j^* is exactly equivalent to at least one of the following conditions being true:

1. The exponents i and i' are both > 0 .
2. The exponent $l \equiv 1 \pmod{2}$.

4.12.6 The E_2 -page

Having computed $\ker j^*$ and $\mathrm{Im} j^*$, we know $A = \mathrm{coker} j^*[1]$ and $B = \ker j^*$ in (2.9). Moreover, since ∂ is a map of spectral sequences, and the fact that on $E_2(Z)$ we have

$$d_3(x_3) = a_3 b_3,$$

shows that we also have this differential on the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence under the identification (2.9).

Therefore all classes $a_3^i b_3^{i'} x_3^l z_3^k$ in $\mathrm{coker} j^*$ with $l \equiv 1 \pmod{2}$ will support a differential:

$$(4.30) \quad d_2(a_3^i b_3^{i'} x_3^l z_3^k) = a_3^{i+1} b_3^{i'+1} x_3^{l-1} z_3^k.$$

Hence all classes with even x_3 -exponent and strictly positive a_3 and b_3 -exponent get hit, which accounts for all classes in $\mathrm{coker} j^*$: they all either hit or get hit by a d_2 . Therefore there cannot be any d_2 -differentials emitting from the B -summand, and $E_3 = B$. Since this is concentrated in the $s = 0$ -row, $E_3 = E_\infty$.

4.12.7 Multiplicative extensions

Since on E_∞ everything is concentrated in the $s = 0$ row, we have

$$\begin{aligned} B &\cong E_\infty^{0,*} \\ &\cong F^0 H^*(BD_8 \times C_2) / F^1 H^*(BD_8 \times C_2) \\ &= H^*(BD_8 \times C_2). \end{aligned}$$

4.12.8 The exponent

We now prove

4.12.2 Proposition *The $\mathcal{E}_{(2)}$ -exponent satisfies*

$$\exp_{\mathcal{E}_{(2)}} \underline{HF}_2 = 2.$$

PROOF Pulling back transitive D_8 -orbits with isotropy in elementary abelian groups along the projection map $D_8 \times C_2$ gives transitive $D_8 \times C_2$ -orbits with isotropy in elementary abelian groups. Therefore pulling back the representation

of D_8 considered in the proof of Corollary 4.4.3 along the projection map $D_8 \times C_2 \rightarrow D_8$ gives a 2-dimensional real representation of $D_8 \times C_2$ with isotropy in $\mathcal{E}_{(2)}$. Hence by Proposition 3.3.3 we have $\exp_{\mathcal{E}_{(2)}} \underline{H}\mathbf{F}_{2D_8 \times C_2} \leq 2$.

Conversely, in the above computation of the $\mathcal{E}_{(2)}$ -homotopy limit spectral sequence converging to $H^*(BD_8 \times C_2; \mathbf{F}_2)$, we saw that $E_2 \neq E_\infty$ (Equation (4.30)). Hence we also have the reverse inequality, and the result follows. \square

Chapter 5

Properties of the \mathcal{F} -homotopy limit spectral sequence for equivariant K -theory

5.1 Elementary abelian 2-groups

5.1.1 Introduction

In [MNN15, App. B], the \mathcal{C} -homotopy limit spectral sequence of KU is computed for $G = C_2^{\times 2}$. The target of this spectral sequence is $\pi_* KU \cong R(G)[\beta^\pm]$, the polynomial ring on the Bott periodicity generator β , with $|\beta| = 2$, and its inverse β^{-1} , with coefficients in the representation ring of G . The E_3 -page is the last page with differentials, hence $\exp_{\mathcal{C}} KU \geq 3$.

In this section, we will establish lower bound on the \mathcal{C} -exponent of KU for all elementary abelian 2-groups. More precisely, we will show that for $G = C_2^{\times n}$ we have $\exp_{\mathcal{C}} KU \geq n$.

The proof can be summarized as follows. Most of the work goes into describing the edge map

$$(5.1) \quad R(C_2^{\times n}) \rightarrow \lim_{\mathcal{C}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}}^0 R(-) \cong E_2^{0,0}$$

of the spectral sequence. The upshot is that the exponent (as a group) of the cokernel of (5.1) is 2^{n-1} (Corollary 5.1.8). A qualitative analysis of the E_2 -page shows that the classes in positive filtration are (at most) 4-torsion, and Bott periodicity implies $E_2^{s,t} = 0$ if $s+t$ is odd. This then implies that there are non-zero differentials on at least $\lceil (n-1)/2 \rceil$ odd pages, whence $\exp_{\mathcal{C}} KU \geq n$ (Proposition 5.1.10).

5.1.2 The exponent of the cokernel of the edge map

We will determine the exponent of the cokernel of the edge map

$$(5.2) \quad R(C_2^{\times n}) \rightarrow \lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} R(-) = E_2^{0,0}$$

of the \mathcal{C} -homotopy limit spectral sequence of KU . We will do this by finding bases of the left and right hand side that diagonalize the edge map.

Splitting of the augmentation ideal

For e the trivial subgroup, identify $R(e) \cong \mathbf{Z}$ by sending the trivial representation to 1. For every group H , the restriction map then gives a map $R(H) \rightarrow \mathbf{Z}$, the kernel of which is called the augmentation ideal of $R(H)$ and is denoted $I(H)$. The map $R(H) \rightarrow \mathbf{Z}$ is naturally split by sending $1 \in \mathbf{Z}$ to the trivial representation.

For the coefficient system $R(-)$ on the orbit category $\mathcal{O}(G)_{\mathcal{F}}$, for every $H \in \mathcal{F}$, the restriction map $R(H) \rightarrow R(e) \cong \mathbf{Z}$ gives a map $R(-) \rightarrow \underline{\mathbf{Z}}$, where $\underline{\mathbf{Z}}$ is the constant coefficient system at \mathbf{Z} . This map is naturally split, by applying the natural splitting to every individual group $H \in \mathcal{F}$.

Combining the two splittings splits the map (5.2) as a direct sum of maps

$$\mathbf{Z} \oplus I(C_2^{\times n}) \rightarrow \lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} \mathbf{Z} \oplus \lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} I(-) \cong \mathbf{Z} \oplus \lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} I(-).$$

This direct sum decomposition reduces describing (5.2) to describing

$$(5.3) \quad I(C_2^{\times n}) \rightarrow \lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} I(-).$$

A basis for $I(C_2^{\times n})$

Denote

$$C_2^{\times n} = \langle s_1, \dots, s_n \mid s_i^2 = e, s_j s_k = s_k s_j \forall i, j, k \rangle.$$

Denote by σ_j the sign representation of $\langle s_j \rangle \cong C_2$, and also of the representation of $C_2^{\times n}$ given by

$$\sigma_j(s_i) = \begin{cases} -1 & \text{if } j = i, \\ 1 & \text{otherwise.} \end{cases}$$

For $J \subset \{1, \dots, n\}$, denote

$$a_J = \prod_{j \in J} (1 - \sigma_j) \in R(C_2^{\times n}).$$

Observe that $a_J \in I(C_2^{\times n})$ if and only if $J \neq \emptyset$. In fact, more is true.

5.1.1 Proposition *The set $A := \{a_J\}_{\emptyset \neq J \subset \{1, \dots, n\}}$ is a basis of $I(C_2^{\times n})$ as a \mathbf{Z} -module.*

PROOF Since $\{1, \sigma_j\}$ is a basis of $R(\langle s_j \rangle)$ ([Ser77, Sec. 5.1]) as a \mathbf{Z} -module, the set $\{1, 1 - \sigma_j\}$ is also basis of $R(\langle s_i \rangle)$. Since $R(C_2^{\times n}) = \bigotimes_i R(\langle s_i \rangle)$ ([Ser77, Thm. 3.2.10]) the set $\{a_J\}_{J \subset \{1, \dots, n\}}$ is a basis of $R(C_2^{\times n})$. Since a_\emptyset generates the \mathbf{Z} -summand of $R(C_2^{\times n}) \cong \mathbf{Z} \oplus I(C_2^{\times n})$ and all $a_J \in I(C_2^{\times n})$ for $J \neq \emptyset$, the result follows. \square

A basis for $\lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} I(-)$

We will now describe a \mathbf{Z} -basis of $\lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} I(-)$. This limit is given by the equalizer

$$\text{eq} \left(\prod_{C \in \mathcal{C}} I(C) \rightarrow I(e) \right),$$

where the map from the product to $I(e)$ comes from the universal property of the product and the restriction maps $I(C) \rightarrow I(e)$ for every C . But since $I(e) = 0$ this is equal to $\prod_{C \in \mathcal{C}'} I(C)$, where \mathcal{C}' denotes the set (not family) of non-trivial cyclic subgroups of $C_2^{\times n}$.

5.1.2 Notation For $J \subset \{1, \dots, n\}$, denote $s_J = \prod_{j \in J} s_j \in C_2^{\times n}$. Then the elements of \mathcal{C}' are in $(1 : 1)$ -correspondence with the non-empty J via $J \leftrightarrow \langle s_J \rangle$.

Denote the sign representation of $\langle s_J \rangle$ by σ_J , and denote $1 - \sigma_J \in I(\langle s_J \rangle)$ by $\bar{\sigma}_J$. Observe that $I(\langle s_J \rangle)$ is \mathbf{Z} -module of rank 1, generated by $\bar{\sigma}_J$.

Denote the projection $\prod_{C \in \mathcal{C}'} I(C) \rightarrow I(\langle s_J \rangle)$ by pr_J , and let b'_J be the element of $\prod_{C \in \mathcal{C}'} I(C)$ satisfying $\text{pr}_K(b'_J) = \bar{\sigma}_J$ if $K = J$ and 0 otherwise.

This discussion shows

5.1.3 Proposition *The set $B' := \{b'_J \mid \emptyset \neq J \subset \{1, \dots, n\}\}$ is a basis of $\lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} I(-)$.*

The map (5.3) is not diagonal with respect to the bases A and B' . To get a diagonal matrix, denote for non-empty $J \subset \{1, \dots, n\}$

$$b_J = \sum_{J \subset K \subset \{1, \dots, n\}} b'_K.$$

5.1.4 Proposition *The set $B = \{b_J\}_{\emptyset \neq J \subset \{1, \dots, n\}}$ is a basis of $\lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}} I(-)$ as a \mathbf{Z} -module.*

PROOF First note that $B \subset \mathbf{Z}B'$. Conversely, by the principle of inclusion-exclusion,

$$b'_J = b_J - \sum_{\substack{J \subset K \\ \#(K-J)=1}} b_K + \sum_{\substack{J \subset K \\ \#(K-J)=2}} b_K - \dots,$$

hence also $B' \subset \mathbf{Z}B$. Proposition 5.1.3 and $\#B' = \#B$ now imply the desired result. \square

The edge map

We now describe the edge map (5.1), which we reduced to describing (5.3), which we will do in terms of the bases A and B .

5.1.5 Warning Although the next proposition is phrased in terms of the bases A (Proposition 5.1.1) and B (Proposition 5.1.4), the proof will also make use of the basis B' (Proposition 5.1.3).

5.1.6 Proposition *The map (5.3) is given on the bases A (Proposition 5.1.1) and B (Proposition 5.1.4) by*

$$(5.4) \quad a_J \mapsto 2^{\#J-1} b_J.$$

PROOF For σ the sign representation of a cyclic group of order 2, we have $(1 - \sigma)^2 = 2\sigma$. Hence for basis elements $b'_J, b'_K \in B'$, we have

$$(5.5) \quad b'_J b'_K = \begin{cases} 2b'_J & \text{if } J = K, \\ 0 & \text{otherwise.} \end{cases}$$

By induction therefore, we also have $(b'_J)^m = 2^{m-1} b'_J$.

First we verify (5.4) on basis elements indexed by singletons:

$$\begin{aligned} a_{\{j\}} &= 1 - \sigma_j \\ &\mapsto \sum_{j \in J \subset \{1, \dots, n\}} b'_j \\ &=: b_{\{j\}}. \end{aligned}$$

Then, by multiplicativity of the edge map, we get for all non-empty $J \subset \{1, \dots, n\}$

$$\begin{aligned} a_J &= \prod_{j \in J} (1 - \sigma_j) \\ &\mapsto \prod_{j \in J} \left(\sum_{j \in K \subset \{1, \dots, n\}} b'_K \right) \end{aligned}$$

By the product structure on $\lim_{\mathcal{O}(C_2^{\times n})^{\text{op}}} I(-)$, all terms of the sum with $K \not\subset J$ will go to zero in the product. Hence we can discard those terms:

$$= \prod_{j \in J} \left(\sum_{J \subset K \subset \{1, \dots, n\}} b'_K \right)$$

The factors of this product do not depend on the index j anymore, so we get:

$$= \left(\sum_{J \subset K \subset \{1, \dots, n\}} b'_K \right)^{\#J}$$

which by Equation (5.5) equals

$$\begin{aligned}
&= \sum_{J \subset K \subset \{1, \dots, n\}} (b'_K)^{\#J} \\
&= 2^{\#J-1} \sum_{J \subset K \subset \{1, \dots, n\}} b'_K \\
&=: 2^{\#J-1} b_J. \quad \square
\end{aligned}$$

5.1.7 Corollary *The cokernel of (5.3) (which equals the cokernel of (5.1)), is*

$$\bigoplus_{\emptyset \neq J \subset \{1, \dots, n\}} \mathbf{Z}/2^{\#J-1} \mathbf{Z}\{b_J\}.$$

5.1.8 Corollary *The exponent of the cokernel of (5.1) is 2^{n-1} .*

5.1.3 A lower bound on $\exp_{\mathcal{C}} KU$

Using Corollary 5.1.8, we now prove a lower bound on $\exp_{\mathcal{C}} KU$. We do this by analyzing the \mathcal{C} -homotopy limit spectral sequence

$$(5.6) \quad E_2^{s,t} = \lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}}^s R(-)[\beta^{\pm}] \Rightarrow R(C_2^{\times n})[\beta^{\pm}].$$

where β is the Bott periodicity generator with (s, t) -degree $|\beta| = (0, 2)$. We need the following lemma, which gives an upper bound on the torsion on the E_2 -page for $s \geq 2$.

5.1.9 Lemma *In the \mathcal{C} -homotopy limit spectral sequence (5.6), $E_2^{s,t}$ is annihilated by 4 for $s \geq 1$.*

PROOF For a family $\mathcal{F} \subset \mathcal{C}$, write $\underline{\mathbf{Z}}[\mathcal{F}]$ for the constant coefficient system $\underline{\mathbf{Z}}$ restricted to $\mathcal{O}(C_2^{\times n})_{\mathcal{F}}$ and then left Kan extended back up to $\mathcal{O}(C_2^{\times n})_{\mathcal{C}}$, as in [MNN15, App. B]. As in [MNN15, B.3], the sum of the counit maps

$$\bigoplus_{C \in \mathcal{C}} \underline{\mathbf{Z}}[\{e, C\}] \rightarrow \underline{\mathbf{Z}}$$

is surjective and yields a short exact sequence of coefficient systems

$$(5.7) \quad 0 \rightarrow \bigoplus_{|\mathcal{C}|-1} \underline{\mathbf{Z}}[\{e\}] \rightarrow \bigoplus_{C \in \mathcal{C}} \underline{\mathbf{Z}}[\{e, C\}] \rightarrow \underline{\mathbf{Z}} \rightarrow 0.$$

This induces a long exact sequence of Ext-groups as in [MNN15, eq. B.6] that we will not spell out, but we describe what we need.

The Ext-groups corresponding to the left term in (5.7) are isomorphic to a direct sum of groups of the form $H^s(BC_2^{\times n}; \mathbf{Z})$, which are 2-torsion for $s \geq 1$ by the Künneth theorem.

The Ext-groups corresponding to the middle term in (5.7) are isomorphic to a direct sum of groups of the form $H^s(BC_2^{\times(n-1)}; R(C_2))$. Here $R(C_2)$ is a

free \mathbf{Z} -module with trivial $C_2^{\times(n-1)}$ -action, hence $H^s(BC_2^{\times(n-1)}; R(C_2))$ is also 2-torsion for $s \geq 1$ by the Künneth theorem.

As in the proof of [MNN15, Thm. B.9], we obtain a short exact sequence

$$0 \rightarrow A^{s,t} \rightarrow E_2^{s,t} \rightarrow B^{s,t} \rightarrow 0$$

where A and B are subquotients of the Ext-groups corresponding to the left and middle term of the short exact sequence (5.7), but with A shifted by 1 in the positive s -direction. In particular, A and B are 2-torsion for $s \geq 2$, and hence E_2 is 4-torsion for $s \geq 2$. \square

5.1.10 Proposition For $G = C_2^{\times n}$, we have $\exp_{\mathcal{C}} KU \geq n$.

PROOF We consider the \mathcal{C} -homotopy limit spectral sequence

$$E_2^{s,t} = \lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}}^s R(-)[\beta^{\pm}] \Rightarrow R(C_2^{\times n})[\beta^{\pm}].$$

where β is the Bott periodicity generator which has (s, t) -degree $(0, 2)$. First, by Lemma 5.1.9, the E_2 -page is qualitatively (made precise after the figure) given by

		*		*		*		*	
s	*		*		*		*		*
		*		*		*		*	
2	*		*		*		*		*
		?		?		?		?	
0	\square		\square		\square		\square		\square
	-4		-2		0		2		$t - s$

Figure 5.1: The E_2 -page qualitatively.

Here qualitatively means that every \square signifies a free \mathbf{Z} -module (not necessarily of rank 1), every $*$ signifies a not necessarily free $\mathbf{Z}/4\mathbf{Z}$ -module (not necessarily of rank 1), every empty square signifies the 0 group and every question mark $?$ signifies that it could be anything.

Second, by Corollary 5.1.8, we have that the edge map

$$R(C_2^{\times n}) \xrightarrow{\widetilde{\text{res}}} \lim_{\mathcal{O}(C_2^{\times n})_{\mathcal{C}}^{\text{op}}}^0 R(-)$$

is injective and has cokernel with exponent 2^{n-1} . That is, there is an x in $E_2^{0,0}$ such that $2^{n-1}x \in \text{Im}(\widetilde{\text{res}})$ but $2^{n-2}x \notin \text{Im}(\widetilde{\text{res}})$.

Combining these two points, we see that $E_2^{0,0}$ will have to support at least differentials on $\lceil \frac{n-1}{2} \rceil$ pages, since all the groups in positive filtration ≥ 2 are 4-torsion, and there is no possibility for a class in the $s = 0$ -line to hit a question mark in the $s = 1$ -line, so taking a kernel gives at least a subgroup of index 4. The even differentials are all 0, and so we get that $d_M \neq 0$ for some $M \geq$

$2\lceil \frac{n-1}{2} \rceil + 1 \geq n$, by the pigeonhole principle. Since $\exp_{\mathcal{C}} KU \leq N$ implies that the spectral sequence degenerates at E_{N+1} ([MNN15, Prop. 2.26]), we get that $\exp_{\mathcal{C}} KU \geq n$, as desired. \square

5.1.11 Remark By distinguishing between n even and odd, the proof in fact shows

$$\exp_{\mathcal{C}} KU \geq \begin{cases} n+1 & \text{if } n \equiv 0 \pmod{2}, \\ n & \text{if } n \equiv 1 \pmod{2}. \end{cases}$$

5.1.4 Odd primes

We do not have a generalization of the structure result of the cokernel like Corollary 5.1.7 to odd primes p yet. However, experimental evidence does suggest a generalization. To describe it, we need the following combinatorial notion.

5.1.12 Definition ([And76], [Com74, Ex. I.16], [Fah12]) Define the **polynomial coefficient** $\binom{x,q}{k}$ by the coefficients of

$$(1 + t + \cdots + t^{q-1})^x = \sum_{k \geq 0} \binom{x,q}{k} t^k.$$

The coefficients for $q = 2$ are also known as binomial coefficients. Polynomial coefficients for $q = 3$ we call trinomial coefficients. For arbitrary q we call them the q -nomial coefficients.

5.1.13 Warning Polynomial coefficients are not the same thing as multinomial coefficients.

5.1.14 Conjecture *Let p be a prime. Set*

$$e_{n,k,p} = \sum_{j=0}^{p-1} \binom{n,p}{(p-1)(k+1)-j}$$

(cf. for $p = 3$ [Inc05]). Then the cokernel of the edge map $R(C_p^{\times n}) \rightarrow E_2^{0,0}$ of the \mathcal{C} -homotopy limit spectral sequence converging to $\pi_* KU$ is isomorphic to

$$\bigoplus_{k=0}^{n-1} (\mathbf{Z}/p^k)^{\oplus e_{n,k,p}}.$$

5.1.15 Remark For $p = 2$, Conjecture 5.1.14 is Corollary 5.1.7.

Experimental evidence for Conjecture 5.1.14 is given in Appendix A.

5.1.16 Corollary (of Conjecture 5.1.14) *For $G = C_p^{\times n}$ the elementary abelian p -group of rank n , we have $\exp_{\mathcal{C}} KU \geq n$.*

PROOF The same as the proof of Proposition 5.1.10. \square

5.1.5 The real case for $p = 2$

For G an elementary abelian 2-group, the complex and real representation rings of G coincide, which we use to give a lower bound on $\exp_{\mathcal{C}} KO$ using real Bott periodicity. The argument is completely analogous to the complex case.

5.1.17 Proposition *For $G = C_2^{\times n}$ the \mathcal{C} -exponent of KO satisfies*

$$\exp_{\mathcal{C}} KO \geq n.$$

PROOF As in the complex case, there need to be at least $\lceil \frac{n-1}{2} \rceil$ non-zero differentials, as again the E_2 -page is 4-torsion for $s \geq 2$. However, real Bott periodicity is 8-fold periodic with 4 non-zero groups in each period, which makes computing a lower bound on the r for which $E_r = E_{\infty}$ a bit harder as it comes down to making a case distinction on the class of n modulo 8 rather than n modulo 2 as in the complex case. Nevertheless, one can do the counting, and the result is:

$$\exp_{\mathcal{C}} KO \geq \begin{cases} 2 + 2\lceil \frac{n-1}{2} \rceil & \text{if } n \equiv 0, 1 \pmod{8}, \\ 2 + 2(\lceil \frac{n-1}{2} \rceil - 1) + 1 & \text{if } n \equiv 2, 3 \pmod{8}, \\ 2 + 2(\lceil \frac{n-1}{2} \rceil - 2) + 3 & \text{if } n \equiv 4, 5 \pmod{8}, \\ 2 + 2(\lceil \frac{n-1}{2} \rceil - 3) + 7 & \text{if } n \equiv 6, 7 \pmod{8}. \end{cases}$$

All these 4 expressions are $\geq n$. □

5.1.18 Remark The proof in fact shows

$$(5.8) \quad \exp_{\mathcal{C}} KO \geq \begin{cases} n + 2 & \text{if } n \equiv 0 \pmod{8}, \\ n + 1 & \text{if } n \equiv 1 \pmod{8}, \\ n + 1 & \text{if } n \equiv 2 \pmod{8}, \\ n & \text{if } n \equiv 3 \pmod{8}, \\ n + 1 & \text{if } n \equiv 4 \pmod{8}, \\ n & \text{if } n \equiv 5 \pmod{8}, \\ n + 3 & \text{if } n \equiv 6 \pmod{8}, \\ n + 2 & \text{if } n \equiv 7 \pmod{8}. \end{cases}$$

The fact that KU is a KO -module implies that KU is a retract of KO and therefore $\exp_{\mathcal{C}} KU \leq \exp_{\mathcal{C}} KO$. Hence the lower bounds on $\exp_{\mathcal{C}} KU$ of Remark 5.1.11 give lower bounds on $\exp_{\mathcal{C}} KO$, but the ones in (5.8) are in all cases at least as good or better.

Appendix A

Sage computations

A.1 Introduction

Let p be a prime, and consider the edge map $R(C_p^{\times n}) \rightarrow E_2^{0,0}$ of the \mathcal{C} -homotopy limit spectral sequence (cf. Section 5.1.2 and Section 5.1.4). Denote the cokernel of this map by $Q_{p,n}$. Using Sage ([S⁺15]), we have computed the isomorphism type of $Q_{p,n}$ for the primes $p = 2, 3, 5, 7, 11$ and various small values of n . The results are summarized in the following five tables. Of course, for $p = 2$ we also have Corollary 5.1.7. We also give the Sage code used for the computation. The computations provide experimental evidence for Conjecture 5.1.14.

A.2 Summary of computations

A.2.1 $p = 2$

The following table gives $Q_{2,n}$ for $1 \leq n \leq 8$. It turns out that for those values of n , $Q_{2,n}$ is the direct sum of groups of the form $(\mathbf{Z}/2^k\mathbf{Z})^{\oplus e_k}$, for some exponents e_k . The entry in the k -th row and n -th column of the table gives the exponent e_k of $\mathbf{Z}/2^k\mathbf{Z}$ in $Q_{2,n}$. An empty entry means that the exponent is zero.

	n	1	2	3	4	5	6	7	8
$\mathbf{Z}/2^k$									
$\mathbf{Z}/1$		1	2	3	4	5	6	7	8
$\mathbf{Z}/2$			1	3	6	10	15	21	28
$\mathbf{Z}/4$				1	4	10	20	35	56
$\mathbf{Z}/8$					1	5	15	35	70
$\mathbf{Z}/16$						1	6	21	56
$\mathbf{Z}/32$							1	7	28
$\mathbf{Z}/64$								1	8
$\mathbf{Z}/128$									1

Table A.1: Exponent e_k in decomposition of $Q_{2,n}$ as direct sum of groups of the form $(\mathbf{Z}/2^k\mathbf{Z})^{\oplus e_k}$.

For instance, from the table we read off that

$$Q_{2,4} \cong \mathbf{Z}/8 \oplus (\mathbf{Z}/4)^{\oplus 4} \oplus (\mathbf{Z}/2)^{\oplus 6} \oplus (\mathbf{Z}/1)^{\oplus 4}.$$

The reason that we write down those trivial groups is that $Q_{2,3}$, for instance, is the quotient of a free abelian group of rank 7. Hence in the quotient “there really are” these trivial groups. Moreover, they fit nicely in the pattern of half of Pascal’s triangle that this table has.

A.2.2 $p = 3$

Also for low values of n , $Q_{3,n}$ is isomorphic to a direct sum of groups of the form $(\mathbf{Z}/3^k)^{\oplus e_k}$ for some exponents e_k . The following table gives these exponents for $1 \leq n \leq 6$ (we go less high because the algorithm takes too long to terminate for higher n).

	n	1	2	3	4	5	6
$\mathbf{Z}/3^k$							
$\mathbf{Z}/1$		2	5	9	14	20	27
$\mathbf{Z}/3$			3	13	35	75	140
$\mathbf{Z}/9$				4	26	96	267
$\mathbf{Z}/27$					5	45	216
$\mathbf{Z}/81$						6	71
$\mathbf{Z}/243$							7

Table A.2: Exponent e_k in decomposition of $Q_{3,n}$ as direct sum of groups of the form $(\mathbf{Z}/3^k)^{\oplus e_k}$.

For example,

$$Q_{3,4} \cong (\mathbf{Z}/27)^{\oplus 5} \oplus (\mathbf{Z}/9)^{\oplus 26} \oplus (\mathbf{Z}/3)^{\oplus 35} \oplus (\mathbf{Z}/1)^{\oplus 14}$$

The pattern in the table seems to be described by pairwise sums of trinomial coefficients, cf. [Inc05].

A.2.3 $p = 5$

We write down the same table as in the previous two sections for the prime 5:

	n	1	2	3	4
$\mathbf{Z}/5^k$					
$\mathbf{Z}/1$		4	14	34	69
$\mathbf{Z}/5$			10	70	285
$\mathbf{Z}/25$				20	235
$\mathbf{Z}/125$					35

Table A.3: Exponent e_k in decomposition of $Q_{5,n}$ as a direct sum of groups of the form $(\mathbf{Z}/5^k)^{\oplus e_k}$.

The pattern of the table seems to be described by 4-term sums of 5-nomial coefficients.

A.2.4 $p = 7$

For the prime 7 the table is

$\mathbf{Z}/7^k$	n	1	2	3
$\mathbf{Z}/1$		6	27	83
$\mathbf{Z}/7$			21	203
$\mathbf{Z}/49$				56

Table A.4: Exponent e_k in decomposition of $Q_{7,n}$ as a direct sum of groups of the form $(\mathbf{Z}/7^k)^{\oplus e_k}$.

The pattern of the table seems to be described by 6-term sums of 7-nomial coefficients.

A.2.5 $p = 11$

For the prime 11 the table is

$\mathbf{Z}/11^k$	n	1	2
$\mathbf{Z}/1$		10	65
$\mathbf{Z}/11$			55

Table A.5: Exponent e_k in decomposition of $Q_{11,n}$ as direct sum of groups of the form $(\mathbf{Z}/11^k)^{\oplus e_k}$.

The pattern of the table seems to be described by 10-term sums of 11-nomial coefficients.

A.3 Sage code

The Sage code implementing the cokernel calculation is

```

1 sage: def first_non_zero_entry(v):
2 ...     r"""
3 ...     Return the first non-zero entry of the row
4 ...     vector 'v'.
5 ...
6 ...     The input is a row vector 'v'. The output
7 ...     is the first non-zero entry of 'v',
8 ...     starting from the right, if it exists.
9 ...     Otherwise the output is None.
10 ...     """
11 ...     for i in range(v.parent().rank()):
12 ...         if v[i] != 0:
13 ...             return v[i]
14 ...     return None
15 sage: def next_p_vector(v,p):
16 ...     r"""
17 ...     Return the next vector modulo p in lexicographic
18 ...     ordering.
```



```

19 ...
20 ...     Input is a prime 'p' and a row vector 'v' with entries
21 ...     between 0 and p-1.
22 ...     Output is the next vector of this type in
23 ...     lexicographic ordering, or None if no such
24 ...     vector exists.
25 ...     """
26 ...     # Start at the right hand side of the row vector 'v'.
27 ...     n = v.parent().rank() - 1
28 ...     # Move from the right to the left.
29 ...     for i in range(v.parent().rank()):
30 ...         # If we can increase the digit ...
31 ...         if v[n-i] < p-1:
32 ...             # ... do it, and return the resulting vector,
33 ...             v[n-i] = v[n-i] + 1
34 ...             return v
35 ...         # otherwise the entry equals 'p-1', hence we need
36 ...         # set the digit to zero and move one digit to
37 ...         # the left
38 ...         elif v[n-i] == p-1:
39 ...             v[n-i] = 0
40 ...     return None
41 sage: def matA(p,n):
42 ...     r"""
43 ...     Return a matrix with generators for all the
44 ...     subgroups of 'C_p^n' isomorphic to 'C_p'.
45 ...
46 ...     Input is a prime 'p' and a natural number 'n'.
47 ...     Output is a matrix 'A' whose columns are
48 ...     precisely generators of all the subgroups
49 ...     of 'C_p^n' isomorphic to 'C_p', namely the
50 ...     vectors that have the shape
51 ...     '(0,0,...,1,*,*,...,*)^T,' where '*' means any
52 ...     entry and '^T' transpose.
53 ...     """
54 ...     # We first do some initialization.
55 ...     # Make 'v' the vector '(0,0,...,0,1)'.
56 ...     v = (ZZ**n)(0)
57 ...     v[n-1] = 1
58 ...     # Make the matrix 'A' the same thing,
59 ...     # but then as a matrix:
60 ...     A = MatrixSpace(ZZ,n,1)(0)
61 ...     A[n-1,0] = 1
62 ...     # Now 'A' already has the current 'v' as
63 ...     # its first column, so we move on to the
64 ...     # next 'v':
65 ...     v = next_p_vector(v,p)
66 ...     # While we haven't gone through all the
67 ...     # vectors ...
68 ...     while v != None:
69 ...         # ... if the current vector is one of our
70 ...         # chosen generators ...
71 ...         if first_non_zero_entry(v) == 1:
72 ...             # ... augment it to A,
73 ...             A = A.augment(v)
74 ...             # otherwise, skip this vector.
75 ...             v = next_p_vector(v,p)
76 ...     # Once we have augmented all the
77 ...     # generators to 'A', return 'A'.

```

```

78 ...     return A
79 sage: def matB(p,n):
80 ...     r"""
81 ...     Return a matrix with columns the
82 ...     non-identity elements of 'C_p^n'.
83 ...
84 ...     Input is a prime p and a natural number n.
85 ...     Output is a matrix whose columns are
86 ...     precisely all the non-identity elements of
87 ...     'C_p^n'.
88 ...     """
89 ...     # First we do some initialization
90 ...     # Set 'v' to be the first non-identity
91 ...     # element in lexicographic ordering.
92 ...     # 'v = (0,0,...,0,1)'.
93 ...     v = (ZZ**n)(0)
94 ...     v[n-1] = 1
95 ...     # Do the same for 'B', but then as a
96 ...     # matrix.
97 ...     B = MatrixSpace(ZZ,n,1)(0)
98 ...     B[n-1,0] = 1
99 ...     # 'B' now already has the current 'v' as
100 ...     # a column, so we move to the next 'v'.
101 ...     v = next_p_vector(v,p)
102 ...     # While we haven't added all elements...
103 ...     while v != None:
104 ...         # add the current one.
105 ...         B = B.augment(v)
106 ...         v = next_p_vector(v,p)
107 ...     return B
108 sage: def redmatmodp(Matrx,p):
109 ...     r"""
110 ...     Return the integer matrix 'Matrx' with all
111 ...     its entries reduced '\mod p'.
112 ...
113 ...     Input is a matrix 'Matrx' with integer values
114 ...     and a prime 'p'. Output is the matrix 'Matrx'
115 ...     with integer entries, reduced '\mod p',
116 ...     hence the output matrix has entries between
117 ...     '0' and 'p-1'.
118 ...     """
119 ...     Matroutput = Matrx.parent()(0)
120 ...     for row in range(Matrx.parent().nrows()):
121 ...         for col in range(Matrx.parent().ncols()):
122 ...             Matroutput[row,col] = mod(Matrx[row,col],p)
123 ...     return Matroutput
124 sage: def matD(C,p,n):
125 ...     r"""
126 ...     Input is a prime p, a natural number n,
127 ...     and a matrix C with entries between 0
128 ...     and p-1. The idea is that all the
129 ...     entries between 1 and p-1 correspond
130 ...     to different basis elements of the
131 ...     representation ring. Therefore, each
132 ...     row of entries between 0 and p-1 will
133 ...     give p-1 rows in the output matrix. If
134 ...     the j-th entry of the row is a number
135 ...     1 <= k <= p-1, then this will give a 1
136 ...     in the k-th row and j-th column (of the

```

```

137 ...     p-1 columns corresponding to this
138 ...     column).
139 ...     """
140 ...     dim = p**n - 1
141 ...     D = MatrixSpace(ZZ,dim,dim)(0)
142 ...     for row in range((p**n - 1)/(p-1)):
143 ...         for col in range(p**n-1):
144 ...             if C[row,col] != 0:
145 ...                 D[C[row,col] + (p-1)*row - 1,col] = 1
146 ...     return D
147 sage: def count_list_elements(L):
148 ...     r"""
149 ...     Count the elements in a list.
150 ...
151 ...     Input is a list. Output is a dictionary with
152 ...     keys the elements of the list, values the
153 ...     number of occurrences.
154 ...     """
155 ...     counts = {}
156 ...     for i in range(len(L)):
157 ...         if L[i] in counts:
158 ...             counts[L[i]] = counts[L[i]] + 1
159 ...         else:
160 ...             counts[L[i]] = 1
161 ...     return counts
162 sage: def calculation_2(p,n):
163 ...     """
164 ...     For the n-th elementary abelian p-group,
165 ...     one can look at the orbit category
166 ...     given by the family of cyclic subgroups.
167 ...     The representation ring functor gives a
168 ...     contravariant functor on this category,
169 ...     and we can compute the limit. The
170 ...     representation ring of the total group
171 ...     maps injectively into this limit. This
172 ...     is in general not surjective, and this
173 ...     function computes the cokernel. The
174 ...     output is a dictionary, whose keys are
175 ...     moduli of the integers, and whose values
176 ...     are the exponents with which the
177 ...     integers modulus the moduli appear in
178 ...     the quotient. For example,
179 ...     calculation_1(2,3) yields
180 ...     {1: 3, 2: 3, 4: 1}, which means that
181 ...     the quotient is isomorphic to
182 ...     {e}^3 x C_2^3 x C_4.
183 ...     """
184 ...     A = matA(p,n)
185 ...     print "Done A"
186 ...     B = matB(p,n)
187 ...     print "Done B"
188 ...     C = A.transpose()*B
189 ...     print "Done C"
190 ...     C = redmatmodp(C,p)
191 ...     print "Done reduction modulo p"
192 ...     D = matD(C,p,n)
193 ...     print "Done D"
194 ...     # The elementary divisors of 'D'
195 ...     # are the elements on the diagonal

```

```
196 ...     # of a Smith Normal form of 'D'.
197 ...     # An elementary divisor 'd'
198 ...     # corresponds to an occurrence of
199 ...     # '\ZZ/d \ZZ' in the quotient.
200 ...     divisors = D.elementary_divisors()
201 ...     print "Done Elementary Divisors"
202 ...     counted = count_list_elements(divisors)
203 ...     return counted
204 sage: calculation_2(2,3)
205 Done A
206 Done B
207 Done C
208 Done reduction modulo p
209 Done D
210 Done Elementary Divisors
211 {1: 3, 2: 3, 4: 1}
```

This last calculation shows, for example, that

$$Q_{2,3} \cong (\mathbf{Z}/1)^{\oplus 3} \oplus (\mathbf{Z}/2)^{\oplus 3} \oplus \mathbf{Z}/4.$$

Bibliography

- [AE81] J. L. Alperin and L. Evens. Representations, resolutions and Quillen's dimension theorem. *J. Pure Appl. Algebra*, 22(1):1–9, 1981. (Cited on page 2.)
- [AM04] Alejandro Adem and R. James Milgram. *Cohomology of finite groups*, volume 309 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2004. (Cited on pages 33, 37, 38, 43, 46, 47, and 48.)
- [And76] Désiré André. Mémoire sur les combinaisons régulières et leurs applications. *Ann. Sci. École Norm. Sup. (2)*, 5:155–198, 1876. (Cited on page 110.)
- [Arm88] M. A. Armstrong. *Groups and symmetry*. Undergraduate Texts in Mathematics. Springer-Verlag, New York, 1988. (Cited on page 29.)
- [Ben93] D. J. Benson. The image of the transfer map. *Arch. Math. (Basel)*, 61(1):7–11, 1993. (Cited on page 2.)
- [Ben04] Dave Benson. Dickson invariants, regularity and computation in group cohomology. *Illinois J. Math.*, 48(1):171–197, 2004. (Cited on page 27.)
- [Bir67] Garrett Birkhoff. *Lattice theory*. Third edition. American Mathematical Society Colloquium Publications, Vol. XXV. American Mathematical Society, Providence, R.I., 1967. (Cited on page 52.)
- [BK72] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin-New York, 1972. (Cited on page 12.)
- [Bor60] Armand Borel. *Seminar on transformation groups*. With contributions by G. Bredon, E. E. Floyd, D. Montgomery, R. Palais. Annals of Mathematics Studies, No. 46. Princeton University Press, Princeton, N.J., 1960. (Cited on page 18.)
- [Bur55] W. Burnside. *Theory of groups of finite order*. Cambridge University Press, 1955. 2d ed. (Cited on page 28.)
- [Car95] Jon F. Carlson. Depth and transfer maps in the cohomology of groups. *Math. Z.*, 218(3):461–468, 1995. (Cited on pages 2 and 64.)

- [CE56] Henri Cartan and Samuel Eilenberg. *Homological Algebra*. Princeton University Press, 1956. (Cited on pages 29 and 46.)
- [Com74] Louis Comtet. *Advanced Combinatorics*. D. Reidel Publishing Company, Dordrecht-Holland/Boston-U.S.A, 1974. (Cited on page 110.)
- [CTVEZ03] Jon F. Carlson, Lisa Townsley, Luis Valeri-Elizondo, and Mucheng Zhang. *Cohomology rings of finite groups*, volume 3 of *Algebras and Applications*. Kluwer Academic Publishers, Dordrecht, 2003. With an appendix: Calculations of cohomology rings of groups of order dividing 64 by Carlson, Valeri-Elizondo and Zhang. (Cited on pages 27, 48, 50, 52, 61, 64, 82, 87, 92, and 93.)
- [Duf81] J. Duflot. Depth and equivariant cohomology. *Comment. Math. Helv.*, 56(4):627–637, 1981. (Cited on page 2.)
- [Eve91] Leonard Evens. *The cohomology of groups*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1991. Oxford Science Publications. (Cited on page 31.)
- [Fah12] N.-E. Fahssi. Polynomial Triangles Revisited. *ArXiv e-prints*, February 2012. <https://arxiv.org/abs/1202.0228>. (Cited on page 110.)
- [GK11] David J. Green and Simon A. King. The computation of the cohomology rings of all groups of order 128. *J. Algebra*, 325:352–363, 2011. (Cited on pages 27 and 28.)
- [GK15] David J. Green and Simon A. King. The cohomology of finite p -groups, 2015. Website. <http://users.minet.uni-jena.de/cohomology/>. (Cited on page 28.)
- [Hus94] Dale Husemoller. *Fibre bundles*, volume 20 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 1994. (Cited on pages 21, 22, and 25.)
- [Ill83] Sören Illman. The equivariant triangulation theorem for actions of compact Lie groups. *Math. Ann.*, 262(4):487–501, 1983. (Cited on page 25.)
- [Inc05] OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, 2005. <http://oeis.org/A123456>. (Cited on pages 110 and 113.)
- [LMSM86] L. G. Lewis, Jr., J. P. May, M. Steinberger, and J. E. McClure. *Equivariant stable homotopy theory*, volume 1213 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986. With contributions by J. E. McClure. (Cited on pages 7 and 9.)
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009. (Cited on page 11.)

- [MM02] M. A. Mandell and J. P. May. Equivariant orthogonal spectra and S -modules. *Mem. Amer. Math. Soc.*, 159(755):x+108, 2002. (Cited on page 4.)
- [MNN15] Akhil Mathew, Niko Naumann, and Justin Noel. Derived induction and restriction theory. *ArXiv e-prints*, July 2015. <http://arxiv.org/abs/1507.06867>. (Cited on pages 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 17, 18, 20, 25, 29, 33, 35, 46, 97, 104, 108, 109, and 110.)
- [MNN17] Akhil Mathew, Niko Naumann, and Justin Noel. Nilpotence and descent in equivariant stable homotopy theory. *Adv. Math.*, 305:994–1084, 2017. (Cited on pages 5, 8, 9, 12, 16, 17, 19, 20, and 25.)
- [PY03] Jonathan Pakianathan and Ergün Yalçın. On nilpotent ideals in the cohomology ring of a finite group. *Topology*, 42(5):1155–1183, 2003. (Cited on page 25.)
- [Qui71a] Daniel Quillen. The Adams conjecture. *Topology*, 10:67–80, 1971. (Cited on pages 33 and 98.)
- [Qui71b] Daniel Quillen. The mod 2 cohomology rings of extra-special 2-groups and the spinor groups. *Mathematische Annalen*, 194:197–212, 1971. (Cited on page 27.)
- [Qui71c] Daniel Quillen. The spectrum of an equivariant cohomology ring. I, II. *Ann. of Math. (2)*, 94:549–572; *ibid.* (2) 94 (1971), 573–602, 1971. (Cited on page 1.)
- [Rus89] David J. Rusin. The cohomology of the groups of order 32. *Math. Comp.*, 53(187):359–385, 1989. (Cited on page 27.)
- [S+15] W. A. Stein et al. *Sage Mathematics Software*. The Sage Development Team, 2015. <http://www.sagemath.org>. (Cited on page 112.)
- [Ser77] Jean-Pierre Serre. *Linear representations of finite groups*. Springer-Verlag, New York-Heidelberg, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42. (Cited on pages 26 and 106.)
- [Sym91] Peter Symonds. The complexity of a module and elementary abelian subgroups: a geometric approach. *Proc. Amer. Math. Soc.*, 113(1):27–29, 1991. (Cited on page 25.)
- [Tot14] Burt Totaro. *Group cohomology and algebraic cycles*, volume 204 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2014. (Cited on pages 50, 52, 86, and 88.)
- [Wil05] Marcel Wild. The groups of order sixteen made easy. *Amer. Math. Monthly*, 112(1):20–31, 2005. (Cited on pages 29 and 30.)