

On the operations of sequences in rings and binomial type sequences

Stefano Barbero, Umberto Cerruti, Nadir Murru

Department of Mathematics G. Peano, University of Turin,
Via Carlo Alberto 10, 10123, Torino, ITALY

stefano.barbero@unito.it, umberto.cerruti@unito.it, nadir.murru@unito.it

Abstract

Given a commutative ring with identity R , many different and interesting operations can be defined over the set H_R of sequences of elements in R . These operations can also give H_R the structure of a ring. We study some of these operations, focusing on the binomial convolution product and the operation induced by the composition of exponential generating functions. We provide new relations between these operations and their invertible elements. We also study automorphisms of the Hurwitz series ring, highlighting that some well-known transforms of sequences (such as the Stirling transform) are special cases of these automorphisms. Moreover, we introduce a novel isomorphism between H_R equipped with the componentwise sum and the set of the sequences starting with 1 equipped with the binomial convolution product. Finally, thanks to this isomorphism, we find a new method for characterizing and generating all the binomial type sequences.

Keywords: binomial type sequence; binomial convolution product; Hurwitz series ring
MSC2010: 11B99, 11T06, 13F25

1 Introduction

Given a commutative unitary ring $(R, +, \cdot)$, where as usual the unit, the additive identity and the set of invertible elements will be denoted by $1, 0$, and R^* , we will examine the set, that we call H_R , of sequences with terms belonging to R . The aim of this paper is to highlight new perspectives in studying the algebraic structures arising when H_R is equipped with the most commonly used and interesting operations between sequences. Specifically, we will provide novel relations between inverse elements with respect to the binomial convolution product (also called Hurwitz product) and with respect to the product defined by the composition of exponential generating functions. Moreover, we will study automorphisms of the Hurwitz series ring and find a new computable isomorphism between the set of sequences starting with 1 equipped with the Hurwitz product and H_R equipped with the componentwise sum. Finally thanks to this isomorphism we will show a straightforward method to generate all the binomial type sequences.

First of all, let us start with some definitions and notation.

Definition 1. *Let us consider the set*

$$H_R = \{(a_n)_{n=0}^{+\infty} = (a_0, a_1, a_2, \dots) : \forall i \geq 0, \quad a_i \in R\}$$

of sequences whose terms belongs to R . We denote with a bold letter the generic element of H_R and we will refer to the $n + 1$ -th term of $\mathbf{a} \in H_R$ with the two equivalent notations $\mathbf{a}[n]$ or a_n .

We also define the following two subsets of H_R

$$H_R^{(n)} = \{\mathbf{a} \in H_R : \mathbf{a} = (a_0, a_1, \dots, a_{n-1})\}$$

the set of sequences having length $n \geq 0$ and the set

$$H_R^m = \{\mathbf{a} \in H_R : \mathbf{a}[i] = 0, i = 0, 1, \dots, m-1\},$$

of sequences having the first m terms equal to zero. We call zero of order m any $\mathbf{a} \in H_R^m$. The exponential generating function (e.g.f.) and ordinary generating function (o.g.f.) related to $\mathbf{a} \in H_R$ will be

$$s_e(\mathbf{a}) = A(t) = \sum_{h=0}^{+\infty} a_h \frac{t^h}{h!}, \quad s_o(\mathbf{a}) = \bar{A}(t) = \sum_{h=0}^{+\infty} a_h t^h,$$

where we denote with s_e and s_o the bijections that map any sequence $\mathbf{a} \in H_R$ to its e.g.f. or o.g.f. respectively.

We define two useful transform acting on sequences $\mathbf{a} \in H_R$.

Definition 2. The transforms λ_- (the shift operator) and $\lambda_{+,u}$, $u \in R$, map any sequence $\mathbf{a} \in H_R$ respectively into the sequences

$$\lambda_-(\mathbf{a}) := (a_1, a_2, \dots), \quad \lambda_{+,u}(\mathbf{a}) := (u, a_0, a_1, \dots).$$

Finally we recall the definition of an important family of partition polynomials.

Definition 3. Let us consider the sequence of variables $X = (x_1, x_2, \dots)$. The complete ordinary Bell polynomials are defined by

$$B_0(X) = 1, \quad \forall n \geq 1 \quad B_n(X) = B_n(x_1, x_2, \dots, x_n) = \sum_{k=1}^n B_{n,k}(X),$$

where $B_{n,k}(X)$ are the partial ordinary Bell polynomials, with

$$B_{0,0}(X) = 1, \quad \forall n \geq 1 \quad B_{n,0}(X) = 0, \quad \forall k \geq 1 \quad B_{0,k}(X) = 0,$$

$$B_{n,k}(X) = B_{n,k}(x_1, x_2, \dots, x_n) = k! \sum_{\substack{i_1+2i_2+\dots+ni_n=n \\ i_1+i_2+\dots+i_n=k}} \prod_{j=1}^n \frac{x_j^{i_j}}{i_j!},$$

or, equivalently,

$$B_{n,k}(X) = B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = k! \sum_{\substack{i_1+2i_2+\dots+(n-k+1)i_{n-k+1}=n-k+1 \\ i_1+i_2+\dots+i_{n-k+1}=k}} \prod_{j=1}^{n-k+1} \frac{x_j^{i_j}}{i_j!},$$

satisfying the equality

$$\left(\sum_{n \geq 1} x_n z^n \right)^k = \sum_{n \geq k} B_{n,k}(X) z^n.$$

The most simple operation we may introduce on H_R is the componentwise sum, i.e., given any $\mathbf{a}, \mathbf{b} \in H_R$, then $\mathbf{a} + \mathbf{b} = \mathbf{c}$, where $c_n = a_n + b_n$, for all $n \geq 0$. In this case, the additive inverse of an element $\mathbf{a} \in H_R$ is obviously the sequence $-\mathbf{a} = (-a_n)_{n=0}^{+\infty}$ and $(H_R, +)$ is a commutative group, whose identity is $\mathbf{0} = (0, 0, 0, \dots)$. On the other hand many different operations, playing the role of a product, can be defined on H_R .

The Hadamard product

Given two sequences $\mathbf{a}, \mathbf{b} \in H_R$, the *Hadamard product* $\mathbf{a} \bullet \mathbf{b}$ is the componentwise product, i.e., $\mathbf{a} \bullet \mathbf{b} = \mathbf{c}$, where $c_n = a_n \cdot b_n$, for all $n \geq 0$. A sequence $\mathbf{a} \in H_R$ is invertible with respect to \bullet if and only if for all $n \geq 0$ $a_n \in R^*$. The identity is clearly the sequence made up with all the elements equal to 1, which we denote by $\mathbf{1}$. This product takes its name since the paper of Hadamard [12]. Some recent and interesting studies on the Hadamard product can be found, e.g., in [1] and [8].

The Hurwitz product

Another well-known and studied operation is the *Hurwitz product* (also called *binomial convolution product*) that we denote with \star . Given $\mathbf{a}, \mathbf{b} \in H_R$, the Hurwitz product is defined as $\mathbf{a} \star \mathbf{b} = \mathbf{c}$, where

$$\forall n \geq 0, \quad c_n = \sum_{h=0}^n \binom{n}{h} a_h b_{n-h}.$$

The identity with respect to the Hurwitz product is the sequence $\bar{\mathbf{1}} = (1, 0, 0, \dots)$. Moreover the Hurwitz product and the product between e.g.f.s of sequences are strictly related. Indeed if $s_e(\mathbf{a}) = A(t)$ and $s_e(\mathbf{b}) = B(t)$ then

$$s_e(\mathbf{a} \star \mathbf{b}) = A(t)B(t) = s_e(\mathbf{a})s_e(\mathbf{b}). \quad (1)$$

Remark 1. If we consider the set $H_R[[t]]$ of the formal power series of the form $\sum_{h=0}^{+\infty} a_h \frac{t^h}{h!}$, then $(H_R[[t]], +, \star)$ is the Hurwitz series ring. The Hurwitz series ring has been extensively studied during the years, latest papers on this subject are, e.g., [13], [3], [9], [4].

An element $\mathbf{a} \in H_R$ is invertible with respect to \star if and only if $a_0 \in R^*$. In the following, we will denote by \mathbf{a}^{-1} such an inverse and with H_R^* the set of the invertible sequences in H_R with respect to the Hurwitz product \star . When it exists, the inverse $\mathbf{b} = \mathbf{a}^{-1}$ can be recursively evaluated using the relations

$$b_0 = a_0^{-1}, \quad b_n = -a_0^{-1} \sum_{h=1}^n \binom{n}{h} a_h b_{n-h}, \quad n \geq 1.$$

Furthermore we easily observe from (1) that the relation between the e.g.f. of $\mathbf{a} = A(t)$ and \mathbf{a}^{-1} is

$$s_e(\mathbf{a}^{-1}) = \frac{1}{A(t)} = (s_e(\mathbf{a}))^{-1}$$

since $s_e(\mathbf{a} \star \mathbf{a}^{-1}) = s_e(\bar{\mathbf{1}}) = 1$. We point out that a closed expression for \mathbf{a}^{-1} has been found in [2] by means of the Bell polynomials. We report this result in the following theorem.

Theorem 1. Let $\mathbf{a} \in H_R^*$, and $\mathbf{b} = \mathbf{a}^{-1}$. Then for all $n \geq 0$, we have

$$b_n = \frac{n! B_n(g_0, g_1, g_2, \dots, g_n)}{a_0}, \quad (2)$$

where

$$\mathbf{g} = (g_n)_{n=0}^{+\infty} = \left(-\frac{a_{n+1}}{a_0(n+1)!} \right)_{n=0}^{+\infty}.$$

The Cauchy product

The Cauchy product $*$ between two sequences $\mathbf{a}, \mathbf{b} \in H_R$, defined as $\mathbf{a} * \mathbf{b} = \mathbf{c}$, where

$$\forall n \geq 0, \quad c_n = \sum_{h=0}^n a_h b_{n-h},$$

is strictly connected to the Hurwitz product. Indeed, the map

$$\gamma : (H_R, +, \star) \rightarrow (H_R, +, \star), \quad \gamma(\mathbf{a})[n] = n! \mathbf{a}[n], n \geq 0 \quad (3)$$

is obviously an isomorphism. Thus all the considerations regarding the Hurwitz product \star can be viewed in terms of the Cauchy product, and vice versa. In the following, we will mainly deal with the Hurwitz product \star .

Compositions of generating functions

Finally, we can define two products by composing ordinary and exponential generating functions. Specifically, given $\mathbf{a}, \mathbf{b} \in H_R$, we define $\mathbf{a} \circ_o \mathbf{b} = \mathbf{c}$ and $\mathbf{a} \circ_e \mathbf{b} = \mathbf{d}$, where

$$\mathbf{c} = s_o^{-1} \left(\sum_{h=0}^{+\infty} a_h \left(\sum_{k=0}^{+\infty} b_k t^k \right)^h \right), \quad \mathbf{d} = s_e^{-1} \left(\sum_{h=0}^{+\infty} \frac{a_h}{h!} \left(\sum_{k=0}^{+\infty} b_k \frac{t^k}{k!} \right)^h \right).$$

Let us observe that these products are defined only if the sequence \mathbf{b} is a zero of order ≥ 1 .

In the following, we will often use the product \circ_e obtained by the composition of exponential generating functions. Thus, we will write \circ instead of \circ_e . It is easy to verify that the sequence $\mathbf{1} = \lambda_{+,0}(\bar{\mathbf{1}}) = (0, 1, 0, 0, \dots)$ is the identity with respect to the product \circ .

A sequence $\mathbf{a} \in H_R$ is invertible with respect to the product \circ if and only if $a_0 = 0$ and $a_1 \in R^*$. In this case, the inverse of \mathbf{a} that we denote by $\mathbf{a}^{(-1)}$ is the sequence in H_R whose terms are

$$\mathbf{a}^{(-1)}[0] = 0, \quad \forall n \geq 1 \quad \mathbf{a}^{(-1)}[n] = \frac{(n-1)!}{a_1^n} \sum_{j=0}^{n-1} (-1)^j \binom{n+j-1}{j} B_{n-1,j}(\bar{a}_1, \dots, \bar{a}_{n-j}), \quad (4)$$

where $\bar{a}_i = \frac{a_{i+1}}{a_1(i+1)!}$, $i \geq 1$ (see, e.g. [5], chapter 11, for a detailed proof).

Given the sequence $\mathbf{a} \in H_R$, with $a_0 \in R^*$, the terms of the sequence \mathbf{a}^{-1} can be explicitly evaluated using (2):

$$\mathbf{a}^{-1} = \left(\frac{1}{a_0}, -\frac{a_1}{a_0^2}, \frac{2a_1^2 - a_0 a_2}{a_0^3}, \frac{-6a_1^3 + 6a_0 a_1 a_2 - a_0^2 a_3}{a_0^4}, \dots \right).$$

On the other hand, the terms of $\lambda_{+,0}(\mathbf{a})^{(-1)}$ can be explicitly evaluated with (4):

$$\lambda_{+,0}(\mathbf{a})^{(-1)} = \left(0, \frac{1}{a_0}, -\frac{a_1}{a_0^3}, \frac{3a_1^2 - a_0 a_2}{a_0^5}, \frac{-15a_1^3 + 10a_0 a_1 a_2 - a_0^2 a_3}{a_0^7}, \dots \right).$$

There is new and simple relation between \mathbf{a}^{-1} and $\lambda_{+,0}(\mathbf{a})^{(-1)}$, i.e., between the inverse elements with respect to the products \star and \circ respectively, as we will show in the next theorem.

Theorem 2. *Let $\mathbf{a} \in H_R$ be a \star -invertible element, i.e., $a_0 \in R^*$, then*

$$\mathbf{a}^{-1} = \lambda_- \left(\lambda_{+,0}(\mathbf{a})^{(-1)} \right) \circ \lambda_{+,0}(\mathbf{a}). \quad (5)$$

Proof. Let $s_e(\mathbf{a}) = A(t)$, then

$$s_e(\lambda_{+,0}(\mathbf{a})) = F(t) = \int A(t) dt.$$

If $s_e(\lambda_{+,0}(\mathbf{a})^{(-1)}) = G(t)$, by definition of inverse with respect to \circ , we have

$$F(G(t)) = G(F(t)) = t.$$

Moreover, let us observe that

$$s_e \left(\lambda_- \left(\lambda_{+,0}(\mathbf{a})^{(-1)} \right) \right) = G'(t) \quad s_e \left(\lambda_- \left(\lambda_{+,0}(\mathbf{a})^{(-1)} \right) \circ \lambda_{+,0}(\mathbf{a}) \right) = G'(F(t)).$$

Thus, we have

$$G'(F(t))F'(t) = 1$$

and, since $F'(t) = A(t)$, we obtain

$$G'(F(t)) = \frac{1}{A(t)},$$

that completes the proof. \square

It seems hard to find a simple relation like (5) describing $\lambda_{+,0}(\mathbf{a})^{(-1)} = \boldsymbol{\lambda}^{(-1)}$ in terms of \mathbf{a}^{-1} . However, we easily find the following recursive expression

$$\forall n \geq 0 \quad \boldsymbol{\lambda}^{(-1)}[n+1] = n! \sum_{k=0}^n \frac{\mathbf{a}^{-1}[k]}{k!} B_{n,k} \left(\bar{\boldsymbol{\lambda}}^{(-1)}[1], \dots, \bar{\boldsymbol{\lambda}}^{(-1)}[n-k+1] \right), \quad (6)$$

where $\bar{\boldsymbol{\lambda}}^{(-1)} = \frac{\boldsymbol{\lambda}^{(-1)}[i]}{i!}$ for all $i \geq 1$. Indeed, using the same notation as in the previous theorem, we know that

$$F'(G(t))G'(t) = 1, \quad F'(G(t)) = A(G(t)).$$

Thus,

$$G'(t) = \frac{1}{A(G(t))}$$

and since

$$\begin{aligned} \frac{1}{A(G(t))} &= \sum_{k=0}^{+\infty} \frac{\mathbf{a}^{-1}[k]}{k!} (G(t))^k = \sum_{n=0}^{+\infty} \left(\sum_{k=0}^n \frac{\mathbf{a}^{-1}[k]}{k!} B_{n,k} \left(\bar{\boldsymbol{\lambda}}^{(-1)}[1], \dots, \bar{\boldsymbol{\lambda}}^{(-1)}[n-k+1] \right) \right) t^n, \\ G'(t) &= \sum_{n=0}^{+\infty} \frac{\boldsymbol{\lambda}^{(-1)}[n+1]}{n!} t^n \end{aligned}$$

we obtain (6).

Automorphisms

Some distributive properties hold for the products \bullet , \star , \circ . We summarize them in the following proposition.

Proposition 1. *Given any $\mathbf{b} \in H_R^0$ and $\mathbf{a}, \mathbf{c} \in H_R$, we have*

- $(\mathbf{a} + \mathbf{c}) \circ \mathbf{b} = (\mathbf{a} \circ \mathbf{b}) + (\mathbf{c} \circ \mathbf{b});$
- $(\mathbf{a} \star \mathbf{c}) \circ \mathbf{b} = (\mathbf{a} \circ \mathbf{b}) \star (\mathbf{c} \circ \mathbf{b}).$

Given the sequence $\boldsymbol{\beta}(r) = (r^n)_{n=0}^{+\infty}$, for any $r \in R$, and any $\mathbf{a}, \mathbf{c} \in H_R$, we have

- $(\mathbf{a} + \mathbf{c}) \bullet \boldsymbol{\beta}(r) = (\mathbf{a} \bullet \boldsymbol{\beta}(r)) + (\mathbf{c} \bullet \boldsymbol{\beta}(r));$
- $(\mathbf{a} \star \mathbf{c}) \bullet \boldsymbol{\beta}(r) = (\mathbf{a} \bullet \boldsymbol{\beta}(r)) \star (\mathbf{c} \bullet \boldsymbol{\beta}(r)).$

Proof. The proof is straightforward remembering that $+$ and \star represent the sum and the product of formal exponential series, respectively. \square

Clearly the product \bullet by $\beta(r)$ may be considered as an endomorphism of the ring $(H_R, +, \star)$. Moreover, we can also introduce a class of endomorphisms in $(H_R, +, \star)$ by means of the product \circ by $\mathbf{b} \in H_R^0$. We explicitly state these important and immediate consequences of the previous proposition

Corollary 1. *For any $z \in R$ and any fixed sequence $\mathbf{b} \in H_R^0$ the maps*

$$\mathcal{H}_{\beta(z)} : (H_R, +, \star) \rightarrow (H_R, +, \star), \quad \mathcal{H}_{\beta(z)}(\mathbf{a}) = \mathbf{a} \bullet \beta(z)$$

$$\mu_{\mathbf{b}} : (H_R, +, \star) \rightarrow (H_R, +, \star), \quad \mu_{\mathbf{b}}(\mathbf{a}) = \mathbf{a} \circ \mathbf{b}$$

are endomorphisms. Furthermore, if $r, \mathbf{b}[1] \in R^*$ then $\mathcal{H}_{\beta(z)}$ and $\mu_{\mathbf{b}}$ are automorphisms whose inverses are, respectively, $\mathcal{H}_{\beta(r^{-1})}$ and $\mu_{\mathbf{b}^{-1}} = \mu_{\mathbf{b}(-1)}$ and the identity automorphism is $\mathcal{H}_{\mathbf{1}} = \mu_{\mathbf{1}}$.

Some interesting classical transforms acting on sequences may be interpreted as special cases of automorphisms $\mu_{\mathbf{b}}$ of $(H_R, +, \star)$ for a suitable choice of \mathbf{b} . Let us summarize these transforms in the following definition.

Definition 4. *Given any sequence $\mathbf{a} \in H_R$, we define*

1. the alternating sign transform \mathcal{E} that maps \mathbf{a} into a sequence $\mathbf{b} = \mathcal{E}(\mathbf{a}) \in H_R$, whose terms are

$$b_n = (-1)^n a_n;$$

2. the Stirling transform \mathcal{S} that maps \mathbf{a} into a sequence $\mathbf{b} = \mathcal{S}(\mathbf{a}) \in H_R$, whose terms are

$$b_n = \sum_{h=0}^n \left\{ \begin{matrix} n \\ h \end{matrix} \right\} a_h,$$

where $\left\{ \begin{matrix} n \\ h \end{matrix} \right\}$ are the Stirling numbers of the second kind (see e.g. [11], chapter 6, for definition and properties of Stirling numbers of first and second kind);

3. the inverse \mathcal{S}^{-1} of the Stirling transform that maps \mathbf{a} into a sequence $\mathbf{b} = \mathcal{S}^{-1}(\mathbf{a}) \in H_R$, whose terms are

$$b_n = \sum_{h=0}^n (-1)^{n-h} \left[\begin{matrix} n \\ h \end{matrix} \right] a_h,$$

where $\left[\begin{matrix} n \\ h \end{matrix} \right]$ are the (unsigned) Stirling numbers of the first kind.

The transform \mathcal{E} is often used for studying properties of sequences, also for its simple but important effect on e.g.f.s, since if $s_e(\mathbf{a}) = A(t)$, then $s_e(\mathcal{E}(\mathbf{a})) = A(-t)$ for all $\mathbf{a} \in H_R$. Clearly, \mathcal{E} is an automorphism of $(H_R, +, \star)$, since we have $\mathcal{E} = \mu_{-\mathbf{1}}$. Furthermore, to clearly identify the sequence \mathbf{b} such that $\mathcal{S} = \mu_{\mathbf{b}}$ we need the following result

Proposition 2. *Given $\mathbf{a} \in H_R$, if $s_e(\mathbf{a}) = A(t)$, then $s_e(\mathcal{S}(\mathbf{a})) = A(e^t - 1)$.*

Proof. Let us recall that $\left\{ \begin{matrix} n \\ h \end{matrix} \right\} = 0$ when $h > n$ and

$$\sum_{n=0}^{+\infty} \left\{ \begin{matrix} n \\ h \end{matrix} \right\} \frac{t^n}{n!} = \frac{(e^t - 1)^h}{h!},$$

see [11] pag. 337. Thus, we have

$$B(t) = \sum_{n=0}^{+\infty} \left(\sum_{h=0}^n \left\{ \begin{matrix} n \\ h \end{matrix} \right\} a_h \right) \frac{t^n}{n!} = \sum_{h=0}^{+\infty} a_h \left(\sum_{n=0}^{+\infty} \left\{ \begin{matrix} n \\ h \end{matrix} \right\} \frac{t^n}{n!} \right) = \sum_{h=0}^{+\infty} a_h \frac{(e^t - 1)^h}{h!} = A(e^t - 1).$$

□

Now, observing that $s_e^{-1}(e^t - 1) = (0, 1, 1, 1, \dots) = \lambda_{+,0}(\mathbf{1})$, we have

$$\mathcal{S} = \mu_{\lambda_{+,0}(\mathbf{1})}.$$

Since from Proposition 2 we also easily find that $s_e(\mathcal{S}^{-1}(\mathbf{a})) = A(\log(t + 1))$, from the formal power series identity

$$\log(t + 1) = \sum_{h=1}^{+\infty} (-1)^{h-1} (h-1)! \frac{t^h}{h!},$$

we have

$$\mathcal{S}^{-1} = \mu_{\lambda_{+,0}(\mathcal{E}(\mathbf{f}))},$$

where $\mathbf{f} = (0!, 1!, 2!, \dots)$ is the sequence of the factorial numbers A000142 in Oeis [16].

2 Isomorphisms

In the following we focus on a subgroup of H_R^* , the set of invertible elements of H_R with respect to the Hurwitz product. Specifically, we consider

$$U_R = \{\mathbf{a} \in H_R^* : a_0 = 1\},$$

and we indicate by $U_R^{(n)}$ the set of sequences in U_R of length n . Furthermore we will deal with products and powers of sequences with respect to the Hurwitz product.

Given $\mathbf{a} \in U_R$ with $s_e(\mathbf{a}) = A(t)$, we let \mathbf{a}^x denote the sequence such that $s_e(\mathbf{a}^x) = (A(t))^x$. The sequence \mathbf{a}^x can be viewed as a sequence of polynomials in the variable x and we use the notation $p^{(\mathbf{a})}(x) = \mathbf{a}^x$ in order to highlight this, i.e.,

$$p^{(\mathbf{a})}(x) = \left(p_n^{(\mathbf{a})}(x) \right)_{n=0}^{+\infty} = (\mathbf{a}^x[n])_{n=0}^{+\infty},$$

where $p_n^{(\mathbf{a})}(x) \in R[x]$. Clearly we also have

$$p^{(\mathbf{a})}(x + y) = p^{(\mathbf{a})}(x) \star p^{(\mathbf{a})}(y) \tag{7}$$

since $(A(t))^{x+y} = (A(t))^x (A(t))^y$. In the next proposition, we will provide a closed form for the coefficients of the polynomials $p_n^{(\mathbf{a})}(x)$.

Proposition 3. *Given the polynomial $p_n^{(\mathbf{a})}(x) = \sum_{j=0}^n c_j x^j$, then*

$$c_j = \sum_{h=0}^{n-j} (-1)^h \frac{n!}{(h+j)!} \begin{bmatrix} h+j \\ j \end{bmatrix} B_{n,h+j}(a_1, \dots, a_{n-h-j+1}), \tag{8}$$

for $j = 0, 1, 2, \dots, n$, where $\begin{bmatrix} k \\ j \end{bmatrix}$ are the (unsigned) Stirling numbers of the first kind.

Proof. Since $a = (1, a_1, a_2, \dots)$ has e.g.f. $A(t) = 1 + \sum_{n=1}^{+\infty} \frac{a_n}{n!} t^n$, we have

$$\begin{aligned} (A(t))^x &= \left(1 + \sum_{n=1}^{+\infty} \frac{a_n}{n!} t^n \right)^x = \sum_{k=0}^{+\infty} \binom{x}{k} \sum_{n \geq k} B_{n,k}(a_1, \dots, a_{n-k+1}) t^n = \\ &= \sum_{n=0}^{+\infty} \frac{1}{n!} \left(n! \sum_{k=0}^n \binom{x}{k} B_{n,k}(a_1, \dots, a_{n-k+1}) \right) t^n. \end{aligned}$$

Thus,

$$p_n^{(\mathbf{a})}(x) = n! \sum_{k=0}^n \binom{x}{k} B_{n,k}(a_1, \dots, a_{n-k+1}). \tag{9}$$

Moreover, from equation 6.13 in [11], we can write

$$\binom{x}{k} = \frac{(-1)^k}{k!} \sum_{j=0}^k (-1)^j \begin{bmatrix} k \\ j \end{bmatrix} x^j \quad (10)$$

and finally, substituting this equality in (9)

$$p_n^{(\mathbf{a})}(x) = \sum_{j=0}^n \left(n! \sum_{k=j}^n B_{n,k}(a_1, \dots, a_{n-k+1}) \frac{(-1)^k}{k!} (-1)^j \begin{bmatrix} k \\ j \end{bmatrix} \right) x^j$$

from which, replacing the index k with $h = k - j$, the thesis follows. \square

Interesting consequences arise studying the sequences $\mathbf{a}^x \in U_R$ with respect to the Hurwitz product. From (8) or evaluating the sequence of formal derivatives of $(A(t))^x$

$$x(A(t))^{x-1}A'(t), x(A(t))^{x-1}A''(t) + (x-1)x(A(t))^{x-2}A'(t)^2, \dots$$

we easily obtain that $p_0^{(\mathbf{a})}(x) = 1$ and $x|p_n^{(\mathbf{a})}(x)$ for $n \geq 1$. Now if we consider, for example, $R = \mathbb{Z}_n$, we have for any $\mathbf{a} \in U_{\mathbb{Z}_n}$, $\mathbf{a}^n = (1, 0, 0, \dots) = \bar{\mathbf{1}}$. Moreover, let p be a prime number, then any element different from $\bar{\mathbf{1}}$ in $U_{\mathbb{Z}_p}$ has period p . Thus $(U_{\mathbb{Z}_p}^{(n)}, \star)$ is a finite group of order p^{n-1} and by the structure theorem for finite abelian groups, we have the following isomorphism

$$(U_{\mathbb{Z}_p}^{(n)}, \star) \simeq (H_{\mathbb{Z}_p}^{(n-1)}, +). \quad (11)$$

Since (11) holds for every prime p we also have for all $n \geq 2$

$$(U_{\mathbb{Z}}^{(n)}, \star) \simeq (H_{\mathbb{Z}}^{(n-1)}, +)$$

and consequently

$$(U_{\mathbb{Z}}, \star) \simeq (H_{\mathbb{Z}}, +). \quad (12)$$

The interesting isomorphism (12) is only a special case of a more general computable isomorphism that we can provide between (U_R, \star) and $(H_R, +)$. Let us consider the infinite set of sequences $\mathcal{B} = \{\mathbf{b}^{(1)}, \mathbf{b}^{(2)}, \mathbf{b}^{(3)}, \dots\}$, where

$$\underline{\mathbf{1}} = \mathbf{b}^{(1)} = (1, 1, 1, 1, \dots), \mathbf{b}^{(2)} = (1, 0, 1, 0, \dots), \mathbf{b}^{(3)} = (1, 0, 0, 1, 0, \dots), \dots$$

i.e., for $i \geq 2$, $\mathbf{b}^{(i)}[i] = 1$ and $\mathbf{b}^{(i)}[k] = 0$ for any $k \neq 0, i$.

Definition 5. We define two elements \mathbf{a} and \mathbf{b} of U_R as independent if $\langle \mathbf{a} \rangle \cap \langle \mathbf{b} \rangle = \bar{\mathbf{1}}$, where $\langle \mathbf{u} \rangle$ is the subgroup of (U_R, \star) generated by $\mathbf{u} \in U_R$.

Clearly the sequences $\mathbf{b}^{(i)}$ are mutually independent, since

$$p^{(\mathbf{b}^{(1)})}(x) = (\mathbf{b}^{(1)})^x = (x^n)_{n=0}^{+\infty}$$

and considering $p^{(\mathbf{b}^{(i)})}(x) = (\mathbf{b}^{(i)})^x$, we surely have

$$\forall i \geq 2, \quad p_1^{(\mathbf{b}^{(i)})}(x) = \dots = p_{i-1}^{(\mathbf{b}^{(i)})}(x) = 0.$$

Thus we may consider the set \mathcal{B} as a basis of (U_R, \star) .

Remark 2. The sequences $p^{(\mathbf{b}^{(i)})}(x)$ can be evaluated in a fast way by the following formulas. We have for $i \geq 2$

$$p_0^{(\mathbf{b}^{(i)})}(x) = 1, \quad p_n^{(\mathbf{b}^{(i)})}(x) = \begin{cases} 0, & n \not\equiv 0 \pmod{i} \\ \binom{x}{k} \frac{(ik)!}{(i!)^k} = \frac{(ik)!}{k!(i!)^k} \prod_{h=1}^k (x-h+1), & n = ki \end{cases}. \quad (13)$$

and when $i = 1$ obviously

$$\forall n \geq 0 \quad p_n^{(\mathbf{b}^{(1)})}(x) = x^n \quad (14)$$

The equalities (13) and (14) easily follow by relation (9) in Proposition 3 and Definition 3 of partial ordinary Bell polynomials, or considering for all $i \geq 1$ the e.g.f.s $s_e \left(\left(\mathbf{b}^{(i)} \right)^x \right)$.

Now we can explicitly define an isomorphism between $\left(H_R^{(n-1)}, + \right)$ and $\left(U_R^{(n)}, \star \right)$ by means of the basis \mathcal{B} .

Theorem 3. Let $\tau^{(n)} : \left(H_R^{(n-1)}, + \right) \rightarrow \left(U_R^{(n)}, \star \right)$ be defined as follows

$$\tau^{(n)} : \mathbf{x} = (x_0, \dots, x_{n-2}) \mapsto \prod_{k=1}^{n-1} \left(\mathbf{b}^{(k,n)} \right)^{x_{k-1}} = \prod_{k=1}^{n-1} p^{(\mathbf{b}^{(k,n)})}(x_{k-1})$$

where $\mathbf{b}^{(k,n)}$ is the sequence $\mathbf{b}^{(k)}$ of length n , $n \geq 2$. Then $\tau^{(n)}$ is an isomorphism.

Proof. Given any $\mathbf{x}, \mathbf{y} \in H_R^{(n-1)}$ as a straightforward consequence of (7) we have

$$\tau^{(n)}(\mathbf{x} + \mathbf{y}) = \tau^{(n)}(\mathbf{x}) \star \tau^{(n)}(\mathbf{y}).$$

Moreover, given any $\mathbf{a} = (1, a_1, \dots, a_{n-1}) \in U_R^{(n)}$ there exists one and only one sequence $\mathbf{x} \in H_R^{(n-1)}$ such that $\tau^{(n)}(\mathbf{x}) = \mathbf{a}$. To prove this claim, first of all we describe explicitly the terms of $\tau^{(n)}(\mathbf{x})$, using equations (13), (14), and the definition of Hurwitz product \star

$$\begin{cases} \tau^{(n)}(\mathbf{x})[0] = 1, \\ \tau^{(n)}(\mathbf{x})[1] = x_0, \\ \tau^{(n)}(\mathbf{x})[m] = m! \sum_{\sum_{i=1}^m l_{j_i} = m} \frac{x_0^{j_1}}{j_1!} \prod_{h=2}^m \frac{(x_{h-1})}{(h!)^{j_h}}, \quad m = 2, \dots, n-1 \end{cases} \quad (15)$$

then, if we consider the equation $\tau^{(n)}(\mathbf{x}) = \mathbf{a}$, we can solve it determining in a unique way every element of \mathbf{x} . Indeed, from (15) is straightforward to observe that for $i = 1, \dots, n-1$ the term x_{i-1} appears for the first time and with power 1 in the expression of $\tau^{(n)}(\mathbf{x})[i]$. Thus we obtain from (15) the following system of equations

$$x_0 = a_1, \quad x_1 + x_0^2 = a_2, \quad x_{i-1} + P(x_0, \dots, x_{i-2}) = \tau^{(n)}(\mathbf{x})[i] = a_i, \quad i = 3, \dots, n-1, \quad (16)$$

where

$$P(x_0, \dots, x_{i-2}) = i! \sum_{\sum_{l=1}^{i-1} l_{j_l} = i} \frac{x_0^{j_1}}{j_1!} \prod_{h=2}^{i-1} \frac{(x_{h-1})}{(h!)^{j_h}}$$

and clearly the unique solution to (16) can be recursively computed. \square

Example 1. Given $\mathbf{x} = (x_0, x_1, x_2, x_3) \in H_R^{(4)}$, we have from (15)

$$\tau^{(5)}(\mathbf{x}) = (1, x_0, x_0^2 + x_1, x_0^3 + 3x_0x_1 + x_2, x_0^4 - 3x_1 + 6x_0^2x_1 + 3x_1^2 + 4x_0x_2 + x_3.)$$

Conversely knowing $\mathbf{a} = (1, a_1, a_2, a_3, a_4) \in U_R^{(5)}$, recursively solving the corresponding equations (16), we get the unique solution $\mathbf{x} \in H_R^{(4)}$ to $\tau^{(5)}(\mathbf{x}) = \mathbf{a}$ given by

$$\begin{aligned} x_0 &= a_1, & x_1 &= a_2 - a_1^2 \\ x_2 &= a_3 - P(x_0, x_1) = a_3 - P(a_1, a_2 - a_1^2) = a_3 + 2a_1^3 - 3a_1a_2 \\ x_3 &= a_4 - P(x_0, x_1, x_2) = a_4 - P(a_1, a_2 - a_1^2, a_3 + 2a_1^3 - 3a_1a_2) = \\ &= a_4 - 3a_1^2 - 6a_1^4 + 3a_2 + 12a_1^2a_2 - 3a_2^2 - 4a_1a_3. \end{aligned}$$

As an immediate consequence of the previous theorem, we can define and evaluate the isomorphism

$$\tau : (H_R, +) \rightarrow (U_R, \star)$$

acting as $\tau^{(n+1)}$ on the first n elements of a sequence $\mathbf{x} \in H_R$, for all $n \geq 1$.

Since R is a generic unitary commutative ring, we point out that all the results we have proved also hold if we consider the ring of polynomials $R[x]$, instead of R , dealing with sequences whose terms are polynomials. In the next section, we will consider binomial type sequences, which are strictly connected with the polynomials $p^{(\mathbf{a})}(x)$ and we will use the isomorphism

$$\tau : (H_{R[x]}, +) \rightarrow (U_{R[x]}, \star)$$

in order to give a method for the construction of all the binomial type sequences.

3 Binomial type sequences

A sequence of polynomials $q(x) = (q_n(x))_{n=0}^{+\infty}$, $q_n(x) \in R[x]$, is called a *binomial type sequence* if $q_0(x) = 1$ and

$$q_n(x+y) = \sum_{h=0}^n \binom{n}{h} q_h(x) q_{n-h}(y).$$

In terms of the Hurwitz product \star the above condition can be restated as

$$q(x+y) = q(x) \star q(y),$$

and clearly $q(x) \in U_{R[x]}$.

The binomial type sequences are a very important subject deeply studied during the years. They were introduced in [17] and used in the theory of umbral calculus. On the other hand, binomial type sequences are widely used in combinatorics and probability. For instance, in [18] the author showed the use of binomial type sequences to solve many "tiling" problems and in [15] the author established new relations between binomial type sequences and Bell polynomials. Further interesting results can be found in [14], [10] and [6]. Moreover, many well-known and useful polynomials are binomial type sequences, like Abel, Laguerre and Touchard polynomials.

It is easy to see, as we have already pointed out in (7), that the polynomial sequences of the kind $p^{(\mathbf{a})}(x)$ are binomial type sequences. Moreover, we have the following result, i.e., any binomial type sequence is of the form $p^{(\mathbf{a})}(x)$ for a certain $\mathbf{a} \in U_R$.

Proposition 4. *If $q(x)$ is a binomial type sequence, then there exists $\mathbf{a} \in U_R$ such that $q(x) = p^{(\mathbf{a})}(x)$.*

Proof. It suffices to prove that $s_e(q(x)) = (A(t))^x$, where $A(t) = s_e(\mathbf{a})$ for a suitable sequence $\mathbf{a} \in U_R$. Let c_i be the coefficient of x in the polynomial $q_i(x)$. Then $s_e(q(x)) = \exp(xf(t))$, where $f(t) = \sum_{n=1}^{\infty} c_n \frac{t^n}{n!}$, as proved in [17]. Thus we have only to choose $\mathbf{a} = s_e^{-1}(\exp(f(t)))$. \square

The result showed in the previous proposition has been also stated in [7]. In the next theorem we highlight a novel characterization of the binomial type sequences by means of the isomorphism τ and the basis \mathcal{B} defined in the previous section.

Theorem 4. *For any $\mathbf{u} = (u_0, u_1, \dots) \in H_R$, let $x\mathbf{u} = (xu_0, xu_1, \dots)$ be a sequence of polynomials in $R[x]$, if we consider the isomorphism*

$$\tau : (H_{R[x]}, +) \rightarrow (U_{R[x]}, \star)$$

then $\tau(x\mathbf{u})$ is a binomial type sequence. Conversely, any binomial type sequence is of the form $\tau(x\mathbf{u})$ for a certain sequence $\mathbf{u} \in H_R$.

Proof. The sequence $\tau(x\mathbf{u})$ satisfies the relation

$$\tau((x+y)\mathbf{u}) = \tau(x\mathbf{u} + y\mathbf{u}) = \tau(x\mathbf{u}) \star \tau(y\mathbf{u})$$

and $\tau(x\mathbf{u})[0] = 1$, thus $\tau(x\mathbf{u})$ is a binomial type sequence. On the other hand, since τ is an isomorphism, given a binomial type sequence $q(x) \in U_{R[x]}$ there exists a unique sequence $r(x) = (r_i(x))_{i=0}^{+\infty} \in H_{R[x]}$ such that

$$\tau^{-1}(q(x)) = r(x).$$

Moreover we must have

$$r(x+y) = r(x) + r(y) \quad (17)$$

indeed

$$r(x+y) = \tau^{-1}(q(x+y)) = \tau^{-1}(q(x) \star q(y)) = \tau^{-1}(q(x)) + \tau^{-1}(q(y)) = r(x) + r(y),$$

The equation (17) is satisfied in $H_{R[x]}$ if and only if

$$\forall i \geq 0 \quad r_i(x) = xu_i, \quad u_i \in R.$$

Hence, $q(x) = \tau(x\mathbf{u})$ for a certain sequence $\mathbf{u} \in H_R$. \square

We would like to point out that the previous theorem give a computable method to evaluate binomial type sequences, starting from any sequence in H_R . For example, if we consider the Fibonacci sequence $\mathbf{F} = (1, 1, 2, 3, 5, 8, 13, \dots)$ then, using (15), we obtain the binomial type sequence

$$\tau(x\mathbf{F}) = (1, x, x^2 + x, x^3 + 3x^2 + 2x, x^4 + 6x^3 + 11x^2, x^5 + 10x^4 + 35x^3 + 20x^2 + 5x, \dots).$$

On the other hand, given any binomial type sequence in $U_{R[x]}$, we have a computable method to find the sequence in H_R that generates it. In fact we provide in the following theorem a recursive formula which gives all the terms of the sequence $\mathbf{u} \in H_R$ such that $\tau(x\mathbf{u}) = q(x)$ for a given binomial type sequence $q(x)$.

Theorem 5. *Let us consider a binomial type sequence $q(x) = (q_n(x))_{n=0}^{+\infty}$, where $q_0(x) = 1$ and $q_n(x) = \sum_{i=1}^n c_{i,n}x^i$. Then we have $\tau(x\mathbf{u}) = q(x)$ if*

$$\forall m \geq 1, \quad u_{m-1} = c_{1,m} + \sum_{k|m, k \neq 1, m} \frac{(-1)^{\frac{m}{k}} m!}{\frac{m}{k} (k!)^{\frac{m}{k}}} u_{k-1}. \quad (18)$$

Proof. From the definition of τ and (15) we have

$$\tau^{(n)}(x\mathbf{u})[0] = 1 = q_0(x), \quad \tau^{(n)}(x\mathbf{u})[1] = xu_0 = c_{1,1}x = q_1(x)$$

and for all $m \geq 2$, the polynomial $q_m(x)$ is equal to

$$\tau^{(n)}(x\mathbf{u})[m] = m! \sum_{\substack{j_1 \\ \sum_{l=1}^m l j_l = m}} \frac{(c_{1,1}x)^{j_1}}{j_1!} \prod_{h=2}^m \frac{\binom{u_{h-1}x}{j_h}}{(h!)^{j_h}}. \quad (19)$$

Clearly $u_0 = c_{1,1}$ and, in order to find the value of u_{m-1} for $m \geq 2$, we only need to find the coefficient of x in (19), since it must be equal to $c_{1,m}$. We observe that in the summation showed in (19) we find x multiplied by a suitable coefficient only when we consider solutions of $\sum_{l=1}^m l j_l = m$ having the form $j_k = \frac{m}{k}$, with $j_l = 0$ for all $l \neq k$, where $k|m$ and $k \neq 1$. Indeed, when $k = m$ we obtain the summand $u_{m-1}x$ and, for all $k|m$, $k \neq 1, m$, we easily find from $\binom{u_{k-1}x}{j_k} \frac{m!}{(k!)^{j_k}}$ the summands

$$\frac{(-1)^{j_k+1} m!}{(k!)^{j_k} j_k!} \begin{bmatrix} j_k \\ 1 \end{bmatrix} (u_{k-1}x) = \frac{(-1)^{j_k+1} m!}{j_k (k!)^{j_k}} u_{k-1}x$$

using (10), since $\begin{bmatrix} j_k \\ 1 \end{bmatrix} = (j_k - 1)!$. Thus we have

$$u_{m-1} + \sum_{k|m, k \neq 1, m} \frac{(-1)^{\frac{m}{k}+1} m!}{\frac{m}{k} (k!)^{\frac{m}{k}}} u_{k-1} = c_{1,m},$$

from which equation (18) easily follows. \square

Remark 3. *As a straightforward consequence we observe that if p is a prime number then $u_{p-1} = c_{1,p}$.*

It would be interesting for further research to focus on the case $R = \mathbb{Z}$, in order to find new relations between integer sequences and polynomial sequences of binomial type. We provide some examples listed in the table below, where, for some well-known binomial type sequences, we give the first terms of the corresponding sequence arising from (18).

| Binomial type polynomials | Expression | Corresponding \mathbf{u} |
|---------------------------|---|--|
| Increasing powers | x^n | $\bar{\mathbf{1}} = (1, 0, 0, 0, \dots)$ |
| Laguerre | $L_n(x) = \sum_{k=0}^n \frac{n!}{k!} \binom{n-1}{k-1} (-x)^k$ | $(-1, -2, -6, -30, -120, -720, -5040, \dots)$ |
| Touchard | $T_n(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k$ | $(1, 1, 1, 4, 1, -19, 1, 771, -559, \dots)$ |
| Abel | $A_n(x, a) = x(x - an)^{n-1}$ | $(1, -2a, 9a^2, -6a - 64a^3, 625a^4, \dots)$ |
| Pochhammer | $(x)_n = x(x+1)(x+2) \cdots (x+n-1)$ | $(1, 1, 2, 9, 24, 110, 720, 5985, 39200, \dots)$ |

As far as we know, none of the sequences \mathbf{u} different from $\bar{\mathbf{1}}$, seems to be already recorded in Oeis [16].

References

- [1] J. P. Allouche, M. M. France, *Hadamard grade of power series*, Journal of Number Theory, Vol. **131**, No. **11**, 2013–2022, 2011.
- [2] S. Barbero, U. Cerruti, N. Murru, *Some combinatorial properties of the Hurwitz series ring*, Ricerche di Matematica, to Appear, <https://doi.org/10.1007/s11587-017-0336-x>, 2017.
- [3] A. Benhissi, *Ideal structure of Hurwitz series rings*, Contributions to Algebra and Geometry, Vol. **48**, No. **1**, 251–256, 2007.
- [4] A. Benhissi, F. Koja, *Basic properties of Hurwitz series rings*, Ricerche di Matematica, Vol. **61**, No. **2**, 255–273, 2012.
- [5] C.A. Charalambides, *Enumerative Combinatorics*, Chapman and Hall, 2002
- [6] A. Di Bucchianico, D. Loeb, *Sequences of binomial type with persistent roots*, Journal of Mathematical Analysis and Applications, Vol. **199**, No. **1**, 39–58, 1996.
- [7] A. Di Bucchianico, *Probabilistic and analytic aspects of the umbral calculus*, Amsterdam, Centrum Wiskunde and Informatica (CWI), 1997.
- [8] J. A. Fill. P. Flajolet, N. Kapur, *Singularity analysis, Hadamard products, and tree recurrences*, J. Comput. Appl. Math., Vol. **174**, 271–313, 2005.
- [9] M. Ghanem, *Some properties of Hurwitz series ring*, International Mathematical Forum, Vol. **6**, No. **40**, 1973–1981, 2007.
- [10] D. Goss, *Polynomials of binomial type and Lucas’ theorem*, Proc. Amer. Math. Soc., Vol. **144**, 1897–1904, 2016.
- [11] R. Graham, D. Knuth, O. Patashnik, *Concrete mathematics: a foundation for computer science*, Addison–Wesley, 2nd Edition, 1994.

- [12] J. Hadamard, *Theorem sur les series entieres*, Acta Math., Vol. **22**, 55–63, 1899.
- [13] W. F. Keigher, F. L. Pritchard, *Hurwitz series as formal functions*, Journal of Pure and Applied Algebra, Vol. **146**, 291–304, 2000.
- [14] V. V. Kisil, *Polynomial sequences of binomial type path integrals*, Annals of Combinatorics, Vol. **6**, No. **1**, 45–56, 2002.
- [15] M. Mihoubi, *Bell polynomials and binomial type sequences*, Discrete Mathematics, Vol. **308**, 2450–2459, 2008.
- [16] The On–Line Encyclopedia of Integer Sequences, <http://oeis.org/>
- [17] G. C. Rota, D. Kahaner, A. Odlyzko, *Finite operator calculus*, J. Math. Anal. Appl., Vol. **42**, 685–760, 1973.
- [18] J. Schneider, *Polynomial sequences of binomial–type arising in graph theory*, The Electronic Journal of Combinatorics, Vol. **21**, No. **1**, P.1.43, 2014.