

ARITHMETIC PROGRESSIONS REPRESENTED BY DIAGONAL TERNARY QUADRATIC FORMS

HAI-LIANG WU AND ZHI-WEI SUN

ABSTRACT. Let $d > r \geq 0$ be integers. For positive integers a, b, c , if any term of the arithmetic progression $\{r + dn : n = 0, 1, 2, \dots\}$ can be written as $ax^2 + by^2 + cz^2$ with $x, y, z \in \mathbb{Z}$, then the form $ax^2 + by^2 + cz^2$ is called (d, r) -universal. In this paper, via the theory of ternary quadratic forms we study the (d, r) -universality of some diagonal ternary quadratic forms conjectured by L. Pehlivan and K. S. Williams, and Z.-W. Sun. For example, we prove that $2x^2 + 3y^2 + 10z^2$ is $(8, 5)$ -universal, $x^2 + 3y^2 + 8z^2$ and $x^2 + 2y^2 + 12z^2$ are $(10, 1)$ -universal and $(10, 9)$ -universal, and $3x^2 + 5y^2 + 15z^2$ is $(15, 8)$ -universal.

1. INTRODUCTION

Let $\mathbb{N} = \{0, 1, 2, \dots\}$. The Gauss-Legendre theorem on sums of three squares states that $\{x^2 + y^2 + z^2 : x, y, z \in \mathbb{Z}\} = \mathbb{N} \setminus \{4^k(8l + 7) : k, l \in \mathbb{N}\}$. A classical topic in the study of number theory asks, given a quadratic polynomial f and an integer n , how can we decide when f represents n over the integers? This topic has been extensively investigated. It is known that for any $a, b, c \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$ the exceptional set

$$E(a, b, c) = \mathbb{N} \setminus \{ax^2 + by^2 + cz^2 : x, y, z \in \mathbb{Z}\}$$

is infinite, see, e.g., [4].

An integral quadratic form f is called *regular* if it represents each integer represented by the genus of f . L. E. Dickson [3, pp. 112-113] listed all the 102 regular ternary quadratic forms $ax^2 + by^2 + cz^2$ together with the explicit characterization of $E(a, b, c)$, where $1 \leq a \leq b \leq c \in \mathbb{Z}^+$ and $\gcd(a, b, c) = 1$. In this direction, W. C. Jagy, I. Kaplansky and A. Schiemann [7] proved that there are at most 913 regular positive definite integral ternary quadratic forms.

By the Gauss-Legendre theorem, for any $n \in \mathbb{N}$ we can write $4n + 1 = x^2 + y^2 + z^2$ with $x, y, z \in \mathbb{Z}$. It is also known that for any $n \in \mathbb{N}$ we can

2010 *Mathematics Subject Classification*. Primary 11E25; Secondary 11B25, 11E20.

Keywords. Arithmetic progressions, representations of integers, ternary quadratic forms.

Supported by the National Natural Science Foundation of China (Grant No. 11571162).

write $2n + 1$ as $x^2 + y^2 + 2z^2$ (or $x^2 + 2y^2 + 3z^2$, or $x^2 + 2y^2 + 4z^2$) with $x, y, z \in \mathbb{Z}$ (see, e.g., Kaplansky [10]). Thus, it is natural to introduce the following definition.

Definition 1.1. Let $d \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$ and $r \in \{0, \dots, d - 1\}$. For $a, b, c \in \mathbb{Z}$, if any $dn + r$ with $n \in \mathbb{N}$ can be written as $ax^2 + by^2 + cz^2$ with $x, y, z \in \mathbb{Z}$, then we say that the ternary quadratic form $ax^2 + by^2 + cz^2$ is (d, r) -universal.

In 2008, A. Alaca, S. Alaca and K. S. Williams [1] proved that there is no binary positive definite quadratic form which can represent all non-negative integers in a residue class. B.-K. Oh [13] showed that for some $U(x, y) \in \mathbb{Q}[x, y]$ the discriminant of any (d, r) -universal positive definite integral ternary quadratic form does not exceed $U(d, r)$.

Z.-W. Sun [17] proved that $x^2 + 3y^2 + 24z^2$ is $(6, 1)$ -universal. Moreover, in 2017 he [18, Remark 3.1] confirmed his conjecture that for any $n \in \mathbb{Z}^+$ and $\delta \in \{0, 1\}$ we can write $6n + 1$ as $x^2 + 3y^2 + 6z^2$ with $x, y, z \in \mathbb{Z}$ and $x \equiv \delta \pmod{2}$. This implies that $4x^2 + 3y^2 + 6z^2$ and $x^2 + 12y^2 + 6z^2$ are $(6, 1)$ -universal. On August 2, 2017 Sun [19] published on OEIS his list (based on his computation) of all possible candidates of (d, r) -universal irregular ternary quadratic forms $ax^2 + by^2 + cz^2$ with $1 \leq a \leq b \leq c$ and $3 \leq d \leq 30$. For example, he conjectured that

$$x^2 + 3y^2 + 7z^2, \quad x^2 + 3y^2 + 42z^2, \quad x^2 + 3y^2 + 54z^2$$

are all $(6, 1)$ -universal, $x^2 + 7y^2 + 14z^2$ is $(7, 1)$ -universal and $x^2 + 2y^2 + 7z^2$ is $(7, r)$ -universal for each $r = 1, 2, 3$. In 2018 L. Pehlivan and K. S. Williams [14] also investigated such problems independently, actually they studied (d, r) -universal quadratic forms $ax^2 + by^2 + cz^2$ with $1 \leq a \leq b \leq c$ and $3 \leq d \leq 11$.

Pehlivan and Williams [14] considered the $(8, 1)$ -universality of $x^2 + 8y^2 + 24z^2$, $x^2 + 2y^2 + 64z^2$ and $x^2 + 8y^2 + 64z^2$ open. However, B. W. Jones and G. Pall [9] proved in 1939 that for any $n \in \mathbb{N}$ we can write

$$8n + 1 = x^2 + 8y^2 + 64z^2 = x^2 + 2(2y)^2 + 64z^2$$

with $x, y, z \in \mathbb{Z}$, and hence $x^2 + 2y^2 + 64z^2$ and $x^2 + 8y^2 + 64z^2$ are indeed $(8, 1)$ -universal. As $8x(x + 1)/2 + 1 = (2x + 1)^2$, the $(8, 1)$ -universality of $x^2 + 8y^2 + 24z^2$ is obviously equivalent to $\{x(x + 1)/2 + y^2 + 3z^2 : x, y, z \in \mathbb{Z}\} = \mathbb{N}$, which was conjectured by Sun [16] and confirmed in [5].

The first part and Part (ii) with $i \in \{2, 3\}$ of the following result were conjectured by Pehlivan and Williams [14], as well as Sun [19].

Theorem 1.1. (i) *The form $2x^2 + 3y^2 + 10z^2$ is $(8, 5)$ -universal.*

(ii) *Let $n \in \mathbb{Z}^+$, $\delta \in \{1, 9\}$ and $i \in \{1, 2, 3\}$. Then $10n + \delta = x_1^2 + 2x_2^2 + 3x_3^2$ for some $(x_1, x_2, x_3) \in \mathbb{Z}^3$ with $2 \mid x_i$.*

Kaplansky [10] showed that there are at most 23 positive definite integral ternary quadratic forms that can represent all positive odd integers (19 for sure and 4 plausible candidates, see also Jagy [6] for further progress). Using one of the 19 forms, we obtain the following result originally conjectured by Sun [19].

Theorem 1.2. *The forms $x^2 + 3y^2 + 14z^2$ and $2x^2 + 3y^2 + 7z^2$ are both $(14, 7)$ -universal.*

Now we turn to study Sun's conjectural $(15, r)$ -universality of some positive definite integral ternary quadratic forms.

Theorem 1.3. (i) *For any $n \in \mathbb{N}$ and $i \in \{1, 2, 3\}$, there exists $(x_1, x_2, x_3) \in \mathbb{Z}^3$ with $3 \mid x_i$ such that $15n + 5 = 2x_1^2 + 3x_2^2 + 5x_3^2$.*

(ii) *The form $x^2 + y^2 + 15z^2$ is $(15, 5r)$ -universal for $r = 1, 2$, and $3x^2 + 3y^2 + 5z^2$ is $(15, 5)$ -universal.*

(iii) *For any $r = 1, 2$, both $x^2 + y^2 + 30z^2$ and $2x^2 + 3y^2 + 5z^2$ are $(15, 5r)$ -universal. Also, the forms $x^2 + 6y^2 + 15z^2$ and $3x^2 + 3y^2 + 10z^2$ are $(15, 10)$ -universal.*

(iv) *The form $x^2 + 2y^2 + 15z^2$ is $(15, 3r)$ -universal for each $r = 1, 2, 3, 4$, and the form $3x^2 + 5y^2 + 10z^2$ is $(15, 3r)$ -universal for $r = 1, 4$.*

Theorem 1.4. (i) *The form $x^2 + 3y^2 + 5z^2$ is $(15, 3r)$ -universal for each $r = 1, 2, 3, 4$. Also, $x^2 + 5y^2 + 15z^2$ is $(15, 3r)$ -universal for $r = 2, 3$.*

(ii) *The form $x^2 + 3y^2 + 15z^2$ is $(15, r)$ -universal for each $r \in \{1, 7, 13\}$. Also, the form $x^2 + 15y^2 + 30z^2$ is $(15, r)$ -universal for $r = 1, 4$, and the form $x^2 + 10y^2 + 15z^2$ is $(15, r)$ -universal for all $r \in \{4, 11, 14\}$.*

(iii) *The form $3x^2 + 5y^2 + 6z^2$ is $(15, r)$ -universal for each $r \in \{8, 11, 14\}$. Also, $3x^2 + 5y^2 + 15z^2$ and $3x^2 + 5y^2 + 30z^2$ are both $(15, 8)$ -universal.*

Remark 1.1. Our proof of Theorem 1.4 relies heavily on the genus theory of quadratic forms as well as the Siegel-Minkowski formula.

We will give a brief overview of the theory of ternary quadratic forms in the next section, and show Theorem 1.1-1.4 in Sections 3-5 respectively.

2. SOME PREPARATIONS

Let

$$f(x, y, z) = ax^2 + by^2 + cz^2 + ryz + szx + txy \quad (2.1)$$

be a positive definite ternary quadratic form with integral coefficients. Its associated matrix is

$$A = \begin{pmatrix} 2a & t & s \\ t & 2b & r \\ s & r & 2c \end{pmatrix}.$$

The discriminant of f is defined by $d(f) := \det(A)/2$.

The following lemma is a fundamental result on integral representations of quadratic forms (cf. [2, pp.129]).

Lemma 2.1. *Let f be a nonsingular integral quadratic form and let m be a nonzero integer represented by f over the real field \mathbb{R} and the ring \mathbb{Z}_p of p -adic integers for each prime p . Then m is represented by some form f^* over \mathbb{Z} with f^* in the same genus of f .*

Now, we introduce some standard notations in the theory of quadratic forms which can be found in [2, 11, 15]. For the positive definite ternary quadratic form f given by (2.1), $\text{Aut}(f)$ denotes the group of integral isometries of f . For $n \in \mathbb{N}$, write

$$r(n, f) := |\{(x, y, z) \in \mathbb{Z}^3 : f(x, y, z) = n\}|$$

(where $|S|$ denotes the cardinality of a set S), and let

$$r(n, \text{gen}(f)) := \sum_{f^* \in \text{gen}(f)} \frac{r(n, f^*)}{|\text{Aut}(f^*)|},$$

where the summation is over a set of representatives of the classes in the genus of f .

We [21] also need our earlier result obtained from the Siegel-Minkowski formula and the knowledge of local densities.

Lemma 2.2. ([21, Lemma 4.1]) *Let f be a positive ternary quadratic form with discriminant $d(f)$. Suppose that $m \in \mathbb{Z}^+$ is represented by $\text{gen}(f)$. Then for each prime $p \nmid 2md(f)$, we have*

$$\frac{r(mp^2, \text{gen}(f))}{r(m, \text{gen}(f))} = p + 1 - \left(\frac{-md(f)}{p} \right), \quad (2.2)$$

where $\left(\frac{\cdot}{p} \right)$ is the Legendre symbol.

3. PROOF OF THEOREM 1.1

Lemma 3.1. *For any $n \in \mathbb{Z}^+$ and $\delta \in \{1, 9\}$, we can write $10n + \delta = x^2 + 2y^2 + 3z^2$ with $x, y, z \in \mathbb{Z}$ and $y^2 + z^2 \neq 0$.*

Proof. By [3, pp.112–113] we can write $10n + \delta = x^2 + 2y^2 + 3z^2$ with $x, y, z \in \mathbb{Z}$; if $10n + \delta$ is not a square then $y^2 + z^2$ is obviously nonzero.

Now suppose that $10n + \delta = m^2$ for some $m \in \mathbb{N}$. As $n > 0$, we have $m > 1$.

Case 1. m has a prime factor $p > 3$.

In this case, by Lemma 2.2 we have

$$r(p^2, x^2 + 2y^2 + 3z^2) = 2 \left(p + 1 - \left(\frac{-6}{p} \right) \right).$$

Hence, $r(m^2, x^2 + 2y^2 + 3z^2) \geq r(p^2, x^2 + 2y^2 + 3z^2) > 2$. Thus, for some $(r, s, t) \in \mathbb{Z}^3$ with $s^2 + t^2 \neq 0$ we have $10n + \delta = m^2 = r^2 + 2s^2 + 3t^2$.

Case 2. $10n + \delta = m^2 = 3^{2k}$ with $k \in \mathbb{Z}^+$.

In this case,

$$10n + \delta = 3^{2k} = (2 \times 3^{k-1})^2 + 2 \times (3^{k-1})^2 + 3 \times (3^{k-1})^2.$$

In view of the above, we have completed the proof. \square

Lemma 3.2. *If $n = 2x^2 + 3y^2 > 0$ with $x, y \in \mathbb{Z}$ and $5 \mid n$, then we can write $n = 2u^2 + 3v^2$ with $u, v \in \mathbb{Z}$ and $5 \nmid uv$.*

Proof. We use induction on $k = \text{ord}_5(\text{gcd}(x, y))$, the 5-adic order of the greatest common divisor of x and y .

When $k = 0$, the desired result holds trivially.

Now let $k \geq 1$ and assume the desired result for smaller values of k . Write $x = 5^k x_0$ and $y = 5^k y_0$, where x_0 and y_0 are integers not all divisible by 5. Then $x_0 + 6y_0$ or $x_0 - 6y_0$ is not divisible by 5. Hence we may choose $\varepsilon \in \{\pm 1\}$ such that $5 \nmid x_0 + 6\varepsilon y_0$. Set $x_1 = 5^{k-1}(x_0 + 6\varepsilon y_0)$ and $y_1 = 5^{k-1}(4x_0 - \varepsilon y_0)$. Then $\text{ord}_5(\text{gcd}(x_1, y_1)) = k - 1$. Note that

$$5^{2k}(2x_0^2 + 3y_0^2) = 5^{2k-2}(2(x_0 + 6\varepsilon y_0)^2 + 3(4x_0 - \varepsilon y_0)^2) = 2x_1^2 + 3y_1^2.$$

So, applying the induction hypothesis we immediately obtain the desired result. \square

Proof of Theorem 1.1. (i) It is easy to see that $8n + 5$ can be represented by the genus of $f(x, y, z) = 2x^2 + 3y^2 + 10z^2$. There are two classes in the

genus of f , and the one not containing f has the representative $g(x, y, z) = 3x^2 + 5y^2 + 5z^2 + 2yz - 2zx + 2xy$. It is easy to verify the following identity:

$$f\left(\frac{x}{2} + y - z, y + z, \frac{x}{2}\right) = g(x, y, z). \quad (3.1)$$

Suppose that $8n + 5 = g(x, y, z)$ for some $x, y, z \in \mathbb{Z}$. Then

$$1 \equiv 8n + 5 = g(x, y, z) \equiv 3x^2 + (y + z)^2 + 2x(y - z) \pmod{4}.$$

Hence $y \not\equiv z \pmod{2}$ and $2 \mid x$. In light of the identity (3.1), $8n + 5$ is represented by f over \mathbb{Z} .

By Lemma 2.1 and the above, $8n + 5$ can be represented by $2x^2 + 3y^2 + 10z^2$ over \mathbb{Z} .

(ii) Let $h(x, y, z) = x^2 + 2y^2 + 3z^2$. By [3, pp.112–113], we can write $10n + \delta = h(x, y, z)$ for some $x, y, z \in \mathbb{Z}$.

We claim that there are $u, v, w \in \mathbb{Z}$ with $u - 2v + 4w \equiv 0 \pmod{5}$ such that $10n + \delta = h(u, v, w)$. Here we handle the case $\delta = 1$. (The case $\delta = 9$ can be handled similarly.)

Case 1. $x^2 \equiv -1 \pmod{5}$.

It is easy to see that $y^2 \equiv 0 \pmod{5}$ and $z^2 \equiv -1 \pmod{5}$, or $y^2 \equiv 1 \pmod{5}$ and $z^2 \equiv 0 \pmod{5}$. When $y^2 \equiv 0 \pmod{5}$ and $z^2 \equiv -1 \pmod{5}$, without loss of generality we may assume that $z \equiv x \pmod{5}$ (otherwise, we may replace z by $-z$). If $y^2 \equiv 1 \pmod{5}$ and $z^2 \equiv 0 \pmod{5}$, then we simply assume $y \equiv -2x \pmod{5}$ without loss of generality. Note that our choice of y and z meets the requirement $x - 2y + 4z \equiv 0 \pmod{5}$.

Case 2. $x^2 \equiv 0 \pmod{5}$.

Clearly, we have $y^2 \equiv -1 \pmod{5}$ and $z^2 \equiv 1 \pmod{5}$. Without loss of generality, we may assume that $y \equiv 2z \pmod{5}$ and hence $x - 2y + 4z \equiv 0 \pmod{5}$.

Case 3. $x^2 \equiv 1 \pmod{5}$.

Apparently, we have $y^2 \equiv z^2 \pmod{5}$. By Lemmas 3.1 and 3.2, we may simply assume that $5 \nmid yz$. When $y^2 \equiv z^2 \equiv x^2 \equiv 1 \pmod{5}$, without loss of generality we may assume that $x \equiv y \equiv -z \pmod{5}$. If $y^2 \equiv z^2 \equiv (2x)^2 \equiv -1 \pmod{5}$, then we may assume that $y \equiv z \equiv 2x \pmod{5}$ without any loss of generality. So, in this case our choice of y and z also meets the requirement $x - 2y + 4z \equiv 0 \pmod{5}$.

In view of the above analysis, we may simply assume $x - 2y + 4z \equiv 0 \pmod{5}$ without any loss of generality. Note that $h(x, y, z) = h(x^*, y^*, z^*)$,

where

$$\begin{aligned} z^* &= \frac{x - 2y + 4z}{5} \not\equiv z \pmod{2}, \\ x^* &= 2y - z + 2z^* \not\equiv x \pmod{2}, \\ y^* &= y - 3z + 3z^* \not\equiv y \pmod{2}. \end{aligned}$$

So we have the desired result in part (ii) of Theorem 1.1. \square

4. PROOFS OF THEOREMS 1.2-1.3

Proof of Theorem 1.2. By [10], we can write $2n + 1 = F(r, s, t)$ with $r, s, t \in \mathbb{Z}$, where $F(x, y, z) = x^2 + 3y^2 + 2yz + 5z^2$. Since

$$(2r - 3t)^2 + 3(r + 2t)^2 + 14s^2 = 7F(r, s, t)$$

and

$$2(s + 3t)^2 + 3(2s - t)^2 + 7r^2 = 7F(r, s, t),$$

we see that $7(2n + 1)$ is represented by the form $x^2 + 3y^2 + 14z^2$ as well as the form $2x^2 + 3y^2 + 7z^2$. \square

Proof of Theorem 1.3. (i) By [3, pp.112–113], we may write $3n + 1 = r^2 + s^2 + 6t^2$ with $r, s, t \in \mathbb{Z}$. One may easily verify the following identities:

$$\begin{aligned} 5(r^2 + s^2 + 6t^2) &= 2(r \pm 3t)^2 + 3(r \mp 2t)^2 + 5s^2 \\ &= 2(s \pm 3t)^2 + 3(s \mp 2t)^2 + 5r^2. \end{aligned}$$

As exactly one of r and s is divisible by 3, one of the the four numbers $r \pm 2t$ and $s \pm 2t$ is a multiple of 3. This proves part (i) of Theorem 1.3.

(ii) Let $r \in \{1, 2\}$. By [3, pp.112–113], for some $x, y, z \in \mathbb{Z}$ we have $3n + r = x^2 + y^2 + 3z^2$. Hence

$$15n + 5r = 5(x^2 + y^2 + 3z^2) = (x + 2y)^2 + (2x - y)^2 + 15z^2.$$

By [3, pp.112–113], we may write $3n + 1 = u^2 + 3v^2 + 3w^2$ with $u, v, w \in \mathbb{Z}$. Thus

$$15n + 5 = 5(u^2 + 3v^2 + 3w^2) = 3(v + 2w)^2 + 3(2v - w)^2 + 5u^2.$$

(iii) Let $r \in \{1, 2\}$. By [3, pp.112–113], there are $u, v, w \in \mathbb{Z}$ such that $3n + r = r^2 + s^2 + 6t^2$. There are two classes in the genus of the form $x^2 + y^2 + 30z^2$, and the one not containing $x^2 + y^2 + 30z^2$ has a representative

$2x^2 + 3y^2 + 5z^2$. Since

$$\begin{aligned} 15n + 5r &= 5(x^2 + y^2 + 6z^2) \\ &= (x + 2y)^2 + (2x - y)^2 + 30z^2 \\ &= 2(x + 3z)^2 + 3(x - 2z)^2 + 5y^2, \end{aligned}$$

we see that $15n + 5r$ is represented by $x^2 + y^2 + 30z^2$ as well as $2x^2 + 3y^2 + 5z^2$.

By [3, pp.112–113], we can write $3n + 2 = 2u^2 + 3v^2 + 3w^2$ with $u, v, w \in \mathbb{Z}$. There are two classes in the genus of $x^2 + 6y^2 + 15z^2$, and the one not containing $x^2 + 6y^2 + 15z^2$ has a representative $3x^2 + 3y^2 + 10z^2$. As

$$\begin{aligned} 15n + 10 &= 5(2u^2 + 3v^2 + 3w^2) \\ &= (2u + 3v)^2 + 6(u - v)^2 + 15w^2 \\ &= 3(u + 2v)^2 + 3(2v - w)^2 + 10u^2, \end{aligned}$$

we see that $15n + 10$ is represented by $x^2 + 6y^2 + 15z^2$ as well as $3x^2 + 3y^2 + 10z^2$.

(iv) Let $r \in \{1, 2, 3, 4\}$. By [3, pp.112–113], there are $x, y, z \in \mathbb{Z}$ such that $5n + r = x^2 + 2y^2 + 5z^2$. Hence

$$15n + 3r = 3(x^2 + 2y^2 + 5z^2) = (x - 2y)^2 + 2(x + y)^2 + 15z^2.$$

Now let $r \in \{1, 4\}$. By [3, pp.112–113], there are $u, v, w \in \mathbb{Z}$ such that $5n + r = u^2 + 5v^2 + 10w^2$. Thus,

$$15n + 3r = 3(u^2 + 5v^2 + 10w^2) = 3u^2 + 5(v - 2w)^2 + 10(v + w)^2.$$

In view of the above, we have completed the proof of Theorem 1.3. \square

5. PROOF OF THEOREM 1.4

Proof of Theorem 1.4(i). Let $r \in \{1, 2, 3, 4\}$. It is easy to see that $15n + 3r$ can be represented by $f_1(x, y, z) = x^2 + 3y^2 + 5z^2$ locally. There are two classes in the genus of f_1 , and the one not containing f_1 has a representative $f_2(x, y, z) = x^2 + 2y^2 + 8z^2 - 2yz$. One may easily verify the following identities:

$$f_1 \left(\frac{x - y - z}{3} - 2z, \frac{x - y - z}{3} + y, \frac{x - y - z}{3} + z \right) = f_2(x, y, z), \quad (5.1)$$

$$f_1 \left(\frac{x + y + z}{3} + 2z, \frac{x + y + z}{3} - y, \frac{x + y + z}{3} - z \right) = f_2(x, y, z). \quad (5.2)$$

Suppose that $15n + 3r = f_2(x, y, z)$ with $x, y, z \in \mathbb{Z}$. Then

$$x^2 - (y + z)^2 \equiv f_2(x, y, z) \equiv 0 \pmod{3},$$

and hence $(x - y - z)/3$ or $(x + y + z)/3$ is an integer. Therefore, by (5.1), (5.2) and Lemma 2.1, we obtain that $x^2 + 3y^2 + 5z^2$ is $(15, 3r)$ -universal.

Now let $r \in \{2, 3\}$. One can easily verify that $15n + 3r$ is represented by the genus of $g_1(x, y, z) = x^2 + 5y^2 + 15z^2$. There are two classes in the genus of g_1 , and the one not containing g_1 has a representative $g_2(x, y, z) = 4x^2 + 4y^2 + 5z^2 + 2xy$. It is easy to verify the identity

$$g_1\left(y + \frac{x + y \mp 5z}{3}, x - \frac{x + y \pm z}{3}, \frac{x + y \pm z}{3}\right) = g_2(x, y, z). \quad (5.3)$$

If $15n + 3r = g_2(x, y, z)$ with $x, y, z \in \mathbb{Z}$, then

$$(x + y)^2 - z^2 \equiv g_2(x, y, z) \equiv 0 \pmod{3}$$

Thus, with the help of (5.3) and Lemma 2.1, we obtain the desired result. \square

Lemma 5.1. (Oh [12]) *Let V be a positive definite ternary quadratic space over \mathbb{Q} . For any isometry $T \in O(V)$ of infinite order,*

$$V_T = \{x \in V : \text{there is a positive integer } k \text{ such that } T^k(x) = x\}$$

is a subspace of V of dimension one, and $T(x) = \det(T)x$ for any $x \in V_T$.

Remark 5.1. Unexplained notations of quadratic space can be found in [2, 11, 15].

Lemma 5.2. *Let $n \in \mathbb{N}$ and $r \in \{1, 7, 13\}$. If we can write $15n + r = f_2(x, y, z) = 3x^2 + 4y^2 + 4z^2 + 2yz$ with $x, y, z \in \mathbb{Z}$, then there are $u, v, w \in \mathbb{Z}$ with $u + 2v - 2w \not\equiv 0 \pmod{3}$ such that $15n + r = f_2(u, v, w)$.*

Proof. Suppose that every integral solution of the equation $f_2(x, y, z) = 15n + r$ satisfies $x + 2y - 2z \equiv 0 \pmod{3}$. We want to deduce a contradiction.

Let

$$T = \begin{pmatrix} 1/3 & 2/3 & -2/3 \\ -2/3 & 2/3 & 1/3 \\ 2/3 & 1/3 & 2/3 \end{pmatrix},$$

and let V be the quadratic space corresponding to f_2 . Since

$$f_2\left(\frac{x + 2y - 2z}{3}, -x + z + \frac{x + 2y - 2z}{3}, x + y - \frac{x + 2y - 2z}{3}\right) = f_2(x, y, z), \quad (5.4)$$

we have $T \in O(V)$. One may easily verify that the order of T is infinite and the space V_T defined in Lemma 5.1 coincides with $\{(0, t, t) : t \in \mathbb{Q}\}$. As $15n + r \neq f_2(0, t, t)$ for any $t \in \mathbb{Z}$, we have $15n + r = f_2(x_0, y_0, z_0)$ for some

$(x_0, y_0, z_0) \in \mathbb{Z}^3 \setminus V_T$. Clearly, the set $\{T^k(x_0, y_0, z_0) : k \geq 0\}$ is infinite and its elements are solutions to the equation $f_2(x, y, z) = 15n + r$. This leads a contradiction since the number of integral representations of any integer by a positive quadratic forms is finite. \square

Lemma 5.3. (Jagy [8]) *If $n = 2x^2 + 2xy + 3y^2$ ($x, y \in \mathbb{Z}$) is a positive integer divisible by 3, then there are $u, v \in \mathbb{Z}$ with $3 \nmid uv$ such that $n = 2u^2 + 2uv + 3v^2$.*

The following lemma is a known result, see, e.g., [8, 9, 12, 18].

Lemma 5.4. *If $n = x^2 + y^2$ ($x, y \in \mathbb{Z}$) is a positive integer divisible by 5, then $n = u^2 + v^2$ for some $u, v \in \mathbb{Z}$ with $5 \nmid uv$.*

Proof of Theorem 1.4(ii). (a) For each $r \in \{1, 7, 13\}$, it is easy to see that $15n + r$ can be represented by the genus of $f_1(x, y, z) = x^2 + 3y^2 + 15z^2$. There are two classes in the genus of $f_1(x, y, z)$, and the one not containing f_1 has a representative $f_2(x, y, z) = 3x^2 + 4y^2 + 4z^2 + 2yz$. One may easily verify the following identities:

$$f_1\left(x - y + z, \frac{x - 2y}{3} - z, \frac{x + y}{3}\right) = f_2(x, y, z), \quad (5.5)$$

$$f_1\left(x + y - z, \frac{x - 2z}{3} - y, \frac{x + z}{3}\right) = f_2(x, y, z). \quad (5.6)$$

Suppose that $15n + r = f_2(x, y, z)$ for some $x, y, z \in \mathbb{Z}$. As $3 \nmid y$ or $3 \nmid z$, when $3 \nmid x$ we may assume that $(x + y)(x + z) \equiv 0 \pmod{3}$ (otherwise we may replace x by $-x$) without loss of generality. If $3 \mid x$ and $y \not\equiv z \pmod{3}$, then $3 \mid yz$ and hence $(x + y)(x + z) \equiv 0 \pmod{3}$. In the remaining case $3 \mid x$ and $y \equiv z \pmod{3}$, we have $x + 2y - 2z \equiv 0 \pmod{3}$; however, we may apply Lemma 5.2 to choose integers $u, v, w \in \mathbb{Z}$ so that $15n + r = f_2(u, v, w)$ and $u + 2v - 2w \not\equiv 0 \pmod{3}$.

In view of the above analysis, there always exist $u, v, w \in \mathbb{Z}$ with $(u + v)(u + w) \equiv 0 \pmod{3}$ such that $15n + r = f_2(u, v, w)$. With the help of (5.5), (5.6), and Lemma 2.1, we obtain the $(15, r)$ -universality of $x^2 + 3y^2 + 15z^2$.

(b) Let $r \in \{1, 4\}$. One can easily verify that $15n + r$ can be represented by $g_1(x, y, z) = x^2 + 15y^2 + 30z^2$ locally. There are two classes in the genus of g_1 , and the one not containing g_1 has a representative $g_2(x, y, z) = 6x^2 + 9y^2 + 10z^2 - 6xy$.

Suppose that $15n + r = g_2(x, y, z)$ with $x, y, z \in \mathbb{Z}$. Clearly $3 \nmid z$. Since $15n + r \neq 10z^2$, by Lemma 5.3 we may assume that x and y are not all divisible by 3. Thus we just need to consider the following two cases.

Case b1. $3 \nmid x$.

When this occurs, without loss of generality, we may assume that $x \equiv -z \pmod{3}$ (otherwise we may replace z by $-z$). In view of the identity

$$g_1\left(x - 3y, \frac{x - 2z}{3}, \frac{x + z}{3}\right) = g_2(x, y, z), \quad (5.7)$$

there are $x^*, y^*, z^* \in \mathbb{Z}$ such that $15n + r = g_1(x^*, y^*, z^*)$.

Case b2. $3 \mid x$ and $3 \nmid y$

In this case, with the help of the identity

$$g_2(x - y, -y, z) = g_2(x, y, z),$$

we return to Case b1 since $x - y \not\equiv 0 \pmod{3}$.

Now applying Lemma Lem2.1 we immediately obtain the $(15, r)$ -universality of $x^2 + 5y^2 + 30z^2$.

(c) For any $r \in \{4, 11, 14\}$, it is easy to verify that $15n + r$ can be represented by $h_1(x, y, z) = x^2 + 10y^2 + 15z^2$ locally. There are two classes in the genus of h_1 , and the one not containing h_1 has a representative $h_2(x, y, z) = 5x^2 + 5y^2 + 6z^2$.

Suppose that $15n + r = h_2(x, y, z)$ with $x, y, z \in \mathbb{Z}$. Since $15n + r \neq 6z^2$, by Lemma 5.4 and the symmetry of x and y , we simply assume $5 \nmid y$ without loss of generality. We claim that we may adjust the signs of x, y, z to satisfy the congruence $(2x + y + 2z)(x - 2y - 2z) \equiv 0 \pmod{5}$.

Case c1. $x^2 \equiv y^2 \pmod{5}$.

Without loss of generality, we may assume $x \equiv y \equiv z \pmod{5}$ if $x^2 \equiv y^2 \equiv z^2 \pmod{5}$, and $x \equiv -y \equiv -2z \pmod{5}$ if $x^2 \equiv y^2 \equiv -z^2 \pmod{5}$. So our claim holds in this case.

Case c2. $x^2 \equiv -y^2 \pmod{5}$.

If $x^2 \equiv -y^2 \equiv z^2 \pmod{5}$, without loss of generality, we may assume that $x \equiv -2y \equiv z \pmod{5}$. If $x^2 \equiv -y^2 \equiv -z^2 \pmod{5}$, we may assume that $x \equiv -2y \equiv 2z \pmod{5}$ without loss of any generality. Thus x, y, z satisfy the desired congruence in our claim.

Case c3. $x^2 \equiv 0 \pmod{5}$.

If $y^2 \equiv z^2 \pmod{5}$, we may assume that $y \equiv -z \pmod{5}$. If $y^2 \equiv -z^2 \pmod{5}$, without loss of generality, we may assume that $y \equiv -2z \pmod{5}$. Clearly, our claim also holds in this case.

In view of the above analysis, there are $x, y, z \in \mathbb{Z}$ with $2x + y + 2z \equiv 0 \pmod{5}$ or $x - 2y - 2z \equiv 0 \pmod{5}$ such that $15n + r = h_2(x, y, z)$. One may easily verify the following identities

$$h_1\left(x - 2y, \frac{2x + y + 2z}{5} - z, \frac{2x + y + 2z}{5}\right) = h_2(x, y, z), \quad (5.8)$$

$$h_1\left(x + 2y, \frac{x - 2y - 2z}{5} + z, \frac{x - 2y - 2z}{5}\right) = h_2(x, y, z). \quad (5.9)$$

With the help of (5.8), (5.9) and Lemma 2.1, the $(15, r)$ -universality of $x^2 + 10y^2 + 15z^2$ is valid. \square

Lemma 5.5. *Let $n \in \mathbb{N}$ and $g_2(x, y, z) = 2x^2 + 8y^2 + 15z^2 - 2xy$. Assume that $15n + 8 = g_2(x, y, z)$ for some $x, y, z \in \mathbb{Z}$ with $y^2 + z^2 \neq 0$. Then $15n + 8 = g_2(u, v, w)$ for some $u, v, w \in \mathbb{Z}$ with $3 \nmid v + w$.*

Proof. Suppose that every integral solution of the equation $g_2(x, y, z) = 15n + 8$ satisfies $y + z \equiv 0 \pmod{3}$. We want to deduce a contradiction.

Let

$$T = \begin{pmatrix} 1 & -2/3 & -2/3 \\ 0 & -1/3 & -4/3 \\ 0 & 2/3 & -1/3 \end{pmatrix}$$

and let V be the quadratic space corresponding to g_2 . Since

$$g_2\left(x + \frac{-2y - 2z}{3}, \frac{-y - 4z}{3}, \frac{2y - z}{3}\right) = g_2(x, y, z).$$

We have $T \in O(V)$. One may easily verify that the order of T is infinite and the space V_T defined in Lemma 5.1 coincides with $\{(t, 0, 0) : t \in \mathbb{Q}\}$. By the assumption in the lemma, we have $15n + 8 = g_2(x_0, y_0, z_0)$ for some $(x_0, y_0, z_0) \in \mathbb{Z}^3 \setminus V_T$. Note that the set $\{T^k(x_0, y_0, z_0) : k \geq 0\}$ is infinite and all elements of this set are solutions to the equation $g_2(x, y, z) = 15n + 8$. This leads to a contradiction since the number of integral representations of any integer by a positive quadratic forms is finite. \square

Proof of Theorem 1.4(iii). (a) Let $r \in \{8, 11, 14\}$. It is easy to see that $15n + r$ can be represented by $f_1(x, y, z) = 3x^2 + 5y^2 + 6z^2$ locally. There are two classes in the genus of f_1 , and the one not containing f_1 has a representative $f_2(x, y, z) = 2x^2 + 6y^2 + 9z^2 + 6yz$.

Suppose that $15n + r = f_2(x, y, z)$ for some $x, y, z \in \mathbb{Z}$. Then $3 \nmid x$.

Case 1. $3 \nmid y$.

In this case, without loss of generality we may assume that $x \equiv -y \pmod{3}$ (otherwise we may replace x by $-x$). In view of the identity

$$f_1\left(\frac{2x-y}{3} - z, -y, \frac{x+y}{3} + z\right) = f_2(x, y, z), \quad (5.10)$$

there are $x^*, y^*, z^* \in \mathbb{Z}$ such that $15n + \delta = f_1(x^*, y^*, z^*)$.

Case 2. $y^2 + z^2 \neq 0$ and $3 \mid y$.

When this occurs, by Lemma 5.3 we may simply assume that $3 \nmid z$. With the help of the identity

$$f_2(x, y+z, -z) = f_2(x, y, z),$$

we return to Case 1.

Case 3. $15n + r = 2m^2$ for some $m \in \mathbb{N}$.

When $15n + r = 2m^2 = 2 \times 2^{2k}$ with $k \geq 1$, we have

$$15n + r = 2 \times 2^{2k} = 3 \times (2^{k-1})^2 + 5 \times (2^{k-1})^2 + 6 \times 0^2.$$

Now suppose that m has a prime factor $p > 5$. By Lemma 2.2 we have

$$r(2p^2, f_1) + r(2p^2, f_2) = 2 \left(p + 1 - \left(\frac{-5}{p} \right) \right) > 10. \quad (5.11)$$

Clearly $r(2p^2, f_1) > 5$ or $r(2p^2, f_2) > 5$. When $r(2p^2, f_1) > 5$, the number $2m^2$ can be represented by f_1 over \mathbb{Z} since $r(2m^2, f_1) \geq r(2p^2, f_1)$. When $r(2p^2, f_2) > 5$, there are $u, v, w \in \mathbb{Z}$ with $v^2 + w^2 \neq 0$ such that $f_2(u, v, w) = 15n + r$. By Lemma 5.3, we return to Case 1 or Case 2.

In view of the above, by applying Lemma 2.1 we get the $(15, r)$ -universality of $3x^2 + 5y^2 + 6z^2$.

(b) It is easy to see that $15n + 8$ can be represented by the genus of $g_1(x, y, z) = 3x^2 + 5y^2 + 15z^2$. There are two classes in the genus of g_1 , and the one not containing g_1 has a representative $g_2(x, y, z) = 2x^2 + 8y^2 + 15z^2 - 2xy$.

Suppose that the equation $15n + 8 = g_2(x, y, z)$ is solvable over \mathbb{Z} . We claim that there are $u, v, w \in \mathbb{Z}$ with $(u+w)(u-v-w) \equiv 0 \pmod{3}$ such that $15n + 8 = g_2(u, v, w)$.

Case 1. $3 \mid x$.

Clearly, $3 \nmid y$. If $3 \nmid z$, without loss of generality we may assume that $z \equiv -y \pmod{3}$ (otherwise we replace z by $-z$). Then $(u, v, w) = (x, y, z)$ meets our purpose.

Case 2. $3 \nmid x$ and $y^2 + z^2 \neq 0$.

In this case, by Lemma 5.5 there are $x', y', z' \in \mathbb{Z}$ with $3 \nmid y' + z'$ such that $15n + 8 = g_2(x', y', z')$. If $x' \equiv y' \pmod{3}$, then by using the identity

$$g_2(x - y, -y, z) = g_2(x, y, z),$$

we return to Case 1. If $x' \not\equiv y' \pmod{3}$, then $3 \mid y'$ and $3 \nmid z'$ since $(x' + y')^2 \equiv 1 \pmod{3}$ and $3 \nmid x'$. Without loss of generality, we may assume that $x' \equiv -z' \pmod{3}$. So $(u, v, w) = (x', y', z')$ meets our purpose.

Case 3. $15n + 8 = 2m^2$ with $m \in \mathbb{N}$.

If $m = 2^k$ for some $k \in \mathbb{Z}^+$, then

$$2m^2 = 3 \times (2^{k-1})^2 + 5 \times (2^{k-1})^2 + 15 \times 0^2.$$

Now suppose that m has a prime factor $p > 5$. By Lemma 2.2, we have

$$r(2p^2, g_1) + r(2p^2, g_2) = 2 \left(p + 1 - \left(\frac{-2}{p} \right) \right) > 10. \quad (5.12)$$

Clearly, $r(2p^2, g_1) > 5$ or $r(2p^2, g_2) > 5$. When $r(2p^2, g_1) > 5$, we have $r(2m^2, g_1) \geq r(2p^2, g_1) > 5$. If $r(2p^2, g_2) > 5$, then there exist $x_0, y_0, z_0 \in \mathbb{Z}$ with $y_0^2 + z_0^2 \neq 0$ such that $15n + 8 = g_2(x_0, y_0, z_0)$. So we are reduced to previous cases.

In view of the proved claim, the (15, 8)-universality of g_1 follows from Lemma 2.1 and the identities

$$g_1 \left(\frac{x - 5z}{3} - y, -y + z, -\frac{x + z}{3} \right) = g_2(x, y, z),$$

$$g_1 \left(\frac{x - y - z}{3} + y + 2z, y - z, \frac{x - y - z}{3} \right) = g_2(x, y, z).$$

(c) One may easily verify that $15n + 8$ can be represented by $h_1(x, y, z) = 3x^2 + 5y^2 + 30z^2$ locally. There are two classes in the genus of h_1 , and the one not containing h_1 has a representative $h_2(x, y, z) = 2x^2 + 15y^2 + 15z^2$.

Suppose that the equation $15n + 8 = h_2(x, y, z)$ for some $x, y, z \in \mathbb{Z}$. In light of Lemma 5.4, we may assume $5 \nmid y$ if $y^2 + z^2 > 0$. We claim that there are $u, v, w \in \mathbb{Z}$ with $(u - v + 2w)(u - 2v + w) \equiv 0 \pmod{5}$ such that $15n + 8 = h_2(u, v, w)$.

Case 1. $y^2 \equiv \varepsilon z^2 \pmod{5}$ and $y^2 + z^2 \neq 0$, where $\varepsilon \in \{\pm 1\}$.

If $y^2 \equiv z^2 \equiv x^2 \pmod{5}$, then we may assume that $x \equiv y \equiv z \pmod{5}$. If $y^2 \equiv z^2 \equiv -x^2 \pmod{5}$, without loss of generality we may assume that $x \equiv -2y \equiv 2z \pmod{5}$. So, $(u, v, w) = (x, y, z)$ meets our requirement in the case $\varepsilon = 1$. The case $\varepsilon = -1$ can be handled similarly.

Case 3. $15n + 8$ is twice a square, say $2m^2$ with $m \in \mathbb{Z}^+$.

When $m = 2^k$ with $k \in \mathbb{Z}^+$, we have

$$15n + 8 = 2 \times 2^{2k} = 3 \times (2^{k-1})^2 + 5 \times (2^{k-1})^2 + 30 \times 0^2.$$

Now assume that m has a prime factor $p > 5$. By Lemma 2.2, we have

$$2r(2p^2, h_1) + r(2p^2, h_2) = 2 \left(p + 1 - \left(\frac{-1}{p} \right) \right) > 10. \quad (5.13)$$

Clearly $r(2p^2, h_1) \geq 4$ or $r(2p^2, h_2) \geq 4$. If $r(2p^2, h_1) \geq 4$, then $r(2m^2, h_1) \geq r(2p^2, h_1) \geq 4$. When $r(2p^2, h_2) \geq 4$, there exist $u, v, w \in \mathbb{Z}$ with $v^2 + w^2 \neq 0$ such that $g_2(u, v, w) = 2m^2$. Thus we are reduced to Case 1.

In view of the proved claim and the identities

$$\begin{aligned} h_2(x, y, z) &= h_1 \left(2y + z, \frac{2x + 3y - z}{5} - z, \frac{x - y + 2z}{5} \right) \\ &= h_1 \left(y + 2z, \frac{2x + y - 3z}{5} + y, \frac{x - 2y + z}{5} \right), \end{aligned}$$

by applying Lemma 2.1 we obtain the (15, 8)-universality of $3x^2 + 5y^2 + 30z^2$. \square

REFERENCES

- [1] A. Alaca, S. Alaca and K. S. Williams, *Arithmetic progressions and binary quadratic forms*, Amer. Math. Monthly **115**(2008), 252–254.
- [2] J. W. S. Cassels, *Rational Quadratic Forms*, Academic Press, London, 1978.
- [3] L. E. Dickson, *Modern Elementary Theory of Numbers*, University of Chicago Press, Chicago, 1939.
- [4] G. Doyle and K. S. Williams, *A positive-definite ternary quadratic form does not represent all positive integers*, Integers **17** (2017), #A41, 19pp (electronic).
- [5] S. Guo, H. Pan and Z.-W. Sun, *Mixed sums of squares and triangular numbers (II)*, Integers **7** (2007), #A56, 5pp (electronic)
- [6] W. C. Jagy, *Five regular or nearly-regular ternary quadratic forms*, Acta Arith, **77** (1996), 361–367.
- [7] W. C. Jagy, I. Kaplansky and A. Schiemann, *There are 913 regular ternary forms*, Mathematika, **44**(1997), 332–341.
- [8] W. C. Jagy, *Integral Positive Ternary Quadratic Forms*, Lecture Notes, 2014.
- [9] B. W. Jones and G. Pall, *Regular and semi-regular positive ternary quadratic forms*, Acta Math. **70**(1939), 165–191.
- [10] I. Kaplansky, *Ternary positive quadratic forms that represent all odd positive integers*, Acta Arith. **70**(1995), 209–214.
- [11] Y. Kitaoka, *Arithmetic of Quadratic Forms*, Cambridge Tracts in Math., Vol. 106, 1993.
- [12] B.-K. Oh, *Ternary universal sums of generalized pentagonal numbers*, J. Korean Math. Soc, **48** (2011) 837–847.
- [13] B.-K. Oh, *Representations of arithmetic progressions by positive definite quadratic forms*, Int. J. Number Theory, **7** (2011), 1603–1614.
- [14] L. Pehlivan and K. S. Williams, *(k, l)-universality of ternary quadratic forms $ax^2 + by^2 + cz^2$* , Integers, **18** (2018), #A20, 44pp (electronic).

- [15] O. T. O'Meara, *Introduction to Quadratic Forms*, Springer-Verlag, New York, 1963.
- [16] Z.-W. Sun, *Mixed sums of squares and triangular numbers*, Acta Arith, **127** (2007), 103–113.
- [17] Z.-W. Sun, *On universal sums of polygonal numbers*, Sci. China Math, **58** (2015), 1367–1396.
- [18] Z.-W. Sun, *On $x(ax+1)+y(by+1)+z(cz+1)$ and $x(ax+b)+y(ay+c)+z(az+d)$* , J. Number Theory **171** (2017), 275–283.
- [19] Z.-W. Sun, Sequence A286885 on OEIS (On-Line Encyclopedia of Integer Sequences), <http://oeis.org/A286885>, August 2, 2018.
- [20] Z.-W. Sun, *Universal sums of three quadratic polynomials*, Sci. China Math., to appear, Doi:10.1007/s11425-017-9354-4. (See also arXiv:1502.03056.)
- [21] H.-L. Wu and Z.-W. Sun, *Some universal quadratic sums over the integers*, preprint, arXiv:1707.06223v4, 2017.

(HAI-LIANG WU) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S REPUBLIC OF CHINA

E-mail address: whl.math@smail.nju.edu.cn

(ZHI-WEI SUN) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S REPUBLIC OF CHINA

E-mail address: zwsun@nju.edu.cn