

Sects and lattice paths over the Lagrangian Grassmannian

Aram Bingham¹ and Özlem Uğurlu²

¹Tulane University, New Orleans; abingham@tulane.edu

²Palm Beach State College, Boca Raton; ugurluo@palmbeachstate.edu

March 19, 2019

Abstract

We examine Borel subgroup orbits in the classical symmetric space of type CI , which are parametrized by skew symmetric (n, n) -clans. We describe bijections between such clans, certain weighted lattice paths, and pattern-avoiding signed involutions, and we give a cell decomposition of the symmetric space in terms of collections of clans called sects. The largest sect with a conjectural closure order is isomorphic (as a poset) to the Bruhat order on partial involutions.

Keywords: Borel orbits, Levi subgroup, Lagrangian Grassmannian, Bruhat order, lattice paths.

MSC: 05A15, 14M15, 14M17

1 Introduction

Let G be a complex reductive group of classical type (SL_n , SO_n , or Sp_{2n}), and θ an automorphism of G of order two. Then we call the fixed point subgroup $L := G^\theta$ a *symmetric subgroup* and G/L a *symmetric space* of classical type. If B is a Borel subgroup of G , then B acts on G/L with finitely many orbits ([11]). The study of Borel orbits and their closures in symmetric spaces imitates and generalizes the study of Borel orbits in flag varieties, bearing comparable combinatorial richness. However, for only three types of classical symmetric spaces, L happens to be a Levi subgroup of a (maximal) parabolic subgroup P ; these are listed in the Table 1 below. This makes it possible to relate the geometry and combinatorics of B -orbits in G/L to those in G/P , via the (B -equivariant) canonical projection map, $\pi : G/L \rightarrow G/P$.

In each of these cases, the homogeneous space G/P parametrizes vector subspaces of \mathbb{C}^n or \mathbb{C}^{2n} which are isotropic with respect to a particular bilinear form, and is often called an

(*isotropic*) *Grassmannian* manifold/variety. Indeed, these symmetric spaces are those which are associated to a polarization of the appropriate vector space; see [7] §11.3.5. The B -orbits of a Grassmannian are called *Schubert cells*, as they are known to give a cell decomposition and an additive basis for (co)homology of the space G/P . Schubert cells can be parametrized by certain lattice paths which are also a tool for understanding their geometry.

Type	Symmetric Pair	B -orbits parametrized by	G/P
<i>AIII</i>	$(SL_{p+q}, S(GL_p \times GL_q))$	(p, q) -clans	$\text{Gr}(p, \mathbb{C}^{p+q})$
<i>CI</i>	(Sp_{2n}, GL_n)	skew-symmetric (n, n) -clans	$\Lambda(n)$
<i>DIII</i>	(SO_{2n}, GL_n)	“type <i>DIII</i> ” (n, n) -clans	$\text{OGr}(n, \mathbb{C}^{2n})$

Table 1: Classical symmetric subgroups which are also spherical Levi subgroups.

B -orbits in G/L are parametrized by objects dubbed *clans* in [10], which have morphed in their development through subsequent works, notably [19], [16], and [4]. In [2], it was shown that π gives G/L the structure of an affine bundle over G/P , and that the pre-images of Schubert cells provide a cell decomposition of G/L . This is used to conclude that the integral Chow rings and cohomology rings of G/L are isomorphic. The type *AIII* case is also treated in detail there, where the pre-images of Schubert cells in the Grassmannian of p -planes in \mathbb{C}^{p+q} , denoted $\text{Gr}_p(\mathbb{C}^{p+q})$, are comprised of collections of (p, q) -clans called *sects*. Each sect contains a unique closed B -orbit, and a unique dense B -orbit. The closures of the dense B -orbits of each sect form a generating set for the integral Chow ring $A^*(G/L)$, akin to Schubert varieties.

In this paper, we apply the ideas of [4] and [2] to the symmetric space of type *CI*, wherein the ambient group is the symplectic group Sp_{2n} and the symmetric subgroup is isomorphic to GL_n . Our first result is Theorem 3.12, which counts Borel orbits in these symmetric spaces by providing a bijection between the parametrizing set of “skew-symmetric” (n, n) -clans and a set of pattern-avoiding signed involutions with known generating function. In Section 4, we describe another bijection of skew symmetric (n, n) -clans with a certain class of *weighted* (n, n) *Delannoy paths*. These are lattice paths in the plane from the origin to the point (n, n) , consisting only of north, east, and northeast diagonal steps, where the diagonal steps can have certain whole number weights.

In Section 5, we describe the sects over the Schubert cells of the Lagrangian Grassmannian $\Lambda(n)$, which is the moduli space of maximal isotropic subspaces of the vector space \mathbb{C}^{2n} with symplectic form Ω . This proceeds in a fashion similar to the type *AIII* case described in [2], where it was also shown that the pre-image of the dense Schubert cell, called the *big sect*, is isomorphic as a poset to the *rook monoid* \mathcal{R}_n with the Bruhat-Chevalley-Renner order. That result relied on a combinatorial description, given by Wyser in [17], of the closure order on B -orbits in the type *AIII* symmetric space.

For a classical symmetric space whose B -orbits are parametrized by a certain family of clans, one can describe the closure poset of clans by the order relation

$$\gamma \leq \tau \iff \overline{Q_\gamma} \subseteq \overline{Q_\tau}$$

for clans γ, τ , with corresponding B -orbits Q_γ, Q_τ . If X is a classical symmetric space of type B, C , or D , then X embeds in some symmetric space X' of type $AIII$. The clans parametrizing B -orbits in X can then be viewed as a subset of the (p, q) -clans parametrizing B' -orbits in X' , where B' is a Borel subgroup of SL_{p+q} and B is the intersection of B' and the relevant symplectic or special orthogonal subgroup. This reflects the fact that a B -orbit in X indexed by a clan γ is exactly the intersection of X with the B' -orbit of X' corresponding to the same clan ([16], Theorem 1.5.8).

From this, one could hope that the closure order on clans in a type B - C - D symmetric space would simply be the restriction of the relevant type $AIII$ closure order. This has been conjectured to be the case in types BI, CI , and CII , but it is known to fail in types DI and $DIII$ ([18], §3.2.2). For this, among other reasons, the analysis for the type $DIII$ symmetric space warrants separate treatment. Nevertheless, there is a *weak order* on clans of a given type, whose order relations are contained within the full closure order, and which can be used to recover the closure order through a simple recursive procedure (see [12]). But the procedure does not appear to easily prove the conjectural closure order for type CI , so we proceed without it.

Our main result of Section 6 says that the big sect over $\Lambda(n)$ with the conjectural closure order is isomorphic, as a poset, to the partial involutions on n letters \mathcal{P}_n with the *Bruhat order* of [1]. The order relations of the latter poset are given by the closure order on congruence orbits of upper triangular matrices acting on symmetric matrices, and they have a convenient combinatorial description which is provided below. Though a clear geometric understanding of this coincidence is wanting, one is led to believe that a similar statement is true about the big sect in the type $DIII$ case as well. In that case, the base space of the bundle is the orthogonal Grassmannian $\text{OGr}(n, \mathbb{C}^{2n})$ of maximally isotropic subspaces with respect to a non-degenerate, symmetric, bilinear form. The specifics of this final case will appear in forthcoming work from the authors.

2 Notation and Preliminaries

All matrix groups in this paper are taken to have entries in the field of complex numbers. Let n be a positive integer. First, we must describe our realization of the type CI symmetric pair (Sp_{2n}, GL_n) , borrowing notation from [16]. Let J_n denote an $n \times n$ matrix with 1's along the anti-diagonal and 0's elsewhere. Let

$$\Omega = \begin{pmatrix} 0 & J_n \\ -J_n & 0 \end{pmatrix}.$$

Then we set

$$G := Sp_{2n} = \{g \in GL_{2n} \mid g^t \Omega g = \Omega\}. \tag{2.1}$$

Let $\text{int}(g) : GL_{2n} \rightarrow GL_{2n}$ denote the map defined by

$$\text{int}(g)(h) = ghg^{-1}.$$

Now define the matrix

$$I_{n,n} := \begin{pmatrix} I_n & 0 \\ 0 & -I_n \end{pmatrix},$$

where I_n denotes the $n \times n$ identity matrix. Then we have an automorphism θ of order two on GL_{2n} defined by $\theta := \text{int}(iI_{n,n})$. Indeed, $(iI_{n,n})^{-1} = -iI_{n,n}$, so if

$$g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

is the $n \times n$ block form of g , we have

$$\theta(g) = \begin{pmatrix} iI_n & 0 \\ 0 & -iI_n \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} -iI_n & 0 \\ 0 & iI_n \end{pmatrix} = \begin{pmatrix} A & -B \\ -C & D \end{pmatrix}.$$

Observe that the restriction of θ to Sp_{2n} induces an order two automorphism on this group as well, since $iI_{n,n} \in Sp_{2n}$. The fixed points of θ must be block diagonal, that is

$$\theta(g) = g \iff g = \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix},$$

while membership in a symplectic group also forces $D = J_n(A^{-1})^t J_n$. Thus, A can be any invertible $n \times n$ matrix, and this completely determines g , so the fixed point subgroup $L := G^\theta$ is isomorphic to GL_n , giving a type CI symmetric pair.

Now, let us give a brief description of our involution notation. The symmetric group of permutations on $[n] := \{1, \dots, n\}$ is denoted by \mathcal{S}_n . If $\pi \in \mathcal{S}_n$, then its one-line notation is the string $\pi_1 \pi_2 \dots \pi_n$, where $\pi_i = \pi(i)$ for $1 \leq i \leq n$. For instance, consider $1(2, 6, 8)(3, 4, 5, 7)9$ which is the standard cycle decomposition of the permutation π from \mathcal{S}_9 whose one-line notation is given by $\pi = 164578329$.

An *involution* is an element of \mathcal{S}_n of order at most 2, and the set of involutions in \mathcal{S}_n is denoted by \mathcal{I}_n . Let $\pi \in \mathcal{I}_n$ be an involution. Since we often need the data of fixed points (1-cycles) of π , we always include them when writing π in cycle notation. Thus, our standard form for π will be

$$\pi = (a_1, b_1)(a_2, b_2) \dots (a_k, b_k) d_1 \dots d_{n-2k},$$

where $a_i < b_i$ for all $1 \leq i \leq k$, $a_1 < a_2 < \dots < a_k$, and $d_1 < \dots < d_{n-2k}$.

Definition 2.1. A *signed (p, q) -involution* is an involution $\pi \in \mathcal{I}_{p+q}$ with an assignment of $+$ and $-$ signs to the fixed points of π such that there are $p - q$ more $+$'s than $-$'s, where $q \leq p$.

For example, $\pi = (1\ 8)(2\ 4)(3^+)(5^-)(6^-)(7^+)$ is a signed $(4, 4)$ -involution. Observe here that p is equal to the number of fixed points in π with a $+$ sign attached plus the number of two-cycles in π , while q is equal to the number of fixed points in π with a $-$ sign attached plus the number of two-cycles in π . Next, we present (p, q) -clans.

Definition 2.2. Let p and q be two positive integers and set $n := p + q$. Suppose that $q \leq p$. A (p, q) -clan $\gamma = c_1 \dots c_n$ is a string of n symbols from $\mathbb{N} \cup \{+, -\}$ such that

1. there are $p - q$ more +’s than -’s;
2. if a natural number appears in γ , then it appears exactly twice.

For example, $12 + 21$ is a $(3, 2)$ -clan and $+1 + 1$ is a $(3, 1)$ -clan. We consider clans γ and γ' to be equivalent if the positions of every pair of matching numbers are the same in each. For example, $\gamma := 1122$ and $\gamma' := 2211$ are the same $(2, 2)$ -clan, since both γ and γ' have matching numbers in the positions $(1, 2)$ and $(3, 4)$.

If $\gamma = c_1 \cdots c_n$, then the *reverse* of γ , denoted by $rev(\gamma)$, is the clan

$$rev(\gamma) = c_n c_{n-1} \cdots c_1.$$

We take $-\gamma$ to be the clan obtained from γ by changing all +’s to -’s, and vice versa.

Definition 2.3. A (p, q) -clan γ is called *skew-symmetric* if

$$\gamma = -rev(\gamma).$$

It is clear that skew-symmetric (p, q) -clans are only possible when $p = q$. Let us illustrate this definition with the following examples.

Example 2.4. Consider the clan $\gamma = + - 123312 + -$, so that $rev(\gamma) = - + 213321 - +$. Since $\gamma = -rev(\gamma)$, it is a skew-symmetric $(5, 5)$ -clan.

The clan $\tau = 1234545321$ is a skew-symmetric $(5, 5)$ -clan as well. In fact, since it has no \pm symbols, $rev(\tau) = -rev(\tau)$.

There is a one-to-one correspondence between (p, q) -clans and signed (p, q) -involutions; detailed proof can be found in [4]. The signed involution associated to a clan γ is the one which has a transposition of the indices of each pair of matching natural numbers in γ , and each fixed point is given the sign found at the corresponding position in γ .

3 Counting Skew-Symmetric (n, n) -Clans

Here, we record the formula for the number of skew-symmetric (n, n) -clans which parametrize the Borel subgroup orbits in the classical symmetric space Sp_{2n}/GL_n ([19], Theorem 3.2.11). Let \mathcal{Z}_n denote the set of all skew-symmetric (n, n) -clans and Z_n denote its cardinality. Let $\zeta_{k,n}$ denote the number of such clans which contain k pairs of natural numbers. In order to count Z_n , first we will count $\zeta_{k,n}$.

To have such a count, we first determine the number of possible ways of placing \pm symbols among $2n - 2k$ spots. By skew-symmetry, it is enough to focus on the first half of the string of length $2n$. Assume that there are j many + signs in the first half. There are k many pairs of natural numbers and $2n - 2k$ of the symbols are \pm signs, half of which must occur among the first n symbols. Therefore, one can conclude that there are

$$\sum_{j=0}^{n-k} \binom{n-k}{j} = 2^{n-k} \tag{3.1}$$

possible ways to place + symbols among those remaining $n - k$ spots. We have just proved the following lemma:

Lemma 3.1. *There are 2^{n-k} ways of placing \pm symbols among $2n - 2k$ spots to obtain skew-symmetric (n, n) -clans with k pairs of matching natural numbers.*

Remark 3.2. Definition 2.3 implies that the number of + signs among the first n symbols of a given skew-symmetric (n, n) -clan must equal to the number of - signs among the last n .

Remark 3.3. The number of pairs of natural numbers in an (n, n) -clan can not exceed n because the inequality $0 \leq 2n - 2k$ gives $0 \leq k \leq n$.

Our next task is to determine the number of possible ways of placing k pairs to build a skew-symmetric (n, n) -clan $\gamma = c_1 \cdots c_n c_{n+1} \cdots c_{2n}$. Together with the lemma, this will yield our main result for this section.

Theorem 3.4. *For every nonnegative integer k with $k \leq n$, we have*

$$\zeta_{k,n} = 2^{n-k} \binom{n}{k} a_k, \quad \text{where} \quad a_k := \sum_{b=0}^{\lfloor k/2 \rfloor} \binom{k}{2b} \frac{(2b)!}{b!}, \quad (3.2)$$

so that

$$Z_n = \sum_{k=0}^n \zeta_{k,n} = \sum_{k=0}^n 2^{n-k} \binom{n}{k} a_k.$$

Proof. Let us first define the following interrelated sets:

$$\begin{aligned} \Pi_{1,1} &:= \{((i, j), (2n + 1 - j, 2n + 1 - i)) \mid 1 \leq i < j \leq n\}, \\ \Pi_{1,2} &:= \{((i, j), (2n + 1 - j, 2n + 1 - i)) \mid 1 \leq i \leq n < j \leq 2n\}, \\ \Pi_1 &:= \Pi_{1,1} \cup \Pi_{1,2}, \\ \Pi_2 &:= \{(i, j) \mid 1 \leq i \leq n < j \leq 2n, i + j = 2n + 1\}. \end{aligned}$$

Given a skew-symmetric (n, n) -clan γ , Π_1 is the set of possible placeholders for two distinct pairs of natural numbers that determine each other in γ . We will refer to these pairs of pairs as *families*, after [10]. The set Π_2 corresponds to the list of possible “stand-alone” pairs in γ .

If (c_i, c_j) is a pair of matching natural numbers in the skew-symmetric clan γ and if (i, j) is an element of Π_2 , then we call (c_i, c_j) a pair of type Π_2 . If $((c_i, c_j), (c_{2n+1-j}, c_{2n+1-i}))$ is a family in a skew-symmetric clan γ and if $((i, j), (2n + 1 - j, 2n + 1 - i)) \in \Pi_{1,s}$ ($s \in \{1, 2\}$), then we call it a family of type $\Pi_{1,s}$.

To illustrate these sets, consider the $(7, 7)$ -clan $1 + 134 - 55 + 342 - 2$. For this clan, $((c_1, c_3), (c_{12}, c_{14}))$ is a type $\Pi_{1,1}$ family, $((c_4, c_9), (c_5, c_{10}))$ is a type $\Pi_{1,2}$ family, and (c_7, c_8) is a Π_2 pair.

Clearly, if γ has b many families and a many Π_2 pairs, then $2b + a = k$ is the total number of pairs in our skew-symmetric clan γ . To see how many different ways there are in which these pairs of indices can be situated in γ , we start by choosing k spots from the first n

positions in $\gamma = c_1 \cdots c_{2n}$. Obviously, this can be done in $\binom{n}{k}$ many different ways, and each choice made in the first half determines the second half of the clan uniquely.

There are $\binom{k}{2b}$ possibilities for choosing which of these k spots will be occupied by symbols from families. Then, we choose b pairs among these $2b$ elements which can be done in $\frac{(2b)!}{b!2^b}$ ways. But each of these pairs can either be in a $\Pi_{1,1}$ family or a $\Pi_{1,2}$, family, so we multiply by these 2^b additional options, resulting in a factor of $\frac{(2b)!}{b!}$. Observe that choosing these pairs is equivalent to choosing (i, j) for the families in $\Pi_{1,1}$ and choosing $(i, 2n + 1 - j)$ for the families in $\Pi_{1,2}$. Once this is done, finally, the remaining $k - 2b$ spots will be filled by the first symbols of the a pairs of type Π_2 . This can be done in only one way.

Therefore, in summary, the number of different ways of placing k pairs to build a skew-symmetric (n, n) -clan γ is given by

$$\binom{n}{k} \sum_{b=0}^{\lfloor \frac{k}{2} \rfloor} \binom{k}{2b} \frac{(2b)!}{b!}, \quad \text{or equivalently,} \quad \binom{n}{k} \sum_{b=0}^{\lfloor \frac{k}{2} \rfloor} \binom{k}{2b} \binom{2b}{b} b!.$$

Combining this with Lemma 3.1 yields the formula for $\zeta_{k,n}$. □

Remark 3.5. By a straightforward calculation, it is easy to check that the recurrence

$$a_k = a_{k-1} + 2(k-1)a_{k-2}, \tag{3.3}$$

holds for all $k \geq 2$, with $a_0 = a_1 = 1$.

Example 3.6. All the possible skew-symmetric $(3, 3)$ clans are:

- - - + + +, + + + - - -, + + - + - -, + - + - + -, - + + - - +, - - + - + +,
 - + - + - +, + - - + + -, 1 + + - - 1, + 1 + - 1 -, + + 1 1 - -, 1 - - + + 1,
 - 1 - + 1 +, - - 1 1 + +, 1 + - + - 1, 1 - + - + 1, - 1 + - 1 +, + 1 - + 1 -,
 + - 1 1 + -, - + 1 1 - +, + 1 1 2 2 -, 1 1 - + 2 2, 1 - 1 2 + 2, - 1 1 2 2 +, 1 2 + - 1 2,
 - 1 2 1 2 +, 1 2 - + 1 2, 1 + 2 1 - 2, + 1 2 1 2 -, 1 + 1 2 - 2, 1 1 + - 2 2, 1 2 + - 2 1,
 1 + 2 2 - 1, + 1 2 2 1 -, 1 2 - + 2 1, 1 - 2 2 + 1, - 1 2 2 1 +, 1 - 2 1 + 2, 1 2 3 3 2 1,
 1 2 3 1 2 3, 1 2 1 3 2 3, 1 2 3 2 3 1, 1 1 2 2 3 3, 1 2 3 3 1 2, 1 2 2 3 3 1.

Thus, there are 45 skew-symmetric $(3, 3)$ clans.

The first few values of Z_n are 1, 3, 11, 45, 201, 963, 4899, 26253, 147345, 862083, 5238459, ... This is also the number of signed involutions on $2n$ letters that are equal to their reverse-complements and avoid the pattern $\bar{2}\bar{1}$ ([13], A083886). We will refer such objects as *restricted involutions* and explain the terminology used to define them in the next subsection.

3.1 Restricted Involutions

In this section, we will show that \mathcal{Z}_n is in one-to-one correspondence with the set of restricted involutions $\hat{\mathcal{I}}_n$, which we introduce below. Our main reference for this section is [20]. Recall

that the elements of the hyperoctahedral group \mathcal{C}_n can be considered as signed permutations and written as $\hat{\pi} = \hat{\pi}(1)\hat{\pi}(2)\dots\hat{\pi}(n)$, in which each of the symbols $1, 2, \dots, n$ appears, possibly barred to indicate a negative. For instance, there are 8 elements in \mathcal{C}_2 , which are explicitly, $12, \bar{1}\bar{2}, \bar{1}2, 1\bar{2}, 21, \bar{2}\bar{1}, \bar{2}1, 2\bar{1}$ in one-line notation.

We define the absolute value notation by $|\hat{\pi}(i)|$ if $\hat{\pi}(i)$ is $\hat{\pi}(i)$ positive and $\overline{\hat{\pi}(i)}$ otherwise. For example, if $\hat{\pi} = \bar{1}2$ then $|\hat{\pi}| = 12$.

Definition 3.7. Let $\hat{\rho} \in \mathcal{C}_k$ and $\hat{\pi} \in \mathcal{C}_n$. We say that $\hat{\pi}$ contains a signed pattern $\hat{\rho}$ or is a $\hat{\rho}$ -containing signed permutation, if there is a sequence of k indices, $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that two conditions hold:

1. $|\hat{\pi}(i_p)| > |\hat{\pi}(i_q)|$ if and only if $|\hat{\rho}(p)| > |\hat{\rho}(q)|$ for all $k \geq p > q \geq 1$;
2. $\hat{\pi}(i_j)$ is signed if and only if $\hat{\rho}(j)$ is signed for all $1 \leq j \leq k$.

A signed permutation $\hat{\pi}$ which does not contain such a pattern $\hat{\rho}$ is said to *avoid* $\hat{\rho}$.

For example, $\hat{\pi} = 21\bar{3}\bar{4} \in \mathcal{C}_4$ contains the signed patterns $\bar{1}\bar{2}$ but does not contain the pattern 12 .

Definition 3.8. A *signed involution on n letters* is an element of \mathcal{C}_n such that its cycle representation contains cycles of either two non-signed symbols or two signed symbols.

We define two simple operations on signed permutations. Given $\hat{\pi} = \hat{\pi}(1)\hat{\pi}(2)\dots\hat{\pi}(n)$ we consider the:

- Reverse permutation: $\text{Rev}(\hat{\pi}) = \hat{\pi}(n)\dots\hat{\pi}(2)\hat{\pi}(1)$
- Complement permutation: $\text{Comp}(\hat{\pi}) = \hat{\rho}(1)\hat{\rho}(2)\dots\hat{\rho}(n)$ where $\hat{\rho}(i) = n + 1 - \hat{\pi}(i)$ if $\hat{\pi}(i) > 0$ and $-(n + 1) - \hat{\pi}(i)$ otherwise.

These operations commute, allowing us to define our set of interest, called restricted involutions.

Definition 3.9. A signed involution $\hat{\pi}$ on $2n$ letters is called a *restricted involution* if it is equal to its reverse complement and avoids the pattern $\bar{2}\bar{1}$. The set of all such involutions for fixed n will be denoted by $\hat{\mathcal{I}}_n$ and its cardinality by \hat{I}_n .

For example, $12, \bar{1}\bar{2}$, and 21 are the restricted involutions of \mathcal{C}_2 , which are the members of $\hat{\mathcal{I}}_1$. We will prove that $\hat{I}_n = Z_n$ by exhibiting an explicit bijection between skew-symmetric (n, n) -clans and the restricted involutions of \mathcal{C}_{2n} . This will allow us to make use of a recurrence relation known for restricted involutions to give a generating function for the number of skew-symmetric (n, n) -clans and the orbits they parametrize.

First, we state the recurrence relation;

Proposition 3.10. *Taking $\hat{I}_0 = 1$ and $\hat{I}_1 = 3$, the number \hat{I}_n satisfies the following recurrence relation;*

$$\hat{\mathcal{I}}_n = 3\hat{\mathcal{I}}_{n-1} + 2(n-1)\hat{\mathcal{I}}_{n-2},$$

for all $n \geq 2$.

Let us make the notion of the *underlying involution* of a clan precise. This is effectively the associated signed (p, q) -involution mentioned at the end of Section 2, but without the signs on the fixed points.

Definition 3.11. For a skew-symmetric (n, n) -clan $\gamma = c_1 c_2 \dots c_{2n}$, the associated underlying involution $\pi \in \mathcal{S}_{2n}$ is defined as follows:

1. $\pi(i) = i$, if c_i is either a $+$ or a $-$ for any i ;
2. $\pi(i) = j$ and $\pi(j) = i$, if $c_i = c_j \in \mathbb{N}$ is a matching pair of natural numbers for any i, j .

The following describes an algorithm for obtaining a restricted involution $\hat{\pi}$ from a skew-symmetric clan γ with underlying involution π by applying signs to the symbols of π .

- (i) If $c_i = +$ (so $c_{2n+1-i} = -$) for $i \leq n$, keep $\hat{\pi}(i) = i$ and $\hat{\pi}(2n+1-i) = 2n+1-i$.
- (ii) If $c_i = -$ (so $c_{2n+1-i} = +$) for $i \leq n$, take $\hat{\pi}(i) = \bar{i}$ and $\hat{\pi}(2n+1-i) = \overline{2n+1-i}$.
- (iii) If $c_i = c_j \in \mathbb{N}$ is a matching pair of natural numbers, then keep $\hat{\pi}(i) = j$ and $\hat{\pi}(j) = i$ for all i, j .

For example, consider the skew-symmetric $(2, 2)$ clan $\gamma = - 1 1 +$. Since $c_1 = -, c_4 = +$ we will have $\hat{\pi}(1) = \bar{1}$ and $\hat{\pi}(4) = \bar{4}$, and having $c_2 = c_3 = 1$ gives us $\hat{\pi}(2) = 3$ and $\hat{\pi}(3) = 2$. It follows that the associated restricted involution is $\bar{1}32\bar{4}$.

Without trouble, this algorithm can be reversed to give a map from restricted involutions to clans. In fact, for each restricted involution, $\hat{\pi}$ will correspond to a unique clan γ . For a given $\hat{\pi} = \hat{\pi}(1)\hat{\pi}(2) \dots \hat{\pi}(2n)$, we define the reverse algorithm as follows.

- (i) If $\hat{\pi}(i) = i$ for any $1 \leq i \leq n$, then $\hat{\pi}(2n+1-i) = 2n+1-i$ because $\hat{\pi} = \text{Rev}(\text{Comp}(\hat{\pi}))$. It follows that the clan γ has $c_i = +$ and $c_{2n+1-i} = -$.
- (ii) If $\hat{\pi}(i) = \bar{i}$ for any $1 \leq i \leq n$, then $\hat{\pi}(2n+1-i) = \overline{2n+1-i}$ because $\hat{\pi} = \text{Rev}(\text{Comp}(\hat{\pi}))$. It follows that the clan γ has $c_i = -$ and $c_{2n+1-i} = +$.
- (iii) If $\hat{\pi}(i) = j$ for any $1 \leq i < j \leq 2n$, then $c_i = c_j = a$ where $a \in \mathbb{N}$.

This algorithm and its reverse are clearly injective, so the fact that each restricted involution in $\hat{\mathcal{I}}_n$ gives a skew-symmetric (n, n) -clan completes the bijection.

Theorem 3.12. *Restricted involutions on $2n$ letters and skew-symmetric (n, n) -clans are in bijection.*

Corollary 3.13. *Taking $Z_0 = 1$ and $Z_1 = 3$, the number of skew-symmetric (n, n) -clans satisfies the recurrence relation*

$$Z_n = 3Z_{n-1} + 2(n-1)Z_{n-2}, \tag{3.4}$$

and has exponential generating function

$$\sum_{n=0}^{\infty} Z_n \frac{x^n}{n!} = e^{3x+x^2}. \tag{3.5}$$

Proof. The recurrence relation appears for restricted involutions as Theorem 4.2 in [8]. The exponential generating function appears in the associated OEIS entry [A083886](#). \square

3.2 Partial Orders on Skew-Symmetric Clans

In this subsection, we depict two important partial orders on \mathcal{Z}_n , namely the *weak* and *full* (or *Bruhat*) closure orders. We refer the reader to [16] for explanation of the notation and a full combinatorial description of the weak order, which is lengthy. The full closure order can be obtained from the weak order by applying the recursive procedure described in [12], which is implicit in the work of Richardson and Springer [14].




Figure 3.1: Closure orders on \mathcal{Z}_2

In the posets, the black labelled edges are those which come from the weak order, while red edges are only present in the Bruhat order.




Figure 3.2: Closure orders on \mathcal{Z}_3

4 A combinatorial interpretation

In this section, we describe a combinatorial set of objects whose cardinality is given by Z_n . Recall that an (n, n) *Delannoy path* is an integer lattice path from $(0, 0)$ to (n, n) in the plane \mathbb{R}^2 consisting only of single north, east, diagonally northeast steps. Alternatively, one can consider strings from the alphabet $\{N, E, D\}$ such that the number of N 's plus the number of D 's is equal to the sum of the E 's and D 's (which is equal to n). We produce an explicit bijection between the set of skew-symmetric (n, n) -clans \mathcal{Z}_n and the set of Delannoy paths with certain labels which are defined as follows.

Definition 4.1. By a *labelled step* we mean a pair (L, l) , where $L \in \{N, E, D\}$ and l is a positive integer such that $l = 1$ if $L = N$ or $L = E$. A *weighted (n, n) Delannoy path* is a word of the form $W := W_1 \dots W_r$, where W_i 's ($i = 1, \dots, r$) are labeled steps $W_i = (L_i, l_i)$ such that

- $L_1 \dots L_r$ is a Delannoy path from $\mathcal{D}(n, n)$;
- if $L_i = D$ and $l_i \neq 1$, then $W_{r+1-i} = (D, 2n + 1 - l_i)$;
- letting $k_i = \#\{j < i \mid l_j \neq 1 \text{ and } j \neq r + 1 - i\}$, then $1 \leq l_i \leq 2n + 1 - 2(i + k_i)$ for $1 \leq i \leq \lfloor \frac{r}{2} \rfloor$.

The set of all weighted (n, n) Delannoy paths is denoted by $\mathcal{L}^w(n, n)$.

Theorem 4.2. *There is a bijection between the set of weighted (n, n) Delannoy paths and the set of skew-symmetric (n, n) -clans. In particular, we have*

$$Z_n = \sum_{W \in \mathcal{L}^w(n, n)} 1.$$

Proof. Let e_n denote the cardinality of $\mathcal{L}^w(n, n)$. We will prove that the sequence of e_n 's obeys the same recurrence as the Z_n 's, and it satisfies the same initial conditions. Let $\gamma = c_1 \dots c_{2n}$ be an arbitrary skew-symmetric (n, n) -clan with associated signed (n, n) -involution

$$\pi = (i_1, j_1) \dots (i_k, j_k) d_1^{s_1} \dots d_{2n-2k}^{s_{2n-2k}}, \quad \text{where } s_1, \dots, s_{2n-2k} \in \{+, -\}.$$

First, we look at the position of $2n$, which gives us four cases. If $2n$ appears as a fixed point with a $+$ sign, 1 must appear as a fixed point with a $-$ sign by the skew-symmetry condition. If this is the case, then we draw first an E -step between $(n-1, n)$ and (n, n) and then an N -step between $(0, 0)$ and $(0, 1)$. We label both of these steps by 1 to turn them into labeled steps. Next, we remove the fixed points 1 and $2n$ from π and shift the remaining symbols down by 1 to obtain a signed $(n-1, n-1)$ involution corresponding to a skew-symmetric clan. There are e_{n-1} possible ways of completing the drawn step steps between $(0, 1)$ and $(n-1, n)$ to an (n, n) Delannoy path.

In a similar manner, in case $2n$ appears as a fixed point with a $-$ sign, 1 must appear as a fixed point with a $+$ sign. Then we draw an N -step between $(n, n-1)$ and (n, n) and

an E -step between $(0, 0)$ and $(1, 0)$, each with weight 1. Then there are again e_{n-1} possible ways of completing these steps between $(1, 0)$ and $(n, n-1)$ to a weighted (n, n) Delannoy path.

Next, consider the case where $2n$ appears in a transposition $(i, 2n)$ coming from a family in the clan γ . Then there is a companion 2-cycle, which is necessarily of the form $(1, j)$ for $j = 2n + 1 - i$. Then we draw a D -step between $(n-1, n-1)$ and (n, n) and label this step by i , and draw another D -step between $(0, 0)$ and $(1, 1)$ and label this step by j . Then we remove the two cycle $(i, 2n)$ as well as its companion $(1, j)$ from π . To get rid of the gaps in the remaining numbers created by the removal of two 2-cycles, we renormalize the remaining entries by appropriately subtracting numbers so that in the resulting object, which we denote by $\pi^{(1)}$, every number from $\{1, \dots, 2n-4\}$ appears exactly once. It is easy to see that we then have a signed $(n-2, n-2)$ -involution which corresponds to a skew-symmetric $(n-2, n-2)$ clan. Now, it can occur that the label of either drawn diagonal step can be any of the $2(n-1)$ numbers from $\{2, \dots, 2n-1\}$, but the choice of one specifies the other as they must add to $2n+1$. Finally, let us note that there are e_{n-2} possible ways to complete these labeled diagonal steps between $(1, 1)$ and $(n-1, n-1)$ to a weighted (n, n) Delannoy path.

As the final case, consider when $2n$ appears in transposition with 1. Then, we draw a D -step between (n, n) and $(n-1, n-1)$ and we label this step by 1. Then we remove the 2-cycle $(1, 2n)$ from π , and to get rid of the gap created, we re-normalize the remaining entries by subtracting 1 from each so that in the resulting object $\pi^{(1)}$, every number from $\{1, \dots, 2n-2\}$ appears exactly once. Finally, let us note that there are e_{n-1} possible weighted paths from $(0, 0)$ to $(n-1, n-1)$ to complete this labeled diagonal step to a weighted (n, n) Delannoy path.

Combining our observations we see that a weighted (n, n) Delannoy path can be obtained (by appending the initial or final weighted steps described above) from an $(n-1, n-1)$ -clan in 3 ways, or from an $(n-2, n-2)$ clan in $2n-2$ different ways. Thus, weighted (n, n) Delannoy paths satisfy the recurrence

$$e_n = 3e_{n-1} + 2(n-1)e_{n-2}. \quad (4.1)$$

We can take $e_0 = 1$, and it is easy to check $e_1 = 3$, where the three paths are $(N, 1)(E, 1)$, $(E, 1)(N, 1)$ and $(D, 1)$. An explicit bijection is achieved by iterating the steps above, removing entries from π to produce weighted steps. This finishes our proof. \square

Let us illustrate our construction by an example.

Example 4.3. Let γ denote the skew-symmetric $(5, 5)$ clan

$$\gamma = + - 123312 + -,$$

and let π denote the corresponding signed involution

$$\pi = (3, 7)(4, 8)(5, 6)1^+2^-9^+10^-.$$

The steps of our constructions are shown in Figure 4.1.




Figure 4.1: Algorithmic construction of the bijection onto weighted Delannoy paths.

One could describe the weak and full closure orders on $B \backslash Sp_{2n} / GL_n$ in terms of weighted lattice paths with appropriate statistics. However, this rephrasing of the order on clans does not seem to be as illuminating as it is for the order on Schubert cells in the (Lagrangian) Grassmannian. Lattice paths associated to Schubert cells determine partition shapes, whence their closure order coincides with the order relations from Young's lattice (given by containment of partition shapes).

5 Sects

In this section, we will verify that the framework of [2], described in the introduction, applies to the CI case.

5.1 Parabolic Subgroups and Levi Factors

In order to present sects for skew-symmetric (n, n) -clans, we must visit the theory of parabolic subgroups of symplectic groups. We refer to [9] for background theory.

Given a vector space V with bilinear form ω , recall that an *isotropic subspace* W is one such that $\omega(\mathbf{u}, \mathbf{v}) = 0$ for all vectors $\mathbf{u}, \mathbf{v} \in W$. If we also use ω to stand for the matrix which represents this bilinear form in some chosen basis, this condition becomes $\mathbf{u}^t \omega \mathbf{v} = \mathbf{0}$. A




Figure 4.2: Weak order on $\mathcal{L}^\omega(2, 2)$

polarization of V is an orthogonal decomposition (with respect to, say, the usual dot product) of V into subspaces which are each isotropic (with respect to ω), that is $V = V_- \oplus V_+$.

Given a vector space V with a bilinear form ω , we define an *isotropic flag* as a sequence of vector spaces

$$\{\mathbf{0}\} \subset V_1 \subset \dots \subset V_r \subset V$$

such that V_i is an isotropic subspace of V for all $1 \leq i \leq r$.

Taking $V = \mathbb{C}^{2n}$, we have a bilinear form given by the same Ω which defines the symplectic group. From [9], Proposition 12.13, the parabolic subgroups of $G = Sp_{2n}$ are precisely the stabilizers of flags which are isotropic with respect to Ω .

Let E_n be the subspace of \mathbb{C}^{2n} generated by standard basis vectors \mathbf{e}_i , $1 \leq i \leq n$. It is easy to check that this is an isotropic subspace of \mathbb{C}^{2n} with respect to Ω , and in fact this subspace is maximally isotropic. Then there is a polarization of \mathbb{C}^{2n} as

$$V = E_n \oplus W,$$

where W is the subspace spanned by $\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{2n}\}$.

The stabilizer of E_n (which is the stabilizer of the the flag $\{\mathbf{0}\} \subset E_n \subset \mathbb{C}^{2n}$) is the parabolic subgroup P consisting of matrices with $n \times n$ block form

$$P = \left\{ p \in G \mid p = \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \right\}, \quad \text{which has Levi factor} \quad L = \left\{ \begin{pmatrix} A & 0 \\ 0 & J_n(A^{-1})^t J_n \end{pmatrix} \mid A \in GL_n \right\}. \quad (5.1)$$

See [9] p. 144 or [6] §8.1 for related discussion. Thus, we see that the Levi subgroup of this parabolic subgroup coincides with the symmetric subgroup presented in Section 2.

The upshot of this coincidence is that we have a G -equivariant projection map

$$\pi : G/L \longrightarrow G/P$$

which we can analyze. Let B be the Borel subgroup of upper triangular matrices in G ([9], page 39). The B -orbits in G/P are Schubert cells, while B -orbits in G/L are indexed by skew symmetric (n, n) -clans. The equivariance of π allows us to ask precisely which clans constitute the pre-image of a particular Schubert cell. We call such a collection of clans the *sect* associated to the Schubert cell.

In the literature, clans usually parametrize L orbits in the isotropic flag variety G/B by encoding the information of how flags in that orbit relate to the polarization. Recall that a *full* isotropic flag F_\bullet in \mathbb{C}^{2n} is a sequence of vector subspaces $\{V_i\}_{i=0}^n$ such that

$$\{\mathbf{0}\} = V_0 \subset V_1 \subset V_2 \subset \dots \subset V_n \tag{5.2}$$

and $\dim V_i = i$ for all i and V_n is a maximal isotropic subspace. We find it convenient to write

$$F_\bullet = \langle \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \rangle$$

to indicate that F_\bullet is the flag with $V_i = \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_i\}$ for all $1 \leq i \leq n$. Any full isotropic flag is canonically extended to a full flag in \mathbb{C}^{2n}

$$\{\mathbf{0}\} \subset V_1 \subset \dots \subset V_{2n-1} \subset V_{2n} = \mathbb{C}^{2n}$$

by assigning

$$V_{2n-i} = V_i^\perp := \{v \in \mathbb{C}^{2n} \mid \Omega(\mathbf{v}, \mathbf{w}) = 0 \ \forall \mathbf{w} \in V_i\},$$

so we may abuse notation slightly by using F_\bullet to refer to either presentation. For example, the standard isotropic full flag is extended as

$$E_\bullet = \langle \mathbf{e}_1, \dots, \mathbf{e}_n, \mathbf{e}_{n+1}, \dots, \mathbf{e}_{2n} \rangle.$$

If $g \in G$ is a matrix whose i^{th} column is a vector \mathbf{v}_i , then one can obtain a full isotropic flag from g by taking $F_\bullet = \langle \mathbf{v}_1, \dots, \mathbf{v}_{2n} \rangle$. For example, the identity matrix I_{2n} gives the standard isotropic flag E_\bullet . All elements of the same (right) B -orbit give the same flag, so one can identify the points of G/B with full isotropic flags with respect to Ω in \mathbb{C}^{2n} .

We must also present a few definitions before describing the process of obtaining orbit-representative flags; see also [2].

Definition 5.1. Given an (n, n) -clan $\gamma = (c_1 \cdots c_{2n})$, one obtains the *default signed clan* associated to γ by assigning to c_i a signature of $-$ and to c_j a signature of $+$ whenever $c_i = c_j \in \mathbb{N}$ and $i < j$. We denote this default signed clan as $\tilde{\gamma} = (\tilde{c}_1 \cdots \tilde{c}_{2n})$.

Remark 5.2. Note that the definition above differs from Definition 3.2.8 of “skew-symmetric signed clan” given in [19], which makes the opposite choice. This will also affect our statement of Yamamoto’s Theorem 3.2.11, given as Theorem 5.4 below.

For example, $\tilde{\gamma} = (+1_-, 2_-, 1_+, 2_+)$ is the default signed clan of $\gamma = (+1212-)$. Every symbol c_i has a signature, which is just the symbol itself in case c_i is $+$ or $-$.

Definition 5.3. Given a default signed clan $\tilde{\gamma}$, define a permutation $\sigma \in \mathcal{S}_{2n}$ which, for $i \leq n$:

- assigns $\sigma(i) = i$ and $\sigma(2n + 1 - i) = 2n + 1 - i$ if c_i is a symbol with signature $+$.
- assigns $\sigma(i) = 2n + 1 - i$ and $\sigma(2n + 1 - i) = i$ if c_i is a symbol with signature $-$.

We call σ the *default permutation* associated to γ .

Note that σ is an involution, and is the σ' which results from choosing $\sigma'' = id$ in the context of [19], Theorem 3.2.11. For instance, $+1212-$ has default permutation 154326 , in one-line notation.

5.2 Sects for Skew-Symmetric Clans

For this subsection, fix B as the Borel subgroup of $G = Sp_{2n}$ consisting of upper triangular matrices, and P and L as defined by the condition 5.1. The following is a specialization of [19] Theorem 3.2.11, which gives representative flags for L -orbits in the isotropic flag variety G/B .

Theorem 5.4. Given a skew-symmetric (n, n) -clan $\gamma = (c_1 \cdots c_{2n})$ with default permutation σ , define a flag $F_\bullet(\gamma) = \langle \mathbf{v}_1, \dots, \mathbf{v}_{2n} \rangle$ by making the following assignments.

- If $c_i = -$, set

$$v_i = \mathbf{e}_{\sigma(i)}$$

- If $c_i = +$, set

$$v_i = \begin{cases} \mathbf{e}_{\sigma(i)} & \text{if } i \leq n, \\ -\mathbf{e}_{\sigma(i)} & \text{if } i > n. \end{cases}$$

- If $c_i = c_j \in \mathbb{N}$ where \tilde{c}_i has signature $-$ and \tilde{c}_j has signature $+$ (that is, $i < j$), then set

$$v_i = \begin{cases} \frac{1}{\sqrt{2}}(\mathbf{e}_{\sigma(i)} + \mathbf{e}_{\sigma(j)}) & \text{if } i \leq n, \\ -\frac{1}{\sqrt{2}}(\mathbf{e}_{\sigma(i)} + \mathbf{e}_{\sigma(j)}) & \text{if } i > n. \end{cases}$$

and

$$v_j = \frac{1}{\sqrt{2}}(\mathbf{e}_{\sigma(i)} - \mathbf{e}_{\sigma(j)}).$$

Then $F_\bullet(\gamma)$ is a representative flag for the L -orbit Q_γ in G/B . Furthermore, if g_γ is the matrix defined by letting v_i be its i^{th} column, then $Q_\gamma = Lg_\gamma B/B$. Matrices/flags obtained in this way from CI clans constitute a full set of representative flags for L -orbits in G/B .

For example, the matrix representative for the clan $\gamma = (+1212-)$ is

$$g_\gamma = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} & 0 \\ 0 & \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} & 0 \\ 0 & \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

In the other direction, one can always recover an isotropic flag F_\bullet from a coset gB by taking F_i to be the span of the first i columns of any matrix in gB .

As in [2], we define the *base clan* of the clan γ as the one obtained by replacing all signed natural numbers in $\tilde{\gamma}$ by their signature. For example, $(+1212-)$ has base clan $(+--++-)$. Notice that the base clan of a skew-symmetric clan remains skew-symmetric, and consists only of $+$ and $-$ symbols. Base clans are in correspondence with closed orbits ([16]).

The following lemma is the major step in identifying the sects.

Lemma 5.5. *Let Q_γ and Q_τ be L -orbits in G/B corresponding to skew symmetric (n, n) -clans γ and τ . Then Q_γ and Q_τ lie in the same P orbit of G/B if and only if γ and τ have the same base clan.*

Proof. First, we prove necessity by showing that if γ has base clan τ , then the representing flags for each L -orbit Q_γ and Q_τ lie in the same P -orbit. Then all clans with base clan τ will lie in the same P -orbit. More precisely, we exhibit parabolic group elements that can iteratively transform the representing flag for τ into the representing flag for γ .

Let $\gamma = (c_1 \cdots c_{2n})$ and $\tau = (t_1 \cdots t_{2n})$, and let $F_\bullet(\gamma) = \langle \mathbf{v}_1, \dots, \mathbf{v}_{2n} \rangle$ and $F_\bullet(\tau) = \langle \mathbf{u}_1, \dots, \mathbf{u}_{2n} \rangle$ be the corresponding flags constructed by Theorem 5.4. As each clan has the same signature at symbols of the same index, they have the same default permutation. Then, we have two kinds of cases to examine.

Case 1: There is a pair of numbers $c_i = c_{2n+1-i}$ with $i \leq n$. Then we have

$$(\mathbf{v}_i, \mathbf{v}_{2n+1-i}) = \left(\frac{1}{\sqrt{2}}(\mathbf{e}_r + \mathbf{e}_{2n+1-r}), \frac{1}{\sqrt{2}}(\mathbf{e}_r - \mathbf{e}_{2n+1-r}) \right) \quad \text{and} \quad (\mathbf{u}_i, \mathbf{u}_{2n+1-i}) = (\mathbf{e}_r, -\mathbf{e}_{2n+1-r})$$

for some $n+1 \leq r \leq 2n$.

Let p^r denote the parabolic group element¹ defined by

$$\begin{aligned} p^r : \mathbf{e}_r &\longmapsto \mathbf{e}_r + \mathbf{e}_{2n+1-r}, \\ \mathbf{e}_i &\longmapsto \mathbf{e}_i \quad \text{for } i \neq r. \end{aligned}$$

Note that each pair of vectors

$$(\mathbf{e}_r + \mathbf{e}_{2n+1-r}, \mathbf{e}_r - \mathbf{e}_{2n+1-r}) \quad \text{and} \quad (\mathbf{e}_r + \mathbf{e}_{2n+1-r}, -\mathbf{e}_{2n+1-r})$$

¹It is easy to check that the matrix of this map satisfies the defining condition 5.1. That the matrix of this map is symplectic according to the condition in 2.1 is routine (if lengthy) linear algebra.

generates the same subspace, so it doesn't matter which one of the vectors $\frac{1}{\sqrt{2}}(\mathbf{e}_r - \mathbf{e}_{2n+1-r})$ or \mathbf{e}_r appears as \mathbf{v}_j in the flag $F_\bullet(\gamma)$. Then, the action of p^r on $F_\bullet(\tau)$ has the effect of taking \mathbf{u}_i to the span of \mathbf{v}_i , making the spans of $(p^r \cdot \mathbf{u}_i, p^r \cdot \mathbf{u}_{2n+1-i})$ and $(\mathbf{v}_i, \mathbf{v}_{2n+1-i})$ the same.

Case 2: In this case, we have $c_i = c_j$ with $i < j \neq 2n+1-i$ so that by skew-symmetry, there is another pair of natural numbers $c_{2n+1-j} = c_{2n+1-i}$. Without loss of generality, we can assume $i \leq n$. In this case, Theorem 5.4 will yield a flag $F_\bullet(\gamma)$ with

$$(\mathbf{v}_i, \mathbf{v}_j) = \left(\frac{1}{\sqrt{2}}(\mathbf{e}_r + \mathbf{e}_{2n+1-s}), \frac{1}{\sqrt{2}}(\mathbf{e}_r - \mathbf{e}_{2n+1-s}) \right)$$

and

$$(\mathbf{v}_{2n+1-j}, \mathbf{v}_{2n+1-i}) = \left(\pm \frac{1}{\sqrt{2}}(\mathbf{e}_s + \mathbf{e}_{2n+1-r}), \frac{1}{\sqrt{2}}(\mathbf{e}_s - \mathbf{e}_{2n+1-r}) \right),$$

where $n < r = \sigma(i)$ and $n < s = \sigma(2n+1-j)$. We also obtain the flag $F_\bullet(\tau)$ with

$$(\mathbf{u}_i, \mathbf{u}_j) = (\mathbf{e}_r, \pm \mathbf{e}_{2n+1-s})$$

and

$$(\mathbf{u}_{2n+1-j}, \mathbf{u}_{2n+1-i}) = (\mathbf{e}_s, -\mathbf{e}_{2n+1-r}).$$

Then define a linear map by

$$\begin{aligned} p^{r,s} : \mathbf{e}_r &\longmapsto \mathbf{e}_r + \mathbf{e}_{2n+1-s}, \\ \mathbf{e}_s &\longmapsto \mathbf{e}_s + \mathbf{e}_{2n+1-r}, \\ \mathbf{e}_i &\longmapsto \mathbf{e}_i \quad \text{for } i \neq r, s. \end{aligned}$$

It is again routine to check that this map defines an element of P , so $p^{r,s} \cdot F_\bullet(\tau)$ is a flag in the same P -orbit. Similar to the previous case, this map takes \mathbf{u}_i to the span of \mathbf{v}_i , \mathbf{u}_{2n+1-j} to the span of \mathbf{v}_{2n+1-j} , yielding pairs with the same span:

$$(p^{r,s} \cdot \mathbf{u}_i, p^{r,s} \cdot \mathbf{u}_j) \quad \text{and} \quad (\mathbf{v}_i, \mathbf{v}_j),$$

and

$$(p^{r,s} \cdot \mathbf{u}_{2n+1-j}, p^{r,s} \cdot \mathbf{u}_{2n+1-i}) \quad \text{and} \quad (\mathbf{v}_{2n+1-j}, \mathbf{v}_{2n+1-i}).$$

Now, after we act on the flag $F_\bullet(\tau)$ by the appropriate element of the form p^r for each pair of natural numbers $c_i = c_{2n+1-i}$ in γ , and the appropriate element $p^{r,s}$ for each family

$$\{c_i = c_j, c_{2n+1-j} = c_{2n+1-i} \mid j \neq 2n+1-i\},$$

then we obtain a flag which is an equivalent presentation of $F_\bullet(\gamma)$. Thus, Q_γ is in the same P -orbit as Q_τ .

To show the converse, it suffices to show that L -orbits corresponding to distinct base clans τ and λ lie in distinct P orbits. Let $\tau = (t_1 \cdots t_{2n}) \neq \lambda = (l_1 \cdots l_{2n})$ and let $F_\bullet(\tau) = \langle \mathbf{u}_1, \dots, \mathbf{u}_{2n} \rangle$ and $F_\bullet(\lambda) = \langle \mathbf{w}_1, \dots, \mathbf{w}_{2n} \rangle$ be flags constructed to represent each orbit using Theorem 5.4. Now let i be the least index such that $t_i \neq l_i$. Without loss of generality,

we may assume that $t_i = -$ and $l_i = +$, and that $i \leq n$. Then we have $\mathbf{u}_i = \mathbf{e}_r$ for some $n+1 \leq r \leq n$ and $\mathbf{w}_i = \mathbf{e}_q$ for some $1 \leq q \leq n$. For these flags to be in the same P -orbit we would need to be able to carry \mathbf{e}_q to a vector with non-zero \mathbf{e}_r component via some $p \in P$. This would force a non-zero entry in the (r, q) -entry of the matrix of p , but since p has a block-diagonal form with zero (i, j) -entry whenever $i > n$ and $j \leq n$, this is impossible. Thus, the L -orbits Q_τ and Q_λ are in distinct P -orbits. □

Note that there is an isomorphism

$$\begin{aligned} \varphi : LgB &\longrightarrow Bg^{-1}L \\ lgb &\longmapsto (lgb)^{-1} \end{aligned} \tag{5.3}$$

for any such double coset. Thus, given a clan γ , we have a map

$$Q_\gamma = Lg_\gamma B/B \longrightarrow Lg_\gamma B \xrightarrow{\varphi} Bg_\gamma^{-1}L \longrightarrow Bg_\gamma^{-1}L/L =: R_\gamma$$

that takes the the L -orbit Q_γ to a unique B -orbit in G/L , which we can safely denote as R_γ since it depends only on the clan γ via g_γ . If we compose this map with the projection $\pi : G/L \rightarrow G/P$, then Lemma 5.5 tells us that the B -orbits of G/L which project to the Schubert cell $Bg_\gamma^{-1}P/P$ are exactly the set of R_τ where τ has the same base clan as γ . This yields the following.

Proposition 5.6. *Let R_γ and R_τ be B orbits in G/L corresponding to clans γ and τ , and let $\pi : G/L \rightarrow G/P$ denote the canonical projection. Then $\pi(R_\gamma) = \pi(R_\tau)$ if and only if γ and τ have the same base clan.*

Having successfully grouped clans according base clan, we find it appropriate to also call the collection of clans with base clan γ the *sect* of γ , denoted Σ_γ . By abuse of terminology, we will also use this term to describe collections of any of the objects (B -orbits, L -orbits, double cosets, etc.) parameterized by Σ_γ .

5.3 Sects and Schubert Cells

As in [3] §3.3, we fix T as the maximal torus of diagonal matrices in G , and then the Weyl group $W = N_G(T)/T$ can be identified with the permutations in the subgroup of the symmetric group \mathcal{S}_{2n} defined by the condition,

$$W \cong \{\sigma \in \mathcal{S}_{2n} \mid \sigma(2n+1-i) = 2n+1-\sigma(i) \text{ for all } i\}.$$

This happens to be isomorphic to the hyperoctahedral group \mathcal{C}_n . A Weyl group element given by a permutation $\sigma \in \mathcal{S}_{2n}$ can also be represented by a matrix $w(\sigma) \in N_G(T)$ where the i^{th} column is $\pm \mathbf{e}_{\sigma(i)}$. Thus, the matrix will be like that of the permutation matrix for σ , only with some -1 entries instead of 1 entries to ensure that it is a symplectic group element.

The choice of B as the Borel subgroup of upper-triangular matrices determines the set of positive roots as

$$\Phi^+ = \{(\varepsilon_i \pm \varepsilon_j), | 1 \leq i < j \leq n\} \cup \{2\varepsilon_i | 1 \leq i \leq n\},$$

where $\pm\varepsilon_i$ denotes the torus character that takes $\text{diag}(t_1, \dots, t_{2n}) \mapsto t_i^{\pm 1}$. We choose simple roots

$$\Delta = \{\alpha_i = \varepsilon_i - \varepsilon_{i+1} | 1 \leq i \leq n-1\} \cup \{\alpha_n = \varepsilon_{2n}\},$$

so that our maximal parabolic subgroup P corresponds to the the subset $\Delta \setminus \{\alpha_n\}$. The Weyl group of P is denoted W_P and is isomorphic to a copy of \mathcal{S}_n with respect to which the minimal coset representatives of W/W_P can be identified as the permutations $\sigma \in W$ which also satisfy

$$\sigma(1) < \sigma(2) < \dots < \sigma(n).$$

The set of minimal coset representatives is denoted W^P . The values $\sigma(1), \dots, \sigma(n)$ become the indices of the standard basis vectors which appear (up to sign) in the first n columns of the matrix $w(\sigma)$.²

Let $[m] := \{1, \dots, m\}$. The coset $w(\sigma)P$ includes all of the \mathcal{S}_n -many permutations of the set

$$\{\mathbf{e}_{\sigma(1)}, \dots, \mathbf{e}_{\sigma(n)}\}$$

by the (right) action of the Levi subgroup on the columns of $w(\sigma)$. So order does not actually matter, and the Schubert cell can be understood as the Borel orbit of the isotropic subspace spanned by $\{\mathbf{e}_{\sigma(1)}, \dots, \mathbf{e}_{\sigma(n)}\}$. Hence, we can make the identification

$$\text{Schubert cells } C_I \longleftrightarrow \left\{ \begin{array}{l} \text{Subsets } I \text{ of } [2n] \text{ such that } |I| = n \\ \text{and if } i \in I \text{ then } 2n+1-i \notin I \end{array} \right\}.$$

Since for each opposite pair of indices we choose just one for membership in the subset I , there are 2^n cells total. Each cell is also associated to a maximal isotropic subspace of the form

$$V_I := \text{span}(\{\mathbf{e}_i | i \in I\}).$$

We can bijectively associate base clans to sets I by defining $\gamma_I = (c_1 \dots c_{2n})$ by

$$c_i = \begin{cases} + & \text{if } i \in I \\ - & \text{if } i \notin I. \end{cases} \quad (5.4)$$

Note that under this assignment, after building the default flag, $F_\bullet(\gamma_I) = \langle v_1, \dots, v_{2n} \rangle$, it easy to check that $\text{span}(\{v_1, \dots, v_n\}) = V_I$. In other words, $C_I = Bg_{\gamma_I}^{-1}P/P$. Finally, we have our promised result.

²It can be arranged so that these columns are exactly standard basis vectors, as in Theorem 5.4.

Theorem 5.7. *Let C_I be a Schubert cell of G/P , and $\pi : G/L \rightarrow G/P$ the natural projection. Associate to I a base clan γ_I as in equation 5.4, and denote the sect of γ_I by Σ_I . If R_γ denotes the B -orbit of G/L associated to the clan γ , then*

$$\pi^{-1}(C_I) = \bigsqcup_{\gamma \in \Sigma_I} R_\gamma. \quad (5.5)$$

Proof. The pre-image of C_I decomposes as a disjoint union of B -orbits R_γ as a consequence of the fact that π is G -equivariant, so in particular π is B -equivariant. It remains to determine for which γ we have $\pi(R_\gamma) = C_I$. Proposition 5.6 tells us that B -orbits R_γ and R_τ project to the same Schubert cell if and only if they are members of the same sect. Thus, we only have to prove that the B -orbit R_{γ_I} projects to the Schubert cell C_I .

We know that $\pi(R_{\gamma_I}) = Bg_{\gamma_I}^{-1}P/P$. Observe that from the construction of g_{γ_I} , if we denote the i^{th} column of $g_{\gamma_I}^{-1}$ as \mathbf{w}_i , then

$$\mathbf{w}_i = \begin{cases} -\mathbf{e}_{\sigma(i)} & \text{if } i \leq n \text{ and } c_i = -, \\ \mathbf{e}_{\sigma(i)} & \text{otherwise.} \end{cases}$$

Because this differs from the columns of g_{γ_I} only possibly by a sign, the first n columns span the same subspace, and in fact

$$g_{\gamma_I}^{-1}P = g_{\gamma_I}P.$$

This implies that

$$\pi(R_{\gamma_I}) = Bg_{\gamma_I}^{-1}P/P = Bg_{\gamma_I}P/P = C_I.$$

□

6 The Big Sect

Here, we are going to analyze the number of clans lying in the largest sect over $\Lambda(n)$, which is denoted by ϵ_n . The set of all such clans will be denoted by \mathcal{E}_n .

A clan lies in the largest sect if and only if it has a base clan in the following form

$$\underbrace{- \cdots -}_{\text{first } n\text{-spots}} - - + + \cdots + + + .$$

This follows from the fact that this base clan is γ_I where $V_I = \text{span}\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{2n}\}$ whose B -orbit C_I is the dense Schubert cell in $\Lambda(n)$, just as it is in $\text{Gr}(n, \mathbb{C}^{2n})$. Hence, a clan lies in the largest sect if and only if its only natural numbers are from $\Pi_{1,2}$ families and Π_2 pairs.

Let α denote the number of $\Pi_{1,2}$ families and β the number of Π_2 pairs in an arbitrary clan $\gamma \in \mathcal{E}_n$. Then $2\alpha + \beta = k$ is the total number of pairs in $\gamma = c_1 \cdots c_n c_{n+1} \cdots c_{2n}$. To see in how many different ways these pairs of indices can be situated in such γ , we start by choosing β spots from the first n positions in γ . Obviously, this can be done in $\binom{n}{\beta}$ many different ways. Then, we count different ways of choosing α pairs within the $n - \beta$ spots remaining (among the first n) to place the families of type $\Pi_{1,2}$. We have $\binom{n-\beta}{2\alpha}$ spots to

choose from, and $\frac{(2\alpha)!}{2^{\alpha}\alpha!}$ different ways to form pairs from these spots. This proves the following corollary.

Corollary 6.1. *The number of clans in the largest sect is given by*

$$\epsilon_n = \sum_{\substack{k=0 \\ 2\alpha+\beta=k}}^n \binom{n}{\beta} \binom{n-\beta}{2\alpha} \frac{(2\alpha)!}{2^{\alpha}\alpha!}, \quad \text{or equivalently,} \quad \epsilon_n = \sum_{\substack{k=0 \\ 2\alpha+\beta=k}}^n \binom{n}{\beta} \frac{(n-\beta)!}{(n-\beta-2\alpha)!2^{\alpha}\alpha!}.$$

The first values of ϵ_n , beginning with $n = 1$, are 2, 5, 14, 43, 142, 499, 1850, \dots . This is in fact the number of self-inverse partial permutations, also known as partial involutions, (A005425), as we show in the next section.

6.1 Partial Involutions

In this section, we will show that the set of all clans in the largest sect \mathcal{E}_n is in one-to-one correspondence with the set of all partial involutions on n elements. We refer to [1] and Chapter 15 of [5] for background theory. Recall that a *partial permutation* is a map

$$\pi : \{1, \dots, m\} \mapsto \{0, \dots, n\}$$

satisfying the following rule:

- if $\pi(i) = \pi(j)$ and $\pi(i) \neq 0$, then $i = j$ for each $1 \leq i, j \leq m$.

A partial permutations matrix π can be represented by an $m \times n$ matrix (x_{ij}) , where x_{ij} is 1 if and only if $\pi(i) = j$, and 0 otherwise. Note that under this convention we view our matrices as acting on vectors from the right. Partial permutations are sometimes called *rook placements*, and in case $m = n$, they form a monoid under matrix multiplication called the *rook monoid* and denoted \mathcal{R}_n .

Definition 6.2. A partial involution on n elements is a partial permutation which is represented by a symmetric $n \times n$ partial permutation matrix. The set of partial involutions on n elements will be denoted by \mathcal{P}_n and its cardinality by P_n .

We will prove that $\epsilon_n = P_n$ by exhibiting an explicit bijection between the clans in the largest sect and the partial involutions. This will allow us to make use of a recursion relationship known for the partial involutions to give a generating function for the sequence of ϵ_n 's. A direct proof of the recurrence using the formula of Corollary 6.1 may be more difficult to achieve. First, we state a well-known recurrence relation for partial involutions.

Proposition 6.3. *Taking $P_0 = 1$, the numbers P_n satisfy the recurrence relation*

$$P_n = 2P_{n-1} + (n-1)P_{n-2}$$

for all $n \geq 1$.

In order to describe the bijection between \mathcal{E}_n and \mathcal{P}_n , let π be a partial involution with matrix (x_{ij}) for $1 \leq i, j \leq n$. We construct a clan $\gamma = c_1 c_2 \dots c_{2n}$ as follows:

- (i) If $x_{ii} = 0$, then $c_i = -$ and $c_{2n+1-i} = +$.
- (ii) If $x_{ii} = 1$ for any $1 \leq i \leq n$, then we have a transposition $(i, 2n+1-i)$ in the underlying involution of γ which yields $c_i = c_{2n+1-i} = a$ for some $a \in \mathbb{N}$.
- (iii) If $x_{ij} = 1$ and $x_{ji} = 1$ (by symmetry), then $(i, 2n+1-j)$ and $(j, 2n+1-i)$ are the corresponding transpositions in the underlying involution for a clan. Thus, $c_i = c_{2n+1-j} = a$ and $c_j = c_{2n+1-i} = b$ for some $a, b \in \mathbb{N}$.

For example, consider the matrix $(x_{ij}) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$. Then since $x_{11} = 1$, we can take

$c_1 = c_6 = 1$ in the corresponding clan. Moreover, since $x_{23} = 1$ and $x_{32} = 1$, we assign $c_2 = c_4 = 2$ and $c_3 = c_5 = 3$. Thus, the corresponding clan is 123231.

This algorithm can be reversed easily. Let us start with a clan $\gamma = c_1 c_2 \dots c_{2n}$ and define its associated partial involution matrix as the one with zeros everywhere except:

- (ii) If $c_i = c_{2n+1-i} = a \in \mathbb{N}$, then $x_{ii} = 1$.
- (iii) If $c_i = c_{2n+1-j} = a \in \mathbb{N}$ for $i \neq j$, then $x_{ij} = x_{ji} = 1$.

This algorithm and its reverse are clearly injective between partial involutions and clans from the largest sect, completing the bijection.

Theorem 6.4. *Partial involutions on n letters and skew-symmetric (n, n) -clans in the largest sect are in bijection.*

Corollary 6.5. *Taking $\epsilon_0 = 1$, the number of clans in the largest sect satisfies the recurrence relation*

$$\epsilon_n = 2\epsilon_{n-1} + (n-1)\epsilon_{n-2},$$

and has exponential generating function

$$\sum_{n=0}^{\infty} \epsilon_n \frac{x^n}{n!} = e^{2x + \frac{x^2}{2}}.$$

Proof. The recurrence follows from Theorem 6.4 and Proposition 6.3. The exponential generating function can be found by a straightforward calculation. \square

Next we describe an order structure on partial involutions. Let B_m^- denote the group of invertible $m \times m$ lower triangular matrices, and B_n the group of invertible upper triangular $n \times n$ matrices. $B_m^- \times B_n$ acts on the set of all $m \times n$ matrices with complex entries by $(a, b) \cdot x = axb^{-1}$. The double cosets of this action are indexed by the partial permutations.

The group B_n also acts on the set of symmetric $n \times n$ matrices $\mathbb{S}(n)$ by the *congruence action*, which is defined by

$$b \cdot s = (b^{-1})^t x b^{-1},$$

for $x \in \mathbb{S}(n)$. The set of partial involutions \mathcal{P}_n parametrize congruence B_n -orbits. In fact, for $\pi \in \mathcal{P}_n$, the congruence B_n -orbit $B_n \cdot \pi$ is exactly the intersection $B_n^- \pi B_n \cap \mathbb{S}(n)$ (see [15], Theorem 3.1). Moreover, the orbit closure order for congruence B_n -orbits is the same as the closure order on $B_n^- \times B_n$ double cosets restricted to the set of partial involutions ([1]). This order can be described in terms of rank-control matrices associated to each partial permutation, which we define presently.

Definition 6.6. Let $\pi = (x_{ij})$ be an $m \times n$ matrix. For each $1 \leq k \leq m$ and $1 \leq l \leq n$, denote by $\pi_{k,l}$ the upper-left $k \times l$ submatrix of π . We denote by $R(\pi)$ the $m \times n$ matrix, whose (k, l) -entry is $\text{rank}(\pi_{k,l})$, and call it the *rank-control matrix* of π .

Example 6.7. If $\pi = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$, then $R(\pi) = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{pmatrix}$.

We then define a partial order on partial permutations by comparing the individual entries of their rank-control matrices, which is to say

$$\pi \preceq \sigma \iff \text{rank}(\pi_{k,l}) \leq \text{rank}(\sigma_{k,l}) \quad \text{for all } k, l.$$

Let C_π and C_σ denote the $B_m^- \times B_n$ -orbits corresponding to partial permutations π and σ . It follows from Theorem 15.31 of [5] that

$$\pi \preceq \sigma \iff \overline{C_\pi} \subseteq \overline{C_\sigma}.$$

Thus, rank-control matrices capture all of the information of orbit closure relationships. Corollary 5.4 of [1] specializes this to say that the restriction of \preceq to \mathcal{P}_n describes closure order on congruence B_n -orbits as well.

6.2 A Conjectural Bruhat Order on CI Clans

In [17], Wyser gives a description of the closure order of $GL_p \times GL_q$ -orbits in the flag variety of GL_{p+q} in terms of statistics on the (p, q) -clans parametrizing these orbits. We recall the statement and notation from that paper.

For any (p, q) -clan $\gamma = c_1 \cdots c_n$ and any i or i, j with $1 \leq i < j \leq n$, we let

- (1) $\gamma(i; +)$ = the total number of $+$ signs and pairs of equal natural numbers occurring among $c_1 \cdots c_i$;
- (2) $\gamma(i; -)$ = the total number of $-$ signs and pairs of equal natural numbers occurring among $c_1 \cdots c_i$;
- (3) $\gamma(i; j)$ = the number of pairs of equal natural numbers $c_s = c_t \in \mathbb{N}$ with $s \leq i < j < t$.

Theorem 6.8 (Theorem 1.2, [17]). *Let γ, τ be (p, q) -clans, and let Y_γ, Y_τ be the corresponding $GL_p \times GL_q$ -orbit closures in the flag variety of GL_{p+q} . Then $\gamma \leq \tau$ (meaning $Y_\gamma \subseteq Y_\tau$) if and only if the following three inequalities hold for all i, j :*

- (1) $\gamma(i; +) \geq \tau(i; +)$;
- (2) $\gamma(i; -) \geq \tau(i; -)$;
- (3) $\gamma(i; j) \leq \tau(i; j)$.

Just as skew-symmetric (n, n) -clans can be viewed as a subset of all (n, n) -clans, the isotropic flag variety $X := Sp_{2n}/B$ can be viewed as a subvariety of the flag variety G'/B' for $G' = GL_{2n}$ and its subgroup B' of upper triangular matrices. Indeed, the L -orbit in Sp_{2n}/B corresponding to the skew-symmetric clan γ is exactly the intersection $Q'_\gamma \cap X$, where Q'_γ is the $L' := GL_n \times GL_n$ -orbit in G'/B' corresponding to γ (see [16]).

It is then natural to wonder if the (full) closure order on skew-symmetric (n, n) -clans is simply the restriction of the closure order on all (n, n) -clans. This is stated as Conjecture 3.6 in [18], where it is reported that the conjecture has been verified computationally up to $n = 7$. The analogous conjecture is, in fact, made for two other classical symmetric pairs: $(SO_{2n+1}, S(O_{2p} \times O_{2q+1}))$ and $(Sp_n, Sp_p \times Sp_q)$. However, neither of the symmetric subgroups in these cases are Levi factors of a parabolic subgroup, so they hold less interest for our purposes.

It is worth noting that the veracity of the conjecture for $(Sp_n, Sp_p \times Sp_q)$ was claimed in [12] on the basis of the recursive process for deducing the full closure order relations from the weak order. However, very little detail is provided, and it is also claimed in that paper that the same holds for (SO_{2n}, GL_n) , though Wyser points out that this fails with a specific example when $n = 4$ ([18], p. 165).

A detailed proof of this conjecture for (Sp_{2n}, GL_n) can likely be achieved via methods similar to [17], but we forego this at present. Instead, we will just assign a partial order on skew-symmetric (n, n) -clans by

$$\gamma \leq_C \tau \iff Y_\gamma \subseteq Y_\tau \quad (\text{so } \gamma \leq \tau),$$

with Y_γ and Y_τ as before, and call \leq_C the *conjectural CI Bruhat order*. With this in hand we can state and prove a final result, analogous to [2], Theorem 1.7.

Theorem 6.9. *As posets $(\mathcal{E}_n, \leq_C) \cong (\mathcal{P}_n, \preceq)$, that is the poset of the largest sect within skew-symmetric (n, n) -clans with the conjectural CI Bruhat order is isomorphic to the poset of partial involutions ordered by congruence B_n -orbit closures.*

Proof. Given a clan $\gamma \in \mathcal{E}_n$, let π^γ denote the partial involution matrix obtained from γ by the algorithm described before Theorem 6.4. We will show that the conditions of Theorem 6.8 on clans γ and τ translate to all of the necessary conditions on the rank-control matrices of π^γ and π^τ , so that the orders are the same. This is achieved simply by identifying the statistics $\gamma(i; +)$, $\gamma(i; -)$ and $\gamma(i; j)$ as ranks of particular submatrices of the associated

partial involution. As in Definition 6.6, let $\pi_{k,l}^\gamma$ denote the northwest $k \times l$ submatrix of π^γ , and $R(\pi^\gamma)$ the rank-control matrix of π^γ so that the (k, l) -entry of $R(\pi^\gamma)$ is $\text{rank } \pi_{k,l}^\gamma$.

First we examine $\gamma(i; +)$. Because $\gamma = c_1 \cdots c_{2n}$ has base clan $- - \cdots - - + + \cdots + +$, the value of $\gamma(i; +)$ is zero for all $i \leq n$, and then because each symbol among the last n in γ is either a $+$ or the second of a natural number pair, we find that $\gamma(i; +) = \tau(i; +)$ for all i from 1 to $2n$. Thus, this statistic contains no ordering information within the largest sect.

Next, consider $\gamma(i; -)$. As i increases from 1 to n , so does $\gamma(i; -)$ unless there is a natural number at c_i . A natural number at c_i indicates a matrix entry of 1 in the i^{th} row of π^γ for $1 \leq i \leq n$. As i increases past n , $\gamma(i; -)$ increases the rest of the way up to n , and a natural number at c_i indicates a matrix entry of 1 in column c_{2n+1-i} . Then,

$$\gamma(i; -) = \begin{cases} i - \text{rank}(\pi_{i,n}^\gamma) & \text{for } i \leq n, \\ n - \text{rank}(\pi_{n,2n+1-i}^\gamma) & \text{for } i > n. \end{cases}$$

Because the ranks of submatrices appear negatively above, $\gamma(i; -) \geq \tau(i; -)$ if and only if $\text{rank}(\pi_{i,n}^\gamma) \leq \text{rank}(\pi_{i,n}^\tau)$ and $\text{rank}(\pi_{n,i}^\gamma) \leq \text{rank}(\pi_{n,i}^\tau)$ for all $1 \leq i \leq n$. This covers the rank conditions along the south and east borders of the matrices.

Next, consider $\gamma(i, j)$. By similar reasoning, it is not hard to see that

$$\gamma(i; j) = \begin{cases} \text{rank}(\pi_{i,n}^\gamma) & \text{for } i < j \leq n \\ \text{rank}(\pi_{i,2n-j}^\gamma) & \text{for } i \leq n < j \\ \text{rank}(\pi_{n,2n-j}^\gamma) & \text{for } n < i < j. \end{cases}$$

The case where $i \leq n < j$ shows that $\gamma(i; j) \leq \tau(i; j)$ for all $1 \leq i, j < n$ if and only if $\text{rank}(\pi_{i,m}^\gamma) \leq \text{rank}(\pi_{i,m}^\tau)$ for all $1 \leq i, m < n$. This covers the rank conditions everywhere else in the partial involution matrices, so we see that $\gamma \leq_C \tau$ if and only if $\pi^\gamma \preceq \pi^\tau$, completing the poset isomorphism. □

Remark 6.10. (\mathcal{E}_n, \leq_C) is also a maximal upper order ideal of the poset of all skew-symmetric (n, n) -clans under \leq_C . Its minimal element is the base clan and the maximal element is $12 \cdots (n-1)nn(n-1) \cdots 21$, which corresponds to the dense B -orbit of G/L . This follows from arguments identical to the proof of Proposition 5.4 in [2].

Remark 6.11. The proof of Theorem 6.9 can be adapted to show that the big sect of the type $AIII$ symmetric space $(GL_n/GL_p \times GL_q)$ is isomorphic to the closure order on $B_p^- \times B_q$ double cosets of the $p \times q$ complex matrices, indexed by partial permutations. This was proved only for the case $p = q$ in [2].

Acknowledgements. We thank Mahir Bilen Can for many constructive suggestions.

References

- [1] Eli Bagno and Yonah Cherniavsky, *Congruence B -orbits and the Bruhat poset of involutions of the symmetric group*, Discrete Mathematics, 312.6 (2012), pp. 1289–1299.
- [2] Bingham, Aram and Can, Mahir Bilen, *Sects*, arXiv preprint arXiv:1810.13159, 2018.
- [3] Billey, Sara and Lakshmibai, Venkatramani and Sarason, Sara, *Singular loci of Schubert varieties*, Springer Science & Business Media, 2000.
- [4] Can, Mahir Bilen and Uğurlu, Özlem, *The genesis of involutions (polarizations and lattice paths)*, Discrete Mathematics, 342 (2019), pp. 201–216.
- [5] Miller, Ezra and Sturmfels, Bernd, *Combinatorial commutative algebra*, Graduate Texts in Mathematics, Springer-Verlag, New York, 227 (2005).
- [6] Garrett, Paul B, *Buildings and classical groups*, CRC Press, 1997.
- [7] Goodman, Roe and Wallach, Nolan R. *Symmetry, representations, and invariants*, Springer, volume =255, 2009.
- [8] Hardt, Andy and Troyka, Justin M., *Restricted symmetric signed permutations*, Pure Mathematics and Applications. P.U.M.A., 23.3 (2012), pp. 179–217.
- [9] Malle, Gunter and Testerman, Donna, *Linear algebraic groups and finite groups of Lie type*, Cambridge University Press, 133 (2011).
- [10] Matsuki, Toshihiko and Oshima, Toshio *Embeddings of discrete series into principal series*, The orbit method in representation theory, Springer, pp. 147–175, 1990.
- [11] Matsuki, Toshihiko, *The orbits of affine symmetric spaces under the action of minimal parabolic subgroups*, Journal of the Mathematical Society of Japan, 31.2 (1979), pp. 331–357.
- [12] William M. McGovern and Peter E. Trapa, *Pattern avoidance and smoothness of closures for orbits of a symmetric subgroup in the flag variety*, Journal of Algebra, 322.8 (2009), pp. 2713–2730.
- [13] Sloane, Neil JA, *Online Encyclopedia of Integer Sequences (OEIS)*, 2018.
- [14] Richardson, RW and Springer, Tonny Albert, *The Bruhat order on symmetric varieties*, Geometriae Dedicata, Springer, 35.1-3 (1990), pp. 389–436.
- [15] Szechtman, Fernando, *Equivalence and congruence of matrices under the action of standard parabolic subgroups*, Electronic Journal of Linear Algebra, 16 (2007), pp. 325–333.
- [16] Benjamin J. Wyser, *Symmetric subgroup orbit closures on flag varieties*, University of Georgia, 2012.

- [17] Benjamin J. Wyser, *The Bruhat order on clans*. J. Algebraic Combinatorics, 44.3 (2016), pp. 495–517.
- [18] Benjamin J. Wyser, *K -orbit closures on G/B as universal degeneracy loci for flagged vector bundles splitting as direct sums*, Geometriae Dedicata, 181 (2016), pp. 137–175.
- [19] Yamamoto, Atsuko, *Orbits in the flag variety and images of the moment map for classical groups I*, Representation Theory of the American Mathematical Society, 1.13 (1997), pp. 329–404.
- [20] W.M.B. Dukes and Toufik Mansour, *Signed Involutions Avoiding 2-Letter Signed Patterns*, Annals of Combinatorics, 11 (2007), pp. 387–403.