

PROOFS OF CONJECTURES ABOUT PATTERN-AVOIDING LINEAR EXTENSIONS

COLIN DEFANT

*Princeton University
Fine Hall, 304 Washington Rd.
Princeton, NJ 08544*

ABSTRACT. After fixing a canonical ordering (or labeling) of the elements of a finite poset, one can associate each linear extension of the poset with a permutation. Some recent papers consider specific families of posets and ask how many linear extensions give rise to permutations that avoid certain patterns. We build off of two of these papers. We first consider pattern avoidance in k -ary heaps, where we obtain a general result that proves a conjecture of Levin, Pudwell, Riehl, and Sandberg in a special case. We then prove some conjectures that Anderson, Egge, Riehl, Ryan, Steinke, and Vaughan made about pattern-avoiding linear extensions of rectangular posets.

1. INTRODUCTION

Let S_n be the set of permutations of $[n] = \{1, \dots, n\}$, which we write as words in one-line notation. Given $\tau = \tau_1 \cdots \tau_m \in S_m$, we say a permutation $\sigma = \sigma_1 \cdots \sigma_n$ *contains the pattern* τ if there exist indices $i_1 < \cdots < i_m$ in $[n]$ such that for all $j, k \in [m]$, we have $\sigma_{i_j} < \sigma_{i_k}$ if and only if $\tau_j < \tau_k$. We say σ *avoids* τ if it does not contain τ . Let $\text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)$ be the set of permutations avoiding the patterns $\tau^{(1)}, \tau^{(2)}, \dots$ (this list of patterns could be finite or infinite).

The study of pattern avoidance in permutations, which began with Knuth's analysis of permutations that are sortable via a stack [19], has grown into a large, thriving area of research in combinatorics [6, 18, 21]. Recently, there has been a great amount of interest in pattern avoidance in other combinatorial objects such as words, inversion sequences, set partitions, and trees [1, 2, 4, 5, 7–17, 20, 22–24, 26, 28]. The papers [3, 5, 14, 20, 28] consider the following general type of problem. Let \mathcal{P} be an n -element poset, and suppose we are given some canonical total ordering of the elements of \mathcal{P} . We can view each linear extension of \mathcal{P} as a bijective labeling of the elements of \mathcal{P} with the elements of $[n]$. If we read these labels in the canonical order, we obtain a permutation. We can then ask how many linear extensions give rise to permutations that avoid certain patterns. An alternative, yet essentially equivalent, formulation of this problem is to consider a canonical *labeling* of \mathcal{P} and then view each linear extension of \mathcal{P} as an ordering. Again, each linear extension gives rise to a permutation, so we can ask the same enumerative questions about pattern avoidance.

E-mail address: cdefant@princeton.edu.

As a first example, we consider the problems introduced in [20]. A *complete k -ary tree* is a rooted tree in which each vertex that is not in the penultimate or the last level has exactly k children and all vertices in the last level are as far left as possible. We make the convention that trees grow up from their roots (as in real life). A *k -ary heap* is a complete k -ary tree whose vertices are bijectively labeled with the elements of $[n]$ so that each vertex is given a label that is smaller than the labels of its children. Associating complete k -ary trees with posets in the natural way, one can view a k -ary heap as a linear extension of the underlying complete k -ary tree. We endow each complete k -ary tree with the breadth-first ordering (traversing the levels from bottom to top with each level traversed from left to right). Reading the labels in a k -ary heap in this order yields a permutation in S_n , which we call the permutation *associated to the k -ary heap*. For example, the left image in Figure 1 shows the unique complete binary tree with 12 vertices. The right image shows a binary heap obtained by labeling this tree. The permutation associated to this binary heap is 132469712511810. In Section 2, we prove a conjecture from [20] concerning binary heaps whose associated permutations avoid the pattern 321. In fact, we will prove a much more general result.




FIGURE 1. The complete binary tree with 12 vertices (left) and a 12-vertex binary heap (right). Reading the labels of this heap in the breadth-first order yields the associated permutation 132469712511810.

The second type of problem we consider deals with the rectangular posets $EN_{s,t}$ studied in [3]. The poset $EN_{s,t}$ has st elements that are labeled in a canonical fashion. The easiest way to define these posets is via examples. The Hasse diagrams of $EN_{3,2}$, $EN_{3,5}$, and $EN_{4,3}$, along with their canonical labelings, are shown in Figure 1. Each linear extension of $EN_{s,t}$ can be viewed as an ordering of the labels, which is a permutation in S_{st} . For example, the linear extensions of $EN_{3,2}$ correspond to the permutations 531642, 536142, 536412, 563142, and 563412. In Section 3, we prove several of the conjectures that the authors of [3] posed about pattern-avoiding linear extensions of these posets.




FIGURE 2. The Hasse diagrams of three rectangular posets. The numbers show the canonical labelings.

2. PATTERN-AVOIDING k -ARY HEAPS

Let $\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)$ denote the set of n -vertex k -ary heaps whose associated permutations avoid the patterns $\tau^{(1)}, \tau^{(2)}, \dots$. The authors of [20] found formulas for $|\mathcal{H}_n^k(\tau^{(1)}, \dots, \tau^{(r)})|$ for every set $\{\tau^{(1)}, \dots, \tau^{(r)}\} \subseteq S_3$ except the singleton set $\{321\}$. In fact, they were unable to explicitly enumerate binary heaps avoiding 321. They did, however, compute $|\mathcal{H}_n^2(321)|$ for $1 \leq n \leq 31$ and prove that

$$2^{n-1} < |\mathcal{H}_n^2(321)| < 4^n.$$

Their data led them to conjecture that¹

$$(1) \quad \lim_{n \rightarrow \infty} |\mathcal{H}_n^2(321)|^{1/n} \text{ exists and is in the interval } (3.66, 4].$$

We will see that this conjecture follows as a special consequence of the main theorem of this section. To state this theorem, we need one additional piece of terminology.

Given $\lambda = \lambda_1 \cdots \lambda_\ell \in S_\ell$ and $\mu = \mu_1 \cdots \mu_m \in S_m$, the *sum* of λ and μ , denoted $\lambda \oplus \mu$, is the permutation in $S_{\ell+m}$ obtained by “placing μ above and to the right of λ .” More formally, the i^{th} entry of $\lambda \oplus \mu$ is

$$(\lambda \oplus \mu)_i = \begin{cases} \lambda_i & \text{if } 1 \leq i \leq \ell; \\ \mu_{i-\ell} + \ell & \text{if } \ell + 1 \leq i \leq \ell + m. \end{cases}$$

A permutation is called *sum indecomposable* if it cannot be written as the sum of two shorter permutations.

Theorem 2.1. *Fix an integer $k \geq 2$ and a nonempty sequence of permutation patterns $\tau^{(1)}, \tau^{(2)}, \dots$. If the permutations $\tau^{(1)}, \tau^{(2)}, \dots$ are all sum indecomposable, then*

$$\lim_{n \rightarrow \infty} |\text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n} = \lim_{n \rightarrow \infty} |\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n}.$$

Remark 2.1. How do we know that the limits in Theorem 2.1 actually exist? Because the permutations $\tau^{(1)}, \tau^{(2)}, \dots$ are sum indecomposable, we have an injective map $S_\ell \times S_m \hookrightarrow S_{\ell+m}$ given by $(\lambda, \mu) \mapsto \lambda \oplus \mu$ for all $\ell, m \geq 1$. This shows that the sequence $(|\text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)|)_{n \geq 1}$ is supermultiplicative, so the limit on the left-hand side exists by Fekete’s lemma. We also have an injective map $\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots) \hookrightarrow \text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)$ sending each k -ary heap to its associated permutation. This shows that $\limsup_{n \rightarrow \infty} |\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n} \leq \lim_{n \rightarrow \infty} |\text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n}$. We will see in the proof of Theorem 2.1 that $\lim_{n \rightarrow \infty} |\text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n} \leq \liminf_{n \rightarrow \infty} |\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n}$, so the limit on the right-hand side exists as well. It is interesting to note that this right-hand limit does not depend on k . ◇

Let $C_n = \frac{1}{n+1} \binom{2n}{n}$ denote the n^{th} Catalan number. It is well known that $|\text{Av}_n(321)| = C_n$. This implies that $\lim_{n \rightarrow \infty} |\text{Av}_n(321)|^{1/n} = 4$. Consequently, we can specialize Theorem 2.1 to obtain the following result. In the special case in which $k = 2$, this settles the conjecture from [20] that is stated in (1).

Corollary 2.1. *If $k \geq 2$, then $\lim_{n \rightarrow \infty} |\mathcal{H}_n^k(321)|^{1/n} = 4$.*

¹The actual conjecture was written as “ $|\mathcal{H}_n^2(321)| \sim c^n$ for some $c \in (3.66, 4)$,” but it is clear from context that the authors really meant what is written in (1).

Proof of Theorem 2.1. Fix an integer $k \geq 2$ and a sequence of sum indecomposable permutation patterns $\tau^{(1)}, \tau^{(2)}, \dots$. As mentioned in Remark 2.1, we must prove that

$$\lim_{n \rightarrow \infty} |\text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n} \leq \liminf_{n \rightarrow \infty} |\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n}.$$

To ease notation, let

$$a(n) = |\text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n}, \quad b(n) = |\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)|^{1/n}, \quad L_a = \lim_{n \rightarrow \infty} a(n), \quad L_b = \liminf_{n \rightarrow \infty} b(n).$$

The proof is trivial if one of the patterns $\tau^{(i)}$ is the permutation $1 \in S_1$, so we may assume otherwise. Because the patterns $\tau^{(1)}, \tau^{(2)}, \dots$ are sum indecomposable, the identity permutation $123 \cdots n$ is in $\text{Av}_n(\tau^{(1)}, \tau^{(2)}, \dots)$ and is the permutation associated to an element of $\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)$. Thus, $L_a, L_b \geq 1$. Fix $\varepsilon \in (0, 1)$, and let $N \geq 1$ be such that $a(n - \lfloor (n-1)/k \rfloor) > L_a - \varepsilon$ and $b(\lfloor (n-1)/k \rfloor) > L_b - \varepsilon$ for every $n \geq N$. Now choose $n \geq N$ such that $b(n) < L_b + \varepsilon$, and put $m = \lfloor (n-1)/k \rfloor$.

Claim: If $\lambda \in S_m$ is a permutation associated to a k -ary heap in $\mathcal{H}_m^k(\tau^{(1)}, \tau^{(2)}, \dots)$ and $\mu \in \text{Av}_{n-m}(\tau^{(1)}, \tau^{(2)}, \dots)$, then $\lambda \oplus \mu$ is a permutation associated to a k -ary heap in $\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)$.

Let us see how this claim implies Theorem 2.1. Suppose by way of contradiction that $L_b < L_a$. Assuming the claim, we have

$$\begin{aligned} (L_b + \varepsilon)^n > b(n)^n &= |\mathcal{H}_n^k(\tau^{(1)}, \tau^{(2)}, \dots)| \geq |\mathcal{H}_m^k(\tau^{(1)}, \tau^{(2)}, \dots)| \cdot |\text{Av}_{n-m}(\tau^{(1)}, \tau^{(2)}, \dots)| \\ &= b(m)^m a(n-m)^{n-m} > (L_b - \varepsilon)^m (L_a - \varepsilon)^{n-m}. \end{aligned}$$

Since $m < n/2$ and $L_b < L_a$, we have $(L_b - \varepsilon)^m (L_a - \varepsilon)^{n-m} > (L_b - \varepsilon)^{n/2} (L_a - \varepsilon)^{n/2}$. Hence, $(L_b + \varepsilon)^2 > (L_b - \varepsilon)(L_a - \varepsilon)$. Letting ε tend to 0 shows that $L_b \geq L_a$, contradicting our assumption that $L_b < L_a$. This completes the proof of Theorem 2.1 assuming the claim.

Now, let λ and μ be as in the claim. Let T be the complete k -ary tree with n vertices, and let v_i be the vertex of T that appears i^{th} in the breadth-first ordering. It is straightforward to check that the vertices v_{m+1}, \dots, v_n are incomparable (meaning none is a descendant of another). It follows that $\lambda \oplus \mu$ is the permutation associated to some k -ary heap in \mathcal{H}_n^k . Because λ and μ avoid the sum indecomposable permutations $\tau^{(1)}, \tau^{(2)}, \dots$, their sum $\lambda \oplus \mu$ must also avoid these permutations. This completes the proof of the claim. \square

3. RECTANGULAR POSETS

An *inversion* of a permutation $\pi = \pi_1 \cdots \pi_n \in S_n$ is a pair (i, j) such that $i < j$ and $\pi_i > \pi_j$. Let $\text{inv}(\pi)$ denote the number of inversions of π . Following [3], we let $\text{EN}_{s,t}(\tau)(q) = \sum_{\pi \in \text{EN}_{s,t}(\tau)} q^{\text{inv}(\pi)}$, where $\text{EN}_{s,t}(\tau)$ is the set of linear extensions of $\text{EN}_{s,t}$ (viewed as permutations of the labels of $\text{EN}_{s,t}$) that avoid the pattern τ . The following three theorems were stated as conjectures in [3]².

Theorem 3.1. *For all $t \geq 1$, we have*

$$\text{EN}_{3,t}(1243)(q) = \frac{q^{3(t^2-t+1)}(1 - q^{2t-1} - 2q^{2t} + q^{3t-1} + q^{3t})}{(1-q)(1-q^2)}.$$

Theorem 3.2. *For all $s \geq 1$, we have*

$$\text{EN}_{s,2}(2143)(q) = q^{(2s-1)(s-1)}(1+q)^{s-1}.$$

²Technically speaking, Theorem 3.1 was stated incorrectly in that article.

Theorem 3.3. *For all $s \geq 1$, we have*

$$\text{EN}_{s,3}(2143)(q) = q^{9\binom{s}{2}} F_s(1/q),$$

where $F_s(r)$ is defined by $F_0(r) = F_1(r) = 1$ and $F_s(r) = (1 + r + 2r^2)F_{s-1}(r) + r^3F_{s-2}(r)$ for $s \geq 2$.

Proof of Theorem 3.1. It is easy to verify this theorem when $t \in \{1, 2\}$, so assume $t \geq 3$. Let $\pi = \pi_1 \cdots \pi_n$ be a linear extension of $\text{EN}_{3,t}(1243)$ (viewed as a permutation of the labels). Note that $2t$ appears before 2 in π because, otherwise, the entries $1, 2, 2t, t$ would form a 1243 pattern. Similarly, $3t$ must appear before $t + 2$, lest the entries $t + 1, t + 2, 3t, 2t$ form a 1243 pattern. It follows that if we remove the entries 1 and $t + 1$ from π , then we will be left with the permutation

$$(2t + 1)(2t + 2) \cdots (3t)(t + 2)(t + 3) \cdots (2t)23 \cdots t.$$

Let i, j be such that $\pi_{i+1} = t + 1$ and $\pi_{j+1} = 1$. We can easily check that the possibilities for i and j are given by $i \in \{1, \dots, t\}$ and $j \in \{i + 1, \dots, 2t\}$. We find that

$$\text{EN}_{3,t}(1243)(q) = \sum_{i=1}^t \sum_{j=i+1}^{2t} q^{3t^2 - 3t + i + j}.$$

This can easily be rewritten as $\frac{q^{3(t^2-t+1)}(1 - q^{2t-1} - 2q^{2t} + q^{3t-1} + q^{3t})}{(1-q)(1-q^2)}$. \square

Proof of Theorem 3.2. Let S be the collection of subsets of $\{1, 3, 5, \dots, 2s - 3\}$, and define $\eta: \text{EN}_{s,2}(2143) \rightarrow S$ by

$$\eta(\pi) = \{i \in \{1, 3, 5, \dots, 2s - 3\} : i + 3 \text{ appears before } i \text{ in } \pi\}.$$

The map η is a bijection. It is now easy to check that

$$\text{EN}_{s,2}(2143)(q) = \sum_{X \subseteq \{1, 3, 5, \dots, 2s - 3\}} q^{(2s-1)(s-1) + |X|} = q^{(2s-1)(s-1)} (1 + q)^{s-1}. \quad \square$$

Proof of Theorem 3.3. Let H_ℓ be the set of $\pi = \pi_1 \cdots \pi_{3\ell} \in \text{EN}_{\ell,3}(2143)$ such that $\pi_2 = 3\ell - 1$. Let $H_\ell(q) = \sum_{\pi \in H_\ell} q^{\text{inv}(\pi)}$. Fix $s \geq 2$, and let $J(r_1, \dots, r_k)$ be the set of permutations $\pi = \pi_1 \cdots \pi_{3s} \in \text{EN}_{s,3}(2143)$ such that $\pi_i = r_i$ for all $i \in \{1, \dots, k\}$. One can check that the sets

$$J(3s - 2, 3s - 1, 3s), \quad J(3s - 2, 3s - 1, 3s - 5, 3s), \quad J(3s - 2, 3s - 5, 3s - 1, 3s), \\ J(3s - 2, 3s - 5, 3s - 1, 3s - 4, 3s), \quad \text{and} \quad J(3s - 2, 3s - 1, 3s - 5, 3s - 4, 3s)$$

partition $\text{EN}_{s,3}(2143)$. Call these sets J_1, J_2, J_3, J_4, J_5 , respectively. We have

$$\sum_{\pi \in J_1} q^{\text{inv}(\pi)} = q^{9(s-1)} \text{EN}_{s-1,3}(2143)(q), \quad \sum_{\pi \in J_2} q^{\text{inv}(\pi)} = q^{9(s-1)-1} \text{EN}_{s-1,3}(2143)(q), \\ \sum_{\pi \in J_3} q^{\text{inv}(\pi)} = q^{9(s-1)-2} \text{EN}_{s-1,3}(2143)(q), \quad \sum_{\pi \in J_4} q^{\text{inv}(\pi)} = q^{9(s-1)-3} H_{s-1}(q),$$

and

$$\sum_{\pi \in J_5} q^{\text{inv}(\pi)} = q^{9(s-1)-2} H_{s-1}(q).$$

This yields

$$(2) \quad \text{EN}_{s,3}(2143)(q) = q^{9(s-1)-2} (1 + q + q^2) \text{EN}_{s-1,3}(2143)(q) + q^{9(s-1)-3} (1 + q) H_{s-1}(q).$$

The sets J_1, J_2, J_5 partition H_s , so

$$(3) \quad H_s(q) = q^{9(s-1)-1} (1 + q) \text{EN}_{s-1,3}(2143)(q) + q^{9(s-1)-2} H_{s-1}(q).$$

Solving the recurrence relations (2) and (3) subject to the initial conditions $\text{EN}_{1,3}(2143)(q) = H_1(q) = 1$, we obtain the desired identity $\text{EN}_{s,3}(2143)(q) = q^{9\binom{s}{2}}F_s(1/q)$. \square

The article [3] also posed several conjectures concerning the polynomials $F_s(q)$. We settle many of these conjectures³ in the following theorem. Since our focus is on the combinatorics of pattern-avoiding linear extensions and not these specific polynomials, we omit some details from the proof. In what follows, let $[q^r]G(q)$ denote the coefficient of q^r in the Laurent series $G(q)$.

Theorem 3.4. *Define the polynomials $F_s(r)$ by $F_0(r) = F_1(r) = 1$ and $F_s(r) = (1+r+2r^2)F_{s-1}(r) + r^3F_{s-2}(r)$ for $s \geq 2$. For $s \geq 2$,*

- *the values of $[q^3]F_s(q)$ are given by OEIS sequence A134465;*
- *the values of $[q^{2s-2}]F_s(q)$ are given by OEIS sequence A098156;*
- *the values of $[q^{s-1}]F_s(q)$ are given by OEIS sequence A116914;*
- *the values of $[q^s]F_s(q)$ are given by OEIS sequence A072547;*
- *the values of $[q^{s+1}]F_s(q)$ are given by OEIS sequence A002054;*
- *the values of $[q^{s+2}]F_s(q)$ are given by OEIS sequence A127531.*

Proof. Let $A(x, q) = \sum_{s \geq 0} F_s(q)x^s$. The recurrence for $F_s(q)$ translates into the identity

$$(4) \quad A(x, q) = \frac{1 - (q + 2q^2)x}{1 - (1 + q + 2q^2)x - q^3x^2}.$$

Computing $\frac{1}{6} \frac{\partial^3}{\partial q^3} A(x, q)$ proves the first bullet point. The remainder of the proof makes use of the method of diagonals, which is discussed in Section 6.3 of [27].

If we view $qA(x/q^2, q)$ as a function of the complex variable q , then

$$\sum_{s \geq 0} ([q^{2s-2}]F_s(q))x^s = [q^{-1}](qA(x/q^2, q)) = \frac{1}{2\pi i} \int_{|q|=\rho} qA(x/q^2, q) dq,$$

where $\rho > 0$ is sufficiently small and the integral is taken over the circle of radius ρ centered at the origin. By the Residue theorem, this is

$$\sum_{j=1}^r \text{Res}_{q=u_j(x)}(qA(x/q^2, q)),$$

where $u_1(x), \dots, u_r(x)$ are the singularities of $qA(x/q^2, q)$ (viewed as functions of x) that tend to 0 as $x \rightarrow 0$. We can explicitly compute that $r = 2$ and that

$$u_1(x) = \frac{x(1+x) + (1-x)\sqrt{x(4+x)}}{2(1-2x)} \quad \text{and} \quad u_2(x) = \frac{x(1+x) - (1-x)\sqrt{x(4+x)}}{2(1-2x)}.$$

Let $U(x, q) = q^2(x - q(1 - 2x))$ and $V(x, q) = x + qx(1 + x) - q^2(1 - 2x)$ so that $qA(x/q^2, q) = U(x, q)/V(x, q)$. Let $V_q(x, q) = \frac{\partial}{\partial q} V(x, q)$. Since $u_1(x)$ and $u_2(x)$ are simple poles of $qA(x/q^2, q)$, we find that

$$\text{Res}_{q=u_j(x)}(qA(x/q^2, q)) = \frac{U(x, u_j(x))}{V_q(x, u_j(x))} \quad \text{for } j \in \{1, 2\}.$$

³We also correct some typos made in the original statements of these conjectures.

We have

$$\sum_{j=1}^2 \operatorname{Res}_{q=u_j(x)}(qA(x/q^2, q)) = \sum_{j=1}^2 \frac{U(x, u_j(x))}{V_q(x, u_j(x))} = \frac{x(1 - 2x + x^2 + x^3)}{(1 - 2x)^2},$$

and this proves the second bullet point.

To prove the third, fourth, fifth, and sixth bullet points, we choose an integer $\ell \leq 2$ and view $q^{-\ell-1}A(x/q, q)$ as a complex function of the variable q . As above, we have

$$\sum_{s \geq 0} ([q^{s+\ell}]F_s(q))x^s = [q^{-1}](q^{-\ell-1}A(x/q, q)) = \sum_{j=1}^t \operatorname{Res}_{q=v_j(x)}(q^{-\ell-1}A(x/q, q)),$$

where $v_1(x), \dots, v_t(x)$ are the singularities of $q^{-\ell-1}A(x/q, q)$ that tend to 0 as $x \rightarrow 0$. Let $Y(x, q) = -1 + x + 2qx$ and $Z(x, q) = x - (1 - x)q + x(2 + x)q^2$ so that $q^{-\ell-1}A(x/q, q) = q^{-\ell}Y(x, q)/Z(x, q)$. If $\ell \leq 0$, then the only singularity of $q^{-\ell-1}A(x/q, q)$ that tends to 0 as $x \rightarrow 0$ is $v_1(x) = \frac{1 - x - \sqrt{1 - 2x - 7x^2 - 4x^3}}{2x(2 + x)}$. If $\ell \in \{1, 2\}$, then there is one other singularity, which is $v_2(x) = 0$. One can check that

$$\operatorname{Res}_{q=0}(q^{-2}A(x/q, q)) = 1 - x^{-1} \quad \text{and} \quad \operatorname{Res}_{q=0}(q^{-3}A(x/q, q)) = 1 + 2x^{-1} - x^{-2}.$$

Note that in each of these expressions, the coefficient of x^s is 0 for every $s \geq 2$. It follows that for all integers $\ell \leq 2$ and $s \geq 2$, the coefficient of x^s in $\sum_{s \geq 0} ([q^{s+\ell}]F_s(q))x^s$ agrees with the coefficient of x^s in $\operatorname{Res}_{q=v_1(x)}(q^{-\ell-1}A(x/q, q))$. We have

$$\operatorname{Res}_{q=v_1(x)}(q^{-\ell-1}A(x/q, q)) = \frac{v_1(x)^{-\ell}Y(x, v_1(x))}{Z_q(x, v_1(x))},$$

where $Z_q(x, q) = \frac{\partial}{\partial q}Z(x, q)$. Computing these expressions explicitly when $\ell \in \{-1, 0, 1, 2\}$ and simplifying, we obtain proofs of the last four bullet points. \square

4. ACKNOWLEDGMENTS

The author was supported by a Fannie and John Hertz Foundation Fellowship and an NSF Graduate Research Fellowship.

REFERENCES

- [1] M. H. Albert, R. E. L. Aldred, M. D. Atkinson, C. Handley, and D. Holton, Permutations of a multiset avoiding permutations of length 3. *European J. Combin.*, **22** (2001), 1021–1031.
- [2] M. Anderson, M. Diepenbroek, L. Pudwell, and A. Stoll, Pattern avoidance in reverse double lists, *Discrete Math. Theor. Comput. Sci.*, **19** (2018), #13.
- [3] D. Anderson, E. S. Egge, M. Riehl, L. Ryan, R. Steinke, Y. Vaughan, Pattern avoiding linear extensions of rectangular posets. *J. Comb.*, **9** (2018), 185–220.
- [4] A. Baxter and L. Pudwell, Ascent sequences avoiding pairs of patterns. *Electron. J. Combin.*, **22** (2015), P1.58.
- [5] D. Bevan, D. Levin, P. Nugent, J. Pantone, L. Pudwell, M. Riehl, and M. L. Tlachac, Pattern avoidance in forests of binary shrubs. *Discrete Math. Theor. Comput. Sci.*, **18** (2016), #8.
- [6] M. Bóna, *Combinatorics of permutations*. CRC Press, 2012.
- [7] P. Brändén and T. Mansour, Finite automata and pattern avoidance in words. *J. Combin. Theory Ser. A*, **110** (2005), 127–145.
- [8] A. Burstein and T. Mansour, Words restricted by patterns with at most 2 distinct letters. *Electron. J. Combin.*, **9.2** (2002) #R3.2.

- [9] S. Corteel, M. A. Martinez, C. D. Savage, and M. Weselcouch, Patterns in inversion sequences I. *Discrete Math. Theor. Comput. Sci.*, **18** (2016).
- [10] M. Dairiko, L. Pudwell, S. Tyner, and C. Wynn, Non-contiguous pattern avoidance in binary trees. *Electron. J. Combin.*, **19** (2012), P22.
- [11] D. Daly and L. Pudwell, Pattern avoidance in the rook monoid. *J. Comb.*, **5** (2014), 471–497.
- [12] C. Defant, Enumeration of stack-sorting preimages via a decomposition lemma. arXiv:1904.02829.
- [13] C. Defant, Fertility, strong fertility, and postorder Wilf equivalence. arXiv:1904.03115.
- [14] C. Defant, Poset pattern-avoidance problems posed by Yakubov. *J. Comb.*, **9** (2018), 233–257.
- [15] C. Defant, Stack-sorting preimages of permutation classes. arXiv:1809.03123.
- [16] C. Defant and N. Kravitz, Stack-sorting for words. arXiv:1809.09158.
- [17] A. Godbole, A. Goyt, J. Herdan, and L. Pudwell, Pattern avoidance in ordered set partitions. *Ann. Comb.*, **18** (2014), 429–445.
- [18] S. Kitaev, Patterns in Permutations and Words. Monographs in Theoretical Computer Science. Springer, Heidelberg, 2011.
- [19] D. E. Knuth, The Art of Computer Programming, volume 1, Fundamental Algorithms. Addison-Wesley, Reading, Massachusetts, 1973.
- [20] D. Levin, L. Pudwell, M. Riehl, and A. Sandberg, Pattern avoidance in k -ary heaps. *Australas. J. Combin.*, **64** (2016), 120–139.
- [21] S. Linton, N. Ruškuc, V. Vatter, Permutation Patterns, London Mathematical Society Lecture Note Series, Vol. 376. Cambridge University Press, 2010.
- [22] T. Mansour and M. Shattuck, Pattern avoidance in inversion sequences. *Pure Math. Appl.*, **25** (2015), 112–129.
- [23] T. Mansour and M. Shattuck, Pattern-avoiding set partitions and Catalan numbers. *Electron. J. Combin.*, **18** (2011–2012), #P34.
- [24] M. A. Martinez and C. D. Savage, Patterns in inversion sequences II: inversion sequences avoiding triples of relations. *J. Integer Seq.*, **21** (2018), Article 18.2.2.
- [25] The On-Line Encyclopedia of Integer Sequences, published electronically at <http://oeis.org>, 2010.
- [26] E. Rowland, Pattern avoidance in binary trees. *J. Combin. Theory Ser. A*, **117** (2010), 741–758.
- [27] R. P. Stanley, Enumerative combinatorics: volume 2. Cambridge University Press, Cambridge, 1999.
- [28] S. Yakubov, Pattern avoidance in extensions of comb-like posets. *J. Comb.*, **6** (2015), 249–272.