

MOCK MODULAR EISENSTEIN SERIES WITH NEBENTYPUS

MICHAEL H. MERTENS, KEN ONO, AND LARRY ROLEN

In celebration of Bruce Berndt's 80th birthday

ABSTRACT. By the theory of Eisenstein series, generating functions of various divisor functions arise as modular forms. It is natural to ask whether further divisor functions arise systematically in the theory of mock modular forms. We establish, using the method of Zagier and Zwegers on holomorphic projection, that this is indeed the case for certain (twisted) “small divisors” summatory functions $\sigma_{\psi}^{\text{sm}}(n)$. More precisely, we show that

$$\mathcal{E}_{\psi}^{+}(\tau) := \frac{1}{\theta_{\psi}(\tau)} \cdot \sum_{n=1}^{\infty} \sigma_{\psi}^{\text{sm}}(n) q^n$$

is a half-integral weight mock modular form, where $\theta_{\psi}(\tau)$ is a Shimura theta function. These include generating functions for combinatorial objects such as the Andrews *spt*-function and the “consecutive parts” partition function. Finally, in analogy with Serre’s result that the weight 2 Eisenstein series is a p -adic modular form, we show that these forms possess canonical congruences with modular forms.

1. INTRODUCTION AND STATEMENT OF RESULTS

In the theory of mock theta functions and its applications to combinatorics as developed by Andrews, Hickerson, Watson, and many others, various formulas for q -series representations have played an important role. For instance, the generating function $R(\zeta; q)$ for partitions organized by their ranks is given by:

$$R(\zeta; q) := \sum_{\substack{n \geq 0 \\ m \in \mathbb{Z}}} N(m, n) \zeta^m q^n = \sum_{n \geq 0} \frac{q^{n^2}}{(\zeta q; q)_n (\zeta^{-1} q; q)_n} = \frac{1 - \zeta}{(q; q)_{\infty}} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{\frac{n(3n+1)}{2}}}{1 - \zeta q^n},$$

where $N(m, n)$ is the number of partitions of n of rank m and $(a; q)_j := \prod_{i=0}^{j-1} (1 - aq^i)$ is the usual q -Pochhammer symbol. The function $R(\zeta; q)$, when ζ is specialized to any root of unity times a certain rational power of q , is a mock modular form on a congruence subgroup; see [11, 24, 38] for discussion and applications. For instance, Ramanujan’s original example of the third order mock theta function $f(q)$ is

$$f(q) := R(-1; q) = \sum_{n \geq 0} \frac{q^{n^2}}{(-q; q)_n^2},$$

which shows that the Fourier coefficients of $f(q)$ enumerate the number of partitions of n with even rank minus the number of partitions of n into odd rank.

Many identities involving partition ranks are subtle and require delicate q -series techniques to prove. Thus, it is interesting to ask for further constructions of mock theta functions that will likely play a future role in the study of ranks and related partition functions. Here we take inspiration from the fact that many classical modular forms are generated by Eisenstein series which, thanks to

2010 *Mathematics Subject Classification.* 11F03, 11F37, 11F30.

The second author thanks the support of the NSF and the Asa Griggs Candler Fund.

their representations as Lambert series, have nice q -expansions in terms of divisor power sums. Such representations have long been a staple in the theory of modular forms and their applications.

To review one classical situation, we first recall some notation. The v -th power divisor functions $\sigma_v(n) := \sum_{d|n} d^v$, for odd $v \geq 3$, arise naturally in the theory of modular forms. Indeed, their generating functions are essentially the classical Eisenstein series $G_k(\tau)$, defined for positive even integers k by

$$G_k(\tau) := \frac{1}{2}\zeta(1-k) + \sum_{n=1}^{\infty} \sigma_{k-1}(n)q^n,$$

where $q := e^{2\pi i\tau}$. In particular, if $k \geq 4$, then $G_k(\tau)$ is a weight k holomorphic modular form for the full modular group $\mathrm{SL}_2(\mathbb{Z})$.

Hecke's well-known theory [21] of modular forms includes many further divisor functions. For example, for a primitive Dirichlet character χ modulo N , consider the complementary divisor functions

$$\sigma_{v,\chi}(n) := \sum_{d|n} \chi(d)d^v \quad \text{and} \quad \sigma'_{v,\chi}(n) := \sum_{d|n} \chi(n/d)d^v.$$

If $k > 2$ and $(-1)^k = \chi(-1)$, then Hecke's work implies that the functions

$$G_{k,\chi}(\tau) := \frac{1}{2}L(1-k, \chi) + \sum_{n=1}^{\infty} \sigma_{k-1,\chi}(n)q^n \quad \text{and} \quad \widehat{G}_{k,\chi}(\tau) := \sum_{n=1}^{\infty} \sigma'_{k-1,\chi}(n)q^n$$

are holomorphic weight k modular forms on $\Gamma_0(N)$ with Nebentypus χ .

The weight $k = 2$ case is particularly interesting. Hecke obtained a modification of $G_2(\tau)$ which establishes that

$$E_2^*(\tau) := 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n)q^n - \frac{3}{\pi\mathfrak{S}(\tau)}$$

is a weight 2 harmonic Maaß form. Therefore, we may view $E_2(\tau) := 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n)q^n$ as a ‘‘mock modular form’’, the holomorphic part of the nonholomorphic modular form $E_2^*(\tau)$. In view of this example, it is natural to ask whether there are further divisor functions which arise naturally in the theory of mock modular forms. We show that this is indeed the case for the generating functions of certain twisted ‘‘small divisors’’ summatory functions. In particular, we shall associate these functions with Shimura's weight $1/2$ and $3/2$ theta functions.

To make this precise, suppose that $\psi \pmod{f_\psi}$ is a primitive Dirichlet character with conductor $f_\psi > 1$. We then define $\lambda_\psi := (1 - \psi(-1))/2$, so that $\lambda_\psi = 0$ (resp. $\lambda_\psi = 1$) if ψ is even (resp. odd). The corresponding Shimura theta function [35] of weight $\lambda_\psi + \frac{1}{2}$ on $\Gamma_0(4f_\psi^2)$ with Nebentypus ψ is defined as

$$(1.1) \quad \theta_\psi(\tau) := \begin{cases} \sum_{n=1}^{\infty} \psi(n)q^{n^2} & \text{if } \lambda_\psi = 0, \\ \sum_{n=1}^{\infty} \psi(n)nq^{n^2} & \text{if } \lambda_\psi = 1. \end{cases}$$

If χ_{-4} is the nontrivial Dirichlet character modulo 4, then we have that $\theta_\psi(\tau) \in M_{\frac{1}{2}}(\Gamma_0(4f_\psi^2), \psi)$ (resp. $S_{\frac{3}{2}}(\Gamma_0(4f_\psi^2), \psi\chi_{-4})$) if $\lambda_\psi = 0$ (resp. $\lambda_\psi = 1$).

We define a twisted ‘‘small divisors’’ summatory function for any non-trivial Dirichlet character ψ . To ease notation, if n is a positive integer, then we define its small divisors by

$$(1.2) \quad D_n := \{d \mid n : 1 \leq d \leq n/d \text{ and } d \equiv n/d \pmod{2}\}.$$

Then the ψ -twisted small divisors summatory function is defined by

$$(1.3) \quad \sigma_{\psi}^{\text{sm}}(n) := \sum_{d \in D_n} \psi \left(\frac{(n/d)^2 - d^2}{4} \right) d.$$

We let $\mathcal{E}_{\psi}^+(\tau)$ denote its generating function divided by $\theta_{\psi}(\tau)$, as in the following expression

$$(1.4) \quad \mathcal{E}_{\psi}^+(\tau) := \frac{1}{\theta_{\psi}(\tau)} \cdot \sum_{n=1}^{\infty} \sigma_{\psi}^{\text{sm}}(n) q^n.$$

It turns out that $\mathcal{E}_{\psi}^+(\tau)$ is a polar mock modular form, the holomorphic part of a weight $\frac{3}{2} - \lambda_{\psi}$ *polar harmonic Maaß form*. Recall that polar harmonic Maaß forms have holomorphic parts which are permitted to have poles in \mathbb{H} (see Ch. 13 of [9]). Its nonholomorphic part is given by the normalized period integral

$$(1.5) \quad \mathcal{E}_{\psi}^-(\tau) := (-1)^{\lambda_{\psi}} \frac{(2\pi)^{\lambda_{\psi} - 1/2} i}{2\Gamma(1/2 + \lambda_{\psi})} \cdot \int_{-\tau}^{i\infty} \frac{\theta_{\psi}^-(z)}{(-i(z + \tau))^{3/2 - \lambda_{\psi}}} dz.$$

Using the constructive method of Zagier and Zwegers (for example, see [4, 10, 23]), where harmonic Maaß forms are constructed via holomorphic projection, we obtain the following theorem.

Theorem 1.1. *Assuming the notation above, we have that $\mathcal{E}_{\psi}(\tau) := \mathcal{E}_{\psi}^+(\tau) + \mathcal{E}_{\psi}^-(\tau)$ is a weight $\frac{3}{2} - \lambda_{\psi}$ polar harmonic weak Maaß form on $\Gamma_0(4f_{\psi}^2)$ with Nebentypus $\overline{\psi} \cdot \chi_{-4}^{\lambda_{\psi}}$.*

Remark. The harmonic Maaß forms $\mathcal{E}_{\psi}(\tau)$ are preimages of $\theta_{\psi}(\tau)$ under the Bruinier-Funke ξ -operator (see Section 5.2 of [9]). In other words, Shimura's theta functions are the *shadows* of these canonical polar harmonic Maaß forms.

Remark. It is clear from the definition that the “mock Eisenstein series” $\mathcal{E}_{\psi}^+(\tau)$ can only have poles in the zeros of $\theta_{\psi}(\tau)$. It is easy to construct a weakly holomorphic modular form $G_{\psi}(\tau)$ of weight 2 for $\Gamma_0(4f_{\psi})$ with trivial Nebentypus such that $\mathcal{E}_{\psi}(\tau) - G_{\psi}(\tau)/\theta_{\psi}(\tau)$ is a harmonic weak Maaß form with no poles in the interior of the upper half-plane.

Remark. An elementary computation reveals that the q -series generating function of $\sigma_{\psi}^{\text{sm}}(n)$ (i.e. without the $1/\theta_{\psi}(\tau)$ factor) can be given in terms of derivatives of Appell-Lerch sums as studied by Zwegers [39, 40]. For a special case of this, see for instance [30, Lemma 2]. The forms obtained in Theorem 1.1 have the pleasing property that both factors depend on ψ in a simple way which is analogous to the standard theory of holomorphic modular forms with Nebentypus.

A natural question arises as to which of the functions $\mathcal{E}_{\psi}^+(\tau)$ from Theorem 1.1 are actually mock modular forms, that is, which don't actually have poles on the upper half-plane. For example, this situation occurs whenever $\theta_{\psi}(\tau)$ has no zeros in the upper half-plane. Due to the infinite product expansion of the Dedekind eta function $\eta(\tau) := q^{\frac{1}{24}} \prod_{n \geq 1} (1 - q^n)$, eta-quotients automatically have no zeros on the upper half-plane. In fact, Rouse and Webb showed in [33] that the *only* modular forms on $\Gamma_0(N)$ with integer Fourier coefficients and no zeros on the upper half-plane are eta quotients. The classification of such theta functions was completed by Mersmann [28] and Lemke Oliver in [26].

Theorem 1.2. *The only nontrivial primitive characters ψ where $\theta_{\psi}(\tau)$ is an eta-quotient (in which case $\mathcal{E}_{\psi}^+(\tau)$ is a mock modular form), are when ψ is in the set of Kronecker characters $\chi_D := \left(\frac{D}{\cdot}\right)$ where D is in the following set of fundamental discriminants:*

$$\{-8, -4, -3, 2, 12, 24\}.$$

Several of the mock Eisenstein series in Theorem 1.2 occur naturally in the theory of partitions. Here we point two particularly well known examples. We let $spt(n)$ denote the Andrews *smallest parts function* [3], which counts the number of smallest parts among partitions of n , whose mock modular properties were first exhibited and exploited by Bringmann [8] and by Folsom and the second author in [18]. Furthermore, we will denote by $cpt(n)$ the total number of parts in all partitions of n into *consecutive integers*. Connections between odd divisors of integers and partitions into consecutive parts have also been noted previously (see works of Hirschhorn and Hirschhorn [22] as well as Sylvester and Franklin [37]).

We summarize these combinatorial examples which arise out of Theorem 1.2 in the following.

Corollary 1.3. *The following are true:*

i) *The functions*

$$\mathcal{E}_{\chi_{12}}^+(\tau) = -2q^{-1} \sum_{n \geq 1} spt(n)q^{24n}$$

and

$$\mathcal{E}_{\chi_{24}}^+(\tau) = -2q^{-1} \sum_{n \geq 1} (-1)^n spt(n)q^{24n}$$

are mock modular forms of weight $3/2$.

ii) *The function*

$$\theta_{\chi_2}(\tau) \cdot \mathcal{E}_{\chi_2}^+(\tau) = 2 \sum_{n \geq 1} (-1)^n cpt(n)q^{8n}$$

is a mixed mock modular form of weight 2.

Remark. There are other prominent examples which have appeared in the literature outside of combinatorial number theory. For instance, in work related to quantum black holes, we find

$$\theta_{\chi_{-4}}(\tau) \cdot \mathcal{E}_{\chi_{-4}}^+\left(\frac{\tau}{8}\right) = \sum_{\substack{r > s > 0 \\ r-s \equiv 1 \pmod{2}}} (-1)^r s q^{\frac{rs}{2}} =: F_2^{(2)}(\tau),$$

where $F_2^{(2)}(\tau)$ is the series which Dabholkar, Murthy, and Zagier computed using holomorphic projection (cf. [13, Section 7.1, Example 4]). They use this form to construct the mock modular form

$$h^{(2)}(\tau) := \frac{24F_2^{(2)}(\tau) - E_2(\tau)}{\eta(\tau)^3},$$

which has shadow $\eta^3(\tau)$, and was first described in works of Eguchi-Taormina [16] and Eguchi-Ooguri-Taormina-Yang [15] in relation to the elliptic genus. It has since played an important role in Mathieu Moonshine [12, 14] and in relation to $SO(3)$ Donaldson invariants of $\mathbb{C}P^2$ [27].

Although the Eisenstein series $E_2(\tau)$ is not modular, Serre [34] proved that it is a p -adic modular form for every prime p . Namely, for every prime power p^t he identified $E_2(\tau) \pmod{p^t}$ as the reduction of a genuine holomorphic modular form. Moreover, he found that such sequences of modular forms, as $t \rightarrow +\infty$, have weights that p -adically converge to 2.

In view of Theorem 1.1, it is natural to ask whether the mock modular Eisenstein series $\mathcal{E}_{\psi}^+(\tau)$ possess similar p -adic properties. In analogy with Serre's construction, we indeed show that these mock modular Eisenstein series satisfy analogous congruences with genuine meromorphic modular forms modulo arbitrary prime powers. Under repeated iteration of the U -operator

$$(1.6) \quad \left(\sum_{n \gg -\infty} a(n)q^n \right) \Big| U(p) := \sum_{n \gg -\infty} a(pn)q^n,$$

we obtain the following result.

Theorem 1.4. *Let $p > 3$ be a prime and $a, b \in \mathbb{N}$. Then there is a weight 2 meromorphic modular form $F_\psi^{(a,b)}(\tau)$ for the group $\Gamma_0(4f_\psi^2 p^m)$, with $m = \max(2a - b, 1 - 2v_p(f_\psi))$, with trivial Nebentypus such that we have the congruence*

$$\left(\theta_\psi(p^{2a}\tau)\mathcal{E}_\psi^+(\tau)\right) |U(p^b) \equiv F_\psi^{(a,b)}(\tau) \pmod{p^{\min(a,b)}}.$$

Remark. In the special case of $spt(n)$, these p -adic properties have been studied previously. Andrews found [3] that the spt -function satisfies the following Ramanujan-type congruences:

$$spt(5n + 4) \equiv 0 \pmod{5}, \quad spt(7n + 5) \equiv 0 \pmod{7}, \quad spt(13n + 6) \equiv 0 \pmod{13},$$

and the second author and Folsom in [18] obtained many further congruences in [32]. Garvan [19] and later Ahlgren and Kim [2] used the modularity properties to study the context of Andrews' congruences. They showed that Andrews' congruences are "explained" by the triviality of the space of cusp forms of level one on and weights $5 + 1, 7 + 1, 13 + 1$, and surmised the general conjecture that no further congruences of this type exist for primes $\ell \geq 11$. As for prime power congruences, results close to p -adic modularity along the lines of Theorem 1.4 have been obtained in [6].

The paper is organized as follows. In Subsection 2.1 we will review the theory of holomorphic projection and establish some preliminary results, which will be exploited in Subsection 2.2 to prove our main results on the construction of the polar mock modular Eisenstein series. We then derive the examples, including $spt(n)$ and $cpt(n)$, where we obtain mock modular forms in Section 3. We conclude in Section 4 with the proof of our p -adic congruence results.

2. MODULARITY PROPERTIES OF NEW DIVISOR FUNCTION GENERATING FUNCTIONS

Here we prove Theorem 1.1. To this end we first recall the important work of Zwegers, which we then manipulate to obtain the desired generating functions.

2.1. Holomorphic projection. Holomorphic projection is now an important tool in the study of non-holomorphic modular forms, in particular harmonic Maaß forms. It was first introduced by Sturm [36] and further developed by Gross and Zagier in their seminal work on derivatives of L -functions [20]. For some applications in the context of mock modular forms and harmonic Maaß forms, the reader may consult for instance [1, 4, 10, 23, 29, 31], among several others. For background on the subject of harmonic Maaß forms, we refer the reader to [9].

Let us first recall the definition of (regularized) holomorphic projection and explicit expressions for holomorphic projection of the product of a harmonic Maaß form with a holomorphic modular form. These can be found e.g. in [23], where a vector-valued version of the same question was considered, or in [29], where Rankin-Cohen brackets of such functions were examined.

The usual holomorphic projection operator is defined for a (not necessarily holomorphic) modular form Φ with Fourier expansion

$$(2.1) \quad \Phi(\tau) = \sum_{n \in \mathbb{Z}} c(n, y) q^n$$

of weight $k \geq 2$ for some subgroup $\Gamma \leq \mathrm{SL}_2(\mathbb{Z})$ with moderate growth towards the cusps as

$$\pi_{hol}(\Phi)(\tau) := c_0 + \sum_{n=1}^{\infty} c_n q^n$$

with

$$(2.2) \quad c_n := \frac{(4\pi n)^{k-1}}{\Gamma(k-1)} \int_0^\infty c(n, y) e^{-4\pi n y} y^{k-2} dy, \quad (n \geq 1),$$

provided that the integral converges, and where c_0 is an appropriate constant term. This operator maps into the space of holomorphic modular forms for $k > 2$ and to the space of holomorphic quasimodular forms for $k = 2$. It also behaves like a projection in that it acts as the identity on holomorphic functions, and it preserves the Petersson inner product, i.e. for every cusp form $g \in S_k(\Gamma)$ we have

$$\langle \Phi, g \rangle = \langle \pi_{hol}(\Phi), g \rangle.$$

This is in fact the origin of the operator's definition.

For many applications, the growth conditions on the function Φ are too restrictive. If instead of moderate growth towards the cusps, one requires that at each cusp \mathfrak{a} there is a polynomial $H_{\mathfrak{a}} \in \mathbb{C}[X]$ such that

$$\Phi - H_{\mathfrak{a}}(q^{-1/h}),$$

where h denotes the width of the cusp \mathfrak{a} , has moderate growth approaching \mathfrak{a} , and requires that the coefficients of Φ satisfy $c(n, y) = O(y^{2-k})$ as $y \rightarrow 0$ uniformly for all $n > 0$, one obtains a *regularized* holomorphic projection operator

$$\pi_{hol}^{reg}(\Phi)(\tau) := H_{i\infty}(q^{-1}) + \sum_{n=1}^{\infty} \tilde{c}_n q^n$$

with

$$(2.3) \quad \tilde{c}_n := \frac{(4\pi n)^{k-1}}{\Gamma(k-1)} \lim_{s \rightarrow 0} \int_0^{\infty} c(n, y) e^{-4\pi n y} y^{k-2-s} dy, \quad (n \geq 1),$$

see [31].

It is a basic fact (see for instance [9, Section 4.2]) that a harmonic Maaß form $f \in H_k(\Gamma)$ of weight $k \neq 1$ naturally splits into a holomorphic and a non-holomorphic part,

$$f(\tau) = f^+(\tau) + f^-(\tau)$$

with

$$(2.4) \quad f^+(\tau) = \sum_{n \gg -\infty} c_f^+(n) q^n \quad \text{and} \quad f^-(\tau) = \frac{(4\pi y)^{1-k}}{1-k} c_f^-(0) + \sum_{n=1}^{\infty} \frac{c_f^-(n)}{n^{k-1} \Gamma(1-k; 4\pi n y)} q^{-n}$$

With this, one obtains the following result.

Proposition 2.1. *Let $f \in H_k(\Gamma)$ be a harmonic Maaß form as in (2.4) with $c_f^-(0) = 0$ and $g \in M_{2-k}(\Gamma)$ with Fourier expansion*

$$g(\tau) = \sum_{n=1}^{\infty} b(n) q^n.$$

Then the regularized holomorphic projection $\pi_{hol}^{reg}(f \cdot g)$ is well-defined and we have

$$\pi_{hol}^{reg}(f \cdot g)(\tau) = f^+(\tau)g(\tau) + \sum_{r=1}^{\infty} c(r) q^r$$

where

$$c(r) = \frac{\Gamma(2-k)}{k-1} \sum_{m=1}^{\infty} \frac{c_f^-(m)}{m^{k-1}} b(m+r) \left[(r+m)^{k-1} - m^{k-1} \right].$$

It is sometimes convenient to define holomorphic projection for translation-invariant functions as done for instance in [10, Lemma 3.1]. Here we consider the following regularization of this definition in order to make it applicable to functions that grow towards the boundary of the upper half-plane. For $T > 1$ let

$$\mathbb{H}_T := \{ \tau \in \mathbb{H} : 1/T < \Im(\tau) < T \}$$

denote the truncated upper half-plane. Note that the group $\mathrm{SL}_2(\mathbb{Z})$ acts on \mathbb{H}_T in the usual way. Following Borcherds' regularization of the Petersson inner product [7] which was used in [31] to define regularized holomorphic projection, one may follow the proofs of [10, Lemmas 3.1-3.3] to show the following result. We leave the details to the reader.

Lemma 2.2. *Let $\Phi : \mathbb{H} \rightarrow \mathbb{C}$ be a translation invariant function with Fourier expansion as in (2.1) and $\kappa > 1$. Then the following are true:*

(1) *We have that $\pi_{hol}^{reg} f(\tau)$ equals the constant term (in s) in the Laurent expansion of*

$$\frac{(\kappa - 1)(2i)^\kappa}{4\pi} \lim_{T \rightarrow \infty} \int_{\mathbb{H}_T} \frac{f(x + iy)y^{k+2s} dx dy}{(\tau - x + iy)^\kappa y^2},$$

provided that both expressions are well-defined.

(2) *If $f : \mathbb{H} \rightarrow \mathbb{C}$ is holomorphic and translation invariant, we have for either of the possible expressions of π_{hol}^{reg} that $\pi_{hol}^{reg} f = f$.*

(3) *For $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ we have*

$$(\pi_{hol}^{reg} f)|_\kappa \gamma = \pi_{hol}^{reg} (f|_\kappa \gamma)$$

From this, one can derive the following result as a generalization of [10, Proposition 3.5].

Proposition 2.3. *Let $f : \mathbb{H} \rightarrow \mathbb{C}$ be a translation invariant function such that Lemma 2.2 (1) is satisfied for $k > 1$. If the weight k holomorphic projection of f vanishes identically and $\xi_k f$ is modular of weight $2 - k$ for some subgroup $\Gamma \leq \mathrm{SL}_2(\mathbb{Z})$, then f is modular of weight k for Γ .*

2.2. Proof of Theorem 1.1.

Proof of Theorem 1.1. It is clear from the definition that \mathcal{E}_ψ is annihilated by the weight $1/2 + \lambda_\psi$ hyperbolic Laplacian away from the singularities, which lie precisely in the zeros of θ_ψ , and that it has the correct growth properties towards the cusps. We are therefore left with showing modularity. The expression for $\pi_{hol}^{reg}(\theta_\psi \mathcal{E}_\psi)$ in Lemma 2.2 (1) is easily seen to be well-defined. A standard computation further reveals that

$$\mathcal{E}_\psi^- = -\frac{1/2 - \lambda}{\Gamma(1/2 + \lambda_\psi)} \sum_{n=1}^{\infty} \frac{\psi(n)n^\lambda}{n^{1-2\lambda_\psi}} \Gamma(\lambda_\psi - 1/2; 4\pi n^2 y) q^{-n^2},$$

wherefore we obtain from Proposition 2.1 that

$$\pi_{hol}(\theta_\psi \mathcal{E}_\psi) = \sum_{n=1}^{\infty} \sigma_\psi^{sm}(n) q^n - \sum_{m>r \geq 1} \psi(mr)(m-r) q^{m^2-r^2},$$

which is easily seen to be identically zero. Since

$$\xi_2(\theta_\psi \mathcal{E}_\psi)(\tau) = 2^{\lambda_\psi - 1/2} i y^{\lambda_\psi + 1/2} |\theta_\psi(\tau)|^2$$

is clearly modular of weight 0 for the group $\Gamma_0(4f_\psi^2)$ with Nebentypus $\chi_{-4}^{\lambda_\psi}$, the claim follows by applying Proposition 2.3. \square

3. PROOF OF THEOREM 1.2 AND COROLLARY 1.3

The proof of Theorem 1.2 follows directly from the characterized list of theta functions which are also eta quotients in [26], as per the discussion preceding the theorem.

Proof of Corollary 1.3. Firstly, note that in part i) , the character χ_{24} is $\left(\frac{2}{n}\right)$ times the first character χ_{12} . Thus, switching from the first example to the second twists the theta function in the denominator of (1.4) by $\left(\frac{2}{n}\right)$, and a similar check can be performed on the character in the sum in (1.3). Thus, it is readily checkable that the extra twist by $(-1)^n$ occurs in this case.

It remains to check that the *spt* and *cpt* functions arise out of these constructions. For both, the mock modularity properties are already known, at least implicitly in the literature. For instance, the mock modularity of the generating function of *spt*(n) is described in [18]), and our Theorem 1.2 establishes the mock modularity of our series $\mathcal{E}_{\chi_{12}}^+(\tau)$. Similarly, for *cpt*(n), the generating function can be shown elementary means to have an Appell-Lerch-type generating function

$$\sum_{n \geq 1} \text{cpt}(n)q^n = \sum_{n \geq 1} \frac{nq^{\frac{n(n+1)}{2}}}{1 - q^n}$$

(see the proof of Joerg Arndt in the comments on the OEIS sequence A204217). This is readily expressible as a theta function times a Taylor coefficient in an elliptic variable of Zwegers' μ -function [39], which by the general theory there is a weight $3/2$ mock modular form. Normalizing by dividing by the theta function, the identity for *cpt*(n) is thus reduced to checking that two weight $3/2$ mock modular forms in known spaces are equal. Thus, to prove the claims, one first needs to check that they live in the same space of mock modular forms, that is, have the same weight, level, and Nebentypus, and that their principal parts at the cusps agree. This check guarantees that the claimed identities are true up to the addition of a cusp form of weight $3/2$, level 576, and character χ_{12} in the case of *spt* and weight $3/2$, level 256, and character χ_2 in the case of *cpt*. This reduces the equality to a finite check, which we omit here. \square

4. PROOF OF THEOREM 1.4

We now turn to the proof of our claim on the behavior of the generating functions modulo powers of primes.

Proof of Theorem 1.4. As remarked after the statement of Theorem 1.1, there is a weakly holomorphic modular form $G_\psi \in M_2^!(\Gamma_0(4f_\psi^2))$ such that $\mathcal{E} - G_\psi/\theta_\psi$ has no poles in the upper half-plane, so we may and will assume this from now on. By Proposition 2.1, we find that

$$(4.1) \quad \pi_{hol}^{reg}(\theta_\psi(p^{2a}\tau)\mathcal{E}_\psi(\tau)) = \theta_\psi(p^{2a}\tau)\mathcal{E}_\psi^+(\tau) + \sum_{r=1}^{\infty} \left(\sum_{\substack{(p^am)^2 - n^2 = r \\ m, n \geq 0}} \psi(mn)(p^am - n) \right) q^r$$

is a weakly holomorphic quasimodular form. This is congruent to a weakly holomorphic modular form $F \in M_2^!(\Gamma_0(4f_\psi^2 p^{2a}))$ because the Eisenstein series E_2 is congruent to the holomorphic modular form $E_2(\tau) - NE_2(N\tau) \in M_2(\Gamma_0(N))$ for any $N > 1$.

Since $(p^am)^2 - n^2 = (p^am - n)(p^am + n)$ factors, we can rewrite the inner sum in the second term in (4.1) as a sum over small divisors of r ,

$$(4.2) \quad \sum_{\substack{(p^am)^2 - n^2 = r \\ m, n \geq 0}} \chi(mn)(p^am - n) = \sum_{\substack{d|r, d \leq r/d \\ d+r/d \equiv 0 \pmod{2p^a} \\ d-r/d \equiv 0 \pmod{2}}} \chi\left(\frac{(r/d)^2 - d^2}{4p^a}\right) d,$$

as in the proof of Theorem 1.1. We now apply the operator $U(p^b)$ to (4.1). This corresponds to replacing r by $p^b r$ in (4.2), and then the condition $d+p^b r/d \equiv 0 \pmod{2p^a}$ forces $d \equiv 0 \pmod{p^{\min(a,b)}}$,

so that we find that

$$\pi_{hol}^{reg}(\theta_\psi(p^{2a}\tau)\mathcal{E}_\psi(\tau))|U(p^b) \equiv \left(\theta_\psi(p^{2a}\tau)\mathcal{E}_\psi^+(\tau)\right)|U(p^b) \equiv F|U(p^b) \pmod{p^{\min(a,b)}}.$$

By [5, Lemma 6 and Lemma 7], $F|U(p^b)$ is modular for $\Gamma_0(4f_\psi^2 p^m)$ for $m = \max(2a - b, 1 - 2v_p(f_\psi))$, as stated in the Theorem. This completes the proof. \square

REFERENCES

- [1] S. Ahlgren and N. Andersen, *Euler-like recurrences for smallest parts functions*, Ramanujan J. **36** (2015), no. 1-2, 237–248.
- [2] S. Ahlgren and B. Kim, *Congruences for a mock modular form on and the smallest parts function*, J. Number Theory **189** (2018), 81–89.
- [3] G. E. Andrews, *The number of smallest parts in the partitions of n* , J. Reine Angew. Math. **624** (2008), 133–142.
- [4] G. E. Andrews, R. C. Rhoades, and S. P. Zwegers *Modularity of the concave composition generating function*, Algebra Number Theory **7** (2013), no. 9, 2103–2139.
- [5] A. O. L. Atkin and J. Lehner, *Hecke operators on $\Gamma_0(m)$* , Math. Ann. **185** (1970), 134–160.
- [6] E. Belmont, H. Lee, A. Musat, S. Trebat-Leder, *ℓ -adic properties of partition functions*, Monatshefte für Mathematik, **173** no. 1 (2014), 1–34.
- [7] R. E. Borcherds, *Automorphic forms with singularities on Grassmannians*, Invent. Math. **132** (1998), 491–562.
- [8] K. Bringmann, *On the explicit construction of higher deformations of partition statistics*, Duke Math J. **144** 195–233.
- [9] K. Bringmann, A. Folsom, K. Ono, and L. Rolén, *Harmonic Maass forms and mock modular forms: Theory and applications*, Amer. Math. Soc., Providence, 2018.
- [10] K. Bringmann, B. Kane, and S. Zwegers, *On a completed generating function of locally harmonic Maass forms*, Compos. Math. **150** (2014), no. 5, 749–762.
- [11] K. Bringmann and K. Ono, *Dyson’s ranks and Maass forms*, Ann. Math. **171** (2010), 419–449.
- [12] M. Cheng, *$K3$ surfaces, $N = 4$ Dyons, and the Mathieu group M_{24}* , Commun. Number Theory Phys. **4** (2010), 623–657.
- [13] A. Dabholkar, S. Murthy, and D. Zagier, *Quantum Black Holes, Wall Crossing, and Mock Modular Forms*, to appear in Cambridge Monographs in Mathematical Physics, 151 pages.
- [14] T. Eguchi, H. Ooguri, and Y. Tachikawa, *Notes on the $K3$ surface and the Mathieu group M_{24}* , Exp. Math. **20** no. 1 (2011), 91–96.
- [15] T. Eguchi, H. Ooguri, S.-K. Yang, and A. Taormina, *Superconformal algebras and string compactification on manifolds with $SU(N)$ Holonomy*, Nucl. Phys. B **315** (1989), 193.
- [16] T. Eguchi and A. Taormina, *On the unitary representation of $N = 2$ and $N = 4$ superconformal algebras*, Phys. Lett. B **210** (1988), 125.
- [17] A. El Basraoui and A. Sebbar, *Zeros of the Eisenstein Series E_2* , Proc. Amer. Math. Soc. **137** no. 7 (2010), 2289–2299.
- [18] A. Folsom and K. Ono, *The spt -function of Andrews*, Proc. Natl. Acad. Sci. USA **105** no. 51 (2008), 20152–20156.
- [19] F. Garvan, *Congruences for Andrews’ spt -function modulo powers of 5, 7, and 13*, Trans. Amer. Math. Soc. **364** no. 9 (2012), 4847–4873.
- [20] B. H. Gross and D. B. Zagier, *Heegner points and derivatives of L -series*, Invent. Math. **84** (1986), 225–320.
- [21] E. Hecke, *Theorie der Eisensteinschen Reihen höherer Stufe und ihre Anwendung auf Funktionentheorie und Arithmetik*, Abh. Math. Sem. Hamburg. Univ. **5** (1927), 199–224; in E. Hecke, *Mathematische Werke*, Vandenhoeck & Ruprecht Göttingen (1959), pp. 461–486.
- [22] M.D. Hirschhorn and P.M. Hirschhorn, *Partitions into consecutive parts*, Math. Mag. **78** no. 5 (2005), 396–397.
- [23] Ö. Imamoglu, M. Raum, and O. Richter. *Holomorphic projections and Ramanujan’s mock theta functions*, Proc. Nat. Acad. Sci. U.S.A. **111.11** (2014), 3961–3967.
- [24] S.-Y. Kang, *Mock Jacobi forms in basic hypergeometric series*, Compos. Math. **145** (2009), no. 3, 553–565.
- [25] W. Kohnen, *On a certain class of modular functions*, Proc. Amer. Math. Soc. **133** (2005), no. 1, 65–70.
- [26] R. Lemke Oliver. *Eta-quotients and theta functions*, Adv. Math. **241** (2013), 1–17.
- [27] A. Malmendier and K. Ono, *$SO(3)$ -Donaldson invariants of $\mathbb{C}P^2$ and mock theta functions*, Geom. Topol., **16** (2012), 1767–1833.
- [28] G. Mersmann. *Holomorphe η -produkte und nichtverschwindende ganze modulformen für $\Gamma_0(n)$* , Diploma Thesis, Univ. of Bonn, 1991.
- [29] M. H. Mertens, *Eichler-Selberg type identities for mixed mock modular forms*, Adv. Math. **301** (2016), 359–382.
- [30] M. H. Mertens, *Mock modular forms and class number relations*, Res. Math. Sci. **1** (2014), Art. 6, 16 pp.

- [31] M. H. Mertens and K. Ono, *Special values of shifted convolution Dirichlet series*, *Mathematika* **62** (2016), no. 1, 47–66.
- [32] K. Ono, *Congruences for the Andrews spt function*, *Proc. Natl. Acad. Sci. USA* **108** no. 2 (2011) 473–476.
- [33] J. Rouse and J.J. Webb, *Spaces of modular forms spanned by eta-quotients*, *Adv. Math.* **272** (2015), 200–224.
- [34] J.-P. Serre, *Formes modulaires et fonctions zêta p-adiques*, (French), *Modular functions of one variable, III* (Proc. Int. Summer School, Univ. Antwerp, 1972), *Springer Lect. Notes Math.* **350**, 1973, pp. 191–268.
- [35] G. Shimura, *On modular forms of half-integral weight*, *Ann. Math.* **97** (1973), 440–481.
- [36] J. Sturm, *Projections of C^∞ automorphic forms*, *Bull. Amer. Math. Soc. (N.S.)* **2** (1980), no. 3, 435–439.
- [37] J. J. Sylvester and F. Franklin, *A constructive theory of partitions, arranged in three acts, an interact and an exodion*, *Amer. J. Math.* **5** (1882), 251–330.
- [38] D. Zagier, *Ramanujan’s mock theta functions and their applications [d’après Zagiers and Bringmann-Ono]*, *Séminaire Bourbaki, 60ème année, 2007-2008*, no. 986, *Astérisque* **326** (2009), Soc. Math. de France, 143–164.
- [39] S. Zagiers, *Mock Theta Functions*, PhD thesis, Universiteit Utrecht, 2002.
- [40] S. Zagiers, *Multivariable Appell functions and nonholomorphic Jacobi forms*, *Res. Math. Sci.* **6** (2019), Art. 16, 15 pp.

MAX-PLANCK-INSITUT FÜR MATHEMATIK BONN, VIVATSGASSE 7, 53111 BONN, GERMANY
E-mail address: `mhmertens@mpim-bonn.mpg.de`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904
E-mail address: `ken.ono691@gmail.com`

DEPARTMENT OF MATHEMATICS, 1420 STEVENSON CENTER, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240
E-mail address: `larry.rolen@vanderbilt.edu`