

A NEW COMBINATORIAL INTERPRETATION OF THE FIBONACCI NUMBERS SQUARED

KENNETH EDWARDS AND MICHAEL A. ALLEN*

ABSTRACT. We consider the tiling of an n -board (a $1 \times n$ array of square cells of unit width) with half-squares ($\frac{1}{2} \times 1$ tiles) and $(\frac{1}{2}, \frac{1}{2})$ -fence tiles. A $(\frac{1}{2}, \frac{1}{2})$ -fence tile is composed of two half-squares separated by a gap of width $\frac{1}{2}$. We show that the number of ways to tile an n -board using these types of tiles equals F_{n+1}^2 where F_n is the n th Fibonacci number. We use these tilings to devise combinatorial proofs of identities relating the Fibonacci numbers squared to one another and to other number sequences. Some of these identities appear to be new.

1. INTRODUCTION

The Fibonacci numbers can be interpreted as the number of ways to tile an $n \times 1$ array of joined 1×1 cells (called an n -board) with 1×1 squares and 2×1 dominoes [3, 2]. More generally, the number of ways to tile an n -board with all the $r \times 1$ r -ominoes up to $r = k$ is the k -step (or k -generalized) Fibonacci number $F_{n+1}^{(k)} = F_n^{(k)} + F_{n-1}^{(k)} + \dots + F_{n-k+1}^{(k)}$, with $F_1^{(k)} = 1$ and $F_{n < 1}^{(k)} = 0$ [2]. In [4] it was shown that it is possible to obtain a combinatorial interpretation of the Tribonacci numbers (the 3-step Fibonacci numbers) as the number of tilings of an n -board using just two types of tiles, namely, squares and $(\frac{1}{2}, 1)$ -fence tiles. A (w, g) -fence tile is composed of two pieces (called posts) of size $w \times 1$ separated by a gap of size $g \times 1$. In [5] a bijection between tiling with $(\frac{1}{2}, g)$ -fence tiles where $g \in \{0, 1, 2, \dots\}$ and strongly restricted permutations was identified and then used to obtain results concerning the permutations in a straightforward way.

Here we show that the number of ways to tile an n -board using half-squares ($\frac{1}{2} \times 1$ tiles) and $(\frac{1}{2}, \frac{1}{2})$ -fences is a Fibonacci number squared. The Fibonacci numbers F_n are given by

$$F_n = F_{n-1} + F_{n-2}, \quad F_0 = 0, \quad F_1 = 1, \quad n \geq 2. \quad (1.1)$$

We use these tilings to formulate combinatorial proofs in the style of those found in [2] of a number of identities relating to the Fibonacci numbers squared.

2. TYPES OF METATILE

When tiling a board with fences one must first determine the types of metatiles since any tiling of the board can be expressed as a tiling using metatiles [4]. A metatiles is a minimal arrangement of tiles that exactly covers an integral number of adjacent cells [4, 5]. When tiling with half-squares (denoted by h) and $(\frac{1}{2}, \frac{1}{2})$ -fence tiles (henceforth referred to simply as fences and denoted by f), the simplest metatiles is two half-squares (h^2) as illustrated in Fig. 1. This is the only metatiles of length 1. The simplest metatiles of length 2 is the bifence (f^2) which is two interlocking fences. Unlike the case of tiling with squares and $(\frac{1}{2}, 1)$ -fences, the filled fence, which is obtained by filling the gap in the fence by a half-square and is denoted by fh , is not a metatiles since it has length $\frac{3}{2}$. However, by adding a half-square to one end we form length-2 metatiles (hfh and fhh). We may also create metatiles of length $2(j+1)$ by adding




FIGURE 1. A 21-board tiled with all possible metatiles of length less than 5: two half-squares h^2 (cell 1), a bifence f^2 (cells 2–3), hfh (cells 4–5), fhh (cells 6–7), $fhfh$ (cells 8–10), hf^2h (cells 11–13), hf^2fh (cells 14–17), fhf^2h (cells 18–21).

j bifences between a half-square and filled fence. Concatenating two filled fences generates a length-3 metatile ($fhfh$) and inserting j bifences between them generates a metatile of length $2j + 3$. Inserting j bifences between two half-squares generates a metatile of length $2j + 1$. Hence there are two metatiles of length l for all $l \geq 3$.

We will refer to a metatile containing both h and f as being *mixed*. Notice that there are two types of mixed metatile of each length larger than 1. This fact is used in the proofs of Identities 4.2 and 4.3. An h is said to be *captured* if it lies between the posts of a single fence, and *free* otherwise. These notions are used in the proof of Identity 4.10. A bifence is said to be free if it is not part of a larger metatile.

3. THE CORRESPONDENCE BETWEEN NUMBERS OF TILINGS AND THE SQUARES OF THE FIBONACCI NUMBERS

Let A_n be the number of ways to tile an n -board using half-squares and fences.

Lemma 3.1.

$$A_n = \delta_{n,0} + A_{n-1} + 3A_{n-2} + 2 \sum_{i=3}^n A_{n-i}, \tag{3.1}$$

where $\delta_{i,j}$ is 1 if $i = j$ and 0 otherwise, and $A_n = 0$ for $n < 0$.

Proof. Following [1, 5], we condition on the last metatile. If the last metatile is of length l there will be A_{n-l} ways to tile the remaining $n - l$ cells. There is one metatile of length 1, three of length 2, and two metatiles of length l for each $l \geq 3$. If $n = l$ there is exactly one tiling corresponding to that final metatile so we make $A_0 = 1$. There is no way to tile an n -board if $n < l$ and so $A_{n < 0} = 0$. \square

Note that the inclusion of the initial value of the sequence via the $\delta_{n,0}$ term is used to simplify the derivations of recurrence relations obtained from recurrence relations with sums containing arbitrarily many terms by subtracting the recursion relation for A_{n-L} for some L . Without it, one would have to consider the $n = L$ case separately.

Theorem 3.2. *The number of ways to tile an n -board using half-squares and $(\frac{1}{2}, \frac{1}{2})$ -fences is F_{n+1}^2 .*

Proof. Subtracting (3.1) with n changed to $n - 1$ from (3.1) gives

$$A_n = \delta_{n,0} - \delta_{n,1} + 2A_{n-1} + 2A_{n-2} - A_{n-3}, \tag{3.2}$$

i.e., $A_n = 2A_{n-1} + 2A_{n-2} - A_{n-3}$ for $n \geq 3$, where $A_0 = A_1 = 1, A_2 = 4$. The Fibonacci numbers squared obey (3.2) with $A_n = F_{n+1}^2$ (see Identity 30 in [2]). \square

4. IDENTITIES

Identity 4.1.

$$F_n^2 = F_{n-1}^2 + 3F_{n-2}^2 + 2 \sum_{i=3}^n F_{n-i}^2, \quad n \geq 2. \quad (4.1)$$

Proof. It follows from Lemma 3.1 and Theorem 3.2. \square

Identity 4.2.

$$F_{n+3}^2 - 1 = \sum_{k=0}^n \left\{ 3F_{k+1}^2 + 2 \sum_{i=1}^k F_i^2 \right\}, \quad n \geq 0. \quad (4.2)$$

Proof. How many ways are there to tile an $(n+2)$ -board using at least 1 fence? Answer 1: $A_{n+2} - 1$ since this corresponds to all tilings except the all- h tiling. Answer 2: condition on the location of the last fence. Suppose this fence lies on cells $k+1$ and $k+2$ ($k = 0, \dots, n$). Either there is a bifence covering these cells and so there are A_k ways to tile the remaining cells, or the cells are at the end of a mixed metatile and so there are $2(A_{k+2-2} + A_{k+2-3} + \dots + A_0)$ ways to tile the remaining cells. Hence, equating the two answers,

$$A_{n+2} - 1 = \sum_{k=0}^n \{3A_k + 2(A_0 + A_1 + \dots + A_{k-1})\}.$$

The identity then follows from Theorem 3.2. \square

Identity 4.3.

$$F_{2n+2}^2 = F_1^2 + \sum_{k=1}^n \left\{ F_{2k+1}^2 + 2 \sum_{i=1}^{2k} F_i^2 \right\}, \quad n \geq 0. \quad (4.3)$$

Proof. How many ways are there to tile an $(2n+1)$ -board? Answer 1: A_{2n+1} . Answer 2: an odd-length board must have at least one h and the final h must be on an odd cell. Condition on the location of the last h . Suppose that the last h is in cell $2k+1$ ($k = 0, \dots, n$). Either it is part of h^2 and so there are A_{2k} ways to tile the remaining cells, or it is part of a mixed metatile and so there are $2(A_{2k+1-2} + A_{2k+1-3} + \dots + A_0)$ ways to tile the remaining cells. In the latter case, evidently, k cannot be zero. Hence

$$A_{2n+1} = \sum_{k=0}^n A_{2k} + 2 \sum_{k=1}^n (A_0 + A_1 + \dots + A_{2k-1}).$$

The identity then follows from Theorem 3.2. \square

Lemma 4.4. *If S_n is the number of ways to tile an n -board using half-squares and $(\frac{1}{2}, \frac{1}{2})$ -fences such that no free bifences occur then*

$$S_n = 2S_{n-1} + S_{n-2}, \quad n \geq 2, \quad (4.4)$$

where $S_0 = S_1 = 1$.

Proof. Conditioning on the last metatile (which can be any metatile except a bifence) gives

$$S_n = \delta_{0,n} + S_{n-1} + 2 \sum_{j=2}^n S_{n-j} \quad (4.5)$$

Subtracting (4.5) with n replaced by $n-1$ from (4.5) gives $S_n = \delta_{0,n} - \delta_{1,n} + 2S_{n-1} + S_{n-2}$ which is equivalent to (4.4). \square

$S_{n \geq 0} = 1, 1, 3, 7, 17, 41, 99, 239, 577, 1393, \dots$ is sequence A001333 in [6].

Identity 4.5.

$$F_{n+1}^2 = S_n + \sum_{k=2}^n F_{k-1}^2 S_{n-k} \tag{4.6}$$

Proof. How many tilings of an n -board contain at least one free bifence? Answer 1: $A_n - S_n$. Answer 2: condition on the location of the last free bifence. The number of tilings when this final free bifence lies on cells $k - 1$ and k (for $k = 2, \dots, n$) is $A_{k-2}S_{n-k}$. Summing over all possible k and equating the two answers gives

$$A_n - S_n = \sum_{k=2}^n A_{k-2}S_{n-k}$$

and the identity follows from Theorem 3.2. □

Lemma 4.6. *If C_n is the number of ways to tile an n -board using half-squares and $(\frac{1}{2}, \frac{1}{2})$ -fences such that no bifences occur then*

$$C_n = C_{n-1} + 2C_{n-2} + C_{n-3}, \quad n \geq 3, \tag{4.7}$$

where $C_0 = C_1 = 1$ and $C_2 = 3$.

Proof. Conditioning on the last metatile which can only be h^2 , fhh , hfh , or $fhfh$ gives $C_n = \delta_{0,n} + C_{n-1} + 2C_{n-2} + C_{n-3}$ which is equivalent to (4.7). □

$C_{n \geq 0} = 1, 1, 3, 6, 13, 28, 60, 129, 277, 595, \dots$ is sequence A002478 in [6] and consists of the even terms of Narayana’s cows sequence (A000930 in [6]).

Identity 4.7.

$$F_{n+1}^2 = C_n + \sum_{k=2}^n F_{k-1}^2 C_{n-k} + \sum_{k=3}^n \sum_{l=3}^k (2 - \delta_{l,3}) F_{k-l+1}^2 C_{n-k}. \tag{4.8}$$

Proof. How many tilings of an n -board contain at least one bifence? Answer 1: $A_n - C_n$. Answer 2: condition on the location of the last metatile containing a bifence. If the metatile is just a bifence, it can occupy cells $k - 1$ and k ($k = 2, \dots, n$). The number of tilings in this case is $A_{k-2}C_{n-k}$. Otherwise the metatile can occupy cells $k - l + 1$ to k ($k = l, \dots, n$) where l is the length of the metatile. The number of tilings for a metatile of length l for a given k is $A_{k-l}C_{n-k}$. There is one $l = 3$ metatile containing a bifence and two for each $l > 3$. Summing over the possible l and k and equating the two answers gives

$$A_n - C_n = \sum_{k=2}^n A_{k-2}C_{n-k} + \sum_{k=3}^n \sum_{l=3}^k (2 - \delta_{l,3}) A_{k-l}C_{n-k}.$$

The identity then follows from Theorem 3.2. □

Lemma 4.8. *If T_n is the number of ways to tile an n -board using half-squares and $(\frac{1}{2}, \frac{1}{2})$ -fences such that no even-length metatiles occur then*

$$T_n = T_{n-1} + T_{n-2} + T_{n-3}, \quad n \geq 3, \tag{4.9}$$

where $T_0 = T_1 = T_2 = 1$.

Proof. Condition on the last metatile of which there is only one of length 1 but two of every other odd length. Thus

$$T_n = \delta_{0,n} + T_{n-1} + 2 \sum_{j=1}^{\lfloor (n-1)/2 \rfloor} T_{n-1-2j} \quad (4.10)$$

with $T_{n<0} = 0$. Subtracting (4.10) with n replaced by $n - 2$ from (4.10) gives $T_n = \delta_{0,n} - \delta_{2,n} + T_{n-1} + T_{n-2} + T_{n-3}$ which is equivalent to (4.9). \square

$T_{n \geq 0} = 1, 1, 1, 3, 5, 9, 17, 31, 57, 105, \dots$ is a Tribonacci sequence (A000213 in [6]).

Identity 4.9.

$$F_{n+1}^2 = T_n + \sum_{k=2}^n \sum_{j=1}^{\lfloor k/2 \rfloor} (2 + \delta_{j,1}) F_{k-2j+1}^2 T_{n-k}. \quad (4.11)$$

Proof. How many tilings of an n -board contain at least one even-length metatile? Answer 1: $A_n - T_n$. Answer 2: condition on the location of the last even-length metatile. The metatile can occupy cells $k - 2j + 1$ to k ($k = 2j, \dots, n$) where $2j$ is the length of the metatile. The number of tilings for a metatile of length $2j$ for a given k is $A_{k-2j} T_{n-k}$. There are three metatiles of length 2 and two for each $j > 1$. Summing over the possible j and k and equating the two answers gives

$$A_n - T_n = \sum_{k=2}^n \sum_{j=1}^{\lfloor k/2 \rfloor} (2 + \delta_{j,1}) A_{k-2j} T_{n-k}.$$

The identity then follows from Theorem 3.2. \square

Identity 4.10.

$$F_{n+1}^2 = 3F_n^2 - F_{n-1}^2 + 2(-1)^n. \quad (4.12)$$

Proof. We find a near bijection between the tilings of an n -board and an $(n-2)$ -board and the tilings of three $(n-1)$ -boards. There is an exact bijection between the first $(n-1)$ -board and tilings of an n -board that end in h^2 . There is a bijection \mathcal{B}_n between the tilings of the n -board that end in a fence and the tilings of the second $(n-1)$ -board if both boards contain at least one h (i.e., neither is an all-bifence tiling). If the n -board ends in a bifence, find the final h . If the final h is captured, replace the filled fence it is in by h . Otherwise replace the free h and the bifence to the right of it by a filled fence. This gives all h -containing $(n-1)$ -board tilings ending with a fence. The tilings ending in h are obtained from the n -board tilings ending in a filled fence by replacing that filled fence by an h . This leaves the tilings of the n -board that end in a free h (but not h^2). Not counting this final h , find the final h in the tiling (as there must be at least one other h) and then obtain the corresponding $(n-1)$ -board by using the same procedure as for n -boards ending in a fence. This generates all tilings of the third $(n-1)$ -board ending in a free h . The bijection between the remaining h -containing tilings of this board (i.e., those ending in a fence) and the all h -containing tilings of an $(n-2)$ -board is simply \mathcal{B}_{n-1} . When n is even, the n and $(n-2)$ -boards both have an all bifence tiling and so $A_n + A_{n-2} = 3A_{n-1} + 2$. When n is odd, the second and third $(n-1)$ -boards have all bifence tilings which do not correspond to any of the h -containing tilings of the n or $(n-2)$ -boards and so we must subtract 2. Thus overall,

$$A_n + A_{n-2} = 3A_{n-1} + 2(-1)^n,$$

and the required identity is obtained from using Theorem 3.2. \square

5. DISCUSSION

Clearly, the tiling given here has the same combinatorial interpretation as tiling an even-length board with squares and $(1, 1)$ -fences. However, in that expanded form of tiling, there are only three metatiles (the square, filled fence, and bifence) and so the identities we obtain here would not arise so naturally. The number of tilings of an odd-length board in this case gives the golden rectangle numbers and so identities relating these to the Fibonacci numbers squared can also be obtained via combinatorial proof.

It should be noted that the occurrence of an infinite number of metatiles is not limited to tiling with fences. For example, if one tiles an n -board with half-squares and squares (S) then the metatiles are $hS^j h$ for $j \geq 0$ (and the number of such tilings is F_{2n+1} since it is equivalent to tiling an even length board with squares and dominoes).

REFERENCES

- [1] A. T. Benjamin, C. R. H. Hanusa, and F. E. Su, *Linear recurrences through tilings and Markov chains*, *Utilitas Mathematica*, **64** (2003), 3–17.
- [2] A. T. Benjamin and J. J. Quinn, *Proofs That Really Count: The Art of Combinatorial Proof*, Mathematical Association of America, Washington, 2003.
- [3] R. C. Brigham, R. M. Caron, P. Z. Chinn, and R. P. Grimaldi, *A tiling scheme for the Fibonacci numbers*, *Journal of Recreational Mathematics*, **28** (1996), 10–16.
- [4] K. Edwards, *A Pascal-like triangle related to the Tribonacci numbers*, *Fibonacci Quarterly*, **46/47** (2008/2009), 18–25.
- [5] K. Edwards and M. A. Allen, *Strongly restricted permutations and tiling with fences*, *Discrete Applied Mathematics*, **187** (2015), 82–90.
- [6] N. J. A. Sloane, *The Online Encyclopedia of Integer Sequences*, Published electronically at <http://oeis.org>, 2010.

MSC2010: 05A19, 11B39

PHYSICS DEPARTMENT, FACULTY OF SCIENCE, MAHIDOL UNIVERSITY, RAMA 6 ROAD, BANGKOK 10400 THAILAND

E-mail address: `kenneth.edw@mahidol.ac.th`

E-mail address: `maa5652@gmail.com` (*corresponding author)