

SCHRÖDER COLORING AND APPLICATIONS

DANIEL BIRMAJER, JUAN B. GIL, JUAN D. GIL, AND MICHAEL D. WEINER

ABSTRACT. We present several bijections, in terms of combinatorial objects counted by the Schröder numbers, that are then used (via coloring) for the construction and enumeration of rational Schröder paths with integer slope, ordered rooted trees, and simple rooted outerplanar maps. On the other hand, we derive partial Bell polynomial identities for the little and large Schröder numbers, which allow us to obtain explicit enumeration formulas.

1. INTRODUCTION

This paper focuses on the little Schröder numbers

$$1, 1, 3, 11, 45, 197, 903, 4279, 20793, 103049, \dots, \quad (1.1)$$

and the role they play for coloring certain combinatorial structures. Consistent with the OEIS, we denote the above sequence by $(s_n)_{n \in \mathbb{N}_0}$. These numbers appeared in a paper by Ernst Schröder [7] from 1870, where he discusses four combinatorial problems regarding parenthesizations. Problem II in [7] lead to (1.1), and it is now well known that $(s_n)_{n \in \mathbb{N}_0}$ enumerates a wide variety of objects, see [6, A001003], [9].

Here we rely on the following three interpretations for s_n with $n \geq 0$:

- (PATH) Number of Schröder paths¹ from $(0, 0)$ to (n, n) with no D-steps on $y = x$.
- (TREE) Number of ordered trees with no vertex of outdegree 1 and having $n + 1$ leaves.
- (POLY) Number of dissections of a convex $(n + 2)$ -gon by nonintersecting diagonals.

Our main computational result is Theorem 2.5, where we give an explicit formula for the evaluation of partial Bell polynomials at s_0, s_1, s_2, \dots . Our proof relies on a combinatorial argument that uses integer compositions and (PATH). In addition, we give a similar formula for when $(s_n)_{n \in \mathbb{N}_0}$ is replaced by the sequence $(r_n)_{n \in \mathbb{N}_0}$ of large Schröder numbers $1, 2, 6, 22, 90, \dots$, see [6, A006318]. These formulas are particularly beneficial when dealing with Bell transformations, as defined in [2].

In Section 3, we consider the set $\mathcal{S}_n(\alpha)$ of rational Schröder paths with slope $\alpha \in \mathbb{N}$, i.e., lattice paths from $(0, 0)$ to $(n, \alpha n)$ with steps E, N, and D, that stay weakly above the line $y = \alpha x$. Our main contribution here is a bijective map that provides an algorithm to generate rational Schröder paths using classical Schröder paths as building blocks. As a consequence, we obtain a formula for the enumeration of $\mathcal{S}_n(\alpha)$ by the number of building blocks used for their construction. In the special case when

¹Lattice paths with steps E = $(1, 0)$, N = $(0, 1)$, and D = $(1, 1)$, staying weakly above $y = x$.

these paths contain the maximal number of blocks of the form D or NE, our formula leads to several sequences listed in the OEIS [6], see Table 1.

Section 4 deals with ordered rooted trees. We consider ordered trees with no vertex of outdegree 1, like in (TREE), and use them to construct ordered rooted trees with a given number of generators.² This is achieved by means of a Dyck path insertion algorithm described in the proof of Theorem 4.1. As a corollary, we obtain a formula for the number of ordered rooted trees with n generators having a prescribed number of nodes of outdegree 1.

Finally, in Section 5, we make a brief excursion into the enumeration of outerplanar maps, cf. [3, 4]. We view polygon dissections as biconnected simple outerplanar maps and use (POLY), together with certain colored Dyck paths, to generate and count simple outerplanar maps by the number of their biconnected components.

All of our proofs rely on explicit combinatorial bijections that involve building blocks counted by the Schröder numbers. The three applications we discussed in this paper represent just a glimpse of the possible ways Schröder-type objects may be used to construct and enumerate more complex combinatorial classes. For example, s_n also counts indecomposable permutations that avoid the patterns 2413 and 3142, and for $n \geq 1$, s_n gives the number of increasing tableaux of shape (n, n) , cf. [6].

In summary, the content of Section 2 provides a toolbox for studying a broad family of sequence transformations of the little and large Schröder numbers.

2. BELL TRANSFORMS OF LITTLE SCHRÖDER NUMBERS

In [2], three of the authors introduced the following family of sequence transformations defined via partial Bell polynomials. Let a, b, c, d be fixed numbers. Given $x = (x_n)_{n \in \mathbb{N}}$, its Bell transform $y = \mathcal{Y}_{a,b,c,d}(x)$ is the sequence defined by

$$y_n = \sum_{k=1}^n \frac{1}{n!} \left[\prod_{j=1}^{k-1} (an + bk + cj + d) \right] B_{n,k}(1!x_1, 2!x_2, \dots) \text{ for } n \geq 1, \quad (2.1)$$

where $B_{n,k}$ denotes the (n, k) -th (exponential) partial Bell polynomial.

For $0 \leq k \leq n$, we have

$$B_{n,k}(z_1, \dots, z_{n-k+1}) = \sum_{\alpha \in \pi(n,k)} \frac{n!}{\alpha_1! \alpha_2! \dots} \left(\frac{z_1}{1!} \right)^{\alpha_1} \left(\frac{z_2}{2!} \right)^{\alpha_2} \dots,$$

where $\pi(n, k)$ denotes the set of multi-indices $\alpha \in \mathbb{N}_0^{n-k+1}$ such that

$$\alpha_1 + \alpha_2 + \dots = k \text{ and } \alpha_1 + 2\alpha_2 + 3\alpha_3 + \dots = n.$$

The polynomial $B_{n,k}$ contains as many monomials as the number of partitions of $[n] = \{1, \dots, n\}$ into k parts.

The definition of $\mathcal{Y}_{a,b,c,d}(x)$ suggests the need for finding a closed formula for $B_{n,k}(1!x_1, 2!x_2, \dots)$, which in most cases is a challenging task. In this section, we

²A *generator* is a leaf or a node with only one child.


will use a combinatorial argument (via colored compositions) to derive a formula for $B_{n,k}(1!s_0, 2!s_1, \dots)$, where $(s_n)_{n \in \mathbb{N}_0}$ is the sequence of little Schröder numbers.

Let $\mathcal{CS}_{n,k}$ be the set of compositions of n with exactly k parts such that part j can take on s_{j-1} colors. Using a formula³ by Hoggatt and Lind [5], we get

$$\#\mathcal{CS}_{n,k} = \frac{k!}{n!} B_{n,k}(1!s_0, 2!s_1, 3!s_2, \dots). \tag{2.2}$$

Lemma 2.3. *There is a bijection between the elements of $\mathcal{CS}_{n+k,k}$ and the set of Schröder paths from $(0, 0)$ to $(n + k - 1, n + k - 1)$ with exactly $k - 1$ diagonal steps on the line $y = x$.*

Proof. Given a Schröder path P from $(0, 0)$ to $(n + k - 1, n + k - 1)$ with $k - 1$ diagonal steps on the line $y = x$, use these diagonal steps to split the paths into k Schröder subpaths P_1, P_2, \dots, P_k with no D-steps on the line $y = x$, including the trivial path of length 0 (i.e., a single lattice point). For example, for $n = 5$ and $k = 4$, we consider the path



and use the three diagonal steps (in orange) to split the path into the four subpaths in black. The first path covers a segment on $y = x$ with 3 lattice points, and the third path covers a segment with 4 lattice points. The second and fourth paths are trivial. With the above path, we then associate the composition $9 = 3 + 1 + 4 + 1$, where part 3 is labeled by the path **NDE** and part 4 is labeled by the path **NNNEEE**.

In general, if P_j covers a segment on $y = x$ with i_j lattice points, we let $\varphi(P_j) = i_j$ and define $\varphi(P)$ to be the composition

$$n + k = i_1 + i_2 + \dots + i_k,$$

where part i_j is labeled by P_j . The map φ is clearly bijective. □

As a consequence of Lemma 2.3, (PATH), and identity (2.2), we get

$$\frac{k!}{(n + k)!} B_{n+k,k}(1!s_0, 2!s_1, \dots) = \sum_{m_1 + \dots + m_k = n} s_{m_1} \cdots s_{m_k}. \tag{2.4}$$

³Written here in terms of Bell polynomials.

Theorem 2.5. *For $1 \leq k < n$, we have*

$$B_{n,k}(1!s_0, 2!s_1, 3!s_2, \dots) = \frac{n!}{(k-1)!} \sum_{j=1}^{n-k} \frac{1}{j} \binom{n-k-1}{j-1} \binom{n+j-1}{j-1}. \quad (2.6)$$

Proof. First of all, note that $s_0 = 1$, and for $n \geq 1$,

$$\begin{aligned} s_n &= \sum_{j=1}^n \frac{1}{j} \binom{n+j}{j-1} \binom{n-1}{j-1} \\ &= \sum_{j=1}^n \binom{n+j}{j-1} \frac{(j-1)! n!}{n! j!} \binom{n-1}{j-1} \\ &= \sum_{j=1}^n \binom{n+j}{j-1} \frac{(j-1)!}{n!} B_{n,j}(1!, 2!, \dots). \end{aligned}$$

Therefore, by a convolution formula given in [2, Section 4],

$$\begin{aligned} \sum_{m_1+\dots+m_k=n} s_{m_1} \cdots s_{m_k} &= k \sum_{j=1}^n \binom{n+j+k-1}{j-1} \frac{(j-1)!}{n!} B_{n,j}(1!, 2!, \dots) \\ &= \sum_{j=1}^n \frac{k}{j} \binom{n-1}{j-1} \binom{n+j+k-1}{j-1}. \end{aligned}$$

Now, identity (2.4) implies

$$\frac{k!}{(n+k)!} B_{n+k,k}(1!s_0, 2!s_1, \dots) = \sum_{j=1}^n \frac{k}{j} \binom{n-1}{j-1} \binom{n+j+k-1}{j-1},$$

and (2.6) follows by replacing n by $n-k$. \square

Using basic properties of the partial Bell polynomials and (2.6), we get:

Corollary 2.7. *If $(r_n)_{n \in \mathbb{N}_0}$ is the sequence of large Schröder numbers $1, 2, 6, 22, \dots$, then for $1 \leq k < n$, we have*

$$\begin{aligned} &B_{n,k}(1!r_0, 2!r_1, 3!r_2, \dots) \\ &= \sum_{\ell=0}^k (-1)^\ell 2^{k-\ell} \binom{n}{\ell} B_{n-\ell, k-\ell}(1!s_0, 2!s_1, 3!s_2, \dots) \\ &= \frac{n!}{(k-1)!} \sum_{\ell=0}^k \sum_{j=1}^{n-k} \frac{(-1)^\ell 2^{k-\ell}}{j} \binom{k-1}{\ell} \binom{n-k-1}{j-1} \binom{n-\ell+j-1}{j-1}. \end{aligned}$$

It is worth noting that some of the most prominent Bell transforms are of the form $\mathcal{Y}_{a,b,-1,1}(x)$ with $a, b \in \mathbb{N}_0$. In such a special case, if $s = (s_n)_{n \in \mathbb{N}_0}$ and $y = \mathcal{Y}_{a,b,-1,1}(s)$, then (2.1) becomes

$$y_n = \sum_{k=1}^n \binom{an + bk}{k-1} \frac{(k-1)!}{n!} B_{n,k}(1!s_0, 2!s_1, \dots).$$

We can then use (2.6) to conclude:

Corollary 2.8. *If $s = (s_n)_{n \in \mathbb{N}_0}$, then the Bell transform $\mathcal{Y}_{a,b,-1,1}(s)$ is given by*

$$y_n = \frac{1}{n} \binom{(a+b)n}{n-1} + \sum_{k=1}^{n-1} \sum_{j=1}^{n-k} \frac{1}{j} \binom{an + bk}{k-1} \binom{n-k-1}{j-1} \binom{n+j-1}{j-1}.$$

3. RATIONAL SCHRÖDER PATHS

For $a \in \mathbb{N}$ and $\mathbf{c} = (c_1, c_2, \dots)$ with $c_j \in \mathbb{N}_0$, let us consider the set of Dyck words of semilength $(a+1)n$ created from strings of the form $P_0 = "d"$ and $P_j = "u^{(a+1)j}d^j"$ for $j = 1, \dots, n$, such that each maximal $(a+1)j$ -ascent substring $u^{(a+1)j}$ may be colored in c_j different ways. Consistent with the notation used in [1], we denote this set by $\mathcal{D}_n^{\mathbf{c}}(a, 1)$.

Let \mathcal{SP}_n^{\wedge} denote the set of Schröder paths from $(0, 0)$ to (n, n) that lie *strictly* above the diagonal $y = x$ except for the diagonal path from $(0, 0)$ to $(1, 1)$. Moreover, let σ be the sequence that enumerates \mathcal{SP}_n^{\wedge} . In particular, $\sigma_1 = 2$ since the two paths D and NE from $(0, 0)$ to $(1, 1)$ both fit the above description, and $\sigma_2, \sigma_3, \sigma_4, \dots$, is precisely the sequence of large Schröder numbers 2, 6, 22, 90, 394, \dots , see [6, A006318]. Note that $\sigma_j = 2s_{j-1}$ for $j \geq 1$.

Recall that $\mathcal{S}_n(\alpha)$ denotes the set of rational Schröder paths with slope $\alpha \in \mathbb{N}$.

Theorem 3.1. *For every $n \in \mathbb{N}$, we have*

$$\#\mathcal{S}_n(\alpha) = \#\mathcal{D}_n^{\sigma}(\alpha - 1, 1).$$

Proof. We will give an explicit bijective map $\xi : \mathcal{S}_n(\alpha) \rightarrow \mathcal{D}_n^{\sigma}(\alpha - 1, 1)$.

Given a rational Schröder path $p \in \mathcal{S}_n(\alpha)$, walk the path from $(n, \alpha n)$ to $(0, 0)$ and construct a colored Dyck path $q \in \mathcal{D}_n^{\sigma}(\alpha - 1, 1)$ as follows:

- if at a point (a, b) on p followed by a D-step, add an α -ascent followed by one down-step to the path q , color the ascent by $\color{blue}{\nearrow}$, and move to the point $(a-1, b-1)$ on p .
- if at a point (a, b) followed by an E-step, draw a line of slope 1 from that point to the next point of the form $(a-j, b-j)$ where the line intersects the path p . The path p_j between these two points is a Schröder path in \mathcal{SP}_j^{\wedge} . Add to the path q an αj -ascent followed by j down-steps, color that ascent by p_j , and move to the point $(a-j, b-j)$ on p .

- if at a point (a, b) followed by an N-step, add a down-step to the path q and move to the point $(a, b - 1)$ on p .

The process stops when $(0, 0)$ has been reached. The path $q = \xi(p)$ created by means of the above algorithm is by construction a Dyck path in $\mathfrak{D}_n^\sigma(\alpha - 1, 1)$. Note that a return to the line $y = \alpha x$ on p corresponds to a return to the x -axis on the Dyck path. For an example, see Figure 1.




FIGURE 1. Example for $\alpha = 2$ and $n = 4$.

The map ξ is reversible. Given a Dyck path $q \in \mathfrak{D}_n^\sigma(\alpha - 1, 1)$, construct a path $p \in \mathcal{S}_n(\alpha)$ as follows. Walk the Dyck path q from right to left. For every down-step that is not part of a block $P_j = u^{\alpha j} d^j$ on q , add an N-step to p , and for every block P_j , add to p the Schröder path labeling the corresponding αj -ascent on P_j . \square

The above bijection provides an algorithm to generate rational Schröder paths in $\mathcal{S}_n(\alpha)$ using classical Schröder paths as building blocks.

By [1, Theorem 2], the number of elements in $\mathfrak{D}_n^\sigma(\alpha - 1, 1)$ having exactly k peaks is given by

$$\begin{aligned} & \binom{(\alpha - 1)n + k}{k - 1} \frac{(k - 1)!}{n!} B_{n,k}(1!\sigma_1, 2!\sigma_2, \dots) \\ &= \binom{(\alpha - 1)n + k}{k - 1} \frac{(k - 1)!}{n!} B_{n,k}(1!(2s_0), 2!(2s_1), \dots) \\ &= \binom{(\alpha - 1)n + k}{k - 1} \frac{(k - 1)!}{n!} 2^k B_{n,k}(1!s_0, 2!s_1, \dots). \end{aligned}$$

Corollary 3.2. *There are*

$$\frac{2^n}{n} \binom{\alpha n}{n - 1} = \frac{2^n}{(\alpha - 1)n + 1} \binom{\alpha n}{n} \quad (3.3)$$

paths in $\mathcal{S}_n(\alpha)$ containing n paths from \mathcal{SP}_1^\wedge (i.e., paths of the form D or NE).

Table 1 shows a few terms of the sequence (3.3) for $\alpha = 1, \dots, 5$.

Slope α	$\frac{2^n}{(\alpha-1)n+1} \binom{\alpha n}{n}$	OEIS
1	2, 4, 8, 16, 32, 64, 128, 256, 512, 1024, ...	A000079
2	2, 8, 40, 224, 1344, 8448, 54912, 366080, ...	A151374
3	2, 12, 96, 880, 8736, 91392, 992256, 11075328, ...	A153231
4	2, 16, 176, 2240, 31008, 453376, 6888960, 107707392, ...	A217360
5	2, 20, 280, 4560, 80960, 1520064, 29680640, 596593920, ...	A217364

TABLE 1. Number of paths in $\mathcal{S}_n(\alpha)$ having n subpaths from \mathcal{SP}_1^\wedge .

Corollary 3.4. *If $1 \leq k < n$, the number of paths in $\mathcal{S}_n(\alpha)$ that can be built with exactly k elements of $\mathcal{SP}^\wedge = \bigcup_n \mathcal{SP}_n^\wedge$ is given by*

$$2^k \binom{(\alpha-1)n+k}{k-1} \sum_{j=1}^{n-k} \frac{1}{j} \binom{n-k-1}{j-1} \binom{n+j-1}{j-1}.$$

Finally, for the cardinality of $\mathcal{S}_n(\alpha)$, we can use [8, Theorem 2.9] to deduce the simple formula

$$\#\mathcal{S}_n(\alpha) = \frac{1}{\alpha n + 1} \sum_{\ell=0}^n \binom{\alpha n + 1}{n - \ell} \binom{\alpha n + \ell}{\ell}.$$

4. ORDERED ROOTED TREES

In this section, we use the interpretation (TREE) to generate and count ordered rooted trees with n generators (leaves or nodes of outdegree 1).

Let \mathcal{ST}_n be the set of ordered trees with no vertex of outdegree 1 and having $n + 1$ leaves. Thus $\#\mathcal{ST}_n = s_n$. Note that every tree in \mathcal{ST}_n has $n + 1$ generators.

Let \mathcal{T}_n be the set of ordered rooted trees with n generators. Our next result shows that every element of $\mathcal{T} = \bigcup_n \mathcal{T}_n$ can be uniquely constructed from elements of the set $\mathcal{ST} = \bigcup_n \mathcal{ST}_n$ via a Dyck path insertion algorithm.

Theorem 4.1. *The set of ordered rooted trees with n generators is in one-to-one correspondence with the set of Dyck paths of semilength n whose ascents of length j , for $j = 1, \dots, n$, may be colored in s_{j-1} different ways.*

Proof. Let $\mathfrak{D}_n^s(1, 0)$ be the set of Dyck paths of semilength n colored by the little Schröder numbers. Because of (TREE), we can consider $\mathfrak{D}_n^s(1, 0)$ to be the set of Dyck paths of semilength n with each j -ascent associated with a tree in \mathcal{ST}_{j-1} .

We will define a map $\psi : \mathfrak{D}_n^s(1, 0) \rightarrow \mathcal{T}_n$ that requires a notion of traversing an ordered tree. When traversing a tree, we stick to the ordering of leaf nodes that would be acquired when performing inorder depth-first search.




TABLE 2. Elements of \mathcal{ST}_n for $n = 0, 1, 2, 3$.

Suppose we have a path $P \in \mathfrak{D}_n^s(1, 0)$ with k peaks. For $1 \leq i \leq k$, let G_i denote the ordered tree associated with the i th ascent of the Dyck path. Starting at the first ascent of P , we construct the ordered tree $\psi(P)$ as follows. We start at the root of the tree G_1 . Suppose that the first ascent of the Dyck path is followed by a j_1 -descent. Then, we traverse G_1 from the root until we reach the j_1 th leaf. At that point, we add an edge between the j_1 th leaf of G_1 and the root of the tree G_2 corresponding to the second ascent. Suppose that the second ascent is followed by a j_2 -descent. From the node that was formerly the j_1 th leaf from the root (and which is no longer a leaf), we continue to traverse the augmented tree until we reach the j_2 th leaf from that node.

This algorithm will always result in an ordered tree with n generators. Note that since P has n up steps, and since every j -ascent corresponds to an ordered tree with j leaves (and no other generators), there are n leaves total in the trees G_1, \dots, G_k . Every leaf in a tree G_i either becomes a leaf in the final tree, or it becomes a node with a single child. Thus the final tree $\psi(P)$ has exactly n generators.

An example that illustrates the above construction is given in Figure 2.




FIGURE 2. Example for $n = 7$.

The inverse map from \mathcal{T}_n to $\mathfrak{D}_n^s(1, 0)$ is straightforward. Given $G \in \mathcal{T}_n$, we construct a Schröder colored Dyck path as follows. Starting at the root of G , we first identify the largest subgraph of G with same root that is an ordered tree with no vertex of outdegree 1. We call this subgraph G_1 . Once G_1 has been specified, we count the number of leaves of G_1 . If G_1 has j leaves, then we begin the Dyck path with a j -ascent and label that first ascent with G_1 . Then, we traverse G_1 until we find a leaf of G_1 that corresponds to a node of outdegree 1 in G . If this leaf is the j_1 th leaf checked, then we add a j_1 -descent to the Dyck path. Then, to find the next ascent, we identify the ordered tree G_2 rooted at the child of the node corresponding to the j_1 th leaf of G_1 . This process continues until the tree G has been completely traversed.

This algorithm gives a Dyck path in $\mathfrak{D}_n^s(1, 0)$. Indeed, every generator in G is counted as both an ascent and a descent, so the path $\psi^{-1}(G)$ has semilength n . Moreover, every j -ascent corresponds to one of the s_{j-1} possible ordered trees with j leaves and no nodes of outdegree 1. Finally, note that the ascent corresponding to a generator of the ordered tree is always added to the path before the descent corresponding to the same generator is added to the path. \square

Corollary 4.2. *As a consequence of [1, Theorem 2] and Theorem 4.1, we get*

$$\#\mathcal{T}_n = \#\mathfrak{D}_n^s(1, 0) = 1 + \sum_{k=1}^{n-1} \sum_{j=1}^{n-k} \frac{1}{j} \binom{n}{k-1} \binom{n-k-1}{j-1} \binom{n+j-1}{j-1}.$$

This gives the sequence 1, 2, 7, 32, 166, 926, 5419, 32816, ..., see [6, A108524].

Moreover, the number of trees in \mathcal{T}_n having exactly $k-1$ nodes of outdegree 1 is equal to the number of paths in $\mathfrak{D}_n^s(1, 0)$ having k peaks. This number is given by

$$\binom{n}{k-1} \frac{(k-1)!}{n!} B_{n,k}(1!s_0, 2!s_1, \dots),$$

which for $1 \leq k < n$ can be written as

$$\binom{n}{k-1} \sum_{j=1}^{n-k} \frac{1}{j} \binom{n-k-1}{j-1} \binom{n+j-1}{j-1}.$$


5. OUTERPLANAR MAPS

Recall that if \mathbf{s} is the sequence $(s_n)_{n \in \mathbb{N}_0}$, $\mathfrak{D}_n^s(1, 1)$ denotes the set of Dyck paths of semilength $2n$ created from down-steps and blocks of the form $P_j = "u^{2j}d^j"$ for $j = 1, \dots, n$, such that each $2j$ -ascent may be colored in s_{j-1} different ways.

In this section, we use the interpretation (POLY) for the little Schröder numbers to generate and count simple outerplanar maps using polygon dissections (seen as biconnected outerplanar maps) put together according to the elements of $\mathfrak{D}_n^s(1, 1)$.

Let \mathcal{B}_n be the set of biconnected (nonseparable) rooted outerplanar maps with $n+2$ vertices. Note that \mathcal{B}_0 consists of the single map $\circ \text{---} \circ$, and for $n \geq 1$, every element of \mathcal{B}_n can be seen as a dissection of a convex $(n+2)$ -gon by nonintersecting diagonals.

Therefore, $\#\mathcal{B}_n = s_n$. For example, \mathcal{B}_1 consists of a single triangle and \mathcal{B}_2 contains the three possible dissections of a square:



By default, we will assume that the elements of \mathcal{B}_n are rooted at the base of the polygon such that the root face is the outer face.

Theorem 5.1. *The set \mathcal{M}_n of simple outerplanar maps with $n + 1$ vertices is in one-to-one correspondence with the set $\mathfrak{D}_n^s(1, 1)$.*

Proof. We will establish a bijective map $\phi : \mathcal{M}_n \rightarrow \mathfrak{D}_n^s(1, 1)$ that relies on the decomposition of a planar map into biconnected components. Given a map $M \in \mathcal{M}_n$ with k biconnected components, we construct a Dyck path $P = \phi(M)$ as follows. Starting at the root vertex v_0 of M , we walk counterclockwise around M , visiting all of its vertices. We let B_1, \dots, B_k be the biconnected components of M , numbered in the order they are visited, and assume that B_j has $i_j + 1$ vertices, $1 \leq i_j \leq n$ for every j . While at v_0 , add a $2i_1$ -ascent to P followed by i_1 down-steps and walk to the next vertex of M (second vertex of the root edge). For each subsequent vertex we visit on B_1 , add a down-step to P until we reach the first vertex on B_2 . At that point, we add a $2i_2$ -ascent to P followed by i_2 down-steps, and move to the next vertex. We then continue adding down-steps to P until we reach the first vertex of the next component of M . When visiting a vertex of a biconnected component that has already been visited, we add a down-step to P . This process ends after we have visited all vertices of M and reach v_0 again. We then label/color the k maximal ascents of P with the corresponding biconnected maps B_1, \dots, B_k , and obtain an element of $\mathfrak{D}_n^s(1, 1)$ with k peaks. An example of the above construction is shown in Figure 3.




FIGURE 3. Example for $n = 6$ and $k = 3$.

The inverse map is clear. Given a Dyck path $P \in \mathfrak{D}_n^s(1, 1)$ with k maximal ascents of lengths $2i_1, \dots, 2i_k$, colored by biconnected rooted maps B_1, \dots, B_k , we construct an outerplanar map as follows. Let $M_1 = B_1$ and let v_0 be the root vertex of B_1 . If the first ascent of P is followed by j_1 down-steps, we walk around M_1 (starting at v_0) until we reach the $(j_1 - i_1 + 1)$ th vertex on M_1 . We call that vertex v_1 and merge

it with the root vertex of B_2 to create a rooted outerplanar map M_2 with the same root edge as M_1 . This extended map has now $i_1 + i_2 + 1$ vertices. We then repeat the process starting now at v_1 . After all coloring biconnected maps have been used as extensions (traversing P from left to right), the resulting rooted outerplanar map M_k will be an element of \mathcal{M}_n . We let $\phi^{-1}(P) = M_k$. \square

Corollary 5.2. *By Theorem 5.1, the number of outerplanar maps in \mathcal{M}_n having exactly k biconnected components is equal to the number of paths in $\mathfrak{D}_n^s(1, 1)$ having k peaks. Thus, by [1, Theorem 2], this number is given by*

$$\binom{n+k}{k-1} \frac{(k-1)!}{n!} B_{n,k}(1!s_0, 2!s_1, \dots), \tag{5.3}$$

which for $1 \leq k < n$ can be written as

$$\binom{n+k}{k-1} \sum_{j=1}^{n-k} \frac{1}{j} \binom{n-k-1}{j-1} \binom{n+j-1}{j-1}.$$

Note that if $k = n$, then (5.3) becomes $\frac{1}{n} \binom{2n}{n-1} = C_n$, and we recover the known fact that planted trees are counted by the Catalan numbers.

Corollary 5.4. *For $n \geq 1$ we have*

$$\#\mathcal{M}_n = \frac{1}{n} \binom{2n}{n-1} + \sum_{k=1}^{n-1} \sum_{j=1}^{n-k} \frac{1}{j} \binom{n+k}{k-1} \binom{n-k-1}{j-1} \binom{n+j-1}{j-1}.$$

This gives the sequence 1, 3, 13, 67, 381, 2307, 14589, 95235, ..., see [6, A064062].

REFERENCES

- [1] D. Birmajer, J. B. Gil, P. R. W. McNamara, and M. D. Weiner, Enumeration of colored Dyck paths via partial Bell polynomials, *Lattice path combinatorics and applications*, 155–165, Dev. Math. **58**, Springer, Cham, 2019.
- [2] D. Birmajer, J. B. Gil, and M. D. Weiner, A family of Bell transformations, *Discrete Math.* **342** (2019), no. 1, 38–54.
- [3] N. Bonichon, C. Gavoille, and N. Hanusse, Canonical decomposition of outerplanar maps and application to enumeration, coding and generation, *J. Graph Algorithms Appl.* **9** (2005), no. 2, 185–204.
- [4] I. Geffner and M. Noy, Counting outerplanar maps, *Electron. J. Combin.* **24** (2) (2017), #P2.3.
- [5] V. E. Hoggatt and D. A. Lind, Fibonacci and binomial properties of weighted compositions, *J. Combin. Theory* **4** (1968), 121–124.
- [6] *The On-Line Encyclopedia of Integer Sequences*, published electronically at <https://oeis.org>, 2019.
- [7] E. Schröder, Vier kombinatorische Probleme, *Z. Math. Phys.* **15** (1870), 361–376.
- [8] J. Schröder, Generalized Schröder numbers and the rotation principle, *J. Integer Seq.* **10** (2007), Article 07.7.7.
- [9] R. P. Stanley, Hipparchus, Plutarch, Schröder, and Hough, *Amer. Math. Monthly* **104** (1997), no. 4, 344–350.

12 DANIEL BIRMAJER, JUAN B. GIL, JUAN D. GIL, AND MICHAEL D. WEINER

NAZARETH COLLEGE, 4245 EAST AVE., ROCHESTER, NY 14618

PENN STATE ALTOONA, 3000 IVYSIDE PARK, ALTOONA, PA 16601

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139

PENN STATE ALTOONA, 3000 IVYSIDE PARK, ALTOONA, PA 16601