The characteristic masses of Niemeier lattices

Gaëtan Chenevier*

February 11, 2020

Abstract

Let L be an integral lattice in the Euclidean space \mathbb{R}^n and W an irreducible representation of the orthogonal group of \mathbb{R}^n . We give an implemented algorithm computing the dimension of the subspace of invariants in W under the isometry group O(L) of L. A key step is the determination of the number of elements in O(L) having any given characteristic polynomial, a datum that we call the *characteristic masses* of L. As an application, we determine the characteristic masses of all the Niemeier lattices, and more generally of any even lattice of determinant ≤ 2 in dimension $n \leq 25$.

For Niemeier lattices, as a verification, we provide an alternative (human) computation of the characteristic masses. The main ingredient is the determination, for each Niemeier lattice L with non-empty root system R, of the G(R)-conjugacy classes of the elements of the "umbral" subgroup O(L)/W(R) of G(R), where G(R) is the automorphism group of the Dynkin diagram of R, and W(R) its Weyl group.

These results have consequences for the study of the spaces of automorphic forms of the definite orthogonal groups in n variables over \mathbb{Q} . As an example, we provide concrete dimension formulas in the level 1 case, as a function of the weight W, up to dimension n = 25.

1. Introduction

1.1. A motivation: dimension of spaces of level 1 automorphic forms for O_n

Let n be an integer $\equiv 0 \mod 8$ and let \mathcal{L}_n be the set of all *even unimodular* lattices in the standard Euclidean space \mathbb{R}^n . A standard example of an

^{*}Université Paris-Saclay, CNRS, Laboratoire de mathématiques d'Orsay, 91405, Orsay, France. During this work, the author has been supported by the C.N.R.S. and by the projects ANR-14-CE25-0002-01 PERCOLATOR and ANR-19-CE40-0015-02 COLOSS.

element of \mathcal{L}_n is the lattice $\mathbf{E}_n = \mathbf{D}_n + \mathbb{Z}_2^1(1, 1, \dots, 1)$, where \mathbf{D}_n denotes the subgroup of elements (x_i) in \mathbb{Z}^n with $\sum_i x_i \equiv 0 \mod 2$. The orthogonal group $O(\mathbb{R}^n)$ naturally acts on \mathcal{L}_n , with finitely many orbits, and we set

(1.1)
$$X_n \stackrel{\text{def}}{=} O(\mathbb{R}^n) \backslash \mathcal{L}_n$$

Representatives of this set X_n have been determined so far for $n \leq 24$ only: we have $X_8 = \{E_8\}$ (Mordell), $X_{16} = \{E_8 \oplus E_8, E_{16}\}$ (Witt) and $|X_{24}| = 24$ (Niemeier): see [KNE57, NI73, VEN80, CS99]. The elements of \mathcal{L}_{24} , to which we shall refer as the *Niemeier lattices*, will play a major role in this paper.

Similarly, for $n \equiv \pm 1 \mod 8$ we define \mathcal{L}_n as the set of all even lattices with covolume $\sqrt{2}$ in \mathbb{R}^n , as well as X_n by the same Formula (1.1). In this case, representatives of X_n are known up to n = 121, this last (and most complicated) case being due to Borcherds [BOR00], and we have

$$|X_1| = |X_7| = |X_9| = 1$$
, $|X_{15}| = 2$, $|X_{17}| = 4$, $|X_{23}| = 32$ and $|X_{25}| = 121$.

For any $n \equiv -1, 0, 1 \mod 8$, and any complex, finite dimensional, continuous, linear representation W of $O(\mathbb{R}^n)$, we consider the complex vector space of W-valued $O(\mathbb{R}^n)$ -equivariant functions on \mathcal{L}_n :

(1.2)
$$M_W(O_n) = \{ f : \mathcal{L}_n \longrightarrow W \mid f(gL) = gf(L) \ \forall L \in \mathcal{L}_n, \ \forall g \in O(\mathbb{R}^n) \}.$$

This space has a natural interpretation as a space of *level* 1 and *weight* W automorphic forms for the orthogonal group scheme O_n of any element of \mathcal{L}_n . In particular, it has a very interesting action of the Hecke ring of O_n (see e.g. [CL19, Sect. 4]), which is a first indication of our interest in it.

If L is a lattice in the Euclidean space \mathbb{R}^n , we denote by $O(L) = \{g \in O(\mathbb{R}^n) \mid gL = L\}$ its (finite) isometry group. If W is a representation of $O(\mathbb{R}^n)$, we denote by $W^{\Gamma} = \{w \in W \mid \gamma w = w, \forall \gamma \in \Gamma\} \subset W$ the subspace of invariants of the subgroup Γ of $O(\mathbb{R}^n)$. Fix representatives L_1, \ldots, L_h of the classes in X_n . Then the map $f \mapsto (f(L_i))$ induces a \mathbb{C} -linear isomorphism

(1.3)
$$M_W(\mathcal{O}_n) \xrightarrow{\sim} \prod_{i=1}^h W^{\mathcal{O}(L_i)}.$$

It follows that $M_W(O_n)$ is finite dimensional. Our main aim in this work, which is of computational flavor, is to explain how to compute dim $M_W(O_n)$ for all $n \leq 25$ and W arbitrary. The special cases n = 7, 8, 9 and n = 16, more precisely their SO-variants¹, had been respectively previously considered in

¹We define $M_W(SO_n)$ by replacing $O(\mathbb{R}^n)$ with $SO(\mathbb{R}^n)$ in (1.2), and W with a representation of $SO(\mathbb{R}^n)$. We have then $M_W(SO_n) \simeq M_{W'}(O_n)$ where W' is the representation of $O(\mathbb{R}^n)$ induced from W [CL19, §4.4.4]. The question of computing dimensions in the SO-case is thus a special case of the same question in the O-case (the one considered here).

[CR12, Ch. 2] and in [CL19, Ch. IX Prop. 5.13]. In a different direction, see Appendix B for an asymptotic formula for dim $M_W(O_n)$ (for any n).

Our main motivation for these computations is the relation between the spaces $M_W(O_n)$ and geometric ℓ -adic representations of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ of Artin conductor 1 (or pure motives over \mathbb{Q} with good reduction everywhere) which follows from the general yoga and point of views of Langlands and Arthur on automorphic representations. This circle of ideas has been studied in great details in the recent works [CL19, CR12], and pursued in [TAI17, CT19a], to which we refer to for further explanations. As a start, the reader may consult the preface of [CL19]. Let us simply say here that in a forthcoming work of Taïbi and the author, we shall use the results of the present paper as an ingredient to extend to higher dimensions d, hopefully up to d = 24, the counting of level 1, algebraic, essentially selfdual cuspidal automorphic representations of GL_d over \mathbb{Q} started in the aforementioned works. One alternative motivating goal of these works is to obtain new information on the size of X_{31} and X_{32} (see e.g. [CL19, Thm. IX.6.1] for a direct proof of the equality $|X_{24}| = 24$ not relying on any lattice computation).

1.2. Dimension of invariants and characteristic masses

Consider now an arbitrary integral lattice L in the standard Euclidean space \mathbb{R}^n of arbitrary dimension n, and a finite dimensional representation W of $O(\mathbb{R}^n)$. Motivated by the previous paragraph, we are interested in algorithms to determine the dimension of the subspace $W^{O(L)} \subset W$ of O(L)-invariants in W. Of course, our requirement will be that these algorithms be efficient for the even lattices of determinant ≤ 2 , as in §1.1.

Obviously, we may and do assume that W is irreducible. It will be convenient to parameterize the isomorphism classes of irreducible complex representations of $O(\mathbb{R}^n)$, following Weyl's original approach [WEYL46], by the n-permissible² (integer) partitions: see Appendix A for a brief reminder of this parameterization and its relation with the highest weight theory for $SO(\mathbb{R}^n)$. This parameterization not only allows to deal with the two connected components of $O(\mathbb{R}^n)$ in a very concise way, but it is also especially relevant for the character formulas we shall use.

We denote by W_{λ} an irreducible representation of $O(\mathbb{R}^n)$ associated with the *n*-permissible partition $\lambda = (\lambda_1 \ge \cdots \ge \lambda_n \ge 0)$. The element $-\mathrm{id}_n$ acts on W_{λ} by the sign $(-1)^{|\lambda|}$, with $|\lambda| = \sum_i \lambda_i$, so $W_{\lambda}^{O(L)}$ vanishes for

²This means that the first two columns of the Young diagram of the partition have at most n boxes in total.

 $|\lambda| \equiv 1 \mod 2$. Our starting point is the trivial formula $\dim W_{\lambda}^{O(L)} = \frac{1}{|O(L)|} \sum_{\gamma \in O(L)} \operatorname{Trace}(\gamma; W_{\lambda})$, that we rewrite as

(1.4)
$$\dim \mathbf{W}_{\lambda}^{\mathbf{O}(L)} = \sum_{P \in \operatorname{Car}_{n}} \operatorname{m}_{\mathbf{O}(L)}(P) \operatorname{Trace}(\mathbf{c}_{P}; \mathbf{W}_{\lambda})$$

where:

(i) $\operatorname{Car}_n \subset \mathbb{Z}[t]$ denotes the (finite) subset of polynomials of degree n which are products of cyclotomic polynomials. This subset is³ also the set of characteristic polynomials of the elements of $O(\mathbb{R}^n)$ preserving some lattice in \mathbb{R}^n . Using the irreducibility of cyclotomic polynomials in $\mathbb{Q}[t]$, it is straightforward to enumerate the elements of Car_n for small n with the help of a computer: see Table 1 for the cardinality of Car_n for $n \leq 27$ (sequence A120963 on [OEIS]).

n	1	2	3	4	5	6	7	8	9
$ Car_n $	2	6	10	24	38	78	118	224	330
n	10	11	12	13	14	15	16	17	18
$ Car_n $	584	838	1420	2002	3258	4514	7134	9754	15010
n	19	20	21	22	23	24	25	26	27
$ Car_n $	20266	30532	40798	60280	79762	115966	152170	217962	283754

Table 1: The cardinality of Car_n for $n \leq 27$.

(ii) For any finite subset $S \subset O(\mathbb{R}^n)$, and any P in $\mathbb{R}[t]$, we denote by $m_S(P)$ the number of elements g in S with $\det(t \operatorname{id}_n - g) = P$, **divided by** |S|. This is an element of $\mathbb{Q}_{>0}$ that we call the mass of P in S. By definition, we have

$$\sum_{P \in \mathbb{R}[t]} \mathbf{m}_S(P) = 1.$$

(iii) For P in $\mathbb{R}[t]$ a monic polynomial of degree n whose complex roots are on the unit circle (e.g. $P \in \operatorname{Car}_n$), we denote by $c_P \subset O(\mathbb{R}^n)$ the unique conjugacy class whose characteristic polynomial is P.

We now discuss the problem of evaluating Formula (1.4). The main unknown, which contains all the required information about L and which does

³Set $\zeta = e^{\frac{2i\pi}{m}}$ for $m \ge 1$. The symmetric bilinear form $(x, y) \mapsto \text{Trace}_{\mathbb{Q}(\zeta)/\mathbb{Q}}(x\overline{y})$ on the free abelian group $L = \mathbb{Z}[\zeta]$ defines a inner product on $L \otimes \mathbb{R}$. The multiplication by ζ is an isometry preserving L, with characteristic polynomial the *m*-th cyclotomic polynomial.

not depend on λ , is of course the collection of masses $m_{O(L)}(P)$ for P in Car_n . This collection will be called *the characteristic masses of* L, or sometimes simply⁴ *the masses of* L, and we will go back to it later. We rather discuss first the question of evaluating, given an arbitrary polynomial P as in (iii), the quantity $\operatorname{Trace}(c_P; W_{\lambda})$. This question does not depend on L.

Evaluation of Trace(c_P ; W_λ). We will use for this the "determinantal" character formula for W_λ proved by Weyl in [WEYL46, Chap. VII §9]. This formula applies to arbitrary elements of $O(\mathbb{R}^n)$, possibly of determinant -1. We found it useful to actually use the following alternative expression proved by Koike and Terada in [KT87] in the spirit of the famous Jacobi-Trudi formula for the Schur polynomials in terms of elementary symmetric polynomials (see Appendix A). Write $t^n P(1/t) = \sum_{i \in \mathbb{Z}} (-1)^i e_i t^i$ (so $e_i = 0$ for i < 0 or i > n). Denote by $\mu_1 \ge \mu_2 \ge \cdots \ge \mu_m$ with $m = \lambda_1$ the partition which is dual to λ , and set $\delta_1 = 0$ and $\delta_j = 1$ for j > 1. Then we have the equality

(1.5)
$$\operatorname{Trace}(\mathbf{c}_P; \mathbf{W}_{\lambda}) = \det(e_{\mu_i - i + j} + \delta_j e_{\mu_i - i - j + 2})_{1 \le i, j \le m}$$

This formula is clearly efficient when $m = \lambda_1$ is small, which suits well for instance the application to $|X_{32}|$ mentioned in §1.1, as it requires all λ 's with $\lambda_1 \leq 4$ for n = 24. Let us note that in this range, the use of the crude degenerate Weyl character formula as in [CR12, §2] would be impracticable as the Weyl group of SO(\mathbb{R}^{24}) is much too big. Actually, the whole tables of invariants obtained in [CR12, §2] for the subgroup of determinant 1 elements in the Weyl groups of type \mathbf{E}_7 , \mathbf{E}_8 and $\mathbf{E}_8 \coprod \mathbf{A}_1$ (with respectively n = 7, 8, 9) can be recomputed essentially instantly using rather Formula (1.5).

Determination of the characteristic masses of L. This is the remaining and most important⁵ unknown. In dimension n as large as 24, it is impossible in general to enumerate the elements of O(L) with a computer, hence to naively list their characteristic polynomials. For instance when Lis a Niemeier lattice then the size of O(L) is always at least 10^{14} , and it is about 10^{30} for $L = E_{24}$. However, those groups have of course much fewer conjugacy classes. Write

$$\operatorname{Conj} \mathcal{O}(L) = \{ \mathcal{C}_i(L) \}_{i \in I}$$

the set of conjugacy classes of O(L). Assuming that we know representatives of the $c_i(L)$, as well as each $|c_i(L)|$, then the enumeration of the characteristic

⁴Beware not to confuse the masses of L in this sense with the mass of the genus of L, which traditionally appears in the study of the Minkowski-Siegel-Smith mass formula.

⁵It is equivalent to determine the finitely many $m_{O(L)}(P)$ for all P in Car_n , and the $\dim W^{O(L)}_{\lambda}$ for all λ , as the $\operatorname{Car}_n \times \Lambda$ -matrix $(\operatorname{Trace}(c_P; W_{\lambda}))_{P,\lambda}$ has rank $|\operatorname{Car}_n|$ for general reasons.

polynomials of O(L) may become straightforward. Of course, if we do not know representatives of $c_i(L)$, but still the trace of the latter in \mathbb{R}^n as well as the power maps on the $c_i(L)$, this may similarly allow to determine the characteristic masses of L.

Example 1.3. (Leech lattice) Consider for instance the case where L = Leech is "the" Leech lattice in \mathbb{R}^{24} . The group O(Leech) is the Conway group Co₀, also denoted 2. Co₁ in the ATLAS p. 180. The character of its natural representation on \mathbb{R}^{24} is the character χ_{102} in the table *loc. cit.* This character, as well as Newton's relations and the power maps of the ATLAS (implemented in GAP), allow to compute the characteristic polynomial of each conjugacy class in O(Leech), hence the characteristic masses of Leech: they are gathered in Table C (see p. 10 below for the notations). Note that despite the huge order $\simeq 8 \cdot 10^{18}$ of O(Leech), this group only has 167 conjugacy classes, and 160 distinct characteristic polynomials. This is actually the minimum for a Niemeier lattice, and makes the table above printable. An interesting consequence of this computation is the observation

$$\frac{1}{|\mathcal{O}(\text{Leech})|} \sum_{g \in \mathcal{O}(\text{Leech})} \det(t \, \mathrm{id}_{24} - g) = t^{24} + t^{16} + t^{12} + t^8 + 1.$$

This asserts the existence of a line of O(Leech)-equivariant alternating gmultilinear form Leech^g $\rightarrow \mathbb{Z}$ for each g in {8, 12, 16, 24}. We refer to [CT19b] for a study of these forms and of the weight 13 pluriharmonic Siegel theta series for $\operatorname{Sp}_{2g}(\mathbb{Z})$ that they allow to construct. The results of this paper suggest several other intriguing constructions to study in the same spirit, for instance whenever a 1 appears as a dimension for $\operatorname{M}_{W_{\lambda}}(O_{24})$ in Table C (the case discussed here corresponding to $\lambda = \emptyset$, 1⁸ and 1¹²).

1.4. Algorithms for computing characteristic masses

Let us give now a first algorithm, called Algorithm A in the sequel, which takes as input the Gram matrix G of some \mathbb{Z} -basis of L and returns for each conjugacy class $c_i(L)$ some representative and its cardinality $|c_i(L)|$, hence in particular the characteristic masses of L. The idea, certainly classical in computational group theory, is to:

A1. Apply the Plesken-Souvignier algorithm [PS97] to G (implemented *e.g.* as qfauto(G) in PARI/GP) to obtain a set \mathcal{G} of generators of O(L),

A2. Choose a (small) finite subset $S \subset L$ stable under O(L), generating $L \otimes \mathbb{R}$, and view O(L) as the subgroup of permutations of S generated by \mathfrak{G} ,

A3. Apply permutation groups algorithms implemented in GAP (such as [HUL00]) to deduce cardinality and representatives of the conjugacy classes of O(L).

A canonical choice of S is the following: for any lattice L set (inductively) $S(L) = M(L) \coprod S(L')$ where M(L) is the subset of elements of L with minimal nonzero length, and where L' is the orthogonal of M(L) in L. The choice S = S(L) has proved efficient enough for us in practice. We will say more about a PARI/GP implementation of the whole algorithm later, when discussing an improvement of it: see §4.3.

Algorithm A is very efficient in small dimension. For instance, when L is a root lattice of type $\mathbf{E}_6, \mathbf{E}_7$ or \mathbf{E}_8 , it returns the characteristic masses of L in a few seconds only.⁶ It turns out that it is still terminates for most of the even lattices of determinant ≤ 2 and dimension ≤ 25 , with running time varying from a few minutes to a few days in dimensions 23, 24 and 25 when terminates. For instance, in the case L = Leech it allows to re-compute Table C from scratch, without relying at all on the ATLAS: it requires about 3 minutes for step A1, nothing for A2, and 42 minutes for A3. On the other hand, it does not terminate for instance on our computer for lattices L in \mathcal{L}_{25} with root system⁷ A₁ D₄ 2D₆ D₈ or A₁ D₆ D₈ D₁₀ (memory issue). Algorithm A is typically very slow (and memory consuming) if either L has too many vectors v of length $v \cdot v = G_{i,i}$ for some $i = 1, \ldots, n$, because of step A1, or if O(L) has too many conjugacy classes, because of step A3. It is also quite sensitive to the choice of Gram matrix G of L in step A1.

In §4, we will explain a significant improvement of Algorithm A when L has a non trivial root system. The basic idea of this Algorithm B is to first write

$$O(L) = W(R) \rtimes O(L)_{\rho}$$

where R is the root system of L, W(R) its Weyl group, ρ a Weyl vector of R and $O(L)_{\rho}$ the stabilizer of ρ in O(L). As we shall see, we may actually reduce the computation of the characteristic masses of L to that of representatives γ_j , and sizes, of the conjugacy classes of the smaller group $O(L)_{\rho}$, an information which can be obtained by replacing O(L) with $O(L)_{\rho}$ in steps A1 and A3 of Algorithm A. There are two ingredients for this reduction. The first is the determination, for each rank r irreducible root system R' of type ADE, of the map $m_S : \operatorname{Car}_r \to \mathbb{Q}$, where S is any coset of W(R') in the full isometry

⁶All the computations in this paper have been made on a processor Intel(R) Xeon(R) CPU E5-2650 v4 @ 2.20GHz with 65 GB of memory. Nevertheless, all the computations involving either Algorithm B, or Algorithm A in small dimension, are equally efficient on our personal computer (processor 1,8 GHz Intel Core i5 with 8 GB of memory).

⁷For $n \leq 25$, it follows from the classification of X_n recalled in §1.1 that two lattices in \mathcal{L}_n are isometric if, and only if, they have isomorphic root systems.

group O(R') of the root system R': see §3 for this step (which does not depend on L). The second is the determination, for each j, of the conjugacy class of γ_j viewed as an element of the automorphism group of the Dynkin diagram of R. See §4 for a detailed discussion of Algorithm B and of its implementation.

Remark 1.5. (Generalizations) In this paper, we use a restricted notion of root which suits well our applications to the lattices in \mathcal{L}_n . A minor modification of Algorithm B allows to consider the most general roots, namely the elements α of a lattice L such that the orthogonal symmetry about α preserves L. In a different direction, it would be useful to extend the algorithms above to the context of hermitian or quaternionic positive definite lattices, possibly over totally real number fields, using the theory of complex or quaternionic reflection groups (see e.g. [COH76, COH80]). That should help extending to higher ranks and weights the computations of dimension spaces of automorphic forms for definite unitary groups (hermitian or quaternionic) started in the literature (e.g. in [LP02, LOE10, DUM13, DEM14, GV14]).

1.6. Main results

Using Algorithm B, it only takes a few seconds to the computer to compute all the characteristic masses of each Niemeier lattices with roots, except in the case (trivial anyway) of D_{24} for which the Plesken-Souvignier algorithm needs about 2 minutes. It is equally efficient in any dimension ≤ 25 : the characteristic masses are computed in a few seconds, except for ten lattices (in dimension 23 or 25) for which it requires less than 5 minutes, and for the lattice $A_1 \oplus$ Leech in \mathcal{L}_{25} (about 35 minutes). We refer to the homepage [CHE19] for the gram matrices we used in our computations. Our main result is then the following.

Theorem. (i) Assume $n \leq 25$. The characteristic masses of all $L \in \mathcal{L}_n$ are those given⁸ in [CHE19].

(ii) The nonzero values of dim $M_{\lambda}(O_{24})$ for $\lambda_1 \leq 3$ are given in Table C.

Table C is deduced from assertion (i) for n = 24 using observation (b) and Formulas (1.4) & (1.5). This step is very efficient: once the masses in (i) are computed, it takes only 5 minutes about to produce this table. The format of the table is as follows. The notation $n_1^{m_1} \dots n_r^{m_r}$ for a partition λ means

⁸They cannot be printed here: there are 53204 polynomials P with $m_{O(L)}(P) \neq 0$ for some L in \mathcal{L}_{24} , that is about half $|Car_{24}|$.

that the diagram of λ has exactly m_i rows of size n_i for $i = 1, \ldots, r$, and no other row. Set $d_{\lambda} = \dim M_{W_{\lambda}}(O_{24})$ and denote by $\operatorname{ass}(\lambda)$ the associate of λ (see §A). The column dim gives the integer d_{λ} in the case $\lambda = \operatorname{ass}(\lambda)$, and the two integers $d_{\lambda} : d_{\operatorname{ass}(\lambda)}$ otherwise. See [CHE19] for more extensive tables, including for instance all λ with $\lambda_1 = 4$ and arbitray $n \leq 25$.

Remark 1.7. Let L be a lattice in \mathbb{R}^n , fix γ in O(L) and write $\det(t - \gamma) = (t-1)^a (t+1)^b Q(t)$ with Q in $\mathbb{Z}[t]$ and $Q(-1)Q(1) \neq 0$. Assuming furthermore $n \equiv -1, 0, 1 \mod 8$ and $L \in \mathcal{L}_n$ then Proposition 3.7 in [CT19a] shows⁹ that for a = 0 (resp. b = 0) the integer Q(1) (resp. Q(-1)) is a square. This constraint is in agreement with our computations.

1.8. A direct computation in the case of Niemeier lattices

In section 5, we will explain an alternative (and human) computation of the characteristic masses of Niemeier lattices. By the results of §3, we are left to determine, for each Niemeier lattice L with non-empty root system R, the G(R)-conjugacy classes of the elements of the subgroup O(L)/W(R) of G(R), where G(R) is the automorphism group of the Dynkin diagram of R. We do so using a tedious case by case analysis.

We found it useful to gather first in section 2 some elementary results about the hyperoctahedral group $H_n = \{\pm 1\}^n \rtimes S_n$. This group is both a typical direct summand of the G(R) above, and closely related to the Weyl groups of type \mathbf{D}_n studied in §3.2. In particular, we introduce and characterize directly in §2.5 and §2.7 a few specific subgroups of H_n that will play a role in the analysis of Niemeier lattices in section 5.

Although more interesting (at least to us) from a mathematical point of view, it will be eventually clear that this nonautomatized method is too complicated to be used systematically: it would even require some work to attack the dimensions 23 and 25 along the same lines. Nevertheless, it provides an important check that the masses returned by the implementation of our algorithms are correct.

AKNOWLEDGEMENTS: We thank Jean Lannes and Olivier Taïbi for useful discussions, the LMO for sharing the machine pascaline, as well as Bill Allombert for answering our questions on PARI/GP.

⁹If det $\gamma = -1$ (so b is odd) and a = 0 (so n is odd), apply the proposition to $-\gamma$.

Contents

1	Introduction	1
2	Preliminaries on the hyperoctahedral groups	11
3	Characteristic masses of root lattices	16
4	An algorithm computing characteristic masses	23
5	The characteristic masses of Niemeier Lattices with roots	26
6	References	34
Α	Irreducible characters of compact orthogonal groups	37
в	An asymptotic formula	39
С	Tables	40

GENERAL NOTATIONS AND CONVENTIONS

In this paper, all group actions will be on the left. We denote by |X| the cardinality of the set X. For $n \ge 1$ an integer, we denote by S_n the symmetric group on $\{1, \ldots, n\}$, by $\operatorname{Alt}_n \subset S_n$ the alternating subgroup, and we set $\mathbb{Z}/n := \mathbb{Z}/n\mathbb{Z}$.

If V is an Euclidean space, we usually denote by $x \cdot y$ its inner product, with associated quadratic form $q: V \to \mathbb{R}$ defined by $q(x) = \frac{x \cdot x}{2}$. A *lattice* in V is a subgroup generated by a basis of V, or equivalently, a discrete subgroup L with finite covolume, denoted covol L.

If L is a lattice in the Euclidean space V, its dual lattice is the lattice L^{\sharp} defined as $\{v \in V \mid v \cdot x \in \mathbb{Z}, \forall x \in L\}$. We say that L is integral (resp. even) if we have $L \subset L^{\sharp}$ (resp. $q(L) \subset \mathbb{Z}$). An even lattice is integral. If L is integral, we have $(\operatorname{covol} L)^2 = |L^{\sharp}/L|$. This integer is also the determinant det L of the Gram matrix $\operatorname{Gram}(e) = (e_i \cdot e_j)_{1 \leq i,j \leq n}$ of any \mathbb{Z} -basis $e = (e_1, \ldots, e_n)$ of L. The orthogonal group of L is the finite group $O(L) = \{\gamma \in O(V), \gamma(L) = L\}$.

In the tables of Appendix C, we use the notation $1^{a_1} 2^{a_2} \dots m^{a_m}$ for the polynomial $\varphi_1^{a_1} \varphi_2^{a_2} \cdots \varphi_m^{a_m}$, where φ_n is the *n*-th cyclotomic polynomial and where the symbol " i^a " is omitted for a = 0, and shorten as "i" for a = 1.

2. Preliminaries on the hyperoctahedral groups

2.1. The hyperoctahedral group

Let $n \ge 1$ be an integer. The symmetric group S_n on the set $\{1, \ldots, n\}$ acts on the elementary abelian 2-group $\{\pm 1\}^n$ by permuting coordinates. The *hyperoctahedral group* on n letters is defined as the semi-direct product

$$\mathbf{H}_n = \{\pm 1\}^n \rtimes \mathbf{S}_n.$$

Equivalently, H_n is the wreath product $\{\pm 1\} \geq S_n$. It is isomorphic to several familiar groups: the Weyl group of a root system of type \mathbf{B}_n or \mathbf{C}_n , the subgroup of monomial matrices in $\mathrm{GL}_n(\mathbb{Z})$, the orthogonal group of the standard unimodular lattice I_n , the subgroup of the symmetric group on $\{\pm 1, \pm 2, \ldots, \pm n\}$ of permutations σ with $\sigma(-i) = -\sigma(i)$ for all i, etc...

In this paper, we will encounter H_n first when discussing $O(D_n)$ and again when studying automorphism groups of isotypic root systems. Certain subgroups of the hyperoctahedral groups will play a role in the study of Niemeier lattices. Here is an example of an interesting subgroup that will occur in the case n = 4. We denote by $\pi : H_n \to S_n$ the canonical projection.

Example 2.2. The group $\operatorname{GL}_2(\mathbb{Z}/3)$ acts on the 8-elements set $(\mathbb{Z}/3)^2 - \{0\}$ by permuting the 4 disjoint pairs of the form $\{v, -v\}$. By the universal property of wreath products, the choice of elements v_1, v_2, v_3, v_4 such that $(\mathbb{Z}/3)^2 - \{0\} = \coprod_i \{v_i, -v_i\}$ defines an embedding $\iota : \operatorname{GL}_2(\mathbb{Z}/3) \longrightarrow \operatorname{H}_4$ (a different choice leading to an H₄-conjugate embedding). We have $\iota(-\operatorname{Id}_2) = -1$; the morphism $\pi \circ \iota$ has kernel $\pm \operatorname{Id}_2$ and induces "the" exceptional isomorphism $\operatorname{PGL}_2(\mathbb{Z}/3) \simeq S_4$. The restriction of $\pi \circ \iota$ to the stabilizer of v_i in $\operatorname{GL}_2(\mathbb{Z}/3)$ is an isomorphism onto the stabilizer ($\simeq S_3$) of i in $\{1, 2, 3, 4\}$.

We end this paragraph with a few notations and remarks about the basic structure of H_n . We denote by ε_i the element of $\{\pm 1\}^n$ whose j^{th} -component is 1 for $j \neq i$ and -1 for j = i. The center of H_n is generated by the element $-1 = \prod_{i=1}^n \varepsilon_i$. The signature $\epsilon : S_n \to \{\pm 1\}$, composed with the natural projection $\pi : H_n \to S_n$, defines a morphism $H_n \to \{\pm 1\}$ that we will still denote by ϵ . Another important morphism $s : H_n \to \{\pm 1\}$ is defined by

(2.1)
$$s(v\sigma) = \prod_{i=1}^{n} v_i$$
, for all $\sigma \in S_n$ and $v = (v_i) \in \{\pm 1\}^n$.

The product character ϵ s coincides with the determinant when we view H_n as a the subgroup of monomial matrices in $GL_n(\mathbb{Z})$.

2.3. Conjugacy classes of H_n

Let Σ be a nonempty subset of $\{1, \ldots, n\}$. A cycle in \mathcal{H}_n with support Σ is an element of the form h = vc, where $c \in \mathcal{S}_n$ permutes transitively the elements of Σ and fixes its complement, and where $v = (v_i) \in \{\pm 1\}^n$ satisfies $v_i = 1$ for $i \notin \Sigma$. Such a cycle has a length l(h) defined as $|\Sigma|$, and a sign s(h) (an element in $\{\pm 1\}$). This sign is also the *i*-th coordinate of $h^{l(h)}$ for any *i* in Σ , and l(h) is the order of *c*. One easily checks that two cycles are conjugate in \mathcal{H}_n if, and only if, they have the same length and the same sign.

Just as for S_n , any element h of H_n may be written as a product of cycles h_i with disjoint supports, this decomposition being unique up to permutation of those cycles. The sum of the lengths of the cycles h_i with $s(h_i) = 1$ (resp. $s(h_i) = -1$) is an integer denoted $n_+(h)$ (resp $n_-(h)$); the collection of the length $l(h_i)$ of those h_i defines a integer partition of $n_+(h)$ (resp. $n_-(h)$) that we denote by $p_+(h)$ (resp. $p_-(h)$). We have $n_+(h) + n_-(h) = n$. The type of h is defined as the couple of integer partitions $(p_+(h), p_-(h))$. Two elements of H_n are conjugate if, and only if, they have the same type.

In the sequel, we will have to determine the type of all the elements of certain specific subgroups $G \subset H_n$. For instance, when G is the group $\iota(\operatorname{GL}_2(\mathbb{Z}/3))$ of Example 2.2, this information is given in Table 2, the row size giving the number of elements of the corresponding type **divided by** |G|:

type	1^{4}	1^4	112	2^2	13	13	4
size	1/48	1/48	1/4	1/8	1/6	1/6	1/4

Table 2: The H₄-conjugacy classes of the elements of $GL_2(\mathbb{Z}/3)$.

In this table, and in others that we will give later, we use standard notations for partitions, and print p_+ in black and p_- in cyan. So the sequence of symbols $1^{a_1} 1^{b_1} 2^{a_2} 2^{b_2} \dots i^{a_i} i^{b_i} \dots$ stands for the couple (p_+, p_-) where p_+ is the partition of $\sum_i a_i$ in a_1 times 1, a_2 times 2, and so on, and p_- is the partition of $\sum_i b_i$ in b_1 times 1, b_2 times 2, and so on. The symbol " i^m " (resp. " i^m ") is omitted for m = 0, and replaced by "i" (resp. "i") for m = 1.

Remark 2.4. Table 2 is easily deduced from the conjugacy classes of $\operatorname{GL}_2(\mathbb{Z}/3)$. To fix ideas, define the embedding ι in Example 2.2 by choosing v_1, v_2, v_3 and v_4 to be respectively $\begin{bmatrix} 0\\1 \end{bmatrix}, \begin{bmatrix} 1\\0 \end{bmatrix}, \begin{bmatrix} 1\\1 \end{bmatrix}$ and $\begin{bmatrix} 1\\-1 \end{bmatrix}$. Then the images under ι of the elements $\pm I_2$, $\begin{bmatrix} 1&0\\0&-1 \end{bmatrix}, \begin{bmatrix} 0&-1\\1&0 \end{bmatrix}, \begin{bmatrix} 1&1\\0&-1 \end{bmatrix}, \pm \begin{bmatrix} 1&0\\0&1 \end{bmatrix}$ and $\pm \begin{bmatrix} 0&1\\1&1 \end{bmatrix}$ of $\operatorname{GL}_2(\mathbb{Z}/3)$ are respectively ± 1 , $\varepsilon_1(34)$, $\varepsilon_2(12)\varepsilon_4(34)$, $\pm \varepsilon_1\varepsilon_4(134)$ and $\pm \varepsilon_2(1342)$. **2.5.** Digression: subgroups of certain wreath products

Let G be a group, X a set equipped with a transitive action of G, and A an abelian group. The group G acts in a natural way on the abelian group A^X of all functions $X \to A$, so we can form the semi-direct product $H := A^X \rtimes G$.

We denote by $\pi : H \to G$ the canonical projection, with kernel ker $\pi = A^X$. We have a "diagonal" map $\delta : A \to A^X$, defined by $\delta(a)(x) = a$ for all a in A and x in X. This map δ is an embedding of G-modules if we view the source A as a trivial G-module: the image of δ is a central subgroup of H. Our aim in this paragraph is to study:

- the set \mathcal{C} of subgroups $C \subset H$ with $\pi(C) = G$ and ker $\pi \cap C = 1$,

- the set \mathfrak{G} of subgroups $\widetilde{G} \subset H$ with $\pi(\widetilde{G}) = G$ and ker $\pi \cap \widetilde{G} = \delta(A)$.

The group H acts both on \mathcal{C} and \mathcal{G} by conjugation. We start with two simple observations:

- For any group morphism $\chi: G \to A$, the set $G^{\chi} := \{\delta(\chi(g)) \cdot g, g \in G\}$ is a subgroup of H isomorphic to G, and G^{χ} is an element of \mathcal{C} .

- There is a natural map $c_2 : \mathcal{G} \to H^2(G, A)$, sending \widetilde{G} in \mathcal{G} to the equivalence class of the central extension $1 \to A \xrightarrow{\delta} \widetilde{G} \xrightarrow{\pi} G \to 1$. Two elements of \mathcal{G} which are *H*-conjugate are also A^X -conjugate, hence define the same class in $H^2(G, A)$.

We fix some $x \in X$ and denote by $G_x \subset G$ the isotropy group of x. For each integer $m \geq 0$, we denote by $\mathbf{r}_m : \mathrm{H}^m(G, A) \longrightarrow \mathrm{H}^m(G_x, A)$ the usual restriction map on the cohomology groups of the trivial G-module A.

- **Proposition 2.6.** (i) For $\chi, \chi' \in \text{Hom}(G, A)$, the subgroups G^{χ} and $G^{\chi'}$ of H are conjugate if, and only if, χ and χ' coincide on G_x .
 - (ii) If \mathbf{r}_1 is surjective then any subgroup $C \in \mathfrak{C}$ is conjugate to G^{χ} for some $\chi \in \operatorname{Hom}(G, A)$.
- (iii) If \mathbf{r}_1 is surjective then the map $\mathbf{c}_2 : H \setminus \mathcal{G} \to H^2(G, A)$ is injective, and its image is the subgroup ker \mathbf{r}_2 of extensions which split over G_x .

Proof. We shall use twice the following classical facts. Let Γ be a group acting on an abelian group V and denote by $\pi : V \rtimes \Gamma \to \Gamma$ the natural projection. Let \mathcal{K} be the set subgroups $K \subset V \rtimes \Gamma$ with $\pi(K) = \Gamma$ and ker $\pi \cap V = 1$. Any $K \in \mathcal{K}$ has the form $\{s(\gamma) \gamma, \gamma \in \Gamma\}$ for a unique 1-cocyle $s \in \mathbb{Z}^1(\Gamma, V)$, that we denote \mathfrak{s}_K . The map $K \mapsto \mathfrak{s}_K, \mathcal{K} \to \mathbb{Z}^1(\Gamma, V)$, is bijective; two elements K, K' in \mathcal{K} are conjugate by an element of V if, and only if, s_K and $s_{K'}$ have the same class in $H^1(\Gamma, V)$. Last but not least, note that K, K' in \mathcal{K} are conjugate by an element of V if, and only if, they are conjugate in $V \rtimes \Gamma$: if we have $K' = gKg^{-1}$ with $g \in V \rtimes \Gamma$, we may write g = vk with $v \in V$ and $k \in K$, and we have $K' = vKv^{-1}$.

We apply this first to $\Gamma = G$ and $V = A^X$. The map $\operatorname{Hom}(G, A) = Z^1(G, A) \to Z^1(G, A^X)$ defined by δ sends χ to the 1-cocycle defining G^{χ} . The choice of $x \in X$ identifies the *G*-module A^X with the co-induced module of the trivial G_x -module A to G. By Shapiro's lemma, we obtain for each integer $m \geq 0$ a natural isomorphism sh : $\operatorname{H}^m(G, A^X) \xrightarrow{\sim} \operatorname{H}^m(G_x, A)$. Concretely, if $f: G^m \to A^X$ is an *m*-cocycle, then $\operatorname{sh}(f)$ is the class of the *m*-cocycle $f': G^m_x \to A$ defined by $f'(g_1, \ldots, g_m) = f(g_1, \ldots, g_m)(x)$. It follows that the composition of the maps

$$\mathrm{H}^{m}(G,A) \xrightarrow{\mathrm{H}^{m}(\delta)} \mathrm{H}^{m}(G,A^{X}) \xrightarrow{\overset{\mathrm{sh}}{\sim}} \mathrm{H}^{m}(G_{x},A)$$

coincides with the map r_m . For m = 1, this proves assertions (i) and (ii).

Let us prove assertion (iii). Let Q be the cokernel of δ . By applying the first paragraph above to $\Gamma = G$ and V = Q, we obtain a natural bijection $c_1 : H \setminus \mathcal{G} \xrightarrow{\sim} H^1(G, Q)$. The long exact sequence of cohomology groups associated to $0 \to A \xrightarrow{\delta} A^X \to Q \to 0$ contains a piece of the form

$$\mathrm{H}^{1}(G,A) \xrightarrow{\mathrm{H}^{1}(\delta)} \mathrm{H}^{1}(G,A^{X}) \longrightarrow \mathrm{H}^{1}(G,Q) \xrightarrow{\eta} \mathrm{H}^{2}(G,A) \xrightarrow{\mathrm{H}^{2}(\delta)} \mathrm{H}^{2}(G,A^{X}).$$

By the second paragraph, the kernel of η is isomorphic to the cokernel of r_1 , and the image of η is the kernel of r_2 . As it is straightforward to check from the definition of c_2 that we have $\eta \circ c_1 = c_2$, this concludes the proof of assertion (iii).

2.7. Applications to H_n

The group H_n is of course the special case of the construction of §2.5 with $G = S_n, X = \{1, \ldots, n\}$ and $A = \{\pm 1\}$ (multiplicative group). The signature ϵ gives rise to the subgroup S_n^{ϵ} of H_n whose elements have the form $\epsilon(\sigma)\sigma$, $\sigma \in S_n$. For any transposition τ in S_n we have $n_-(\epsilon(\tau)\tau) = n - 2$, whereas $n_-(\sigma) = 0$ for all σ in S_n : this shows that S_n^{ϵ} is not conjugate to S_n in H_n for n > 2 (a fact which also follows from assertion (i) below).

Proposition 2.8. (i) Let G be a subgroup of H_n of order n! with $\pi(G) = S_n$. Then G is either conjugate to S_n or to S_n^{ϵ} . Moreover, S_n and S_n^{ϵ} are conjugate in H_n if, and only if, we have $n \leq 2$.

- (ii) Let G be a subgroup of H_n of order 2n! with $\pi(G) = S_n$. Then -1 is in G and exactly one of the following properties holds:
 - (a) G is conjugate to $\{\pm 1\} \cdot S_n$,
 - (b) n = 2 and $G \simeq \mathbb{Z}/4$,
 - (c) n = 4 and G is conjugate to the group $\operatorname{GL}_2(\mathbb{Z}/3)$ embedded in H_4 as in Example 2.2.

Proof. Note first that in case (ii), $\{\pm 1\}^n \cap G$ is a normal subgroup of order 2 of G, hence it is central and generated by −1 by the assumption $\pi(G) = S_n$. The stabilizer of n in S_n is naturally identified with S_{n-1} , with the convention $S_0 = 1$. The signature ϵ is a generator of $H^1(S_n, \{\pm 1\})$, so the restriction map $H^1(S_n, \{\pm 1\}) \rightarrow H^1(S_{n-1}, \{\pm 1\})$ is clearly surjective, and bijective for $n \neq 2$. Moreover, we know form Schur that the restriction map $r_2 : H^2(S_n, \{\pm 1\}) \rightarrow$ $H^2(S_{n-1}, \{\pm 1\})$ is surjective as well for all n > 1, and that the dimension of the $\mathbb{Z}/2$ -vector space $H^2(S_n, \{\pm 1\})$ is 2 for $n \geq 4$, 1 for n = 3 and 2, and 0 for n = 1 [SCH1911]. The kernel of r_2 is thus 0 for $n \neq 2, 4$, isomorphic to $\mathbb{Z}/2$ otherwise. We conclude by Proposition 2.6 and Example 2.2.

Remark 2.9. The natural map $\mathrm{H}^{i}(\mathrm{Alt}_{4}, \mathbb{Z}/2) \to \mathrm{H}^{i}(\mathrm{Alt}_{3}, \mathbb{Z}/2)$ is $0 \to 0$ for i = 1 and $\mathbb{Z}/2 \to 0$ for i = 2. By Proposition 2.6 (iii), there is thus a unique conjugacy class of nonsplit central extensions of Alt_{4} by $\{\pm 1\}$ in H_{4} (or in $\{\pm 1\}^{4} \rtimes \mathrm{Alt}_{4}$). As Alt_{4} does not embed in $\mathrm{GL}_{2}(\mathbb{Z}/3)$, one such extension is the inverse image of Alt_{4} in the extension described in Example 2.2.

We now give another example. As is well-known, the group S_5 has a unique isomorphism class of transitive actions on the set $\{1, \ldots, 6\}$, obtained from the conjugation action on its 6 subgroups of order 5. We fix such an action and consider the associated semi-direct product $\{\pm 1\}^6 \rtimes S_5$, as in §2.5. We have a defined *loc. cit.* a set \mathcal{G} of subgroups of $\{\pm 1\}^6 \rtimes S_5$ which are central extensions of S_5 by $\{\pm 1\}$.

Proposition 2.10. The set \mathfrak{G} is the disjoint union of two conjugacy classes: the one of the split extension $\{\pm 1\} \cdot S_5$, and another one consisting of nonsplit extensions which are split over the alternating subgroup Alt₅ of S_5 .

Proof. Let $N \subset S_5$ be the normalizer of the subgroup $S = \langle (12345) \rangle$. Then N is the semi-direct product of $\langle (2354) \rangle \simeq \mathbb{Z}/4$ by $S \simeq \mathbb{Z}/5$, so we have $\mathrm{H}^i(N,\mathbb{Z}/2) \simeq \mathbb{Z}/2$ for each $i \geq 0$ and the restriction map $\mathrm{H}^1(S_5,\mathbb{Z}/2) \longrightarrow \mathrm{H}^1(N,\mathbb{Z}/2)$ is an isomorphism. We observe from the presentation given by

Schur of the two Schur-covers of S_5 that they are non split over the subgroups of S_5 containing a double transposition, such as N or Alt₅. This implies that the kernel of the restriction map $H^2(S_5, \mathbb{Z}/2) \longrightarrow H^2(N, \mathbb{Z}/2)$ is generated by the remaining nonzero class in $H^2(S_5, \mathbb{Z}/2)$, namely the one which splits over Alt₅ (recall $H^2(Alt_5, \mathbb{Z}/2) \simeq \mathbb{Z}/2$), and we conclude by Proposition 2.6. \Box

A homomorphism $S_5 \to S_6$ as above can alternatively be constructed from the natural action of $PGL_2(\mathbb{Z}/5) \simeq S_5$ on the projective line $P^1(\mathbb{Z}/5)$. The action of $GL_2(\mathbb{Z}/5)$ on the 12-elements set $((\mathbb{Z}/5)^2 - \{0\})/\{\pm 1\}$ permutes the 6 disjoint pairs of the form $\{v, 2v\}$, which defines a natural conjugacy class of embeddings

(2.2)
$$\iota: \operatorname{GL}_2(\mathbb{Z}/5)/\{\pm I_2\} \longrightarrow \{\pm 1\}^6 \rtimes S_5.$$

The group $\iota(\operatorname{GL}_2(\mathbb{Z}/5)/\{\pm I_2\})$ belongs to the second class of Proposition 2.10 (recall $\operatorname{PSL}_2(\mathbb{Z}/5) \simeq \operatorname{Alt}_5$). The map ι is explicit enough to allow the computation of the conjugacy classes of the elements of $\iota(\operatorname{GL}_2(\mathbb{Z}/5)/\{\pm I_2\})$ viewed as a subgroup of $\operatorname{H}_6 \supset \{\pm 1\}^6 \rtimes S_5$: they are gathered in Table 3.

	1^{6}										
size	1/240	1/240	1/16	1/16	1/12	1/12	1/12	1/4	1/10	1/10	1/6

Table 3: The H₆-conjugacy classes of the elements of $GL_2(\mathbb{Z}/5)/\{\pm I_2\}$.

3. Characteristic masses of root lattices

3.1. Root systems and root lattices

Let V be an Euclidean space. By a root of V we mean an element $\alpha \in V$ with $\alpha \cdot \alpha = 2$; we denote by R(V) the set of roots of V (a sphere). For each $\alpha \in R(V)$, the orthogonal reflection about α is an element s_{α} of O(V), given by the formula $s_{\alpha}(x) = x - (\alpha \cdot x) \alpha$.

An **ADE** root system in V is a finite set $R \subset \mathbb{R}(V)$ generating V as a real vector space, and such that for all $\alpha, \beta \in R$ we have $\alpha \cdot \beta \in \mathbb{Z}$ and $s_{\alpha}(\beta) \in R$. In particular, R is a root system in the sense of Bourbaki [BOU81b], and each irreducible component of R is of type \mathbf{A}_n with $n \ge 1$, \mathbf{D}_n with $n \ge 4$, or \mathbf{E}_n with n = 6, 7, 8. The root lattice of R is the lattice $Q(R) \stackrel{\text{def}}{=} \sum_{\alpha \in R} \mathbb{Z} \alpha \subset V$ generated by R. This is an even lattice, and we have the important equality

(3.1)
$$\mathbf{R}(V) \cap \mathbf{Q}(R) = R.$$

If $L \subset V$ is any integral lattice, we denote by $R(L) = L \cap R(V)$ the set of roots of L. It follows at once from the definitions that R(L) is an **ADE** root system in the Euclidean subspace U of V generated by R(L). We say that L is a root lattice if R(L) generates L as an abelian group, *i.e.* if we have L = Q(R(L)) (hence U = V). By definitions and (3.1), the map $R \mapsto Q(R)$ is a bijection between the set of **ADE** root systems of V and the set of root lattices of V, whose inverse is $L \mapsto R(L)$.

We shall always use a bold font to denote an isomorphism class of root systems, and reserve the normal font for a root lattice with the corresponding root system. For instance, if for $n \ge 2$ we set $D_n = \{(x_i) \in \mathbb{Z}^n, \sum_i x_i \equiv 0 \mod 2\}$ as in the introduction, then $R(D_n)$ is a root system of type \mathbf{D}_n in the standard Euclidean space \mathbb{R}^n . We have also defined *loc. cit.* the lattice E_n for $n \equiv 0 \mod 8$. It is easy to check $R(E_n) = R(D_n)$ for n > 8 and that $R(E_8)$ is of type \mathbf{E}_8 . We choose in an arbitrary way root lattices A_n for $n \ge 1$, as well as E_6 and E_7 , whose root systems are of type \mathbf{A}_n , \mathbf{E}_6 and \mathbf{E}_7 .

Let L be an integral lattice in V and set R = R(L). The s_{α} with α in Rgenerate a subgroup of O(L) called the *Weyl group* of L, and denoted W(L). This is a normal subgroup of O(L), and we denote by G(L) = O(L)/W(L)the quotient group. Assume first that L is the root lattice Q(R); in this case we also set W(R) := W(L), O(R) := O(L) (this latter group is also denoted A(R) by Bourbaki) and G(R) = G(L). As is well-known, G(R) is isomorphic to the automorphism group of the Dynkin diagram of R, and we have

(3.2)
$$G(R) \simeq \begin{cases} 1 & \text{for } R \simeq \mathbf{A}_1, \mathbf{E}_7, \mathbf{E}_8 \\ S_3 & \text{for } R \simeq \mathbf{D}_4, \\ \mathbb{Z}/2 & \text{otherwise.} \end{cases}$$

Moreover, W(R) permutes the positive root systems¹⁰ R^+ of R, or equivalently the Weyl vectors¹¹ of R, in a simply transitive way. Let us now go back to the case of an arbitrary L. The set R is a root system in the Euclidean space U generated by R, and the restriction $\sigma \mapsto \sigma_{|U}$ induces a morphism $O(L) \rightarrow$ O(R) and an isomorphism $W(L) \xrightarrow{\sim} W(R)$. It follows that O(L) permutes the Weyl vectors of R, and that W(L) permutes them simply transitively. So for any Weyl vector ρ of R, the stabilizer $O(L)_{\rho}$ of ρ in O(L) is naturally isomorphic to G(L) and we have

(3.3)
$$W(L) \cap O(L)_{\rho} = 1$$
, $O(L) = W(L) \cdot O(L)_{\rho}$ and $W(L) \simeq W(R)$,

so that O(L) is the semi-direct product of $O(L)_{\rho}$ by W(L).

¹⁰Recall that a positive root system in R is a subset of the form $\{\alpha \in R, \varphi(\alpha) > 0\}$ where $\varphi : V \to \mathbb{R}$ is a linear form with $0 \notin \varphi(R)$.

¹¹A Weyl vector of R is a vector of the form $\rho = \frac{1}{2} \sum_{\alpha \in R^+} \alpha$ for R^+ a positive root system of R. In particular we have $2\rho \in Q(R)$.

3.2. Characteristic masses of irreducible root lattices

In this paragraph, we let $R \subset V$ be an **ADE** root system and L = Q(R). Set $n = \dim V$. Our aim is to determine the characteristic masses of O(L)and, more generally, the map $m_S : \operatorname{Car}_n \to \mathbb{Q}_{\geq 0}$ where S is any subset of the form $\sigma W(L)$ with $\sigma \in O(L)$ (see §1.2 (ii) for the definition of m_S). We assume first R irreducible, and argue case by case.

(A) CASE $R \simeq \mathbf{A}_n$ WITH $n \ge 1$.

We may assume that V is the hyperplane of sum 0 vectors in \mathbb{R}^{n+1} and $R = \{\pm(\epsilon_i - \epsilon_j), 1 \le i < j \le n+1\}$, where $\epsilon_1, \ldots, \epsilon_{n+1}$ denotes the canonical basis of \mathbb{R}^{n+1} , and $L = A_n$. The group W(A_n) may be identified with the symmetric group S_{n+1}, acting on V by permuting coordinates.

Let S denote the set of integer sequences $\underline{m} = (m_i)_{i\geq 1}$ with $m_i \geq 0$ for each *i*, and $m_i = 0$ for *i* big enough. Let $\mathcal{A}_n \subset S$ denote the subset of \underline{m} such that $\sum_i i m_i = n + 1$. For any \underline{m} in \mathcal{A}_n , the elements of S_{n+1} whose cycle decomposition contains m_i cycles of length *i* for each *i* form a single conjugacy class $C_{\underline{m}} \subset S_{n+1}$. We have furthermore $|C_{\underline{m}}| = (n+1)!/n_{\underline{m}}$ with

$$\mathbf{n}_{\underline{m}} = \prod_{i} m_i! \; i^{m_i}.$$

The characteristic polynomial of $C_{\underline{m}}$ acting on V is

$$P_{\underline{m}} = (t-1)^{-1} \prod_{i} (t^{i} - 1)^{m_{i}}$$

since \mathbb{R}^{n+1}/V is the trivial representation of S_{n+1} . The following trivial lemma even shows that we have $P_m \neq P_{\underline{m'}}$ for $\underline{m} \neq \underline{m'}$.

Lemma 3.3. The polynomials $t^{l} - 1$, with $l \ge 1$, are \mathbb{Z} -linearly independent in the multiplicative group of the field $\mathbb{Q}(t)$.

As a consequence, we obtain the:

Corollary 3.4. For $n \ge 1$, we have $m_{W(A_n)}(P_{\underline{m}}) = 1/n_{\underline{m}}$ for \underline{m} in A_n , and $m_{W(A_n)}(P) = 0$ for all other P in Car_n .

As is easily seen, the element $-1 = -id_V$ is in $W(A_n)$ if and only if n = 1, and we have $O(A_n) = W(A_n) \cup -W(A_n)$ (this fits of course Formula (3.2)). The map m_S for the coset $S = -W(A_n)$ is deduced from $m_{W(A_n)}$ by the following trivial lemma: **Lemma 3.5.** Let S be a finite subset of O(V) with dim V = n. Then for all $P \in \mathbb{R}[t]$ we have $m_{-S}(P) = m_S(Q)$ with $Q(t) = (-1)^n P(-t)$.

(D) CASE $R \simeq \mathbf{D}_n$ WITH $n \ge 3$.

We may assume $V = \mathbb{R}^n$, $R = \{\pm \epsilon_i \pm \epsilon_j, 1 \leq i < j \leq n\}$ where $\epsilon_1, \ldots, \epsilon_n$ denote again the canonical basis of V, and $L = D_n$. The lattice D_n is the largest even sublattice of the standard lattice

$$\mathbf{I}_n = \mathbb{Z}^n = \bigoplus_{i=1}^n \mathbb{Z} \epsilon_i,$$

and thus $O(I_n)$ is a subgroup of $O(D_n)$. This group $O(I_n)$ is nothing else than the hyperoctahedral group H_n already introduced in §2.1: we have

$$\mathcal{O}(\mathcal{I}_n) = \mathcal{H}_n = \{\pm 1\}^n \rtimes \mathcal{S}_n$$

where S_n (resp. $\{\pm 1\}^n$) act on \mathbb{R}^n by permuting coordinates (resp. sign changes). As is well-known, $W(D_n)$ is the index 2 subgroup ker s of $O(I_n)$ (recall s is defined by Formula (2.1)). By (3.2) we also have

(3.4)
$$O(I_n) = O(D_n)$$
 for $n \neq 4$ and $G(D_4) \simeq S_3$ (triality).

The conjugacy classes of \mathcal{H}_n have been recalled in §2.3. Let $\mathcal{D}_n \subset \mathbb{S} \times \mathbb{S}$ be the subset of $(\underline{m}^+, \underline{m}^-)$ with $\sum_i i (m_i^+ + m_i^-) = n$. For any $(\underline{m}^+, \underline{m}^-)$ in \mathcal{D}_n the elements of \mathcal{H}_n whose cycle decomposition contains m_i^+ (resp. m_i^-) cycles of length *i* with sign +1 (resp. -1) for each *i* form a single conjugacy class $\mathcal{C}_{\underline{m}^+,\underline{m}^-} \subset \mathcal{H}_n$. We easily check $|\mathcal{C}_{\underline{m}^+,\underline{m}^-}| = 2^n n! / n_{\underline{m}^+,\underline{m}^-}$ with

$$\mathbf{m}_{\underline{m}^+,\underline{m}^-} = \prod_i m_i^+! m_i^-! (2i)^{m_i^+ + m_i^-},$$

and $s(C_{\underline{m}^+,\underline{m}^-}) = (-1)^{|\underline{m}^-|}$ where we have set $|\underline{m}| = \sum_i m_i$ for $\underline{m} \in S$. The characteristic polynomial of $C_{\underline{m}^+,\underline{m}^-}$ acting on V is

$$\mathbf{P}_{\underline{m}^+,\underline{m}^-} = \prod_i (t^i - 1)^{m_i^+} (t^i + 1)^{m_i^-} = \prod_i (t^i - 1)^{m_i^+ - m_i^- + m_{i/2}^-},$$

where we have set $m_{i/2}^- = 0$ for i odd, and used for $i \ge 1$ the relation $(t^i - 1)(t^i + 1) = (t^{2i} - 1)$. In contrast with the \mathbf{A}_n case, we may thus have $\mathbf{P}_{\underline{m}^+,\underline{m}^-} = \mathbf{P}_{\underline{n}^+,\underline{n}^-}$ for distinct $(\underline{m}^+,\underline{m}^-)$ and $(\underline{n}^+,\underline{n}^-)$ in \mathcal{D}_n . This leads us to introduce the subset

$$\mathcal{D}'_n = \{ (\underline{m}^+, \underline{m}^-) \in \mathcal{D}_n \mid m_i^+ m_i^- = 0 \text{ for all } i \ge 1 \}.$$

Lemma 3.3 shows that we have $P_{\underline{m}^+,\underline{m}^-} \neq P_{\underline{n}^+,\underline{n}^-}$ for $(\underline{m}^+,\underline{m}^-) \neq (\underline{n}^+,\underline{n}^-)$ in \mathcal{D}'_n . We reduce to \mathcal{D}'_n as follows. Consider the following map $\phi : \mathcal{D}_n \to \mathcal{D}_n$:

(i) if
$$(\underline{m}^+, \underline{m}^-) \in \mathcal{D}'_n$$
 set $\phi(\underline{m}^+, \underline{m}^-) = (\underline{m}^+, \underline{m}^-)$,

(ii) otherwise there is a smallest $j \geq 1$ with $m_j^+ m_j^- \neq 0$ and we set $\phi(\underline{m}^+, \underline{m}^-) = (\underline{n}^+, \underline{n}^-)$ with $(n_i^+, n_i^-) = (m_i^+, m_i^-)$ for $i \neq j$ or $i \neq 2j$, and with $(n_j^+, n_j^-) = (m_j^+ - 1, n_j^+ - 1)$ and $(n_{2j}^+, n_{2j}^-) = (m_{2j}^+ + 1, m_{2j}^-)$.

It is clear that we have $P_{\phi(\underline{m}^+,\underline{m}^-)} = P_{\underline{m}^+,\underline{m}^-}$ for all $(\underline{m}^+,\underline{m}^-)$ in \mathcal{D}_n , and that for each $m = (\underline{m}^+,\underline{m}^-) \in \mathcal{D}_n$ the sequence $m, \phi(m), \phi^2(m), \ldots$ is eventually constant and equal to some element of \mathcal{D}'_n , that we denote by $\psi(m)$.

Corollary 3.6. Let $\sigma \in O(I_n)$. For all $(\underline{m}^+, \underline{m}^-)$ in \mathcal{D}'_n we have

$$\mathbf{m}_{\sigma \mathbf{W}(\mathbf{D}_n)}(\mathbf{P}_{\underline{m}^+,\underline{m}^-}) = \sum \frac{1}{\mathbf{n}_{\underline{n}^+,\underline{n}^-}}$$

the sum being over all the $(\underline{n}^+, \underline{n}^-)$ in \mathcal{D}_n with $\psi(\underline{n}^+, \underline{n}^-) = (\underline{m}^+, \underline{m}^-)$ and $(-1)^{|\underline{m}^-|} = \mathbf{s}(\sigma)$. We have $\mathbf{m}_{\sigma W(\mathbf{D}_n)}(P) = 0$ for all other P in Car_n .

We have $G(D_4) \simeq S_3$ so it remains to determine $m_{\sigma W(D_4)}$ for the 6 possible classes $\sigma W(D_4)$. A first general reduction is the following lemma:

Lemma 3.7. Let *L* be an integral lattice, as well as elements σ_1, σ_2 in O(L) whose images in G(L) are conjugate. Then we have $m_{\sigma_1W(L)} = m_{\sigma_2W(L)}$.

Proof. Write $\sigma_2 = \gamma \sigma_1 \gamma^{-1} w_0$ with γ in O(L) and w_0 in W(L). For $w \in W(L)$ we have $\det(t - \sigma_2 w) = \det(t - \sigma_1 \gamma^{-1} w_0 w \gamma)$. We conclude as $w \mapsto \gamma^{-1} w_0 w \gamma$ is a bijection of the normal subgroup W(L) of O(L).

In particular, $m_{\sigma W(D_4)}$ is already given by Lemma 3.6 whenever the image of σ in $G(D_4) \simeq S_3$ has order 1 or 2, and does not depend on σ if this image has order 3. There are many ways to determine $m_{\sigma W(D_4)}$ in this latter case. One way is to consider first the set

$$R' = \{v \in D_4, v \cdot v = 4\} = \{\pm 2\epsilon_i \mid i = 1, \dots, 4\} \cup \{\sum_{i=1}^4 \pm \epsilon_i\}$$

and observe that we have $\alpha \cdot x \in 2\mathbb{Z}$ for all $\alpha \in R'$ and $x \in D_4$. In particular, $\frac{1}{\sqrt{2}}R'$ is a root system (of type \mathbf{D}_4) in \mathbb{R}^4 and we have $W(\frac{1}{\sqrt{2}}R') \subset O(D_4)$. The two roots $\alpha = \sqrt{2}\epsilon_1$ and $\beta = \frac{1}{\sqrt{2}}(\epsilon_1 + \epsilon_2 + \epsilon_3 + \epsilon_4)$ are in $\frac{1}{\sqrt{2}}R'$ with $\alpha \cdot \beta = 1$, and the order 3 element

of $O(D_4)$ does not belong to $W(D_4)$. It is trivial to enumerate with a computer the $2^3 4! = 384$ elements of $\sigma_0 W(D_4)$ and to list their characteristic polynomials. We obtain:

Corollary 3.8. Let σ be an element of order 3 in $G(D_4)$. The $m_{\sigma W(D_4)}(P)$ with $P \in Car_4$ are given by Table¹² 9.

In fact, the reasoning above can be pushed a little further: it turns out that $R'' = R \coprod R'$ is a root system of type \mathbf{F}_4 in \mathbb{R}^4 (not **ADE** of course) and that we have $W(R'') = \langle W(R), W(\frac{1}{\sqrt{2}}R') \rangle = O(D_4)$. But the conjugacy classes of W(S), with S any irreducible root system of exceptional type, have been listed and studied in a conceptual way by Carter in [CAR72], including their characteristic polynomials (see p. 22 & 23 *loc. cit.*). The map $m_{O(D_4)}$ may be deduced in particular from Table 8 of [CAR72]. The map $m_{\sigma_0 W(D_4)}$ follows then from the equality $m_{O(D_4)} = -\frac{1}{3} m_{W(D_4)} + \frac{1}{2} m_{O(I_4)} + \frac{1}{3} m_{\sigma_0 W(D_4)}$.

Remark 3.9. Assume R is an irreducible root system. It follows from (3.2) that two elements of G(R) are conjugate if and only if they have the same order, which is always 1, 2 or 3. In particular, Lemma 3.7 shows that for σ in O(R) the map $m_{\sigma W(R)}$ only depends on the order of σ in G(R).

(E) CASES $R \simeq \mathbf{E}_n$ WITH n = 6, 7 AND 8.

The aforementioned results of Carter also allow to deduce $m_{W(E_n)}$ for n = 6,7 and 8 (using Tables 9, 10 and 11 *loc. cit*). Alternatively, and as a useful check, these masses can also be computed directly using a variant of the Algorithm A explained in §1.4. Indeed, choosing a positive system $R^+ \subset R$, we may view W(R) as the subgroup of O(V) generated by the *n* reflections s_{α} , with α a simple root in R^+ . As W(R) acts faithfully and transitively on R, it is also the subgroup of the permutation group of R generated by these *n* permutations s_{α} , with |R| = 72 (case n = 6), |R| = 126 (case n = 7) or |R| = 240 (case n = 8). Applying GAP's ConjugacyClasses algorithm to this permutation group, we obtain representatives and cardinalities of the conjugacy classes of W(R), and it only remains to compute their characteristic polynomials. All in all, these computations only take a few seconds for the computer. Both methods lead to the:

Corollary 3.10. For n = 6, 7 and 8, the $m_{W(E_n)}(P)$ with $P \in Car_n$ are given by Tables 10, 11 and 12.

Note that for n = 7,8 we have $O(E_n) = W(E_n)$ (no non trivial diagram automorphism). For n = 6, we have $O(E_6) = W(E_6) \coprod -W(E_6)$, but the map $m_{-W(E_6)}$ is deduced from $m_{W(E_6)}$ using Lemma 3.5.

¹²The notation for polynomials used in Table 9 is explained on p. 10.

3.11. The non irreducible case

Assume now R is a non necessarily irreducible **ADE** root system in V, set L = Q(R) and fix σ in O(R). Our aim is to give a formula for $m_{\sigma W(R)}$. Write R as the disjoint union of its irreducible components $R = \prod_{i \in I} R_i$. We have

$$L = \bigoplus_{i \in I}^{\perp} \mathcal{Q}(R_i) \text{ and } \mathcal{W}(R) = \prod_{i \in I} \mathcal{W}(R_i).$$

The element σ induces a permutation of the set $\{R_i \mid i \in I\}$ of irreducible components of R. We write $\sigma = c_1 c_2 \cdots c_r$ the cycle decomposition of this permutation. For each $j = 1, \ldots, r$, we choose an irreducible component S_j of R in the support of c_j , denote by $s_j = \dim Q(S_j)$ the rank of S_j and by l_j the length of the cycle c_j . For each j we have $\sigma^{l_j}(S_j) = S_j$ and we denote by τ_j the restriction of σ^{l_j} to $Q(S_j)$; so τ_j is an element of $O(S_j)$.

Proposition 3.12. In the setting above, we have for all P in Car_n

$$\mathbf{m}_{\sigma \mathbf{W}(R)}(P) = \sum_{(P_1,\dots,P_r)} \prod_{j=1}^r \mathbf{m}_{\tau_j \mathbf{W}(S_j)}(P_j)$$

summing over all $(P_1, \ldots, P_r) \in \operatorname{Car}_{s_1} \times \cdots \times \operatorname{Car}_{s_r}$ with $\prod_{j=1}^r P_j(t^{l_j}) = P(t)$.

The first ingredient in the proof is the following trivial lemma.

Lemma 3.13. For i = 1, 2, let V_i be an Euclidean space and $\Gamma_i \subset O(V_i)$ a finite subset. Set $V = V_1 \perp V_2$ and view $\Gamma = \Gamma_1 \times \Gamma_2$ as a subset of O(V). For all monic polynomials P in $\mathbb{R}[t]$ of degree dim V we have

$$m_{\Gamma}(P) = \sum_{(P_1, P_2)} m_{\Gamma_1}(P_1) m_{\Gamma_2}(P_2),$$

the sum being over the (P_1, P_2) , with $P_i \in \mathbb{R}[t]$ monic of degree dim V_i , and with $P_1P_2 = P$.

Proof. (of Proposition 3.12) Applying Lemma 3.13, we may and do assume r = 1, *i.e.* that σ permutes transitively the irreducible components of R. In this case, we simply write (S, s, l, τ) instead of (S_1, s_1, l_1, τ_1) . We may also assume that we have $I = \{0, \ldots, l-1\}$ and $R_i = \sigma^i(S)$ for $0 \le i < l$. In particular, we have $W(R) = \prod_{0 \le i < l} \sigma^i W(S) \sigma^{-i}$. Choose a \mathbb{Z} -basis $e = (e_1, \ldots, e_s)$ of Q(S) and consider the following \mathbb{Z} -basis of L:

$$f = (e_1, \ldots, e_s, \sigma(e_1), \ldots, \sigma(e_s), \ldots, \sigma^{l-1}(e_1), \ldots, \sigma^{l-1}(e_s)).$$

For all $w = (\sigma^i w_i \sigma^{-i})_{0 \le i \le l}$ in W(R), the matrix of σw in the basis f is

$$(3.5) \qquad \begin{pmatrix} & & & MW_{l-1} \\ W_0 & & & \\ & W_1 & & \\ & & \ddots & \\ & & & W_{l-2} \end{pmatrix}$$

where W_i is the matrix of $w_i \in W(S)$ in the basis e, and M is the matrix of $\tau = \sigma^l$ in the basis e. By Lemma 3.14 below, it follows that the multiset of polynomials det $(t - \sigma w)$ (counted with their multiplicities) when w varies in W(R), coincides with that of polynomials det $(t^s - \tau w_{l-1}w_{l-2}\cdots w_0)$ when the l-tuple $(w_0, w_1, \ldots, w_{l-1})$ varies in $W(S)^l$. As W(S) is a group, this multiset is also $|W(S)|^{l-1}$ times the multiset of the det $(t^s - \tau w)$ when w varies in W(S), and we are done.

Lemma 3.14. For any $W_0, W_1, \ldots, W_{l-1}$ and M in $M_s(\mathbb{C})$, the characteristic polynomial of the matrix (3.5) of size sl is $det(t^s - MW_{l-1}W_{l-2}\cdots W_0)$.

Proof. By continuity, we may assume $W_i \in \operatorname{GL}_s(\mathbb{C})$ for each *i*. Up to conjugating (3.5) by the diagonal matrix $(1, W_0, W_1W_0, \ldots, W_{l-2} \cdots W_1W_0)$ we may assume $W_0 = W_1 = \cdots = W_{l-1} = \operatorname{id}_s$. But in this case, the entries of (3.5) commute and we conclude by [ING37] and the following well-known fact (applied to $Q = t^s - a$): the characteristic polynomial of the companion matrix of a given monic polynomial Q is the polynomial Q itself. \Box

4. An algorithm computing characteristic masses

4.1. Algorithm B

Consider the following algorithm, which takes as input an integral lattice L in the standard Euclidean space $V = \mathbb{R}^n$:

B1. Compute the root system R = R(L), a positive root system $R^+ \subset R$ and the associate Weyl vector $\rho = \frac{1}{2} \sum_{\alpha \in R^+} \alpha$.

B2. Determine a set \mathcal{G} of generators of the stabilizer $O(L)_{\rho}$ of ρ in O(L).

B3. Compute the set S = S(L) defined on p. 7 and view $O(L)_{\rho}$ as the subgroup of permutations of S generated by G.

B4. Use permutation groups algorithms to determine the sizes $(m_j)_{j \in J}$ and representatives $(\gamma_j)_{j \in J}$ of the conjugacy classes $(c_j)_{j \in J}$ of $O(L)_{\rho}$.

B5. Compute the set Irr(R) of irreducible components of R, the isomorphism class of each such component, and a basis of the orthogonal R^{\perp} of R in V.

B6. For each j in J, compute:

– the characteristic polynomial P_j of γ_j on R^\perp ,

– a set of representatives $\operatorname{Irr}_j \subset \operatorname{Irr}(R)$ of the orbits for the action of γ_j on $\operatorname{Irr}(R)$,

- for each $S \in Irr_j$, the size l_S of its γ_j -orbit and the order $d_S \in \{1, 2, 3\}$ of the permutation $\gamma_i^{l_S}$ of S.

B7. For each $(S, d_S) \in Irr(R) \times \{1, 2, 3\}$ found in B6, compute $m_{\tau W(S)}$ using the results of §3.2, where τ is any element of order d_S in O(S)/W(S) (see Remark 3.9).

B8. Using Proposition 3.12 and step B7, deduce for each j in J the map $m_{\gamma_j W(R)}$.

B9. For each j in J, define $M_j : \operatorname{Car}_n \to \mathbb{Q}$ by setting $M_j(P) = m_{\gamma_j W(R)}(Q)$ if we have $P = QP_j$, and $M_j(P) = 0$ otherwise.

B10. Return $\frac{\sum_{j \in J} m_j M_j}{\sum_{j \in J} m_j}$.

We will say more about each step of this algorithm in §4.3. Recall from (3.3) that we have a semi-direct product $O(L) = W(L) \rtimes O_{\rho}(L)$ and that the restriction to the subspace $U = Q(R) \otimes \mathbb{R}$ of V induces a morphism res : $O(L) \to O(R)$, an isomorphism $W(L) \xrightarrow{\sim} W(R)$ and a morphism $O(L)_{\rho} \to O(R)_{\rho}$. Together with (3.2), this explains why the elements d_S introduced in the step B6 are indeed in $\{1, 2, 3\}$. Moreover, the more correct notation for $\gamma_j W(R)$ in B8 should be $\operatorname{res}(\gamma_j) W(R)$. For j in J, we have $M_j = m_{\gamma_j W(L)}$ as W(L) acts trivially on R^{\perp} . Last but not least, Lemma 3.7 shows

$$\mathbf{m}_{\mathcal{O}(L)} = \frac{\sum_{j \in J} m_j \ \mathbf{m}_{\gamma_j \mathbf{W}(L)}}{\sum_{j \in J} m_j}$$

We have proved the:

Proposition 4.2. Algorithm B returns $m_{O(L)}$.

4.3. Precisions and an implementation

We now discuss more precisely the steps of Algorithm B, as well as some aspects of our implementation: see [CHE19] for the source code and a documentation of the PARI/GP function masses_calc (requiring GAP) that we developped. Its input is a Gram matrix G of the lattice L, which is thus viewed as the lattice \mathbb{Z}^n equipped with the inner product defined by G.

B1. Apply the Fincke-Pohst algorithm [FP85] to G to compute $R \subset \mathbb{Z}^n$. In PARI's implementation, qfminim(G)[3] returns a set $T \subset \mathbb{Z}^n$ with $T \cup -T =$

R consisting of all the elements of R lying in a certain half-space of \mathbb{R}^n : this is a positive system, and we simply choose $R^+ = T$.

B2. Let b be the Gram matrix of the Z-valued bilinear form $(x, y) \mapsto 4(\rho \cdot x)(\rho \cdot y)$ in the canonical basis of \mathbb{Z}^n . Apply the Plesken-Souvignier algorithm [PS97] to the pair of matrices (G, b). This is implemented in PARI/GP as qfauto([G,b])[2] (following Souvignier's C code). It returns a set $\mathcal{G}' \subset \operatorname{GL}_n(\mathbb{Z})$ of generators of the subgroup of O(L) whose elements g satisfy $g\rho = \pm \rho$. For each $g \in \mathcal{G}'$, determine the sign ϵ_g with $g\rho = \epsilon_g \rho$. Define¹³ \mathcal{G} as the set of $\epsilon_a g$ with $g \in \mathcal{G}'$.

B3. Apply recursively the Fincke-Pohst algorithm to find $S \subset \mathbb{Z}^n$, as explained on p. 7. Choose arbitrarily an ordering $\psi : S \xrightarrow{\sim} \{1, \ldots, N\}$. For each g in \mathcal{G} , compute the permutation $\sigma_g = \psi \circ g \circ \psi^{-1}$ in the symmetric group S_N . For later use, also extract a basis $S_0 \subset S$ of \mathbb{R}^n .

B4. Apply GAP's ConjugacyClasses algorithm to the subgroup H of S_N generated by the σ_g with g in \mathcal{G} . It returns a list of representatives $(r_j)_{j\in J}$ of the conjugacy classes of H, as well as their cardinalities $(m_j)_{j\in J}$. Each r_j is a permutation of $\{1, \ldots, N\}$. Using the subset S_0 introduced in B3, compute the matrix $\gamma_j \in \operatorname{GL}_n(\mathbb{Z})$ of the element of $O(L)_\rho$ corresponding to r_j under the natural isomorphism $H \simeq O(L)_\rho$.

B5. Compute first the basis B of the root system R associated to R^+ , using $B = \{\alpha \in R^+ \mid \alpha \cdot \rho = 1\}$. Define a graph with set of vertices B, and with an edge between $b, b' \in B$ if and only if we have $b \cdot b' \neq 0$. Determine the connected components $B = \prod_{i \in I} B_i$ of this graph. For i in I define R_i^+ as the subset of elements α in R^+ with $\alpha \cdot B_i \neq 0$. We have $\operatorname{Irr}(R) = \{R_i \mid i \in I\}$. The isomorphism class of the **ADE** root system $R_i = R_i^+ \cup -R_i^+$ is uniquely determined by its rank $|B_i|$ and its cardinality $2|R_i^+|$.

B6. Use $i \mapsto R_i$ to identify I with $\operatorname{Irr}(R)$. Compute the Weyl vector $\rho_i = \frac{1}{2} \sum_{\alpha \in R_i^+} \alpha$ of R_i for each i in I. Fix $j \in J$. There is a unique permutation τ_j of I such that $\gamma_j(\rho_i) = \rho_{\tau_j(i)}$ for all i in I. Compute τ_j and determine its cycle decomposition.

Steps B7-B10 are theoretically straightforward. Nevertheless, the efficient implementation of these steps depends on the way the maps m_S are represented: see the documentation in [CHE19] for more about the (imperfect) way

¹³An alternative (cleaner) method to compute the stabilizer in O(L) of a given element x of L would be to simply add the condition $v_i \cdot x = b_i \cdot x$ for all $i \leq k$ in the definition of a k-partial automorphism in §3 of [PS97], as well as a similar constraint in the definition of their fingerprint in §4 *loc. cit.* The main advantage of the trick we use is that we do not have to modify the code of the PARI port of Souvignier's program.

we proceed in gp. In the end, masses_calc(G) returns the vector [a, b, c, d] where:

- a is the vector of all [P, m] with P in Car_n and $m = m_{O(L)}(P)$ with $m \neq 0$, - b is the isomorphism class of the root system R(L),

-c is the vector $(c_j)_{j\in J}$ where c_j encodes both the cycle decomposition of γ_j on $\operatorname{Irr}(R)$ and the integers d_S for each S in Irr_j ,

-d is the vector $(d_j)_{j\in J}$ with $d_j = [P_j, m_j/M]$ and $M = \sum_{j\in J} m_j$.

5. The characteristic masses of Niemeier Lattices with roots

The aim of this section is to explain a way to determine the characteristic masses of the Niemeier lattices with roots which does not use the computationally heavy steps B1 and B4 in Algorithm B, by rather determining directly the information of step B6 (and then using of course the elementary results of §3). We will use for this the case by case descriptions of these lattices given by Venkov [VEN80] or Conway and Sloane [CS99, Ch. 16], based on the classical connections between lattices and codes [CS99, EBE02], and study their automorphism groups in slightly more details than what we could find in the literature. To keep this section short, we assume some familiarity with Niemeier lattices and mostly follow the exposition in [CL19, Chap. 2.3] to which we refer for more details.

5.1. Linking modules, Venkov modules and even unimodular lattices

(a) A (quadratic) linking module¹⁴ is a finite abelian group A equipped with a quadratic map $q: A \longrightarrow \mathbb{Q}/\mathbb{Z}$ whose associated symmetric \mathbb{Z} -bilinear map $b(x, y) := q(x + y) - q(x) - q(y), A \times A \longrightarrow \mathbb{Q}/\mathbb{Z}$, is a perfect pairing. The isometry group of A is denoted O(A). If $I \subset A$ is a subgroup, we denote by I^{\perp} the orthogonal of I with respect to b. We say that I is *isotropic* if we have q(I) = 0 (this is usually stronger than $I \subset I^{\perp}$). We say that I is a Lagrangian if it is isotropic and if we have $I = I^{\perp}$ (or equivalently $|A| = |I|^2$).

(b) A Venkov module is a linking module A equipped with a (set theoretic) map qm : $A \to \mathbb{Q}_{\geq 0}$ such that for all $a \in A$ we have qm $(a) \equiv q(a) \mod \mathbb{Z}$, qm(0) = 0 and qm(a) > 0 for $a \neq 0$. Venkov modules form an additive category Ven in an obvious way; in particular we have an obvious notion of orthogonal direct sum of such objects, denoted \oplus . A root of a Venkov module A is an element $a \in A$ such that qm(a) = 1.

¹⁴Such a module is also called a qe-module in [CL19, Chap. 2.3].

(c) Assume L is an even lattice in the Euclidean space V. Recall that we set $q(x) = \frac{x \cdot x}{2}$ for $x \in V$. The finite abelian group L^{\sharp}/L , equipped with the well-defined quadratic map (that we shall still denote by q) $L^{\sharp}/L \longrightarrow \mathbb{Q}/\mathbb{Z}$, $x + L \mapsto q(x) \mod \mathbb{Z}$, is a linking module that we shall denote by res L (sometimes also called the *discriminant group* or *glue group* of L). This linking module has a canonical structure of Venkov module defined by

$$\operatorname{qm}(x) = \inf_{y \in x+L} \operatorname{q}(y).$$

Let $\pi : L^{\sharp} \to \operatorname{res} L$ be the canonical projection. The map $I \mapsto \pi^{-1}I$ is a bijection between the set of isotropic subspaces I of res L and the set of even lattices of V containing L. In this bijection, $\pi^{-1}I$ is unimodular if, and only if, I is a Lagrangian. Moreover, we have $\operatorname{R}(\pi^{-1}I) = \operatorname{R}(L)$ if, and only if, I does not contain any root of res L.

(d) We now focus on the case L = Q(R) with R an **ADE** root system in V. In this case $Q(R)^{\sharp}$ is called the *weight* lattice of R and we set res R = res L. The group O(R) naturally acts on res R, with W(R) acting trivially, so we have a morphism $G(R) \to \text{Aut}_{\text{Ven}}(\text{res } R)$. The Venkov module res R is the orthogonal sum of the res S with S an irreducible component of R. Assume now R is irreducible. Canonical representatives for the nonzero elements of res R are given by the so-called *minuscule weights* of R, that we denote by ϖ_i following Bourbaki's conventions [BOU81b] for the indices. A key property is $qm(\varpi_i + Q(R)) = q(\varpi_i)$ (see Table 5.1). We also have $G(R) \xrightarrow{\sim} \text{Aut}_{\text{Ven}}(\text{res } R)$. In particular, the element -id of O(R) is in W(R) if, and only if, res R is a $\mathbb{Z}/2$ -vector space.

R	\mathbf{A}_n	\mathbf{D}_n, n even	$\mathbf{D}_n, n \text{ odd}$	\mathbf{E}_6	\mathbf{E}_7	\mathbf{E}_8
$\operatorname{res} R$	$\mathbb{Z}/(n+1)$	$\mathbb{Z}/2 imes \mathbb{Z}/2$	$\mathbb{Z}/4$	$\mathbb{Z}/3$	$\mathbb{Z}/2$	0
min.wts	$\varpi_i, i = 1, \dots, n$	$\varpi_n, \varpi_1, \varpi_{n-1}$	$\varpi_n, \varpi_1, \varpi_{n-1}$	ϖ_1, ϖ_6	ϖ_7	
class	$i \bmod n+1$	$\omega, 1, \overline{\omega}$	$1,2,3 \bmod 4$	$1,2 \bmod 3$	$1 \bmod 2$	
$^{\rm qm}$	$\frac{i(n+1-i)}{2(n+1)}$	$\frac{n}{8}, \frac{1}{2}, \frac{n}{8}$	$\frac{n}{8}, \frac{1}{2}, \frac{n}{8}$	$\frac{2}{3}, \frac{2}{3}$	$\frac{3}{4}$	

Table 4: The Venkov module res R for R an irreducible **ADE** root system.

Remark 5.2. In the case $R \simeq \mathbf{D}_n$ with n even, some authors (e.g. [CS99]) identify the $\mathbb{Z}/2$ -vector space res R with the finite field $\mathbb{F}_4 = \{0, 1, \omega, \overline{\omega}\}$. Using this identification, the automorphism group of res R, in Ven, is generated by the Frobenius $f(x) = x^2$, as well as $m(x) = \omega x$ for n = 4 (triality).

5.3. The Niemeier lattices with roots

Niemeier and Venkov have shown that $L \mapsto R(L)$ induces a bijection between the isomorphism classes of Niemeier lattices with roots, and the isomorphism classes of equi-Coxeter¹⁵ **ADE** root systems in \mathbb{R}^{24} . Fix such a root system R in \mathbb{R}^{24} . By §5.1, there is thus a unique O(R)-orbit of Lagrangians I in res R containing no root (the "codes"). For any such I, the associated Niemeier lattice with root system R is $L = \pi^{-1}I$, we have $|\operatorname{res} R| = |I|^2$ and G(L) = O(L)/W(R) is the stabilizer of $I \subset \operatorname{res} R$ in G(R). In particular, the conjugacy class of G(L) in G(R) does not depend on the choice of I.

Goal: For each of the 23 possible isomorphism classes of R, determine the G(R)-conjugacy class of the elements of G(L) (with their multiplicity).

This is exactly the information actually needed to apply Proposition 3.12 to each coset $\sigma W(R)$ in O(L). We will use information on G(L) given by Venkov [VEN80] and Conway-Sloane [CS99, Table 16.1] (see also [ER082]), such as their order and a composition series. Note that those G(L) are also exactly the *umbral groups* studied in [CDH14].¹⁶ We may assume that the decomposition of R as a union of its irreducible components has the form¹⁷

$$R = N_1 R_1 N_2 R_2 \dots N_g R_g,$$

with $R_i \not\simeq R_j$ for $i \neq j$, and $N_i \geq 1$ for all *i*. We have natural decompositions

res
$$R = \bigoplus_{i} (\operatorname{res} R_i)^{N_i}$$
, $G(R) = \prod_{i} G(N_i R_i)$ and $G(N_i R_i) = G(R_i) \wr S_{N_i}$.

By (3.2), each $G(N_iR_i)$ is naturally isomorphic either to the symmetric group S_{N_i} , to the hyperoctahedral group H_{N_i} , or to $T_{N_i} := S_3 \wr S_{N_i}$ in the exceptional case $R_i \simeq \mathbf{D}_4$. The natural exact sequence $1 \to \prod_i G(R_i)^{N_i} \to G(R) \to \prod_i S_{N_i} \to 1$ induces an exact sequence $1 \to G_1(L) \to G(L) \to G_2(L) \to 1$. The orders of $G_1(L)$ and $G_2(L)$ are given in [CS99, Table 16.1]. Moreover, the image of $G_2(L)$ in S_{N_i} is always a transitive subgroup for each *i*.

We denote by $\eta \in G(L)$ the class of the element -id of O(L). It is a central element which does not depend on the choice of I, and satisfies $\eta^2 = 1$. Its

¹⁵A root system R is called equi-Coxeter if its irreducible components have the same Coxeter number, then called the Coxeter number of R and denoted h(R). The Coxeter numbers of \mathbf{A}_n , \mathbf{D}_n , \mathbf{E}_6 , \mathbf{E}_7 and \mathbf{E}_8 are respectively n + 1, 2n - 2, 12, 18 and 30.

¹⁶Although we will not use it, as this not the information we need, let us mention that the character tables of umbral groups have been listed in the appendix 2 *loc. cit.* (and computed using GAP).

¹⁷This is a short notation for $\coprod_{i=1}^{g} \coprod_{i=1}^{N_i} R_i$.

image in $G(N_iR_i)$ is trivial if R_i has type \mathbf{A}_1 , \mathbf{D}_{2n} , \mathbf{E}_7 or \mathbf{E}_8 , and equal to the element -1 of $G(N_iR_i) = \mathbf{H}_{N_i}$ otherwise (§2.1 and Table 5.1). An inspection of Table [CS99, Table 16.1] shows that we always have $G_1(L) = \langle \eta \rangle$, except in the case $R \simeq \mathbf{D}_4^6$ for which we have $G_1(L) \simeq \mathbb{Z}/3$ (and $\eta = 1$).

NOTATION: A conjugacy class $C \subset G(R)$ has the form $\prod_i C_i$ where C_i is a conjugacy class in $G_i = G(N_iR_i)$. So C is uniquely determined by the collection (t_i) where t_i is the type of C_i : a partition of N_i in the case $G_i = S_{N_i}$, a couple of partitions as in §2.3 in the case $G_i = H_{N_i}$, and similarly a triple of partitions in the case $G_i = T_{N_i}$. In this last case, and as in §2.3, we use the sequence of symbols $\cdots i^{a_i} i^{b_i} i^{c_i} \cdots$ to denote the conjugacy class whose elements have a cycle decomposition with a_i (resp. b_i , c_i) cycles of length iwhose i-th power has order 1 (resp. 2, 3), with same conventions as *loc.cit*.

We now start the description of the G(R)-conjugacy classes of the elements of G(L). In the non trivial cases, we list their type and give the number of elements of any given type divided by |G(L)| (the *size* of the type):

- $R \simeq \mathbf{D}_{24}$, $\mathbf{D}_{16} \mathbf{E}_8$, \mathbf{A}_{24} , $\mathbf{A}_{17} \mathbf{E}_7$, $\mathbf{A}_{15} \mathbf{D}_9$ and $\mathbf{A}_{11} \mathbf{D}_7 \mathbf{E}_6$. We have $\mathbf{G}_2(L) = 1$, so $\mathbf{G}(L) = \mathbf{G}_1(L) = \langle \eta \rangle$.
- $R \simeq 3\mathbf{E}_8$. We have $G(R) = S_3$ and res R = 0, so $G(L) = G(R) = S_3$.
- $R \simeq 2\mathbf{D}_{12}$. We have $G(R) = H_2$, $G_1(L) = 1$ and $G_2(L) = S_2$. We may take for I the subgroup $\{0, (1, \omega), (\omega, 1), (\overline{\omega}, \overline{\omega})\}$ (note $q(I) = \{0, 2, 3\}$). For this I, G(L) is the natural subgroup S_2 of H_2 .
- $R \simeq \mathbf{D}_{10} 2\mathbf{E}_7$. We have $\mathbf{G}(R) = \mathbf{H}_1 \times \mathbf{H}_2$, $\mathbf{G}_1(L) = 1$ and $\mathbf{G}_2(L) = \mathbf{S}_2$. We may take $I = \{0, (\omega, 1, 0), (\overline{\omega}, 0, 1), (1, 1, 1)\}$ (note qm $(I) = \{0, 2\}$). So $\mathbf{G}(L) \simeq \mathbb{Z}/2$ is generated by the element $(\varepsilon_1, (12))$ of $\mathbf{G}(R)$, whose type is (1, 2).
- $R \simeq 3\mathbf{D}_8$. We have $G(R) = H_3$, $G_1(L) = 1$ and $G_2(L) = S_3$. We may take for I the subgroup generated by the \mathfrak{S}_3 -orbit of $(1, 1, \omega)$ (it contains $(0, \overline{\omega}, \overline{\omega}), (\omega, \omega, \omega)$ and we have |I| = 8 and $qm(I) = \{0, 2, 3\}$). For this I, G(L) is the natural subgroup S_3 of H_3 .
- $R \simeq 2\mathbf{A}_{12}$. We have $G(R) = H_2$, $\eta = -1$ and $G_2(L) = S_2$. We may take $I = \langle a \rangle \simeq \mathbb{Z}/13$ with a = (1,5) (note qm(I) = {0,2,3}). The order 4 element $\sigma = \varepsilon_1(12)$ of H₂ satisfies $\sigma a = -5a$, hence generates G(L). The type of the elements of G(L) are thus 1^2 , 1^2 and 2, with respective size 1/4, 1/4 and 1/2.

- $R \simeq 4\mathbf{E}_6$. We have $G(R) = H_4$, $\eta = -1$ and $G_2(L) = S_4$: G(L) is a central extension of S_4 by $\mathbb{Z}/2$. Let I be the Lagrangian of res $R \simeq (\mathbb{Z}/3)^4$ with $\pi^{-1}I = L$. The stabilizer of I in $O(\operatorname{res} R)$ is the semidirect product of $GL(I) \simeq GL_2(\mathbb{Z}/3)$ and of a $\mathbb{Z}/3$ -vector space. The natural morphism $G(L) \to GL(I)$ is thus injective, hence bijective. In particular, G(L) is not isomorphic to $S_4 \times \mathbb{Z}/2$ and we are in case (ii) (c) of Proposition 2.8: G(L) is H_4 -conjugate to the subgroup of Example 2.2. The H_4 -conjugacy classes of G(L) are thus given by Table 2.
- $R \simeq 4\mathbf{D}_6$. We have $G(R) = H_4$, $G_1(L) = 1$ and $G_2(L) = S_4$. We claim that G(L) does not contain the natural subgroup S_4 of H_4 . Indeed, assume that the Lagrangian I of res $R = (\text{res } D_6)^4$ defining L is stable under S_4 . For any $x = (x_1, x_2, x_3, x_4) \in I$ and any $1 \le i \ne j \le 4$, we have x + (ij)x in I, hence $2qm(x_i + x_j)$ is either 0 or an integer ≥ 2 . This forces $x_i + x_j = 0$ by Table 5.1, hence $|I| \le 4$: a contradiction. By Proposition 2.8, the subgroup $G(L) \subset H_4$ is thus H_4 -conjugate to the subgroup S_4^ϵ , and we are done.
- $R \simeq 2\mathbf{A}_9 \mathbf{D}_6$. We have $G(R) = H_2 \times H_1$, $\eta = (-1, 1)$ and $G_2(L) = S_2 \times S_1$. We may take for I the subgroup generated by the elements $a = (2, 4, 0), b = (5, 0, \omega)$ and $c = (0, 5, \overline{\omega})$, of respective orders 5, 2 and 2 (check qm(I) = {0, 2, 3}). Observe that I is stable by the element $\sigma = (\varepsilon_2(12), -1)$ of G(R): we have $\sigma(a) = (4, -2, 0) = 2a, \sigma(b) = (0, -5, \overline{\omega}) = c$ and $\sigma(c) = (5, 0, \omega) = b$. This shows $G(L) = \langle \sigma \rangle \simeq \mathbb{Z}/4$, the types of its elements being $(1^2, 1), (1^2, 1)$ and (2, 1), with respective size 1/4, 1/4 and 1/2.
- $R \simeq 3\mathbf{A}_8$. We have $G(R) = H_3$, $\eta = -1$ and $G_2(L) = S_3$. By Proposition 2.8, G(L) is H₃-conjugate to the subgroup $\{\pm 1\} \cdot S_3$ of H₃.
- $R \simeq 2\mathbf{A}_7 2\mathbf{D}_5$. We have $G(R) = H_2 \times H_2$, $\eta = (-1, -1)$ and $G_2(L) = S_2 \times S_2$. We may take for I the subgroup generated by the elements a = (1, 1, 1, 2) and b = (1, -1, 2, 1) of order 8 (check qm $(I) = \{0, 2, 3\}$). Note that I is stable under $\sigma_1 = ((12), \varepsilon_2)$ and $\sigma_2 = (\varepsilon_2, (12))$: we have $\sigma_1(a) = a, \sigma_1(b) = (-1, 1, 2, 3) = -b$, and σ_2 exchanges a and b. This shows $G(L) = \langle \sigma_1, \sigma_2 \rangle$ (dihedral of order 8), with types $(1^2, 1^2)$, $(1^2, 1^2)$ of size 1/8, and types (2, 11), (11, 2) and (2, 2) of size 1/4.
- $R \simeq 4\mathbf{A}_6$. We have $G(R) = H_4$, $\eta = -1$ and $|G_2(L)| = 12$. So G(L) is a central extension of $G_2(L) = \text{Alt}_4$ by $\mathbb{Z}/2$. It has an injective morphism to $GL(I) = GL_2(\mathbb{Z}/7)$ (same argument as for $4\mathbf{E}_6$): this is a non split

extension. By Remark 2.9, the types of the elements of G(L) follow thus from Table 2: they are 1^4 , 1^4 , 2^2 , 13 and 13, with respective sizes 1/24, 1/24, 1/4, 1/3 and 1/3.

• $R \simeq 6\mathbf{D}_4$. We have $G(R) = T_6$, $|G_1(L)| = 3$ and $G_2(L) = S_6$. We identify res D_4 with \mathbb{F}_4 as in Remark 5.2. Following Conway and Sloane, I is an *hexacode* in \mathbb{F}_4^6 . By [CS99, §11.2], we may choose for I the \mathbb{F}_4 -vector space generated by the K-orbit of $(\omega, \overline{\omega}, \omega, \overline{\omega}, \omega, \overline{\omega})$, where K is the subgroup of Alt₆ preserving $\{\{1, 2\}, \{3, 4\}, \{5, 6\}\}$. For this choice, G(L)contains K and $G_1(L)$ is generated by the element (m, m, m, m, m, m)of $\operatorname{Aut}_{\mathbb{Z}/2}(\mathbb{F}_4)^6 = (S_3)^6$. Two other elements of G(L) are for instance (f, f, f, f, f, f)(12) and $(1, 1, 1, 1, m^2, m)(123)$ (for the latter, recall that $(\omega, \omega, \overline{\omega}, \overline{\omega}, 1, 1)$ is in I). As K, (12) and (123) generate S_6 , a straightforward computation allows to list the types of the elements of G(L): we obtain Table 5.

type	1^{6}	1^{6}	$1^{4} 2$	$1^2 2^2$	$1^2 2^2$	2^{3}	11^23	3^{2}
size	1/2160	1/1080	1/48	1/48	1/24	1/48	1/18	1/18
type	$1^2 4$	24	24	15	15	6	123	

Table 5: The T_6 -conjugacy classes of the elements of $3.S_6$.

• $R \simeq 4\mathbf{A}_5 \mathbf{D}_4$. We have $\mathbf{G}(R) = \mathbf{H}_4 \times \mathbf{T}_1$, $\eta = (-1, 1)$ and $\mathbf{G}_2(L) = \mathbf{S}_4 \times \mathbf{S}_1$. We identify res \mathbf{D}_4 with \mathbb{F}_4 as in Remark 5.2. The first projection $\mathrm{pr}_1 : \mathbf{G}(L) \to \mathbf{H}_4$ is injective, and its image $\mathbf{H}(L)$ is a central extension of \mathbf{S}_4 by $\mathbb{Z}/2$. By Proposition 2.8, $\mathbf{H}(L)$ is either conjugate to $\{\pm 1\} \times \mathbf{S}_4$ or to the group $\mathrm{GL}_2(\mathbb{Z}/3)$ embedded as in Example 2.2. By [CS99, §11.2], we may take for I the subgroup generated by the σ -orbit of a = (2, 0, 2, 4, 0) and $b = (3, 3, 0, 0, \overline{w})$, where $\sigma = ((2 \ 3 \ 4), m)$. In order to determine $\mathbf{H}(L)$ it is enough to find the unique $\pm v \in \{\pm 1\}^4$ such that $v(12) \in \mathbf{H}(L)$. Note that 2I is the $\mathbb{Z}/3$ -vector space generated by (2, 0, 2, 4, 0) and (2, 4, 0, 2, 0). We deduce $v = \pm(1, 1, 1, -1)$: we have $\mathbf{G}(L) \simeq \mathbf{H}(L) \simeq \mathbf{GL}_2(\mathbb{Z}/3)$. On the other hand, the second projection $\mathrm{pr}_2 : \mathbf{G}(L) \to \mathbf{T}_1$ is trivial on η hence factors through a morphism $\mu : \mathbf{S}_4 \to \mathbf{S}_3$. We have $\sigma \in \mathbf{G}(L)$ and $\mathrm{pr}_2(\sigma) = m$ has order 3: μ is "the" classical surjective morphism from \mathbf{S}_4 to \mathbf{S}_3 . The types of $\mathbf{G}(L)$ are thus immediately deduced from Table 2.

- $R \simeq 6\mathbf{A}_4$. We have $G(R) = H_6$, $\eta = -1$ and $G_2(L)$ is a transitive subgroup of S_6 of order 120, so $G_2(L)$ is isomorphic to S_5 and G(L) is a central extension of S_5 by $\mathbb{Z}/2$ in H_6 . We claim that G(L)does not contain the triple transposition $\tau = (12)(34)(56)$. Indeed, otherwise the Lagrangian I defining L would be invariant by τ . Set $I^{\pm} = \{x \in I, \tau(x) = \pm x\}$. A nonzero element of I^+ has the form (a, a, b, b, c, c) with 2qm(a) + 2qm(b) + 2qm(c) an integer $\neq 1$. This forces $\{\pm a, \pm b, \pm c\} = \mathbb{Z}/5$ since $qm(\operatorname{res} A_4) = \{0, \frac{2}{5}, \frac{3}{5}\}$. But the nondegenerate conic $a^2 + b^2 + c^2 = 0$ in $(\mathbb{Z}/5)^3$ contains all those vectors: we have $\dim_{\mathbb{Z}/5} I^+ \leq 1$. A similar argument shows $\dim_{\mathbb{Z}/5} I^- \leq 1$. This is a contradiction as $I = I^+ \oplus I^-$ has dimension 3, hence the claim. By Proposition 2.10 and the discussion after this proposition, G(L) is H_6 -conjugate to the image of the map (2.2). The type of its elements are thus given by Table 3.
- $R \simeq 8A_3$. We have $G(R) = H_8$, $\eta = -1$ and $|G_2(L)| = 1344$. In this case, $I \subset \operatorname{res} R = (\mathbb{Z}/4)^8$ is a so-called *octacode* [CS99]. The subgroup C := I/2I a Hamming code in res $R \otimes \mathbb{Z}/2 = (\mathbb{Z}/2)^8$ and $G_2(L)$ is the automorphism group of this code. In particular, C is included in the hyperplane H of $(\mathbb{Z}/2)^8$ defined by $\sum_i x_i = 0$ and V := H/C is a hyperplane in the 4-dimensional $\mathbb{Z}/2$ -vector space W := $(\mathbb{Z}/2)^8/C$. The (easy) theory of Hamming codes shows that the map $\iota: \{1, \ldots, 8\} \to W$, sending j to the class of the canonical basis element δ_i of $(\mathbb{Z}/2)^8$, is injective with image an affine hyperplane under V, and identifies $G_2(L)$ with the affine group of $\{1, \ldots, 8\}$ for this affine structure. In particular, $G_2(L)$ is isomorphic to $GA_3(\mathbb{Z}/2) = (\mathbb{Z}/2)^3 \rtimes$ $GL_3(\mathbb{Z}/2)$. To go further we choose some I: following [CS99, Table 16.1] we take the subgroup generated by the c-orbit of the element (3, 2, 0, 0, 1, 0, 1, 1) where c is the 7-cycle (2345678) (we have qm(I) = $\{0, 2, 3, 4\}$). With this choice of I, we have $c \in G(L)$ and checks that $\tau = (346)(587)$ and $\sigma = \varepsilon_3 \varepsilon_6 \varepsilon_7 \varepsilon_8 (23)(4568)$ lie in G(L) as well. The images in $G_2(L)$ of $1, c, c^{-1}, \tau, \sigma$ and σ^2 are representatives of the conjugacy classes of the stabilizer $G_2(L)_1$ of 1 in $\{1, \ldots, 8\}$, with resp. sizes 1/168, 1/7, 1/7, 1/3, 1/4 and 1/8 (recall $G_2(L)_1 \simeq GL_3(\mathbb{Z}/2)$). But c, τ and σ belong to the stabilizer $G(L)_1$ of $1 \in \{\pm 1\}^8$ in G(L): the natural map $G(L) \to G_2(L)$ induces an isomorphism $G(L)_1 \xrightarrow{\sim} G_2(L)$ $G_2(L)_1$. An inspection of C shows that the translation by the class of $\delta_1 - \delta_2$ in V is the element (12)(37)(45)(68) of S₈. One deduces from these information representatives of the conjugacy classes of G(L): their types are gathered in Table 6.

type	18	$1^2 1^2 2^2$	2^{4}	2^{4}	$1^{2}3^{2}$	$1^{2}3^{2}$	1124	4^{2}	4^{2}	26	17	17
size	1/2688	1/32	1/192	1/32	1/12	1/12	1/8	1/16	1/8	1/6	1/7	1/7

Table 6: The H₈-conjugacy classes of the nontrivial elements of $2.\text{GA}_3(\mathbb{Z}/2)$.

• $R \simeq 12\mathbf{A}_2$. We have $G(R) = H_{12}$, $\eta = -1$ and $G_2(L)$ is isomorphic to the Mathieu group M_{12} . The Lagrangian $I \subset \operatorname{res} R = (\mathbb{Z}/3)^{12}$ is a *ternary Golay code*, whose automorphism group G(L) is the central extension of M_{12} by $\mathbb{Z}/2$ denoted $2.M_{12}$ in the ATLAS. We know since Frobenius [FR004, p. 11] the cycle decompositions, and cardinality, of all the conjugacy classes of M_{12} . The inverse image in $2.M_{12}$ of such a class c is the union of one or two conjugacy classes $c' \cup -c'$, the cycle decomposition of c' being the same as that of c except that each cycle of c now has a sign to be determined. It is an amusing exercise¹⁸ to extract these signs from the lines χ_2 and χ_{18} , and from the power maps, of the character table of $2.M_{12}$ in the ATLAS. We obtain Table 7.¹⁹

type	1^{12}	2^6	$1^{4}2^{4}$	$1^{4}2^{4}$	$1^{3}3^{3}$	$1^{3}3^{3}$	3^4	3^{4}	$2^{2}4^{2}$	$1^2 1^2 4^2$
size	1/190080	1/240	1/384	1/384	1/108	1/108	1/72	1/72	1/32	1/32
type	$1^{2}5^{2}$	$1^{2}5^{2}$	6^2	1236	1236	48	1128	210	1 1 1	1 11
size	1/20	1/20	1/12	1/12	1/12	1/8	1/8	1/10	1/11	1/11

Table 7: The H_{12} -conjugacy classes of the nontrivial elements of $2.M_{12}$.

• $R \simeq 24\mathbf{A}_1$. We have $G(R) = S_{24}$ and G(L) is a Mathieu group M_{24} . The cycle decompositions and cardinality of the conjugacy classes of M_{24} are given by Frobenius in [FR004, p. 12-13]: see Table 8.

Comparison with the output of Algorithm B. For each of the 23 root systems R above, we verified that the types and sizes of the G(R)-conjugacy

¹⁸That such an exercise is possible follows from the following fact: if we have an equality of polynomials $\prod_i (t^i - 1)^{a_i} (t^i + 1)^{b_i} = \prod_i (t^i - 1)^{a'_i} (t^i + 1)^{b'_i}$ with $a_i + b_i = a'_i + b'_i$ for each i, then $a_i = a'_i$ and $b_i = b'_i$ for each i (use $t^i + 1 = (t^{2i} - 1)/(t^i - 1)$ and Lemma 3.3).

¹⁹An alternative way to proceed is to use the description of $2.M_{12}$ given by Hall in [HAL62], as the automorphism group of a 12×12 Hadamard matrix (a subgroup of H_{12}). Using the 4 generators given by Hall *loc. cit.*, and applying GAP's ConjugacyClasses algorithm to the permutation group on 24 letters they generate, we confirm Table 7.

type	$1^8 2^8$	2^{12}	$1^{6} 3^{6}$	3^{8}	$2^4 4^4$	$1^4 2^2 4^4$	4^{6}	$1^{4} 5^{4}$	$1^2 2^2 3^2 6^2$	6^{4}
mass	1/21504	1/7680	1/1080	1/504	1/384	1/128	1/96	1/60	1/24	1/24
type	$1^{3}7^{3}$	1^2248^2	$2^2 10^2$	$1^2 11^2$	24612	12^{2}	12714	13515	321	123
mass	1/21	1/16	1/20	1/11	1/12	1/12	1/7	2/15	2/21	2/23

Table 8: The S_{24} -conjugacy classes of the nontrivial elements of M_{24} .

classes of G(L) found are exactly those returned (from scratch, and in a few seconds!) by Algorithm B (components 3 and 4 returned by masses_calc, see §4.3). The natural isomorphism $O(L)_{\rho} \simeq G(L)$ and Algorithm B provide thus a rather useful tool to study the groups G(L).

6. References

- [BOR00] R. Borcherds, Classification of positive definite lattices, Duke Math. J. 105, 525– 567 (2000).
- [BOU81b] N. Bourbaki, Éléments de mathématique, Groupes et algèbres de Lie, Chapitres 4, 5 et 6, Masson, Paris (1981).
- [CAR72] R. W. Carter, Conjugacy classes in the Weyl group, Compositio Math. 25, 1–59 (1972).
- [CHE19] G. Chenevier, Characteristic Masses of Lattices, http://gaetan.chenevier.perso.math.cnrs.fr/charmasses/index.html.
- [CC09] G. Chenevier & L. Clozel, Corps de nombres peu ramifiés et formes automorphes autoduales, Journal of the A.M.S. 22 Vol 2, 467-519 (2009).
- [CL19] G. Chenevier & J. Lannes, Automorphic forms and even unimodular lattices, Ergebnisse der Mathematik und ihrer Grenzgebiete 69, Springer Verlag, xxi + 417 pp. (2019).
- [CR12] G. Chenevier & D. Renard, Level one algebraic cusp form of classical groups of small rank, Mem. Amer. Math. Soc. 1121, 128 p. (2015).
- [CT19a] G. Chenevier & O. Taïbi, Discrete series multiplicities and level 1 algebraic cusp forms, to appear in Publ. Math. I.H.É.S., 62 p.
- [CT19b] G. Chenevier & O. Taïbi, Siegel modular forms of weight 13 and the Leech lattice, arXiv preprint 1907.08781, 30 p. (2019).

- [CDH14] M. Cheng, J. Duncan & J. Harvey, Umbral moonshine and the Niemeier lattices, Research in the Mathematical Sciences 1:3 (2014).
- [COH76] A. Cohen, Finite complex reflection groups, Ann. Sci. É.N.S. 9, 379–436 (1976).
- [COH80] A. Cohen, Finite quaternionic reflection groups, Journal of Algebra 64, 293–324 (1980).
- [CON69a] J. Conway, A group of order 8,315,553,613,086,720,000, Bull. London Math. Soc. 1, 79–88 (1969).
- [CS99] J.H. Conway & N. J. A. Sloane, Sphere packings, lattices and groups, Grundlehren der Math. Wissenschaften 290, Springer-Verlag, New York (1999).
- [CS88] J.H. Conway & N. J. A. Sloane, Low-dimensional lattices. IV. The mass formula, Twenty-three constructions for the Leech lattice, Proc. Roy. Soc. London 419 (1988).
- [DEM14] L. Dembélé, On the computation of algebraic modular forms on compact inner forms of GSp₄, Math. Comp. 83, no. 288, p. 1931-1950 (2014).
- [DUM13] N. Dummigan, A simple trace formula for algebraic modular forms, Experimental Math. 22, 123–131 (2013).
- [EBE02] W. Ebeling, Lattices and Codes, A Course Partially Based on Lectures by F. Hirzebruch, 2nd ed., Adv. Lect. in Math., Vieweg, 203 pp. (2002).
- [ER082] V. A. Erokhin, Groups of automorphisms of 24-dimensional even unimodular lattices. (in Russian), Zap. Naučn. Sem. Leningrad Otdel. Mat. Inst. Steklov. (LOMI) 116 (1982), 68–73, 162.
- [FP85] U. Fincke & M. Pohst, Improved methods for calculating vectors of short lenght in a lattice, including a complexity analysis, Math. of Computation 44, 463–471 (1985).
- [FR004] G. Frobenius, Über die Charaktere der mehrfach transitiven Gruppen, Preussische Akademie der Wissenschaften Berlin 16, 558-571 Reichsdr. (1904).
- [FH91] W. Fulton & J. Harris, Representation theory. A first course, Springer Verlag, GTM 129, xvi + 551 pp. (1991).
- [GAP] The GAP group, GAP Groups, Algorithms, and Programming, Version 4.10.2, https://www.gap-system.org (2019).
- [GV14] M. Greenberg & J. Voight, Lattice methods for algebraic modular forms on classical groups, in Computations with modular forms, eds. Gebhard Boeckle and Gabor Wiese, Contrib. Math. Comput. Sci., vol. 6, Springer, Berlin, 147-179 (2014).
- [HAL62] M. Hall, Note on the Mathieu group M_{12} , Arch. Math. XIII, 334–240 (1962).

- [HUL00] A. Hulpke, Conjugacy classes algorithms in finite permutation groups via homomorphic images, Mathematics of Computation 69, no 232, 1633–1651 (2000).
- [ING37] M. H. Ingraham, A note on determinants, Bull. Amer. Math. Soc. 43, no 8, 579– 580 (1937).
- [KNE57] M. Kneser, Klassenzahlen definiter quadratischer formen, Archiv der Math. 8, 241–250 (1957).
- [KT87] K. Koike & I. Terada, Young-diagrammatic methods for the representation theory of the classical groups of type B_n, C_n, D_n, Journal of Algebra 107, 466–511 (1987).
- [Kos61] B. Kostant, Lie Algebra Cohomology and the Generalized Borel-Weil Theorem, Ann. Math. 74, 320–387 (1961).
- [LP02] J. Lansky & D. Pollack, Hecke algebras and automorphic forms, Compositio Math. 130, 21–48 (2002).
- [LOE10] D. Loeffler, Explicit Calculations of Automorphic Forms for Definite Unitary Groups, LMS Journal of Comp. and Math. 11, 326–342 (2010).
- [N173] H.-V. Niemeier, Definite quadratische Formen der Dimension 24 und Diskriminante 1, J. Number Theory 5, 142–178 (1973).
- [OEIS] OEIS Foundation Inc. (2019), The On-Line Encyclopedia of Integer Sequences, https://oeis.org/A120963.
- [PARI/GP] The PARI Group, PARI/GP version 2.5.0, Bordeaux, 2014, http://pari.math.u-bordeaux.fr/.
- [PS97] W. Plesken & B. Souvignier, Computing isometries of lattices, J. Symbolic Computations 24, 327-334 (1997).
- [SCH1911] J. Schur, Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene linear Substitutionen, J. Reine Angew. Math. 139, 155– 250 (1911).
- [SER70] J.-P. Serre, Cours d'arithmétique, P.U.F., Paris (1970).
- [TAI17] O. Taïbi, Dimensions of spaces of level one automorphic forms for split classical groups using the trace formula, Ann. Sci. E.N.S. 50, no 2, 269–344 (2017).
- [VEN80] B. Venkov, On the classification of integral even unimodular 24-dimensional quadratic forms, Chapter 18 in [CS99].
- [WEN01] R. Wendt, Weyl's character formula for non-connected Lie groups and orbital theory for twisted affine Lie algebras, Journal of Functional Analysis 180, no 1, 31–65 (2001).
- [WEYL46] H. Weyl, The Classical Groups, Princeton University Press xiii + 320 p. (1946).

Appendix A. Irreducible characters of compact orthogonal groups

Let $n \geq 1$ be an integer. We denote by O(n) the isometry group of the standard Euclidean space $V = \mathbb{R}^n$. We know since Weyl that the complex, irreducible, continuous representations of the compact group O(n) are all defined over \mathbb{R} and parameterized in a natural way by the *n*-permissible (integer) partitions λ . In this section, we recall this parameterization and discuss formulas for the irreducible characters due to Weyl and Koike-Terrada.

A. 1 The *n*-permissible partitions

Recall that a partition λ is a non-increasing integer sequence $\lambda_1 \geq \lambda_2 \geq \cdots$ with $\lambda_i \geq 0$ for all $i \geq 1$ and $\lambda_i = 0$ for i big enough. We also say that λ is a partition of the integer $|\lambda| := \sum_i \lambda_i$. The *diagram* of λ is the Young diagram whose *i*-th row has λ_i boxes for each $i \geq 1$. The *dual* of λ is the partition λ^* defined by $\lambda_i^* = |\{j \geq 1 \mid \lambda_j \geq i\}|$ (with "transpose" diagram).

Following Weyl, the partition λ is called *n*-permissible if the first two columns of its diagram contain at most *n* boxes, or equivalently if we have $\lambda_1^* + \lambda_2^* \leq n$. If λ is *n*-permissible, there is a unique *n*-permissible partition μ with $\lambda_i^* = \mu_i^*$ for i > 1 and $\lambda_1^* + \mu_1^* = n$, called the *associate* of λ and denoted $\operatorname{ass}(\lambda)$. The map $\lambda \mapsto \operatorname{ass}(\lambda)$ is an involution of the set of *n*-permissible integer partitions. An partition λ is called *n*-positive if we have $\lambda_1^* \leq n/2$ (hence $\lambda_i = 0$ for

i > n/2). If λ is *n*-admissible but not *n*-positive, then ass (λ) is *n*-positive.

A. 2 Weyl's construction

For any integer $d \ge 0$, we consider following Weyl the kernel $K_d(V)$ of the direct sum of the d(d-1)/2 contraction maps²⁰ $c_{i,j} : V^{\otimes d} \to V^{\otimes (d-2)}$, defined for $1 \le i < j \le d$ by $c_{i,j}(v_1 \otimes v_2 \otimes \cdots \otimes v_d) = (v_i \cdot v_j) v_1 \otimes v_2 \otimes \cdots \otimes \hat{v_i} \otimes \cdots \otimes \hat{v_j} \otimes \cdots \otimes v_d$. This kernel has a natural linear action of $O(n) \times \mathfrak{S}_d$, hence decomposes as

$$\mathcal{K}_d(V) \simeq \bigoplus_{\{\lambda \mid |\lambda| = d\}} \mathcal{K}_\lambda(V) \otimes \mathcal{R}_\lambda$$

where R_{λ} is "the" irreducible representation of \mathfrak{S}_d classically parameterized by λ , and $K_{\lambda}(V)$ is a real representation of O(n). Set $W_{\lambda} = K_{\lambda}(V) \otimes \mathbb{C}$.

²⁰All tensor products are taken over \mathbb{R} in §A. 2.

Weyl shows that W_{λ} is either 0 or an irreducible representation of O(n)[WEYL46, Thm. 5.7.D]. Moreover, W_{λ} is nonzero if and only if λ is *n*permissible [WEYL46, Thm. 5.7.A & C]. Moreover, he shows that $\lambda \to W_{\lambda}$ is a bijection between the set of *n*-permissible partitions and the isomorphism classes of irreducible representations of O(n) [WEYL46, Thm. 5.7.H & 7.9.B]. The element -1_n clearly acts as multiplication by $(-1)^d$ on W_{λ} . Weyl shows

(1.1)
$$W_{\operatorname{ass}(\lambda)} \simeq W_{\lambda} \otimes \det$$

and studies the restriction of W_{λ} to the index two subgroup $SO(n) \subset O(n)$ in Chap. V.9 & VII.9. We may assume λ is *n*-positive. There are two cases: (i) $\lambda \neq \operatorname{ass}(\lambda)$. The restriction of W_{λ} to SO(n) is then irreducible with highest weight $\sum_{i \leq n/2} \lambda_i \varepsilon_i$, using the classical notations of Bourbaki [BOU81b, Pl. IV]. Moreover, the natural action of $O(n)/SO(n) = \mathbb{Z}/2$ on the highest weight lines of W_{λ} is trivial (and non trivial on those of $W_{\operatorname{ass}(\lambda)} \otimes \mathbb{C}$).

(ii) $\lambda = \operatorname{ass}(\lambda)$. This forces $n \equiv 0 \mod 2$ and $\lambda_{n/2} > 0$. The restriction of W_{λ} to $\operatorname{SO}(n)$ is then the sum of the two irreducible representations, conjugate under O(n), with highest weights $\pm \lambda_{n/2} \varepsilon_{n/2} + \sum_{i=1}^{n/2-1} \lambda_i \varepsilon_i$.

A. 3 Character formulas

Weyl gives a determinantal formula for the character of W_{λ} in [WEYL46, Theorem 7.9.A]. Contrary to the standard so-called Weyl character formula, which applies to any connected compact Lie groups, that formula equally applies to elements in any of the two connected components²¹ of O(n). Assume g is in O(n) and write $\det(1-tg)^{-1} = \sum_{i \in \mathbb{Z}} p_i t^i$ in $\mathbb{Z}[[t]]$ (so $p_i = 0$ for i < 0). Weyl shows *loc. cit.* that for any *n*-permissible partitions λ we have

(1.2)
$$\operatorname{Trace}(g; W_{\lambda}) = \det (p_{\lambda_i - i + j} - p_{\lambda_i - i - j})_{1 \le i, j \le \lambda_1^*}.$$

If we write $det(1 + tg) = \sum_{i \in \mathbb{Z}} e_i t^i$ (so $e_i = 0$ for i < 0 or i > n), and set $\delta_1 = 0$ and $\delta_j = 1$ for j > 1, then [KT87, Theorem 2.3.3 (6)] implies

(1.3)
$$\det (p_{\lambda_i - i + j} - p_{\lambda_i - i - j})_{1 \le i, j \le \lambda_1^*} = \det (e_{\lambda_i^* - i + j} + \delta_j e_{\lambda_i^* - i - j + 2})_{1 \le i, j \le \lambda_1},$$

See also the equivalence of (ii) and (iv) in [FH91, Cor. A.46] for a direct alternative proof of this equality.

 $^{^{21}}$ Let us mention that there exists also a variant of the Weyl character formula which applies to the irreducible characters of non connected compact Lie groups: see e.g. [KOS61, WEN01].

Remark A. 4. In the case $\lambda_1=0$, or equivalently $|\lambda| = 0$ or λ is the empty diagram, then W_{λ} is the trivial representation and both determinants above are indeed 1 by convention. Moreover, the formula $e_{n-i} = (\det g)e_i$ for $i \in \mathbb{Z}$ shows that the determinant on the right-hand side of (1.3) is multiplied by det g if λ is replaced by $\operatorname{ass}(\lambda)$ (it amounts to multiply by det g the first line of the matrix inside the determinant), in agreement with Formula (1.1).

Appendix B. An asymptotic formula

Proposition B. 1. Let L be a lattice in the Euclidean space \mathbb{R}^n and λ an *n*-permissible partition with $|\lambda| \equiv 0 \mod 2$. Then we have

$$\dim \mathbf{W}_{\lambda}^{\mathbf{O}(L)} \sim \frac{2}{|\mathbf{O}(L)|} \dim \mathbf{W}_{\lambda}$$

for $\lambda \to \infty$, in the sense that $\lambda_i - \lambda_{i+1} \to +\infty$ for each $1 \le i \le n/2$.

Proof. As we have $\lambda \to \infty$ we may assume λ is positive and $\lambda_{[n/2]} > 0$. Denote by V_{λ} the irreducible constituent of $(W_{\lambda})_{|SO(n)}$ with highest weight $\sum_{i \le n/2} \lambda_i \varepsilon_i$. Set $SO(L) = O(L) \cap SO(n)$. If n is odd, we have $(W_{\lambda})_{|SO(n)} = V_{\lambda}$, $O(L) = \{\pm id\} \times SO(L)$ and $W_{\lambda}^{O(L)} = V_{\lambda}^{SO(L)}$. If n is even, then $(W_{\lambda})_{|SO(n)}$ is the direct sum of V_{λ} and of its outer conjugate V'_{λ} , and W_{λ} is induced from V_{λ} : we have thus $W_{\lambda}^{O(L)} = V_{\lambda}^{SO(L)}$ in the case $O(L) \neq SO(L)$ and $W_{\lambda}^{O(L)} = V_{\lambda}^{SO(L)} \oplus (V'_{\lambda})^{SO(L)}$ otherwise. We conclude from the degenerate form of Weyl's character formula for SO(n) given in [CC09, Prop. 1.9].

Assume now $n \equiv -1, 0, 1 \mod 8$ and set $\mu_n = \sum_{[L] \in \mathbf{X}_n} \frac{1}{|\mathcal{O}(L)|}$. The mass formula of Minkowski-Siegel-Smith asserts that we have $\mu_n = |\frac{\mathbf{B}_{n/2}}{n} \prod_{j=1}^{n/2-1} \frac{\mathbf{B}_{2j}}{4j}|$ for $n \equiv 0 \mod 8$, and $\mu_n = |\prod_{j=1}^{(n-1)/2} \frac{\mathbf{B}_{2j}}{4j}|$ for $n \equiv \pm 1 \mod 8$, where the \mathbf{B}_m are the Bernouilli numbers [CS88].

Corollary B. 2. For $n \equiv -1, 0, 1 \mod 8$, $|\lambda| \equiv 0 \mod 2$ and $\lambda \to \infty$ we have $\dim M_{W_{\lambda}}(O_n) \sim 2 \mu_n \dim W_{\lambda}$.

For instance, in the case n = 24 of main interest here we have $\mu_{24} \approx 8 \cdot 10^{-15}$, quite a small number compared to $|X_{24}| = 24$, and of course we expect dim $M_{W_{\lambda}}(O_{24})$ to be small for small values of λ .

Appendix C. Tables

P	m	P	m	P	m	P	m	P	m	P	m	P	m
3^{2}	1/24	6^{2}	1/24	$1^{2}3$	1/12	$2^{2}6$	1/12	123	1/4	12	1/4	126	1/4

Table 9: The 7 nonzero $m = m_{\sigma W(D_4)}(P)$ for P in Car₄, where σ in $G(D_4)$ has order 3.

P	m	Р	m	P	m	P	m	Р	m	Р	m	Р	m	Р	m	Р	m
1^{6}	1/51840	3^{3}	1/648	$1^{2}3^{2}$	1/108	$12^{3}4$	1/96	123^{2}	1/36	$1^{3}24$	1/32	3 1 2	1/12	$1^{2}5$	1/10	128	1/8
$1^{5}2$	<i>'</i>		'					$1^{2}2^{2}6$							1/10		
$1^{2}2^{4}$	1/1152	$1^{4}2^{2}$	1/192	$1^{3}2^{3}$	1/96	$1^{3}23$	1/36	$2^{2}36$	1/36	$1^2 2^2 4$	1/16	1246	1/12	9	1/9		

Table 10: The 25 nonzero $m = m_{W(E_6)}(P)$ for P in Car_6 .

P	m	P	m	P	m	P	m	P	m	P	m	P	m	P	m	P	m
17	1/2903040	$1^{5}2^{2}$	1/3072	$1^{4}2^{3}$	13/9216	$1^{3}2^{2}6$	1/288	$23^{2}6$	1/144	$1^{3}5$	1/60	$1^{2}246$	1/48	135	1/30	19	1/18
2^{7}	1/2903040	$1^{2}2^{5}$	1/3072	$1^{3}2^{4}$	13/9216	$12^{4}6$	1/288	136^2	1/144	$2^{3}10$	1/60	12^246	1/48	2610	1/30	218	1/18
$1^{6}2$	1/46080	$1 3^3$	1/1296	1^424	1/384	$1^2 2 4^2$	1/256	$1^{3}2^{2}3$	1/96	$1^{3}2^{2}4$	7/384	$1^{2}28$	1/32	1312	1/24	$1^2 2 3 6$	1/18
$1 2^{6}$	1/46080	26^{3}	1/1296	$12^{4}4$	1/384	$1 \ 2^2 4^2$	1/256	$1^{2}2^{3}6$	1/96	$1^2 2^3 4$	7/384	148	1/32	2612	1/24	12^236	1/18
$1^{5}3$	1/4320	$1^{3}4^{2}$	1/768	$1^{4}23$	1/288	$1^{3}3^{2}$	1/216	$1^2 2 3^2$	1/72	$1^2 2 3 4$	1/48	$12^{2}8$	1/32	$1^{2}25$	1/20	17	1/14
$2^{5}6$	1/4320	$2^{3}4^{2}$	1/768	$1^{2}2^{3}3$	1/288	$2^{3}6^{2}$	1/216	12^26^2	1/72	12^234	1/48	248	1/32	$12^{2}10$	1/20	214	1/14

Table 11: The 54 nonzero $m = m_{W(E_7)}(P)$ for P in Car₇.

P	m	Р	m	P	m	Р	m	Р	m	P	m	P	m	Р	m
18	1/696729600	$2^{4}4^{2}$	1/18432	$3^{2}6^{2}$	1/1728	$1^2 3 4^2$	1/576	1236^2	1/288	$12^{3}10$	1/120	$2^{2}610$	1/60	$1^{2}7$	1/28
2^{8}	1/696729600	$1^{5}24$	1/15360	$1^3 2 4^2$	1/1536	$1^{3}2^{3}3$	1/576	$1^2 2^2 6^2$	1/288	$1^2 2^2 4^2$	9/1024	39	1/54	127	1/28
$1^{7}2$	1/5806080	$12^{5}4$	1/15360	$12^{3}4^{2}$	1/1536	12^334	1/576	$6^{2}12$	1/288	$1^{2}9$	1/108	618	1/54	1214	1/28
$1 2^{7}$	1/5806080	$1^5 2 3$	1/8640	$1^{4}5$	1/1200	$1^3 2 4 6$	1/576	$1^{3}2^{3}4$	19/4608	$2^{2}18$	1/108	1238	1/48	$2^{2}14$	1/28
$1^{6}3$	1/311040	$12^{5}6$	1/8640	$2^{4}10$	1/1200	$1^{3}2^{3}6$	1/576	8 ²	1/192	$1^2 2^2 3 4$	1/96	12312	1/48	24	1/24
$2^{6}6$	1/311040	$1^{2}3^{3}$	1/7776	$1^4 2^2 3$	1/1152	$2^{2}4^{2}6$	1/576	$1^3 2 3 4$	1/192	$1^2 2^2 4 6$	1/96	1268	1/48	12412	1/24
$1^{6}2^{2}$	1/184320	$2^{2}6^{3}$	1/7776	$1^{2}2^{4}6$	1/1152	$1^3 2 \ 3^2$	1/432	12^346	1/192	$1^{2}2^{2}5$	1/80	12612	1/48	20	1/20
$1^{2}2^{6}$	1/184320	$1^{2}2^{4}3$	1/6912	$2^{2}3^{2}6$	1/864	12^36^2	1/432	$1^2 3 12$	1/144	$1^2 2^2 10$	1/80	1245	1/40	1248	5/64
3^{4}	1/155520	$1^4 2^2 6$	1/6912	$1^2 3 6^2$	1/864	$1^{3}28$	1/384	$1^3 2 3 6$	1/144	$4^{2}12$	1/72	12410	1/40		
6^{4}	1/155520	$1^{4}2^{4}$	37/221184	$1^4 2^2 4$	1/768	$12^{3}8$	1/384	12^336	1/144	12346	1/72	129	1/36		
4^4	1/46080	$1^{4}3^{2}$	1/2592	124^{3}	1/768	$1^2 2^2 3^2$	1/288	$2^{2}612$	1/144	$1^{2}2^{2}8$	1/64	$1\ 2\ 18$	1/36		
$1^{4}4^{2}$	1/18432	123^{3}	1/2592	$1^{2}2^{4}4$	1/768	$3^{2}12$	1/288	$1^{2}48$	1/128	$1^2 3 5$	1/60	$1^2 2^2 3 6$	1/36		
$1^{5}2^{3}$	1/18432	126^3	1/2592	5^{2}	1/600	12^{2}	1/288	$2^{2}48$	1/128	$1\ 2\ 3\ 5$	1/60	15	1/30		
$1^{3}2^{5}$	1/18432	$2^4 6^2$	1/2592	10^{2}	1/600	123^26	1/288	$1^{3}25$	1/120	12610	1/60	30	1/30		

Table 12: The 106 nonzero $m = m_{W(E_8)}(P)$ for P in Car₈.

		Table It	. The f	JU HOHZEI	0 m - 1	IIO(Leech)(I) 101 1	$111 \text{ Oat}_{24}.$			
P	m	Р	m	P	m	Р	m	P	m	Р	m
1^{24}	1/8315553613086720000	$2^{6}3^{4}6^{5}$	1/311040	$3^4 6^4 12^2$	1/4608	$1^4 2^4 4^4 8^2$	1/768	$4^2 8^2 12 24$	1/144	1^23515^2	1/60
2^{24}	1/8315553613086720000	$1^8 2^8 4^4$	1/294912	$3^{3}9^{3}$	1/3888	12^224^2	1/576	$1^4 1 1^2$	1/132	$1^2 3 5 15 30$	1/60
3^{12}	1/2690072985600	$3^4 12^4$	1/276480	$6^{3}18^{3}$	1/3888	$1^4 2^4 3^3 4^2 6 12$	1/576	$2^4 2 2^2$	1/132	$2^{2}6101530$	1/60
6^{12}	1/2690072985600	$6^4 12^4$	1/276480	$1^4 3^2 4^4 12^2$	1/3456	$1^4 2^2 3^3 6^2 12^2$	1/576	1^24816^2	1/128	$1^2 2^2 3 5 6 10 15$	1/60
$1^{16}2^8$	1/178362777600	$4^{4}8^{4}$	1/92160	$1^6 2^6 3^3 6^3$	1/3456	$2^2 3^2 4^2 6 12^3$	1/576	2^24816^2	1/128	$2^{2}61030^{2}$	1/60
$1^8 2^{16}$	1/178362777600	$1^{8}5^{4}$	1/72000	$2^4 4^4 6^2 12^2$	1/3456	$1^2 2^2 3 4^4 6 12^2$	1/576	$7^{2}21$	1/126	$1^2 2^2 3 5 6 10 30$	1/60
4^{12}	1/2012774400	$2^8 10^4$	1/72000	$1^{6}7^{3}$	1/2352	$1^234^26^212^3$	1/576	$14^{2}42$	1/126	$1^2 4^2 7 28$	1/56
$1^{8}4^{8}$	1/743178240	86	1/48384	$2^{6}14^{3}$	1/2352	$1^4 2^4 3^4 6^4$	1/576	$15\ 60$	1/120	$2^{2}4^{2}1428$	1/56
$2^{8}4^{8}$	1/743178240	$1^8 2^4 3^4 6^2$	1/41472	$1^2 2^2 4^2 8^4$	1/2048	$1^2 2^4 3^2 6^3 12^2$	1/576	30 60	1/120	52	1/52
$1^{12}2^{12}$	1/389283840	$1^4 2^8 3^2 6^4$	1/41472	$3^4 15^2$	1/1800	$1^4 2^4 3 4^2 6^3 12$	1/576	$1^4 2^2 7^2 14$	1/112	$1^2 2^2 3 6 8^2 24$	1/48
$1^{12}3^{6}$	1/117573120	7^{4}	1/35280	$6^4 30^2$	1/1800	21^{2}	1/504	$1^{2}2^{4}714^{2}$	1/112	$1^2 2^2 3^2 4 6 12 24$	1/48
$2^{12}6^{6}$	1/117573120	14^{4}	1/35280	$1^{6}2^{4}4^{3}8^{2}$	1/1536	42^{2}	1/504	$1^2 2^2 3 9^2 18$	1/108	$1^2 2^2 3 4 6^2 12 24$	1/48
$1^{6}3^{9}$	1/25194240	$1^4 2^2 3^6 6^3$	1/31104	$1^4 2^6 4^3 8^2$	1/1536	3^269^218	1/432	1^2369^218	1/108	$1^2 2^2 3^2 4^2 6^2 12^2$	1/48
$2^{6}6^{9}$	1/25194240	$1^2 2^4 3^3 6^6$	1/31104	$4^4 12^4$	1/1440	36^2918^2	1/432	$1^2 2^2 6 9 18^2$	1/108	$1^2 3^2 7 21$	1/42
3^86^4	1/19906560	$1^{4}5^{5}$	1/30000	20^{3}	1/1200	$3^2 12 24^2$	1/384	2^236918^2	1/108	$2^{2}6^{2}1442$	1/42
$3^4 6^8$	1/19906560	$2^4 10^5$	1/30000	$4^4 20^2$	1/1200	$6^2 12 24^2$	1/384	$3^2 6^2 12^2 24$	1/96	$20\ 40$	1/40
5^{6}	1/6048000	$5^4 10^2$	1/19200	$1^4 5^3 10^2$	1/1200	$5^{2}15^{2}$	1/360	84	1/84	$1^2 2^2 5^2 10\ 20$	1/40
10^{6}	1/6048000	$5^{2}10^{4}$	1/19200	$1^4 2^4 5^3 10$	1/1200	$10^2 30^2$	1/360	391236	1/72	$1^2 2^2 5 10^2 20$	1/40
$1^{10}2^{6}4^{4}$	1/1474560	$1^6 2^6 4^6$	1/15360	$2^4 5^2 10^3$	1/1200	13^{2}	1/312	$8^{2}24^{2}$	1/72	$1^{2}2^{2}4^{2}51020$	1/40
$1^{6}2^{10}4^{4}$	1/1474560	$1^4 4^2 8^4$	1/12288	$1^4 2^4 5 10^3$	1/1200	26^{2}	1/312	6121836	1/72	39	1/39
$1^4 2^4 4^8$	1/1179648	$2^{4}4^{2}8^{4}$	1/12288	$1^4 3 9^3$	1/972	$1^4 2^4 3^2 4^2 6^2 12$	1/288	35	1/70	78	1/39
$1^{8}3^{8}$	1/1088640	15^{3}	1/10800	$1^2 3^2 9^3$	1/972	$15^2 30$	1/240	70	1/70	56	1/28
$2^{8}6^{8}$	1/1088640	30^{3}	1/10800	$2^4 6 18^3$	1/972	1530^{2}	1/240	$3^{2}33$	1/66	$1^{2}23$	1/23
12^{6}	1/483840	$3^2 6^2 12^4$	1/9216	$2^{2}6^{2}18^{3}$	1/972	$1^4 3^2 5^2 15$	1/180	$6^{2}66$	1/66	$2^{2}46$	1/23
$1^6 2^6 3^5 6$	1/311040	$1^4 2^4 8^4$	1/6144	$5^2 20^2$	1/960	$2^4 6^2 10^2 30$	1/180	$1^2 2^2 8 16^2$	1/64	$1^2 2^2 11 22$	1/22
$1^6 3^5 6^4$	1/311040	$1^2 3^3 4^2 12^3$	1/5184	$10^{2}20^{2}$	1/960	28 ²	1/168	20.60	1/60		
$1^{6}2^{6}36^{5}$	1/311040	$2^{2}4^{2}6^{3}12^{3}$	1/5184	24^{3}	1/864	$1^4 2^4 5^2 10^2$	1/160	$12^{2}60$	1/60		

Table 13: The 160 nonzero $m = m_{O(\text{Leech})}(P)$ for P in Car₂₄.

λ	dim	λ	dim	λ	dim	λ	dim	λ	dim
	24:1	3^{2}	4:0	$3^{3}2^{4}1$	181:0	$3^{4}2^{8}$	148	$3^{6}2^{6}$	276
1^{8}	1:1	$3^{2}1^{2}$	19:0	$3^3 2^4 1^3$	97:0	$3^{5}1$	27:0	$3^{7}1$	174:0
1^{12}	1	$3^{2}2$	3:0	$3^3 2^4 1^5$	1	$3^{5}1^{3}$	94:1	$3^{7}1^{3}$	333:17
2	9:0	$3^{2}21^{2}$	19:0	$3^{3}2^{5}1$	251:1	$3^{5}1^{5}$	20:0	$3^{7}1^{5}$	211
2^{2}	27:0	$3^{2}2^{2}$	15:0	$3^3 2^5 1^3$	120 : 1	$3^{5}1^{7}$	1	$3^{7}21$	512:17
2^{3}	26:0	$3^2 2^2 1^2$	50:0	$3^3 2^6 1$	265:1	$3^{5}21$	140:0	$3^{7}21^{3}$	801 : 342
2^{4}	43:0	$3^{2}2^{3}$	18:0	$3^3 2^6 1^3$	100	$3^{5}21^{3}$	242:1	$3^7 2^2 1$	905:253
$2^{4}1^{8}$	1	$3^2 2^3 1^2$	58:0	$3^{3}2^{7}1$	219:51	$3^{5}21^{5}$	82:1	$3^7 2^2 1^3$	927
2^{5}	35:0	$3^{2}2^{4}$	46:0	$3^{3}2^{8}1$	134	$3^{5}2^{2}1$	308:0	$3^7 2^3 1$	1042:683
$2^{5}1^{4}$	1:1	$3^2 2^4 1^2$	97:0	3^4	28:0	$3^5 2^2 1^3$	417:1	$3^7 2^4 1$	675
2^{6}	67:1	$3^2 2^4 1^6$	1	$3^{4}1^{2}$	28:0	$3^5 2^2 1^5$	87	3^{8}	191:34
$2^{6}1^{4}$	1:1	$3^{2}2^{5}$	48:0	$3^{4}1^{4}$	53:1	$3^{5}2^{3}1$	546:1	$3^{8}1^{2}$	476:137
$2^{6}1^{6}$	1	$3^2 2^5 1^2$	91:0	$3^{4}1^{8}$	1	$3^5 2^3 1^3$	551:111	$3^{8}1^{4}$	530
2^{7}	42:0	$3^{2}2^{6}$	97:0	$3^{4}2$	30:0	$3^{5}2^{4}1$	672:58	$3^{8}2$	327:51
2^{8}	69:1	$3^2 2^6 1^2$	123:1	$3^{4}21^{2}$	80:0	$3^5 2^4 1^3$	525	3^821^2	881:552
$2^{8}1^{4}$	1	$3^2 2^6 1^4$	1	$3^{4}21^{4}$	51:0	$3^{5}2^{5}1$	659:325	$3^{8}2^{2}$	660 : 333
2^{9}	37:0	$3^{2}2^{7}$	70:0	$3^{4}21^{6}$	1:1	$3^{5}2^{6}1$	398	$3^8 2^2 1^2$	1047
2^{10}	48:0	$3^2 2^7 1^2$	74:0	$3^4 2^2$	112:1	3^{6}	36:0	$3^{8}2^{3}$	500:364
2^{11}	11:0	$3^{2}2^{8}$	104:0	$3^4 2^2 1^2$	202:1	$3^{6}1^{2}$	217:1	$3^{8}2^{4}$	346
2^{12}	37	$3^2 2^8 1^2$	86	$3^4 2^2 1^4$	132:2	$3^{6}1^{4}$	180:0	$3^{9}1$	307:133
31	1:0	$3^{2}2^{9}$	39:8	$3^4 2^2 1^6$	1	$3^{6}1^{6}$	91	$3^{9}1^{3}$	496
321	7:0	$3^2 2^{10}$	54	$3^{4}2^{3}$	155:0	$3^{6}2$	79:0	$3^{9}21$	651:491
$32^{2}1$	11:0	$3^{3}1$	8:0	$3^4 2^3 1^2$	291:0	$3^{6}21^{2}$	474:0	$3^9 2^2 1$	542
$32^{3}1$	31:0	$3^{3}1^{3}$	6:0	$3^4 2^3 1^4$	126:1	$3^{6}21^{4}$	367:61	3^{10}	158:121
$32^{4}1$	33:0	$3^{3}21$	25:0	$3^{4}2^{4}$	293:1	$3^{6}2^{2}$	270:0	$3^{10}1^2$	406
$32^{5}1$	56:0	$3^{3}21^{3}$	33:0	$3^4 2^4 1^2$	432:1	$3^{6}2^{2}1^{2}$	902:93	$3^{10}2$	177:160
$32^{6}1$	61:0	$3^{3}2^{2}1$	67:0	$3^4 2^4 1^4$	156	$3^{6}2^{2}1^{4}$	551	$3^{10}2^2$	161
$32^{7}1$	63:0	$3^3 2^2 1^3$	49:0	$3^{4}2^{5}$	270:0	$3^{6}2^{3}$	386:16	$3^{11}1$	93
$32^{8}1$	59:0	$3^3 2^2 1^5$	1:1	$3^4 2^5 1^2$	387 : 75	$3^{6}2^{3}1^{2}$	988:418	3^{12}	74
$32^{8}1^{3}$	1	$3^{3}2^{3}1$	122:0	$3^4 2^6$	380 : 73	$3^{6}2^{4}$	563:197		
$32^{9}1$	53:0	$3^3 2^3 1^3$	102:1	$3^4 2^6 1^2$	362	$3^{6}2^{4}1^{2}$	948		
$32^{10}1$	18	$3^3 2^3 1^5$	1:1	$3^{4}2^{7}$	192:89	$3^{6}2^{5}$	371 : 286		
					42				

Table 14: The nonzero dimensions of $M_{W_{\lambda}}(O_{24})$ for $\lambda_1 \leq 3$.