

COMBINATORICS OF QUASI-HEREDITARY STRUCTURES

MANUEL FLORES, YUTA KIMURA, AND BAPTISTE ROGNERUD

ABSTRACT. A quasi-hereditary algebra is an Artin algebra together with a partial order on its set of isomorphism classes of simple modules which satisfies certain conditions. In this article we investigate all the possible choices that yield to quasi-hereditary structures on a given algebra, in particular we introduce and study what we call the poset of quasi-hereditary structures. Our techniques involve certain quiver decompositions and idempotent reductions. For a path algebra of Dynkin type \mathbb{A} , we provide a full classification of its quasi-hereditary structures. For types \mathbb{D} and \mathbb{E} , we give a counting method for the number of quasi-hereditary structures. In the case of a hereditary incidence algebra, we present a necessary and sufficient condition for its poset of quasi-hereditary structures to be a lattice.

CONTENTS

1. Introduction	1
2. Preliminaries	5
2.1. Adapted posets	5
2.2. Quasi-hereditary algebras	7
2.3. Poset of quasi-hereditary structures	8
3. Quasi-hereditary structures and deconcatenations	10
3.1. Deconcatenations at a sink or a source	10
3.2. Characteristic tilting modules and deconcatenations	12
4. Path algebras of type \mathbb{A}	14
4.1. Path algebra of an equioriented quiver of type \mathbb{A}	14
4.2. Path algebras of type \mathbb{A} : general case	16
5. Path algebras of types \mathbb{D} and \mathbb{E}	17
5.1. Idempotent reduction	17
5.2. Decomposition of the set of quasi-hereditary structures	19
5.3. Counting quasi-hereditary structures	20
6. Lattice of quasi-hereditary structures	22
6.1. Lattice properties and incidence algebras	22
6.2. Quasi-hereditary structures of full subposets	25
6.3. Quasi-hereditary structures of diamond-free posets	29
Acknowledgments	32
References	32

1. INTRODUCTION

Quasi-hereditary algebras were introduced by Scott in [Sco87] as an algebraic axiomatization of the theory of rational representations of semisimple algebraic groups. They were generalized to the concept of highest weight categories soon after in [CPS88] allowing infinitely many simple objects. They form a huge class of algebras and categories which appear in many areas of modern representation theory, including complex semisimple Lie algebras, Schur algebras, algebras of global dimension at most two and many more.

2010 *Mathematics Subject Classification*. Primary 16G20, 16W99, 05E10.

Key words and phrases. Quasi-hereditary algebra, lattice, tilting module.

The first author has been supported by the Consejo Nacional de Ciencia y Tecnología, México, (CONACYT), the Deutscher Akademischer Austauschdienst (DAAD), and the Bielefeld University through the Bielefelder Nachwuchsfonds programme. The second author has been supported by the Alexander von Humboldt Foundation in the framework of an Alexander von Humboldt Professorship endowed by the German Federal Ministry of Education and Research.

A quasi-hereditary algebra is a pair $(A, (I, \triangleleft))$ where A is an Artin algebra and (I, \triangleleft) is a partially ordered set indexing the isomorphism classes of simple A -modules which satisfies additional properties involving the filtrations of the projective modules by the so-called standard modules (see Definition 2.11). In particular, an algebra is *not intrinsically* a quasi-hereditary algebra since the standard modules heavily depend on the partial order (I, \triangleleft) .

In the early examples of highest weight categories and quasi-hereditary algebras, the partial orders were easy to choose because there were related to the classical combinatorics of weights and roots and we already knew that these partial orderings were related to the representation theory of our examples. However, there are instances of quasi-hereditary algebras where there is no natural choice for the partial ordering and even if there is such a natural choice, one may wonder about the other possible orderings.

To our knowledge, there are two known results in this direction. The first, which is due to Dlab and Ringel, asserts that an Artin algebra is hereditary if and only if it is quasi-hereditary for any total ordering on I (see [DR89, Theorem 1] for more details). The second, due to Coulembier, says that if an algebra has a simple preserving duality, then it has at most one ‘quasi-hereditary structure’ (see [Cou19, Theorem 2.1.1] for more details, the notion of quasi-hereditary structure is explained below).

The main objective of this article is to continue the systematic study of all the possible choices of partially ordered sets that yield to quasi-hereditary structures on a given Artin algebra. We start by known and easy remarks that will allow us to better define our objectives, and then give the foundations for our investigations. The organisation of this article can be summarised chronologically as follows.

The first main issue when one tries to classify all the partial orders \triangleleft on I such that $(A, (I, \triangleleft))$ is a quasi-hereditary algebra is that there are too many of them. However, as it has been proved by Conde (see [Con16, Proposition 1.4.12]), that if $(A, (I, \triangleleft))$ is a quasi-hereditary algebra, then the partial order \triangleleft is an *adapted poset to A* in the sense of Dlab and Ringel (see Definition 2.1). The definition of adapted order is rather technical, but it reduces significantly the number of partial orders that we have to consider.

The next step is to realize that we do not want to classify adapted partial orders, but we want to classify equivalence classes for an appropriate equivalence relation. Indeed, as it can be seen in Definition 2.11, the only place where the partial order (I, \triangleleft) is involved is in the construction of the standard modules. As a consequence, if $(A, (I, \triangleleft_1))$ and $(A, (I, \triangleleft_2))$ are two quasi-hereditary algebras with same set of standard modules, it is then natural to say that the partial orders \triangleleft_1 and \triangleleft_2 are *equivalent*, and we write $\triangleleft_1 \sim \triangleleft_2$. We call the equivalence classes of this relation the *quasi-hereditary structures* on the algebra A , and these are exactly what we want to classify. We denote by $\text{qh.str}(A)$ the set of quasi-hereditary structures on the algebra A .

This equivalence relation was introduced by Dlab and Ringel in [DR92] where they proved that any quasi-hereditary structure contains *at least one total ordering*. This key fact is the reason why in most of the references the authors usually assume the ordering on I to be total. However, for our purpose, we should not restrict ourselves to total orderings since one equivalence class may contain *many different* total orderings. For example, when A is a semisimple algebra with n isomorphism classes of simple modules, it has a unique quasi-hereditary structure which contains $n!$ total orderings.

Let $(A, (I, \leq))$ be a quasi-hereditary algebra with set of standard and costandard modules denoted by Δ and ∇ . We define subsets $\text{Dec}(\triangleleft)$ and $\text{Inc}(\triangleleft)$ of I^2 as follows:

$$\text{Dec}(\triangleleft) := \{(i, j) \in I^2 \mid [\Delta(j) : S(i)] \neq 0\}, \quad \text{Inc}(\triangleleft) := \{(i, j) \in I^2 \mid [\nabla(j) : S(i)] \neq 0\}.$$

Then, we show in Lemma 2.8 that the transitive cover \triangleleft_m of the relation $(\text{Dec}(\triangleleft) \cup \text{Inc}(\triangleleft))$ is equivalent to \triangleleft and we prove in Proposition 2.9 that it is the unique minimal element (with respect to the inclusion of relations) in the quasi-hereditary structure containing \triangleleft . These partial orders were already considered by Coulembier under the name of *essential orderings* (see [Cou19, Definition 1.2.5]). Here, we call them *minimal adapted orders*. The relations $\text{Dec}(\triangleleft)$ and $\text{Inc}(\triangleleft)$ are called the *decreasing* and *increasing* relations of \triangleleft . These names have been chosen to highlight the relationship of our work and the work of Châtel, Pilaud and Pons on the weak order of integer posets (see [CPP19]). It turns out that the combinatorics of the quasi-hereditary structures on an equioriented quiver of type \mathbb{A} is the same as their combinatorics of the ‘‘Tamari order element poset’’ (see Section 4). Moreover, in the case of a quiver algebra, these names are particularly relevant since an increasing relation follows the paths of the quiver, and a decreasing one goes against them.

In addition to standard and costandard modules, a quasi-hereditary algebra has a distinguished tilting module called the *characteristic tilting module* (see Proposition 2.16). This leads to the following easy proposition.

Proposition 1.1 (Lemma 2.18). *Let \triangleleft_1 and \triangleleft_2 be two partial orders on I such that $(A, (I, \triangleleft_1))$ and $(A, (I, \triangleleft_2))$ are quasi-hereditary algebras. Then the following statements are equivalent.*

- (1) $\triangleleft_1 \sim \triangleleft_2$.
- (2) $(\triangleleft_1)_m = (\triangleleft_2)_m$.
- (3) $T_1 \cong T_2$, where T_i is the characteristic tilting module of $(A, (I, \triangleleft_i))$ for $i = 1, 2$.

It is now clear that we should see the set of quasi-hereditary structures on A as a subposet of its poset of tilting modules in the sense of Happel and Unger (see [HU05; RS91] for more details).

The poset of tilting modules is most of the time infinite, but the poset of quasi-hereditary structures is always finite since any quasi-hereditary structure contains at least one total order. In Example 2.24 we show that the weak order on the set of permutations on n letters can be realized as the poset of quasi-hereditary structures on the path algebra of a suitable orientation of a complete graph. The Tamari lattices are obtained by considering path algebras of an equioriented orientation of a type \mathbb{A} Dynkin diagram. More precisely, we prove the following result in Section 4.

Theorem 1.2 (Theorem 4.6). *Let $n \in \mathbb{N}$, and A_n be an equioriented quiver of type \mathbb{A} with n vertices, and $\Lambda_n = \mathbf{k}A_n$. Then, there are explicit bijections between*

- (1) *Minimal adapted posets to Λ_n ,*
- (2) *Binary trees with n vertices,*
- (3) *Isomorphism classes of tilting modules over Λ_n .*

Moreover, the poset of quasi-hereditary structures on Λ_n is isomorphic to the Tamari lattice.

This implies that the number of quasi-hereditary structures is given by the n -th Catalan number $c_n = \frac{1}{n+1} \binom{2n}{n}$. Furthermore, any tilting module for this algebra is a characteristic tilting module.

The understanding of the other orientations of a type \mathbb{A} quiver is obtained as a consequence of a more general argument involving sources and sinks of a quiver.

A (iterated) *deconcatenation* of a quiver Q is a disjoint union $Q^1 \sqcup \dots \sqcup Q^\ell$ of full subquivers Q^i of Q satisfying certain properties, see Subsection 3.1 for details. The quivers Q^i are obtained by deconcatenating Q at sources or sinks. Then in Section 3 we prove the following result.

Theorem 1.3 (Theorem 3.7). *Let $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^\ell$ be an iterated deconcatenation of Q . Let A be a factor algebra of $\mathbf{k}Q$ modulo some admissible ideal and $A^i := A/\langle e_u \mid u \in Q_0 \setminus Q_0^i \rangle$. Then we have an isomorphism of posets*

$$\text{qh.str}(A) \longrightarrow \prod_{i=1}^{\ell} \text{qh.str}(A^i).$$

Under the isomorphism, we have an explicit construction of the characteristic tilting A -modules in terms of the characteristic tilting A^i -modules, see Theorem 3.14.

A quiver Q whose underlying graph is of type \mathbb{A} has an iterated deconcatenation $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^\ell$ such that each Q^i is an equioriented quiver of type \mathbb{A} . Therefore as a corollary we obtain a classification of the quasi-hereditary structures on the path algebras of type \mathbb{A} .

Theorem 1.4 (Theorems 4.8, 4.9). *Let Q be a quiver whose underlying graph is of type \mathbb{A} . Let $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^\ell$ be an iterated deconcatenation of Q such that each Q^i is an equioriented quiver of type \mathbb{A}_{n_i} for some $n_i \in \mathbb{Z}_{\geq 1}$. Then we have an isomorphism of posets*

$$\text{qh.str}(\mathbf{k}Q) \longrightarrow \prod_{i=1}^{\ell} \text{qh.str}(\Lambda_{n_i}).$$

Moreover, if \triangleleft is a minimal adapted order to $\mathbf{k}Q$ and $T = \bigoplus_{i \in I} T(i)$ is the characteristic tilting module associated to \triangleleft , then for vertices $i, j \in Q_0$, a simple $\mathbf{k}Q$ -module associated to j is a composition factor of $T(i)$ if and only if $j \triangleleft i$ holds.

In Section 5 we count the number of quasi-hereditary structures on the path algebras of Dynkin types \mathbb{D} and \mathbb{E} . For the quivers of type \mathbb{D} , by Theorem 1.3 and the duality argument of Lemma 2.22, it is enough to consider only two orientations of \mathbb{D}_n , with $n \geq 4$, that is, the quivers D_1 and D_2 defined below.

Theorem 1.5 (Lemmas 5.8, 5.9). *Let $n \geq 3$. Then,*

- (1) $|\text{qh.str}(\mathbf{k}D_1)| = 2c_n - 3c_{n-1}$,
- (2) $|\text{qh.str}(\mathbf{k}D_2)| = 3c_{n-1} - c_{n-2}$,

where c_n is the n -th Catalan number and

$$D_1 = 1 \rightarrow 2 \rightarrow \cdots \rightarrow n-2 \begin{array}{l} \nearrow n-1 \\ \searrow n \end{array}, \quad D_2 = 1 \rightarrow 2 \rightarrow \cdots \rightarrow n-2 \begin{array}{l} \nearrow n-1 \\ \nwarrow n \end{array}.$$

The type \mathbb{E} case is explained in Example 5.10.

So far, in all our examples, we have found a lattice of quasi-hereditary structures: in the complete graph case, we obtained the weak order and for quivers of type \mathbb{A} we obtained products of Tamari lattices. As it can be seen in Proposition 6.11, this is no longer true in the case of affine quivers of type \mathbb{A} . The question of understanding which Artin algebras have a lattice of quasi-hereditary structures is both natural and very intriguing. We have the feeling that this will only happen in a few cases but our progress is rather modest: in Section 6 we prove the following result.

Theorem 1.6 (Theorem 6.1). *Let Q be a finite quiver whose underlying graph is a tree. Then $\text{qh.str}(\mathbf{k}Q)$ is a lattice if and only if Q does not have the following quiver as a subquiver for any $n \geq 4$:*

$$\begin{array}{c} 1 \searrow \\ 2 \nearrow \end{array} \begin{array}{c} 3 \rightarrow 4 \rightarrow \cdots \rightarrow n-1 \\ \nearrow n \\ \searrow n+1 \end{array}.$$

When it is a lattice, we give an explicit description of the partial orders which represent a meet or a join of given two quasi-hereditary structures, see Subsection 6.3. As a corollary of this theorem, we see that the set of quasi-hereditary structures on a path algebra of Dynkin type is always a lattice. Curiously, Theorem 1.6 does not seem to have a nice generalization to the setting of finite acyclic quivers. However, it seems to generalize to the setting of incidence algebras of a finite poset. We denote by Z_n the partially ordered set whose Hasse diagram is a zig-zag orientation of an affine quiver of type \mathbb{A}_n (see diagram 6.2) and we propose the following conjecture.

Conjecture 1.7. Let (P, \leq) be a finite poset. Then, the poset of quasi-hereditary structures on the incidence algebra of (P, \leq) is a lattice if and only if Z_n is not isomorphic to a full subposet of (P, \leq) for any $n \geq 4$.

Actually, the setting of incidence algebras is not only a good setting for a generalization of our result, it also simplifies our arguments. So we prove Theorem 1.6 as a corollary of the following theorem.

Theorem 1.8 (Theorem 6.5). *Let (P, \leq) be a finite poset. We assume that the incidence algebra $A(P)$ of (P, \leq) is hereditary. Then, the poset of quasi-hereditary structures on $A(P)$ is a lattice if and only if Z_n is not isomorphic to a full subposet of (P, \leq) for any $n \geq 4$.*

The relationship between Theorem 1.6 and Theorem 1.8 is explained in Remark 6.9.

Experiments were carried out using the GAP-package QPA [QPA19] and SageMath [Sag19]. The code is available online [FR20].

Notation and conventions. Throughout this paper, \mathbf{k} denotes a field of arbitrary characteristic. A subcategory always means a full subcategory which is closed under isomorphisms. An Artin algebra is an R -algebra A such that R is a commutative Artinian ring and A is finitely generated as an R -module. For an Artin algebra, we usually deal with finitely generated right modules. We denote by $\text{mod } A$ the category of finitely generated right A -modules. For an element or subset X of A , we denote by $\langle X \rangle$ the two-sided ideal of A generated by X . For a quiver Q , let $\mathbf{k}Q$ be the path algebra of Q . For two arrows α, β of Q , if the terminating vertex of α is equal to the starting vertex of β , then $\alpha\beta$ indicates the composite of α with β . An order means a partially ordered set.

2. PRELIMINARIES

In this section, we give some definitions and preliminary results which we use throughout this paper.

2.1. Adapted posets. We first consider the so-called *adapted partial orders* introduced by Dlab and Ringel in [DR92] on the set of isomorphism classes of simple modules over an Artin algebra A .

Let (I, \triangleleft) be a finite partially ordered set with $\{S(i)\}_{i \in I}$ being a complete set of representatives of isomorphism classes of simple A -modules. For $i \in I$, we denote by $P(i)$ a projective cover of $S(i)$ and by $I(i)$ an injective envelope of $S(i)$. If M is an A -module, we denote by $[M : S(i)]$ the Jordan-Hölder multiplicity of $S(i)$ in M .

Let Θ be a class of A -modules which is closed under isomorphisms. We denote by $\mathcal{F}(\Theta)$ the subcategory of $\text{mod } A$ consisting of all A -modules which have a Θ -filtration, that is, an A -module M for which there exists a chain of submodules

$$0 = M_n \subset M_{n-1} \subset \cdots \subset M_1 \subset M_0 = M$$

such that M_i/M_{i+1} belongs to Θ . Let $\Delta(i)$ be the largest quotient of $P(i)$ whose composition factors $S(j)$ are such that $j \triangleleft i$. Similarly let $\nabla(i)$ be the largest submodule of $I(i)$ whose composition factors $S(j)$ are such that $j \triangleleft i$. Set $\Delta = \{\Delta(i)\}_{i \in I}$ and $\nabla = \{\nabla(i)\}_{i \in I}$. The module $\Delta(i)$ (resp. $\nabla(i)$) is called the *standard* (resp. *costandard*) module with *weight* i .

If $M \in \mathcal{F}(\Delta)$, the number of times that $\Delta(i)$ appears in a Δ -filtration of M does not depend on the choice of the filtration. We denote it by $(M : \Delta(i))$. Similarly, the number of times that $\nabla(i)$ appears in $N \in \mathcal{F}(\nabla)$ is independent of the choice of the filtration and we denote it by $(N : \nabla(i))$.

Definition 2.1 (Dlab-Ringel). Let A be an Artin algebra with set of isomorphism classes of simple modules $\{S(i)\}_{i \in I}$. A partial order \triangleleft on I is *adapted* to A if for every A -module M with top $S(i)$ and socle $S(j)$, where i and j are incomparable, there is $k \in I$ with $i \triangleleft k$ and $j \triangleleft k$ and $[M : S(k)] \neq 0$.

Lemma 2.2 (Dlab-Ringel). *Let A be an Artin algebra with set of isomorphism classes of simple modules $\{S(i)\}_{i \in I}$. A partial order \triangleleft on I is adapted to A if and only if for every A -module M with top $S(i)$ and socle $S(j)$, where i and j are incomparable, there is $k \in I$ with $i \triangleleft k$ or $j \triangleleft k$ and $[M : S(k)] \neq 0$.*

Proof. See [DR92] at the bottom of page 3. □

As a nice property of adapted posets, we have the following result.

Lemma 2.3 (Dlab-Ringel). *Let (I, \triangleleft_1) be an adapted poset to A . Let \triangleleft_2 be a refinement of \triangleleft_1 . For $l = 1, 2$ let $\Delta_l(i)$ be the standard module with weight i for the poset \triangleleft_l . Then*

$$\Delta_1(i) = \Delta_2(i) \quad \forall i \in I.$$

Proof. This is stated without proof at the top of page 4 of [DR92]. We sketch the proof for the convenience of the reader. It is clear that there is a surjective map $\psi_j: \Delta_2(j) \rightarrow \Delta_1(j)$. If this is not an isomorphism, we consider i such that $S(i)$ is a composition factor of $\Delta_2(j)$ at the top of the kernel of ψ_j . Then, there is a module M which is a non-split extension of $\Delta_1(j)$ and $S(j)$. Since \triangleleft_1 is adapted to A one of the following holds.

- (1) i and j are comparable.
- (2) They are not comparable and there is a composition factor $S(k)$ of $\Delta_1(j)$ such that $j \triangleleft k$ and $i \triangleleft k$.

Since the composition factors in $\Delta_1(j)$ are smaller than j with respect to \triangleleft_1 , the second condition cannot occur. For the first condition, since $i \triangleleft_2 j$ and \triangleleft_2 is a refinement of \triangleleft_1 the only possibility is that $i \triangleleft_1 j$ and this contradicts the fact that $S(i)$ is not a composition factor of $\Delta_1(j)$. □

By duality we have a similar result for costandard modules.

Definition 2.4. Let A be an Artin algebra with set of isomorphism classes of simple modules $\{S(i)\}_{i \in I}$. Let \triangleleft_1 and \triangleleft_2 be two partial orders on I . Then \triangleleft_1 is *equivalent* to \triangleleft_2 , denoted by $\triangleleft_1 \sim \triangleleft_2$, if $\Delta_1 = \Delta_2$ and $\nabla_1 = \nabla_2$.

This equivalence relation is compatible with the notion of adapted poset in the following sense.

Lemma 2.5. *Let A be an Artin algebra and I be a set indexing the isomorphism classes of simple A -modules. Let \triangleleft_1 be an adapted poset to A . If \triangleleft is equivalent to \triangleleft_1 , then \triangleleft is adapted to A .*

Proof. Let M be an indecomposable module with simple top $S(i)$ and simple socle $S(j)$ with i and j incomparable with respect to \triangleleft . Since M has simple top $S(i)$ it is a quotient of $P(i)$. We denote by $U(i)$ the kernel of the projection from $P(i)$ to $\Delta(i)$. Since i and j are incomparable, the module $S(j)$ is a composition factor of $U(i)$. Then, there is a composition factor $S(k)$ which is at the top of $U(i)$ and which is also a composition factor of M . We denote by N a non-split extension of $S(k)$ and $\Delta(i)$.

Since \triangleleft_1 is an adapted poset we see that $i \triangleleft_1 k$ and by transitivity if $S(e)$ is a composition factor of $\Delta(i)$ then $e \triangleleft_1 k$. So the largest submodule of N cogenerated by $S(k)$ is a submodule of $\nabla_1(k)$. Since $\nabla(k) = \nabla_1(k)$ and $S(i)$ is a composition factor of this module, we see that $i \triangleleft k$ and the result follows from Lemma 2.2 \square

Partial orders on a finite set I can be ordered by inclusion of their sets of relations. This gives a ‘poset of posets’ over I where the minimal element is the equality relation on I and the maximal elements are the total orders. The poset obtained by taking the intersection of the relations of two given posets \triangleleft_1 and \triangleleft_2 is called the *intersection* of \triangleleft_1 and \triangleleft_2 .

Lemma 2.6. *Let A be an Artin algebra with I a set indexing a complete set of isomorphism classes of simple A -modules. Let \triangleleft_1 and \triangleleft_2 be two adapted orders in the same equivalence class. Then the following statements hold.*

- (1) *The intersection of \triangleleft_1 and \triangleleft_2 is an adapted order in the same equivalence class.*
- (2) *In each equivalence class of adapted posets to A there is a unique minimal poset.*

Proof. It is clear that (1) implies (2). Let \triangleleft_1 and \triangleleft_2 be two posets in the same equivalence class. We denote by $\Delta = \Delta_1 = \Delta_2$ the corresponding set of standard modules. We let \triangleleft_{int} be the intersection of \triangleleft_1 and \triangleleft_2 and we denote by Δ_{int} the corresponding set of standard modules.

Let $i \in I$. By definition $\Delta_{int}(i)$ is the largest quotient of $P(i)$ whose composition factors are $S(j)$ such that $j \triangleleft_1 i$ and $j \triangleleft_2 i$. So $\Delta_1(i)$ surjects onto $\Delta_{int}(i)$. If they are not isomorphic, at the top of the kernel there is a simple module $S(j)$ such that $j \triangleleft_1 i$ but j is not smaller than i for \triangleleft_2 . This contradicts $\Delta_1(i) = \Delta_2(i)$. So, we have $\Delta_{int} = \Delta$ and by a dual argument, we see that $\nabla_{int} = \nabla$ and the poset \triangleleft_{int} is equivalent to the posets \triangleleft_1 and \triangleleft_2 . The result follows from Lemma 2.5 \square

Remark 2.7. Lemma 2.6 is used in the codes of the first and last authors [FR20].

For a partial order \triangleleft on I , let Δ be the set of standard A -modules and ∇ the set of costandard A -modules associated to \triangleleft . We define subsets $\text{Dec}(\triangleleft)$ and $\text{Inc}(\triangleleft)$ of I^2 as follows:

$$\text{Dec}(\triangleleft) := \{(i, j) \in I^2 \mid [\Delta(j) : S(i)] \neq 0\}, \quad \text{Inc}(\triangleleft) := \{(i, j) \in I^2 \mid [\nabla(j) : S(i)] \neq 0\}.$$

Clearly, $\text{Dec}(\triangleleft)$ and $\text{Inc}(\triangleleft)$ depend on only the equivalence class of \triangleleft . For $i, j \in I$, we write $i \triangleleft^D j$ if $(i, j) \in \text{Dec}(\triangleleft)$ and write $i \triangleleft^I j$ if $(i, j) \in \text{Inc}(\triangleleft)$. For $i \in I$, we have $i \triangleleft^D i$ and $i \triangleleft^I i$.

For a subset R of I^2 , we denote by R^{tc} the transitive closure of R . Then the following lemma is easy to prove.

Lemma 2.8. *Let $\triangleleft_m = (\text{Dec}(\triangleleft) \cup \text{Inc}(\triangleleft))^{\text{tc}}$.*

- (1) *If $i \triangleleft_m j$, then $i \triangleleft j$ holds.*
- (2) *\triangleleft_m is a partial order on I .*

Proof. By definition, $i \triangleleft^D j$ implies $i \triangleleft j$, and $i \triangleleft^I j$ implies $i \triangleleft j$. Thus the assertions hold. \square

Then we have the following proposition.

Proposition 2.9. *Let A be an Artin algebra and I be a set indexing the isomorphism classes of simple A -modules. For an adapted partial order \triangleleft on I , let $\triangleleft_m = (\text{Dec}(\triangleleft) \cup \text{Inc}(\triangleleft))^{\text{tc}}$.*

- (1) *The partial orders \triangleleft and \triangleleft_m are equivalent.*
- (2) *\triangleleft_m is the unique minimal partial order among partial orders \triangleleft' on I with $\triangleleft \sim \triangleleft'$.*

Proof. (1) We denote by Δ_m the set of standard A -modules associated to \triangleleft_m . Fix $i \in I$. Since all composition factors $S(k)$ of $\Delta_m(i)$ satisfies $k \triangleleft_m i$, hence $k \triangleleft i$ holds. We see that $\Delta_m(i)$ is a quotient

of $\Delta(i)$. Conversely if $S(k)$ is a composition factor of $\Delta(i)$, then by construction we have $k \triangleleft_m i$. Since this holds for all the composition factors of $\Delta(i)$, we see that $\Delta(i)$ is a quotient of $\Delta_m(i)$.

By a dual argument, we see that the set of costandard modules for \triangleleft and \triangleleft_m are equal.

(2) Since $\text{Dec}(\triangleleft) = \text{Dec}(\triangleleft')$ and $\text{Inc}(\triangleleft) = \text{Inc}(\triangleleft')$ hold, the assertion follows from Lemma 2.8. \square

Definition 2.10. Let A be an Artin algebra. A partial order of the form \triangleleft_m for an adapted order \triangleleft to A is called a *minimal adapted order*.

2.2. Quasi-hereditary algebras. Let A be an Artin algebra, and (I, \triangleleft) be a finite partially ordered set indexing the set $\{S(i)\}_{i \in I}$ of isomorphism classes of simple A -modules.

Definition 2.11 (Cline-Parshall-Scott [CPS88]). The pair $(A, (I, \triangleleft))$ is a *quasi-hereditary algebra* if:

- (1) $[\Delta(i) : S(i)] = 1$.
- (2) $P(i) \in \mathcal{F}(\Delta)$.
- (3) $(P(i) : \Delta(i)) = 1$, and $(P(i) : \Delta(j)) \neq 0$ implies $i \triangleleft j$.

In [DR92], Dlab and Ringel restrict themselves to adapted posets in order to work with total orders (since an adapted poset is always equivalent to a total order). Using the next result by Conde, we see that this is not a restriction.

Lemma 2.12 ([Con16]). *If $(A, (I, \triangleleft))$ is a quasi-hereditary algebra, then (I, \triangleleft) is adapted to A .*

Proof. The proof is a part of [Con16, Proposition 1.4.12]. We sketch it for the convenience of the reader. Let M be a module with simple top $S(i)$ and simple socle $S(j)$. Since M is a quotient of $P(i)$, $S(j)$ is a composition factor of $P(i)$. Because the algebra is quasi-hereditary $P(i)$ has a Δ -filtration. So $S(j)$ must be a composition factor of a standard module $\Delta(k)$ with $i \triangleleft k$ by Definition 2.11 (3). If $k = i$ then $j \triangleleft i$. If $k \neq i$ and $S(j)$ is at the top of $\Delta(k)$ we have $i \triangleleft j$. Finally, if $S(j)$ is not at the top of $\Delta(k)$ we have $i \triangleleft k$ and $j \triangleleft k$. \square

We can now considerably simplify the definition of quasi-hereditary algebras.

Proposition 2.13. *Let A be an Artin algebra with an adapted poset (I, \triangleleft) indexing the isomorphism classes of simple A -modules. Then $(A, (I, \triangleleft))$ is a quasi-hereditary algebra if and only if*

- (1) (I, \triangleleft) is adapted to A .
- (2) For every $i \in I$, $[\Delta(i) : S(i)] = 1$.
- (3) $P(i) \in \mathcal{F}(\Delta)$ for $i \in I$.

Proof. See [DR92, Theorem 1]. \square

If an Artin algebra A is hereditary, then any adapted order automatically defines a quasi-hereditary algebra.

Proposition 2.14. *If A is hereditary, then for any adapted order (I, \triangleleft) to A , the pair $(A, (I, \triangleleft))$ is a quasi-hereditary algebra.*

Proof. By Lemma 2.3, we may assume that \triangleleft is a total order. Then the assertion follows from [DR89, Theorem 1]. \square

We have the following well known property of quasi-hereditary algebras.

Lemma 2.15. *Let $(A, (I, \triangleleft))$ be a quasi-hereditary algebra, and assume that i is a maximal element of (I, \triangleleft) . We denote by $\triangleleft|_{I \setminus \{i\}}$ the restriction of \triangleleft on $I \setminus \{i\}$. Then $(A/\langle e_i \rangle, \triangleleft|_{I \setminus \{i\}})$ is a quasi-hereditary algebra such that the standard $A/\langle e_i \rangle$ -module with weight $k \in I \setminus \{i\}$ corresponds to the standard A -module with weight k .*

Proof. Since i is maximal, it is clear that the poset $(I \setminus \{i\}, \triangleleft|_{I \setminus \{i\}})$ is an order ideal. The subcategory of $\text{mod } A$ consisting of all A -modules with composition factors $S(j)$ with $j \in I \setminus \{i\}$ coincides with $\text{mod}(A/\langle e_i \rangle)$. Then the Lemma is a consequence of [CPS88, Theorem 3.5]. \square

For a subcategory \mathcal{C} of $\text{mod } A$, we denote by \mathcal{C}^\perp (${}^\perp \mathcal{C}$, respectively) the full subcategory of $\text{mod } A$ consisting of modules M with $\text{Ext}_A^i(\mathcal{C}, M) = 0$ ($\text{Ext}_A^i(M, \mathcal{C}) = 0$, respectively) for any $i > 0$ and $C \in \mathcal{C}$. For a module M , we denote by $\text{add } M$ the full subcategory of $\text{mod } A$ consisting of direct summands of the direct sum of finitely many copies of M . For simplicity, write $(\text{add } M)^\perp = M^\perp$ and so on.

We recall some properties of quasi-hereditary algebras which we use in this paper. For a quasi-hereditary algebra $(A, (I, \triangleleft))$, we have the following equalities, see [Rin91, Theorems 4, 4*]:

$$(2.1) \quad \mathcal{F}(\Delta) = {}^\perp \mathcal{F}(\nabla), \quad \mathcal{F}(\nabla) = \mathcal{F}(\Delta)^\perp.$$

An A -module T is called a *tilting module* if it satisfies the following three conditions:

- (i) the projective dimension of T is finite,
- (ii) $\text{Ext}_A^i(T, T) = 0$ for all $i > 0$,
- (iii) there exists an exact sequence $0 \rightarrow A \rightarrow T_0 \rightarrow T_1 \rightarrow \cdots \rightarrow T_\ell \rightarrow 0$, with $T_i \in \text{add } T$.

Let M be an A -module and \mathcal{C} be a subcategory of $\text{mod } A$. A *left \mathcal{C} -approximation* of M is a morphism $f: M \rightarrow C$ such that $C \in \mathcal{C}$ and the map $\text{Hom}_A(f, C'): \text{Hom}_A(C, C') \rightarrow \text{Hom}_A(M, C')$ is surjective for any $C' \in \mathcal{C}$. A *right \mathcal{C} -approximation* of M is defined dually.

Proposition 2.16 ([Rin91]). *Let $(A, (I, \triangleleft))$ be a quasi-hereditary algebra. For each $i \in I$, there exists an indecomposable A -module $T(i)$ and short exact sequences*

$$0 \rightarrow \Delta(i) \xrightarrow{f} T(i) \rightarrow X(i) \rightarrow 0, \quad 0 \rightarrow Y(i) \rightarrow T(i) \xrightarrow{g} \nabla(i) \rightarrow 0,$$

where $X(i)$ belongs to $\mathcal{F}(\Delta(j) \mid j \triangleleft i, j \neq i)$ and $Y(i)$ belongs to $\mathcal{F}(\nabla(j) \mid j \triangleleft i, j \neq i)$ such that

- (1) f is a left $\mathcal{F}(\nabla)$ -approximation of $\Delta(i)$.
- (2) g is a right $\mathcal{F}(\Delta)$ -approximation of $\nabla(i)$.
- (3) $T = \bigoplus_{i \in I} T(i)$ is a tilting A -module satisfying $\text{add } T = \mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$.
- (4) $\mathcal{F}(\Delta) = {}^\perp T$ and $\mathcal{F}(\nabla) = T^\perp$ hold.

A tilting module $T = \bigoplus_{i \in I} T(i)$ in Proposition 2.16 is called a *characteristic tilting module* for a quasi-hereditary algebra $(A, (I, \triangleleft))$.

If the standard modules and the costandard modules have small projective or injective dimension, then categories $\mathcal{F}(\Delta)$ and $\mathcal{F}(\nabla)$ have good properties. We only refer a statement about $\mathcal{F}(\Delta)$, the statement for $\mathcal{F}(\nabla)$ is dual.

Lemma 2.17 ([Rin10, Appendix]). *Let $(A, (I, \triangleleft))$ be a quasi-hereditary algebra. Then $\mathcal{F}(\Delta)$ is closed under submodules if and only if the projective dimension of any standard module is at most one.*

2.3. Poset of quasi-hereditary structures. Let I be a set indexing the set of isomorphism classes of simple A -modules. Let \triangleleft_1 and \triangleleft_2 be two partial orders on I . If $(A, (I, \triangleleft_1))$ is a quasi-hereditary algebra and $\triangleleft_2 \sim \triangleleft_1$ then $(A, (I, \triangleleft_2))$ is also a quasi-hereditary algebra since the definition of quasi-hereditary algebra only depends on the set of standard modules. In this case, we can give various characterizations of this equivalence relation.

Lemma 2.18. *Let \triangleleft_1 and \triangleleft_2 be two partial orders on I such that $(A, (I, \triangleleft_1))$ and $(A, (I, \triangleleft_2))$ are two quasi-hereditary algebras. Then the following statements are equivalent.*

- (1) $\triangleleft_1 \sim \triangleleft_2$.
- (2) $\Delta_1 = \Delta_2$.
- (3) $\nabla_1 = \nabla_2$.
- (4) $\mathcal{F}(\Delta_1) = \mathcal{F}(\Delta_2)$.
- (5) $\mathcal{F}(\nabla_1) = \mathcal{F}(\nabla_2)$.
- (6) $T_1 \cong T_2$ where T_i is the characteristic tilting module of $(A, (I, \triangleleft_i))$ for $i = 1, 2$.

Proof. We show (4) implies (2). For each $i \in I$, let $K(i)$ be the sum of the kernels of non-zero surjective maps $P(i) \rightarrow X$ with $X \in \mathcal{F}(\Delta_1)$. Then $\Delta_1(i) = P(i)/K(i)$ holds by the proof of [Rin91, Corollary 4]. We have $\Delta_1(i) = P(i)/K(i) = \Delta_2(i)$ by the assumption. Dually, (5) implies (3).

The other equivalences are induced from equation (2.1) and Proposition 2.16. \square

Definition 2.19. Let A be an Artin algebra with an adapted poset (I, \triangleleft) indexing the isomorphism classes of simple A -modules. The equivalence class of \triangleleft with respect to \sim is a *quasi-hereditary structure* on A provided $(A, (I, \triangleleft))$ is a quasi-hereditary algebra.

We denote by $\text{qh.str}(A)$ the set of all quasi-hereditary structures on A and denote by $[\triangleleft]$ a quasi-hereditary structure represented by \triangleleft .

By Lemma 2.18, the equivalence class of \triangleleft only depends on its characteristic tilting module. It is then natural to order the quasi-hereditary structures in the following way. Let $[\triangleleft_1]$ and $[\triangleleft_2]$ be two quasi-hereditary structures on A with respective set of standard modules Δ_1 and Δ_2 , then we set $[\triangleleft_1] \preceq [\triangleleft_2]$ if $\mathcal{F}(\Delta_2) \subseteq \mathcal{F}(\Delta_1)$. By this ordering, we regard $(\text{qh.str}(A), \preceq)$ as a poset.

As we see in the next lemma, this ordering is induced from a partial ordering on tilting modules [HU05; RS91].

Lemma 2.20. *Let $[\triangleleft_1]$ and $[\triangleleft_2]$ be two quasi-hereditary structures on A and T_i be the characteristic tilting module of (A, \triangleleft_i) for $i = 1, 2$. The following statements are equivalent.*

- (1) $[\triangleleft_1] \preceq [\triangleleft_2]$.
- (2) $\mathcal{F}(\nabla_1) \subseteq \mathcal{F}(\nabla_2)$.
- (3) $T_1^\perp \subseteq T_2^\perp$.

Proof. The assertions follow from $\mathcal{F}(\nabla) = \mathcal{F}(\Delta)^\perp$, $\mathcal{F}(\Delta) = {}^\perp\mathcal{F}(\nabla)$, $\mathcal{F}(\nabla) = T^\perp$ and $\mathcal{F}(\Delta) = {}^\perp T$ for a characteristic tilting module T . \square

Lemma 2.21. *Let $[\triangleleft_1]$ and $[\triangleleft_2]$ be two quasi-hereditary structures on A .*

- (1) *The following are equivalent.*
 - (a) *There is a surjective morphism $\Delta_2(i) \rightarrow \Delta_1(i)$ for all $i \in I$.*
 - (b) $\text{Dec}(\triangleleft_1) \subseteq \text{Dec}(\triangleleft_2)$.
- (2) *The following are equivalent.*
 - (c) *There is an injective morphism $\nabla_2(i) \rightarrow \nabla_1(i)$ for all $i \in I$.*
 - (d) $\text{Inc}(\triangleleft_2) \subseteq \text{Inc}(\triangleleft_1)$.
- (3) *If $[\triangleleft_1] \preceq [\triangleleft_2]$ holds, then the statements in (1) and (2) hold. Conversely, if A is a hereditary algebra, then statements in (1) or (2) imply $[\triangleleft_1] \preceq [\triangleleft_2]$.*

Proof. (1), (2) By the definitions of standard and costandard modules, (a) is equivalent to (b), and (c) is equivalent to (d).

(3) It is clear that $\mathcal{F}(\Delta_2) \subseteq \mathcal{F}(\Delta_1)$ implies (a). Assume that A is hereditary and (a) holds. We prove by induction on (I, \triangleleft_2) that all standard modules in Δ_2 are Δ_1 -filtered. If i is minimal in (I, \triangleleft_2) , then $\Delta_2(i)$ is isomorphic to $S(i)$. Since there is a surjection from $\Delta_2(i)$ to $\Delta_1(i)$ and $\Delta_1(i)$ is non-zero module, the surjection is an isomorphism. For $i \in I$, assume that $\Delta_2(j)$ is Δ_1 -filtered for any j which is strictly smaller than i with respect to \triangleleft_2 . By assumption there is an exact sequence $0 \rightarrow X \rightarrow \Delta_2(i) \rightarrow \Delta_1(i) \rightarrow 0$ for some module X . Since A is hereditary, the category $\mathcal{F}(\Delta_2)$ is closed under submodules. Thus X is $\{\Delta_2(j)\}$ -filtered, where j is strictly smaller than i with respect to \triangleleft_2 . By induction hypothesis, X is Δ_1 -filtered and so is $\Delta_2(i)$. \square

Note that if I is a set indexing the simple A -modules, then I also indexes simple A^{op} -modules. Since A is an Artin R -algebra, there exists a duality between $\text{mod } A$ and $\text{mod } A^{\text{op}}$ induced from $D = \text{Hom}_R(-, E)$, where E is an injective envelop of the direct sum of representatives of all simple R -modules up to isomorphisms. For a partial order \triangleleft on I , we denote by Δ^{op} (by ∇^{op} , respectively) the set of standard (costandard, respectively) A^{op} -modules associated to the partial order \triangleleft . This induces the following lemma.

Lemma 2.22. *Let \triangleleft be a partial order on I . Then the following statements hold.*

- (1) *We have $\Delta^{\text{op}}(i) \cong D\nabla(i)$ and $\nabla^{\text{op}}(i) \cong D\Delta(i)$ for any $i \in I$. In particular, $(A, (I, \triangleleft))$ is a quasi-hereditary algebra if and only if $(A^{\text{op}}, (I, \triangleleft))$ is a quasi-hereditary algebra.*
- (2) *An assignment $\triangleleft \mapsto \triangleleft$ induced an anti-isomorphism between $\text{qh.str}(A)$ and $\text{qh.str}(A^{\text{op}})$.*

Proof. A dual of a simple A -module indexed by i is a simple A^{op} -module indexed by i . Thus the isomorphisms in (1) are obtained from the definition of (co)standard modules. By (1), the assignment induces a bijection from $\text{qh.str}(A)$ to $\text{qh.str}(A^{\text{op}})$. By the isomorphisms in (1), we have $D\mathcal{F}(\Delta) = \mathcal{F}(\nabla^{\text{op}})$. This implies that the map reverses the ordering on quasi-hereditary structures. \square

Example 2.23. Let Q be a finite acyclic quiver and $\mathbf{k}Q$ be the path algebra of Q . Then $\text{qh.str}(\mathbf{k}Q)$ always admits both a unique maximal element and a unique minimal element. In fact, it is easy to construct a total order \triangleleft on Q_0 such that all corresponding standard modules are projective by Lemma 2.15. Then $[\triangleleft]$ is a unique maximal element in $\text{qh.str}(\mathbf{k}Q)$.

Example 2.24. Let K_n be a quiver such that the set of vertices is $I = \{1, 2, \dots, n\}$ and there is a unique arrow from i to j whenever $i > j$. In particular, the underlying graph of K_n is a complete graph. It is easy to see that any adapted order to $\mathbf{k}K_n$ is a total order on I , and two distinct total orders on I induce different quasi-hereditary structures on $\mathbf{k}K_n$.

Let S_n be the permutation group on I . For $v, w \in S_n$, we write $v \leq_S w$ if every pair i, j in I such that $i < j$ and $w^{-1}(i) > w^{-1}(j)$ also satisfies $v^{-1}(i) > v^{-1}(j)$. It is well known that this \leq_S induces a partial order on S_n called the weak (Bruhat) order (we refer to [BB05, Proposition 3.1.3] for the proof and to [BB05, Chapter 3] for historical comments on the weak order.)

We have a bijection from S_n to $\text{qh.str}(\mathbf{k}K_n)$ by $w \mapsto w(1) \triangleleft w(2) \triangleleft \dots \triangleleft w(n)$. By Lemma 2.21, this bijection preserves orderings. Thus we have the following isomorphism of posets

$$(S_n, \leq_S) \cong (\text{qh.str}(\mathbf{k}K_n), \preceq).$$

3. QUASI-HEREDITARY STRUCTURES AND DECONCATENATIONS

3.1. Deconcatenations at a sink or a source. Throughout this section, let Q be a finite connected quiver and v be a sink or a source of Q .

Definition 3.1. A *deconcatenation* of Q at a sink or a source v is a disjoint union $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^\ell$ of full subquivers Q^i of Q satisfying the following properties:

- (1) each Q^i is a connected full subquiver of Q having a vertex v ,
- (2) $Q_0 = (Q_0^1 \setminus \{v\}) \sqcup \dots \sqcup (Q_0^\ell \setminus \{v\}) \sqcup \{v\}$ and $Q_0^i \cap Q_0^j = \{v\}$ hold, and
- (3) there are no arrows between u and w in Q , where $u \in Q_0^i \setminus \{v\}$ and $w \in Q_0^j \setminus \{v\}$ for $1 \leq i \neq j \leq \ell$.

Example 3.2. Let $Q = 1 \rightarrow 2 \leftarrow 3 \leftarrow 4 \rightarrow 5$. Then we have two deconcatenations of Q .

$$(1 \rightarrow 2) \sqcup (2 \leftarrow 3 \leftarrow 4 \rightarrow 5), \quad (1 \rightarrow 2 \leftarrow 3 \leftarrow 4) \sqcup (4 \rightarrow 5).$$

Moreover, the former has a deconcatenation at 4 and the latter has a deconcatenation at 2, and the resulting quiver is the same as follows:

$$(1 \rightarrow 2) \sqcup (2 \leftarrow 3 \leftarrow 4) \sqcup (4 \rightarrow 5).$$

In this section, for a given deconcatenation $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^\ell$ of Q , we compare quasi-hereditary structures on algebras whose Gabriel quivers are Q and Q^i . It is easy to see that if $Q^1 \sqcup Q^2$ is a deconcatenation of Q at a vertex v , and $Q^3 \sqcup Q^4$ is a deconcatenation of Q^2 at the vertex v , then $Q^1 \sqcup Q^3 \sqcup Q^4$ is a deconcatenation of Q . Therefore, we consider a deconcatenation which is the disjoint union of two full subquivers.

Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or a source v . Let A be a factor algebra of $\mathbf{k}Q$ modulo some admissible ideal. For each $\ell = 1, 2$, let

$$A^\ell := \frac{A}{\langle e_u \mid u \in Q_0 \setminus Q_0^\ell \rangle}.$$

Thus we have a surjective morphism of algebras $A \rightarrow A^\ell$, and there exists a fully faithful exact functor $\text{mod } A^\ell \rightarrow \text{mod } A$. By this functor, we regard $\text{mod } A^\ell$ as a full subcategory of $\text{mod } A$. Therefore, an A -module M is an A^ℓ -module if and only if $Me_u = 0$ for any $u \in Q_0 \setminus Q_0^\ell$. For a vertex $i \in Q_0^\ell$, let $P^\ell(i)$, $I^\ell(i)$, $S^\ell(i)$ be the indecomposable projective, indecomposable injective and the simple A^ℓ -module associated to the vertex i , respectively. The following lemma is easy and we omit the proof.

Lemma 3.3. *Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or a source v . Let $\ell = 1, 2$.*

- (1) *For any vertex $i \in Q_0^\ell$, we have $S(i) \cong S^\ell(i)$.*
- (2) *For any $i \in Q_0^\ell \setminus \{v\}$, we have $P(i) \cong P^\ell(i)$ and $I(i) \cong I^\ell(i)$.*
- (3) *If v is a sink, then we have $S(v) \cong P(v) \cong P^\ell(v)$ for $\ell = 1, 2$.*
- (4) *If v is a source, then we have $S(v) \cong I(v) \cong I^\ell(v)$ for $\ell = 1, 2$.*
- (5) *Let M be a non-zero A -module. If both of the top and the socle of M are simple, then one of $M \in \text{mod } A^1$ or $M \in \text{mod } A^2$ holds.*
- (6) *Let $M \in \text{mod } A^\ell$ and $i \in Q_0$. If $[M : S(i)] \neq 0$, then $i \in Q_0^\ell$ holds.*

Let \triangleleft be a partial order on Q_0 . By restricting this order, we have a partial order $\triangleleft|_{Q_0^\ell}$ on Q_0^ℓ for $\ell = 1, 2$. We first compare standard and costandard modules associated to these orders.

Lemma 3.4. *Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or a source v . Let \triangleleft be a partial order on Q_0 and Δ (∇ , respectively) the set of standard (costandard, respectively) A -modules associated to \triangleleft . Let $\ell = 1, 2$. We denote by Δ^ℓ (∇^ℓ , respectively) the set of standard (costandard, respectively) A^ℓ -modules associated to $\triangleleft|_{Q_0^\ell}$. Then we have the following statements.*

- (1) *For any $i \in Q_0^\ell \setminus \{v\}$, we have $\Delta(i) \cong \Delta^\ell(i)$ and $\nabla(i) \cong \nabla^\ell(i)$.*
- (2) *If v is a sink, then we have $S(v) \cong \Delta(v) \cong \Delta^\ell(v)$.*
- (3) *If v is a source, then we have $S(v) \cong \nabla(v) \cong \nabla^\ell(v)$.*
- (4) *If \triangleleft defines a quasi-hereditary structure on A , then $\triangleleft|_{Q_0^\ell}$ defines a quasi-hereditary structure on A^ℓ for each $\ell = 1, 2$.*

Proof. (1) Since $P(i) \cong P^\ell(i)$, a composition factor $S(j)$ of $P(i)$ satisfies $j \in Q_0^\ell$. Since \triangleleft^ℓ is a restriction of \triangleleft , for any $j \in Q_0^\ell$, $j \triangleleft i$ if and only if $j \triangleleft^\ell i$. This implies that $\Delta^\ell(i) \cong \Delta(i)$. Similarly, we have $\nabla^\ell(i) \cong \nabla(i)$. (2) (3) The assertions are clear. (4) Assume that v is a sink. Let $i \in Q_0^\ell$. By (1), $\Delta(i) \cong \Delta^\ell(i)$ holds. Thus we have $[\Delta^\ell(i) : S^\ell(i)] = 1$ by Lemma 3.3. Since any composition factor of $P(i)$ is a simple A^ℓ -module, a Δ -filtration of $P(i)$ in $\text{mod } A$ gives a Δ^ℓ -filtration of $P^\ell(i)$ in $\text{mod } A^\ell$. Clearly, this filtration satisfies the axiom (3) of Definition 2.11. Thus $\triangleleft|_{Q_0^\ell}$ defines a quasi-hereditary structure on A^ℓ . The assertion also holds in the case where v is a source by a similar argument. \square

Set $\bar{1} = 2$ and $\bar{2} = 1$. Next we construct a partial order on Q_0 from partial orders on Q_0^ℓ . Let \triangleleft^ℓ be partial orders on Q_0^ℓ for $\ell = 1, 2$. Then we have a partial order $\triangleleft = \triangleleft(\triangleleft^1, \triangleleft^2)$ on Q_0 as follows: for $i, j \in Q_0$, $i \triangleleft j$ if one of the following two statements holds:

- (1) $i, j \in Q_0^\ell$ and $i \triangleleft^\ell j$ holds for some ℓ ,
- (2) $i \in Q_0^{\bar{\ell}}, j \in Q_0^{\bar{\ell}}$, $i \triangleleft^\ell v$ and $v \triangleleft^\ell j$ hold.

Lemma 3.5. *Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or a source v . Let \triangleleft^ℓ be a partial order on Q_0^ℓ and Δ^ℓ (∇^ℓ , respectively) the set of standard (costandard, respectively) A^ℓ -modules associated to \triangleleft^ℓ for $\ell = 1, 2$. We denote by Δ (∇ , respectively) the set of standard (costandard, respectively) A -modules associated to $\triangleleft = \triangleleft(\triangleleft^1, \triangleleft^2)$. Let $\ell = 1, 2$. Then we have the following statements.*

- (1) *For any $i \in Q_0^\ell \setminus \{v\}$, we have $\Delta(i) \cong \Delta^\ell(i)$ and $\nabla(i) \cong \nabla^\ell(i)$.*
- (2) *If v is a sink, then we have $S(v) \cong \Delta(v) \cong \Delta^\ell(v)$.*
- (3) *If v is a source, then we have $S(v) \cong \nabla(v) \cong \nabla^\ell(v)$.*
- (4) *If \triangleleft^ℓ defines a quasi-hereditary structure on A^ℓ for both $\ell = 1, 2$, then $\triangleleft = \triangleleft(\triangleleft^1, \triangleleft^2)$ defines a quasi-hereditary structure on A .*

Proof. (1) Since $P(i) \cong P^\ell(i)$, a composition factor $S(j)$ of $P(i)$ satisfies $j \in Q_0^\ell$. Then by the definition of \triangleleft , we have that, for a composition factor $S(j)$ of $P(i)$, $j \triangleleft i$ holds if and only if $j \triangleleft^\ell i$ holds. This implies the assertion. This implies that $\Delta(i) \cong \Delta^\ell(i)$. (2) (3) The assertions are clear. (4) Assume that v is a sink. Let $i \in Q_0^\ell$. By (1), we have $\Delta(i) \cong \Delta^\ell(i)$. In particular, we have $[\Delta(i) : S(i)] = 1$. By Lemma 3.3, $P(i) \cong P^\ell(i)$ holds for $i \in Q_0^\ell$. Since \triangleleft^ℓ defines a quasi-hereditary structure on A^ℓ , $P(i)$ has a Δ^ℓ -filtration. Since the number $(P(i) : \Delta(j))$ does not depend on the choice of the filtration, the axiom (3) of Definition 2.11 is satisfied. If v is a source, then the dual argument works. \square

Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or a source v . Let A be a factor algebra of $\mathbf{k}Q$ modulo some admissible ideal. By Lemmas 3.4 and 3.5, we have the following map.

$$\Phi: \text{qh.str}(A) \longrightarrow \text{qh.str}(A^1) \times \text{qh.str}(A^2), \quad [\triangleleft] \mapsto \left([\triangleleft|_{Q_0^1}], [\triangleleft|_{Q_0^2}] \right).$$

We also have an inverse map

$$\Psi: \text{qh.str}(A^1) \times \text{qh.str}(A^2) \longrightarrow \text{qh.str}(A), \quad ([\triangleleft^1], [\triangleleft^2]) \mapsto [\triangleleft(\triangleleft^1, \triangleleft^2)].$$

For two posets (A, \leq_A) , (B, \leq_B) and $(a_1, b_1), (a_2, b_2) \in A \times B$, we write $(a_1, b_1) \leq (a_2, b_2)$ if $a_1 \leq_A a_2$ and $b_1 \leq_B b_2$ hold. Then $(A \times B, \leq)$ is a poset, called the *product poset*.

Proposition 3.6. *The map Φ is an isomorphism of posets and its inverse is Ψ .*

Proof. Consider the product poset on $\text{qh.str}(A^1) \times \text{qh.str}(A^2)$. Then the assertion directly follows from Lemmas 2.18, 3.4 and 3.5, since (co)standard modules are preserved by the maps. \square

Let $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^\ell$ be a deconcatenation of Q at a vertex v . If $Q^{\ell+1} \sqcup \dots \sqcup Q^m$ is a deconcatenation of Q^ℓ at a vertex u , then we have a disjoint union $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^{\ell-1} \sqcup Q^{\ell+1} \sqcup \dots \sqcup Q^m$ of full subquivers of Q , and so on for each connected quiver Q^i . We call a disjoint union $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^\ell$ of full subquivers of Q obtained by iterated operations as above an *iterated deconcatenation* of Q .

Then we have the following theorem.

Theorem 3.7. *Let $Q^1 \sqcup Q^2 \sqcup \dots \sqcup Q^\ell$ be an iterated deconcatenation of Q at sink or source vertices. Let A be a factor algebra of $\mathbf{k}Q$ modulo some admissible ideal and $A^i := A/\langle e_u \mid u \in Q_0 \setminus Q_0^i \rangle$. Then we have an isomorphism of posets*

$$\mathrm{qh.str}(A) \longrightarrow \prod_{i=1}^{\ell} \mathrm{qh.str}(A^i),$$

which is given by $[\triangleleft] \mapsto \left([\triangleleft|_{Q_0^i}]\right)_{i=1}^{\ell}$.

Proof. By applying Proposition 3.6 iteratively, we have the assertion. \square

Example 3.8. Let $Q = 1 \rightarrow 2 \leftarrow 3 \leftarrow 4 \rightarrow 5$. We have an isomorphism of posets

$$\mathrm{qh.str}(\mathbf{k}Q) \longrightarrow \mathrm{qh.str}(\Lambda_2) \times \mathrm{qh.str}(\Lambda_3) \times \mathrm{qh.str}(\Lambda_2),$$

where Λ_n is the path algebra of an equioriented quiver of type \mathbb{A}_n . We study $\mathrm{qh.str}(\Lambda_n)$ precisely in Section 4.

We use the following lemma later.

Lemma 3.9. *Let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink or a source v . Let A be a factor algebra of $\mathbf{k}Q$ modulo some admissible ideal. Let \triangleleft ($\triangleleft^1, \triangleleft^2$ respectively) be a partial order on Q_0 (Q_0^1, Q_0^2 , respectively) defining a quasi-hereditary structure on A (A^1, A^2 , respectively). Then the following statements hold.*

- (1) *If \triangleleft is a minimal adapted order, then both $\triangleleft|_{Q_0^1}$ and $\triangleleft|_{Q_0^2}$ are minimal adapted orders.*
- (2) *If \triangleleft^1 and \triangleleft^2 are minimal adapted orders, then $\triangleleft(\triangleleft^1, \triangleleft^2)$ is a minimal adapted order.*

In particular, if \triangleleft is minimal, then $\triangleleft = \triangleleft(\triangleleft|_{Q_0^1}, \triangleleft|_{Q_0^2})$ holds.

Proof. We show only (1) and the last assertion. The assertion (2) is shown similarly. We show that $\triangleleft_1 := \triangleleft|_{Q_0^1}$ is a minimal adapted order. Let \triangleleft'_1 be a partial order on Q_0^1 such that $\triangleleft'_1 \sim \triangleleft_1$. Let $i, j \in Q_0^1$ and assume that $i \triangleleft'_1 j$ holds. This implies that $i \triangleleft j$. By Proposition 3.6, we have $\triangleleft' := \triangleleft(\triangleleft'_1, \triangleleft|_{Q_0^2}) \sim \triangleleft$. Since \triangleleft is minimal, $i \triangleleft' j$ holds. By the definition of \triangleleft' , we have $i \triangleleft'_1 j$. The last assertion follows from a uniqueness of a minimal adapted order. \square

3.2. Characteristic tilting modules and deconcatenations. In this subsection, we compare characteristic tilting modules via deconcatenations. Throughout this subsection let $Q^1 \sqcup Q^2$ be a deconcatenation of Q at a sink v . We denote by A a factor algebra of $\mathbf{k}Q$ modulo some admissible ideal. Let $A^\ell = A/\langle e_u \mid u \in Q_0 \setminus Q_0^\ell \rangle$ for $\ell = 1, 2$. Since A^1 and A^2 are factor algebras of A , we regard A^1 -modules and A^2 -modules as A -modules.

Let \triangleleft be a partial order on Q_0 defining a quasi-hereditary structure on A and $\triangleleft^\ell := \triangleleft|_{Q_0^\ell}$ for $\ell = 1, 2$. By Lemma 3.4, \triangleleft^ℓ defines a quasi-hereditary structure on A^ℓ . We denote by $\Delta^*(i)$, $\nabla^*(i)$, $T^*(i)$ a standard module, a costandard module and an indecomposable direct summand of the characteristic tilting module of a quasi-hereditary algebra $(A^*, (Q_0^*, \triangleleft^*))$ for $* = \emptyset, 1, 2$. Note that $\nabla(v)$ is obtained by the push-out of $\nabla^1(v) \leftarrow S(v) \rightarrow \nabla^2(v)$. By Lemma 3.4, we have that $\mathcal{F}(\Delta^\ell)$ is contained in $\mathcal{F}(\Delta)$ and $\mathcal{F}(\nabla^\ell(u) \mid u \neq v)$ is contained in $\mathcal{F}(\nabla)$ for $\ell = 1, 2$. Since v is a sink, $\Delta^1(v) = \Delta^2(v) = S(v)$ is a simple projective A^* -module for $* = \emptyset, 1, 2$.

Let $e^\ell = \sum_{u \in Q_0^\ell} e_u$. Then it is easy to see that $A^\ell \cong e^\ell A e^\ell$ as algebras. Thus we have an exact functor $\mathrm{mod} A \rightarrow \mathrm{mod} A^\ell$ defined by $M \mapsto M e^\ell$. Moreover, we have $\Delta(i) e^\ell \cong \Delta^\ell(i)$ if $i \in Q_0^\ell$ and $\Delta(i) e^\ell = 0$ if $i \notin Q_0^\ell$. This restricts to functors $\mathcal{F}(\Delta) \rightarrow \mathcal{F}(\Delta^\ell)$ and $\mathcal{F}(\nabla) \rightarrow \mathcal{F}(\nabla^\ell)$, since the functor is exact and $\nabla(v) e^\ell = \nabla^\ell(v)$ holds. Note that by regarding $M e^\ell$ as an A -module via a map $A \rightarrow A^\ell \cong e^\ell A e^\ell$, we have an injective morphism $M e^\ell \rightarrow M$ of A -modules.

For $i \in Q_0^1$, let $T^1(i)$ be an indecomposable direct summand of the characteristic tilting A^1 -module. We denote by m the length of an $S(v)$ -socle $\text{soc}_v(T^1(i))$ of $T^1(i)$, that is, the maximal direct summand of $\text{soc}(T^1(i))$ which is a direct sum of copies of $S(v)$. So there exists an injective morphism $s: S(v)^{\oplus m} \rightarrow T^1(i)$. Recall that by Proposition 2.16 we have a short exact sequence

$$0 \rightarrow S(v) \xrightarrow{f} T^2(v) \rightarrow X^2(v) \rightarrow 0,$$

where $X^2(v)$ belongs to $\mathcal{F}(\Delta^2(j) \mid j \triangleleft v, j \neq v)$. Consider the following push-out diagram.

$$(3.1) \quad \begin{array}{ccc} S(v)^{\oplus m} & \xrightarrow{s} & T^1(i) \\ f^{\oplus m} \downarrow & & \downarrow \beta \\ T^2(v)^{\oplus m} & \longrightarrow & U(i) \end{array}$$

In Theorem 3.14 we show that $U(i)$ is isomorphic to $T(i)$. Since the cokernel of β is isomorphic to $X^2(v)^{\oplus m}$, $U(i)$ belongs to $\mathcal{F}(\Delta)$. We have the following proposition.

Proposition 3.10. *The push-out $U(i)$ belongs to $\mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$. Moreover, we have $U(i)e^1 \cong T^1(i)$ and $U(i)e^2 \cong T^2(v)^{\oplus m}$.*

Proof. The diagram induces a short exact sequence $0 \rightarrow S(v)^{\oplus m} \rightarrow T^2(v)^{\oplus m} \oplus T^1(i) \rightarrow U(i) \rightarrow 0$. By multiplying $U(i)$ on the right by e^1 and e^2 , we have $U(i)e^1 \cong T^1(i)$ and $U(i)e^2 \cong T^2(v)^{\oplus m}$.

We show that $U(i)$ belongs to $\mathcal{F}(\nabla)$. By [DR92, Theorem 1], $\mathcal{F}(\nabla) = \{Y \in \text{mod } A \mid \text{Ext}_A^1(\Delta, Y) = 0\}$ holds. Therefore we show that $\text{Ext}_A^1(\Delta(j), U(i)) = 0$ for any $j \in Q_0$. Indeed, if $j = v$, then $\text{Ext}_A^1(\Delta(v), U(i)) = 0$ holds, since $\Delta(v)$ is a projective A -module. Take a short exact sequence

$$(3.2) \quad 0 \rightarrow U(i) \rightarrow X \xrightarrow{g} \Delta(j) \rightarrow 0$$

in $\text{mod } A$. Assume that $j \in Q_0^1 \setminus \{v\}$. We have a short exact sequence $0 \rightarrow U(i)e^1 \rightarrow Xe^1 \xrightarrow{ge^1} \Delta(j)e^1 \rightarrow 0$ in $\text{mod}(e^1Ae^1)$. Since $U(i)e^1 \cong T^1(i)$ and $\Delta(j)e^1 \cong \Delta^1(j)$, this sequence splits. Thus there exists an (e^1Ae^1) -morphism $h: \Delta(j)e^1 \rightarrow Xe^1$ such that $ge^1 \circ h = \text{id}$. By Lemma 3.4, $\Delta(j) = \Delta(j)e^1$ holds. So a map $h: \Delta(j) \rightarrow X$ given by $d \mapsto h(d)$ is well-defined. We show that h is a morphism of A -modules. Let $a \in A$ and $d \in \Delta(j)$. Note that $da = de^1ae^1$ holds, since $\Delta(j) = \Delta(j)e^1$ holds. Then we have that $h(da) = h(d)e^1ae^1$. On the other hand, since $\text{im}(h) \subset Xe^1$, we have $h(d)a = h(d)e^1ae^1 + h(d)e^1a(1 - e^1)$. However, $e^1A(1 - e^1) = 0$ holds, since $Q^1 \sqcup Q^2$ is a deconcatenation at a sink. It is easy to see that $g \circ h = \text{id}$, that is, the short exact sequence (3.2) splits. If $j \in Q_0^2 \setminus \{v\}$, then by using the fact that $U(i)e^2 \cong T^2(v)^{\oplus m}$, $\Delta(j) = \Delta(j)e^2 \cong \Delta^2(j)$ and $e^2A(1 - e^2) = 0$, we can show that g is a split morphism. \square

Lemma 3.11. *In the diagram (3.1), the morphism β is a right $\mathcal{F}(\nabla)$ -approximation of $T^1(i)$.*

Proof. By Proposition 3.10, $U(i) \in \mathcal{F}(\nabla)$. Let $M \in \mathcal{F}(\nabla)$ and $g: T^1(i) \rightarrow M$ be a morphism. Since the image of gs is a direct sum of copies of $S(v)$, the image is contained in Me^2 . Since $Me^2 \in \mathcal{F}(\nabla^2)$ and $S(v) \rightarrow T^2(v)$ is a left $\mathcal{F}(\nabla^2)$ -approximation of $S(v)$, there exists a morphism $h: T^2(v)^{\oplus m} \rightarrow M$ such that $hf^{\oplus m} = gs$. Because the diagram is a push-out diagram, there exists a morphism $g': U(i) \rightarrow M$ such that $g'\beta = g$. \square

Recall that $\Delta(i) = \Delta^1(i)$ holds.

Lemma 3.12. *Let $f_1: \Delta(i) \rightarrow T^1(i)$ be a morphism obtained by Proposition 2.16. Then the composite $\Delta(i) \xrightarrow{f_1} T^1(i) \xrightarrow{\beta} U(i)$ is a left $\mathcal{F}(\nabla)$ -approximation of $\Delta(i)$.*

Proof. We denote by $\iota: Me^1 \rightarrow M$ the inclusion map. Let $M \in \mathcal{F}(\nabla)$ and $h: \Delta(i) \rightarrow M$ be a morphism. Since $\Delta^1(i) = \Delta(i)$ holds, there exists a morphism $\iota: \Delta(i) \rightarrow Me^1$ with $h = \iota h'$. Since f_1 is a left $\mathcal{F}(\nabla^1)$ -approximation of $\Delta(i)$ and $Me^1 \in \mathcal{F}(\nabla^1)$, there exists a morphism $f'_1: T^1(i) \rightarrow Me^1$ with $h' = f'_1 f_1$. By Lemma 3.11, there exists a morphism $f'': T^{(\ell)}(i) \rightarrow M$ with $\iota f'_1 = f'' \beta_\ell$. Then we have

$$h = \iota h' = \iota f'_1 f_1 = f''_1 \beta.$$

$$\begin{array}{ccccc}
 & \Delta(i) & \xrightarrow{f_1} & T^1(i) & \xrightarrow{\beta} & U(i) \\
 & \swarrow h' & & \searrow f'_1 & & \\
 & & \downarrow h & & & \\
 Me^1 & \xrightarrow{\iota} & M & \xleftarrow{f''_1} & &
 \end{array}$$

□

Let R be a ring with a surjective ring morphism $R \rightarrow R/I$. It is easy to see that if R/I is a local ring and I is contained in the radical of R , then R is a local ring. The following lemma implies that $T^{(\ell)}(i)$ is indecomposable.

Lemma 3.13. *Let A be a finite dimensional algebra and S be a simple A -module. Let X, Y be A -modules such that the S -socles are $S^{\oplus m}$ and S , respectively, with $m \geq 1$. Consider the following push-out diagram of A -modules*

$$\begin{array}{ccc}
 S^{\oplus m} & \xrightarrow{a} & X \\
 \downarrow b & & \downarrow \beta \\
 Y^{\oplus m} & \xrightarrow{\alpha} & Z
 \end{array}$$

such that a and b are inclusion morphisms associated to S -socles. Assume that the following conditions hold:

- (1) there exist no morphisms from X to the cokernel of b ,
- (2) there exist no morphisms from Y to the cokernel of a ,
- (3) β is a left (add Z)-approximation of X ,
- (4) X and Y are indecomposable.

Then Z is indecomposable.

Proof. We show that the endomorphism algebra of Z is a local algebra. By the assumption (1), we have a morphism of algebras $R_X: \text{End}_A(Z) \rightarrow \text{End}_A(X)$ given by $R_X(\phi) = \phi|_X$. By the assumption (3), this R_X is surjective. We have an exact sequence

$$0 \rightarrow \ker(R_X) \rightarrow \text{End}_A(Z) \rightarrow \text{End}_A(X) \rightarrow 0.$$

By the assumption (4), $\text{End}_A(X)$ is a local algebra. Thus it is enough to show that $\ker(R_X)$ is contained in the radical of $\text{End}_A(Z)$.

Let ϕ be an endomorphism of Z with $R_X(\phi) = 0$. We show that ϕ belongs to the radical of $\text{End}_A(Z)$. By the assumption (2), we have a morphism of algebras $R_{Y^{\oplus m}}: \text{End}_A(Z) \rightarrow \text{End}_A(Y^{\oplus m})$ given by $R_{Y^{\oplus m}}(\phi) = \phi|_{Y^{\oplus m}}$. The endomorphism $R_{Y^{\oplus m}}(\phi)$ is presented by a $(m \times m)$ -matrix with entries in $\text{End}_A(Y)$. Since the S -socle of Y is S and $\phi|_{S^{\oplus m}} = (\phi|_X)|_{S^{\oplus m}} = 0$, all the entries of a matrix presentation of $R_{Y^{\oplus m}}(\phi)$ are not isomorphisms and so these morphisms belongs to the radical of a local algebra $\text{End}_A(Y)$. This implies that $R_{Y^{\oplus m}}(\phi)$ belongs to the radical of $\text{End}_A(Y^{\oplus m})$. In particular, $R_{Y^{\oplus m}}(\phi)$ is nilpotent. Let $f \in \text{End}_A(Z)$ be any morphism. Then $(f\phi)|_{S^{\oplus m}} = 0$ holds since $R_X(\phi) = 0$. Thus by the same argument, $R_{Y^{\oplus m}}(f\phi)$ is nilpotent. By the universality of the push-out diagram, $f\phi$ is nilpotent for any $f \in \text{End}_A(Z)$. This implies that $\text{id}_Z - f\phi$ is an isomorphism, and so ϕ belongs to the radical of $\text{End}_A(Z)$. □

Then we have the following result.

Theorem 3.14. *For $i \in Q_0^1$, let m be the length of an $S(v)$ -socle of $T^1(i)$. Then the push-out $U(i)$ of $T^2(v)^{\oplus m} \leftarrow S(v)^{\oplus m} \rightarrow T^1(i)$ as (3.1) is isomorphic to $T(i)$.*

Proof. By Lemmas 3.11, 3.12 and 3.13, $U(i)$ is indecomposable. By Lemma 3.12, $U(i)$ gives a left $\mathcal{F}(\nabla)$ -approximation of $\Delta(i)$. Therefore, $U(i)$ is isomorphic to $T(i)$. □

4. PATH ALGEBRAS OF TYPE \mathbb{A}

4.1. Path algebra of an equioriented quiver of type \mathbb{A} . Let $A_n = 1 \rightarrow 2 \rightarrow \cdots \rightarrow n$ be an equioriented quiver of type \mathbb{A} . It was first noticed by Gabriel that tilting modules over $\Lambda_n := \mathbf{k}A_n$

are counted by the n -th Catalan number $c_n = \frac{1}{n+1} \binom{2n}{n}$ [Gab81]. In this subsection we show that the number of different quasi-hereditary structures on the path algebra Λ_n coincides with c_n .

Recall that *binary trees* can be defined inductively as follows. A *binary tree* T is either the empty set or a tuple (r, L, R) where r is a singleton set, called the root of T , and L and R are two binary trees. The empty set has no vertex but has one leaf. The set of leaves of $T = (r, L, R)$ is the disjoint union of the set of leaves of L and R . The *size* of the tree is its number of vertices (equivalently the number of leaves minus 1). It is classical that binary trees are counted by the Catalan numbers.

A *binary search tree* is a binary tree labeled by integers such that if a vertex x is labeled by k , then the vertices of the left subtree (resp. right subtree) of x are labeled by integers less than or equal (resp. superior) to k . If T is a binary tree with n vertices, there is a unique labeling of the vertices by each of the integers $1, 2, \dots, n$ that makes it a binary search tree. This procedure is sometimes called the *in-order traversal* of the tree or simply as the in-order algorithm (recursively visit left subtree, root and right subtree). The first vertex visited by the algorithm is labeled by 1, the second by 2 and so on, see Fig. 1.




FIGURE 1. Binary search tree of size 5 labelled by the in-order algorithm.

Let T be a binary tree of size n viewed as a binary search tree. Then T induces a poset \triangleleft_T on $\{1, 2, \dots, n\}$ by setting $i \triangleleft_T j$ if i labels a vertex in the subtree of the vertex labelled by j .

Proposition 4.1. *Let \triangleleft be a partial order on $\{1, 2, \dots, n\}$. Then there is a binary tree T such that $\triangleleft = \triangleleft_T$ if and only if*

- (1) *For every $i < j$ incomparable with respect to \triangleleft , there exists k such that $i < k < j$ and $i \triangleleft k$ and $j \triangleleft k$.*
- (2) *For every $i < j < k$, if $i \triangleleft k$ then $j \triangleleft k$ and if $k \triangleleft i$ then $j \triangleleft i$.*

Proof. See [CPP19, Proposition 2.21]. □

Remark 4.2. Condition (1) is equivalent to the following weaker condition: for every $i < j$ incomparable there exists k such that $i < k < j$ and $i \triangleleft k$ or $j \triangleleft k$.

Lemma 4.3. (1) *Let T be a binary tree of size n . Then \triangleleft_T is an adapted poset to Λ_n .*

- (2) *If \triangleleft is an adapted poset to Λ_n , then there is a binary tree T such that $\triangleleft \sim \triangleleft_T$.*

Proof. We first show (1). The indecomposable Λ_n -modules can be identified with usual intervals in $\{1, 2, \dots, n\}$. Then the assertion follows from Proposition 4.1.

We next show (2). Let \triangleleft be an adapted poset to Λ_n and let \triangleleft_m the minimal adapted poset equivalent to \triangleleft . Then by Proposition 2.9, we have $\triangleleft_m = (\text{Dec}(\triangleleft) \cup \text{Inc}(\triangleleft))^{tc}$. We may assume that $\triangleleft = \triangleleft_m$. The relation $j \triangleleft i$ is *decreasing* if $[\Delta(i) : S(j)] \neq 0$. Since $P(i)$ has basis the set of paths starting at i , this implies that there is a path from i to j . Our choice of orientation implies that $i \leq j$. In addition for every $i \leq k \leq j$, the simple module $S(k)$ is a composition factor of $\Delta(i)$, so $k \triangleleft i \in \text{Dec}(\triangleleft)$. This proves that the decreasing relations, and by a similar argument the increasing relations, satisfy the assertions (2) of Proposition 4.1. Conversely a relation $j \triangleleft i$, such that $i \leq j$ and any $k \in [i, j]$ satisfies $k \triangleleft i$, is decreasing. It is easy to see that the relation obtained as transitivity of two decreasing (increasing) relations is also decreasing (increasing). Moreover if $i \triangleleft j \in \text{Inc}(\triangleleft)$ and $j \triangleleft k \in \text{Dec}(\triangleleft)$ with $k \neq j$, then we have $k < i$, because if $k \in [i, j]$, then $k \triangleleft j$ which contradicts the antisymmetry of \triangleleft . For $x \in [k, i]$ we have $x \triangleleft k$, so $i \triangleleft k$ is decreasing. With a similar argument, the relation obtained as transitivity of a decreasing relation and an increasing one is increasing. This shows that $\triangleleft_m = \text{Dec}(\triangleleft) \cup \text{Inc}(\triangleleft)$ and that every relation satisfies (2) of Proposition 4.1. Because \triangleleft_m is adapted it satisfies (1), so there is a binary tree T such that $\triangleleft_m = \triangleleft_T$. □

Proposition 4.4. *Let $n \in \mathbb{N}$. The map sending a binary tree T to the equivalence class of the adapted poset \triangleleft_T is a bijection between the set of binary trees of size n and the set of equivalence classes of adapted posets for Λ_n .*

Proof. We already know that this map is surjective, we need to see that it is injective. For that we explain how we can recover the tree for the set of standard and costandard modules.

Let T be a binary tree. Then \triangleleft_T is an adapted poset to Λ_n and $(\Lambda_n, \triangleleft_T)$ is a quasi-hereditary algebra. It is easy to see that the composition factors of the $\Delta(i)$ are indexed by the elements in the right subtree of i and the composition factors of $\nabla(i)$ are indexed the elements of the left subtree. It follows that two different trees induce two non-equivalent posets. \square

Lemma 4.5. *Let T be a binary tree of size n . Then \triangleleft_T is a minimal adapted order to Λ_n .*

Proof. Let \triangleleft' be an adapted poset to Λ_n such that $\triangleleft_T \sim \triangleleft'$. Then there exists a binary tree T' such that $\triangleleft' \sim \triangleleft_{T'}$, by Lemma 4.3. Thus $T = T'$ by Proposition 4.4, and the proof of Lemma 4.3 (2) shows that \triangleleft_T is extended by \triangleleft' , which shows the claim. \square

If \triangleleft is an adapted order to Λ_n , the pair $(\Lambda_n, \triangleleft)$ is a quasi-hereditary algebra so it has a characteristic tilting module T which is characterized by $\text{add}(T) = \mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$. Since Λ_n is a hereditary algebra, the module T is a tilting module. Moreover, the tilting module only depends on the equivalence class of the partial order. So we have a map char from the set of equivalence classes of adapted partial orders to the set of isomorphism classes of tilting modules for Λ_n which sends the equivalence class of \triangleleft to the characteristic tilting module of $(\Lambda_n, \triangleleft)$.

Theorem 4.6. *We have a commutative diagram of bijections*

$$\begin{array}{ccc}
 & \{ \text{Binary trees of size } n \} & \\
 \begin{array}{c} \swarrow \\ T \mapsto \triangleleft_T \end{array} & & \searrow \phi \\
 \{ \text{Adapted partial orders to } \Lambda_n \} / \sim & \xrightarrow{\text{char}} & \{ \text{Tilting modules over } \Lambda_n \} / \cong
 \end{array}$$

where ϕ is the classical bijection between binary trees and tilting modules for Λ_n .

Proof. In the proof of Proposition 4.4 we determined the set of standard and costandard modules from the binary tree T' . We claim that the indecomposable direct summand $T(i)$ of the characteristic tilting module T is the indecomposable module with composition factors indexed by the interval consisting of i and the label of its subtrees (left and right). Since the map ϕ sends T' to the module constructed in this way (see [Hil06, Section 9]), the proof follows from this claim.

By induction on the size of the subtrees we show that the module $T(i)$ is in $\mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$. This is clear for the subtrees of size one since in this case $T(i) = S(i) = \Delta(i) = \nabla(i)$.

In the general case, if i_l (resp. i_r) denotes the left (resp. right) child of i we have two exact sequences

$$0 \rightarrow \Delta(i) \rightarrow T(i) \rightarrow T(i_r) \rightarrow 0$$

and

$$0 \rightarrow T(i_l) \rightarrow T(i) \rightarrow \nabla(i) \rightarrow 0.$$

If i has not left (resp.) right child then we let $T(i_l) = 0$ (resp. $T(i_r) = 0$) and we still have the two exact sequences. By induction $T(i_r) \in \mathcal{F}(\Delta)$ and $T(i_l) \in \mathcal{F}(\nabla)$, so $T(i) \in \mathcal{F}(\Delta) \cap \mathcal{F}(\nabla)$. The results follows. \square

Corollary 4.7. *Let $n \geq 1$. Then $\text{qh.str}(\Lambda_n)$ and the Tamari lattice of size n are isomorphic as partially ordered sets.*

Proof. As explained in the proof of Lemma 4.3 our notion of decreasing and increasing relations coincide with the one of [CPP19]. The result is then a consequence of [CPP19, Proposition 41]. Note that char is a morphism of posets by Lemma 2.20. Moreover, it is well known that ϕ is also a morphism of posets if the set of binary trees is endowed with the usual partial order induced by rotations. \square

4.2. Path algebras of type \mathbb{A} : general case. Let Q be a quiver whose underlying graph is of type \mathbb{A}_n and $I = Q_0$. In this subsection, we classify all the quasi-hereditary structures on $\mathbf{k}Q$ and its characteristic tilting modules.

Theorem 4.8. *Let $Q^1 \sqcup Q^2 \sqcup \cdots \sqcup Q^\ell$ be an iterated deconcatenation of Q such that each Q^i is an equioriented quiver of type \mathbb{A}_{n_i} for some $n_i \in \mathbb{Z}_{\geq 1}$. Then there is a bijection*

$$\text{qh.str}(\mathbf{k}Q) \longrightarrow \prod_{i=1}^{\ell} \text{qh.str}(\Lambda_{n_i})$$

given by $[\triangleleft] \mapsto ([\triangleleft|_{Q_0^i}])_{i=1}^{\ell}$. Moreover, if \triangleleft is a minimal adapted order, then there exists a binary tree T_i of size n_i such that $\triangleleft|_{Q_0^i} = \triangleleft|_{T_i}$, for each $1 \leq i \leq \ell$.

Proof. The bijection follows from Theorem 3.7. The second assertion is consequence of Lemmas 3.9 and 4.5 and Proposition 4.4. \square

Since any indecomposable $\mathbf{k}Q$ -module is determined by its composition factor, the following theorem gives a complete construction of characteristic tilting modules from minimal adapted orders.

Theorem 4.9. *Let \triangleleft be a minimal adapted order to $\mathbf{k}Q$ and $T = \bigoplus_{i \in I} T(i)$ be the characteristic tilting module associated to \triangleleft . Then for vertices $i, j \in I$, $S(j)$ is a composition factor of $T(i)$ if and only if $j \triangleleft i$ holds.*

Proof. The only if part follows from Proposition 2.16. Assume that Q has an iterated deconcatenation with ℓ components such that each quiver is an equioriented quiver of type \mathbb{A} . We show the if part by an induction on ℓ . If $\ell = 1$, then the assertion holds by Theorem 4.6. Assume that $\ell > 1$. We have a deconcatenation $Q^1 \sqcup Q^2$ of Q at a vertex $v \in Q_0$ such that Q^1 is an equioriented quiver of type \mathbb{A} , and Q^2 has an iterated deconcatenation with $\ell - 1$ components such that each quiver is an equioriented quiver of type \mathbb{A} . By Lemma 3.9, the adapted orders $\triangleleft|_{Q_0^1}$ and $\triangleleft|_{Q_0^2}$ are also minimal. If v is a source, then apply Lemma 2.22, and we may assume that v is a sink. For $k = 1, 2$ and $i \in Q_0^k$, let $T^k(i)$ be the indecomposable direct summand of the characteristic tilting $\mathbf{k}Q^k$ -module associated to an adapted order $\triangleleft|_{Q_0^k}$. Let $e^k = \sum_{u \in Q_0^k} e_u$ and $\bar{1} = 2, \bar{2} = 1$. If $i \in Q_0^k$, then by Proposition 3.10 we have $T(i)e^k \cong T^{\bar{k}}(v)$, $T(i)e^{\bar{k}} \cong T^k(v)$ if $[T(i) : S(v)] \neq 0$ and $T(i)e^{\bar{k}} = 0$ if $[T(i) : S(v)] = 0$. The assertion follows by induction on ℓ and Theorem 4.6. \square

We finish this section by giving a concrete example.

Example 4.10. Let $A = \mathbf{k}Q$, where $Q = 1 \rightarrow 2 \leftarrow 3 \leftarrow 4 \rightarrow 5$ as before. Then we have that

$$\begin{array}{ccccc} & & 3 & & 5 \\ & & \nearrow \searrow & & \uparrow \\ 1 & & 2 & 4 & 4 \\ \uparrow & & & & \\ 2 & & & & \end{array}$$

are the Hasse diagrams of some minimal adapted posets to $\mathbf{k}(1 \rightarrow 2)$, $\mathbf{k}(2 \leftarrow 3 \leftarrow 4)$ and $\mathbf{k}(4 \rightarrow 5)$, respectively. Then, by Lemma 3.9 we have that the concatenation of the last Hasse diagrams

$$\begin{array}{ccccc} & & 3 & & 5 \\ & & \nearrow \searrow & & \nearrow \\ 1 & & 2 & 4 & 4 \end{array}$$

is the corresponding minimal adapted poset to A . In this case $|\text{qh.str}A| = 20$. In Fig. 2 we depict the Hasse diagram of $\text{qh.str}(A)$. The vertices correspond to minimal adapted orders to A which represent all the quasi-hereditary structures on A . Note that if a total order is a minimal adapted order, then it is the unique element in its equivalence class.

5. PATH ALGEBRAS OF TYPES \mathbb{D} AND \mathbb{E}

In this section, we count the number of quasi-hereditary structures on $\mathbf{k}Q$ for a quiver Q of Dynkin type \mathbb{D} and \mathbb{E} .

5.1. Idempotent reduction. Let A be a finite dimensional algebra and $\{S(i)\}_{i \in I}$ the set of isomorphism classes of simple A -modules. Fix $i \in I$ and a corresponding idempotent e_i in A . For a partial order \triangleleft on $I \setminus \{i\}$, we denote by \triangleleft' a partial order on I such that $\triangleleft'|_{I \setminus \{i\}} = \triangleleft$ and i is a unique maximal element.

Lemma 5.1. *If \triangleleft be an adapted order to $A/\langle e_i \rangle$, then \triangleleft' is an adapted order to A .*




FIGURE 2. Poset of quasi-hereditary structures on $A = \mathbf{k}(1 \rightarrow 2 \leftarrow 3 \leftarrow 4 \rightarrow 5)$.

Proof. Let M be an A -module such that $\text{top } M \cong S(j)$ and $\text{soc } M \cong S(k)$ for some $j, k \in I \setminus \{i\}$, and j, k are incomparable with respect to \triangleleft' . If M is an $A/\langle e_i \rangle$ -module, then nothing to show, since \triangleleft is an adapted order to $A/\langle e_i \rangle$. On the other hand, if $Me_i \neq 0$, then $j \triangleleft' i, k \triangleleft' i$ implies that \triangleleft' is adapted to A . \square

Lemma 5.2. *Let $\triangleleft_1, \triangleleft_2$ be partial orders on $I \setminus \{i\}$ which define quasi-hereditary structures on $A/\langle e_i \rangle$. Assume that both $\triangleleft'_1, \triangleleft'_2$ define quasi-hereditary structures on A . Then $\triangleleft_1 \sim \triangleleft_2$ if and only if $\triangleleft'_1 \sim \triangleleft'_2$.*

Proof. By Lemma 2.15, it is clear that $\triangleleft'_1 \sim \triangleleft'_2$ implies $\triangleleft_1 \sim \triangleleft_2$. Since i is a unique maximal element in I , $\Delta'_1(i) = P(i) = \Delta'_2(i)$ holds. Thus the converse is also true. \square

Then we concentrate on path algebras. Let Q be a finite acyclic quiver and fix $i \in Q_0$. By Lemmas 5.1 and 5.2, we have the following well-defined injective map:

$$\iota_i: \text{qh.str}(\mathbf{k}Q/\langle e_i \rangle) \longrightarrow \text{qh.str}(\mathbf{k}Q) \quad [\triangleleft] \mapsto [\triangleleft'].$$

Lemma 5.3. *For a finite acyclic quiver Q , we have*

$$\text{qh.str}(\mathbf{k}Q) = \bigcup_{i \in Q_0} \text{Im}(\iota_i),$$

and each $\text{Im}(\iota_i)$ bijectively corresponds to $\text{qh.str}(\mathbf{k}Q/\langle e_i \rangle)$.

Proof. Let \triangleleft be a partial order on Q_0 which defines a quasi-hereditary structure on $\mathbf{k}Q$. Then we may assume that \triangleleft is a total order with a unique maximal element i . Then we have a total order $\triangleleft|_{Q_0 \setminus \{i\}}$ on $Q_0 \setminus \{i\}$ and we have $\iota_i([\triangleleft|_{Q_0 \setminus \{i\}}]) = [\triangleleft]$. \square

5.2. Decomposition of the set of quasi-hereditary structures. Throughout this subsection, let $r, s, t \in \mathbb{Z}_{\geq 1}$ and $Q = Q(r, s, t)$ the following quiver:

$$(5.1) \quad \begin{array}{ccccccc} & & & & & b_1 & \longrightarrow & \cdots & \longrightarrow & b_s \\ & & & & & \nearrow & & & & \\ a_r & \longrightarrow & a_{r-1} & \longrightarrow & \cdots & \longrightarrow & a_1 & \longrightarrow & a_0 & \\ & & & & & \searrow & & & & \\ & & & & & c_1 & \longrightarrow & \cdots & \longrightarrow & c_t \end{array}$$

Recall that we have $\mathbf{qh.str}(\mathbf{k}Q) = \bigcup_i \mathbf{lm}(\iota_i)$, where i runs over all the vertices of the quiver Q . Each $\mathbf{lm}(\iota_i)$ bijectively corresponds to $\mathbf{qh.str}(\mathbf{k}Q/\langle e_i \rangle)$. Let $Q_0^b := \{b_k \mid 1 \leq k \leq s\}$ and $Q_0^c := \{c_k \mid 1 \leq k \leq t\}$. We begin with the following lemma.

Lemma 5.4. *Let i, j be vertices of Q with $i \neq j$. If $\mathbf{lm}(\iota_i) \cap \mathbf{lm}(\iota_j) \neq \emptyset$, then we have $i \in Q_0^b$ and $j \in Q_0^c$.*

Proof. Let $\mathfrak{q} \in \mathbf{lm}(\iota_i) \cap \mathbf{lm}(\iota_j)$ and denote by Δ the set of standard $\mathbf{k}Q$ -modules associated to \mathfrak{q} . For each $\ell = i, j$, by the definition of ι_ℓ , there exist a partial order \triangleleft'_ℓ on Q_0 such that $[\triangleleft'_\ell] = \mathfrak{q}$ and ℓ is a unique maximal element of \triangleleft'_ℓ . By extending \triangleleft'_ℓ and Lemma 2.3, we may assume that \triangleleft'_ℓ is totally ordered for each $\ell = i, j$. We have $\Delta(i) = P(i)$ and $\Delta(j) = P(j)$.

Assume that $P(i)$ and $P(j)$ have a common composition factor $S(k)$. Then we can take $k = i$ or $k = j$ because of the shape of the quiver Q . If $k = i$, then $\Delta(j) = P(j)$ implies that $i \triangleleft'_i j$, which is a contradiction. Similarly, $k = j$ induces a contradiction. Thus $P(i)$ and $P(j)$ have no common composition factor. This implies the assertion. \square

Then we observe the intersection $\mathbf{lm}(\iota_i) \cap \mathbf{lm}(\iota_j)$ for $i \in Q_0^b$ and $j \in Q_0^c$. For each vertex $i \in Q_0$, we have a fully faithful functor $\text{mod}(\mathbf{k}Q/\langle e_i \rangle) \rightarrow \text{mod } \mathbf{k}Q$. By this functor, we regard $\mathbf{k}Q/\langle e_i \rangle$ -modules as $\mathbf{k}Q$ -modules.

Lemma 5.5. *Let $i \in Q_0^b$, $j \in Q_0^c$ and $\mathfrak{q}_i \in \mathbf{qh.str}(\mathbf{k}Q/\langle e_i \rangle)$. The following statements are equivalent.*

- (1) *There exists a partial order \triangleleft on $Q_0 \setminus \{i\}$ such that $[\triangleleft] = \mathfrak{q}_i$ and a vertex j is a maximal element by \triangleleft .*
- (2) $\iota_i(\mathfrak{q}_i) \in \mathbf{lm}(\iota_j)$.

Proof. Clearly, (1) implies (2). Conversely, assume that there exists $\mathfrak{q}_j \in \mathbf{qh.str}(\mathbf{k}Q/\langle e_j \rangle)$ with $\iota_i(\mathfrak{q}_i) = \iota_j(\mathfrak{q}_j)$. For $\ell = i, j$, let \triangleleft_ℓ be a partial order on $Q_0 \setminus \{\ell\}$ such that $[\triangleleft_\ell] = \mathfrak{q}_\ell$. We have $[\triangleleft'_\ell] = \iota_\ell(\mathfrak{q}_\ell)$. We denote by Δ the standard $\mathbf{k}Q$ -modules with respect to $\iota_i(\mathfrak{q}_i) = \iota_j(\mathfrak{q}_j)$. Since $\iota_i(\mathfrak{q}_i) = \iota_j(\mathfrak{q}_j)$ is represented by both \triangleleft'_i and \triangleleft'_j , for any $k \in Q_0 \setminus \{i, j\}$, $\Delta(k)$ does not have composition factors $S(i)$ and $S(j)$, and $\Delta(i) = P(i)$, $\Delta(j) = P(j)$ hold.

We denote by \triangleleft^i the partial order on $Q_0 \setminus \{i\}$ obtained by restricting \triangleleft'_j to $Q_0 \setminus \{i\}$. We show that $[\triangleleft^i] = \mathfrak{q}_i$ holds. Let Δ_i and Δ^i the standard $\mathbf{k}Q/\langle e_i \rangle$ -modules of \triangleleft_i and \triangleleft^i , respectively. We show that $\Delta_i(k) = \Delta(k) = \Delta^i(k)$ holds for any $k \in Q_0 \setminus \{i\}$. Since $\iota_i(\mathfrak{q}_i)$ is represented by \triangleleft'_i , we have $\Delta_i(k) = \Delta(k)$ for any $k \in Q_0 \setminus \{i\}$ by Lemma 2.15. Let $k \in Q_0 \setminus \{i\}$. Since j is maximal by \triangleleft^i , $i \in Q_0^b$ and $j \in Q_0^c$, $\Delta^i(j) = P(j) = \Delta(j)$ holds. Assume that $k \neq j$. Again since j is maximal by \triangleleft^i , $\Delta^i(k)$ does not have a composition factor $S(j)$. Then we can show that for any $v \in Q_0 \setminus \{i, j\}$, $S(v)$ is a composition factor of $\Delta^i(k)$ if and only if $S(v)$ is a composition factor of $\Delta(k)$. In fact, since \triangleleft^i is obtained by restricting \triangleleft'_j , $\Delta^i(k)$ is a factor of $\Delta(k)$. Namely, the only if part holds. On the other hand, if $S(v)$ is a composition factor of $\Delta(k)$, then there exists a (unique) path p in Q from k to v such that each vertex u of the path p satisfies $u \triangleleft'_j k$. Since $S(i)$ is not a composition factor of $\Delta(k)$, the path p does not factor through a vertex i . Thus the path p is a path in $Q \setminus \{i\}$. This implies that $S(v)$ is a composition factor of $\Delta^i(k)$. We have $\Delta(k) = \Delta^i(k)$. \square

For $1 \leq k \leq t$, we denote by \mathcal{C}_k a subset of $\mathbf{qh.str}(\mathbf{k}Q/\langle e_{c_k} \rangle)$ consisting of \mathfrak{q} such that there exists no partial order \triangleleft on $Q_0 \setminus \{c_k\}$ such that $[\triangleleft] = \mathfrak{q}$ and \triangleleft has one of b_1, \dots, b_s as a maximal element. Then by Lemma 5.5 we have the following proposition.

Proposition 5.6. *We have a bijection*

$$\bigsqcup_{i=0}^r \text{qh.str} \left(\frac{\mathbf{k}Q}{\langle e_{a_i} \rangle} \right) \sqcup \bigsqcup_{j=1}^s \text{qh.str} \left(\frac{\mathbf{k}Q}{\langle e_{b_j} \rangle} \right) \sqcup \bigsqcup_{k=1}^t \mathcal{C}_k \longrightarrow \text{qh.str}(\mathbf{k}Q),$$

where the map is given by ι_i .

5.3. Counting quasi-hereditary structures. In this subsection, we count the number of quasi-hereditary structures on path algebras of Dynkin type \mathbb{D} . Let $Q = Q(r, s, t)$ be the quiver defined in Eq. (5.1). We consider type \mathbb{D}_n for an integer $n \geq 4$. By taking the opposite quiver and Theorem 3.7, it is enough to study the following two cases.

$$Q(n-3, 1, 1) : \quad a_{n-3} \longrightarrow \cdots \longrightarrow a_1 \longrightarrow a_0 \begin{array}{l} \nearrow b_1 \\ \searrow c_1 \end{array}$$

$$Q(1, n-3, 1) : \quad a_1 \longrightarrow a_0 \begin{array}{l} \nearrow b_1 \longrightarrow \cdots \longrightarrow b_{n-3} \\ \searrow c_1 \end{array}$$

In both cases, $t = 1$, so we study \mathcal{C}_1 for both cases. We denote by c_n the n -th Catalan number $c_n = \frac{1}{n+1} \binom{2n}{n}$. We begin with the following easy observation.

Lemma 5.7. *Let $Q = n \rightarrow \cdots \rightarrow 2 \rightarrow 1$ be an equioriented quiver of type \mathbb{A}_n . For $1 \leq i \leq n$, let $\text{qh}(i)$ be a subset of $\text{qh.str}(\mathbf{k}Q)$ consisting of \mathfrak{q} which is represented by a partial order having i as a maximal element. Then the following statements hold.*

- (1) *Any adapted order to $\mathbf{k}Q$ has a unique maximal element.*
- (2) *The image of the map $\iota_i: \text{qh.str}(\mathbf{k}Q/\langle e_i \rangle) \rightarrow \text{qh.str}(\mathbf{k}Q)$ corresponds to $\text{qh}(i)$, that is, $\text{Im}(\iota_i) = \text{qh}(i)$. In particular, we have a decomposition $\text{qh.str}(\mathbf{k}Q) = \bigsqcup_{i=1}^n \text{Im}(\iota_i)$.*

Proof. (1) By Theorem 4.6, any adapted order to $\mathbf{k}Q$ is an extension of the adapted order induced from a binary tree. Clearly, each partial order induced from binary trees has a unique maximal element. (2) directly follows from the definition of ι_i . \square

Lemma 5.8. *Let $Q = Q(n-3, 1, 1)$. Then we have the following equalities.*

- (1) $|\mathcal{C}_1| = c_{n-1} - c_{n-2}$
- (2) $|\text{qh.str}(\mathbf{k}Q)| = 2c_n - 3c_{n-1}$

Proof. (1) Recall that \mathcal{C}_1 a subset of $\text{qh.str}(\mathbf{k}Q/\langle e_{c_1} \rangle)$ consisting of \mathfrak{q} such that there exist no partial orders on $Q_0 \setminus \{c_1\}$ which represent \mathfrak{q} and have b_1 as a maximal element. Therefore, by Lemma 5.7, we have $\mathcal{C}_1 = \text{qh.str}(\mathbf{k}Q/\langle e_{c_1} \rangle) \setminus \text{Im}(\iota_{b_1})$. Therefore, we have $|\mathcal{C}_1| = |\text{qh.str}(\mathbf{k}Q/\langle e_{c_1} \rangle)| - |\text{Im}(\iota_{b_1})| = |\text{qh.str}(\mathbf{k}Q/\langle e_{c_1} \rangle)| - |\text{qh.str}(\mathbf{k}Q/\langle e_{c_1}, e_{b_1} \rangle)| = c_{n-1} - c_{n-2}$.

(2) We show the equality by an induction on n . Although we are considering \mathbb{D} type, it is easy to see that the equality holds for $n = 3$. Assume that $n > 3$. We have the following equalities.

$$(5.2) \quad |\text{qh.str}(\mathbf{k}Q)| = \left| \bigsqcup_{i=0}^{n-3} \text{qh.str} \left(\frac{\mathbf{k}Q}{\langle e_{a_i} \rangle} \right) \right| + \left| \text{qh.str} \left(\frac{\mathbf{k}Q}{\langle e_{b_1} \rangle} \right) \right| + |\mathcal{C}_1|$$

$$(5.3) \quad = \sum_{i=0}^{n-3} c_{n-i-3}(2c_{i+2} - 3c_{i+1}) + c_{n-1} + c_{n-1} - c_{n-2}$$

$$(5.4) \quad = 2(c_n - c_{n-1} - c_{n-2}) - 3(c_{n-1} - c_{n-2}) + 2c_{n-1} - c_{n-2} \\ = 2c_n - 3c_{n-1}.$$

For the first and the second equalities (5.2) and (5.3), we use the statement (1), Theorem 4.6, Proposition 5.6, Lemma 5.7, and an inductive hypothesis. For the third equality (5.4), we use the well known equality $c_{n+1} = \sum_{i=0}^n c_{n-i}c_i$. \square

Lemma 5.9. *Let $Q = Q(1, n-3, 1)$. Then the following equalities hold.*

- (1) $|\mathcal{C}_1| = c_{n-2} + c_{n-3}$
- (2) $|\text{qh.str}(\mathbf{k}Q)| = 3c_{n-1} - c_{n-2}$

Proof. (1) Recall that \mathcal{C}_1 is a subset of $\text{qh.str}(\mathbf{k}Q/\langle e_{c_1} \rangle)$ consisting of \mathfrak{q} such that there exist no partial orders on $Q_0 \setminus \{c_1\}$ which represent \mathfrak{q} and have one of b_1, \dots, b_{n-3} as a maximal element. Therefore, by Lemma 5.7, we have $|\mathcal{C}_1| = |\text{qh.str}(\mathbf{k}Q/\langle e_{c_1}, e_{a_1} \rangle)| + |\text{qh.str}(\mathbf{k}Q/\langle e_{c_1}, e_{a_0} \rangle)| = c_{n-2} + c_{n-3}$.

(2) We show the equality by an induction on n . For $n = 3$, the equality holds. Assume that $n > 3$. We have the following equalities.

$$\begin{aligned} |\text{qh.str}(\mathbf{k}Q)| &= \left| \bigsqcup_{i=1}^{n-3} \text{qh.str} \left(\frac{\mathbf{k}Q}{\langle e_{b_i} \rangle} \right) \right| + \left| \text{qh.str} \left(\frac{\mathbf{k}Q}{\langle e_{a_1} \rangle} \right) \right| + \left| \text{qh.str} \left(\frac{\mathbf{k}Q}{\langle e_{a_0} \rangle} \right) \right| + |\mathcal{C}_1| \\ &= \sum_{i=1}^{n-3} (3c_{i+1} - c_i)c_{n-i-3} + 2c_{n-2} + c_{n-3} + c_{n-2} + c_{n-3} \\ &= 3(c_{n-1} - c_{n-3} - c_{n-2}) - (c_{n-2} - c_{n-3}) + 3c_{n-2} + 2c_{n-3} \\ &= 3c_{n-1} - c_{n-2}. \end{aligned}$$

All equalities are induced by a similar way as those of the proof of Lemma 5.8. \square

We find the following correspondence to OEIS sequences [OEI20].

Q		Formula	OEIS
$Q(n-3, 1, 1)$	$ \mathcal{C}_1 $	$c_{n-1} - c_{n-2}$	A000245($n-2$)
	$ \text{qh.str}(\mathbf{k}Q) $	$2c_n - 3c_{n-1}$	A070031($n-2$)
$Q(1, n-3, 1)$	$ \mathcal{C}_1 $	$c_{n-2} + c_{n-3}$	A005807($n-3$)

The sequence $(3c_{n-1} - c_{n-2})_{n \geq 2}$ is not listed in [OEI20] until the publication of this article.

Example 5.10. Here we give a list of the numbers of quasi-hereditary structures on path algebras of Dynkin types $\mathbb{E}_6, \mathbb{E}_7$ and \mathbb{E}_8 . By taking the opposite quiver and Theorem 3.7, it is enough to calculate the following cases of $Q = Q(r, s, t)$.

- (1) For \mathbb{E}_6 , $(r, s, t) = (1, 2, 2)$ or $(2, 2, 1)$.
- (2) For \mathbb{E}_7 , $(r, s, t) = (1, 3, 2), (2, 3, 1)$ or $(3, 2, 1)$.
- (3) For \mathbb{E}_8 , $(r, s, t) = (1, 4, 2), (2, 4, 1)$ or $(4, 2, 1)$.

	(r, s, t)	$ \mathcal{C}_1 $	$ \mathcal{C}_2 $	$ \text{qh.str}(\mathbf{k}Q) $
\mathbb{E}_6	$(1, 2, 2)$	7	19	106
	$(2, 2, 1)$	23	0	130
\mathbb{E}_7	$(1, 3, 2)$	19	52	322
	$(2, 3, 1)$	66	0	416
	$(3, 2, 1)$	76	0	453
\mathbb{E}_8	$(1, 4, 2)$	56	154	1020
	$(2, 4, 1)$	202	0	1368
	$(4, 2, 1)$	255	0	1584

For example, let $Q = Q(1, 3, 2)$. Then $|\mathcal{C}_1|$, $|\mathcal{C}_2|$ and $|\mathbf{qh.str}(\mathbf{k}Q)|$ are obtained as follows:

$$\begin{aligned}
|\mathcal{C}_1| &= |\mathbf{qh.str}(\mathbf{k}Q/\langle e_{c_1} \rangle)| - \sum_{i=1}^3 |\mathbf{qh.str}(\mathbf{k}Q/\langle e_{c_1}, e_{b_i} \rangle)| \\
&= c_5 - c_2c_2 - c_3 - c_4 = 19, \\
|\mathcal{C}_2| &= |\mathbf{qh.str}(\mathbf{k}Q/\langle e_{c_2} \rangle)| - \sum_{i=1}^3 |\mathbf{qh.str}(\mathbf{k}Q/\langle e_{c_2}, e_{b_i} \rangle)| \\
&= (3c_5 - c_4) - c_3c_2 - (3c_3 - c_2) - (3c_4 - c_3) = 52, \\
|\mathbf{qh.str}(\mathbf{k}Q)| &= \sum_{i=0}^1 |\mathbf{qh.str}(\mathbf{k}Q/\langle e_{a_i} \rangle)| + \sum_{i=1}^3 |\mathbf{qh.str}(\mathbf{k}Q/\langle e_{b_i} \rangle)| + |\mathcal{C}_1| + |\mathcal{C}_2| \\
&= c_3c_2 + c_4c_3 + c_4c_2 + (3c_4 - c_3) + |\mathbf{qh.str}(\mathbf{k}Q(1, 2, 2))| + 19 + 52 \\
&= 322.
\end{aligned}$$

6. LATTICE OF QUASI-HEREDITARY STRUCTURES

Throughout this section, all algebras are defined over a fixed field \mathbf{k} .

6.1. Lattice properties and incidence algebras. We denote by \tilde{D}_n the following quiver,

$$(6.1) \quad \begin{array}{ccccccc} & \circ & & & & & \circ \\ & \searrow & & & & & \nearrow \\ & & \circ & \longrightarrow & \circ & \longrightarrow & \circ & \longrightarrow & \circ & \longrightarrow & \circ \\ & \nearrow & & & & & \searrow & & & & \circ \\ \circ & & & & & & & & & & \circ \end{array}$$

where the number of vertices is $n + 1$.

The main result of this section is the following theorem.

Theorem 6.1. *Let Q be a finite quiver whose underlying graph is a tree. Then the set $\mathbf{qh.str}(\mathbf{k}Q)$ of quasi-hereditary structures on $\mathbf{k}Q$ is a lattice if and only if Q does not have a quiver \tilde{D}_n as a subquiver for any $n \geq 4$.*

Curiously, this characterization does not seem to have a natural generalization to the setting of general acyclic quivers. However, a tree quiver Q can be naturally seen as a partial order (Q_0, \rightsquigarrow) where $x \rightsquigarrow y$ if there is a (necessarily unique) path from x to y . Then, the path algebra of Q is isomorphic to the incidence algebra of (Q_0, \rightsquigarrow) . Our experimentation suggests that this may be a good setting for a generalization of Theorem 6.1.

Let $n \in \mathbb{N}$ be an even integer and Z_n be the following zigzag orientation of an affine Dynkin diagram of type \mathbb{A} .

$$(6.2) \quad \begin{array}{ccccccc} & & 1 \circ & \longrightarrow & \circ & 2 & \\ & & \swarrow & & & \nwarrow & \\ & n \circ & & & & & \circ 3 \\ & & \uparrow & & & & \downarrow \\ & n-1 \circ & & & & & \circ 4 \\ & & \swarrow & & & & \dots \\ & & n-2 \circ & & & & \end{array}$$

We may view Z_n as a quiver, or as the Hasse diagram of a poset. Even for tree quivers, the language of posets is more flexible: for example, it is easy to see that Z_4 is not a subquiver of \tilde{D}_4 but it is a (full) subposet of $(\tilde{D}_4, \rightsquigarrow)$.

We remind the reader that (P, \leq_P) is a *subposet* of (Q, \leq_Q) if $P \subseteq Q$ and the inclusion $P \hookrightarrow Q$ is a morphism of posets. Now we recall the following central definitions for this section.

Definition 6.2. Let (P, \leq_P) and (Q, \leq_Q) be two posets and $\phi: P \rightarrow Q$ a morphism of posets. Then,

- (1) The map ϕ is *full* if for any $p_1, p_2 \in P$, $p_1 \leq_P p_2$ holds if and only if $\phi(p_1) \leq_Q \phi(p_2)$ holds.
- (2) If ϕ is full and injective, the image of ϕ is called a *full subposet* of (P, \leq_P) .

We remark that some authors call *weak* subposets our subposets, and full subposets are also known as *induced* subposets.

Let (P, \leq) be a poset. For $i \leq j$ in P , an interval $[i, j]$ is always considered with respect to \leq , that is, $[i, j] := \{k \in P \mid i \leq k \leq j\}$.

Definition 6.3. Let (P, \leq) be a finite poset. The *incidence algebra* of P over \mathbf{k} , denoted by $A(P)$ is the \mathbf{k} -vector space with basis the set of intervals $[i, j]$ in the poset P with multiplication induced by $[i, j] \cdot [k, l] = [i, l]$ if $j = k$, and 0 otherwise.

Then, we propose the following generalization of Theorem 6.1.

Conjecture 6.4. Let (P, \leq) be a finite poset. Then, the poset of quasi-hereditary structures on the incidence algebra of (P, \leq) is a lattice if and only if Z_n is not isomorphic to a full subposet of (P, \leq) for any $n \geq 4$.

Actually, the setting of incidence algebras is not only a good setting for a generalization of our result, it also simplifies our arguments. So we prove Theorem 6.1 as a corollary of the following theorem.

Theorem 6.5. *Let (P, \leq) be a finite poset. We assume that the incidence algebra $A(P)$ of (P, \leq) is hereditary. Then, the poset of quasi-hereditary structures on $A(P)$ is a lattice if and only if Z_n is not isomorphic to a full subposet of (P, \leq) for any $n \geq 4$.*

Proof. The only if part follows from Proposition 6.11 and Theorem 6.21. The if part follows from Theorem 6.22. \square

We prove Theorem 6.21 in Subsection 6.2 and show Theorem 6.22 in Subsection 6.3. For Theorems 6.1 and 6.5 we need the following Lemmas. The precise comparison between Theorem 6.1 and Theorem 6.5 is given in Remark 6.9.

Lemma 6.6. *Let Q be a finite quiver whose underlying graph is a tree. Then the following statements hold.*

- (1) Z_n is not a full subposet of (Q_0, \rightsquigarrow) for any even $n \geq 6$.
- (2) Z_4 is not a full subposet of (Q_0, \rightsquigarrow) if and only if \tilde{D}_n is not a full subquiver of Q for any $n \geq 4$.

Proof. (1) Assume that there exists a subset $Z = \{1, 2, \dots, n\}$ of Q_0 such that $(Z, \rightsquigarrow|_Z)$ is isomorphic to Z_n . So we have paths in Q of the form:

$$n \leftarrow 1 \rightsquigarrow 2 \leftarrow \dots \rightsquigarrow n-2 \leftarrow n-1 \rightsquigarrow n.$$

For the rest of this proof, integers are considered modulo n . For each odd k , there exist a vertex $k' \in Q_0$ and three paths $k \rightsquigarrow k'$, $k' \rightsquigarrow (k-1)$ and $k' \rightsquigarrow (k+1)$ such that $k' \rightsquigarrow (k-1)$ and $k' \rightsquigarrow (k+1)$ have no common vertices except k' . Similarly, for each even k , there exist a vertex k' and three paths $(k-1) \rightsquigarrow k'$, $(k+1) \rightsquigarrow k'$ and $k' \rightsquigarrow k$ such that $(k-1) \rightsquigarrow k'$ and $(k+1) \rightsquigarrow k'$ have no common vertices except k' . In particular, we have the following paths:

$$\begin{array}{ccccccccccc} n' & \leftarrow & 1 & \rightsquigarrow & 2' & \leftarrow & 3 & \rightsquigarrow & \dots & \rightsquigarrow & (n-2)' & \leftarrow & (n-1) & \rightsquigarrow & n' \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\ n & \leftarrow & 1' & \rightsquigarrow & 2 & \leftarrow & 3' & \rightsquigarrow & \dots & \rightsquigarrow & (n-2) & \leftarrow & (n-1)' & \rightsquigarrow & n \end{array}$$

Let k be an odd integer. Since Q is tree, both k' and $(k+1)'$ appear in the path $k \rightsquigarrow (k+1)$. If there exists a path from $(k+1)'$ to k' , then there exists a path from $(k+2)$ to $(k-1)$. This is a contradiction, since Z_n is a full subposet of (Q_0, \rightsquigarrow) and $n \geq 6$. Therefore we assume that there exists a path from k' to $(k+1)'$. Similarly, for even k , we assume that there exists a path from $(k+1)'$ to k' . We have paths as follows:

$$n' \leftarrow 1' \rightsquigarrow 2' \leftarrow \dots \rightsquigarrow (n-2)' \leftarrow (n-1)' \rightsquigarrow n'.$$

Because of the construction of k' , any two adjacent paths have no common vertices. This is a contradiction, since Q is tree.

(2) Assume that \tilde{D}_n is a full subquiver of Q for some n . Let $Z = \{1, 2, 3, 4\}$ be the four vertices of \tilde{D}_n such that each of them is a sink or a source. Then $(Z, \rightsquigarrow|_Z)$ is isomorphic to Z_4 . Conversely, assume that Z_4 is a full subposet of (Q_0, \rightsquigarrow) . Then the same argument in the proof of (1) implies that \tilde{D}_n is a full subquiver of Q for some n . \square

We first observe under which conditions an incidence algebra is hereditary and we collect some basic properties of such an incidence algebra. For that, we recall the next concept.

Let (P, \leq) be a finite poset. We say that P is *diamond-free* if it does not have four elements a, b, c , and d forming a diamond suborder with $a \leq b \leq d$ and $a \leq c \leq d$ and with b and c incomparable. The Hasse diagram of this type of poset is also called a *multi-tree*.

Lemma 6.7. *Let (P, \leq) be a finite poset. Then*

- (1) *The algebra $A(P)$ is hereditary if and only if P is a diamond-free poset.*
- (2) *If (Q, \leq) is a full subposet of (P, \leq) and $A(P)$ is a hereditary algebra, then $A(Q)$ is a hereditary algebra.*

Proof. Recall that the incidence algebra of a finite poset is isomorphic to the quotient of the path algebra of its Hasse quiver modulo the relations of total commutativity. As a consequence, if (P, \leq) is a diamond-free poset, there is no relation to mod out, so the incidence algebra is hereditary. Conversely, assume that there are a, b, c and d in P such that $a \leq b, c \leq d$ and b and c are incomparable with respect to \leq . Let M be the submodule of P_a supported by the elements larger or equal than b or c . Its top is S_b and S_c , so its projective cover is $P_b \oplus P_c$. However S_d appears twice in this direct sum but only once in P_a . So P_a has a submodule which is not projective and the algebra is not hereditary. The second statement of the lemma is a direct consequence of the first. \square

Lemma 6.8. *Let (P, \leq) be a connected finite poset which is diamond-free and has no Z_n as full subposet for any $n \geq 4$. Then, its Hasse diagram is a tree.*

Proof. We only sketch the proof. Assume that there is a cycle in the unoriented Hasse diagram of (P, \leq) . Since (P, \leq) is a poset, it is neither an oriented cycle nor a 3-cycle. The cycle has at least one source and one sink. If it has exactly one source and one sink, then it is not diamond-free. So, it has at least two sources and two sinks, and the number of sources is equal to the number of sinks. Then, the subset consisting of all the sources and sinks of the cycle induces a full subposet isomorphic to Z_n for some $n \geq 4$. \square

Remark 6.9. As a consequence of Lemmas 6.6, 6.8 and Theorem 6.5 we see that:

- (1) An *hereditary* incidence algebra of a finite poset (P, \leq) has a *lattice* of quasi-hereditary structures if and only if the Hasse quiver of (P, \leq) is a tree which does not have \tilde{D}_n as subquivers for any $n \geq 4$.
- (2) On the other hand, Theorem 6.5 is slightly stronger than Theorem 6.1 since it allows to treat more general quivers such as 6.2.

We list easy properties about the incidence algebra of a diamond-free poset. All the properties are easy to show and we omit the proof, and we use the following lemma without referring.

Lemma 6.10. *Let (P, \leq) be a finite diamond-free poset and $i, j \in P$. The following statements hold.*

- (1) *If $i \leq j$, then $([i, j], \leq)$ is a total order. In other words, in the Hasse quiver of (P, \leq) there is a unique path from i to j .*
- (2) *If $i \leq j$, then there exists an $A(P)$ -module with simple top $S(i)$ and simple socle $S(j)$, and such that a simple module $S(k)$ is a composition factor of it if and only if $i \leq k \leq j$ holds.*
- (3) *Let L be a submodule of an indecomposable projective $A(P)$ -module. Then any non-zero indecomposable direct summand of L has a simple top.*

We end this subsection by showing the following key result.

Proposition 6.11. *For an even integer $n \geq 4$, the poset of quasi-hereditary structures on the path algebra $\mathbf{k}Z_n$ is not a lattice.*

Proof. We assume that the quiver is labeled as in (6.2). This means that the sources of the quiver are labeled by odd integers and the sinks by even integers. Moreover we view the labels as elements of $\mathbb{Z}/n\mathbb{Z}$ in order to have a cyclic labeling.

The projective indecomposable $\mathbf{k}Z_n$ -modules indexed by sinks are simple and the projective indecomposable module indexed by a source i has top $S(i)$ and socle $S(i-1) \oplus S(i+1)$.

Let x and y be the labels of two distinct sources of Z_n . For $s \in \{x, y\}$ we consider the total order \triangleleft_s defined by

$$s \triangleleft_s s+1 \triangleleft_s \dots \triangleleft_s s-1$$

These two orderings are total, so they are adapted and since the algebra is hereditary they represent two quasi-hereditary structures. For $s \in \{x, y\}$, we denote by Δ_s the corresponding set of standard modules. The standard modules indexed by sinks are simple, the standard module indexed by s is also simple and the standard modules indexed by the other sources v are uniserial with top $S(v)$ and socle $S(v-1)$. Since x and y are distinct, we see that the orderings \triangleleft_x and \triangleleft_y represent two different quasi-hereditary structures.

For a sink i we let $\theta(i) = S(i)$ and for a source i we let $\theta(i)$ be the uniserial module with top $S(i)$ and socle $S(i-1)$ and we denote by Θ the set consisting of the modules $\theta(i)$ for $i \in \{1, 2, \dots, n\}$.

For $s \in \{x, y\}$, we have $\mathcal{F}(\Theta) \subseteq \mathcal{F}(\Delta)$. Indeed, for $i \neq s$ we have $\Delta_s(i) = \theta(i)$ and $\theta(s)$ clearly has a Δ_s -filtration. Conversely, we show by induction on the length of M that if $M \in \mathcal{F}(\Delta_x) \cap \mathcal{F}(\Delta_y)$, then $M \in \mathcal{F}(\Theta)$. This is clearly true if M is simple. Let

$$0 = M_0 \subset M_1 \subset \dots \subset M_{n-1} \subset M_n = M$$

be a Δ_x -filtration of M . Then we have $M/M_{n-1} \cong \Delta_x(i)$ for some $i \in \{1, 2, \dots, n\}$. If $i \neq x$ then $\Delta_x(i) \in \mathcal{F}(\Delta_y)$. Since M is also in $\mathcal{F}(\Delta_y)$ and this category is closed under kernel of epimorphism, we obtain that $M_{n-1} \in \mathcal{F}(\Delta_x) \cap \mathcal{F}(\Delta_y)$. By induction, the module M_{n-1} is in $\mathcal{F}(\Theta)$. Since the category $\mathcal{F}(\Theta)$ is closed under extensions and $\Delta_x(i) \in \mathcal{F}(\Theta)$, we conclude that $M \in \mathcal{F}(\Theta)$. If $i = x$, then M has the simple module $S(x)$ as direct summand of its top. So there is a Δ_y -filtration of M where the standard module $\Delta_y(x)$ appears as first factor. That is, there is $N \in \mathcal{F}(\Delta_y)$ such that $M/N \cong \Delta_y(x)$. Since $x \neq y$, the module $\Delta_y(x)$ is Δ_x -filtered and we use the same argument as above to conclude that $M \in \mathcal{F}(\Theta)$.

Let us remark that there is no adapted order that have Θ as set of standard modules. Indeed, let us assume that there is such an ordering \triangleleft . The shape of the modules implies that if i is an odd integer, we have $i-1 \triangleleft i$ and $i+1 \not\triangleleft i$. Since \triangleleft is adapted, we must have $i \triangleleft i+1$. It follows that $n \triangleleft 1$ and $1 \triangleleft 2 \triangleleft 3 \triangleleft \dots \triangleleft n$ contradicting the antisymmetry of \triangleleft .

Let us finish the proof by showing that if the join \triangleleft_J of \triangleleft_x and \triangleleft_y in the poset of quasi-hereditary structures exists, it must have Θ as set of standard modules, contradicting the fact that no adapted order have this set as set of standard modules.

For this, we consider $z = x+1$ and $w = y+1$ the sinks that are respectively after x and y in the cyclic ordering of Z_n . As before, following the cyclic orders with starting point x or y gives two total orderings \triangleleft_z and \triangleleft_t where the minimal elements are respectively z and t and the maximal elements are respectively x and y . For $s \in \{z, t\}$ the standard modules Δ_s are as follows: the standard modules indexed by sinks are simple, the standard module indexed by the source $s-1$ (which is respectively x and y) are projective and the standard modules indexed another source i are uniserial with top $S(i)$ and socle $S(i-1)$. It is straightforward to check that $\mathcal{F}(\Delta_s) \subseteq \mathcal{F}(\Delta_x) \cap \mathcal{F}(\Delta_y)$ for $s \in \{z, t\}$.

Let us now assume that \triangleleft_x and \triangleleft_y have a join \triangleleft_J in the poset of quasi-hereditary structures. Then we have

$$\mathcal{F}(\Delta_z) \cup \mathcal{F}(\Delta_t) \subseteq \mathcal{F}(\Delta_J) \subset \mathcal{F}(\Delta_x) \cap \mathcal{F}(\Delta_y) = \mathcal{F}(\Theta).$$

Since every module $\theta(i)$ appears in Δ_z or Δ_t , we have $\mathcal{F}(\Theta) = \mathcal{F}(\Delta_J)$. For $i \in \{1, 2, \dots, n\}$, we see $\theta(i) \in \mathcal{F}(\Delta_J)$ implies that $\Delta_J(i)$ is a quotient of $\theta(i)$ since the only module in Θ with top $S(i)$ is $\theta(i)$. Similarly $\Delta_J(i) \in \mathcal{F}(\Theta)$ implies that $\theta(i)$ is a quotient of $\Delta_J(i)$. So, $\Theta = \Delta_J$ and if the join of \triangleleft_x and \triangleleft_y exists, it has Θ as set of standard modules. \square

6.2. Quasi-hereditary structures of full subposets. In this subsection, we construct a morphism between the sets of quasi-hereditary structures on the incidence algebras of two finite posets such that one is a full subposet of the other (Proposition 6.18). This map will be used to prove the ‘only if’ part of Theorem 6.5, see Theorem 6.21.

We fix, for all this subsection, the following setting and notation: Let (P, \leq) be a finite poset and (Q, \leq) be a full subposet. We fix \triangleleft an adapted poset to the incidence algebra of Q . We let $R = P \setminus Q$. The relation \leq induces a structure of poset on R .

We consider the binary relation $\widehat{\triangleleft}$ on P defined as follows:

- (1) For $r_1, r_2 \in R$, $r_1 \widehat{\triangleleft} r_2$ if and only if $r_1 \leq r_2$.
- (2) For $q_1, q_2 \in Q$, $q_1 \widehat{\triangleleft} q_2$ if and only if $q_1 \triangleleft q_2$.

In other words $\widehat{\triangleleft}$ restricts as \leq on R and \triangleleft on Q . There is *no* relations of the form $q \widehat{\triangleleft} r$ with $r \in R$ and $q \in Q$. The relations $r \triangleleft q$ are of two possible shapes:

- (3) If $r \leq q$ then $r \widehat{\triangleleft} q$ if and only if there exists $q_1 \in [r, q] \cap Q$ such that $q_1 \triangleleft q$ and $[r, q_1] \cap Q = \{q_1\}$.
- (4) If $q \leq r$ then $r \widehat{\triangleleft} q$ if and only if there exists $q_1 \in [q, r] \cap Q$ such that $q_1 \triangleleft q$ and $[q_1, r] \cap Q = \{q_1\}$.

These two conditions may look unnatural for now, but we will see in Lemma 6.17 that they induce a natural set of standard modules.

Lemma 6.12. *The transitive cover of $\widehat{\triangleleft}$ is a partial order on P .*

Proof. For this proof we use the symbol $\widehat{\triangleleft}$ for the relation described above and $\widehat{\triangleleft}^{tc}$ for its transitive cover. Let us verify that it is an antisymmetric relation. Let $x, y \in P$ with $x \widehat{\triangleleft}^{tc} y$. Then, there is $a_1, \dots, a_n \in P$ such that

$$x \widehat{\triangleleft} a_1 \widehat{\triangleleft} \dots \widehat{\triangleleft} a_n \widehat{\triangleleft} y.$$

If there is $i \in \{1, \dots, n\}$ such that $a_i \in Q$, then a_{i+1}, \dots, a_n and y are in Q and $a_i \triangleleft y$ by transitivity of the partial order \triangleleft on the elements of Q . So, if $x \in Q$, then $y \in Q$ and we have $x \triangleleft y$. If $y \widehat{\triangleleft}^{tc} x$, then by the same argument $y \triangleleft x$ and $y = x$ since \triangleleft is antisymmetric. If $x \in R$ and there exists $i \in \{1, 2, \dots, n\}$ such that $a_i \in Q$, then $y \in Q$ and $y \widehat{\triangleleft}^{tc} x$.

If $x \in R$ and all the a_i 's are in R , then $x \leq y$. If $y \widehat{\triangleleft}^{tc} x$, then there exist b_1, \dots, b_r in P such that $y \widehat{\triangleleft} b_1 \widehat{\triangleleft} b_2 \dots \widehat{\triangleleft} b_r \widehat{\triangleleft} x$. Since $x \in R$, all the b_i 's are in R and $y \leq x$. So $x = y$ by antisymmetry of \leq . This shows that $\widehat{\triangleleft}^{tc}$ is antisymmetric and it is by construction reflexive and transitive. \square

From now on the symbol $\widehat{\triangleleft}$ is used for the partial order of Lemma 6.12.

Lemma 6.13. *Let (P, \leq) be a finite poset and (Q, \leq) be a full subposet of (P, \leq) . Let \triangleleft be an adapted poset to $A(Q)$. Then $\widehat{\triangleleft}$ is an adapted poset to $A(P)$.*

Proof. Let $x, y \in P$ such that $x \leq y$. If x and y are both in R , then $x \widehat{\triangleleft} y$. If x and y are both in Q , there exists $q_1 \in [x, y] \cap Q$ such that $x \triangleleft q_1$ and $y \triangleleft q_1$. Since $\widehat{\triangleleft}$ coincides with \triangleleft on the elements of Q , we have $x \widehat{\triangleleft} q_1$ and $y \widehat{\triangleleft} q_1$.

We assume that $x = r \in R$ and $y = q \in Q$ and r and q are incomparable with respect to $\widehat{\triangleleft}$. Let $q_1 \in Q$ such that $r \leq q_1 \leq q$ and q_1 is minimal for this property. Then $r \widehat{\triangleleft} q_1$. Now the poset \triangleleft is adapted so there exists $q_2 \in [q_1, q]$ such that $q_1 \triangleleft q_2$ and $q \triangleleft q_2$. By transitivity, we have $r \widehat{\triangleleft} q_2$ and $q \widehat{\triangleleft} q_2$. The case $x \in Q$ and $y \in R$ is similar. \square

Let us describe the increasing and decreasing relations for $\widehat{\triangleleft}$.

Lemma 6.14. *Let $r \in R$ and $q \in Q$. Then*

- (1) $r \widehat{\triangleleft} q$ is increasing if and only if $r \leq q$ and

$$\forall q_1 \in [r, q] \cap Q, q_1 \triangleleft q.$$

- (2) $r \widehat{\triangleleft} q$ is decreasing if and only if $q \leq r$ and

$$\forall q_1 \in [q, r] \cap Q, q_1 \triangleleft q.$$

Proof. We only prove the second point since the proof of the first is similar. If $r \widehat{\triangleleft} q$ is decreasing, then we have $q \leq r$ and $\forall x \in [q, r]$, we have $x \widehat{\triangleleft} q$. This is in particular true for $x \in Q$. Since $\widehat{\triangleleft}$ restricts as \triangleleft on the elements of Q , we have $x \triangleleft q$. Conversely assume that $q \leq r$ and $\forall q_1 \in [q, r] \cap Q, q_1 \triangleleft q$. Let $x \in [q, r]$. Let $q_1 \in Q$ such that $q \leq q_1 \leq x$ and q_1 is maximal for this property. Then, $q_1 \triangleleft q$ by hypothesis, and $x \widehat{\triangleleft} q$ by definition of $\widehat{\triangleleft}$. \square

Remark 6.15. We see that our rather technical definition of $\widehat{\triangleleft}$ leads to natural increasing and decreasing relations: they are the relations that are increasing or decreasing when forgetting the elements which are not in Q .

To describe precisely the standard modules associated with this partial order we consider the functors in the right part of the classical idempotent recollement associated to the idempotent $e_Q = \sum_{q \in Q} [q, q] \in A(P)$ corresponding to Q .

Lemma 6.16. *The algebra $e_Q A(P) e_Q$ is isomorphic to $A(Q)$.*

Proof. The incidence algebra of P over \mathbf{k} has basis the set of intervals $[x, y]$ such that $x \leq_P y$ are two elements of P . Multiplying on the right and the left by e_Q , we have a basis consisting of intervals $[x, y]$ such that $x, y \in Q$ and $x \leq_P y$. Since Q is a full subposet of P , this is exactly the set of intervals $[x, y]$ such that $x, y \in Q$ and $x \leq_Q y$. \square

There is a ‘restriction’ functor $(-)_e_Q: \text{mod } A(P) \rightarrow \text{mod } A(Q)$ which sends an $A(P)$ -module M to $M e_Q$. It has a left adjoint $L = - \otimes_{A(Q)} e_Q A(P)$. We recall without proofs the following well-known, and easy to check, properties of these functors:

- (1) The functor $(-)_e_Q$ is exact.
- (2) The functor L is right exact, fully faithful and sends projective modules to projective modules.
- (3) $(-)_e_Q \circ L \cong \text{Id}_{\text{mod } A(Q)}$.

Lemma 6.17. *Let (P, \leq) be a finite poset and (Q, \leq) be a full subposet. Let \triangleleft be an adapted poset to $A(Q)$. We denote by Δ its set of standard modules. We denote by $\widehat{\Delta}$ the set of standard modules for $A(P)$ corresponding to $\widehat{\triangleleft}$. Then,*

- (1) *If $r \in P \setminus Q$, then $\widehat{\Delta}(r) = S(r)$.*
- (2) *If $q \in Q$, then $\widehat{\Delta}(q) = L(\Delta(q))$.*

Proof. Let $r \in R$. If $x \widehat{\triangleleft} r$, then $x \in R$ and $x \leq r$. Since the composition factors of $P(r)$ are indexed by the elements larger than r in P , we have the first point.

Let $q \in Q$. We know that the composition factors of $\widehat{\Delta}(q)$ are indexed by the elements $x \in P$ such that $x \triangleleft q$ is a decreasing relation. Lemma 6.14 gives us the decreasing relations. It remains to compute the composition factors of $L(\Delta(q))$ and check that the two sets coincide. Recall that $\Delta(q)$ is the largest quotient of $P(q)$ having composition factors indexed by elements q' such that $q' \triangleleft q$. Let \mathfrak{m} be the set of q' such that $q \leq q'$, $q' \not\triangleleft q$ and q' is minimal for this property. Then,

$$\bigoplus_{x \in \mathfrak{m}} P(x) \rightarrow P(q) \rightarrow \Delta(q) \rightarrow 0,$$

is the beginning of a (minimal) projective presentation of $\Delta(q)$. Applying the functor L we get that

$$\bigoplus_{x \in \mathfrak{m}} P(x) \rightarrow P(q) \rightarrow L(\Delta(q)) \rightarrow 0,$$

is the beginning of a projective presentation of $L(\Delta(q))$. Let us also recall that all the non-zero morphisms between projective indecomposable modules are injective. So, to compute the composition factors of $L(\Delta(q))$ we have to compute the composition factors of $P(q)$ which are not in $P(x)$ for $x \in \mathfrak{m}$. If $S(y)$ is a composition factor of $P(x)$ for $x \in \mathfrak{m}$, then $q \leq x \leq y$ and $x \not\triangleleft q$. Conversely we assume that there is $q_1 \in Q$ such that $q \leq q_1 \leq y$ and $q_1 \not\triangleleft q$. If q_1 is minimal for this property $q_1 \in \mathfrak{m}$. If not, let q_2 such that $q \leq q_2 \leq q_1$ such that $q_2 \not\triangleleft q$ and q_2 is minimal for this property. Then $q_2 \in \mathfrak{m}$ and $S(y)$ is a composition factor of $P(q_2)$. This shows that the composition factors of $L(\Delta(q))$ are indexed by the elements x such that $q \leq x$ and $\forall q_1 \in [q, x] \cap Q$, $q_1 \triangleleft q$. The result now follows from Lemma 6.14. \square

Proposition 6.18. *Let (P, \leq) be a finite poset such that $A(P)$ is a hereditary algebra. Let (Q, \leq) be a full subposet of Q . Then $\triangleleft \mapsto \widehat{\triangleleft}$ induces a well-defined full embedding of posets from $\text{qh.str}(A(Q))$ to $\text{qh.str}(A(P))$.*

Proof. Since the algebra $A(P)$ is hereditary, all the adapted orders give rise to quasi-hereditary structures.

Lemma 6.17 implies that the mapping is compatible with the equivalence relation. Lemma 6.17 and $(-)_e_Q \circ L \cong \text{Id}_{\text{mod } A(Q)}$ implies that it is injective. Lemma 2.21 and right exactness of L implies that it is a morphism of posets and $(-)_e_Q \circ L \cong \text{Id}_{\text{mod } A(Q)}$ implies that it is full. \square

Definition 6.19. Let $\phi: (O_1, \leq_1) \rightarrow (O_2, \leq_2)$ be a morphism of posets. Then ϕ is *interval-preserving* if for every $x, y \in O_1$ and $z \in O_2$ we have $\phi(x) \leq_2 z \leq_2 \phi(y)$ if and only if there is $z' \in O_1$ such that $x \leq_1 z' \leq_1 y$ and $\phi(z') = z$.

Proposition 6.20. *The morphism $[\triangleleft] \mapsto [\widehat{\triangleleft}]$ from $\text{qh.str}(A(Q))$ to $\text{qh.str}(A(P))$ of Proposition 6.18 is interval-preserving.*

Proof. Let \triangleleft be an adapted order such that $[\triangleleft_1] \preceq [\triangleleft] \preceq [\triangleleft_2]$. We use the following notation: $\Delta_1, \nabla_1, \Delta, \nabla, \Delta_2, \nabla_2$ for the set of standard and costandard modules respectively induced by $[\triangleleft_1], [\triangleleft]$ and $[\triangleleft_2]$.

Using Lemma 2.21, for $x \in P$ we have:

$$\Delta_2(x) \twoheadrightarrow \Delta(x) \twoheadrightarrow \Delta_1(x),$$

and

$$\nabla_2(x) \hookrightarrow \nabla(x) \hookrightarrow \nabla_1(x).$$

In particular for $r \in R := P \setminus Q$, we have $\Delta(r) = S(r)$ and there is no composition factor of ∇_r indexed by an element of Q since the elements of Q are not smaller for $\widehat{\triangleleft}_1$ than the elements of R .

For $q \in Q$, the beginning of a minimal projective presentation of $\Delta(q)$ is

$$(6.3) \quad \bigoplus_{y \in \mathfrak{m}} P(y) \rightarrow P(q) \rightarrow \Delta(q) \rightarrow 0,$$

where \mathfrak{m} is the set of $y \in P$ such that $q \leq y$, $y \not\triangleleft q$ and y is minimal (with respect to \leq) for this property. Since \triangleleft is an adapted order, we have $q \triangleleft y$ or there is $q < x < y$ such that $q \triangleleft x$ and $y \triangleleft x$. The second possibility contradicts the minimality of y , so we have $q \triangleleft y$. Moreover, by minimality of y , if $q < x < y$, we have $x \triangleleft q \triangleleft y$. This implies that the relation $q \triangleleft y$ is *increasing*. In other words, the simple module $S(q)$ is a composition factor of $\nabla(y)$. As explained above the composition factors of the costandard modules indexed by the elements of R are in R , so we must have $y \in Q$. Then, by the presentation (6.3) and an isomorphism $L(e_Q A(P) e_Q) \cong e_Q A(P)$, we obtain $\Delta(q) = L(\Delta(q) e_Q)$.

In order to finish the proof, we have to show that there is a quasi-hereditary structure on $A(Q)$ which has $\{\Delta(q) e_Q ; q \in Q\}$ as set of standard modules. The obvious candidate is $\triangleleft|_Q$ the restriction of the poset \triangleleft to the elements of Q . For $q \in Q$, we denote by $\Delta'(q)$ the standard module induced by the poset $\triangleleft|_Q$. Then the beginning of a minimal projective presentation of $\Delta'(q)$ is

$$\bigoplus_{q' \in \mathfrak{m}'} P(q') \rightarrow P(q) \rightarrow \Delta'(q) \rightarrow 0,$$

where \mathfrak{m}' is the set of $q' \in Q$ such that $q \leq q'$, $q' \not\triangleleft q$ and q' is minimal (with respect to \leq) for this property. As explained above, this set equals \mathfrak{m} . Since the functor $(-)_e_Q$ is exact, by applying it to the exact sequence (6.3), we get $\Delta'(q) = \Delta(q) e_Q$.

Now we check that $[\triangleleft|_Q]$ is a quasi-hereditary structure for $A(Q)$. For $q \in Q$, we let

$$0 \subset M_0 \subset M_1 \subset \cdots \subset M_n = P(q)$$

be a Δ -filtration. Since the functor $(-)_e_Q$ is exact we have:

$$0 \subset M_0 e_Q \subset M_1 e_Q \subset \cdots \subset M_n e_Q = P(q)$$

For $i \in \{1, \dots, n\}$ we have $(M_i e_Q)/(M_{i-1} e_Q) \cong \Delta(x)$ for $x \in P$. If $x \in R$, we have $\Delta(x) = S(x)$ and $\Delta(x) e_Q = 0$. So $M_i e_Q = M_{i-1} e_Q$. In this case, we remove M_{i-1} from the sequence of submodules. By induction, we end up with

$$0 \subset M_{i_0} e_Q \subset M_{i_1} e_Q \subset \cdots \subset M_{i_r} e_Q = P(q)$$

$M_{i_j} e_Q / M_{i_{j-1}} e_Q \cong \Delta(q) e_Q$ for some $q \in Q$. This proves that $[\triangleleft|_Q]$ is a quasi-hereditary structure for $A(Q)$ and $[\triangleleft] = ([\widehat{\triangleleft|_Q}])$.

Finally we have to check that $[\triangleleft_1] \leq [\triangleleft|_Q] \leq [\triangleleft_2]$. This is an easy consequence of Lemma 2.21 and Lemma 6.7 \square

Theorem 6.21. *Let (P, \leq) be a finite poset such that $A(P)$ is a hereditary algebra. Let (Q, \leq) be a full subposet of (P, \leq) such that $\text{qh.str}(A(Q))$ is not a lattice. Then $\text{qh.str}(A(P))$ is not a lattice.*

Proof. Let $[\triangleleft_1]$ and $[\triangleleft_2]$ be two elements of $\mathbf{qh.str}(A(Q))$. It is easy to check that \leq^{op} induces a quasi-hereditary structure on $A(P)$ with simple modules as co-standard modules. As a consequence, it is the greatest element of $\mathbf{qh.str}(A(Q))$. Since $\mathbf{qh.str}(A(Q))$ has a greatest element, the set $U = \{[\triangleleft] : [\triangleleft_1] \preceq [\triangleleft] \text{ and } [\triangleleft_2] \preceq [\triangleleft]\}$ is non empty. We assume that the two elements have no join, then this set has at least two minimal elements $[\triangleleft_x]$ and $[\triangleleft_y]$. Using Proposition 6.18, we see that $[\widehat{\triangleleft}_i] \preceq [\widehat{\triangleleft}_s]$ in $\mathbf{qh.str}(A(P))$ for $i = 1, 2$ and $s = x, y$. So if $[\widehat{\triangleleft}_1]$ and $[\widehat{\triangleleft}_2]$ have a join $[\triangleleft']$, then $[\triangleleft']$ is larger than $[\widehat{\triangleleft}_i]$ for $i = 1, 2$ and smaller than $[\widehat{\triangleleft}_s]$ for $s = x, y$. Looking at $[\triangleleft'] \in [[\widehat{\triangleleft}_1], [\widehat{\triangleleft}_x]]$, Proposition 6.20 implies that there is $[\triangleleft] \in [[\triangleleft_1], [\triangleleft_x]]$ such that $[\triangleleft'] = [\triangleleft]$. We can repeat the same arguments with the other intervals involving $[\triangleleft']$ and since $[\triangleleft] \mapsto [\widehat{\triangleleft}]$ is injective, we see that $[\triangleleft]$ is larger than both $[\triangleleft_1]$, $[\triangleleft_2]$ and smaller than $[\triangleleft_x]$ and $[\triangleleft_y]$. This is a contraction since $[\triangleleft_x]$ and $[\triangleleft_y]$ are incomparable and minimal with respect to the property of being larger than $[\triangleleft_1]$ and $[\triangleleft_2]$. If all the pairs of elements of $\mathbf{qh.str}(A(Q))$ have a join, but two elements have no meet, the proof is dual. \square

6.3. Quasi-hereditary structures of diamond-free posets. In this subsection we prove the reciprocity of Theorem 6.5.

Theorem 6.22. *Let (P, \leq) be a finite poset such that $A(P)$ is a hereditary algebra. If Z_n is not a full subposet of (P, \leq) for any $n \geq 4$, then $\mathbf{qh.str}(A(P))$ is a lattice and the join and the meet of two quasi-hereditary structures $[\triangleleft_1]$ and $[\triangleleft_2]$ are represented by the following partial orders*

$$\begin{aligned} \triangleleft_1 \wedge \triangleleft_2 &= \left((\text{Dec}(\triangleleft_1) \cap \text{Dec}(\triangleleft_2)) \cup \text{Inc}(\triangleleft_1) \cup \text{Inc}(\triangleleft_2) \right)^{\text{tc}}, \\ \triangleleft_1 \vee \triangleleft_2 &= \left(\text{Dec}(\triangleleft_1) \cup \text{Dec}(\triangleleft_2) \cup (\text{Inc}(\triangleleft_1) \cap \text{Inc}(\triangleleft_2)) \right)^{\text{tc}}. \end{aligned}$$

Our strategy for the proof is as follows: first we prove that the candidate for the meet is a partial order (this is the technical part, Proposition 6.27) and then we prove that it induces a quasi-hereditary structure and check that it gives a meet of $[\triangleleft_1]$ and $[\triangleleft_2]$ in the poset of quasi-hereditary structures (Proposition 6.30). Since $\mathbf{qh.str}(A(Q))$ is a finite poset with a greatest element, this is enough to prove that it is a lattice.

For the rest of this subsection, we fix a finite poset (P, \leq) and fix two adapted orders \triangleleft_1 and \triangleleft_2 on P . Let

$$\triangleleft' = \left((\text{Dec}(\triangleleft_1) \cap \text{Dec}(\triangleleft_2)) \cup \text{Inc}(\triangleleft_1) \cup \text{Inc}(\triangleleft_2) \right)^{\text{tc}}.$$

We denote by Δ_1, Δ_2 the set of standard $A(P)$ -modules associated to \triangleleft_1 and \triangleleft_2 , respectively. For $i, j \in P$, we write $i \triangleleft^D j$ if $(i, j) \in \text{Dec}(\triangleleft_1) \cap \text{Dec}(\triangleleft_2)$ and write $i \triangleleft^{I_m} j$ if $(i, j) \in \text{Inc}(\triangleleft_m)$ for $m = 1, 2$.

For $i, j \in P$ we have $j \triangleleft^D i$ if and only if $i \leq j$ and every $k \in [i, j]$ satisfies $k \triangleleft_1 i$ and $k \triangleleft_2 i$. Similarly, for $m = 1, 2$, $j \triangleleft^{I_m} i$ holds if and only if $i \leq j$ and every $k \in [i, j]$ satisfies $k \triangleleft_m i$.

The following lemma is obvious, but it is important to realize that without the diamond-free hypothesis it is false.

Lemma 6.23. *Let (P, \leq) be a diamond-free poset. Then \triangleleft^D is a transitive relation.*

Proof. Assume that $k \triangleleft^D j \triangleleft^D i$ for $i \leq j \leq k \in P$. Let $x \in [i, k]$. Since P is diamond-free, x is either smaller than j or larger than j . If x is smaller than j , then $x \triangleleft_m i$ holds for $m = 1, 2$. If x is larger than j , then $x \triangleleft_m j \triangleleft_m i$ holds for $m = 1, 2$. This implies $k \triangleleft^D i$. \square

Lemma 6.24. *Let $i, j \in P$. Assume that $i \leq j$ and any $k \in [i, j] \setminus \{j\}$ satisfies $k \triangleleft^D i$. Then one of $j \triangleleft^D i$, $i \triangleleft^{I_1} j$ or $i \triangleleft^{I_2} j$ holds.*

Proof. We used the argument that we proved in the proof of Proposition 6.20: if j is minimal for \leq such that $j \not\triangleleft_s i$, then $i \triangleleft_1 j \in \text{Inc}(\triangleleft_s)$ for $s = 1, 2$. So if $j \not\triangleleft^D i$, then either $j \triangleleft_1 i$ or $j \triangleleft_2 i$ holds. In the first case we have $i \triangleleft^{I_1} j$ holds and in the second case $i \triangleleft^{I_2} j$ holds. \square

Lemma 6.25. *Assume that there exists a sequence of elements of P such that*

$$i_0 \triangleleft^{I_{x(0)}} i_1 \triangleleft^{I_{x(1)}} \dots \triangleleft^{I_{x(\ell-1)}} i_\ell,$$

where $x(k) \in \{1, 2\}$ for $k = 0, \dots, \ell - 1$. If $i_\ell \triangleleft^D i_0$ holds, then $i_0 = i_1 = \dots = i_\ell$ holds.

Proof. Since $i_\ell \triangleleft^D i_0$ holds, we have $i_0 \leq i_\ell$. On the other hand, we have $i_0 \leq i_1 \leq \dots \leq i_\ell$. Since $i_1 \in [i_0, i_\ell]$ we have $i_1 \triangleleft_1 i_0$ and $i_1 \triangleleft_2 i_0$. By antisymmetry of \triangleleft_1 and \triangleleft_2 , we have $i_1 = i_0$. By induction on ℓ , we have the assertion. \square

In Lemma 6.26 and Proposition 6.27, we show that \triangleleft' is a partial order on P if Z_n is not a full subposet of (P, \leq) for any $n \geq 4$.

Lemma 6.26. *Let (P, \leq) be a finite poset and ℓ be an odd integer with $\ell \geq 3$. Assume that (P, \leq) does not have Z_n as a full subposet for any even $n \geq 4$. If there exist relations in P of the form:*

$$i_0 \geq i_1 \leq i_2 \geq \cdots \leq i_{\ell-1} \geq i_\ell \leq i_0,$$

then there exists $k \in \{0, 1, \dots, \ell\}$ such that we have at least one of the following $i_k \leq i_{k+2}$, $i_{k+2} \leq i_k$ where k considered modulo $\ell + 1$.

Proof. In this proof the labels are considered modulo $\ell + 1$. We prove the result by induction on $\ell \geq 3$.

We need a precise inductive hypothesis: let $Z = \{a_0, \dots, a_n\}$ be a subposet consisting of $n + 1$ consecutive elements of $\{i_0, \dots, i_\ell\}$ and which induces a subposet isomorphic to Z_{n+1} . Then, there is $k \in \{0, 1, \dots, n+1\}$ such that $a_k \leq a_{k+2}$ or $a_{k+2} \leq a_k$ and $\{a_k, a_{k+1}, a_{k+2}\}$ is a set of three consecutive elements of $\{i_0, \dots, i_\ell\}$.

If $n = 3$, then we have a subposet isomorphic to Z_4 which by hypothesis is not full. So, we must have a relation between a_0 and a_2 or a relation between a_1 and a_3 .

We assume now that $n \geq 5$ and that we have a subposet $Z = \{a_0, \dots, a_n\}$ isomorphic to Z_{n+1} . Without loss of generality we can assume that $a_i \leq a_{i+1}$ when i is odd. Since it is not full, there are two indices k and j with $k \notin \{j-1, j, j+1\}$ and a relation $a_k \leq a_j$ in (P, \leq) . If k is even, then $a_{k+1} \leq a_k \leq a_j$ and $a_{k-1} \leq a_k \leq a_j$. Moreover at least one of a_{k+1} and a_{k-1} is not in the set $\{a_{j-1}, a_j, a_{j+1}\}$. Replacing a_k by this element, we may assume k to be odd. Similarly, if j is even, then we have $a_k \leq a_j \leq a_{j+1}$ and $a_k \leq a_j \leq a_{j-1}$ and for at least one $j' \in \{j-1, j+1\}$, we have $k \notin \{j'-1, j', j'+1\}$. Changing a_j by $a_{j'}$, we may assume that j' is even.

The relation $a_k \leq a_j$ splits Z into two subposets Z_1 and Z_2 which are isomorphic to Z_{m_1} and Z_{m_2} for some integers m_1 and m_2 . By assumption k is odd and j is even, so we have $4 \leq m_1 < n$ and $4 \leq m_2 < n$. Precisely, Z_1 is the set consisting of the elements after or equal to a_k and before or equal to a_j in the cyclic ordering and Z_2 consists of the elements after or equal to a_j and before or equal to a_k in the cyclic ordering. One of Z_1 or Z_2 satisfies the hypothesis of consecutive elements and by induction we have the result. \square

Proposition 6.27. *Let (P, \leq) be a diamond-free poset. Assume that (P, \leq) does not have Z_n as a full subposet. Then for any two adapted orders \triangleleft_1 and \triangleleft_2 on P , a binary relation*

$$\triangleleft' = \left((\text{Dec}(\triangleleft_1) \cap \text{Dec}(\triangleleft_2)) \cup \text{Inc}(\triangleleft_1) \cup \text{Inc}(\triangleleft_2) \right)^{\text{tc}}$$

is a partial order on P .

Proof. By the definition, \triangleleft' is reflexive and transitive. We show that \triangleleft' is antisymmetric. It is enough to show the following claim: if there a sequence of vertices of P such that

$$(6.4) \quad i_0 \triangleleft^{X_0} i_1 \triangleleft^{X_1} \cdots \triangleleft^{X_{\ell-1}} i_\ell = i_0,$$

where $X_k \in \{D, I_1, I_2\}$ for $k = 0, \dots, \ell - 1$, then we have $i_0 = i_1 = \cdots = i_{\ell-1}$.

By Lemma 6.23 \triangleleft^D is transitive, so we may assume that there is no k with $X_k = X_{k+1} = D$. Let d be the number of $k = 0, \dots, \ell - 1$ such that $X_k = D$. We prove the claim by an induction on d . Assume that $d = 0$, then we have $i_0 \leq i_1 \leq \cdots \leq i_{\ell-1} \leq i_0$, so $i_0 = i_1 = \cdots = i_{\ell-1}$. Assume that $d = 1$. Without loss of generality, we may assume that $X_0 = D$. By applying Lemma 6.25 to $i_1 \triangleleft^{X_1} \cdots \triangleleft^{X_{\ell-1}} i_\ell = i_0$, we have $i_0 = i_1 = \cdots = i_{\ell-1}$.

Assume that $d > 1$. Without loss of generality, we may assume that $X_0 = D$. There exists a function $\delta: \{0, 1, \dots, d-1\} \rightarrow \{0, \dots, \ell-1\}$ such that $X_{\delta(k)} = D$, δ is strictly increasing and $\delta(0) = 0$. Then we have a sequence

$$i_0 \triangleleft^D i_1 \triangleleft^{X_1} \cdots \triangleleft^{X_{\delta(k)-1}} i_{\delta(k)} \triangleleft^D i_{\delta(k)+1} \triangleleft^{X_{\delta(k)+1}} \cdots \triangleleft^{X_{\ell-1}} i_\ell = i_0,$$

and this sequence induces the following relations in (P, \leq) .

$$i_0 \geq i_1 \leq i_{\delta(1)} \geq i_{\delta(1)+1} \leq \cdots \geq i_{\delta(d-1)+1} \leq i_\ell = i_0.$$

Since there is no k with $X_k = X_{k+1} = D$, and $d > 1$, we have $\delta(d-1) + 1 \geq 3$. By applying Lemma 6.26, without loss of generality, there is at least one of the following relation:

$$i_0 \leq i_{\delta(1)}, \quad i_{\delta(1)} \leq i_0, \quad i_1 \leq i_{\delta(1)+1}, \quad i_{\delta(1)+1} \leq i_1.$$

If $i_0 \leq i_{\delta(1)}$ holds, then $i_0 \in [i_1, i_{\delta(1)}]$ holds since (P, \leq) is a diamond-free poset. Then there exists an integer $m \in \{1, \dots, \delta(1) - 1\}$ such that $i_0 \in [i_m, i_{m+1}]$. Since $i_m \triangleleft^{X_m} i_{m+1}$ and $X_m \in \{I_1, I_2\}$, $i_0 \triangleleft^{X_m} i_{m+1}$ holds. By removing vertices i_1, i_2, \dots, i_m from (6.4), we have a sequence

$$i_0 \triangleleft^{X_m} i_{m+1} \triangleleft^{X_{m+1}} \dots \triangleleft^{X_{\ell-1}} i_{\ell} = i_0.$$

In this sequence, the number of k such that $X_k = D$ is $d - 1$. Thus by inductive hypothesis, we have $i_0 = i_{m+1} = \dots = i_{\ell-1}$. Applying Lemma 6.25 to the sequence $i_1 \triangleleft^{X_1} i_2 \triangleleft^{X_2} \dots \triangleleft^{X_m} i_{m+1} = i_0$, we have the claim.

If we have the relation $i_{\delta(1)} \leq i_0$, then the elements $i_2, i_3, \dots, i_{\delta(1)}$ appear in $[i_1, i_0]$. In particular, $i_{\delta(1)} \triangleleft^D i_1$ holds. By applying Lemma 6.25 to a sequence $i_1 \triangleleft^{X_1} i_2 \triangleleft^{X_2} \dots \triangleleft^{X_{\delta(1)-1}} i_{\delta(1)}$, we have $i_1 = i_2 = \dots = i_{\delta(1)}$. By identifying $i_1 = i_2 = \dots = i_{\delta(1)}$ in the sequence (6.4), since \triangleleft^D is transitive and by an inductive hypothesis, we have the claim.

If we have the relation $i_1 \leq i_{\delta(1)+1}$, then there exists an integer $m \in \{1, \dots, \delta(1) - 1\}$ such that $i_{\delta(1)+1} \in [i_m, i_{m+1}]$. We have $i_{\delta(1)+1} \triangleleft^{X_m} i_{m+1}$. By applying Lemma 6.25 to a sequence

$$i_{\delta(1)+1} \triangleleft^{X_m} i_{m+1} \triangleleft^{X_{m+1}} \dots \triangleleft^{X_{\delta(1)-1}} i_{\delta(1)},$$

we have $i_{m+1} = \dots = i_{\delta(1)} = i_{\delta(1)+1}$. By an inductive hypothesis, we have the claim.

If we have the relation $i_{\delta(1)+1} \leq i_1$, then $i_1 \in [i_{\delta(1)+1}, i_{\delta(1)}]$. In particular, $i_1 \triangleleft^D i_{\delta(1)+1}$ holds. Then by an inductive hypothesis, we have the claim.

We proved the claim and it induces that \triangleleft' is antisymmetric. \square

We show that the partial order \triangleleft' gives a meet of $[\triangleleft_1]$ and $[\triangleleft_2]$.

Lemma 6.28. *Let (P, \leq) be a diamond-free poset and $i, j \in P$. Assume that \triangleleft' is a partial order on P . Let Δ' be the set of standard $A(P)$ -modules associated to \triangleleft' . Then $S(j)$ is a composition factor of $\Delta'(i)$ if and only if $S(j)$ is a composition factor of both $\Delta_1(i)$ and $\Delta_2(i)$.*

Proof. The if part is easy. If $j = i$, then the assertion is clear. So let $j \neq i$. Assume that $S(j)$ is a composition factor of $\Delta'(i)$. Then $i \leq j$ and any $k \in [i, j]$ satisfies $k \triangleleft' i$. Let m be the maximal element $k \in [i, j]$ such that $i_{k'} \triangleleft^D i$ holds for any $i \leq k' \leq k$. If $k = j$, then $S(j)$ is a composition factor of both $\Delta_1(i)$ and $\Delta_2(i)$.

Suppose that $k \neq j$. By Lemma 6.24, the minimal element l in $[k, j]$ satisfies either $i \triangleleft^{I_1} l$ or $i \triangleleft^{I_2} l$. This implies $i = l$, since \triangleleft' is antisymmetric. This is a contradiction, since i never equals to l . Therefore we have $k = j$ and the assertion holds. \square

Lemma 6.29. *Assume that (P, \leq) is a diamond-free poset and that \triangleleft' is a partial order on P . Let Δ' be the set of standard $A(P)$ -modules associated to \triangleleft' . Then $\mathcal{F}(\Delta_1) \cup \mathcal{F}(\Delta_2) \subset \mathcal{F}(\Delta')$ holds.*

Proof. We show that $\Delta_1(i)$ belongs to $\mathcal{F}(\Delta')$ for any $i \in P$. It is enough to show that for a submodule K of $\Delta_1(i)$, if $\Delta_1(i)/K$ belongs to $\mathcal{F}(\Delta')$, then there exists a surjective morphism from K to a product of some $\Delta'(j)$'s. Assume that L is an indecomposable direct summand of K . Since $A(P)$ is hereditary, L has a simple top $S(j)$ for some $j \in P$. Let $k \in P$. We show that $S(k)$ is a composition factor of L if $S(k)$ is a composition factor of $\Delta'(j)$. Since $A(P)$ is hereditary, we have that $S(k)$ is a composition factor of L if and only if $S(k)$ is a composition factor of $\Delta_1(i)$ and $j \leq k$ holds.

Since L is a submodule of $\Delta_1(i)$, $i \leq j$ holds and any $u \in [i, j]$ satisfies $u \triangleleft_1 i$. Assume that $S(k)$ is a composition factor of $\Delta'(j)$. Then $j \leq k$ holds and any $v \in [j, k]$ satisfies $v \triangleleft' j$. By Lemma 6.28, such v satisfies $v \triangleleft_1 j$. Thus for the interval $[i, k]$, we have that any $u \in [i, k]$ satisfies $u \triangleleft_1 i$. Thus $S(k)$ is a composition factor of $\Delta_1(i)$, and therefore $S(k)$ is a composition factor of L . \square

Proposition 6.30. *Let (P, \leq) be a diamond-free poset. For given two quasi-hereditary structures $[\triangleleft_1]$ and $[\triangleleft_2]$ on $A(P)$, assume that \triangleleft' is a partial order on P . Let Δ' be the set of standard $A(P)$ -modules associated to \triangleleft' . Then we have the following statements.*

- (1) *The partial order \triangleleft' induces a quasi-hereditary structure on $A(P)$.*
- (2) *The meet of $[\triangleleft_1]$ and $[\triangleleft_2]$ in $\text{qh.str}(A(P))$ exists and is represented by \triangleleft' .*

Proof. (1) We show that $(A(P), (P, \triangleleft'))$ satisfies (1), (2) and (3) of Definition 2.4. By the definition and Lemma 6.29, (1) and (2) hold. Clearly $(P(i) : \Delta'(i)) = 1$ holds for any $i \in P$. Assume that

$(P(i) : \Delta'(j)) \neq 0$ for $j \leq i \in P$. We show $i \triangleleft' j$. Since $P(i)$ belongs to $\mathcal{F}(\Delta')$, there exist the following sequence of submodules of $P(i)$:

$$M_{\ell+1} \subset M_\ell \subset \cdots \subset M_0 = P(i)$$

such that $M_k/M_{k+1} \cong \Delta'(i_k)$ for $k = 0, \dots, \ell$, $i_0 = i$ and $i_\ell = j$. Let i'_k be the maximal element in $[i_{k-1}, i_k] \setminus \{i_k\}$ for $k = 1, \dots, \ell$. We have a disjoint union of intervals

$$[i, j] = [i_0, i'_1] \sqcup [i_1, i'_2] \sqcup \cdots \sqcup [i_{\ell-1}, i'_\ell] \sqcup \{i_\ell\}.$$

For each $k = 0, \dots, \ell - 1$ and any $m \in [i_k, i'_{k+1}]$, $S(m)$ is a composition factor of $\Delta'(i_k)$. Therefore by Lemma 6.28, we can apply Lemma 6.24 to each interval $[i_{k-1}, i_k] = [i_{k-1}, i'_k] \sqcup \{i_k\}$. We have that $i_{k-1} \triangleleft^{I*} i_k$ for each $k = 1, \dots, \ell$ and some $*$ $\in \{1, 2\}$. Namely, we have $i \triangleleft' j$.

(2) Let \triangleleft_3 be an adapted order on P with $[\triangleleft_3] \preceq [\triangleleft_1]$ and $[\triangleleft_3] \preceq [\triangleleft_2]$. Let Δ_3 be the standard modules associated to \triangleleft_3 . By Lemma 2.20, any composition factor of $\Delta_3(i)$ is a composition factor of both $\Delta_1(i)$ and $\Delta_2(i)$. By Lemma 6.28, there exists a surjection from $\Delta'(i)$ to $\Delta_3(i)$. Therefore, we have $[\triangleleft_3] \preceq [\triangleleft']$. \square

Then we complete the proof of Theorem 6.22.

Proof of Theorem 6.22. Let (P, \leq) be a diamond-free poset such that Z_n is not a full subposet for any $n \geq 4$. For two quasi-hereditary structures $[\triangleleft_1]$ and $[\triangleleft_2]$, consider the following two binary relations on P :

$$\begin{aligned} \triangleleft' &= \left((\text{Dec}(\triangleleft_1) \cap \text{Dec}(\triangleleft_2)) \cup \text{Inc}(\triangleleft_1) \cup \text{Inc}(\triangleleft_2) \right)^{\text{tc}}, \\ \triangleleft'' &= \left(\text{Dec}(\triangleleft_1) \cup \text{Dec}(\triangleleft_2) \cup (\text{Inc}(\triangleleft_1) \cap \text{Inc}(\triangleleft_2)) \right)^{\text{tc}}. \end{aligned}$$

By Propositions 6.27 and 6.30, \triangleleft' defines a quasi-hereditary structure on $A(P)$ and $[\triangleleft']$ is a meet of $[\triangleleft_1]$ and $[\triangleleft_2]$.

Let B be the incidence algebra of the opposite poset (P, \leq^{op}) of (P, \leq) . It is easy to see that the opposite algebra of $A(P)$ is isomorphic to B . Let Dec_B and Inc_B be a decreasing and a increasing over B , respectively. By Lemma 2.22 (1), for a partial order \triangleleft on P and $i, j \in P$, we have

- $(i, j) \in \text{Dec}(\triangleleft)$ if and only if $(j, i) \in \text{Inc}_B(\triangleleft)$.
- $(i, j) \in \text{Inc}(\triangleleft)$ if and only if $(j, i) \in \text{Dec}_B(\triangleleft)$.

Therefore, we have the following equality of binary relations on P :

$$\triangleleft'' = \left((\text{Dec}_B(\triangleleft_1) \cap \text{Dec}_B(\triangleleft_2)) \cup \text{Inc}_B(\triangleleft_1) \cup \text{Inc}_B(\triangleleft_2) \right)^{\text{tc}}.$$

By applying Propositions 6.27 to B , \triangleleft'' is a partial order on P . By Lemma 2.22, we have an anti-isomorphism of posets between $\text{qh.str}(A(P))$ and $\text{qh.str}(B)$ which is induced from an identity map of the set of partial orders on P . Thus \triangleleft_m defines a quasi-hereditary structure $[\triangleleft_m]_B$ on B for $m = 1, 2$. By Proposition 6.30, \triangleleft'' represents a meet of $[\triangleleft_1]_B$ and $[\triangleleft_2]_B$ in $\text{qh.str}(B)$. This implies that \triangleleft'' represents a join of $[\triangleleft_1]$ and $[\triangleleft_2]$ in $\text{qh.str}(A(P))$. \square

ACKNOWLEDGMENTS

The first author is indebted to his advisor, Henning Krause, for his guidance and for suggesting the study of quasi-hereditary structures as PhD research project, resulting the classification of quasi-hereditary structures on path algebras of type \mathbb{A} as part of his doctoral thesis.

REFERENCES

- [BB05] A. Björner and F. Brenti. “Combinatorics of Coxeter groups”. Vol. 231. New York, NY: Springer, 2005, pp. xiv + 363.
- [CPP19] G. Châtel, V. Pilaud, and V. Pons. “The weak order on integer posets”. In: *Algebr. Comb.* 2.1 (2019), pp. 1–48.
- [CPS88] E. T. Cline, B. J. Parshall, and L. L. Scott. “Finite dimensional algebras and highest weight categories.” In: *J. Reine Angew. Math.* 391 (1988), pp. 85–99.
- [Con16] T. Conde. “On certain strongly quasihereditary algebras”. PhD thesis. University of Oxford, 2016.

- [Cou19] K. Coulembier. “The classification of blocks in BGG category \mathcal{O} ”. In: *arXiv e-prints* (Mar. 2019). arXiv: [1903.02717](https://arxiv.org/abs/1903.02717).
- [DR89] V. Dlab and C. M. Ringel. “Quasi-hereditary algebras”. In: *Illinois J. Math.* 33.2 (1989), pp. 280–291.
- [DR92] V. Dlab and C. M. Ringel. “The module theoretical approach to quasi-hereditary algebras”. In: *Representations of algebras and related topics (Kyoto, 1990)*. Vol. 168. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 1992, pp. 200–224.
- [FR20] M. Flores and B. Rognerud. *Quasi-hereditary structures, 2020, Research code (Version 1.0)*. 2020. URL: https://webusers.imj-prg.fr/~baptiste.rognerud/en/quasi_hereditary_structures.html.
- [Gab81] P. Gabriel. “Un jeu? Les nombres de Catalan”. In: *Uni Zürich, Mitteilungsblatt des Rektorats* 6 (1981), pp. 4–5.
- [HU05] D. Happel and L. Unger. “On a partial order of tilting modules”. In: *Algebr. Represent. Theory* 8.2 (2005), pp. 147–156.
- [Hil06] L. Hille. “On the volume of a tilting module”. In: *Abh. Math. Sem. Univ. Hamburg* 76 (2006), pp. 261–277.
- [OEI20] OEIS Foundation Inc. (2020). *The On-Line Encyclopedia of Integer Sequences*. URL: <https://oeis.org/>.
- [QPA19] The QPA-team. *QPA - Quivers, path algebras and representations - a GAP package, Version 1.30*. 2019. URL: <https://folk.ntnu.no/oyvinso/QPA/>.
- [RS91] C. Riedtmann and A. Schofield. “On a simplicial complex associated with tilting modules”. In: *Comment. Math. Helv.* 66.1 (1991), pp. 70–78.
- [Rin10] C. M. Ringel. “Iyama’s finiteness theorem via strongly quasi-hereditary algebras”. In: *J. Pure Appl. Algebra* 214.9 (2010), pp. 1687–1692.
- [Rin91] C. M. Ringel. “The category of modules with good filtrations over a quasi-hereditary algebra has almost split sequences”. In: *Math. Z.* 208.2 (1991), pp. 209–223.
- [Sag19] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 8.8)*. 2019. URL: <https://www.sagemath.org>.
- [Sco87] L. L. Scott. “Simulating algebraic geometry with algebra. I. The algebraic theory of derived categories”. In: *The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986)*. Vol. 47. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 1987, pp. 271–281.

MANUEL FLORES: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, 33501 BIELEFELD, GERMANY
E-mail address: mflores@math.uni-bielefeld.de

YUTA KIMURA: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, 33501 BIELEFELD, GERMANY
E-mail address: ykimura@math.uni-bielefeld.de

BAPTISTE ROGNERUD: UNIVERSITÉ PARIS DIDEROT, INSTITUT DE MATHÉMATIQUES DE JUSSIEU - PARIS RIVE GAUCHE,
 75013 PARIS, FRANCE
E-mail address: baptiste.rognerud@imj-prg.fr