

Distribution of peak heights modulo k and double descents on k -Dyck paths

ALEXANDER BURSTEIN
 Department of Mathematics
 Howard University
 Washington, DC 20059, USA
 aburstein@howard.edu

September 2, 2020

Abstract

We show that the distribution of the number of peaks at height i modulo k in k -Dyck paths of a given length is independent of $i \in [0, k-1]$ and is the reversal of the distribution of the total number of peaks. Moreover, these statistics, together with the number of double descents, are jointly equidistributed with any of their permutations. We also generalize this result to generalized Motzkin paths and generalized ballot paths.

Given a positive integer k , a k -Dyck path of down-size n is a lattice path from $(0, 0)$ to $((k+1)n, 0)$ with unit steps $u = (1, 1)$ (an *up-step*) and $d = (1, -k)$ (a *down-step*) that stays in the first quadrant. This is a generalization of the well-known Dyck paths (see e.g. [3]), where $k = 1$ and the down-size is called the *semilength*. Note that a k -Dyck path of down-size n contains exactly n down-steps and kn up-steps.

A ud block in a k -Dyck path is called a *peak*, and the *height* of that peak is defined as the second coordinate of the left endpoint of the d step in the peak. Given a peak ud , we also call the vertex between the u and d steps a peak. Similarly, a dd block (and the vertex in the middle of it) is called a *double descent*.

In Section 1 of this paper we will consider a group of statistics related to the heights of peaks in k -Dyck paths, and prove that they not only have the same distribution, but also have the same joint distribution as any permutation of themselves. Furthermore, these statistics are naturally related to the number-of- i -th-children statistics on $(k+1)$ -ary trees for $i = 1, 2, \dots, k+1$, which are easily seen to have the same joint distribution as any of their permutations.

In Section 2 we will extend and generalize the results of Section 1 to include level steps (e.g. generalized Motzkin and Schröder paths) and some paths that do not end at their initial height (e.g. generalized ballot paths).

Key words and phrases: k -Dyck path, peak height.

2020 *Mathematics Subject Classification:* 05A15 (Primary) 05A19, 05A05 (Secondary).

1 Main results

Let \mathcal{P}_n^k be the set of k -Dyck paths of down-size $n \geq 0$. Note that the rightmost peak of any path $P \in \mathcal{P}_n^k$ is followed only by down-steps and thus is at height $0 \pmod k$. Therefore, we will only consider non-rightmost peaks of paths in \mathcal{P}_n^k .

We define the following statistics on the \mathcal{P}_n^k . Given a k -Dyck path $P \in \mathcal{P}_n^k$, for each $i = 0, 1, \dots, k-1$, let $\text{pk}(P)$ be the total number of non-rightmost peaks in P , and let $\text{pk}_i(P)$ be the number of non-rightmost peaks in P that are at heights $i \pmod k$. (Of course, the rightmost peak of P is always at height $0 \pmod k$, since it is followed only by down-steps, but we will need the values of the above statistics on some subpaths of P as well, where height modulo k of the rightmost peak may be different.) We also define $\text{dd}(P)$ to be the number of double descents in P . Note that every d step of P follows either a u step or a d step, and thus $\text{dd}(P) = n - 1 - \text{pk}(P)$ for $n \geq 1$ (we subtract 1 for the rightmost peak of P). Then we have the following result.

Theorem 1.1. *The $(k+1)$ -statistic $(\text{pk}_0, \text{pk}_1, \dots, \text{pk}_{k-1}, \text{dd})$ on \mathcal{P}_n^k is jointly equidistributed with any of its permutations.*

We obtain this result by giving a recursive bijection between k -Dyck paths and $(k+1)$ -ary trees that maps the above statistics on k -Dyck paths to the number of $(i+1)$ -st children statistics on $(k+1)$ -ary trees for $i = 0, 1, \dots, k-1, k$.

An m -ary tree is a rooted ordered tree in which each node has at most m children assigned positions in $\{1, 2, \dots, m\}$ from left to right, not necessarily consecutively.

Let \mathcal{T}_n^m be the set of m -ary trees on n vertices, $n \geq 1$. Define the following statistics on \mathcal{T}_n^m . Given an m -ary tree $T \in \mathcal{T}_n^m$, let $e_i(T)$ be the number of i th children in T , i.e. nodes that are children in position i from the left, for $1 \leq i \leq m$. Equivalently, $e_i(T)$ is the number of i th edges in T , i.e. edges from a vertex to its child in position i .

Proposition 1.2. *The m -statistic (e_1, e_2, \dots, e_m) on \mathcal{T}_n^m is jointly equidistributed with any of its permutations.*

Proof. This is almost immediately obvious. It is enough to recursively permute the subtrees of each node of T starting from the root according to the permutation of the statistics e_i , $i = 1, \dots, m$. We will leave the details of this bijection to the reader. \square

We can now connect Proposition 1.2 to Theorem 1.1. We will begin by defining two operations on paths.

Definition 1.3. Given a k -Dyck path Q starting at height 0, define its i -lifting $Q^{(i)}$ as the path with the same sequence of steps but starting at height i .

Note that any k -Dyck path Q with the maximal down-step suffix of length $n \geq 1$ has a unique decomposition as

$$Q = Q_0^{(0)} u Q_1^{(1)} u \dots u Q_{kn-1}^{(kn-1)} u \underbrace{\text{dd} \dots d}_n \quad (1.1)$$

where each Q_j , $0 \leq j \leq kn-1$, is a k -Dyck path.

Definition 1.4. Let Q be a k -Dyck path with the maximal down-step suffix of length $n \geq 1$ and with the decomposition as in (1.1). Define a permutation π_n on the set $[0, kn-1]$ by

$$\pi_n = (k-1, \dots, 1, 0) (2k-1, \dots, k+1, k) \dots (kn-1, \dots, k(n-1)+1, k(n-1))$$

in cycle notation. Define the *cyclic shift* $\kappa(Q)$ of Q by replacing each $Q_j^{(j)}$ with $Q_{\pi_n(j)}^{(j)}$, where $j \in [0, kn - 1]$. In other words,

$$Q_j^{(j)} \mapsto \begin{cases} Q_{j+k-1}^{(j)} & \text{if } j \equiv 0 \pmod{k}, \\ Q_{j-1}^{(j)} & \text{otherwise.} \end{cases}$$

Note that κ is a bijection on \mathcal{P}_n^k since π_n is a bijection, and that κ^k is the identity map on \mathcal{P}_n^k for any n .

Lemma 1.5. *For any k -Dyck path Q with the maximal down-step suffix of length $n \geq 1$, the map $\lambda : Q^{(1)} \mapsto \kappa(Q)$ preserves peak heights modulo k in each segment Q_j , $0 \leq j \leq kn - 1$, as well as the number of double descents in Q .*

We will call $\lambda(Q^{(1)}) = \kappa(Q)$ the *cyclic lowering* of $Q^{(1)}$.

Proof. Decomposing Q as in (1.1) and lifting it by 1 unit, we see that each segment Q_j in $Q^{(1)}$ starts at height $j + 1$. On the other hand, the starting height of the segment Q_j in $\kappa(Q)$ is $j - k + 1$ if $j \equiv k - 1 \pmod{k}$, or $j + 1$ if $j \not\equiv k - 1 \pmod{k}$. In other words, the height of peaks either does not change or decreases by k , and thus remains the same modulo k .

Thus, λ preserves the statistics pk_i ($0 \leq i \leq k - 1$) on Q , and hence their sum pk . But $pk + dd = m - 1$ on \mathcal{P}_m^k for any $m \geq 0$, and therefore λ preserves the number of double descents as well. \square

Applying cyclic lowering several times, we obtain an easy corollary of Lemma 1.5.

Corollary 1.6. *For any k -Dyck path Q with the maximal down-step suffix of length $n \geq 1$, the map $\lambda^i : Q^{(i)} \mapsto \kappa^i(Q)$ on k -Dyck paths preserves peak heights modulo k in each block Q_j , $0 \leq j \leq kn - 1$.*

We will call $\lambda^i(Q^{(i)}) = \kappa^i(Q)$ the *cyclic i -lowering* of $Q^{(i)}$. Note that $\kappa^i(Q)$ permutes blocks Q_j as follows:

$$Q_j^{(j)} \mapsto \begin{cases} Q_{j+k-i}^{(j)} & \text{if } j \equiv 0, 1, \dots, i-1 \pmod{k}, \\ Q_{j-i}^{(j)} & \text{if } j \equiv i, i+1, \dots, k-1 \pmod{k}. \end{cases}$$

[As an aside, we note that this is not a unique way to define λ so as to preserve non-rightmost peak heights modulo k . Another possible way, among many, is to cyclically shift the whole sequence of subpaths Q_j , i.e. to define the permutation π_n as the cycle $(kn - 1, kn - 2, \dots, 1, 0)$. In that case, κ would require less block moves but would have order kn , which depends on n .]

Theorem 1.7. *The statistics $(pk_0, pk_1, \dots, pk_{k-2}, pk_{k-1}, dd)$ on \mathcal{P}_n^k and $(e_1, e_2, \dots, e_{k+1})$ on \mathcal{T}_n^{k+1} are equidistributed.*

Proof. To prove this theorem, we give a bijection $\psi : \mathcal{P}^k \rightarrow \mathcal{T}^{k+1}$ that maps peaks at heights i modulo k ($i = 0, 1, \dots, k - 1$) of a k -Dyck path P of down-size n to $(i + 1)$ -st children of a $(k + 1)$ -ary tree $\psi(P)$ on n vertices, and maps double descents of P onto $(k + 1)$ -st (i.e. rightmost) children in $\psi(P)$.

To do this, first label the peaks and double descents of P (i.e. the left endpoints of all down steps of P) from left to right as follows:

- label the rightmost peak by r ,

- label the left endpoint of a down edge i_j if it is the j th non-rightmost peak at height i modulo k , where $0 \leq i \leq k-1$, and
- label the left endpoint of a down edge d_j if it is the j th double descent of P .

For a tree $T \in \mathcal{T}^{k+1}$, let T_i its i -th subtree and write $T = (T_1, T_2, \dots, T_{k+1})$. Then define the map ψ as follows: let $\psi(\emptyset) = \emptyset$ (i.e. the one-point path corresponds to the empty tree), and for $P \neq \emptyset$, define $\psi(P)$ recursively so that, for $i = 0, 1, \dots, k$, the $(i+1)$ -th subtree of $\psi(P)$ is $\psi(P)_{i+1} = \psi(\kappa^i(P_i))$. In other words,

$$\psi(P) = (\psi(P_0), \psi(\kappa(P_1)), \psi(\kappa^2(P_2)), \dots, \psi(\kappa^{k-1}(P_{k-1}), \psi(\kappa^k(P_k))), \quad (1.2)$$

and we recall that $\kappa^k(P_k) = P_k$. Now label the vertices of $\psi(P)$ recursively as follows:

- label the root of $\psi(P)$ with r ,
- for $i = 0, 1, \dots, k-1$, label the $(i+1)$ -th child of the root of $\psi(P)$ (i.e. the root of $\psi(\kappa^i(P_i))$) with the label of the rightmost peak of $P_i^{(i)}$ (which is at height i modulo k).
- label the $(k+1)$ -st (i.e. the rightmost) child of the root of $\psi(P)$ (i.e. the root of $\psi(P_k)$) with the label of the right endpoint of $P_k^{(k)}$ (which is a double descent if $P_k \neq \emptyset$, since it is preceded and followed by a down-step). Recall that the rightmost peak of $P_k^{(k)}$ is also the rightmost peak of P , and hence its label has already been used for the root of $\psi(P)$.

Now, proceeding inductively for each $i = 0, 1, \dots, k$, label the remaining (nonroot) vertices of each $\psi(\kappa^i(P_i))$ with the labels of the corresponding non-rightmost peaks or double descents of $P_i^{(i)}$. Note that the cyclic i -lowering λ^i may permute positions of peaks and double descents of $P_i^{(i)}$ but preserves peak heights modulo k . Therefore, all $(i+1)$ -st children in $\psi(P)$ are labeled either i_j (for some j) if $0 \leq i \leq k-1$, or d_j (for some j) if $i = k$ (see Figure 1.1 and Example 1.8).

In particular, if P has down-size n , then $\psi(P)$ has n vertices. Moreover,

$$(pk_0, pk_1, \dots, pk_{k-1}, dd)(P) = (e_1, e_2, \dots, e_{k+1})(\psi(P)). \quad \square$$

Finally, we see that Proposition 1.2 and Theorem 1.7 together imply Theorem 1.1.

Example 1.8. Figure 1.1 gives an example of the application of map ψ of Theorem 1.7, together with the corresponding mapping of the peaks and double descents of a path P to the nodes of the corresponding tree $\psi(P)$. We label the peaks of P as described in Theorem 1.7. In our example, $k = 2$, $n = 10$, and

$$P = \underbrace{uuduuuuuudd}_{P_0} \underbrace{u u u d u u u u u d d d}_{P_1^{(1)}} \underbrace{u u u d}_{P_2^{(2)}} d.$$

Then $\kappa^2 = \text{id}$ and $\psi(P) = (\psi(P_0), \psi(\kappa(P_1)), \psi(\kappa^2(P_2))) = (\psi(P_0), \psi(\kappa(P_1)), \psi(P_2))$. For $P_1 = u u d u u u u u d d$, we use the other decomposition,

$$P_1 = \underbrace{u u d}_{P_{1,0}} \underbrace{u}_{P_{1,1}^{(1)}} \underbrace{\emptyset}_{P_{1,2}^{(2)}} \underbrace{u}_{P_{1,3}^{(3)}} \underbrace{u u d}_{P_{1,3}^{(3)}} d d.$$




Figure 1.1: An example of the application of map ψ , together with the corresponding mapping of the peaks and double descents of a 2-Dyck path P to the nodes of the ternary tree $\psi(P)$.

Since the $ud\dots d$ suffix of P_1 contains 2 d 's, we have that κ permutes the blocks $P_{1,j}$, $j = 0, 1, 2, 3$, according to $\pi_2 = (10)(32)$, i.e.

$$\kappa(P_1) = \underbrace{\emptyset}_{P_{1,1}^{(0)}} \underbrace{u \text{ u u d}}_{P_{1,0}^{(1)}} \underbrace{u \text{ u u d}}_{P_{1,3}^{(2)}} \underbrace{u \text{ u d}}_{P_{1,2}^{(3)}} = \text{uuu duuu duudd}.$$

Note also that $P_0 = P_1$ in this example, but the subtrees $\psi(P)_1 = \psi(P_0)$ and $\psi(P)_2 = \psi(\kappa(P_1))$ of $\psi(P)$ corresponding to P_0 and P_1 , respectively, are not equal as ordered trees.

As a direct consequence of Theorem 1.7, we have the following corollary.

Corollary 1.9. *The statistics $pk_0, pk_1, \dots, pk_{k-1}, dd$ are equidistributed on \mathcal{P}_n^k for $n \geq 0$.*

Moreover, the fact that $dd(P) = n - 1 - pk(P)$ for $P \in \mathcal{P}_n^k$, $n \geq 1$, implies that the distribution of each of $pk_0, pk_1, \dots, pk_{k-1}, dd$ on \mathcal{P}_n^k is the reversed distribution of pk on \mathcal{P}_n^k .

Example 1.10. Letting $k = 1$, we obtain that (dd, pk) and (pk, dd) are equidistributed on Dyck paths. In fact, the well-known bijection $\eta(\emptyset) = \emptyset$, $\eta(P_0 u P_1 d) \mapsto \eta(P_1) u \eta(P_0) d$ on Dyck paths (more precisely, an involution) interchanges the dd and pk statistics. Since the distribution of pk is given by the Narayana numbers $N(n, r) = \frac{1}{n} \binom{n}{r} \binom{n}{r-1}$, $1 \leq r \leq n$ (see A001263 [2]), this yields (yet another) bijective proof that $N(n, r) = N(n, n - r)$.




Figure 1.2: Peak heights modulo 2 and double descents in 2-Dyck paths of down-size 3. A non-rightmost peak at height $i \bmod 2$ ($i \in \{0, 1\}$) is labeled with i . A double descent is labeled with d . The rightmost peak is unlabeled. Below each path P is the value of $(pk_0, pk_1, dd)(P)$.

Example 1.11. Letting $k = 2$, we see that dd , pk_1 (odd-height non-rightmost peaks), and pk_0 (even-height non-rightmost peaks) are equidistributed on 2-Dyck paths, also known as *ternary paths* (see Figure 1.2). The distribution of each of pk_0 , pk_1 , and dd is given by A120986 [2], and the distribution of $pk = pk_0 + pk_1 = n - 1 - dd$ is given by A108767 [2].

The preceding results enable us now to calculate the joint distribution on \mathcal{P}_n^k ($n \geq 1$) of the statistic $(pk_0, pk_1, \dots, pk_{k-2}, pk_{k-1}, dd)$.

Corollary 1.12. *The number of paths $P \in \mathcal{P}_n^k$ ($n \geq 1$) such that $(pk_0, pk_1, \dots, pk_{k-1}, dd)(P) = (r_0, r_1, \dots, r_{k-1}, r_k)$, where $\sum_{i=0}^k r_i = n - 1$, is*

$$\frac{1}{n} \prod_{i=0}^k \binom{n}{r_i}.$$

Moreover, for each statistic $st \in \{pk_0, pk_1, \dots, pk_{k-1}, dd\}$, the number of paths $P \in \mathcal{P}_n^k$ ($n \geq 1$) such that $st(P) = r$ is

$$\frac{1}{n} \binom{n}{r} \binom{kn}{n-1-r},$$

whereas the number of paths $P \in \mathcal{P}_n^k$ ($n \geq 1$) such that $\text{pk}(P) = r + 1$ is

$$\frac{1}{n} \binom{n}{n-1-r} \binom{kn}{r} = \frac{1}{n} \binom{n}{r+1} \binom{kn}{r},$$

where $0 \leq r \leq n - 1$.

Note that $\sum_{i=0}^k r_i = n - 1$ in Corollary 1.12 since a down-step must be preceded by an up-step or another down-step, so that $\sum_{i=0}^k r_i$ counts all the down-steps of P except the one immediately after its rightmost peak.

Proof. Let $f = f(x; q_0, q_1, \dots, q_{k-1}, q_k)$ be the generating function over the nonempty k -Dyck paths defined by

$$\begin{aligned} f &= f(x; q_0, q_1, \dots, q_{k-1}, q_k) = \sum_{n=1}^{\infty} \sum_{P \in \mathcal{P}_n^k} x^n \left(\prod_{i=0}^{k-1} q_i^{\text{pk}_i(P)} \right) q_k^{\text{dd}(P)} \\ &= \sum_{n=1}^{\infty} \sum_{T \in \mathcal{T}_n^{k+1}} x^n \left(\prod_{i=0}^k q_i^{e_{i+1}(T)} \right). \end{aligned} \quad (1.3)$$

Then Proposition 1.7 and its proof imply that

$$f = x \prod_{i=0}^k (q_i f + 1). \quad (1.4)$$

The number of paths $P \in \mathcal{P}_n^k$ such that $(\text{pk}_0, \text{pk}_1, \dots, \text{pk}_{k-1}, \text{dd})(P) = (r_0, r_1, \dots, r_{k-1}, r_k)$ is the coefficient of f at $x^n \prod_{i=0}^k q_i^{r_i}$. Applying Lagrange inversion to (1.4), we get that

$$\left[x^n \prod_{i=0}^k q_i^{r_i} \right] f = \frac{1}{n} \left[f^{n-1} \prod_{i=0}^k q_i^{r_i} \right] \left(\prod_{i=0}^k (q_i f + 1) \right)^n = \frac{1}{n} \prod_{i=0}^k \binom{n}{r_i},$$

which is the number of paths we want to enumerate.

The second and third statement of the corollary can be derived similarly by setting all but one $q_i = 1$ and noting that $\text{pk} = n - 1 - \text{dd}$ on \mathcal{P}_n^k . \square

2 Extensions and generalizations

2.1 Paths with level steps

Theorem 1.1 can be extended to generalized colored Motzkin and Schröder paths as follows. Let $A \subseteq \mathbb{N}$, and let $c = (c_a)_{a \in \mathbb{N}}$ be a vector of nonnegative integers such that $c_a = 0$ exactly when $a \notin A$. Consider a path P from $(0, 0)$ back to height 0 with unit steps $u = (1, 1)$, $d = (1, -k)$, and $l_{a,b} = (a, 0)$ of color b , where $1 \leq b \leq c_a$, for all $a \in A$. We also include the case $A = \{\infty\}$ and $c_\infty = 1$ to be able to exclude all level steps so as to generate the k -Dyck paths considered above. We will call such paths (k, A, c) -paths, as this generalizes (k, a) -paths defined in [1] for $A = \{a\}$ and $c_a = 1$, and let $\mathcal{P}_n^{k,A,c}$ be the set of (k, A, c) -paths of length n . We also define the length of up,

down, and level steps by $|u| = 1$, $|d| = 1$, and $|l_{a,b}| = a$, and the length $|P|$ of a path P as the sum of the lengths of its steps.

On such a (k, A, c) -path P , call the blocks dd and $l_{a,b}d$ *weak double descents*, and call blocks ud and $ul_{a,b}$, as well as the single step $l_{a,b}$ if it is the leftmost step of P , *weak peaks* (in each case here, $a \in A$ and $b \in [c_a]$). Extending our peak height definition for k -Dyck paths, let the height of a weak peak be the second coordinate of the left endpoint of the down or level step in the peak. Furthermore, on (k, A, c) -paths P , define the following statistics:

- $\text{wpk}_i(P)$, the number of non-rightmost weak peaks of P at heights $i \bmod k$, where $i = 0, 1, \dots, k-1$,
- $\text{wpk}(P) = \sum_{i=0}^{k-1} \text{wpk}_i(P)$, the total number of non-rightmost peaks of P ,
- $\text{wdd}(P)$, the number of weak double descents of P .

Then we have the following unique decomposition:

$$P = L \quad \text{or} \quad P = P_0^{(0)} u P_1^{(1)} u \dots u P_{k-1}^{(k-1)} u P_k^{(k)} d L,$$

where each P_i is a (k, A, c) -path of down-size n_i (such that $1 + \sum_{i=0}^k n_i = n$, the down-size of P) and L is a block of level steps of allowed lengths and colors. Furthermore, as before, for each path P_i we have the unique decomposition

$$P_i = P_{i,0}^{(0)} u P_{i,1}^{(1)} u \dots u P_{i,k m_i - 1}^{(k m_i - 1)} u D_i,$$

where each $P_{i,j}$ is a (k, A, c) -path and D_i is block of down-steps and level steps (i.e. no up-steps) containing exactly m_i down-steps and possibly some level steps of any allowed lengths and colors. Then it is easy to see that, for each $i \in [0, k-1]$, the cyclic i -lowering map λ_i that permutes the subpaths $P_{i,j}$ in the same way as in Corollary 1.6, preserves weak peak heights modulo k as well as the number of weak peaks and weak double descents in $P_i^{(i)}$. The preceding discussion implies the following result.

Theorem 2.1. *The $(k+1)$ -statistic $(\text{wpk}_0, \text{wpk}_1, \dots, \text{wpk}_{k-1}, \text{wdd})$ on $\mathcal{P}_n^{k,A,c}$ is jointly equidistributed with any of its permutations.*

It is a pretty straightforward exercise to adapt the bijective proof of Theorem 1.1 to prove Theorem 2.1, and we will leave it to the reader. Instead, we will give a short generating function argument similar to that given in Corollary 1.12 so as not to repeat essentially the same proof with minor variations.

Proof. Note that letting $A = \infty$ yields Theorem 1.1. Let $A \neq \infty$ and let $c_A(x) = \sum_{a \in A} c_a x^a$, $\mathcal{P}^{k,A,c} = \cup_{n=0}^{\infty} \mathcal{P}_n^{k,A,c}$, and

$$f_{k,A,c} = f_{k,A,c}(x; q_0, q_1, \dots, q_{k-1}, q_k) = \sum_{P \in \mathcal{P}^{k,A,c} \setminus \{\emptyset\}} x^{|P|} \left(\prod_{i=0}^{k-1} q_i^{\text{wpk}_i(P)} \right) q_k^{\text{wdd}(P)}. \quad (2.1)$$

Then, from the path decompositions above and the cyclic i -lowering maps, we obtain the functional equation

$$f_{k,A,c} = c_A(x)(f_{k,A,c} + 1) + x^{k+1} \prod_{i=0}^k (q_i f_{k,A,c} + 1), \quad (2.2)$$

Now notice that Equation (2.2) is symmetric in the variables q_0, q_1, \dots, q_k . □




Figure 2.1: Weak peaks and weak double descents in Motzkin paths of length 5. A non-rightmost weak peak is labeled with 0 (the only remainder modulo $k = 1$). A weak double descent is labeled with d . The rightmost weak peak is unlabeled. Below each path P is the value of $(\text{wpk}, \text{wdd})(P)$.

Example 2.2. Letting $k = 1$, we obtain the joint equidistribution of (wdd, wpk) and (wpk, wdd) on Motzkin paths by setting $A = \{1\}$, $c_1 = 1$ (see Figure 2.1 for an example), and on Schröder paths by setting $A = \{2\}$, $c_2 = 1$.

2.2 Ballot paths

We can also generalize Theorem 1.1 to left prefixes of k -Dyck paths starting at $(0,0)$ and ending at some height $m \geq 0$. Define a (k, m) -ballot path of down-size n to be a path from $(0,0)$ to $((k+1)n + m, m)$ with steps $u = (1,1)$ and $d = (1,-k)$. Let $\mathcal{P}^{k,m}$ be the set of (k, m) -ballot paths, and let $\mathcal{P}_n^{k,m}$ be the set of (k, m) -ballot paths of down-size n .

Let $m \equiv r \pmod{k}$, so that $m = \ell k + r$ for some integers $\ell \geq 0$ and $0 \leq r \leq k-1$. Observe that a (k, m) -ballot path is a prefix of some k -Dyck path, and moreover, for $m \not\equiv 0 \pmod{k}$, it can only be extended on the right to a k -Dyck path by using at least some up-steps. Thus, for $m \not\equiv 0 \pmod{k}$, any suffix extending a (k, m) -ballot path to a k -Dyck path must contain a peak. If $m \equiv 0 \pmod{k}$, then the rightmost peak is necessarily at height $0 \pmod{k}$, so peaks at heights $i \not\equiv 0 \pmod{k}$ are not rightmost. Hence, we need to modify our peak statistics and let $\text{pk}_i^*(P)$ be the number of peaks in (k, m) -ballot path P at heights $i \pmod{k}$ (so these statistics may count the rightmost peak as well).

Theorem 2.3. Let $m, \ell \geq 0$ and $0 \leq r \leq k-1$ be integers such that $m = \ell k + r$. Then for any $0 \leq r \leq k-1$,

- the $(r+1)$ -statistic $(\text{pk}_0^*, \dots, \text{pk}_r^*)$ on $\mathcal{P}_n^{k,m}$ is jointly equidistributed with any of its permutations.
- the $(k-r-1)$ -statistic $(\text{pk}_{r+1}^*, \dots, \text{pk}_{k-1}^*)$ on $\mathcal{P}_n^{k,m}$ is jointly equidistributed with any of its permutations.

Proof. Note that any (k, m) -ballot path P can be uniquely decomposed as

$$P = P_0^{(0)} u P_1^{(1)} u \dots u P_{m-1}^{(m-1)} u P_m^{(m)}, \quad (2.3)$$

where each P_i , $i = 0, 1, \dots, m$, is a k -Dyck path, on which the multistatistics $(\text{pk}_0, \dots, \text{pk}_r)$ and $(\text{pk}_{r+1}, \dots, \text{pk}_{k-1})$ are jointly equidistributed with any of their respective permutations. The only

peaks not counted by the unstarred statistics are the rightmost peaks. Note that the rightmost peak of each subpath $P_i^{(i)}$ is at height $i \bmod k$. Thus, $\ell + 1$ of these subpaths, if nonempty, have their rightmost peak at heights in $[0, r]$ modulo k , whereas only ℓ of these subpaths, if nonempty, have their rightmost peak at heights in $[r + 1, k - 1]$ modulo k . Thus, for $i = 0, 1, \dots, k - 1$,

$$\begin{aligned} pk_i^*(P) &= \sum_{j=0}^m pk_i^*(P_j^{(j)}) = \begin{cases} \sum_{j=0}^m pk_i(P_j^{(j)}) + \sum_{j=0}^{\ell} [P_{kj+i} \neq \emptyset], & \text{if } 0 \leq i \leq r, \\ \sum_{j=0}^m pk_i(P_j^{(j)}) + \sum_{j=0}^{\ell-1} [P_{kj+i} \neq \emptyset], & \text{if } r + 1 \leq i \leq k - 1, \end{cases} \\ &= \begin{cases} \sum_{j=0}^m pk_i(\kappa^j(P_j)) + \sum_{j=0}^{\ell} [P_{kj+i} \neq \emptyset], & \text{if } 0 \leq i \leq r, \\ \sum_{j=0}^m pk_i(\kappa^j(P_j)) + \sum_{j=0}^{\ell-1} [P_{kj+i} \neq \emptyset], & \text{if } r + 1 \leq i \leq k - 1, \end{cases} \end{aligned} \quad (2.4)$$

where κ is the cyclic shift map defined on page 2 and, as before, $[\cdot]$ is the Iverson bracket.

The distribution of these statistics can be computed as well. Let

$$g_{k,m} = g_{k,m}(x; q_0, q_1, \dots, q_k) = \sum_{n=0}^{\infty} \sum_{P \in \mathcal{P}_n^{k,m}} x^n \left(\prod_{i=0}^{k-1} q_i^{pk_i^*(P)} \right) q_k^{dd(P)}, \quad (2.5)$$

then, from (2.4),

$$g_{k,m} = g_{k,\ell k+r} = \left(\prod_{i=0}^r (q_i f + 1) \right)^{\ell+1} \left(\prod_{i=r+1}^{k-1} (q_i f + 1) \right)^{\ell}, \quad (2.6)$$

where the function f is defined in (1.3) and satisfies (1.4). The multipliers q_i in (2.6) correspond to the fact that the rightmost peak of a nonempty $P_{kj+i}^{(kj+i)}$ is at height i modulo k . From (1.4), f is symmetric in q_0, \dots, q_k , and therefore it is clear from (2.6) that $g_{k,\ell k+r}$ is symmetric in q_0, \dots, q_r as well as in q_{r+1}, \dots, q_{k-1} . \square

Applying Lagrange inversion to (2.6) as in Corollary 1.12 yields after some routine manipulations that the number of paths $P \in \mathcal{P}_n^{k,\ell k+r}$ with $(pk_0^*, \dots, pk_{k-1}^*, dd)(P) = (s_0, \dots, s_{k-1}, s_k)$, where $\sum_{i=0}^k s_i = n \geq 1$, is

$$\begin{aligned} &\left[x^n \prod_{i=0}^r q_i^{s_i} \right] g_{k,\ell k+r} \\ &= \frac{1}{n} \left(\frac{\ell+1}{n+\ell+1} \sum_{i=0}^r s_i + \frac{\ell}{n+\ell} \sum_{i=r+1}^{k-1} s_i \right) \left(\prod_{i=0}^r \binom{n+\ell+1}{s_i} \right) \left(\prod_{i=r+1}^{k-1} \binom{n+\ell}{s_i} \right) \binom{n}{s_k}. \end{aligned} \quad (2.7)$$

Note that letting $\ell = 0$ and $r = 0$ in (2.7) yields the result of Corollary 1.12, since (r_0, r_1, \dots, r_k) in Corollary 1.12 is $(s_0 - 1, s_1, \dots, s_k)$ in the notation of (2.7).

Example 2.4. Letting $k = 2$, $\ell = 0$, $r = 1$, we see that statistics (pk_0^*, pk_1^*) and (pk_1^*, pk_0^*) are equidistributed on $(2, 1)$ -ballot paths of down-size n . In particular, statistics pk_0^* (even-height peaks) and pk_1^* (odd-height peaks) are equidistributed on those paths. For an example of this, see the last 7 paths in Figure 1.2 (rightmost three in row 2 and all of row 3), delete the rightmost ud from each of those, and ignore the dd statistic.

The above result can be generalized for the weak versions of the corresponding statistics.

Let A and c be defined as for (k, A, c) -paths. Define a (k, A, c, m) -path of down-size n to be a path from height 0 to height $m \geq 0$ with steps $u = (1, 1)$, $d = (1, -k)$, and any level steps $l_{a,b} = (a, 0)$ of color b , for $a \in A$ and $b \in [c_a]$. Note that a (k, A, c, m) -path of down-size n (i.e. with n down-steps) has $kn + m$ up-steps. Let $\mathcal{P}_n^{k,A,c,r}$ be the set of such paths, and let $\mathcal{P}^{k,A,c,m} = \bigcup_{n=0}^{\infty} \mathcal{P}_n^{k,A,c,m}$. Note that $\mathcal{P}_n^{k,A,c} = \mathcal{P}_n^{k,A,c,0}$, and therefore, $\mathcal{P}^{k,A,c} = \mathcal{P}^{k,A,c,0}$. Let $\text{wpk}_i^*(P)$ ($i = 0, 1, \dots, k-1$) be the number of (possibly rightmost) weak peaks of P at heights $i \bmod k$. Then we have the following result.

Theorem 2.5. *Let $m, \ell \geq 0$ and $0 \leq r \leq k-1$ be integers such that $m = \ell k + r$. Then for any $0 \leq r \leq k-1$,*

- *the $(r+1)$ -statistic $(\text{wpk}_0^*, \dots, \text{wpk}_r^*)$ on $\mathcal{P}_n^{k,A,c,m}$ is jointly equidistributed with any of its permutations.*
- *the $(k-r-1)$ -statistic $(\text{wpk}_{r+1}^*, \dots, \text{wpk}_{k-1}^*)$ on $\mathcal{P}_n^{k,A,c,m}$ is jointly equidistributed with any of its permutations.*

Proof. Note that any (k, A, c, m) -path P can be uniquely decomposed as

$$P = P_0^{(0)} u P_1^{(1)} u \dots P_{m-1}^{(m-1)} u P_m^{(m)}, \quad (2.8)$$

where each P_i , $i = 0, 1, \dots, m$, is a (k, A, c) -path. Define

$$g_{k,A,c,m} = g_{k,A,c,m}(x; q_0, q_1, \dots, q_k) = \sum_{P \in \mathcal{P}^{k,A,c,r}} x^{|P|} \left(\prod_{i=0}^{k-1} q_i^{\text{wpk}_i^*(P)} \right) q_k^{\text{wdd}(P)}, \quad (2.9)$$

then the decomposition (2.3) implies, as in the proof of Theorem 2.3, that

$$g_{k,A,c,m} = g_{k,A,c,\ell k+r} = x^m \left(\prod_{i=0}^r (q_i f_{k,A,c} + 1) \right)^{\ell+1} \left(\prod_{i=r+1}^{k-1} (q_i f_{k,A,c} + 1) \right)^{\ell},$$

where $f_{k,A,c}$ is defined in (2.1) and satisfies (2.2), which is symmetric in q_0, \dots, q_k . Therefore, $g_{k,A,c,\ell k+r}$ is symmetric in q_0, \dots, q_r and in q_{r+1}, \dots, q_{k-1} . \square

ACKNOWLEDGEMENT

The author would like to thank Lou Shapiro and Sergey Kirgizov for their valuable advice on improving the presentation of the results.

References

- [1] T. Mansour, M. Shattuck, Counting humps and peaks in generalized Motzkin paths, *Discrete Appl. Math.* **161** (2013), 2213–2216.
- [2] N.J.A. Sloane, ed., The On-Line Encyclopedia of Integer Sequences, published electronically at <https://oeis.org>, 2020.
- [3] R.P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, 1999.