

RESTRICTED 132-DUMONT PERMUTATIONS

Toufik Mansour ¹

Department of Mathematics, Chalmers University of Technology, S-41296 Göteborg, Sweden

toufik@math.chalmers.se

ABSTRACT

In [D] Dumont showed that certain classes of permutations on n letters are counted by the Genocchi numbers. In particular, Dumont showed that the $(n+1)$ st Genocchi number is the number of permutations on $2n$ letters with the following properties: (1) each even integer must be followed by a smaller integer (this rule disallows the sequence from ending with an even integer), (2) each odd integer is either followed by a larger integer or is final in the sequence. We call such permutations by *Dumont permutations of the first kind*. In this paper we study the number of Dumont permutations of the first kind on n letters avoiding the pattern 132 and avoiding (or containing exactly once) an arbitrary pattern on k letters. In several interesting cases the generating function depends only on k .

Keywords: Dumont permutations, restricted permutations, generating functions.

1. INTRODUCTION

Classical patterns. Let $\alpha \in \mathfrak{S}_n$ and $\tau \in \mathfrak{S}_k$ be two permutations. We say that α *contains* τ if there exists a subsequence $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that $(\alpha_{i_1}, \dots, \alpha_{i_k})$ is order-isomorphic to τ ; in such a context τ is usually called a *pattern*. We say that α *avoids* τ , or is τ -*avoiding*, if such a subsequence does not exist. The set of all τ -avoiding permutations in \mathfrak{S}_n is denoted $\mathfrak{S}_n(\tau)$. For an arbitrary finite collection of patterns T , we say that α avoids T if α avoids any $\tau \in T$; the corresponding subset of \mathfrak{S}_n is denoted $\mathfrak{S}_n(T)$.

While the case of permutations avoiding a single pattern has attracted much attention, the case of multiple pattern avoidance remains less investigated. In particular, it is natural, as the next step, to consider permutations avoiding pairs of patterns τ_1, τ_2 . This problem was solved completely for $\tau_1, \tau_2 \in \mathfrak{S}_3$ (see [SS]), for $\tau_1 \in \mathfrak{S}_3$ and $\tau_2 \in \mathfrak{S}_4$ (see [W]), and for $\tau_1, \tau_2 \in \mathfrak{S}_4$ (see [Bo1, Km] and references therein). Several recent papers [CW, MV1, Kr, MV2, MV3, MV4] deal with the case $\tau_1 \in \mathfrak{S}_3, \tau_2 \in \mathfrak{S}_k$ for various pairs τ_1, τ_2 . Another natural question is to study permutations avoiding τ_1 and containing τ_2 exactly t times. Such a problem for certain $\tau_1, \tau_2 \in \mathfrak{S}_3$ and $t = 1$ was investigated in [R], and for certain $\tau_1 \in \mathfrak{S}_3, \tau_2 \in \mathfrak{S}_k$ in [RWZ, MV1, Kr]. The tools involved in these papers include generating trees, continued fractions, Chebyshev polynomials, and Dyck words. Also, the tools involved in these papers include many sequences, for example sequence of Catalan numbers, Fibonacci numbers, and Pell numbers.

¹Research financed by EC's IHRP Programme, within the Research Training Network "Algebraic Combinatorics in Europe", grant HPRN-CT-2001-00272

We denote the n th *Catalan number* by $C_n = \frac{1}{n+1} \binom{2n}{n}$. The generating function for the Catalan numbers is denoted by $C(x)$, that is, $C(x) = \sum_{n \geq 0} C_n x^n = \frac{1-\sqrt{1-4x}}{2x}$.

Generalized patterns. In [BS] introduced generalized permutation patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation. We write a classical pattern with dashes between any two adjacent letters of the pattern, say 1342, as 1-3-4-2, and if we write, say 24-3-1, then we mean that if this pattern occurs in permutation $\pi \in \mathfrak{S}_n$, then the letters in the permutation π that correspond to 2 and 4 are adjacent (see [C]). For example, the permutation $\pi = 35421$ has only two occurrences of the pattern 23-1, namely the subsequences 352 and 351, whereas π has four occurrences of the pattern 2-3-1, namely the subsequences 352, 351, 342, and 341.

Claesson [C] presented a complete solution for the number of permutations avoiding any single generalized pattern of length three with exactly one adjacent pair of letters. Claesson and Mansour [CM] presented a complete solution for the number of permutations avoiding any double generalized patterns of length three with exactly one adjacent pair of letters. Kitaev [Ki] investigate simultaneous avoidance of two or more 3-letters generalized patterns without internal dashes. Later, Mansour [M1, M2] (for more details see [M3]) presented a general approach to study the number of permutations avoiding 1-3-2 and avoiding (or containing exactly once) an arbitrary generalized pattern.

Dumont permutations. Dumont [D] showed that certain classes of permutations in \mathfrak{S}_n are counted by the Genocchi numbers (see [SP, Sequence A001469(M3041)]). Dumont showed that the $(n+1)$ st Genocchi number is the number of permutations in \mathfrak{S}_{2n} with the following properties: (1) each even integer must be followed by a smaller integer (this rule disallows the sequence from ending with an even integer), (2) each odd integer is either followed by a larger integer or is final in the sequence. We call such permutations by *Dumont permutations of the first kind*. For example, 2143, 3421, and 4213 are the all Dumont permutations of the first kind of length 4.

Dumont [D] defined another type of permutations in \mathfrak{S}_n and showed that the $(n+1)$ st Genocchi number is the number of permutations in \mathfrak{S}_{2n} with the following properties: (1) $\pi_i < i$ for any even position i , (2) $\pi_i \geq i$ for any odd position i . We call such permutations by *Dumont permutations of the second kind*. For example, 2143, 3142, 4132 are the all Dumont permutations of the second kind of length 4.

Remark 1.1. Let $\pi \in \mathfrak{S}_n$ be any Dumont permutation of the second kind; since $\pi_2 < 2$ we get $\pi_2 = 1$. Hence, it is easy to see that there are no Dumont permutations of the second kind in $\mathfrak{S}_n(132)$ for all $n \geq 4$. So, in this paper we discuss only the case of Dumont permutations of the first kind.

We define for all $r \geq 2$,

$$(1.1) \quad Q_r(x) = 1 + \frac{x^2 Q_{r-1}(x)}{1 - x^2 Q_{r-2}(x)}.$$

We denote the solution of Recurrence 1.1 with $Q_0(x) = 0$ and $Q_1(x) = 1$ by $F_r(x)$, and we denote the solution of Recurrence 1.1 with $Q_0(x) = Q_1(x) = 1$ by $G_r(x)$. For example, $F_2(x) = 1 + x^2$, $F_3(x) = \frac{1+x^4}{1-x^2}$, $G_2(x) = \frac{1}{1-x^2}$, and $G_3(x) = \frac{1-x^2+x^4}{(1-x^2)^2}$. Evidently, $F_r(x)$ and $G_r(x)$ are rational functions in x^2 , and for all $r \geq 1$,

$$(1.2) \quad F_r(x) = 1 + \sum_{j=1}^{r-1} \frac{x^{2j}}{\prod_{m=r-1-j}^{r-2} (1 - x^2 F_m(x))} \text{ and } G_r(x) = 1 + \sum_{j=1}^{r-1} \frac{x^{2j}}{\prod_{m=r-1-j}^{r-2} (1 - x^2 G_m(x))}.$$

Example 1.2. Using Recurrence 1.1 it is easy to see that

$$F_4(\sqrt{x}) = \sum_{n \geq 0} (f_{n+2} + f_n - 2)x^n \text{ and } G_4(\sqrt{x}) = 1 + x + \sum_{n \geq 2} (3 \cdot 2^{n-2} - 1)x^n,$$

where f_n is the n th Fibonacci number.

Organization of the paper. In this paper we use generating function techniques to study those Dumont permutations of the first kind which avoid 132 and avoid (or contain exactly once) an arbitrary pattern on k letters. In several interesting cases the generating function depends only on k .

The paper is organized as follows. The case of Dumont permutations of the first kind avoiding both 132 and τ is treated in Section 2. We present a simple structure for any Dumont permutation of the first kind avoiding 132. This structure can be obtained explicitly for several interesting cases, including classical patterns and generalized patterns. This allows us to find explicitly some statistics on Dumont permutations of the first kind which avoid 132. The case of avoiding 132 and containing another pattern τ exactly once is treated in Section 3. Again, we find explicitly the generating function for several interesting cases of τ , including classical patterns and generalized patterns.

Most of the explicit solutions obtained in Sections 2-4 involve the generating functions $F_k(x)$ and $G_k(x)$.

2. DUMONT PERMUTATIONS OF THE FIRST KIND WHICH AVOID 132 AND ANOTHER PATTERN

Let $\mathfrak{D}_\tau^{(1)}(n)$ denote the number of Dumont permutations of the first kind in $\mathfrak{S}_n(132, \tau)$, and let $\mathfrak{D}_\tau^{(1)}(x) = \sum_{n \geq 0} \mathfrak{D}_\tau^{(1)}(n)x^n$ be the corresponding generating function. In this section we describe a method for enumerating Dumont permutations of the first kind which avoid 132 and another pattern and we use our method to enumerate $\mathfrak{D}_\tau^{(1)}(n)$ for various τ . We begin with an observation concerning the structure of the Dumont permutations of the first kind avoiding 132 which holds immediately from definitions.

Proposition 2.1. For any $\pi \in \mathfrak{D}_n^{(1)}(132)$ such that $\pi_j = n$, there holds one of the following assertions:

1. if n is odd number then $\pi = (\pi', n)$, where $\pi' \in \mathfrak{D}_{n-1}^{(1)}(132)$;
2. if n is even number then $\pi = (\pi', n, \pi'')$ such that π' is a Dumont permutation of the first kind on the numbers $n-j+1, n-j+2, \dots, n-1$, π'' is nonempty Dumont permutation of the first kind on the numbers $1, 2, \dots, n-j$, and $j = 1, 2, 4, \dots, n-2$.

2.1. $\tau = \emptyset$. As a corollary of Proposition 2.1 we find an explicit formula for the number of 132-avoiding Dumont permutations of the first kind in \mathfrak{S}_n .

Theorem 2.2. The generating function for the number of 132-avoiding Dumont permutations of the first kind in \mathfrak{S}_n is given by $(1+x)C(x^2)$. In other words, the number of 132-avoiding Dumont permutations of the first kind in \mathfrak{S}_n is given by $C_{[n/2]}$, which is the $[n/2]$ th Catalan number.

Proof. By Proposition 2.1, we have two possibilities for block decomposition of an arbitrary $\pi \in \mathfrak{D}_n^{(1)}(132)$. Let us write an equation for $\mathfrak{D}_\emptyset^{(1)}(x)$. The contribution of the first decomposition above equals

$$\sum_{n \geq 0} \mathfrak{D}_\emptyset^{(1)}(2n+1)x^{2n+1} = x \sum_{n \geq 0} \mathfrak{D}_\emptyset^{(1)}(2n)x^{2n},$$

equivalently,

$$(2.1) \quad \mathfrak{D}_{\emptyset}^{(1)}(x) - \mathfrak{D}_{\emptyset}^{(1)}(-x) = x(\mathfrak{D}_{\emptyset}^{(1)}(x) + \mathfrak{D}_{\emptyset}^{(1)}(-x)).$$

The contribution of the second decomposition above equals

$$\sum_{n \geq 1} \mathfrak{D}_{\emptyset}^{(1)}(2n)x^{2n} = \sum_{n \geq 1} \mathfrak{D}_{\emptyset}^{(1)}(2n-1)x^{2n} + \sum_{n \geq 1} \sum_{j=0}^n \mathfrak{D}_{\emptyset}^{(1)}(2j+1)\mathfrak{D}_{\emptyset}^{(1)}(2n+2-2j)x^{2n},$$

equivalently,

$$(2.2) \quad \begin{aligned} & \mathfrak{D}_{\emptyset}^{(1)}(x) + \mathfrak{D}_{\emptyset}^{(1)}(-x) - 2 = \\ & = x(\mathfrak{D}_{\emptyset}^{(1)}(x) - \mathfrak{D}_{\emptyset}^{(1)}(-x)) + \frac{x}{2}(\mathfrak{D}_{\emptyset}^{(1)}(x) - \mathfrak{D}_{\emptyset}^{(1)}(-x))(\mathfrak{D}_{\emptyset}^{(1)}(x) + \mathfrak{D}_{\emptyset}^{(1)}(-x) - 2). \end{aligned}$$

By putting $\mathfrak{D}_{\emptyset}^{(1)}(x) = (1+x)A(x)$ in Equations 2.1 and 2.2 it is easy to see that $A(x) = C(x^2)$. \square

2.2. A classical pattern $\tau = 12\dots k$. Let us start by the following example.

Example 2.3. By definitions we have $\mathfrak{D}_1^{(1)}(x) = 1$ and $\mathfrak{D}_{12}^{(1)}(x) = 1 + x + x^2$.

The case of varying k is more interesting. As an extension of Example 2.3, let us consider the case $\tau = 12\dots k$.

Theorem 2.4. Let $A_k(x) = \frac{1}{2}(\mathfrak{D}_{12\dots k}^{(1)}(x) + \mathfrak{D}_{12\dots k}^{(1)}(-x))$ and $B_k(x) = \frac{1}{2}(\mathfrak{D}_{12\dots k}^{(1)}(x) - \mathfrak{D}_{12\dots k}^{(1)}(-x))$ for all $k \geq 0$. Then

$$A_k(x) = F_k(x), \quad B_k(x) = xF_{k-1}(x), \quad \text{and} \quad \mathfrak{D}_{12\dots k}^{(1)}(x) = F_k(x) + xF_{k-1}(x).$$

Proof. Using the same arguments as in the proof of Theorem 2.2 we get

$$\mathfrak{D}_{12\dots k}^{(1)}(x) - \mathfrak{D}_{12\dots k}^{(1)}(-x) = x(\mathfrak{D}_{12\dots (k-1)}^{(1)}(x) + \mathfrak{D}_{12\dots (k-1)}^{(1)}(-x)),$$

and

$$\begin{aligned} & \mathfrak{D}_{12\dots k}^{(1)}(x) + \mathfrak{D}_{12\dots k}^{(1)}(-x) - 2 = x(\mathfrak{D}_{12\dots k}^{(1)}(x) - \mathfrak{D}_{12\dots k}^{(1)}(-x)) + \\ & + \frac{x}{2}(\mathfrak{D}_{12\dots (k-1)}^{(1)}(x) - \mathfrak{D}_{12\dots (k-1)}^{(1)}(-x))(\mathfrak{D}_{12\dots k}^{(1)}(x) + \mathfrak{D}_{12\dots k}^{(1)}(-x) - 2). \end{aligned}$$

The rest is easy to check by the definitions of A_k and B_k . \square

Example 2.5. Theorem 2.4, for $k = 3$, yields $\mathfrak{D}_{123}^{(1)}(x) = \frac{1+x+x^4-x^5}{1-x^2}$. In other words, the number of 132-avoiding Dumont permutation of the first kind in $\mathfrak{S}_n(123)$ is given by $1 + (-1)^n$ for all $n \geq 4$, and 1 for $n = 0, 1, 2, 3$. An another example, Theorem 2.4, for $k = 4$, yields $\mathfrak{D}_{1234}^{(1)}(x) = \frac{1+2x+x^2+2x^6+x^7+x^8}{(1+x)(1-x^2-x^4)}$. In other words, the number of 132-avoiding Dumont permutation of the first kind in $\mathfrak{S}_n(1234)$ is $f_{n/2+2} + f_{n/2} - 2$ if n is even number, otherwise 2 for all $n \geq 2$, where f_n is the n th Fibonacci number.

As an extension of Theorem 2.4, let us define

$$\mathfrak{A}(x_1, x_2, x_3, \dots) = \sum_{\pi \in \mathfrak{D}^{(1)}} \prod_{j \geq 1} x_1^{12\dots j(\pi)},$$

where $\mathfrak{D}^{(1)}$ is the set of all Dumont permutations of the first kind of all sizes including the empty permutation, and $\tau(\pi)$ is the number of occurrences of τ in π . Let

$$\begin{aligned} A^{(1)}(x_1, x_2, x_3, \dots) &= \frac{1}{2}(\mathfrak{A}(x_1, x_2, x_3, \dots) + \mathfrak{A}(-x_1, x_2, x_3, \dots)), \\ B^{(1)}(x_1, x_2, x_3, \dots) &= \frac{1}{2}(\mathfrak{A}(x_1, x_2, x_3, \dots) - \mathfrak{A}(-x_1, x_2, x_3, \dots)). \end{aligned}$$

Using the same arguments as in the proof of Theorem 2.4, we obtain the following.

Theorem 2.6. *We have*

$$A^{(1)}(x_1, x_2, x_3, \dots) = 1 + \frac{x_1^2 A^{(1)}(x_1 x_2, x_2 x_3, x_3 x_4, \dots)}{1 - x_1^2 x_2 A^{(1)}(x_1 x_2^2 x_3, x_2 x_3^2 x_4, x_3 x_4^2 x_5, \dots)},$$

and

$$B^{(1)}(x_1, x_2, x_3, \dots) = x_1 A^{(1)}(x_1 x_2, x_2 x_3, x_3 x_4, \dots).$$

As an application to Theorem 2.6, for $x_1 = x$ and $x_j = 1$, $j \geq 2$, we get that

$$B^{(1)}(x, 1, 1, \dots) = x A^{(1)}(x, 1, 1, \dots),$$

and

$$\begin{aligned} A^{(1)}(x, 1, 1, \dots) &= \frac{1}{1 - \frac{x^2}{1 - \frac{x^2}{1 - \frac{x^2}{\ddots}}}} = C(x^2). \end{aligned}$$

Hence, we have $\mathfrak{D}_{\mathcal{O}}^{(1)}(x) = (1+x)C(x^2)$ (see Theorem 2.2).

Another application for Theorem 2.6 is the number of right to left maxima. Let $\pi \in \mathfrak{S}_n$, π_i is a *right to left maxima* if $\pi_i > \pi_j$ for all $i < j$. We denote the number of right to left maxima of π by $rlm(\pi)$. In [BCS, Proposition 5] proved

$$lrm(\pi) = \sum_{j \geq 1} 12 \dots j(\pi)(-1)^{j-1}.$$

Therefore,

$$\sum_{\pi \in \mathfrak{D}^{(1)}} x^{|\pi|} y^{rlm(\pi)} = \mathfrak{A}(xy, y^{-1}, y, y^{-1}, \dots)$$

together with Theorem 2.6 and $A^{(1)}(x, 1, 1, \dots) = C(x^2)$ we get

$$\sum_{\pi \in \mathfrak{D}^{(1)}} x^{|\pi|} y^{rlm(\pi)} = 1 + xC(x^2)y + \sum_{n \geq 2} x^{2n-2} C^{n-1}(x^2)y^n.$$

Corollary 2.7. *The generating function for the number of Dumont permutations of the first kind avoiding 132 and having exactly k right to left maxima is given by $x^{2k-2} C^{k-1}(x^2)$ for all $k \geq 2$, and $x^k C^k(x^2)$ for $k = 0, 1$.*

2.3. A classical pattern $\tau = 2134 \dots k$. Similarly as in Theorem 2.4, we obtain the case $\tau = 2134 \dots k$.

Theorem 2.8. *For all $k \geq 2$,*

$$\mathfrak{D}_{213 \dots k}^{(1)}(x) = G_{k-1}(x) + xG_{k-2}.$$

Example 2.9. *Theorem 2.8 for $k = 3, 4$ yields $\mathfrak{D}_{213}^{(1)}(x) = \frac{1+x-x^3}{1-x}$ and $\mathfrak{D}_{2134}^{(1)}(x) = \frac{1+x-x^2-x^3+x^4}{(1-x^2)^2}$.*

2.4. A generalized pattern 12-3- \cdots - k . In this subsection we use the notation of generalized patterns (see Section 1). For example, we write the classical pattern 132 as 1-3-2.

By definitions, we get $\mathfrak{D}_{12}^{(1)}(x) = 1 + x + x^2$. So, by the same arguments as in the proof of Theorem 2.4, together with

$$\mathfrak{D}_{12}^{(1)}(x) = \mathfrak{D}_{1-2}^{(1)}(x) = 1 + x + x^2,$$

we obtain the following.

Theorem 2.10. *For all $k \geq 1$,*

$$\mathfrak{D}_{12-3-\cdots-k}^{(1)}(x) = \mathfrak{D}_{1-2-3-\cdots-k}^{(1)}(x) = F_k(x) + xF_{k-1}(x).$$

A comparison of Theorem 2.4 with Theorem 2.10 suggests that there should exist a bijection between the sets $\mathfrak{S}_n(1\text{-}3\text{-}2, 12\text{-}3\text{-}\cdots\text{-}k)$ and $\mathfrak{S}_n(1\text{-}3\text{-}2, 1\text{-}2\text{-}3\text{-}\cdots\text{-}k)$. However, we failed to produce such a bijection, and finding it remains a challenging open question.

Now, let us define

$$\mathfrak{B}(x_1, x_2, x_3, \dots) = \sum_{\pi \in \mathfrak{D}^{(1)}} x_1^{1(\pi)} \prod_{j \geq 2} x_1^{12-3-\cdots-j(\pi)},$$

where $\mathfrak{D}^{(1)}$ is the set of all Dumont permutations of the first kind of all sizes including the empty permutation, and $\tau(\pi)$ is the number of occurrences of τ in π . Let

$$\begin{aligned} A^{(2)}(x_1, x_2, x_3, \dots) &= \frac{1}{2}(\mathfrak{B}(x_1, x_2, x_3, \dots) + \mathfrak{B}(-x_1, x_2, x_3, \dots)), \\ B^{(2)}(x_1, x_2, x_3, \dots) &= \frac{1}{2}(\mathfrak{B}(x_1, x_2, x_3, \dots) - \mathfrak{B}(-x_1, x_2, x_3, \dots)). \end{aligned}$$

Using the same arguments as those in the proof of Theorem 2.4, we get

Theorem 2.11.

$$A^{(2)}(x_1, x_2, x_3, \dots) = 1 + \frac{x_1^2(1 - x_2 + x_2 A^{(2)}(x_1, x_2 x_3, x_3 x_4, \dots))}{1 - x_1^2 x_2(1 - x_2 x_3 + x_2 x_3 A^{(2)}(x_1, x_2 x_3^2 x_4, x_3 x_4^2 x_5, \dots))},$$

and

$$B^{(2)}(x_1, x_2, x_3, \dots) = x_1 - x_1 x_2 + x_1 x_2 A^{(2)}(x_1, x_2 x_3, x_3 x_4 \dots).$$

Let $\pi \in \mathfrak{S}_n$; we say π_j is a *rise* for π if $\pi_j < \pi_{j+1}$ for all $j = 1, 2, \dots, n-1$. We denote the number of rises of π by $rises(\pi)$. By definitions, we have

$$\sum_{\pi \in \mathfrak{D}^{(1)}} x^{|\pi|} y^{rises(\pi)} = x - xy + (1 + xy) A^{(2)}(x, y, 1, 1, \dots),$$

so an application for Theorem 2.11 we get

Corollary 2.12. *The generating function $\sum_{\pi \in \mathfrak{D}^{(1)}} x^{|\pi|} y^{rises(\pi)}$ is given by*

$$\frac{1 + xy - 2x^2y + 2x^2y^2 - (1 + xy)\sqrt{1 - 4x^2y}}{2x^2y^2}.$$

In other words, the generating function for Dumont permutations of the first kind avoiding 1-3-2 with exactly k rises is given by $C_k x^{2k+1} + C_{k+1} x^{2k+2}$ for all $k \geq 1$, and $1 + x + x^2$ for $k = 0$, where C_m is the m th Catalan number.

2.5. A generalized pattern $\tau = 21\text{-}3\text{-}\cdots\text{-}k$. In this subsection, we use the notation of generalized patterns (see Section 1). For example, we write the classical pattern 132 as 1-3-2.

By definitions, we get $\mathfrak{D}_{21}^{(1)}(x) = 1 + x$. So, by the same arguments as in the proof of Theorem 2.4 together with

$$\mathfrak{D}_{21}^{(1)}(x) = \mathfrak{D}_{2-1}^{(1)}(x) = 1 + x,$$

we obtain the following.

Theorem 2.13. *For all $k \geq 2$,*

$$\mathfrak{D}_{21\text{-}3\text{-}\cdots\text{-}k}^{(1)}(x) = \mathfrak{D}_{2-1\text{-}3\text{-}\cdots\text{-}k}^{(1)}(x) = G_{k-1}(x) + xG_{k-2}(x).$$

A comparison of Theorem 2.8 with Theorem 2.13 suggests that there should exist a bijection between the sets $\mathfrak{S}_n(1\text{-}3\text{-}2, 21\text{-}3\text{-}\cdots\text{-}k)$ and $\mathfrak{S}_n(1\text{-}3\text{-}2, 2\text{-}1\text{-}3\text{-}\cdots\text{-}k)$. However, we failed to produce such a bijection, and finding it remains a challenging open question.

Now, let us define

$$\mathfrak{C}(x_1, x_2, x_3, \dots) = \sum_{\pi \in \mathfrak{D}^{(1)}} x_1^{1(\pi)} \prod_{j \geq 2} x_1^{21\text{-}3\text{-}\cdots\text{-}j(\pi)},$$

where $\mathfrak{D}^{(1)}$ is the set of all Dumont permutations of the first kind of all sizes including the empty permutation, and $\tau(\pi)$ is the number of occurrences of τ in π . Let

$$\begin{aligned} A^{(3)}(x_1, x_2, x_3, \dots) &= \frac{1}{2}(\mathfrak{C}(x_1, x_2, x_3, \dots) + \mathfrak{C}(-x_1, x_2, x_3, \dots)), \\ B^{(3)}(x_1, x_2, x_3, \dots) &= \frac{1}{2}(\mathfrak{C}(x_1, x_2, x_3, \dots) - \mathfrak{C}(-x_1, x_2, x_3, \dots)). \end{aligned}$$

Using the same arguments as in the proof of Theorem 2.4, we get the following.

Theorem 2.14. *We have*

$$A^{(3)}(x_1, x_2, x_3, \dots) = 1 + \frac{x_1^2 x_2 A^{(3)}(x_1, x_2 x_3, x_3 x_4, \dots)}{1 - x_1^2 x_2 A^{(3)}(x_1, x_2 x_3^2 x_4, x_3 x_4^2 x_5, \dots)},$$

and

$$B^{(3)}(x_1, x_2, x_3, \dots) = x_1 A^{(3)}(x_1, x_2 x_3, x_3 x_4 \dots).$$

Let $\pi \in \mathfrak{S}_n$; we say that π_j is a *descent* for π if $\pi_j > \pi_{j+1}$ for all $j = 1, 2, \dots, n-1$. We denote the number of descents of π by $\text{descents}(\pi)$. By definitions, we have

$$\sum_{\pi \in \mathfrak{D}^{(1)}} x^{|\pi|} y^{\text{descents}(\pi)} = (1+x) A^{(3)}(x, y, 1, 1, \dots),$$

therefore an application for Theorem 2.14 we get

Corollary 2.15. *The generating function $\sum_{\pi \in \mathfrak{D}^{(1)}} x^{|\pi|} y^{\text{descents}(\pi)}$ is given by $(1+x)C(x^2y)$. In other words, the generating function for Dumont permutations of the first kind avoiding 1-3-2 with exactly k descents is given by $C_k x^{2k+1} + C_k x^{2k+2}$ for all $k \geq 0$, where C_m is the m th Catalan number.*

2.6. A classical pattern $\tau = 23\dots k1$. Again, Proposition 2.1 gives a complete answer for $\tau = 23\dots k1$.

Theorem 2.16. *For all $k \geq 3$,*

$$\mathfrak{D}_{23\dots k1}^{(1)}(x) = 1 + x + \frac{x^2(1+x)}{1-x^2-x^2F_{k-3}(x)}.$$

Proof. Using the same arguments as in the proof of Theorem 2.2 we get

$$\mathfrak{D}_{23\dots k1}^{(1)}(x) - \mathfrak{D}_{23\dots k1}^{(1)}(-x) = x(\mathfrak{D}_{23\dots k1}^{(1)}(x) + \mathfrak{D}_{23\dots k1}^{(1)}(-x)),$$

and

$$\begin{aligned} \mathfrak{D}_{23\dots k1}^{(1)}(x) + \mathfrak{D}_{23\dots k1}^{(1)}(-x) - 2 &= x(\mathfrak{D}_{23\dots k1}^{(1)}(x) - \mathfrak{D}_{23\dots k1}^{(1)}(-x)) + \\ &+ \frac{x}{2}(\mathfrak{D}_{12\dots(k-2)}^{(1)}(x) - \mathfrak{D}_{12\dots(k-2)}^{(1)}(-x))(\mathfrak{D}_{23\dots k1}^{(1)}(x) + \mathfrak{D}_{23\dots k1}^{(1)}(-x) - 2). \end{aligned}$$

The rest is easy to check by the definitions of $F_k(x)$ together with Theorem 2.4. \square

Example 2.17. *Theorem 2.16, for $k = 5$, yields $\mathfrak{D}_{23451}^{(1)}(x) = \frac{(1+x)(1-x^2-x^4)}{1-2x^2-x^4}$. In other words, the number of Dumont permutation of the first kind in $\mathfrak{S}_n(132, 23451)$ is given by $P_{[n/2]}$, which is the $[n/2]$ th Pell number for all $n \geq 2$.*

3. DUMONT PERMUTATIONS OF THE FIRST KIND WHICH AVOID 132 AND CONTAIN ANOTHER PATTERN EXACTLY ONCE

Let $\mathfrak{D}_{\tau;r}^{(1)}(n)$ denote the number of Dumont permutations of the first kind in $\mathfrak{S}_n(132)$ containing τ exactly r times, and let $\mathfrak{D}_{\tau;r}^{(1)}(x) = \sum_{n \geq 0} \mathfrak{D}_{\tau;r}^{(1)}(n)x^n$ be the corresponding generating function.

3.1. A classical pattern $\tau = 12\dots k$.

Theorem 3.1. *Let*

$$A_k(x) = \frac{x^2}{1-x^2F_{k-2}(x)}A_{k-1}(x) + \frac{x^4F_{k-1}(x)}{(1-x^2F_{k-2}(x))^2}A_{k-2}(x)$$

for all $k \geq 2$, where $A_1(x) = 0$ and $A_2(x) = x^4$. Then for all $k \geq 2$

$$\mathfrak{D}_{12\dots k;1}^{(1)}(x) = A_k(x) + xA_{k-1}(x).$$

Proof. By Proposition 2.1, we have two possibilities for block decomposition of an arbitrary π in $\mathfrak{D}_n^{(1)}(132)$. Let us write an equation for $\mathfrak{D}_{12\dots k;1}^{(1)}(x)$. The contribution of the first decomposition above is

$$\sum_{n \geq 0} \mathfrak{D}_{12\dots k;1}^{(1)}(2n+1)x^{2n+1} = x \sum_{n \geq 0} \mathfrak{D}_{12\dots(k-1);1}^{(1)}(2n)x^{2n},$$

equivalently

$$(3.1) \quad \mathfrak{D}_{12\dots k;1}^{(1)}(x) - \mathfrak{D}_{12\dots k;1}^{(1)}(-x) = x(\mathfrak{D}_{12\dots(k-1);1}^{(1)}(x) + \mathfrak{D}_{12\dots(k-1);1}^{(1)}(-x)).$$

The contribution of the second decomposition above is

$$\begin{aligned} \sum_{n \geq 1} \mathfrak{D}_{12\dots k;1}^{(1)}(2n)x^{2n} &= \sum_{n \geq 1} \mathfrak{D}_{12\dots k;1}^{(1)}(2n-1)x^{2n} + \\ &\quad + \sum_{n \geq 1} \sum_{j=0}^n \mathfrak{D}_{12\dots (k-1);1}^{(1)}(2j+1)\mathfrak{D}_{12\dots k;0}^{(1)}(2n+2-2j)x^{2n} + \\ &\quad + \sum_{n \geq 1} \sum_{j=0}^n \mathfrak{D}_{12\dots (k-1);0}^{(1)}(2j+1)\mathfrak{D}_{12\dots k;1}^{(1)}(2n+2-2j)x^{2n}, \end{aligned}$$

equivalently

$$\begin{aligned} (3.2) \quad \mathfrak{D}_{12\dots k;1}^{(1)}(x) + \mathfrak{D}_{12\dots k;1}^{(1)}(-x) &= x(\mathfrak{D}_{12\dots k;1}^{(1)}(x) - \mathfrak{D}_{12\dots k;1}^{(1)}(-x)) + \\ &\quad + \frac{x}{2}(\mathfrak{D}_{12\dots (k-1);1}^{(1)}(x) - \mathfrak{D}_{12\dots (k-1);1}^{(1)}(-x))(\mathfrak{D}_{12\dots k;0}^{(1)}(x) + \mathfrak{D}_{12\dots k;0}^{(1)}(-x) - 2) + \\ &\quad + \frac{x}{2}(\mathfrak{D}_{12\dots (k-1);0}^{(1)}(x) - \mathfrak{D}_{12\dots (k-1);0}^{(1)}(-x))(\mathfrak{D}_{12\dots k;1}^{(1)}(x) + \mathfrak{D}_{12\dots k;1}^{(1)}(-x)). \end{aligned}$$

Using Theorem 2.4, Equation 3.1, Equation 3.2, and Definition 1.1, we get the desired result. \square

Example 3.2. *Theorem 3.1 for $k = 3$ we get*

$$\mathfrak{D}_{123;1}^{(1)}(x) = \frac{x^5(1+x-x^2)}{1-x^2},$$

and for $k = 4$ we get

$$\mathfrak{D}_{1234;1}^{(1)}(x) = \frac{x^7(1+x-3x^2+2x^3+3x^4+3x^5-x^6+x^7)}{(1-x^2)(1-x^2-x^4)^2}.$$

As an extension of Theorem 3.1, let us consider the case $r \geq 1$. Theorem 2.6, for given k and r , yields an explicit formula for $\mathfrak{D}_{12\dots k;r}^{(1)}(x)$. For example, for $k = 3$ and $r = 0, 1, 2, 3, 4$, we have the following.

Theorem 3.3. *We have*

- (i) $\mathfrak{D}_{123;0}^{(1)}(x) = \frac{1+x+x^4-x^5}{1-x^2};$
- (ii) $\mathfrak{D}_{123;1}^{(1)}(x) = \frac{x^5(1+x-x^2)}{1-x^2};$
- (iii) $\mathfrak{D}_{123;2}^{(1)}(x) = \frac{x^5(1+x^2)(1+2x-2x^2-x^3+x^4)}{(1-x^2)^2};$
- (iv) $\mathfrak{D}_{123;3}^{(1)}(x) = \frac{x^7(1+x-x^2+x^3-x^4-x^5+x^6)}{(1-x^2)^2};$
- (v) $\mathfrak{D}_{123;4}^{(1)}(x) = \frac{x^9(1+x^2)(-1-3x+3x^2+3x^3-3x^4-x^5+x^6)}{(1-x^2)^2}.$

3.2. **A classical pattern $\tau = 2134\dots k$.** Similarly to Theorem 3.1, we have

Theorem 3.4. *Let*

$$A_k(x) = \frac{x^2}{1-x^2G_{k-2}(x)}A_{k-1}(x) + \frac{x^4G_{k-1}(x)}{(1-x^2G_{k-2}(x))^2}A_{k-2}(x)$$

for all $k \geq 4$, where $A_1(x) = A_2(x) = x^2$ and $A_3(x) = \frac{x^4}{1-x^2}$. Then, for all $k \geq 2$,

$$\mathfrak{D}_{213\dots k;1}^{(1)}(x) = A_k(x) + xA_{k-1}(x).$$

3.3. A generalized patterns $\tau = 12\text{-}3\text{-}\cdots\text{-}k$ and $\tau = 21\text{-}3\text{-}\cdots\text{-}k$. Similarly to Theorem 3.1, we get

Theorem 3.5. *Let*

$$A_k(x) = \frac{x^2}{1-x^2 F_{k-2}(x)} A_{k-1}(x) + \frac{x^4 F_{k-1}(x)}{(1-x^2 F_{k-2}(x))^2} A_{k-2}(x)$$

for all $k \geq 4$, where $A_1(x) = x^2$ and $A_2(x) = 2x^4$. Then, for all $k \geq 2$,

$$\mathfrak{D}_{12\text{-}3\text{-}\cdots\text{-}k;1}^{(1)}(x) = A_k(x) + x A_{k-1}(x).$$

As an extension of Theorem 3.5, let us consider the case $r \geq 1$. Theorem 2.11, for given k and r , yields an explicit formula for $\mathfrak{D}_{12\text{-}3\text{-}\cdots\text{-}k;r}^{(1)}(x)$. For example, for $k = 3$ and $r = 0, 1, 2, 3, 4$, we have the following.

Theorem 3.6. *We have*

- (i) $\mathfrak{D}_{12\text{-}3;0}^{(1)}(x) = \frac{1+x+x^4-x^5}{1-x^2};$
- (ii) $\mathfrak{D}_{12\text{-}3;1}^{(1)}(x) = \frac{x^5(2+3x-4x^2-x^3+2x^4)}{(1-x^2)^2};$
- (iii) $\mathfrak{D}_{12\text{-}3;2}^{(1)}(x) = \frac{x^7(2+2x-6x^2-x^3+6x^4+x^5-2x^6)}{(1-x^2)^3};$
- (iv) $\mathfrak{D}_{12\text{-}3;3}^{(1)}(x) = \frac{x^7(3+5x-10x^2-9x^3+10x^4+3x^5+4x^7-5x^8-x^9+2x^{10})}{(1-x^2)^4};$
- (v) $\mathfrak{D}_{12\text{-}3;4}^{(1)}(x) = \frac{x^9(5+5x-23x^2-7x^3+40x^4-x^5-30x^6+5x^7+5x^8-x^9+5x^{10}+x^{11}-2x^{12})}{(1-x^2)^5}.$

Similarly to Theorem 3.1, we have

Theorem 3.7. *Let*

$$A_k(x) = \frac{x^2}{1-x^2 G_{k-2}(x)} A_{k-1}(x) + \frac{x^4 G_{k-1}(x)}{(1-x^2 G_{k-2}(x))^2} A_{k-2}(x)$$

for all $k \geq 4$, where $A_1(x) = A_2(x) = x^2$, $A_3(x) = \frac{x^4}{1-x^2}$, and $A_4(x) = \frac{x^6(2-x^2)}{(1-x^2)^3}$. Then, for all $k \geq 2$,

$$\mathfrak{D}_{21\text{-}3\text{-}\cdots\text{-}k;1}^{(1)}(x) = A_k(x) + x A_{k-1}(x).$$

As an extension of Theorem 3.7, let us consider the case $r \geq 1$. Theorem 2.14, for given k and r , yields an explicit formula for $\mathfrak{D}_{21\text{-}3\text{-}\cdots\text{-}k;r}^{(1)}(x)$. For example, for $k = 3$ and $r = 0, 1, 2, 3, 4$, we have the following.

Theorem 3.8. *We have*

- (i) $\mathfrak{D}_{21\text{-}3;0}^{(1)}(x) = \frac{1+x+x^4-x^5}{1-x^2};$
- (ii) $\mathfrak{D}_{21\text{-}3;1}^{(1)}(x) = \frac{x^3(1+x-x^2)}{1-x^2};$
- (iii) $\mathfrak{D}_{21\text{-}3;2}^{(1)}(x) = \frac{x^5(1+2x-2x^2-x^3+x^4)}{(1-x^2)^2};$

$$(iv) \quad \mathfrak{D}_{21-3;3}^{(1)}(x) = \frac{x^5(1+x-x^2+x^3-x^4-x^5+x^6)}{(1-x^2)^2};$$

$$(v) \quad \mathfrak{D}_{21-3;4}^{(1)}(x) = \frac{x^7(1+2x-2x^2-2x^5+2x^6+x^7-x^8)}{(1-x^2)^3}.$$

4. FURTHER RESULTS

Here we present three different directions to generalize the results of the previous sections. The first of these directions is to consider one occurrence of the classical pattern 132. For example, the following result is true.

Theorem 4.1. *There does not exist a Dumont permutation of the first kind containing 132 (classical pattern) exactly once.*

Proof. Let $\pi = (\pi', n, \pi'')$ be a Dumont permutation of the first kind of length n , which contain the pattern 132 exactly once. It is easy to see that there does not exist a Dumont permutation of the first kind where $n = 0, 1, 2, 3$. Suppose $n \geq 4$, and let us assume by induction on n that there does not exist a Dumont permutation of the first kind of length $m \leq n-1$ containing 132 exactly once. To prove this property for n , let us consider the following two cases together with using Proposition 2.1: n is either an even number, or n is an odd number.

1. Let n be an odd number. Since π is a Dumont permutation of the first kind, we get $\pi'' = \emptyset$, so π contains 132 exactly once if and only if π' contains 132 exactly once.
2. Let n be an even number. Since π is a Dumont permutation of the first kind we have $\pi'' \neq \emptyset$. Now, let us consider two cases: either n does not appear in the occurrence of 132, or n does it.
 - (a) Let the occurrence of 132 does not contain the element n . So, every element of π' greater than every element of π'' . Therefore, either π' is a Dumont permutation of the first kind of length $m \leq n-2$ contains 132 exactly once, or π'' is a Dumont permutation of the first kind of length $m \leq n-1$ contains 132 exactly once.
 - (b) Let the occurrence of 132 contains the element n . So, $\pi = (\pi', a, n, \pi'', a+1, \pi''')$ (see [MV4]) such that $\pi_p = n$ and $\pi_q = a+1$, where every element of π' is greater than every element of π'' and every element of π'' is greater than every element of π''' . Since n is even number and maximal in π we have that a is an odd number, so $a+1$ is an even number. Therefore, by using Proposition 2.1 we get that p, q are even numbers, (π', a) is of odd length, and π'' is of even length. On the other hand, $q = p+1+|\pi''|$, so q is an odd number, a contradiction.

Hence, by induction on n we get the desired result. \square

The second direction is to consider more than one additional restriction. For example, the following result is true.

Theorem 4.2. *Let $k \geq 2$. The generating function for the number of Dumont permutations of the first kind in $\mathfrak{S}_n(1-3-2, 1-2-3 \cdots -k, 2-1-3 \cdots -k)$ is given by*

$$G_{k-1}(x) + xG_{k-2}(x).$$

A comparison of Theorem 4.2 with Theorem 2.8 suggests that there should exist a bijection between the sets $\mathfrak{S}_n(1-3-2, 2-1-3 \cdots -k)$ and $\mathfrak{S}_n(1-3-2, 1-2-3 \cdots -k, 2-1-3 \cdots -k)$. However, we failed to produce such a bijection, and finding it remains an open question.

The third direction is to consider another 3-letter pattern instead of 1-3-2.

Theorem 4.3. *The number of Dumont permutation of the second kind in $\mathfrak{S}_n(3\text{-}2\text{-}1)$ is the same as the number of Dumont permutation of the first kind in $\mathfrak{S}_n(2\text{-}3\text{-}1)$ (or in $\mathfrak{S}_n(3\text{-}1\text{-}2)$) which is equal to $C_{[n/2]}$.*

Acknowledgments: The author are grateful to S. Kitaev for his careful reading of the manuscript.

REFERENCES

- [Bo1] M. Bóna, The permutation classes equinumerous to the smooth class, *Electron. J. Combin.* **5** (1998) #R31.
- [BS] E. Babson and E. Steingrímsson, Generalized permutation patterns and a classification of the Mahonian statistics, *Séminaire Lotharingien de Combinatoire*, B44b:18pp, (2000).
- [BCS] P. Brändén, A. Claesson, and E. Steingrímsson, Continued fractions and increasing subsequences in permutations, *Discr. Math.*, to appear.
- [C] A. Claesson, Generalised pattern avoidance, *European Journal of Combinatorics*, **22** (2001) 961–973.
- [CM] A. Claesson and T. Mansour, Permutations avoiding a pair of generalized patterns of length three with exactly one dash, preprint CO/0107044.
- [CW] T. Chow and J. West, Forbidden subsequences and Chebyshev polynomials, *Discr. Math.* **204** (1999) 119–128.
- [D] D. Dumont, Interprétations combinatoires des nombres de Genocchi, *Duke Math. J.* **41** (1974), 305–318.
- [Ki] S. Kitaev, Multi-avoidance of generalised patterns, to appear in Discrete Mathematics.
- [Km] D. Kremer, Permutations with forbidden subsequences and a generalized Schröder number, *Discr. Math.* **218** (2000) 121–130.
- [Kr] C. Krattenthaler, Permutations with restricted patterns and Dyck paths, *Adv. in Applied Math.* **27** (2001), 510–530.
- [M1] T. Mansour, Continued fractions and generalized patterns, *European Journal of Combinatorics*, **23:3** (2002), 329–344.
- [M2] T. Mansour, Restricted 1-3-2 permutations and generalized patterns, *Annals of Combinatorics* **6** (2002), 1–12.
- [M3] T. Mansour, Continued fractions, statistics, and generalized patterns, *Ars Combinatoria*, to appear (2002), preprint CO/0110040.
- [MV1] T. Mansour and A. Vainshtein, Restricted permutations, continued fractions, and Chebyshev polynomials *Electron. J. Combin.* **7** (2000) #R17.
- [MV2] T. Mansour and A. Vainshtein, Restricted 132-avoiding permutations, *Adv. Appl. Math.* **126** (2001), 258–269.
- [MV3] T. Mansour and A. Vainshtein, Layered restrictions and Chebychev polynomials (2000), *Annals of Combinatorics, Annals of Combinatorics* **5** (2001), 451–458.
- [MV4] T. Mansour and A. Vainshtein, Restricted permutations and Chebyshev polynomials, *Séminaire Lotharingien de Combinatoire* **47** (2002), Article B47c.
- [R] A. Robertson, Permutations containing and avoiding 123 and 132 patterns, *Discrete Mathematics and Theoretical Computer Science*, **3** (1999) 151–154.
- [RWZ] A. Robertson, H. Wilf, and D. Zeilberger, Permutation patterns and continuous fractions, *Electron. J. Combin.* **6** (1999) #R38.
- [SS] R. Simion, F.W. Schmidt, Restricted Permutations, *Europ. J. of Combinatorics* **6** (1985), 383–406.
- [SP] N.J.A. Sloane and S. Plouffe, *The Encyclopedia of Integer Sequences*, Academic Press, New York (1995).
- [W] J. West, Generating trees and forbidden subsequences, *Discr. Math.* **157** (1996), 363–372.