

Knots, Links and Tangles

STEVEN FINCH

August 8, 2003

We start with some terminology from differential topology [1]. Let C be a circle and $n \geq 2$ be an integer. An **immersion** $f : C \rightarrow \mathbb{R}^n$ is a smooth function whose derivative never vanishes. An **embedding** $g : C \rightarrow \mathbb{R}^n$ is an immersion that is one-to-one. It follows that $g(C)$ is a manifold but $f(C)$ need not be (f is only locally one-to-one, so consider the map that twists C into a figure eight).

A **knot** is a smoothly embedded circle in \mathbb{R}^3 ; hence a knot is a closed spatial curve with no self-intersections. Two knots J and K are **equivalent** if there is a homeomorphism $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ taking J onto K . This implies that the complements $\mathbb{R}^3 - J$ and $\mathbb{R}^3 - K$ are homeomorphic as well.

A **link** is a compact smooth 1-dimensional submanifold of \mathbb{R}^3 . The connected components of a link are disjoint knots, often with intricate intertwinings. Two links L and M are **equivalent** if, likewise, there is a homeomorphism $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ taking L onto M .

We can project a knot or a link into the plane in such a way that its only self-intersections are transversal double points. Ambiguity is removed by specifying at each double point which arc passes over and which arc passes under. Over all possible such projections of K or L , determine one with the minimum number of double points; this defines the **crossing number** of K or L .

There is precisely 1 knot with 0 crossings (the circle), 1 knot with 3 crossings (the trefoil), and 1 knot with 4 crossings. Note that, although the left-hand trefoil T_L is not ambiently isotopic (i.e., deformable) to the right-hand trefoil T_R , a simple reflection about a plane gives T_R as a homeomorphic image of T_L . Under our definition of equivalence, chiral pairs as such are counted only once.

There are precisely 2 knots with 5 crossings, and 5 knots with 6 crossings. In particular, there is no homeomorphism $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ taking the granny knot $T_L \# T_L$ onto the square knot $T_L \# T_R$, where $\#$ denotes the connected sum of manifolds [2, 3]. (See Figure 1.) Also, there are precisely 8 knots with 7 crossings, and 25 knots with 8 crossings.

A link L is **splittable** if we can embed a plane in \mathbb{R}^3 , disjoint from L , that separates one or more components of L from other components of L . There are precisely 1, 0, 1, 1, 3, 4, 15 nonsplittable links with 0, 1, 2, 3, 4, 5, 6 crossings, respectively.

⁰Copyright © 2003 by Steven R. Finch. All rights reserved.




Figure 1: Four famous knots (T_L and T_R are prime and equivalent; $T_L \# T_R$ and $T_L \# T_L$ are composite and distinct).




Figure 2: All two-component prime links with crossing number ≤ 5 .

A knot K or nonsplittable link L is **prime** if it is not a circle and if, for any plane P that intersects K or L transversely in exactly two points, P slices off merely an unknotted arc away from the rest. (See Figure 2.) Otherwise it is **composite**. For example, $T_L \# T_L$ and $T_L \# T_R$ are composite knots, each being nontrivial connected sums of knots. Every knot decomposes as a unique connected sum of prime knots [4].

People have known for a long time that there exist non-equivalent links with homeomorphic complements [5, 6]. This cannot happen for knots, as recently proved by Gordon & Luecke [7, 8].

Let B denote the compact unit ball in \mathbb{R}^3 and ∂B denote its boundary. A **tangle** U is a smooth 1-dimensional submanifold of B meeting ∂B transversely at the four points

$$NE = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0\right), \quad NW = \left(\frac{-1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0\right), \quad SW = \left(\frac{-1}{\sqrt{2}}, \frac{-1}{\sqrt{2}}, 0\right), \quad SE = \left(\frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{2}}, 0\right)$$




Figure 3: All prime alternating tangles with crossing number ≤ 3 .

and meeting ∂B nowhere else. Thus U is a union of two smoothly embedded line segments in B with distinct endpoints on ∂B , together with an arbitrary number of smoothly embedded circles in the interior of B , all disjoint but often intertwined. Two tangles U and V are **(strongly) equivalent** if there is a homeomorphism $B \rightarrow B$ that takes U onto V , is orientation-preserving on B , and leaves ∂B fixed pointwise. The crossing number of a tangle is defined via projections as before. Tangles form the building blocks of knots and links [9, 10, 11]; the first precise asymptotic enumeration results discovered in this subject concerned tangles (as we shall soon see).

A tangle is **trivial** if it is only the union of the two line segments $NW-NE$ and $SW-SE$, or the union of the two line segments $SW-NW$ and $SE-NE$. A tangle U is **prime** if it is not trivial; if, for any sphere S in B that is disjoint from U , no portion of U is enclosed by S ; and if, for any sphere S in B that intersects U transversely in exactly two points, S encloses merely an unknotted arc of U . (See Figures 3 and 4.)

Finally, a knot, link or tangle is **alternating** if, for some projection, as we proceed along any connected component in the projection plane from beginning to end, the sequence of underpasses and overpasses is strictly alternating. The first non-alternating knots appear with crossing number ≥ 8 . General references on knot theory include [12, 13, 14, 15, 16, 17].




Figure 4: Five of the 4-crossing prime alternating tangles; the other five are obtained by rotating through 90° (and switching crossings to maintain the convention that the NW strand is an underpass).

0.1. Prime Alternating Tangles. Let a_n denote the number of prime alternating tangles with n crossings (up to strong equivalence) and let $A(x) = \sum_{n=1}^{\infty} a_n x^n$ be the corresponding generating function. Then [18]

$$A(x) = x + 2x^2 + 4x^3 + 10x^4 + 29x^5 + 98x^6 + 372x^7 + 1538x^8 + 6755x^9 + 30996x^{10} + \dots$$

satisfies the equation

$$A(x)(1+x) - A(x)^2 - (A(x)+1)r(A(x)) - x - 2\frac{x^2}{1-x} = 0$$

where the algebraic function $r(x)$ is defined by

$$r(x) = \frac{(1-4x)^{\frac{3}{2}} + (2x^2 - 10x - 1)}{2(x+2)^3} - \frac{2}{1+x} - x + 2$$

Further, $A(x)$ satisfies the irreducible quintic equation

$$\begin{aligned} 0 = & (x^4 - 2x^3 + x^2)A(x)^5 + (8x^4 - 14x^3 + 8x^2 - 2x)A(x)^4 + \\ & (25x^4 - 16x^3 - 14x^2 + 8x + 1)A(x)^3 + (38x^4 + 15x^3 - 30x^2 - x + 2)A(x)^2 + \\ & (28x^4 + 36x^3 - 5x^2 - 12x + 1)A(x) + (8x^4 + 17x^3 + 8x^2 - x) \end{aligned}$$

Sundberg & Thistlethwaite [19] proved the above remarkable formulas, as well as the following asymptotics:

$$a_n \sim \frac{3\alpha}{4\sqrt{\pi}} n^{-\frac{5}{2}} \lambda^{n-\frac{3}{2}} \sim \frac{3}{4} \sqrt{\frac{\beta}{\pi}} n^{-\frac{5}{2}} \lambda^n$$

where

$$\alpha = \frac{5^{\frac{7}{2}}}{3^5 \sqrt{2}} \sqrt{\frac{(21001 + 371\sqrt{21001})^3}{(17 + 3\sqrt{21001})^5}} = 3.8333138762\dots$$

$$\beta = \alpha^2 \lambda^{-3} = 0.0632356411\dots$$

and

$$\lambda = \frac{101 + \sqrt{21001}}{40} = 6.1479304437\dots$$

A completely different approach to the solution of this problem appears in [20].

Let \hat{a}_n denote the number of n -crossing prime alternating tangles with exactly two components. That is, no circles are allowed. A two-component tangle is also known as a **knot with four external legs**. The sequence [18, 21, 22]

$$\{\hat{a}_n\}_{n=1}^{\infty} = \{1, 2, 4, 8, 24, 72, 264, 1074, 4490, 20296, 92768, \dots\}$$

is believed to possess a leading term of the form $\hat{\lambda}^n$ with $\hat{\lambda} < \lambda$, but more intensive analysis is needed to compute $\hat{\lambda}$.

0.2. Prime Alternating Links. Let b_n denote the number of prime alternating links with n crossings (up to equivalence), then the sequence [23, 24]

$$\{b_n\}_{n=1}^{\infty} = \{0, 1, 1, 2, 3, 8, 14, 39, 96, 297, 915, 3308, 12417, \dots\}$$

satisfies the following asymptotics [25]:

$$b_n \sim \frac{3}{16\gamma} \sqrt{\frac{\beta}{\pi}} n^{-\frac{7}{2}} \lambda^n$$

where

$$\gamma = \frac{1}{2} \left(\frac{371}{\sqrt{21001}} - 1 \right) = 0.7800411357\dots$$

and λ, β are as before. This is a somewhat more precise result than that proved in [19].

Let c_n denote the number of prime links with n crossings (including both alternating and non-alternating links), then we have [23, 26, 27]

$$\{c_n\}_{n=1}^{\infty} = \{0, 1, 1, 2, 3, 9, 16, 50, 132, 452, 1559, \dots\}$$

The value c_{12} is not known. Stoimenow [28], building on Ernst & Sumners [29] and Welsh [30], proved that

$$4 \leq \liminf_{n \rightarrow \infty} c_n^{1/n} \leq \limsup_{n \rightarrow \infty} c_n^{1/n} \leq \frac{\sqrt{13681} + 91}{20} = 10.3982903484\dots$$

but further improvements in the upper bound are likely. The two-component analogs [23]

$$\begin{aligned}\{\hat{b}_n\}_{n=1}^\infty &= \{0, 1, 0, 1, 1, 3, 6, 14, 42, 121, 384, 1408, 5100, 21854, \dots\} \\ \{\hat{c}_n\}_{n=1}^\infty &= \{0, 1, 0, 1, 1, 3, 8, 16, 61, 185, 638 \dots\}\end{aligned}$$

also await study.

0.3. Prime Alternating Knots. Let d_n denote the number of prime alternating knots with n crossings (up to equivalence), then the sequence [31]

$$\{d_n\}_{n=1}^\infty = \{0, 0, 1, 1, 2, 3, 7, 18, 41, 123, 367, 1288, 4878, 19536, \dots\}$$

is more difficult and only *conjectured* to satisfy the following asymptotics [32]:

$$d_n \sim \eta \cdot n^\xi \cdot \kappa^n$$

where

$$\xi = -\frac{\sqrt{13} + 1}{6} - 3 = -3.7675918792\dots$$

Thistlethwaite [33] proved that

$$\limsup_{n \rightarrow \infty} d_n^{1/n} < \lambda$$

and further claimed that $\lim_{n \rightarrow \infty} d_n^{1/n}$ exists. If the conjectured asymptotic form for d_n is true, it would follow that $\kappa < \lambda$. Again, more intensive analysis is needed to compute κ . Might it be true that $\kappa = \hat{\lambda}$ [22]?

Let e_n denote the number of prime knots with n crossings (including both alternating and non-alternating knots), then we have [31]

$$\{e_n\}_{n=1}^\infty = \{0, 0, 1, 1, 2, 3, 7, 21, 49, 165, 552, 2176, 9988, 46972, \dots\}$$

The value e_{17} is not known. Welsh [30] proved that

$$2.68 \leq \liminf_{n \rightarrow \infty} e_n^{1/n}$$

and clearly Stoimenow's upper bound 10.40 applies to the limit superior. Sharper bounds for both $\{c_n\}$ and $\{e_n\}$ would be good to see.




Figure 5: All closed planar curves with crossing number ≤ 2 .

0.4. Planar Curves. Here are enumeration problems that seem to be even more complicated than those in knot theory [34, 35, 36, 37, 38]. A **closed planar curve** is a smoothly immersed circle in \mathbb{R}^2 whose only self-intersections are transversal double points. Define an equivalence relation between closed planar curves in the same manner as between knots, with the additional condition that the homeomorphism $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ is orientation-preserving. (See Figure 5.)

An **open planar curve** is a smoothly immersed line in \mathbb{R}^2 , given by $h : \mathbb{R} \rightarrow \mathbb{R}^2$, whose only self-intersections are transversal double points and which satisfies $h(x) = (x, 0)$ for all sufficiently large $|x|$. Such a curve is also known as a **knot with two external legs**. Define an equivalence relation between open planar curves in the same manner as between closed planar curves. Note that, unlike closed curves, open curves are oriented from the initial point $(-\infty, 0)$ to the final point $(\infty, 0)$. (See Figure 6.)

Let p_n and q_n denote the number of n -crossing closed curves and open curves, respectively. The sequences [39, 40]

$$\{p_n\}_{n=0}^\infty = \{1, 2, 5, 20, 82, 435, 2645, 18489, 141326, 1153052, 9819315, \dots\}$$

$$\{q_n\}_{n=0}^\infty = \{1, 2, 8, 42, 260, 1796, 13396, 105706, 870772, 7420836, 65004584, \dots\}$$

are *conjectured* to satisfy the following asymptotics [32]:

$$p_n \sim \frac{1}{4}q_n \sim \omega \cdot n^\theta \cdot \mu^n$$

where $\theta = \xi + 1 = -2.7675918792\dots$. Numerically, we have $\mu = 11.4\dots$ [22]. There is a great amount of work to be done in this area.




Figure 6: All open planar curves with crossing number ≤ 2 .

REFERENCES

- [1] V. Guillemin and A. Pollack, *Differential Topology*, Prentice-Hall, 1974, pp. 13–18, 27–32; MR 50 #1276.
- [2] H. Seifert, Verschlingungsinvarianten, *Sitzungsber. Preuss. Akad. Wiss.* 26 (1933) 811–823.
- [3] R. H. Fox, On the complementary domains of a certain pair of inequivalent knots, *Proc. Konink. Nederl. Akad. Wetensch. Ser. A* 55 (1952) 37–40; *Indag. Math.* 14 (1952) 37–40; MR 13,966c.
- [4] H. Schubert, Die eindeutige Zerlegbarkeit eines Knotens in Primknoten, *Sitzungsberichte der Heidelberger Akademie der Wissenschaften Mathematisch-Naturwissenschaftliche Klasse* (1949) n. 3, 57–104; MR 11,196f.
- [5] J. H. C. Whitehead, On doubled knots, *J. London Math. Soc.* 12 (1937) 63–71.
- [6] D. Rolfsen, *Knots and Links*, Publish or Perish, 1976, p. 49; MR 58 #24236.
- [7] C. M. Gordon and J. Luecke, Knots are determined by their complements, *J. Amer. Math. Soc.* 2 (1989) 371–415; *Bull. Amer. Math. Soc.* 20 (1989) 83–87; MR 90a:57006a–b.

- [8] B. A. Cipra, To have and have knot: When are knots alike? *Science* 241 (1988) 1291–1292; MR 89k:57009.
- [9] J. H. Conway, An enumeration of knots and links, and some of their algebraic properties, *Computational Problems in Abstract Algebra*, Proc. 1967 Oxford conf., ed. J. Leech, Pergamon Press, 1970, pp. 329–358; MR 41 #2661.
- [10] W. B. R. Lickorish, Prime knots and tangles, *Trans. Amer. Math. Soc.* 267 (1981) 321–332; MR 83d:57004.
- [11] M. Eudave Muñoz, Primeness and sums of tangles, *Trans. Amer. Math. Soc.* 306 (1988) 773–790; MR 89g:57005.
- [12] R. H. Crowell and R. H. Fox, *Introduction to Knot Theory*, Springer-Verlag, 1977; MR 26 #4348.
- [13] M. Thistlethwaite, Knot tabulations and related topics, *Aspects of Topology*, ed. I. M. James and E. H. Kronheimer, Cambridge Univ. Press, 1985, pp. 1–76; MR 86j:57004.
- [14] G. Burde and H. Zieschang, *Knots*, de Gruyter, 1985; MR 87b:57004.
- [15] C. C. Adams, *The Knot Book: An Elementary Introduction to the Mathematical Theory of Knots*, W. H. Freeman, 1994; MR 94m:57007.
- [16] W. B. R. Lickorish, *An Introduction to Knot Theory*, Springer-Verlag, 1997; MR 98f:57015.
- [17] J. Hoste, M. Thistlethwaite, and J. Weeks, The first 1,701,936 knots, *Math. Intelligencer* v. 20 (1998) n. 4, 33–48; MR 99i:57015.
- [18] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A047051 and A067646.
- [19] C. Sundberg and M. Thistlethwaite, The rate of growth of the number of prime alternating links and tangles, *Pacific J. Math.* 182 (1998) 329–358; MR 98m:57013.
- [20] P. Zinn-Justin and J.-B. Zuber, Matrix integrals and the counting of tangles and links, *Discrete Math.* 246 (2002) 343–360; math-ph/9904019; MR 2003i:57019.
- [21] P. Zinn-Justin and J.-B. Zuber, On the counting of colored tangles, *J. Knot Theory Ramifications* 9 (2000) 1127–1141; math-ph/0002020; MR 2002b:57008.

- [22] J. L. Jacobsen and P. Zinn-Justin, A transfer matrix approach to the enumeration of knots, *J. Knot Theory Ramifications* 11 (2002) 739–758; math-ph/0102015; MR 2003f:57014.
- [23] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A049344, A048952, A048953, A059739, A059741, and A086771.
- [24] M. Thistlethwaite, Prime Unoriented Alternating Links to 19 Crossings, http://www.math.utk.edu/~morwen/png/link_stats.png.
- [25] S. Kunz-Jacques and G. Schaeffer, The asymptotic number of prime alternating links, presentation at *Formal Power Series and Algebraic Combinatorics (FPSAC)* conf., Tempe, Arizona, 2001.
- [26] D. Bar-Natan, The Hoste-Thistlethwaite Link Table, <http://www.math.toronto.edu/~drorbn/KAtlas/Links/index.html>.
- [27] R. G. Scharein, Number of Prime Links, <http://www.pims.math.ca/knotplot/manual/PrimeLinks.html>.
- [28] A. Stoimenow, On the number of links and link polynomials, *Quart. J. Math.*, to appear (2003).
- [29] C. Ernst and D. W. Sumners, The growth of the number of prime knots, *Math. Proc. Cambridge Philos. Soc.* 102 (1987) 303–315; MR 88m:57006.
- [30] D. J. A. Welsh, On the number of knots and links, *Sets, Graphs and Numbers*, Proc. 1991 Budapest conf., Colloq. Math. Soc. János Bolyai 60, North-Holland, 1992, pp. 713–718; MR 94f:57010.
- [31] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A002863 and A002864.
- [32] G. Schaeffer and P. Zinn-Justin, On the asymptotic number of plane curves and alternating knots, math-ph/0304034.
- [33] M. Thistlethwaite, On the structure and scarcity of alternating links and tangles, *J. Knot Theory Ramifications* 7 (1998) 981–1004; MR 99k:57031.
- [34] V. I. Arnold, *Topological Invariants of Plane Curves and Caustics*, Amer. Math. Soc., 1994; MR 95h:57003.
- [35] V. I. Arnold, Plane curves, their invariants, perestroikas and classifications, *Singularities and Bifurcations*, ed. V. I. Arnold, Adv. Soviet Math. 21, Amer. Math. Soc., 1994, pp. 33–91; MR 95m:57009.

- [36] S. M. Guseĭn-Zade, On the enumeration of curves from infinity to infinity, *Singularities and Bifurcations*, ed. V. I. Arnold, Adv. Soviet Math. 21, Amer. Math. Soc., 1994, pp. 189–198; MR 96d:57029.
- [37] S. M. Guseĭn-Zade and F. S. Duzhin, On the number of topological types of plane curves (in Russian), *Uspekhi Mat. Nauk* v. 53 (1998) n. 3, 197–198; Engl. transl. in *Russian Math. Surveys* 53 (1998) 626–627; MR 99i:57001.
- [38] S. K. Lando, On enumeration of unicursal curves, *Differential and Symplectic Topology of Knots and Curves*, ed. S. Tabachnikov, Amer. Math. Soc., 1999, pp. 77–81; MR 2000m:57012.
- [39] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A008983 and A054993.
- [40] F. S. Duzhin, Number of Topological Types of Plane Curves with Simple Double Points, <http://www.pdmi.ras.ru/arnsem/dataprogram/allcurve.txt>.