

Enumerating alcohols and other classes of chemical molecules, an example of Polya theory

Frederic Chyzak
(Version of January 13, 1997)

Alkanes are a simple class of chemical compounds. They are generically described by the chemical formula $C_n H_{2n+2}$. First examples for small n are methane ($n = 1$), ethane ($n = 2$), propane ($n = 3$), butane ($n = 4$), a.s.o. For a given n however, there exist several different *isomers*, i.e., different structures of bonds between atoms. In chemistry, there is much interest in knowing the number, or better yet the list, of such isomers. Alcohols are obtained from alkanes by replacing a hydrogen atom by an *OH* group. It follows that they are isomorphic to carbon chains with a distinguished node, or again to alkyl radicals $C_n H_{2n+1}$, which are alkanes with a missing hydrogen atom. If we disregard geometrical constraints (i.e., if we consider *structural* isomers only, and not *conformational* isomers), this leads to a pure graph-theoretical problem: how many rooted trees are there with n internal nodes, where each internal node has degree 4?

In this session, we thus consider *rooted* trees, so that we count and enumerate alkyls, with generic formula $C_n H_{2n+1}$. The combinatorics also corresponds to simple alcohols $C_n H_{2n+1} OH$, organo-metallic compounds $C_n H_{2n+1} X$, and any other monosubstituted alkanes. We next treat the cases of disubstituted and trisubstituted alkanes. We develop the study of our models using the package [Combstruct](#).

```
> with(combstruct);  
[allstructs, count, draw, finished, gfeqns, gfsolve, iterstructs, nextstruct, prog_gfeqns,  
 prog_gfsolve, prog_gfsolve]
```

Enumerations of such classes of chemical compounds are part of Polya theory. We refer to the book by G. Polya and R. C. Read [*Combinatorial Enumeration of Groups, Graphs, and Chemical Compounds*, (1987), Springer-Verlag] for more extensive results.

■ Monosubstituted alkanes, $C_n H_{2n+1} X$

In this section, we study monosubstituted alkanes, i.e., *rooted* trees, first without any constraint, next according to the *height*.

■ General alkyls

■ Definition

An alkyl radical can be viewed as a carbon atom linked to at most 3 alkyl radicals. Thus, we only take into account hydrogen atoms implicitly. There is no loss of information, since hydrogen atoms can always be recovered from the carbon skeleton. This yields the class equation $Alkyl = Carbon (E + Alkyl + Alkyl^2 + Alkyl^3)$, which we map into the following grammar:

```
> gramm_Alkyl:=Alkyl=Prod(Carbon,Set(Alkyl,card<=3)),Carbon=Atom:  
 specs_Alkyl:=[Alkyl,{gramm_Alkyl},unlabelled]:
```

Note that since the Set construct denotes multisets, i.e., sets with repetitions, a carbon atom of an alkyl is allowed to be bound to two copies of the same subtree (but the order of the subtrees does not matter).

Define the size of an alkyl as the number of carbon atoms it contains. We compute the number of alkyls of a given size using [combstruct\[count\]](#).

```
> seq(count(specs_Alkyl, size=i), i=0..50);
```

0, 1, 1, 2, 4, 8, 17, 39, 89, 211, 507, 1238, 3057, 7639, 19241, 48865, 124906,
321198, 830219, 2156010, 5622109, 14715813, 38649152, 101821927,
269010485, 712566567, 1891993344, 5034704828, 13425117806, 35866550869,
95991365288, 257332864506, 690928354105, 1857821351559, 5002305607153,
13486440075669, 36404382430278, 98380779170283, 266158552000477,
720807976831447, 1954002050661819, 5301950692017063,
14398991611139217, 39137768751465752, 106465954658531465,
289841389106439413, 789642117549095761, 2152814945971655556,
5873225808361331954, 16033495247557039074, 43797554941937577760

This series appears as the entry **M1146** ("quartic planted trees with n nodes") in the book by N. J. A. Sloane and S. Plouffe [*The Encyclopedia of Integer Sequences*, (1995), Academic Press].

Here is an example of an alkyl with 6 carbon atoms, obtained by the command `combstruct[draw]`.

```
> alk:=draw(specs_Alkyl,size=6);
alk := Prod(Carbon, Set(Prod(Carbon, Set(
    Prod(Carbon, Set(Prod(Carbon, E)))), Prod(Carbon, Set(Prod(Carbon, E)))))))
```

The following procedure rewrites an alkyl into a more readable way.

```
> nice:=proc(alk)
   eval(subs({Epsilon=NULL,Carbon=C,Prod=proc() global H;
   [args] end,Set=proc() args end},alk)) end:
> nice(alk);
[C,[C,[C,[C]], [C,[C]]]]
```

The following procedure computes the size of a given alkyl.

```
> size:=proc(alk) option remember;
   1+convert(map(size,op(2,alk)), '+') end:
size(Prod(Carbon,Epsilon)):=1:
```

The following procedure computes the height of a given alkyl.

```
> height:=proc(alk) option remember;
1+max(op(map(height,op(2,alk)))) end:
height(Prod(Carbon,Epsilon)):=1:
```

Here is an alkyl with 50 carbon atoms, its nice representation and height.

```
> alk:=draw(specs_Alkyl,size=50):
```

```
> nice(alk) ;
```

```

    [C, [C, [C, [C, [C]], [C, [C], [C]]]]], [C, [C, [C, [C]]], [C, [C, [C]]]]]]]
]
> height(alk) ;

```

13

Empirical study

Drawing

By drawing several random structures, we can study probabilistic properties of alkyls. For instance, the following is a probabilistic estimate of their height on average:

```

> for i to 10 do
  ho[i]:=height(draw(specs_Alkyl,size=50)) od;
  ho1:=18
  ho2:=15
  ho3:=19
  ho4:=13
  ho5:=13
  ho6:=15
  ho7:=13
  ho8:=16
  ho9:=14
  ho10:=17
> add(ho[i],i=1..10)/10. ;
15.30000000

```

In the same way, we get a probabilistic estimate of their standard deviation:

```

> sqrt(add((ho[i]-")^2,i=1..10)/10) ;
2.051828453

```

Exhaustive enumeration

The command [combstruct\[draw\]](#) permits us to draw *one* structure at random. We can also generate *all* alkyls of a given size, using [combstruct\[allstructs\]](#), so as to compute the mean of a particular parameter exactly, or to count all those with a particular property. For instance, the height of trees cannot be represented in the class of combinatorial structures when using [Combstruct](#). For instance, by computing all alkyls of size 5, we get the distribution of height for these alkyls (in their nice representation).

```

> allstructs(specs_Alkyl,size=5) :
> map(nice,) ;
[[C, [C, [C], [C, [C]]]], [C, [C], [C], [C, [C]]], [C, [C], [C, [C], [C]]],
 [C, [C, [C, [C, [C]]]]], [C, [C, [C], [C], [C]]], [C, [C, [C, [C], [C]]]],
 [C, [C], [C, [C, [C]]]], [C, [C, [C]], [C, [C]]]]
> sort(map(height, ""));

```

[3, 3, 3, 3, 4, 4, 4, 5]

[Here we count 4 alkyls of size 5 and height 3, 3 alkyls of size 5 and height 4, and 1 alkyl of size 5 and height 5.

[By the same method, we get the exact mean and standard deviation of the height for small sizes.

```
> esd:=proc(n) local i,as,mean;
    as:=map(height,allstructs(specs_Alkyl,size=n));
    mean:=evalf(convert(as,'+')/nops(as));

    nops(as),mean,evalf(sqrt(add((i-mean)^2,i=as))/nops(as))
end:
> for i from 2 to 6 do i=esd(i) od;
    2=(1, 2., 0)
    3=(2, 2.500000000, .3535533906)
    4=(4, 3., .3535533905)
    5=(8, 3.625000000, .2460627461)
    6=(17, 4.117647059, .2017630413)
```

[We could go up to $i = 9$ in less than 2 minutes.

Alkyls according to their height

Grammar

[We define the class Alkyl_height_n to be the class of alkyls of height *at most* n . An alkyl of height at most n can be viewed as a carbon atom linked to at most three alkyls of height at most $n - 1$, according to the equation

$$\text{Alkyl_height}_n = \text{Carbon} (\text{E} + \text{Alkyl_height}_{n-1} + \text{Alkyl_height}_{n-1}^2 + \text{Alkyl_height}_{n-1}^3).$$

```
> gramm_ltd_height:=proc(n) option remember;

Alkyl_height[n]:=Prod(Carbon,Set(Alkyl_height[n-1],card<=3)),gramm_ltd_height(n-1)
end:
gramm_ltd_height(1):=Alkyl_height[1]:=Prod(Carbon,Epsilon
n),Carbon=Atom:
specs_ltd_height:=proc(n) option remember;
    [Alkyl_height[n],{gramm_ltd_height(n)},unlabelled]
end:
```

[The following procedure rewrites an alkyl into a more readable way.

```
> nice:=proc(alk)
    eval(subs({Epsilon=NULL,Carbon=C,Prod=proc() global H;
    [args] end,Set=proc() args end},alk)) end:
```

[The following procedures compute the size and height of a given alkyl.

```
> size:=proc(alk) option remember;
    1+convert(map(size,op(2,alk)), '+') end:
size(Prod(Carbon,Epsilon)):=1:
```

```

    > height:=proc(alk) option remember;
      1+max(op(map(height,op(2,alk)))) end:
    height(Prod(Carbon,Epsilon)):=1:
[ For instance, we compute the height of a random alkyl of size 10 and height at most 5.
[ > alk:=draw(specs_ltd_height(5),size=10):
[ > nice(alk);
[ [C,[C],[C,[C]], [C,[C,[C], [C], [C,[C]]]]]
[ > size(alk),height(alk);
[ 10, 5

```

[In this section, we proceed to compute a table of the number of alkyls according to their size and height. The first method is by generating all structures. Next, we use generating functions to extend the table.

Generating all structures

[The following procedure remembers all the alkyls of a given size and with bounded height.

```

    > list_all_st:=proc(d,s) option remember;
      allstructs(specs_ltd_height(d),size=s) end:

```

[An alkyl of height h has a size at most $\frac{3^h - 1}{2}$. Therefore, to produce all alkyls with

[height at most h_{max} , we need to produce all alkyls with size up to $s_{max} = \frac{3^{h_{max}} - 1}{2}$.

```

    > s[max]:=(3^h[max]-1)/2;

```

$$s_{max} := \frac{1}{2} 3^{h_{max}} - \frac{1}{2}$$

[To begin with, we enumerate all alkyls with height at most 3.

```

    > h[max]:=3;



```

$$h_{max} := 3$$

```

    > for i from 1 to s[max] do
      i,nops(list_all_st(h[max],i)),map(nice,list_all_st(h[max],i)) od;

```



```

8, 4, [[C, [C], [C, [C], [C]], [C, [C], [C]]],  

[C, [C, [C], [C]], [C, [C], [C, [C]]], [C, [C], [C, [C]]], [C, [C], [C, [C], [C]]],  

[C, [C, [C]], [C, [C]], [C, [C], [C]]]]]  

9, 4, [[C, [C], [C, [C], [C]], [C, [C], [C], [C]]],  

[C, [C, [C]], [C, [C]], [C, [C], [C, [C]]],  

[C, [C, [C]], [C, [C], [C]], [C, [C], [C]]],  

[C, [C, [C], [C], [C]], [C, [C], [C], [C]]]]]  

10, 3, [[C, [C, [C]], [C, [C], [C]], [C, [C], [C], [C]]],  

[C, [C], [C, [C], [C]], [C, [C], [C, [C]]],  

[C, [C, [C], [C]], [C, [C], [C]], [C, [C], [C]]]]]  

11, 2, [[C, [C, [C]], [C, [C], [C]], [C, [C], [C], [C]]],  

[C, [C, [C], [C]], [C, [C], [C]], [C, [C], [C], [C]]]]]  

12, 1, [[C, [C, [C], [C]], [C, [C], [C], [C]], [C, [C], [C], [C], [C]]]]]  

13, 1, [[C, [C, [C], [C], [C]], [C, [C], [C], [C], [C]], [C, [C], [C], [C], [C]]]]]

```

In this way, we have obtained the truncation of the bivariate generating function of alkyls with size marked by z and height by u .

```

> enum_BGF:=map(series,series(convert(map(proc(s,z,u)
z^size(s)*u^height(s)
end,map(op,[seq(list_all_st(h[max],i),i=1..s[max])]),z,
u),`+`),z,infinity),u,infinity);
enum_BGF:=(u) z+(u^2) z^2+(u^2+u^3) z^3+(u^2+2 u^3) z^4+(4 u^3) z^5+(4 u^3) z^6
+(5 u^3) z^7+(4 u^3) z^8+(4 u^3) z^9+(3 u^3) z^10+(2 u^3) z^11+(u^3) z^12+(u^3) z^13

```

Generating functions

`combstruct[gfeqns]` returns a system of functional equations satisfied by the generating functions of related combinatorial structures. In the case of the alkyls with maximum height above, we get the following triangular system.

```

> gfeqns(op(2..3,specs_ltd_height(4)),z);
[ Carbon(z)=z, Alkyl_height2(z)=Carbon(z)  $\left( 1 + \text{Alkyl\_height}_1(z) \right.$ 
 $\left. + \frac{1}{2} \text{Alkyl\_height}_1(z^2) + \frac{1}{2} \text{Alkyl\_height}_1(z)^2 + \frac{1}{3} \text{Alkyl\_height}_1(z^3) \right.$ 
 $\left. + \frac{1}{2} \text{Alkyl\_height}_1(z) \text{Alkyl\_height}_1(z^2) + \frac{1}{6} \text{Alkyl\_height}_1(z)^3 \right) \text{Alkyl\_height}_3(z)$ 
=Carbon(z)  $\left( 1 + \text{Alkyl\_height}_2(z) + \frac{1}{2} \text{Alkyl\_height}_2(z^2) + \frac{1}{2} \text{Alkyl\_height}_2(z)^2 \right.$ 
 $\left. + \frac{1}{3} \text{Alkyl\_height}_2(z^3) + \frac{1}{2} \text{Alkyl\_height}_2(z) \text{Alkyl\_height}_2(z^2) \right)$ 

```

$$\begin{aligned}
& + \frac{1}{6} Alkyl_height_2(z)^3 \Big) Alkyl_height_1(z) = \text{Carbon}(z), Alkyl_height_4(z) = \\
& \text{Carbon}(z) \left(1 + Alkyl_height_3(z) + \frac{1}{2} Alkyl_height_3(z^2) + \frac{1}{2} Alkyl_height_3(z)^2 \right. \\
& + \frac{1}{3} Alkyl_height_3(z^3) + \frac{1}{2} Alkyl_height_3(z) Alkyl_height_3(z^2) \\
& \left. + \frac{1}{6} Alkyl_height_3(z)^3 \right) \Big] \\
> gfsol := \text{gfsolve}(\text{op}(2..3, \text{specs_ltd_height}(4)), z); \\
gfsol := \{ Alkyl_height_3(z) = \\
z + 5z^7 + 4z^9 + 3z^{10} + z^{13} + z^2 + 2z^{11} + z^{12} + 4z^6 + 4z^8 + 2z^3 + 3z^4 + 4z^5, \\
Alkyl_height_4(z) = 12z^6 + 31z^8 + 47z^9 + 137z^{12} + z + 184z^{13} + 70z^{10} + 99z^{11} \\
+ 7z^5 + 20z^7 + 300z^{15} + 498z^{18} + 594z^{20} + 453z^{26} + 570z^{24} + 369z^{16} + 614z^{21} \\
+ 378z^{27} + 181z^{30} + z^{39} + 56z^{33} + 12z^{36} + 624z^{22} + 239z^{14} + 3z^{38} + 6z^{37} \\
+ 37z^{34} + 20z^{35} + 128z^{31} + 312z^{28} + 238z^{29} + 89z^{32} + 601z^{23} + 514z^{25} \\
+ 432z^{17} + 551z^{19} + z^{40} + z^2 + 2z^3 + 4z^4, Alkyl_height_1(z) = z, \text{Carbon}(z) = z, \\
Alkyl_height_2(z) = z + z^2 + z^3 + z^4 \}
\end{aligned}$$

In particular, we have obtained a truncation of the bivariate generating function of all alkyls (i.e., with no constraint on height). In this series, u marks the height. It extends the previous truncation *enum_BGF*.

$$\begin{aligned}
> \text{BGF} := \text{map}(\text{series}, \text{series}(\text{eval}(\text{subs}(Alkyl_height[0]=0, \text{gfsol} \\
1, \text{add}(u^h * (\text{Alkyl_height}[h] - \text{Alkyl_height}[h-1])(z), h=1..4) \\
))), z, \text{infinity}), u, \text{infinity}); \\
BGF := (u)z + (u^2)z^2 + (u^2 + u^3)z^3 + (u^2 + 2u^3 + u^4)z^4 + (4u^3 + 3u^4)z^5 + \\
(4u^3 + 8u^4)z^6 + (5u^3 + 15u^4)z^7 + (4u^3 + 27u^4)z^8 + (4u^3 + 43u^4)z^9 + \\
(3u^3 + 67u^4)z^{10} + (2u^3 + 97u^4)z^{11} + (u^3 + 136u^4)z^{12} + (u^3 + 183u^4)z^{13} + \\
(239u^4)z^{14} + (300u^4)z^{15} + (369u^4)z^{16} + (432u^4)z^{17} + (498u^4)z^{18} + (551u^4)z^{19} + \\
(594u^4)z^{20} + (614u^4)z^{21} + (624u^4)z^{22} + (601u^4)z^{23} + (570u^4)z^{24} + \\
(514u^4)z^{25} + (453u^4)z^{26} + (378u^4)z^{27} + (312u^4)z^{28} + (238u^4)z^{29} + (181u^4)z^{30} + \\
(128u^4)z^{31} + (89u^4)z^{32} + (56u^4)z^{33} + (37u^4)z^{34} + (20u^4)z^{35} + (12u^4)z^{36} + \\
(6u^4)z^{37} + (3u^4)z^{38} + (u^4)z^{39} + (u^4)z^{40}
\end{aligned}$$

This is made explicit on the following normalized difference: each entry starts with a term in u^4 , denoting alkyls with height at least 4.

$$\begin{aligned}
> \text{map}(\text{series}, \text{series}(\text{BGF} - \text{enum_BGF}, z, \text{infinity}), u, \text{infinity}); \\
(u^4)z^4 + (3u^4)z^5 + (8u^4)z^6 + (15u^4)z^7 + (27u^4)z^8 + (43u^4)z^9 + (67u^4)z^{10} \\
+ (97u^4)z^{11} + (136u^4)z^{12} + (183u^4)z^{13} + (239u^4)z^{14} + (300u^4)z^{15} + \\
(369u^4)z^{16} + (432u^4)z^{17} + (498u^4)z^{18} + (551u^4)z^{19} + (594u^4)z^{20} + (614u^4)z^{21}
\end{aligned}$$

$$\begin{aligned}
& z^{21} + (624 u^4) z^{22} + (601 u^4) z^{23} + (570 u^4) z^{24} + (514 u^4) z^{25} + (453 u^4) z^{26} + \\
& (378 u^4) z^{27} + (312 u^4) z^{28} + (238 u^4) z^{29} + (181 u^4) z^{30} + (128 u^4) z^{31} + (89 u^4) \\
& z^{32} + (56 u^4) z^{33} + (37 u^4) z^{34} + (20 u^4) z^{35} + (12 u^4) z^{36} + (6 u^4) z^{37} + (3 u^4) z^{38} \\
& + (u^4) z^{39} + (u^4) z^{40}
\end{aligned}$$

Table of the number of alkyls according to size and height

Calculations with respect to different heights are much more efficient than the method of exhaustive enumeration. This makes it possible for us to set up the table of the number of alkyls according to size and height in a few minutes:

```

> h[max]:=5;
          hmax := 5
> gfsol:=gfsolve(op(2..3,specs_ltd_height(h[max])),z);
gfsol := {Alkyl_height5(z) = 16 z6 + 63 z8 + 121 z9 + 749 z12 + z + 1344 z13
          + 225 z10 + 415 z11 + 8 z5 + 33 z7 + 39922778 z93 + 552046535 z84
          + 261713408 z87 + 12677964 z96 + 4129 z15 + 20354 z18 + 55706 z20 + 872727 z26
          + 364555 z24 + 7106 z16 + 90628 z21 + 1328545 z27 + 4393287 z30 + 89755449 z39
          + 13204526 z33 + 36095102 z36 + 145729 z22 + 2365 z14 + 66951451 z38
          + 49418998 z37 + 18657905 z34 + 26088244 z35 + 6407683 z31 + 2000536 z28
          + 2980554 z29 + 9246830 z32 + 231801 z23 + 567206 z25 + 12104 z17 + 33883 z19
          + 119063149 z40 + z2 + 2 z3 + z121 + 3994067586 z69 + 2532213546 z75
          + 1323523938 z51 + 2848892771 z57 + z120 + 3993437445 z62 + 2576241555 z56
          + 3116054839 z58 + 4227813312 z64 + 519559381 z46 + 1121537006 z50
          + 1235964517 z80 + 4132169661 z63 + 1030602778 z81 + 109250394 z90 + 27 z117
          + 3469135 z99 + 25584 z108 + 338 z114 + 3353 z111 + 808547 z102 + 417359802 z45
          + 158476 z105 + 1545167948 z52 + 779843029 z48 + 202984042 z42
          + 2038928979 z54 + 3606730433 z60 + 1710204982 z78 + 4277663720 z66
          + 3356383912 z72 + 331715843 z44 + 2249523074 z76 + 2815570112 z74
          + 4134086103 z68 + 3813878148 z70 + 4 z4 + 848181768 z82 + 688882553 z83
          + 436439448 z85 + 340324807 z86 + 198431393 z88 + 148315264 z89
          + 79298004 z91 + 56695196 z92 + 27675367 z94 + 18885015 z95 + 8372800 z97
          + 5435626 z98 + 2174395 z100 + 1338790 z101 + 479339 z103 + 278280 z104
          + 88204 z106 + 48126 z107 + 13348 z109 + 6744 z110 + 1604 z112 + 758 z113
          + 154 z115 + 62 z116 + 10 z118 + 4 z119 + 3371001341 z59 + 3816383212 z61
          + 4276971739 z65 + 4229607116 z67 + 3599155257 z71 + 3092740855 z73
          + 1973755292 z77 + 1463215476 z79 + 156284730 z41 + 260865858 z43
          + 639939517 z47 + 940236752 z49 + 1784589063 z53 + 2304400735 z55,

```

$\text{Alkyl_height}_3(z) =$
 $z + 5z^7 + 4z^9 + 3z^{10} + z^{13} + z^2 + 2z^{11} + z^{12} + 4z^6 + 4z^8 + 2z^3 + 3z^4 + 4z^5,$
 $\text{Alkyl_height}_4(z) = 12z^6 + 31z^8 + 47z^9 + 137z^{12} + z + 184z^{13} + 70z^{10} + 99z^{11}$
 $+ 7z^5 + 20z^7 + 300z^{15} + 498z^{18} + 594z^{20} + 453z^{26} + 570z^{24} + 369z^{16} + 614z^{21}$
 $+ 378z^{27} + 181z^{30} + z^{39} + 56z^{33} + 12z^{36} + 624z^{22} + 239z^{14} + 3z^{38} + 6z^{37}$
 $+ 37z^{34} + 20z^{35} + 128z^{31} + 312z^{28} + 238z^{29} + 89z^{32} + 601z^{23} + 514z^{25}$
 $+ 432z^{17} + 551z^{19} + z^{40} + z^2 + 2z^3 + 4z^4, \text{Alkyl_height}_1(z) = z, \text{Carbon}(z) = z,$
 $\text{Alkyl_height}_2(z) = z + z^2 + z^3 + z^4 \}$

> $\text{BGF} := \text{map}(\text{series}, \text{series}(\text{eval}(\text{subs}(\text{Alkyl_height}[0]=0, \text{gfso}\text{l}, \text{add}(\text{u}^{\text{hh}} * (\text{Alkyl_height}[\text{hh}] - \text{Alkyl_height}[\text{hh}-1])(\text{z}), \text{hh}=1..h[\text{max}]))), \text{z}, \text{infinity}), \text{u}, \text{infinity}) ;$

$BGF := (u)z + (u^2)z^2 + (u^2 + u^3)z^3 + (u^2 + 2u^3 + u^4)z^4 + (4u^3 + 3u^4 + u^5)z^5 +$
 $(4u^3 + 8u^4 + 4u^5)z^6 + (5u^3 + 15u^4 + 13u^5)z^7 + (4u^3 + 27u^4 + 32u^5)z^8 +$
 $(4u^3 + 43u^4 + 74u^5)z^9 + (3u^3 + 67u^4 + 155u^5)z^{10} + (2u^3 + 97u^4 + 316u^5)z^{11}$
 $+ (u^3 + 136u^4 + 612u^5)z^{12} + (u^3 + 183u^4 + 1160u^5)z^{13} + (239u^4 + 2126u^5)$
 $z^{14} + (300u^4 + 3829u^5)z^{15} + (369u^4 + 6737u^5)z^{16} + (432u^4 + 11672u^5)z^{17} +$
 $(498u^4 + 19856u^5)z^{18} + (551u^4 + 33332u^5)z^{19} + (594u^4 + 55112u^5)z^{20} +$
 $(614u^4 + 90014u^5)z^{21} + (624u^4 + 145105u^5)z^{22} + (601u^4 + 231200u^5)z^{23} +$
 $(570u^4 + 363985u^5)z^{24} + (514u^4 + 566692u^5)z^{25} + (453u^4 + 872274u^5)z^{26} +$
 $(378u^4 + 1328167u^5)z^{27} + (312u^4 + 2000224u^5)z^{28} + (238u^4 + 2980316u^5)$
 $z^{29} + (181u^4 + 4393106u^5)z^{30} + (128u^4 + 6407555u^5)z^{31} +$
 $(89u^4 + 9246741u^5)z^{32} + (56u^4 + 13204470u^5)z^{33} + (37u^4 + 18657868u^5)z^{34}$
 $+ (20u^4 + 26088224u^5)z^{35} + (12u^4 + 36095090u^5)z^{36} + (6u^4 + 49418992u^5)$
 $z^{37} + (3u^4 + 66951448u^5)z^{38} + (u^4 + 89755448u^5)z^{39} + (u^4 + 119063148u^5)z^{40}$
 $+ (156284730u^5)z^{41} + (202984042u^5)z^{42} + (260865858u^5)z^{43} +$
 $(331715843u^5)z^{44} + (417359802u^5)z^{45} + (519559381u^5)z^{46} + (639939517u^5)$
 $z^{47} + (779843029u^5)z^{48} + (940236752u^5)z^{49} + (1121537006u^5)z^{50} +$
 $(1323523938u^5)z^{51} + (1545167948u^5)z^{52} + (1784589063u^5)z^{53} +$
 $(2038928979u^5)z^{54} + (2304400735u^5)z^{55} + (2576241555u^5)z^{56} +$
 $(2848892771u^5)z^{57} + (3116054839u^5)z^{58} + (3371001341u^5)z^{59} +$
 $(3606730433u^5)z^{60} + (3816383212u^5)z^{61} + (3993437445u^5)z^{62} +$
 $(4132169661u^5)z^{63} + (4227813312u^5)z^{64} + (4276971739u^5)z^{65} +$
 $(4277663720u^5)z^{66} + (4229607116u^5)z^{67} + (4134086103u^5)z^{68} +$
 $(3994067586u^5)z^{69} + (3813878148u^5)z^{70} + (3599155257u^5)z^{71} +$

$$\begin{aligned}
& (3356383912 u^5) z^{72} + (3092740855 u^5) z^{73} + (2815570112 u^5) z^{74} + \\
& (2532213546 u^5) z^{75} + (2249523074 u^5) z^{76} + (1973755292 u^5) z^{77} + \\
& (1710204982 u^5) z^{78} + (1463215476 u^5) z^{79} + (1235964517 u^5) z^{80} + \\
& (1030602778 u^5) z^{81} + (848181768 u^5) z^{82} + (688882553 u^5) z^{83} + \\
& (552046535 u^5) z^{84} + (436439448 u^5) z^{85} + (340324807 u^5) z^{86} + (261713408 u^5) \\
& z^{87} + (198431393 u^5) z^{88} + (148315264 u^5) z^{89} + (109250394 u^5) z^{90} + \\
& (79298004 u^5) z^{91} + (56695196 u^5) z^{92} + (39922778 u^5) z^{93} + (27675367 u^5) z^{94} \\
& + (18885015 u^5) z^{95} + (12677964 u^5) z^{96} + (8372800 u^5) z^{97} + (5435626 u^5) z^{98} \\
& + (3469135 u^5) z^{99} + (2174395 u^5) z^{100} + (1338790 u^5) z^{101} + (808547 u^5) z^{102} + \\
& (479339 u^5) z^{103} + (278280 u^5) z^{104} + (158476 u^5) z^{105} + (88204 u^5) z^{106} + \\
& (48126 u^5) z^{107} + (25584 u^5) z^{108} + (13348 u^5) z^{109} + (6744 u^5) z^{110} + (3353 u^5) \\
& z^{111} + (1604 u^5) z^{112} + (758 u^5) z^{113} + (338 u^5) z^{114} + (154 u^5) z^{115} + (62 u^5) z^{116} \\
& + (27 u^5) z^{117} + (10 u^5) z^{118} + (4 u^5) z^{119} + (u^5) z^{120} + (u^5) z^{121}
\end{aligned}$$

In the following table, the entry at row r and column c is the number of alkyls of size r and height c :

```
> matrix( [[` `, seq(`height =
` .hh, hh=1..h[max])], seq(`size =
` .ss, seq(coeff(coeff(BGF, z, ss), u, hh), hh=1..h[max])], ss=
1..s[max])]);
```

	<i>height = 1</i>	<i>height = 2</i>	<i>height = 3</i>	<i>height = 4</i>	<i>height = 5</i>
<i>size = 1</i>	1	0	0	0	0
<i>size = 2</i>	0	1	0	0	0
<i>size = 3</i>	0	1	1	0	0
<i>size = 4</i>	0	1	2	1	0
<i>size = 5</i>	0	0	4	3	1
<i>size = 6</i>	0	0	4	8	4
<i>size = 7</i>	0	0	5	15	13
<i>size = 8</i>	0	0	4	27	32
<i>size = 9</i>	0	0	4	43	74
<i>size = 10</i>	0	0	3	67	155
<i>size = 11</i>	0	0	2	97	316
<i>size = 12</i>	0	0	1	136	612
<i>size = 13</i>	0	0	1	183	1160
<i>size = 14</i>	0	0	0	239	2126
<i>size = 15</i>	0	0	0	300	3829
<i>size = 16</i>	0	0	0	369	6737
<i>size = 17</i>	0	0	0	432	11672
<i>size = 18</i>	0	0	0	498	19856
<i>size = 19</i>	0	0	0	551	33332
<i>size = 20</i>	0	0	0	594	55112
<i>size = 21</i>	0	0	0	614	90014
<i>size = 22</i>	0	0	0	624	145105
<i>size = 23</i>	0	0	0	601	231200
<i>size = 24</i>	0	0	0	570	363985
<i>size = 25</i>	0	0	0	514	566692
<i>size = 26</i>	0	0	0	453	872274
<i>size = 27</i>	0	0	0	378	1328167
<i>size = 28</i>	0	0	0	312	2000224
<i>size = 29</i>	0	0	0	238	2980316
<i>size = 30</i>	0	0	0	181	4393106
<i>size = 31</i>	0	0	0	128	6407555
<i>size = 32</i>	0	0	0	89	9246741
<i>size = 33</i>	0	0	0	56	13204470
<i>size = 34</i>	0	0	0	37	18657868
<i>size = 35</i>	0	0	0	20	26088224
<i>size = 36</i>	0	0	0	12	36095090
<i>size = 37</i>	0	0	0	6	49418992
<i>size = 38</i>	0	0	0	3	66951448
<i>size = 39</i>	0	0	0	1	89755448
<i>size = 40</i>	0	0	0	1	119063148
<i>size = 41</i>	0	0	0	0	156284730
<i>size = 42</i>	0	0	0	0	202984042
<i>size = 43</i>	0	0	0	0	260865858
<i>size = 44</i>	0	0	0	0	331715843

[[A (huge) table for $h_{max} = 7$ could be computed in less than 10 minutes.

Disubstituted alkanes, $C_n H_{2n} X Y$

Enumerating disubstituted alkanes $C_n H_{2n} X Y$ is equivalent to enumerating monosubstituted alkyls $C_n H_{2n} X$. The latter can generically be viewed as a carbon atom linked to one monosubstituted alkyl and at least 2 nonsubstituted alkyls. This yields the class equation
 $S1_Alkyl_X = Carbon\ S1_Alkyl_X(E + Alkyl) + Carbon\ X(E + Alkyl + Alkyl^2)$.

```
> gramm_S1_Alkyl:=S1_Alkyl[X]=Union(Prod(Carbon,S1_Alkyl[X],Set(Alkyl,card<=2)),Prod(Prod(Carbon,X),Set(Alkyl,card<=2))),X=Epsilon;
> specs_S1_Alkyl:=[S1_Alkyl[X],{gramm_S1_Alkyl,gramm_Alkyl},unlabelled];
> seq(count(specs_S1_Alkyl,size=i),i=0..50);
0, 1, 2, 5, 12, 31, 80, 210, 555, 1479, 3959, 10652, 28760, 77910, 211624, 576221,
1572210, 4297733, 11767328, 32266801, 88594626, 243544919, 670228623,
1846283937, 5090605118, 14047668068, 38794922293, 107215238057, 296501478704,
820476261295, 2271726458263, 6293333029156, 17443168163416, 48370062636654,
134190690985978, 372435833881578, 1034078866908394, 2872232726571749,
7980695109514561, 22182422656423849, 61676117449283837, 171537091915110029,
477227744594009504, 1328048856698095447, 3696729316849207130,
10292748327630264925, 28664895623718825161, 79849131533514081701,
222477780725979665937, 620005805241494744835, 1728199288005906578667
```

This series appears as the entry **M1418** ("paraffins with n carbon atoms") in the book by N. J. A. Sloane and S. Plouffe [*The Encyclopedia of Integer Sequences*, (1995), Academic Press].

Of course, there are more monosubstituted alkyls than nonsubstituted ones. We give the ratios number of monosubstituted alkyls/number of alkyls for small sizes:

```
> seq([i=evalf(count(specs_S1_Alkyl,size=i)/count(specs_Alkyl,size=i))],i=1..50);
[1 = 1.], [2 = 2.], [3 = 2.500000000], [4 = 3.], [5 = 3.875000000], [6 = 4.705882353],
[7 = 5.384615385], [8 = 6.235955056], [9 = 7.009478673], [10 = 7.808678501],
[11 = 8.604200323], [12 = 9.407916258], [13 = 10.19897892], [14 = 10.99859675],
[15 = 11.79210069], [16 = 12.58714553], [17 = 13.38032304], [18 = 14.17376379],
[19 = 14.96597929], [20 = 15.75825478], [21 = 16.54987862], [22 = 17.34135391],
[23 = 18.13247884], [24 = 18.92344500], [25 = 19.71418351], [26 = 20.50478793],
[27 = 21.29523810], [28 = 22.08557742], [29 = 22.87580605], [30 = 23.66594591],
[31 = 24.45600192], [32 = 25.24598688], [33 = 26.03590630], [34 = 26.82576826],
[35 = 27.61557771], [36 = 28.40534018], [37 = 29.19505975], [38 = 29.98474048],
[39 = 30.77438565], [40 = 31.56399832], [41 = 32.35358114], [42 = 33.14313651],
[43 = 33.93266655], [44 = 34.72217319], [45 = 35.51165815], [46 = 36.30112299],
[47 = 37.09056911], [48 = 37.87999780], [49 = 38.66941024], [50 = 39.45880747]
```

Here is an example of a monosubstituted alkyl with 6 carbon atoms, obtained by the command `combstruct[draw]`.

```
> alk:=draw(specs_S1_Alkyl,size=6);
alk := Prod(Carbon, Prod(Carbon,
    Prod(Prod(Carbon, X), Set(Prod(Carbon, Set(Prod(Carbon, E)))))),
    Set(Prod(Carbon, E))), E)
```

The following procedure rewrites a monosubstituted alkyl into a more readable way.

```
> nice:=proc(alk)
    subs( [C,X]=CX, eval(subs({Epsilon=NULL, Carbon=C, Prod=proc()
        [args] end, Set=proc() args end}, alk)) ) end:
> nice(alk);
[C, [C, [CX, [C, [C]]], [C]]]
```

The following procedures compute the size and height of a given monosubstituted alkyl.

```
> size:=proc(alk) option remember;
    1+convert(map(op,map2(map,size,[op(2..-1,alk)])), '+') end:
size(Carbon):=1:
size(X):=0:
size(Epsilon):=0:
> height:=proc(alk) option remember;
    'if'(nops(alk)=2,1+max(op(map(height,op(2,alk)))),1+max(height(op(2,alk)),op(map(height,op(3,alk))))) end:
height(Carbon):=1:
height(X):=0:
height(Epsilon):=0:
```

Here is a monosubstituted alkyl with 50 carbon atoms, its nice representation and height.

```
> alk:=draw(specs_S1_Alkyl,size=50):
> nice(alk);
[C, [C,
    [C, [C, [C, [CX], [C], [C, [C], [C, [C, [C, [C]]], [C, [C, [C]], [C, [C]]]]]]], [C, [C, [C], [C, [C, [C, [C]]], [C, [C, [C], [C, [C]]]]]]], [C, [C, [C], [C, [C, [C, [C, [C]]], [C, [C, [C], [C, [C]]]]]]], [C, [C, [C, [C, [C, [C, [C]]], [C, [C, [C, [C, [C]]], [C, [C, [C, [C]]]]]]]]], [C, [C, [C, [C, [C, [C, [C]]], [C, [C, [C, [C, [C]]], [C, [C, [C, [C]]]]]]]]]
```

```
> height(alk);
```

11

■ Trisubstituted alkanes, $C_n H_{2n-1} XYZ$

Enumerating trisubstituted alkanes $C_n H_{2n-1} XYZ$ is equivalent to enumerating disubstituted alkyls $C_n H_{2n-1} XY$. In this section, we assume X , Y and Z to be distinct. The grammar is more involved than in the disubstituted case: we have to distinguish several cases, according to which of X and Y go into subtrees, and into which subtrees. The corresponding class equation is $S2_Alkyl_{X,Y} = Carbon\ S2_Alkyl_{X,Y}(E + Alkyl + Alkyl^2)$
 $+ Carbon\ (X + S1_Alkyl_X)\ (Y + S1_Alkyl_Y)\ (E + Alkyl)$.

```
> gramm_S2_Alkyl:=S2_Alkyl[X,Y]=Union(Prod(Carbon,S2_Alkyl[X,Y]
,Set(Alkyl,card<=2)),Prod(Carbon,Union(S1_Alkyl[X],X),Union(S
1_Alkyl[Y],Y),Set(Alkyl,card<=1))):
```

```

> specs_S2_Alkyl:=[S2_Alkyl[X,Y],{gramm_S2_Alkyl,gramm_S1_Alkyl
,op(subs(X=Y,[gramm_S1_Alkyl])),gramm_Alkyl},unlabelled];
> seq(count(specs_S2_Alkyl,size=i),i=0..50);
0, 1, 4, 13, 42, 131, 402, 1218, 3657, 10899, 32298, 95257, 279844, 819390, 2392392,
6967956, 20250974, 58744089, 170118980, 491913999, 1420493862, 4096940530,
11803172152, 33970257473, 97678027311, 280624328431, 805587723862,
2310919999992, 6624670101196, 18978908257258, 54340562045429,
155503251237194, 444766664162993, 1271498880014923, 3633315536811959,
10377791606909654, 29630012602096393, 84565516906270186, 241267111088603729,
688104455854536297, 1961866816555420391, 5591785495039589114,
15933226976278838204, 45387390825197706053, 129255934882453809489,
368005885500527402163, 1047497232155611335438, 2980913638772831767811,
8481029051770110331198, 24124279361703485318308, 68607333785448471672444

```

This series extends the entry **M3466** ("paraffins with n carbon atoms") in the book by N. J. A. Sloane and S. Plouffe [*The Encyclopedia of Integer Sequences*, (1995), Academic Press].

Here is an example of a disubstituted alkyl with 6 carbon atoms, obtained by the command `combstruct[draw]`.

```

> alk:=draw(specs_S2_Alkyl,size=6);
alk := Prod(Carbon, Prod(Carbon, X,
Prod(Carbon, Prod(Prod(Carbon, Y), Set(Prod(Carbon, E))), Set(Prod(Carbon, E))), E
,E)

```

The following procedure rewrites a disubstituted alkyl into a more readable way.

```

> nice:=proc(alk)
subs({[C,X]=CX, [C,Y]=CY, [C,X,Y]=CXY}, eval(subs({Epsilon=NULL,
Carbon=C, Prod=proc() [args] end, Set=proc() args end}, alk)))
end;
> nice(alk);
[C, [C, X, [C, [CY, [C]], [C]]]]

```

The following procedures compute the size and height of a given disubstituted alkyl.

```

> size:=proc(alk) option remember;
`if`(nops(alk)=2,1+convert(map(size,op(2,alk)), '+'),1+convert
(map(size,[op(2..-2,alk)]), '+')+convert(map(size,op(-1,alk)),
'+')) end;
size(Carbon):=1:
size(X):=0:
size(Y):=0:
size(Epsilon):=0:
> height:=proc(alk) option remember;
`if`(nops(alk)=2,1+max(op(map(height,op(2,alk)))),1+max(op(ma
p(height,[op(2..-2,alk)]),op(map(height,op(-1,alk))))) end:
height(Carbon):=1:
height(X):=0:
height(Y):=0:
height(Epsilon):=0:

```

[Here is a disubstituted alkyl with 50 carbon atoms, its nice representation and height.

```
> alk:=draw(specs_S2_Alkyl,size=50):
> nice(alk);
[C,[C,[C,[C,[C,[C,[C,[C,[C,[C,
    [C,[C,[CX],[C,[C,[C],[C]]], [C,[C,[C,[C]]]], [C,[C]], [C,[C],[C]]],
    [C],[C,[C]]]]], [C,[C],[C]]], [C,[C,[C]], [C,[C],[C]]]], [C,[C]]],
    [C,[CY,[C,[C,[C,[C]]]]], [C,[C]]]]
> height(alk);
```

17

■ Trisubstituted alkanes, $C_n H_{2n-1} X_2 Y$

Enumerating trisubstituted alkanes $C_n H_{2n-1} X_2 Y$ is equivalent to enumerating disubstituted alkyls $C_n H_{2n-1} X_2$. In this section, we assume X and Y to be distinct. The class equation is

$$S2\text{-}Alkyl_{X,X} = Carbon \ S2\text{-}Alkyl_{X,X} (E + Alkyl + Alkyl^2)$$

$$+ Carbon (S1\text{-}Alkyl_X^2 + S1\text{-}Alkyl_X X + X^2) (E + Alkyl).$$

```
> gramm_S2b_Alkyl:=S2_Alkyl[X,X]=Union(Prod(Carbon,S2_Alkyl[X,X]
    ,Set(Alkyl,card<=2)),Prod(Carbon,Union(Prod(S1_Alkyl[X],S1_A
    lkyl[X]),Prod(S1_Alkyl[X],X),Prod(X,X)),Set(Alkyl,card<=1))):
```

```
> specs_S2b_Alkyl:=[S2_Alkyl[X,X],{gramm_S2b_Alkyl,gramm_S1_Alk
    yl,gramm_Alkyl},unlabelled]:
> seq(count(specs_S2b_Alkyl,size=i),i=0..50);
```

```
0, 1, 3, 9, 29, 91, 282, 864, 2624, 7905, 23665, 70458, 208813, 616400, 1813354,
5318673, 15559072, 45410239, 132258752, 384493609, 1115907585, 3233782193,
9358293913, 27048189854, 78087437216, 225198736949, 648826102943,
1867669679538, 5371673739750, 15437695238654, 44334467825097,
127235245949015, 364921130790146, 1046003809731397, 2996578878003288,
8580064056382390, 24555042887500969, 70240586991176211, 200837035211971305,
574007829828207160, 1639908132470162451, 4683360193435350915,
13370263334951108655, 38156995934449004558, 108859702271798238723,
310474064975088473530, 885227243624407302887, 2523254862611498946312,
7190343827493770143147, 20484498978131112323407, 58343548311292541719012
```

This series does not appear in the book by N. J. A. Sloane and S. Plouffe [*The Encyclopedia of Integer Sequences*, (1995), Academic Press], nor do its first four differences. Comparing to the entry **M2838** ("tertiary alcohols with n carbon atoms") in this book, we check that there are always more compounds $C_n H_{2n-1} X_2 Y$ than compounds $C_n H_{2n-1} X_3$ (for $X \neq Y$).

Here is an example of a disubstituted alkyl with 6 carbon atoms, obtained by the command `combstruct[draw]`.

```
> alk:=draw(specs_S2b_Alkyl,size=6);
alk := Prod(Carbon, Prod(Prod(Carbon, Prod(Prod(Carbon, X), Set(Prod(Carbon, E)))),
```

```
| Set(Prod(Carbon, E), Prod(Carbon, E))), X), E)
```

| The following procedure rewrites a disubstituted alkyl into a more readable way.

```
> nice:=proc(alk)
   subs({ [C, X]=CX, [C, X, X]=CX[2] }, eval(subs({Epsilon=NULL, Carbon=
   C, Prod=proc() [args] end, Set=proc() args end}, alk))) end;
> nice(alk);
```

```
[C, [[C, [CX, [C]], [C], [C]], X]]
```

| Here are the 9 disubstituted compounds $C_n H_{2n-1} X_2 Y$ for $n = 3$:

```
> map(nice, allstructs(specs_S2b_Alkyl, size=3));
[[C, [[CX], X], [C]], [C, [C, [X, X]], [C]], [C, [C, [X, X], [C]]], [C, [[CX, [C]], X]],
 [C, [[C, [CX]], X]], [C, [C, [C, [X, X]]]], [C, [C, [[CX], X]]], [C, [X, X], [C, [C]]],
 [C, [[CX], [CX]]]]
```

Conclusion: multiply substituted alkyls

In the previous sections, we have enumerated the substituted compounds $C_n H_{2n+1} X$, $C_n H_{2n} XY$, $C_n H_{2n-1} XYZ$ and $C_n H_{2n-1} X_2 Y$. We could in principle enumerate the class S_{p_1, \dots, p_t} of compounds obtained after substituting p_1 hydrogen atoms by $X^{(1)}$ atoms, p_2 hydrogen atoms by $X^{(2)}$ atoms, ..., p_t hydrogen atoms by $X^{(t)}$ atoms, and one hydrogen atom by Y (so as to plant the trees). Doing so would require to define the class S_{q_1, \dots, q_t} for each $q_1 \leq p_1$, ..., $q_t \leq p_t$, and for each $q_1 \leq p_1$, ..., $q_t \leq p_t$, to write a recursion involving partitions into 4 parts of the multiset $\{X^{(1)} (q_1 \text{ times}), \dots, X^{(t)} (q_t \text{ times})\}$. When the q_i 's are given, those partitions can be computed by a call to [combstruct\[allstructs\]](#). It follows that we would describe and generate the grammar for multiply substituted alkyls in terms of the grammar for partitions into 4 parts!