

Chern Classes of Tautological Sheaves on Hilbert Schemes of Points on Surfaces

Manfred Lehn

Introduction

Hilbert schemes $X^{[n]}$ of n -tuples of points on a complex projective manifold X are natural compactifications of the configuration spaces of unordered distinct n -tuples of points on X . Their geometry is determined by the geometry of X itself and the geometry of the ‘punctual’ Hilbert schemes of all zero-dimensional subschemes in affine space that are supported at the origin. Thus one is naturally led to the following problem:

Determine explicitly the geometric or topological invariants of the Hilbert schemes $X^{[n]}$ such as the Betti numbers, the Hodge numbers, the Chern numbers, the cohomology ring, from the corresponding data of the manifold X itself.

This problem is most attractive when X is a surface, since then the Hilbert schemes are themselves irreducible projective manifolds, by a result of Fogarty [9], whereas for higher dimensional varieties the Hilbert schemes are in general neither irreducible nor smooth nor pure of expected dimension.

In the surface case, the answer to the problem above for the Betti numbers was first given by Göttsche in [11]. The answer turns out to be particularly beautiful (cf. Theorem 2.1 below). The problem for the Hodge numbers was solved by Sörgel and Göttsche [12]. For a different approach to both results see [3]. The answer for the Chern classes will be implicitly given in a forthcoming paper by Ellingsrud, Göttsche and the author [4].

The question for the ring structure of the cohomology is more difficult. In general, $X^{[2]}$ is the quotient of the blow-up of $X \times X$ along the diagonal by the canonical involution that exchanges the factors. Thus the case of interest is $H^*(X^{[n]})$, $n \geq 3$. The ring structure was found for $(\mathbb{P}^2)^{[3]}$ by Ellingsrud and Strømme [5], and for $X^{[3]}$, X smooth projective of arbitrary dimension, by Fantechi and Göttsche [8]. In another direction, Ellingsrud and Strømme [6] gave generators for $H^*((\mathbb{P}^2)^{[n]}, \mathbb{Z})$, n arbitrary, and an implicit description of the relations.

Vafa and Witten [27] remarked that Göttsche’s Formula for the Betti numbers is identical with the Poincaré series of a Fock space modelled on the cohomology of X . Nakajima [21] succeeded in giving a geometric construction of such a Fock space structure on the cohomology of the Hilbert schemes, leading to a natural ‘explanation’ of Göttsche’s result. Similar results have been announced by Grojnowski [13].

Following the presentation of Grojnowski, this can be made more precise as follows: sending a pair (ξ', ξ'') of subschemes of length n' and n'' , respectively, and of disjoint support to their union $\xi' \cup \xi''$ defines a rational map

$$m : X^{[n']} \times X^{[n'']} \dashrightarrow X^{[n'+n'']}.$$

This map induces linear maps on the rational cohomology

$$m_{n', n''} : H^*(X^{[n']}; \mathbb{Q}) \otimes H^*(X^{[n'']}; \mathbb{Q}) \longrightarrow H^*(X^{[n'+n'']}; \mathbb{Q})$$

and

$$m^{n', n''} : H^*(X^{[n'+n'']}; \mathbb{Q}) \longrightarrow H^*(X^{[n']}; \mathbb{Q}) \otimes H^*(X^{[n'']}; \mathbb{Q}).$$

If we let $\mathbb{H} := \bigoplus_n H^*(X^{[n]}; \mathbb{Q})$, then these maps define a multiplication and a comultiplication

$$m_* : \mathbb{H} \otimes \mathbb{H} \longrightarrow \mathbb{H}, \quad m^* : \mathbb{H} \longrightarrow \mathbb{H} \otimes \mathbb{H},$$

which make \mathbb{H} a commutative and cocommutative bigraded Hopf algebra. The result of Nakajima and Grojnowski says that this Hopf algebra is isomorphic to the graded symmetric algebra of the vector space $H^*(X; \mathbb{Q}) \otimes t\mathbb{Q}[t]$.

More explicitly, Nakajima constructed linear maps¹

$$q_n : H^*(X; \mathbb{Q}) \longrightarrow \text{End}_{\mathbb{Q}}(\mathbb{H}), \quad n \in \mathbb{Z},$$

and proved that they satisfy the ‘oscillator’ or ‘Heisenberg’ relations

$$[q_n(\alpha), q_m(\beta)] = (-1)^n \cdot n \cdot \delta_{n+m} \cdot \int_X \alpha \beta \cdot \text{id}_{\mathbb{H}}.$$

Here the commutator is to be taken in a graded sense.

The multiplication and the comultiplication of \mathbb{H} are not obviously related to the quite different ring structure of \mathbb{H} , which is given by the usual cup product on each direct summand $H^*(X^{[n]}; \mathbb{Q})$. (Strictly speaking, \mathbb{H} contains a countable number of idempotents $1_{X^{[n]}} \in H^0(X^{[n]}; \mathbb{Q})$ but not a unit unless we pass to some completion).

This paper attempts to relate the Hopf algebra structure and the cup product structure. More precisely:

Let F be locally free sheaf of rank r on X . Attaching to a point $\xi \in X^{[n]}$, i.e. a zero-dimensional subscheme $\xi \subset X$, the \mathbb{C} -vector space $F \otimes \mathcal{O}_{\xi}$ defines a locally free sheaf $F^{[n]}$ of rank rn on $X^{[n]}$. The Chern classes of all sheaves on $X^{[n]}$ of this type generate a subalgebra $\mathcal{A} \subset \mathbb{H}$. We will describe a purely algebraic algorithm to determine the action of \mathcal{A} on \mathbb{H} in terms of the \mathbb{Q} -basis of \mathbb{H} provided by Nakajima’s results. We collect the Chern classes of all sheaves $F^{[n]}$ for a given sheaf F into operators

$$\text{ch}(F) : \mathbb{H} \rightarrow \mathbb{H}, \quad \text{c}(F) : \mathbb{H} \rightarrow \mathbb{H}$$

and geometrically compute the commutators of these operators with the ‘standard operators’ defined by Nakajima.

¹Our presentation differs in notations and conventions slightly from Nakajima’s.

A central rôle is played by the operator $\mathfrak{d} := \mathfrak{q}_1(\mathcal{O}_X)$, which — up to a factor $(-1/2)$ — can also be interpreted as the intersection with the ‘boundaries’ of the Hilbert schemes, i.e. the divisors $\partial X^{[n]} \subset X^{[n]}$ of all tuples ξ which have a multiple point somewhere. The derivative of any operator $\mathfrak{f} \in \text{End}(\mathbb{H})$ is defined by $\mathfrak{f}' := [\mathfrak{d}, \mathfrak{f}]$. Our main technical result then says that

$$\mathfrak{q}'_n(\alpha) = \frac{n}{2} \sum_{\nu} \mathfrak{q}_{\nu} \mathfrak{q}_{n-\nu} \delta(\alpha) + n \frac{|n| - 1}{2} \mathfrak{q}_n(K\alpha), \quad (1)$$

where $\delta : H^*(X; \mathbb{Q}) \rightarrow H^*(X; \mathbb{Q}) \otimes H^*(X; \mathbb{Q})$ is the map induced by the diagonal embedding and K is the canonical class of X . An immediate algebraic consequence of this relation is

$$[\mathfrak{q}'_n(\alpha), \mathfrak{q}_m(\beta)] = -nm \cdot \mathfrak{q}_{n+m}(\alpha\beta) \quad (2)$$

for $n, m > 0$. By induction one concludes that the operators \mathfrak{q} and \mathfrak{d} suffice to generate all \mathfrak{q}_n , $n \geq 1$.

The commutator of the Chern character operator $\text{ch}(F)$ with the standard operator \mathfrak{q}_1 can be expressed in terms of higher derivatives of \mathfrak{q} :

$$[\text{ch}(F), \mathfrak{q}_1(\alpha)] = \sum_{n \geq 0} \frac{1}{n!} \mathfrak{q}_1^{(n)}(\text{ch}(F)\alpha). \quad (3)$$

Equations (1), (2) and (3) together give a complete description of the action of \mathcal{A} on \mathbb{H} . Here are two applications:

1. We give a general algebraic solution to Donaldson’s question for the integral N_n of the top Segre class of the bundles $L^{[n]}$ associated to a line bundle L for any n and explicitly compute N_n for $n \leq 7$.

2. We prove the following formula conjectured by Göttsche: If L is a line bundle on X then

$$\sum_{n \geq 0} c(L^{[n]})z^n = \exp \left(\sum_{m \geq 1} \frac{(-1)^{m-1}}{m} \mathfrak{q}_m(c(L))z^m \right).$$

This paper is organised as follows: In Section 1 we recall the basic geometric notions used in the later parts. Section 2 provides an introduction to Nakajima’s results. Section 3 contains the core of this paper: we first define Virasoro operators \mathfrak{L}_n in analogy to the standard construction and show how these arise geometrically. We then introduce the operator \mathfrak{d} and compute the derivative of \mathfrak{q}_n . Finally, in Section 4 we apply these results to compute the action of the Chern classes of tautological bundles.

Discussions with A. King were important to me in clarifying and understanding the picture that Nakajima draws in his very inspiring article. I am very grateful to G. Ellingsrud for all the things I learned from his talks and conversations with him

about Hilbert schemes. To some extent the results in this article are a reflection on an induction method entirely due to him. Most of the research for this paper was carried out during my stay at the SFB 343 of the University of Bielefeld. On various occasions I was allowed to lecture on Hilbert schemes and their cohomology in the seminar of the algebraic geometry group in Bielefeld: it is a pleasure to thank S. Bauer, R. Brussee and T. Zink for their willingness to listen attentively and critically even to not yet fully correct preliminary results. I owe special thanks to S. Bauer for his continuous encouragement, interest and support.

Bielefeld, 8 October, 1997.

Manfred Lehn

Contents

1 Preliminaries	5
1.1 Symmetric products	5
1.2 Hilbert schemes and Hilbert-Chow morphism	6
1.3 Hilbert schemes of smooth surfaces	7
1.4 Incidence schemes	8
2 The structure of the cohomology	11
2.1 Correspondences	12
2.2 Nakajima's Main Theorem	13
3 The boundary operator	15
3.1 Virasoro generators	15
3.2 The boundary of the Hilbert scheme	21
3.3 The derivative of q_n	24
3.4 The vertex operator, completion of the proof	31
4 Towards the ring structure of \mathbb{H}	34
4.1 Tautological sheaves	34
4.2 The line bundle case	36
4.3 Top Segre classes	40
References	44

1 Preliminaries

In this section we introduce the basic notations that will be used throughout the paper and collect a number of results from the literature, mostly without proof.

1.1 Symmetric products

Let Y be a quasi-projective scheme over \mathbb{C} . The symmetric group \mathfrak{S}_n acts on Y^n by permutation of the factors, and there exists a geometric quotient $\pi : Y^n \rightarrow S^n Y$ for this action. $S^n Y$ is again quasi-projective, and if Y is irreducible (reduced, integral or normal) then the same is true for $S^n Y$. Moreover, this construction is functorial: any morphism $f : Y' \rightarrow Y$ induces a morphism $S^n f : S^n Y' \rightarrow S^n Y$.

It follows from the theorem on elementary symmetric functions that $S^n \mathbb{A}^1 \cong \mathbb{A}^n$. Consequently, the symmetric products of smooth curves are again smooth. On the other hand, if Y is a smooth variety of dimension greater than one, then $S^n Y$ is singular for $n > 1$.

By a result of Grothendieck [15], the natural map

$$\pi^* : H^*(S^n Y; \mathbb{Q}) \longrightarrow H^*(Y^n; \mathbb{Q}) \cong H^*(Y; \mathbb{Q})^{\otimes n}$$

is an isomorphism onto the subring of invariant elements under the action of \mathfrak{S}_n . From this Macdonald computed the following formula for the Betti numbers of $S^n Y$ by a purely algebraic argument:

Theorem 1.1 (Macdonald [20]) — *The Betti numbers of the symmetric products are given by the formula*

$$\sum_{n \geq 0} \sum_{i \geq 0} b_i(S^n Y) t^i q^n = \prod_{i=0}^{2 \dim(Y)} (1 - (-1)^i t^i q)^{-(-1)^i b_i(Y)}.$$

□

There is another property of the symmetric product, which is important for the definition of the Hilbert-Chow morphism. Consider the following set-valued contravariant functor $\mathcal{M}_n(Y)$ on the category of locally Noetherian \mathbb{C} -schemes:

Let S be a \mathbb{C} -scheme, and let $p : S \times Y \rightarrow S$ be the projection. Then $\mathcal{M}_n(Y)(S)$ is the set of all isomorphism classes of coherent sheaves F on $S \times Y$, where F is S -flat, $p : \text{Supp}(F) \rightarrow S$ is a finite map, and $p_* F$ is locally free of rank n . If $f : S' \rightarrow S$ is a \mathbb{C} -morphism, then $\mathcal{M}_n(Y)(f) : \mathcal{M}(Y)(S) \rightarrow \mathcal{M}_n(Y)(S')$ maps the class of F to $f_Y^* F$. Here and in the following we write f_Y instead of $f \times \text{id}_Y$.

Grothendieck [14] asserted that there is a natural transformation $\mathcal{M}_n(Y) \rightarrow S^n Y$ sending any zero-dimensional sheaf F to its weighted support. This means that for any $[F] \in \mathcal{M}_n(Y)(S)$ there is a classifying morphism $\Phi_F : S \rightarrow S^n Y$ such that $\Phi_F(s) = \sum_{y \in Y} \ell(F_{s,y}) \cdot y$ for all $s \in S$, where for any coherent sheaf \mathcal{G} we let $\ell(\mathcal{G}_y)$

denote the length of the stalk \mathcal{G}_y as a module over $\mathcal{O}_{Y,y}$. Moreover, $\Phi_{f_Y^*F} = \Phi_F \circ f$ for any $f : S' \rightarrow S$.

This was first proved by Iversen [18] using the technique of linear determinants. In fact, if Y is normal then $S^n Y$ corepresents the functor $\mathcal{M}_n(Y)$ (cf. [16, Ex. 4.3.6]).

1.2 Hilbert schemes and Hilbert-Chow morphism

Throughout this paper, the term ‘Hilbert scheme’ will always refer to Hilbert schemes of zero-dimensional subschemes.

Let Y be a quasi-projective scheme over \mathbb{C} . The Hilbert functor is the following set-valued functor on the category of locally Noetherian \mathbb{C} -schemes:

Let $\text{Hilb}(Y, n)(S)$ be the set of all closed subschemes $Z \subset S \times Y$ such that the projection $p : Z \rightarrow S$ is flat and finite of degree n . If $f : S' \rightarrow S$ is a \mathbb{C} -morphism, the induced map is given by pull-back: $Z \mapsto f_Y^{-1}(Z) = S' \times_S Z$.

Grothendieck [14] showed that $\text{Hilb}(Y, n)$ is represented by a quasi-projective scheme $Y^{[n]}$. If Y is projective, $Y^{[n]}$ is projective as well.

‘Functoriality’ in Y is limited to a few cases: if $f : Y' \rightarrow Y$ is an (open, closed) immersion, then there is a natural (open, closed) immersion

$$f^{[n]} : (Y')^{[n]} \rightarrow Y^{[n]},$$

defined by taking the image of subschemes under f . Moreover, suppose that $f : Y' \rightarrow Y$ is an étale (surjective) morphism. Let $U \subset (Y')^{[n]}$ denote the open subset of all subschemes $\xi \subset Y'$ such that the set-theoretic support of ξ is mapped injectively to Y . Then taking images under f defines an étale (surjective) morphism $U \rightarrow Y^{[n]}$.

For small values of n there are explicit descriptions of $Y^{[n]}$: Clearly, $Y^{[0]}$ is a reduced point, $Y^{[1]} \cong Y$, and $Y^{[2]}$ is the quotient for the \mathfrak{S}_2 -action on the blow-up of $Y \times Y$ along the diagonal. Proceeding by induction, it is not difficult to see that all Hilbert schemes $Y^{[n]}$ are connected if Y is connected.

Observe that there is a natural transformation of functors

$$\text{Hilb}(Y, n) \longrightarrow \mathcal{M}_n(Y)$$

which sends a subscheme $Z \subset S \times Y$ to its structure sheaf $\mathcal{O}_Z \in \text{Coh}(S \times Y)$. As $\text{Hilb}(Y, n)$ is represented by $Y^{[n]}$, this transformation induces a morphism of schemes

$$\rho : Y^{[n]} \rightarrow S^n Y,$$

the *Hilbert-Chow* morphism. On a point $[v] \in Y^{[n]}$, i.e. a subscheme $v \subset Y$, this morphism is given by

$$\rho([v]) = \sum_{y \in Y} \ell(\mathcal{O}_{v,y}) \cdot y.$$

For example, if C is a smooth curve, then $\rho : C^{[n]} \rightarrow S^n C$ is an isomorphism.

1.3 Hilbert schemes of smooth surfaces

From now on, let X denote a smooth irreducible projective surface. The basic geometry of the Hilbert schemes of points on surfaces is governed by two theorems due to Fogarty [9] and Briançon [1].

Theorem 1.2 (Fogarty) — $X^{[n]}$ is a $2n$ -dimensional smooth irreducible projective variety.

Here is a short sketch of the proof: projectivity is due to Grothendieck. He also showed that the Zariski tangent space of $X^{[n]}$ at a point ξ is canonically isomorphic to $\text{Hom}(\mathcal{I}_\xi, \mathcal{O}_\xi)$. Since we already know that $X^{[n]}$ is connected, it therefore suffices to show that $\text{hom}(\mathcal{I}_\xi, \mathcal{O}_\xi) = 2n$ for all $\xi \in X^{[n]}$. This can be done using Serre duality and the Hirzebruch-Riemann-Roch Theorem applied to the groups $\text{Ext}^i(\mathcal{O}_\xi, \mathcal{O}_\xi)$. \square

Remark 1.3 — We already mentioned that $C^{[n]}$ is smooth for smooth curves. Computing the dimension of the tangent space one can show that $Y^{[3]}$ is smooth for a smooth variety Y of any dimension. On the other hand, $Y^{[n]}$ is singular if $\dim(Y) > 2$ and $n > 3$.

Fix a point $p \in X$ and let $X_p^{[n]} \subset X^{[n]}$ denote the closed subset of all subschemes $\xi \subset X$ with $\text{Supp}(\xi) = \{p\}$ (with the reduced induced subscheme structure). This is indeed a closed subset, as it is the fibre $\rho^{-1}(np)$ of the Hilbert-Chow morphism over the point $np \in S^n X$.

Let $(\mathcal{O}, \mathfrak{m})$ denote the local ring of X at p . Since any point $\xi \in X_p^{[n]}$ may be considered as a subscheme of $\text{Spec}(\mathcal{O}/\mathfrak{m}^n)$, and since $\mathcal{O}/\mathfrak{m}^n \cong \mathbb{C}[x, y]/(x, y)^n$, all schemes $X_p^{[n]}$ — for varying X and p — are (non-canonically) isomorphic. Clearly, $X_p^{[1]} = \{p\}$ and $X_p^{[2]} = \mathbb{P}(T_p X^\vee)$, moreover it is not too difficult to see that $X_p^{[3]}$ is isomorphic to the projective cone over the twisted cubic $\mathcal{C} \subset \mathbb{P}^3$, the vertex of the cone corresponding to the subscheme $\text{Spec}(\mathcal{O}/\mathfrak{m}^2)$. It is not accidental that in these examples the dimension of $X_p^{[n]}$ increases by one in each step:

Theorem 1.4 (Briançon) — For all $n \geq 1$, $X_p^{[n]}$ is an irreducible variety of dimension $n - 1$. \square

For a proof see [1]. A new proof with a more geometric and conceptual argument was recently given by Ellingsrud and Strømme [7].

Briançon's Theorem emphasises the importance of curvilinear schemes: recall that a zero-dimensional subscheme $\xi \subset X$ is called *curvilinear* at $x \in X$, if ξ_x is contained in some smooth curve $C \subset X$. Equivalently, one might say that $\mathcal{O}_{\xi, x}$ is isomorphic to the \mathbb{C} -algebra $\mathbb{C}[z]/(z^\ell)$, where $\ell = \ell(\xi_x)$. Hence ξ is curvilinear at x if ξ_x is either empty, a reduced point, or if $\dim T_x \xi = 1$. From this criterion it is clear, that in any flat family of zero-dimensional subschemes the points in the base space which correspond to curvilinear subschemes form an open subset.

In particular, we may consider the open subset $X_{p,curv}^{[n]} \subset X_p^{[n]}$. This set has a very nice structure:

Lemma 1.5 — *If $n \geq 2$, then the morphism*

$$t : X_{p,curv}^{[n]} \longrightarrow \mathbb{P}(T_p X^\vee), [\xi] \mapsto [T_p \xi]$$

is a bundle morphism with affine fibres \mathbb{A}^{n-2} . In particular, $X_{p,curv}^{[n]}$ is an irreducible smooth variety of dimension $n - 1$.

Proof. Let $x, y \in \mathcal{O}_{X,p}$ be local coordinates and consider the open subset $U = \{(y + \alpha_1 x) \mid \alpha_1 \in \mathbb{C}\} \subset \mathbb{P}(T_p X^\vee)$. Then there is an isomorphism $\mathbb{A}^{n-1} \rightarrow t^{-1}(U)$ sending the $(n - 1)$ -tuple $(\alpha_1, \dots, \alpha_{n-1})$ to the subsheaf corresponding to the ideal $(y + \alpha_1 x + \dots + \alpha_{n-1} x^{n-1}) + \mathcal{I}_p^n$. \square

As a consequence of this lemma we see that Briançon's Theorem is equivalent to saying that $X_{p,curv}^{[n]}$ is dense in $X_p^{[n]}$. This is a very important information: curvilinear subschemes are far easier to handle than any of the others. They contain only one subscheme for any given smaller length, any small deformation of a curvilinear subscheme is again locally curvilinear etc.

Generalising the definition of $X_p^{[n]}$ slightly, let $\Delta \subset S^m X$ denote the diagonal, and let $X_0^{[n]} := \rho^{-1}(\Delta)$, endowed with the reduced induced subscheme structure. Thus $X_0^{[n]}$ consists of all subschemes $\xi \subset X$ of length n which are supported at *some* point in X . The fibres of the surjective morphism $\rho : X^{[n]} \rightarrow X$ are the schemes $X_p^{[n]}$ considered above. In fact, a choice of regular parameters near a point p leads to a trivialisation of the morphism $\rho : X^{[n]} \rightarrow X$ near p , i.e. ρ is a fibre bundle for the Zariski topology.

As an immediate consequence of Briançon's Theorem we get

Corollary 1.6 — $X_0^{[n]}$ is an irreducible variety of dimension $n + 1$. \square

Note that $X_p^{[n]}$ and $X_0^{[n]}$ have complementary dimensions as subvarieties in $X^{[n]}$. Their homological intersection is therefore zero-dimensional. However, the inclusion $X_p^{[n]} \subset X_0^{[n]}$ complicates the computation of the intersection product. The following result was obtained by Ellingsrud and Strømme [7] by an inductive geometric argument:

Theorem 1.7 (Ellingsrud, Strømme) — $\deg([X_p^{[n]}] \cdot [X_0^{[n]}]) = (-1)^{n-1} \cdot n$. \square

1.4 Incidence schemes

Since $X^{[n]}$ represents the functor $\text{Hilb}^n(X)$, there is a *universal family* of subschemes

$$\Xi_n \subset X^{[n]} \times X.$$

Again, for small values of n there are explicit descriptions: Ξ_0 is empty, Ξ_1 is the diagonal in $X \times X$, and Ξ_2 is the blow-up $\text{Bl}_\Delta(X \times X)$ of the diagonal in $X \times X$. The identification is given by the quotient map $\text{Bl}_\Delta(X \times X) \rightarrow X^{[2]} = \text{Bl}_\Delta(X \times X)/\mathcal{G}_2$ and any of the two projections $\text{Bl}_\Delta(X \times X) \rightarrow X$.

Assume that $n' > n > 0$. Then there is a uniquely determined closed subscheme $X^{[n',n]} \subset X^{[n']} \times X^{[n]}$ with the property that any morphism

$$f = (f_1, f_2) : T \rightarrow X^{[n']} \times X^{[n]}$$

factors through $X^{[n',n]}$ if and only if $f_{2,X}^{-1}(\Xi_n) \subset f_{1,X}^{-1}(\Xi_{n'})$. Closed points in $X^{[n',n]}$ correspond to pairs (ξ', ξ) of subschemes with $\xi \subset \xi'$. Let

$$X^{[n']} \xleftarrow{p_1} X^{[n',n]} \xrightarrow{p_2} X^{[n]}$$

denote the two projections. Then $X^{[n',n]}$ parametrises two flat families

$$p_{2,X}^{-1}(\Xi_n) \subset p_{1,X}^{-1}(\Xi_{n'}).$$

Consider the corresponding exact sequence

$$0 \rightarrow \mathcal{I}_{n',n} \rightarrow p_{1,X}^* \mathcal{O}_{\Xi_{n'}} \rightarrow p_{2,X}^* \mathcal{O}_{\Xi_n} \rightarrow 0. \quad (4)$$

The ideal sheaf $\mathcal{I}_{n',n}$ is a coherent sheaf on $X^{[n',n]} \times X$ which is flat over $X^{[n',n]}$ and fibrewise zero-dimensional of length $n' - n$. It therefore induces a classifying morphism to the symmetric product, analogously to the Hilbert-Chow morphism, which we will also denote by

$$\rho : X^{[n',n]} \rightarrow S^{n'-n} X.$$

As before let $X_0^{[n',n]} := \rho^{-1}(\Delta)$, where $\Delta \subset S^{n'-n} X$ is the small diagonal. A point in $X_0^{[n',n]}$ is a triple (ξ', x, ξ) with $\xi \subset \xi'$ and $\text{Supp}(\mathcal{I}_{\xi/\xi'}) = \{x\}$.

We may decompose $X_0^{[n',n]}$ into locally closed subsets Z_ℓ , $\ell \geq 0$, with

$$Z_\ell := \{(\xi', x, \xi) \mid \ell(\xi_x) = \ell\}.$$

Lemma 1.8 — Z_0 and Z_1 are irreducible of dimension $n + n' + 1$ and $n + n'$, respectively, and $\dim(Z_\ell) < n + n'$ for all $\ell > 1$. Moreover, Z_1 is contained in the closure of Z_0 .

Proof. If $\ell = 0$ or 1 , the map $(\xi', x, \xi) \mapsto (\xi - \xi_x, \xi'_x)$ is an open immersion

$$Z_\ell \longrightarrow X^{[n-\ell]} \times X_0^{[n'-n+\ell]}.$$

It follows from Briançon's Theorem that Z_ℓ is irreducible and

$$\dim(Z_\ell) = 2(n - \ell) + (n' - n + \ell + 1) = n + n' + 1 - \ell.$$

For $\ell \geq 2$ consider the embedding

$$Z_\ell \longrightarrow X^{[n-\ell]} \times (X_0^{[\ell]} \times_X X_0^{[n'-n+\ell]}), \quad (\xi', x, \xi) \mapsto (\xi - \xi_x, \xi_x, \xi'_x).$$

In fact, the image of Z_ℓ is contained in a *proper* closed subset of the target variety: For *either* ξ_x^ℓ is curvilinear, in which case there is only a unique subscheme $\xi \subset \xi_x^\ell$ of length ℓ , *or* ξ_x^ℓ is not curvilinear and therefore contained in a proper closed subset of $X_0^{[n'-n+\ell]}$. Now, the variety on the right hand side has dimension

$$2(n - \ell) + (\ell + 1) + (n' - n + \ell + 1) - 2 = n + n'.$$

Finally, a general point in Z_1 is of the form $(\zeta \cup \eta, x, \zeta \cup \{x\})$ where η is a curvilinear subscheme supported at x and disjoint from ζ . Now it is easy to deform η to a subscheme $\{x\} \cup \eta'$ with η' supported at a point $x' \neq x$. Hence a general point of Z_1 deforms into Z_0 . \square

Definition 1.9 — For any pair of nonnegative integers define subvarieties

$$E^{[n',n]}, Q^{[n',n]} \subset X^{[n']} \times X \times X^{[n]}$$

as follows: if $n' > n > 0$ let $Q^{[n',n]}$ and $E^{[n',n]}$ be the closure of Z_0 and Z_1 , respectively. Moreover, $Q^{[n',0]} := X_0^{[n']}$, $E^{[n',0]} := \emptyset$ and $Q^{[n,n]} := \emptyset$, whereas $E^{[n,n]} := \{(\xi, x, \xi) | x \in \xi\} \cong \Xi_n$. On the other hand, if $n \geq n'$, let $Q^{[n',n]} = T(Q^{[n,n']})$ and $E^{[n',n]} = T(E^{[n,n']})$ under the twist

$$T : X^{[n]} \times X \times X^{[n']} \rightarrow X^{[n']} \times X \times X^{[n]}.$$

By construction $Q^{[n,n']}$ and $E^{[n,n']}$ are empty or irreducible varieties of dimension $n + n' + 1$ and $n + n'$, respectively.

Let us return to the particular case $n' - n = 1$, the most basic of all incidence situations: consider the projectivisation $\sigma : \mathbb{P}(\mathcal{I}_{\Xi_n}) \rightarrow X^{[n]} \times X$. It is an easy exercise to see that there is a natural isomorphism $\mathbb{P}(\mathcal{I}_{\Xi_n}) \cong X^{[n+1,n]}$ such that the diagram

$$\begin{array}{ccc} \mathbb{P}(\mathcal{I}_{\Xi_n}) & \xrightarrow{\cong} & X^{[n+1,n]} \\ \sigma \searrow & & (p_2, \rho) \swarrow \\ & & X^{[n]} \times X \end{array}$$

commutes.

Theorem 1.10 (Ellingsrud, Strømme [7]) — *The incidence scheme $X^{[n+1,n]}$ is an irreducible variety.*

An immediate corollary is the following: there is a natural closed immersion $\text{Bl}_{\Xi_n}(X^{[n]} \times X) \rightarrow \mathbb{P}(\mathcal{I}_{\Xi_n})$; since both are irreducible varieties, this must be an isomorphism. The exceptional divisor E is precisely the variety $E^{[n+1,n]}$ defined above. Hence in this situation we may write the sequence (4) as

$$0 \rightarrow (\text{id}, \rho)_* \mathcal{O}_{X^{[n+1,n]}}(-E) \rightarrow p_{1,X}^* \mathcal{O}_{\Xi_{n+1}} \rightarrow p_{2,X}^* \mathcal{O}_{\Xi_n} \rightarrow 0. \quad (5)$$

In fact, the incidence scheme is smooth. This has independently be proved by Ellingsrud, Tikhomirov and Cheah. The proofs are unpublished.

2 The structure of the cohomology

As before, let X be a smooth irreducible projective surface. By Fogarty's Theorem the Hilbert schemes $X^{[n]}$ are projective manifolds of real dimension $4n$. The motivating problem in this study is to understand the cohomology rings $H^*(X^{[n]})$ in terms of the cohomology ring $H^*(X)$.

As far as the vector space structure of the cohomology is concerned, i.e. if we only ask for the dimensions of the graded pieces of the cohomology, this problem was solved by Göttsche [11]. The answer is given by the following beautiful formula for the Betti numbers.

Theorem 2.1 (Göttsche) — *The Betti numbers $b_i(X^{[n]})$ are determined by the Betti numbers $b_j(X)$. More precisely, the following formula holds:*

$$\sum_{n \geq 0} \sum_{i \geq 0} b_i(X^{[n]}) t^i q^n = \prod_{m > 0} \prod_{j \geq 0} (1 - (-1)^j t^{2m-2+j} q^m)^{-(-1)^j b_j(X)}$$

Göttsche's original proof uses the Weil Conjectures [11]. For a different approach see [3].

Among other things one learns from this formula that it is a good idea to consider all Hilbert schemes simultaneously. This will become even more striking through Nakajima's method which we will review in the next sections. As a preparation we collect a few definitions:

Definition 2.2 — Let $\mathbb{H} := \bigoplus_{n,i \geq 0} \mathbb{H}^{n,i}$ denote the double graded vector space with components $\mathbb{H}^{n,i} = H^i(X^{[n]}; \mathbb{Q})$. Since $X^{[0]}$ is a point, $\mathbb{H}^{0,0} = \mathbb{Q}$. The unit in $H^0(X^{[0]}; \mathbb{Q})$ is called the 'vacuum vector' and denoted by $\mathbf{1}$.

A linear map $f : \mathbb{H} \rightarrow \mathbb{H}$ is homogeneous of bidegree (ν, ι) if $f(\mathbb{H}^{n,i}) \subset \mathbb{H}^{n+\nu, i+\iota}$ for all n and i . If $f, f' \in \text{End}(\mathbb{H})$ are homogeneous linear maps of bidegree (ν, ι) and (ν', ι') , respectively, their commutator is defined by

$$[f, f'] = f \circ f' - (-1)^{\nu \cdot \iota'} f' \circ f.$$

We use the notation $|\alpha|$, $|f|$ etc. to denote the cohomological degree of homogeneous cohomology classes, homogeneous linear maps etc.

Setting

$$(\alpha, \beta) := \int_{X^{[n]}} \alpha \beta$$

for any $\alpha, \beta \in H^*(X^{[n]}; \mathbb{Q})$ defines a non-degenerate (anti)symmetric bilinear form on $H^*(X^{[n]}; \mathbb{Q})$ and hence on \mathbb{H} . For any homogeneous linear map $f : \mathbb{H} \rightarrow \mathbb{H}$ its adjoint f^\dagger is characterised by the relation

$$(f(\alpha), \beta) = (-1)^{|f| \cdot |\alpha|} (\alpha, f^\dagger(\beta)).$$

Clearly, $(f \circ g)^\dagger = g^\dagger \circ f^\dagger$.

2.1 Correspondences

Let Y_1 and Y_2 be smooth projective varieties, and let u be a class in the Chow group $A_n(Y_1 \times Y_2)$. (We tacitly assume rational coefficients. This will not always be necessary. On the other hand, we are not interested in integrality questions for the moment, and hence will not pay attention to this problem). The image of u in $H_{2n}(Y_1 \times Y_2)$ will be denoted by the same symbol. u induces a homogeneous linear map

$$u_* : H^i(Y_2) \rightarrow H^{i+2(\dim Y_1 - n)}(Y_1), \quad y \mapsto PD^{-1} p_{1*}(u \cap p_2^* y),$$

where $PD : H^*(Y_1) \rightarrow H_*(Y_1)$ is the Poincaré duality map.

Assume that Y_3 is another smooth projective variety, and $v \in A_m(Y_2 \times Y_3)$. Let p_{ij} be the projection from $Y_1 \times Y_2 \times Y_3$ to the factors $Y_i \times Y_j$, and consider the element

$$w := p_{13*}(p_{12}^* u \cdot p_{23}^* v) \in A_{n+m-\dim Y_2}(Y_1 \times Y_3).$$

Then

$$w_* = u_* \circ v_*.$$

See [10, Ch. 16] for details.

Suppose $U \subset Y_1 \times Y_2$ and $V \subset Y_2 \times Y_3$ are closed subschemes such that $u \in A_*(U)$ and $v \in A_*(V)$. Let

$$W := p_{13}(p_{12}^{-1}(U) \cap p_{23}^{-1}(V))$$

Then the class w defined above is already defined in $A_*(W)$.

The following type of arguments will often show up in the sequel: one shows that the dimension of W is smaller than the degree of w , which forces w to be zero; or that there is at most one irreducible component W_0 of W of maximal dimension with ‘correct’ dimension $\dim(W_0) = \deg(w)$. In this case one must have $w = \mu \cdot [W_0]$ and it suffices to determine the multiplicity μ .

Let $T : Y_1 \times Y_2 \rightarrow Y_2 \times Y_1$ exchange the factors. Then a Chow cycle u induces two maps

$$u_* : H^*(Y_2) \rightarrow H^*(Y_1) \quad \text{and} \quad (Tu)_* : H^*(Y_1) \rightarrow H^*(Y_2)$$

which are related by the formula

$$\int_{Y_1} u_*(\alpha) \cdot \beta = \int_{Y_2} \alpha \cdot (Tu)_*(\beta).$$

This follows directly from the projection formula. Thus $(Tu)_* = u_*^\dagger$.

The following operators were introduced by Nakajima [21]. The study of their properties is the major theme of this article. We take the liberty to change the notations and sign conventions.

Recall that we defined (1.9) subvarieties

$$Q^{[n_1, n_2]} \subset X^{[n_1]} \times X \times X^{[n_2]}$$

of dimension $n_1 + n_2 + 1$. Their fundamental classes are cycles

$$[Q^{[n_1, n_2]}] \in A_{n_1+n_2+1}(X^{[n_1]} \times X \times X^{[n_2]}).$$

Let the projections to the factors be denoted by p_1 , ρ and p_2 .

Definition 2.3 (Nakajima) — Define linear maps

$$q_\ell : H^*(X; \mathbb{Q}) \longrightarrow \text{End}(\mathbb{H}), \quad \ell \in \mathbb{Z},$$

as follows: assume first that $\ell \geq 0$. For $\alpha \in H^*(X; \mathbb{Q})$ and $y \in H^*(X^{[n]}; \mathbb{Q})$ let

$$q_\ell(\alpha)(y) := [Q^{[n+\ell, n]}]_*(\alpha \otimes y) = PD^{-1}p_{1*}([Q^{[n+\ell, n]}] \cap (\rho^*\alpha \cdot p_2^*y)).$$

The operators for negative indices then are determined by the relation

$$q_{-\ell}(\alpha) := (-1)^\ell q_\ell(\alpha)^\dagger.$$

By definition, $q_\ell(\alpha)$ is a homogeneous linear map of bidegree $(\ell, 2\ell - 2 + |\alpha|)$. Moreover, $q_0 = 0$, and if $\ell > 0$, the operator $q_\ell(\alpha)^\dagger$ is induced by the subvarieties $Q^{[n, n+\ell]}$, $n \geq 0$.

2.2 Nakajima's Main Theorem

In this section we review the main result of [21] and some of the immediate consequences. Similar results have been announced by Grojnowski [13].

Theorem 2.4 (Nakajima) — *For any integers n and m and cohomology classes α and β , the operators $q_n(\alpha)$ and $q_m(\beta)$ satisfy the following ‘oscillator relations’:*

$$[q_n(\alpha), q_m(\beta)] = n \cdot \delta_{n+m} \cdot \int_X \alpha\beta \cdot \text{id}_{\mathbb{H}}.$$

□

Here and in the following we adopt the convention that δ_ν equals 1 if $\nu = 0$ and is zero else, and that any integral $\int_Z \alpha$ is zero if $\deg(\alpha) \neq \dim_{\mathbb{R}}(Z)$.

In [21] Nakajima only showed that the commutator relation hold with some universal nonzero constant instead of the coefficient n . The correct value was first computed directly by Ellingsrud and Strømme [7]: up to a sign factor, which depends on our convention, this number is the intersection number of Theorem 1.7. Briefly afterwards, Nakajima gave a different proof using ‘vertex operators’ [22].

Consider the vector spaces

$$W_+ := H^*(X; \mathbb{Q}) \otimes t\mathbb{Q}[t] \quad \text{and} \quad W_- := H^*(X; \mathbb{Q}) \otimes t^{-1}\mathbb{Q}[t^{-1}].$$

Define a non-degenerate skew-symmetric pairing on the vector space $W := W_- \oplus W_+$ by

$$\{\alpha \otimes t^n, \beta \otimes t^m\} := n \cdot \delta_{n+m} \cdot \int_X \alpha \beta.$$

Note that we are taking the expression ‘skew-symmetric’ in a graded sense:

$$\{\alpha \otimes t^n, \beta \otimes t^m\} = -(-1)^{|\alpha| \cdot |\beta|} \{\beta \otimes t^m, \alpha \otimes t^n\}.$$

The *Heisenberg algebra* is the quotient of the tensor algebra $\mathcal{T}W$ by the two-sided ideal I generated by the expressions $[v, w] - \{v, w\} \cdot 1$ with $v, w \in W$:

$$\mathcal{H} := \mathcal{T}W/I.$$

\mathcal{H} is the (restricted) tensor product of countably many copies of Clifford algebras arising from $H^{odd}(X; \mathbb{Q})$ and countably many copies of Weyl algebras arising from $H^{even}(X; \mathbb{Q})$. As W_+ is isotropic with respect to the skew-form $\{, \}$, the subalgebra in \mathcal{H} generated by W_+ is the symmetric algebra S^*W_+ (taken again in a graded sense). This becomes a double graded vector space if we define the bidegree of $\alpha \otimes t^p$ as $(n, 2n - 2 + |\alpha|)$.

Using these notations, Nakajima’s Theorem can be rephrased by saying: Sending $\alpha \otimes t^n \in W$ to $\mathfrak{q}_n(\alpha) \in \text{End}(\mathbb{H})$ defines a representation of \mathcal{H} on \mathbb{H} .

The subspace W_- of monomials of negative degree annihilates the vacuum vector $\mathbf{1} \in \mathbb{H}$ for obvious degree reasons. Hence there is an embedding

$$S^*W_+ \cong \mathcal{H}/\mathcal{H} \cdot W_- \xrightarrow{-\mathbf{1}} \mathcal{H} \cdot \mathbf{1} \subset \mathbb{H}.$$

It is not difficult to check that the Poincaré series of S^*W_+ equals the right hand side of Göttsche’s formula. This implies:

Corollary 2.5 (Nakajima) — *The action of \mathcal{H} on \mathbb{H} induces a module isomorphism $S^*W_+ \rightarrow \mathbb{H}$. In particular, \mathbb{H} is irreducible and generated by the vacuum vector.* \square

In fact, this can be strengthened as follows:

Consider the rational map $a : X^{[n]} \times X^{[m]} \dashrightarrow X^{[n+m]}$ which is defined on the open subset of all pairs (ξ, ξ') with disjoint support by $a(\xi, \xi') := \xi \cup \xi'$. This rational map induces homomorphisms

$$a_* : H^*(X^{[n]}; \mathbb{Q}) \otimes H^*(X^{[m]}; \mathbb{Q}) \longrightarrow H^*(X^{[n+m]}; \mathbb{Q})$$

and

$$a^* : H^*(X^{[n+m]}; \mathbb{Q}) \longrightarrow H^*(X^{[n]}; \mathbb{Q}) \otimes H^*(X^{[m]}; \mathbb{Q})$$

and hence

$$a_* : \mathbb{H} \otimes \mathbb{H} \longrightarrow \mathbb{H} \quad \text{and} \quad a^* : \mathbb{H} \rightarrow \mathbb{H} \otimes \mathbb{H}.$$

Corollary 2.6 (Nakajima, Grojnowski) — *The homomorphism a^* and a_* endow \mathbb{H} with the structure of a Hopf algebra. If S^*W_+ is given the canonical Hopf algebra structure of the symmetric product, then $S^*W_+ \rightarrow \mathbb{H}$ is an isomorphism of Hopf algebras.* \square

3 The boundary operator

This section contains the main technical results of the paper. The key to our solution of the Chern class problem is the introduction of the boundary operator $\mathfrak{d} \in \text{End}(\mathbb{H})$. This is done in 3.2. We begin with the discussion of related topics and ingredients for later proofs.

3.1 Virasoro generators

Starting from the basic generators \mathfrak{q}_n and the fundamental oscillator relations we will define the corresponding Virasoro generators \mathfrak{L}_n in analogy to the procedure in conformal field theory. We will then give concrete geometric interpretations for these generators.

Let $\delta : H^*(X) \rightarrow H^*(X \times X) = H^*(X) \otimes H^*(X)$ be the push-forward map associated to the diagonal embedding. Equivalently, this is the linear map adjoint to the cup-product map. If $\delta(\alpha) = \sum_i \alpha'_i \otimes \alpha''_i$, we will write $\mathfrak{q}_n \mathfrak{q}_m \delta(\alpha)$ for $\sum_i \mathfrak{q}_n(\alpha'_i) \mathfrak{q}_m(\alpha''_i)$.

Definition 3.1 — Define operators $\mathfrak{L}_n : H^*(X; \mathbb{Q}) \rightarrow \text{End}(\mathbb{H})$, $n \in \mathbb{Z}$, as follows:

$$\mathfrak{L}_n := \frac{1}{2} \sum_{\nu \in \mathbb{Z}} \mathfrak{q}_\nu \mathfrak{q}_{n-\nu} \delta, \quad \text{if } n \neq 0$$

and

$$\mathfrak{L}_0 := \sum_{\nu > 0} \mathfrak{q}_\nu \mathfrak{q}_{-\nu} \delta.$$

Remark 3.2 — i) The sums that appear in the definition are formally infinite. However, as operators on any fixed vector in \mathbb{H} , only finitely many of them are nonzero. Hence the sums are locally finite and the operators \mathfrak{L}_n are well-defined. $\mathfrak{L}_n(\alpha)$ is homogeneous of bidegree $(n, 2n + |\alpha|)$

ii) Using the physicists' normal order convention

$$: \mathfrak{q}_n \mathfrak{q}_m : := \begin{cases} \mathfrak{q}_n \mathfrak{q}_m & \text{if } n \geq m, \\ \mathfrak{q}_m \mathfrak{q}_n & \text{if } n \leq m, \end{cases}$$

the operators \mathfrak{L}_n can be uniformly expressed as

$$\mathfrak{L}_n = \frac{1}{2} \sum_{\nu \in \mathbb{Z}} : \mathfrak{q}_\nu \mathfrak{q}_{n-\nu} : \delta.$$

Theorem 3.3 — The operators \mathfrak{L}_n and \mathfrak{q}_m satisfy the following commutator relations:

1. $[\mathfrak{L}_n(\alpha), \mathfrak{q}_m(\beta)] = -m \cdot \mathfrak{q}_{n+m}(\alpha\beta)$.
2. $[\mathfrak{L}_n(\alpha), \mathfrak{L}_m(\beta)] = (n-m) \cdot \mathfrak{L}_{n+m}(\alpha\beta) - \frac{n^3-n}{12} \delta_{n+m} \cdot \int_X \alpha\beta \cdot \text{id}_{\mathbb{H}}$.

Proof. Assume first that $n \neq 0$. For any classes α and β with

$$\delta(\alpha) = \sum_i \alpha'_i \otimes \alpha''_i$$

we have

$$\begin{aligned} [\mathfrak{q}_\nu(\alpha'_i) \mathfrak{q}_{n-\nu}(\alpha''_i), \mathfrak{q}_m(\beta)] &= \mathfrak{q}_\nu(\alpha'_i) [\mathfrak{q}_{n-\nu}(\alpha''_i), \mathfrak{q}_m(\beta)] \\ &\quad + (-1)^{|\beta| \cdot |\alpha''_i|} [\mathfrak{q}_\nu(\alpha'_i), \mathfrak{q}_m(\beta)] \mathfrak{q}_{n-\nu}(\alpha''_i) \\ &= (-m) \delta_{n+m-\nu} \cdot \mathfrak{q}_{n+m}(\alpha'_i) \cdot \int_X \alpha''_i \beta \\ &\quad + (-1)^{|\beta| \cdot |\alpha|} (-m) \delta_{\nu+m} \cdot \int_X \beta \alpha'_i \cdot \mathfrak{q}_{n+m}(\alpha''_i). \end{aligned}$$

If we sum up over all ν and i , we get

$$2[\mathfrak{L}_n(\alpha), \mathfrak{q}_m(\beta)] = \sum_\nu [\mathfrak{q}_\nu \mathfrak{q}_{n-\nu} \delta(\alpha), \mathfrak{q}_m(\beta)] = (-m) \cdot \mathfrak{q}_{n+m}(\gamma)$$

with

$$\gamma = pr_{1*}(\delta(\alpha) \cdot pr_2^*(\beta)) + (-1)^{|\beta| \cdot |\alpha|} \cdot pr_{2*}(pr_1^*(\beta) \cdot \delta(\alpha)) = 2 \cdot \alpha\beta.$$

Similarly, for $\nu > 0$,

$$[\mathfrak{q}_\nu \mathfrak{q}_{-\nu} \delta(\alpha), \mathfrak{q}_m(\beta)] = -m \cdot \mathfrak{q}_m(\alpha\beta) \cdot (\delta_{m-\nu} + \delta_{m+\nu}).$$

Thus summing up over all $\nu > 0$ we find again

$$[\mathfrak{L}_0(\alpha), \mathfrak{q}_m(\beta)] = -m \cdot \mathfrak{q}_m(\alpha\beta).$$

This proves the first part of the theorem.

As for the second part, assume first that $n \geq 0$. In order to avoid case considerations let us agree that $\mathfrak{q}_{\frac{N}{2}}$ is zero if N is odd. Then we may write:

$$\mathfrak{L}_m = \frac{1}{2} \mathfrak{q}_{\frac{m}{2}}^2 \delta + \sum_{\mu > \frac{m}{2}} \mathfrak{q}_\mu \mathfrak{q}_{m-\mu} \delta.$$

By the first part of the theorem we have

$$[\mathfrak{L}_n(\alpha), \mathfrak{q}_\mu \mathfrak{q}_{m-\mu} \delta(\beta)] = \left(-\mu \mathfrak{q}_{n+\mu} \mathfrak{q}_{m-\mu} + (\mu - m) \mathfrak{q}_\mu \mathfrak{q}_{n+m-\mu} \right) \delta(\alpha\beta).$$

In the following calculation we suppress α, β and δ up to the very end. Summing up over all $\mu \geq 0$, we get:

$$\begin{aligned} [\mathfrak{L}_n, \mathfrak{L}_m] &= -\frac{m}{4} (\mathfrak{q}_{n+\frac{m}{2}} \mathfrak{q}_{\frac{m}{2}} + \mathfrak{q}_{\frac{m}{2}} \mathfrak{q}_{n+\frac{m}{2}}) \\ &\quad + \sum_{\mu > \frac{m}{2}} (\mu - m) \mathfrak{q}_\mu \mathfrak{q}_{n+m-\mu} + \sum_{\mu > \frac{m}{2}} (-\mu) \mathfrak{q}_{n+\mu} \mathfrak{q}_{m-\mu} \\ &= -\frac{m}{4} (\mathfrak{q}_{n+\frac{m}{2}} \mathfrak{q}_{\frac{m}{2}} + \mathfrak{q}_{\frac{m}{2}} \mathfrak{q}_{n+\frac{m}{2}}) \\ &\quad + \sum_{\mu > \frac{m}{2}} (\mu - m) \mathfrak{q}_\mu \mathfrak{q}_{n+m-\mu} + \sum_{\mu > n+\frac{m}{2}} (n - \mu) \mathfrak{q}_\mu \mathfrak{q}_{n+m-\mu} \end{aligned}$$

Hence

$$\begin{aligned}
[\mathfrak{L}_n, \mathfrak{L}_m] - (n-m) \sum_{\mu > \frac{n+m}{2}} \mathfrak{q}_\mu \mathfrak{q}_{n+m-\mu} &= -\frac{m}{4} (\mathfrak{q}_{n+\frac{m}{2}} \mathfrak{q}_{\frac{m}{2}} + \mathfrak{q}_{\frac{m}{2}} \mathfrak{q}_{n+\frac{m}{2}}) \\
&+ \sum_{\frac{m}{2} < \mu \leq \frac{m+n}{2}} (\mu-m) \mathfrak{q}_\mu \mathfrak{q}_{m+n-\mu} \\
&- \sum_{\frac{n+m}{2} < \mu \leq n+\frac{m}{2}} (n-\mu) \mathfrak{q}_\mu \mathfrak{q}_{n+m-\mu}
\end{aligned}$$

Now split off the summands corresponding to the indices $\mu = \frac{m+n}{2}$ and $\mu = n + \frac{m}{2}$ from the sums. Substituting $n+m-\mu$ for μ in the second sum on the right hand side, we are left with the expression:

$$[\mathfrak{L}_n, \mathfrak{L}_m] - (n-m) \mathfrak{L}_{n+m} = -\frac{m}{4} [\mathfrak{q}_{\frac{m}{2}}, \mathfrak{q}_{n+\frac{m}{2}}] + \sum_{\frac{m}{2} < \mu < \frac{n+m}{2}} (\mu-m) [\mathfrak{q}_\mu, \mathfrak{q}_{n+m-\mu}]$$

The right hand side is zero unless $n+m=0$. Hence we see that

$$[\mathfrak{L}_n(\alpha), \mathfrak{L}_m(\beta)] = (n-m) \mathfrak{L}_{n+m}(\alpha\beta) + \delta_{n+m} \cdot \int_X \alpha\beta \cdot N,$$

where N is the number

$$N = \sum_{0 < \nu < \frac{n}{2}} \nu(\nu-n) \quad \text{if } n \text{ is odd,}$$

and

$$N = \sum_{0 < \nu < \frac{n}{2}} \nu(\nu-n) - \frac{n^2}{8} \quad \text{if } n \text{ is even.}$$

An easy computation shows that in both cases N equals $(n-n^3)/12$. \square

Recall the definition of the varieties $E^{[n, n']} \subset X^{[n]} \times X \times X^{[n']}$ in (1.9).

Definition 3.4 — Let ℓ be a nonnegative integer and let

$$\epsilon_\ell : H^*(X) \rightarrow \text{End}(\mathbb{H})$$

be the linear map

$$\epsilon_\ell(\alpha)(y) = [E^{[n+\ell, n]}]_* (\alpha \otimes y) = PD^{-1} p_{1*} ([E^{[n+\ell]}] \cap (\rho^* \alpha \cdot p_2^* y))$$

for $\alpha \in H^*(X; \mathbb{Q})$ and $y \in H^*(X^{[n]}; \mathbb{Q})$.

The following theorem gives a ‘finite’ geometric interpretation of the infinite sums which define the Virasoro operators.

Theorem 3.5 — *Let n be a nonnegative integer.*

1.

$$[\epsilon_n(\alpha), \mathfrak{q}_m(\beta)] = \begin{cases} m \cdot \mathfrak{q}_{n+m}(\alpha\beta) & \text{if } m > 0 \text{ or } m < -n. \\ 0 & \text{else.} \end{cases}$$

2.

$$\epsilon_n + \mathfrak{L}_n = \frac{1}{2} \sum_{0 < \nu < n} \mathfrak{q}_\nu \mathfrak{q}_{n-\nu} \delta.$$

Proof. Ad 1: Assume first that $m \geq 1$. To simplify the notations we introduce the short-hand

$$X^{[n_1],[n_2],\dots,[n_k]} := X^{[n_1]} \times X^{[n_2]} \times \dots \times X^{[n_k]}$$

Suppose $\ell \geq 0$, and consider the following diagram

$$\begin{array}{ccccc} X^{[\ell+n+m],[1],[\ell+m]} & \xleftarrow{p_{123}} & X^{[\ell+n+m],[1],[\ell+m],[1],[\ell]} & \xrightarrow{p_{345}} & X^{[\ell+m],[1],[\ell]} \\ & & \downarrow p_{1245} & & \\ & & X^{[\ell+n+m],[1],[1],[\ell]} & & \end{array}$$

The product operator $\epsilon_n \mathfrak{q}_m$ is induced by the class

$$z := p_{1245*} (p_{123}^* [E^{[\ell+m+n,\ell+m]}] \cdot p_{345}^* [Q^{[\ell+m,\ell]}]) \in A_{2\ell+n+m+1}(Z')$$

where

$$\begin{aligned} Z' &:= p_{1245} (p_{123}^{-1} (E^{[\ell+m+n,\ell+m]}) \cap p_{345}^{-1} (Q^{[\ell+m,\ell]})) \\ &\subset Z := \{(\xi', x, y, \xi) \mid \exists \eta : \xi' - \eta = nx, \eta - \xi = my, x \in \eta\} \end{aligned}$$

Here the notation $\eta - \xi = my$ should comprise the conditions: ξ is a subscheme of η , and the ideal sheaf of ξ in η is of length m and is supported at y etc.

Similarly, the operator $\mathfrak{q}_n \epsilon_m$ is induced by a class $v \in A_{2\ell+m+n+1}(V')$ with

$$V' \subset V := \{(\xi', x, y, \xi) \mid \exists \eta' : \xi' - \eta' = mx, \eta' - \xi = ny, y \in \xi\}.$$

Moreover, if $T : X^{[\ell+m+n],[1],[1],[\ell]} \longrightarrow X^{[\ell+m+n],[1],[1],[\ell]}$ exchanges the two copies of X in the middle, then the commutator $[\epsilon_n, \mathfrak{q}_m]$ is induced by $z - T(v)$.

Now observe that off the diagonal $\{x = y\} \subset X^{[\ell+m+n],[1],[1],[\ell]}$ the subsets Z and $T(V)$ are equal. Moreover, there is only one component of (maximal possible) dimension $2\ell + n + m + 1$. It is easy to see that this component has multiplicity 1 both in z and $T(v)$: the intersection

$$p_{123}^{-1} (E^{[\ell+m+n,\ell+m]}) \cap p_{345}^{-1} (Q^{[\ell+m,\ell]})$$

is transversal over a general point in this component of Z , and maps injectively into Z . Thus the only contributions to $z - T(v)$ may arise from the diagonal part. Now

$$V \cap \{x = y\} = \{(\xi', x, x, \xi) \mid \xi' - \xi = (n+m)x, x \in \xi\}.$$

We have seen earlier (1.8) that this set has dimension $\leq 2\ell + n + m$ and hence may be disregarded. On the other hand

$$Z \cap \{x = y\} = \{(\xi', x, x, \xi) \mid \xi' - \xi = (n + m)x\}.$$

Again using 1.8 we see that this set has only one component D of (maximal) dimension $2\ell + n + m + 1$. Moreover, this component is the image of the embedding

$$\iota : Q^{[\ell+n+m, \ell]} \rightarrow X^{[\ell+n+m], [1], [1], [\ell]}, (\xi', x, \xi) \mapsto (\xi', x, x, \xi).$$

Let $\alpha, \beta \in H^*(X; \mathbb{Q})$ and $y \in H^*(X^{[\ell]}; \mathbb{Q})$. Then we have

$$\begin{aligned} & p_{1*}([D] \cap p_{23}^*(\alpha \otimes \beta) \cdot p_4^*y) \\ &= p_{1*}(\iota_*[Q^{[\ell+n+m, \ell]}] \cap p_{23}^*(\alpha \otimes \beta) \cdot p_4^*y) \\ &= p_{1*}([Q^{[\ell+n+m, \ell]}] \cap \iota^*(p_{23}^*(\alpha \otimes \beta) \cdot p_4^*y)) \\ &= p_{1*}([Q^{[\ell+n+m, \ell]}] \cap p_2^*(\alpha\beta) \cdot p_3^*y) \end{aligned}$$

This shows that

$$[\epsilon_n(\alpha), \mathfrak{q}_m(\beta)] = \mu \cdot \mathfrak{q}_{n+m}(\alpha\beta)$$

for some integer μ . Hence it remains to compute the multiplicity μ of $[D]$ in z . To this end we pick a general point $d \in D$ and inspect the intersection of $\bar{p}_{23}^{-1}(E^{[\ell+n+m, \ell]})$ and $p_{345}^{-1}(Q^{[\ell+m, \ell]})$ along the fibre $p_{1245}^{-1}(d)$.

A general point in D is of the form

$$d = (\xi', x, x, \xi) \quad \text{with} \quad \xi' = \xi \cup \zeta,$$

where ζ is a curvilinear subscheme of X of length $n + m$, supported in a single point x which is disjoint from ξ . Since ζ is curvilinear, there is a unique subscheme $\eta \subset \zeta$ of length m , and hence $p_{1245}^{-1}(d)$ consists of the single point

$$d' = (\xi \cup \zeta, x, \xi \cup \eta, x, \xi)$$

Near d' the varieties $X^{[\ell+m+n], [1], [\ell+m], [1], [\ell]}$ and $X^{[\ell], [\ell], [\ell]} \times X^{[m+n], [1], [m], [1]}$ are locally isomorphic; and similarly $E^{[\ell+m+n, \ell+m]}$ to $X^{[\ell]} \times E^{[m+n, m]}$ and $Q^{[\ell+m, \ell]}$ to $X^{[\ell]} \times X_0^{[m]}$. Thus we may split off the factors $X^{[\ell]}$ from the geometric picture. In the end this amounts to saying that we may assume without loss of generality that $\ell = 0$.

Moreover, the calculation is local in X , so that we may assume that $X = \mathbb{A}^2 = \text{Spec}\mathbb{C}[z, w]$ and $\mathcal{I}_\zeta = (w, z^{n+m})$, $\mathcal{I}_\eta = (w, z^m)$ and $\mathcal{I}_x = (w, z)$. Then d' has an affine neighbourhood $\cong \mathbb{A}^{4m+2n+4}$ in $X^{[n+m], [1], [m], [1]}$ with coordinate functions

$$a_0, \dots, a_{n+m-1}, b_0, \dots, b_{n+m-1}, w_1, z_1, c_0, \dots, c_{m-1}, d_0, \dots, d_{m-1}, w_2, z_2,$$

which parametrises quadrupels (ζ, x, η, y) of subschemes in X given by the ideals

$$(w - g_1(z), f_1(z)), \quad (w - w_1, z - z_1), \quad (w - g_2(z), f_2(z)), \quad (w - w_2, z - z_2),$$

where

$$f_1(z) = a_0 + a_1z + \dots + z^{n+m}, \quad g_1(z) = b_0 + b_1z + \dots + b_{n+m-1}z^{n+m-1}$$

and

$$f_2(z) = c_0 + c_1z + \dots + z^m, \quad g_2(z) = d_0 + d_1z + \dots + z^m.$$

Now (η, y) belongs to $X_0^{[m]}$, i.e. $\text{Supp}(\eta) = \{y\}$, if and only if

$$f_2(z) = (z - z_2)^m \text{ and } w_2 = g_2(z_2). \quad (6)$$

And (ζ, x, η) belongs to $Q^{[n+m, m]}$ if and only if the following three conditions are satisfied: $\eta \subset \zeta$, i.e.

$$g_1(z) = g_2(z) + f_2(z) \cdot h(z) \text{ and } f_1(z) = f_2(z) \cdot k(z) \quad (7)$$

with polynomials h and k of degree $n - 1$ and n , respectively; the ideal sheaf $\mathcal{I}_{\eta/\zeta}$ is supported at x , i.e.

$$k(z) = (z - z_1)^m \text{ and } w_1 = g_1(z_1) \quad (8)$$

and finally, x must be contained in η , which imposes the condition

$$f_2(z_1) = 0 \quad (9)$$

One easily checks that the equations (6) - (8) cut out a smooth subvariety which projects isomorphically to the affine space $\text{Spec } \mathbb{C}[\tilde{x}_1, z_2, b_0, \dots, b_{n+m-1}]$. Moreover, in these coordinates the last condition (9) simply reads $(\tilde{x}_1 - z_2)^m = 0$. Hence the multiplicity μ equals the exponent m .

Next, we consider the case $[\mathfrak{e}_n, \mathfrak{q}_{-m}]$ with $0 \leq m \leq n$. There is nothing to prove if $m = 0$. Hence assume that $m > 0$. Dimension arguments similar to the ones above show that the cycle v which induces the commutator $[\mathfrak{q}_{-m}, \mathfrak{e}_n]$ must be supported on the closed subsets

$$V := \{(\xi, x, x, \zeta) \mid \xi \supset \zeta \ni x, \xi - \zeta = (n + m)x\} \subset X^{[\ell+n-m], [1], [1], [\ell]}, \quad \ell \geq 0.$$

The cycle v has degree $2\ell + n - m + 1$, so that it suffices to show that $\dim(V) \leq 2\ell + n - m$. This follows from Lemma 1.8.

It remains to consider the case $[\mathfrak{e}_n, \mathfrak{q}_m]$ with $m < -n$. A dimension check of the set-theoretic support of the intersection cycle shows that we must have

$$[\mathfrak{e}_n(\alpha), \mathfrak{q}_{-m}(\beta)] = \mu \cdot \mathfrak{q}_{n-m}(\alpha\beta)$$

for some integer μ , independently of α and β . To determine μ , we proceed algebraically and take the commutator with $\mathfrak{q}_{m-n}(1)$:

$$[[\mathfrak{e}_n(\alpha), \mathfrak{q}_{-m}(\beta)], \mathfrak{q}_{m-n}(1)] = \mu \cdot [\mathfrak{q}_{n-m}(\alpha\beta), \mathfrak{q}_{m-n}(1)] = \mu(n - m) \int_X \alpha\beta \cdot \text{id}_{\mathbb{H}}.$$

On the other hand, combining the Jacobi identity, the oscillator relations and the first part of the proof yields

$$\begin{aligned}
[[\mathfrak{e}_n(\alpha), \mathfrak{q}_{-m}(\beta)], \mathfrak{q}_{m-n}(1)] &= [[\mathfrak{e}_n(\alpha), \mathfrak{q}_{m-n}(1)], \mathfrak{q}_{-m}(\alpha)] \\
&= (m-n)[\mathfrak{q}_m(\alpha), \mathfrak{q}_{-m}(\beta)] \\
&= m(m-n) \int_X \alpha\beta \cdot \text{id}_{\mathbb{H}}.
\end{aligned}$$

It follows that $\mu = -m$.

Ad 2: Consider the difference $\eta := \mathfrak{e}_n(\alpha) + \mathfrak{L}_n(\alpha) - \frac{1}{2} \sum_{\nu=1}^{n-1} \mathfrak{q}_\nu \mathfrak{q}_{n-\nu} \delta(\alpha)$. Comparing the expressions in 3.3 and part 1 of the theorem we see that η commutes with all operators \mathfrak{q}_m , $m \in \mathbb{Z}$. Since \mathbb{H} is a simple \mathcal{N} -module, η must be a scalar (in some algebraic extension of \mathbb{Q}), which is impossible: if $n > 0$, then η has non-trivial bidegree $(n, 2n + |\alpha|)$, and if $n = 0$, it is easy to see directly that $\eta \cdot 1 = 0$. \square

Remark 3.6 — In particular, the operator $\mathfrak{L}_0(\alpha)$ has the following geometric interpretation: the universal family $\Xi_n \subset X^{[n]} \times X$ induces a homomorphism

$$[\Xi_n]_* : H^*(X; \mathbb{Q}) \longrightarrow H^*(X^{[n]}; \mathbb{Q}),$$

and

$$\mathfrak{L}_0(\alpha)(y) = [\Xi_n]_*(\alpha) \cdot y \quad \text{for all } y \in H^*(X^{[n]}, \mathbb{Q}).$$

In particular, if we insert $\alpha = 1_X$, we get

$$\mathfrak{L}_0(1_X)(y) = n \cdot y \text{ for all } y \in H^*(X^{[n]}; \mathbb{Q}).$$

Thus $\mathfrak{L}_0(1_X)$ is the ‘energy’ or ‘counting’ operator, that measures with which ‘energy level’, i.e. how many points we are dealing. This can, of course, also be deduced directly from the definition of \mathfrak{L}_0 .

3.2 The boundary of the Hilbert scheme

For any partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_s > 0)$ of n the tuples $\sum_{1 \leq i \leq s} \lambda_i x_i$, $x_i \in X$, form a locally closed subset $S_\lambda^n X$ in $S^n X$. Let $X_\lambda^{[n]} = \rho^{-1}(S_\lambda^n X)$. It follows from Briançon’s Theorem that $X_\lambda^{[n]}$ is irreducible and

$$\dim(X_\lambda^{[n]}) = \sum_{1 \leq i \leq s} (\lambda_i + 1) = n + s.$$

The generic open stratum is $X_{(1,1,\dots,1)}^{[n]}$. It corresponds to the configuration space of unordered n -tuples of pairwise distinct points. Furthermore, there is precisely one stratum of codimension 1, namely $X_{(2,1,\dots,1)}^{[n]}$.

If $\lambda = (\lambda_1, \dots, \lambda_s)$ and $\mu = (\mu_1, \dots, \mu_{s'})$ are partitions of n , then $X_\mu^{[n]}$ is contained in the closure of $X_\lambda^{[n]}$ if and only if there is a surjection

$$\varphi : \{1, \dots, s\} \rightarrow \{1, \dots, s'\}$$

such that $\mu_j = \sum_{i \in \varphi^{-1}(j)} \lambda_i$ for all j . It follows that

$$\partial X^{[n]} := \bigcup_{\lambda \neq (1, \dots, 1)} X_\lambda^{[n]} = \overline{X_{(2,1,\dots,1)}^{[n]}}$$

is an irreducible divisor in $X^{[n]}$. As it is the complement of the configuration space in $X^{[n]}$ we might and will call it the *boundary* of $X^{[n]}$.

Lemma 3.7 — *Let $E \subset X^{[n+1,n]}$ be the exceptional divisor. Then*

$$p_1^* \partial X^{[n+1]} - p_2^* \partial X^{[n]} = 2 \cdot E.$$

Proof. Points in $X^{[n+1,n]}$ are triples (ξ', x, ξ) with $\xi \subset \xi'$ and $\mathcal{I}_{\xi/\xi'} \cong k(x)$, and $p_1^{-1}(\partial X^{[n+1]})$ consists of those triples such that there is a point $y \in X$ with $\ell(\xi'_y) \geq 2$. Now either $y = x$, in which case $\ell(\xi'_x) = \ell(\xi'_x) - 1 \geq 1$ and therefore $(\xi', x, \xi) \in E$, or $y \neq x$, in which case $\ell(\xi'_y) = \ell(\xi'_y) \geq 2$ so that $(\xi', x, \xi) \in p_2^{-1}(\partial X^{[n]})$. Hence set-theoretically, we have $p_1^{-1}(\partial X^{[n+1]}) = p_2^{-1}(\partial X^{[n]}) \cup E$. We must check the multiplicities.

Off the exceptional divisor E we have $X^{[n+1,n]} \setminus E = X^{[n]} \times X \setminus \Xi_n$, which embeds as an open subset into $X^{[n+1]}$. Clearly, $(X^{[n]} \times X \setminus \Xi_n) \cap \partial X^{[n+1]} = p_1^* \partial X^{[n]}$. Thus $p_2^* \partial X^{[n+1]} - p_1^* \partial X^{[n]} = \mu \cdot E$. In order to compute the multiplicity μ we pick a general point in E which is of the form $(\eta \cup \zeta, x, \eta \cup \{x\})$, where $x \notin \eta$ and ζ has length 2 and is supported at x . Without loss of generality we may assume that η is empty, i.e., that $n = 1$. But then $X^{[2,1]}$ is the blow-up of $X \times X$ along the diagonal, E is the exceptional divisor, and $p_2 : X^{[2,1]} \rightarrow X^{[2]}$ is the quotient map for the action of \mathfrak{S}_2 on the blow-up. In this picture E is the ramification divisor, $\partial X^{[2]}$ is the branching divisor, and the ramification order is 2. Hence indeed, $\mu = 2$. \square

We will need a different description of the divisor $\partial X^{[n]}$ in sheaf theoretic terms.

Let $p : \Xi_n \rightarrow X^{[n]}$ be the projection, and define sheaves

$$\mathcal{O}_X^{[n]} := p_*(\mathcal{O}_{\Xi_n}) \in \text{Coh}(X^{[n]}).$$

As p is flat and finite of degree n , $\mathcal{O}_X^{[n]}$ is locally free of rank n . The fibre at a point $\xi \in X^{[n]}$ is the \mathbb{C} -vector space underlying the algebra \mathcal{O}_ξ .

Lemma 3.8 — $c_1(\mathcal{O}_X^{[n]}) = -\frac{1}{2} [\partial X^{[n]}]$.

Proof. Consider the following incidence scheme with the natural projections:

$$X^{[n+1]} \longleftarrow X^{[n+1,n]} \longrightarrow X^{[n]}.$$

We have seen earlier in 1.4 that $\mathcal{I}_{n+1,n} = (id, \rho)_* \mathcal{O}_{X^{[n+1,n]}}(-E)$ and hence that $p_* \mathcal{I}_{n+1,n} = \mathcal{O}_X^{[n+1,n]}(-E)$. This shows

$$p_1^* c_1(\mathcal{O}_X^{[n+1]}) - p_2^* c_1(\mathcal{O}_X^{[n]}) = -E.$$

On the other hand, by Lemma 3.7,

$$p_1^* \partial X^{[n+1]} - p_2^* \partial X^{[n]} = 2 \cdot E.$$

Therefore, if we put $\gamma_n := c_1(\mathcal{O}_X^{[n]}) + \frac{1}{2} \partial X^{[n]}$, we get $p_2^* \gamma_n = p_1^* \gamma_{n+1}$. Now $\gamma_0 = \gamma_1 = 0$, since $\mathcal{O}_X^{[1]} \cong \mathcal{O}_{X^{[1]}}$ and $\partial X = \emptyset$. Assume by induction that $\gamma_n = 0$. It follows that $p_1^* \gamma_{n+1} = 0$, and since $p_1 : X^{[n+1,n]} \rightarrow X^{[n+1]}$ is generically finite and surjective, we must have $\gamma_{n+1} = 0$ as well. \square

Definition 3.9 — Let $\mathfrak{d} : \mathbb{H} \rightarrow \mathbb{H}$ be the homogeneous linear map of bidegree $(0, 2)$ given by

$$\mathfrak{d}(x) := c_1(\mathcal{O}_X^{[n]}) \cdot x = -\frac{1}{2} \left[\partial X^{[n]} \right] \cdot x \quad \text{for all } x \in H^*(X^{[n]}).$$

For any endomorphism $\mathfrak{f} \in \text{End}(\mathbb{H})$ its derivative is $\mathfrak{f}' := [\mathfrak{d}, \mathfrak{f}]$. As usual, we write $\mathfrak{f}^{(n)} := (\text{ad } \mathfrak{d})^n(\mathfrak{f})$ for the higher derivatives.

It follows directly from the definition of the commutator that $\mathfrak{f} \mapsto \mathfrak{f}'$ is a derivation, i.e. for any two operators $\mathfrak{a}, \mathfrak{b} \in \text{End}(\mathbb{H})$ the ‘Leibniz rule’ holds:

$$(\mathfrak{a}\mathfrak{b})' = \mathfrak{a}'\mathfrak{b} + \mathfrak{a}\mathfrak{b}' \quad \text{and} \quad [\mathfrak{a}, \mathfrak{b}]' = [\mathfrak{a}', \mathfrak{b}] + [\mathfrak{a}, \mathfrak{b}'].$$

Moreover, if $\mathfrak{f} : H^*(X^{[\ell]}) \rightarrow H^*(X^{[n]})$ is a homogeneous linear map, then $|\mathfrak{f}'| = |\mathfrak{f}| + 2$, so that \mathfrak{f} and \mathfrak{f}' have the same parity. Furthermore,

$$(\mathfrak{f}')^\dagger = -(\mathfrak{f}^\dagger)'$$

Indeed,

$$\begin{aligned} \int_{X^{[n]}} \mathfrak{f}'(y) \cdot z &= \int_{X^{[n]}} \mathfrak{f}(y) \cdot \mathfrak{d}(z) - \mathfrak{f}(\mathfrak{d}(y)) \cdot z \\ &= (-1)^{|y| \cdot |\mathfrak{f}|} \int_{X^{[\ell]}} y \cdot \mathfrak{f}^\dagger(\mathfrak{d}(z)) - y \cdot \mathfrak{d}\mathfrak{f}^\dagger(z) \\ &= -(-1)^{|y| \cdot |\mathfrak{f}|} \int_{X^{[\ell]}} y \cdot (\mathfrak{f}^\dagger)'(z). \end{aligned}$$

Let $n' > n$ be nonnegative integers, and consider the incidence variety $X^{[n',n]} \subset X^{[n']} \times X^{[n]}$. Recall the definition of the ideal sheaf $\mathcal{I}_{n',n}$ and the exact sequence

$$0 \rightarrow \mathcal{I}_{n',n} \rightarrow p_{1,X}^* \mathcal{O}_{\Xi_{n'}} \rightarrow p_{2,X}^* \mathcal{O}_{\Xi_n} \rightarrow 0.$$

Then $p_*(\mathcal{I}_{n',n})$ is a locally free sheaf of rank $n' - n$ on $X^{[n',n]}$.

In a certain sense, the following lemma simply is a reformulation of the definition of the derivative.

Lemma 3.10 — Let $u_* : H^*(X^{[n]}; \mathbb{Q}) \rightarrow H^*(X^{[n']}; \mathbb{Q})$ be the induced linear map associated to a class $u \in A_*(X^{[n', n]})$. Then

$$(u_*)' = (c_1(p_*(\mathcal{I}_{n', n})) \cdot u)_*$$

Proof. Let $y \in H^*(X^{[n]}; \mathbb{Q})$. Then

$$\begin{aligned} (u_*)'(y) &= \mathfrak{d}(u_*(y)) - u_*(\mathfrak{d}(y)) \\ &= c_1(p_*\mathcal{O}_{\Xi_{n'}}) \cdot PD^{-1}p_{1*}(u \cdot p_2^*y) \\ &\quad - PD^{-1}p_{1*}(u \cdot p_2^*(c_1(p_*\mathcal{O}_{\Xi_n}) \cdot y)) \\ &= PD^{-1}p_{1*}((p_1^*c_1(p_*\mathcal{O}_{\Xi_{n'}}) - p_2^*c_1(p_*\mathcal{O}_{\Xi_n})) \cdot u \cdot p_2^*y) \\ &= v_*(y) \end{aligned}$$

with $v = (p_1^*c_1(p_*\mathcal{O}_{\Xi_{n'}}) - p_2^*c_1(p_*\mathcal{O}_{\Xi_n})) \cdot u$, and

$$\begin{aligned} p_1^*c_1(p_*\mathcal{O}_{\Xi_{n'}}) - p_2^*c_1(p_*\mathcal{O}_{\Xi_n}) &= c_1(p_*p_{1, X}^*\mathcal{O}_{\Xi_{n'}}) - c_1(p_*p_{2, X}^*\mathcal{O}_{X_n}) \\ &= c_1(p_*\mathcal{I}_{n', n}). \end{aligned}$$

□

3.3 The derivative of q_n

In order to understand the intersection behaviour of the boundary $\partial X^{[n]}$ we need to know how the operator \mathfrak{d} commutes with the basic operators q_n , in other words: we need to compute the derivative of q_n .

The following theorem is the main technical theorem of this paper. It describes the derivative of the operator q_n in two ways: By its action on any of the other basic operators, and as a polynomial expression in the basic operators.

Let K denote the canonical class of the surface X .

Theorem 3.11 — For all $n, m \in \mathbb{Z}$ and $\alpha, \beta \in H^*(X; \mathbb{Q})$ the following holds:

1. $[q_n'(\alpha), q_m(\beta)] = -nm \cdot \left\{ q_{n+m}(\alpha\beta) + \frac{|n|-1}{2} \delta_{n+m} \cdot \int_X K\alpha\beta \cdot \text{id}_{\mathbb{H}} \right\}$.
2. $q_n'(\alpha) = n \cdot \mathfrak{L}_n(\alpha) + \frac{n(|n|-1)}{2} q_n(K\alpha)$.

Corollary 3.12 — The operators \mathfrak{d} and $q_1(\alpha)$, $\alpha \in H^*(X)$, suffice to generate \mathbb{H} from the vacuum **1**. □

Proof of the theorem. The second assertion is an immediate consequence of the first: by Nakajima's relations 2.4 and the relations 3.3 we see that

$$\begin{aligned} [n \cdot \mathfrak{L}_n(\alpha) + \frac{n(|n|-1)}{2} q_n(K\alpha), q_m(\beta)] &= \\ -nm \cdot q_{n+m}(\alpha\beta) + \delta_{n+m} \frac{n^2(|n|-1)}{2} \int_X K\alpha\beta \cdot \text{id}_{\mathbb{H}}. \end{aligned}$$

Hence the difference of d'_n and the expression on the right hand side in the theorem commutes with all operators q_m , $m \in \mathbb{Z}$. Since \mathbb{H} is an irreducible \mathcal{N} -module, it follows from Schur's Lemma that this difference is given by multiplication with a scalar (say, after passage to some algebraic closure of \mathbb{Q}). But this is impossible for degree reasons: the bidegree of $d'_n(\alpha)$ is $(n, 2n + |\alpha|)$. (The case $n = 0$ being trivial anyhow.)

The proof of the first assertion has two parts of quite different nature: We need to distinguish the cases $n + m \neq 0$ and $n + m = 0$ and deal with them separately.

Proposition 3.13 — $[q'_n(\alpha), q_m(\beta)] = -nm \cdot q_{n+m}(\alpha\beta)$ for any two integers n, m with $n + m \neq 0$ and cohomology classes $\alpha, \beta \in H^*(X)$.

Proof. Step 1: Assume that n and m are positive. We proceed as in the proof of Theorem 3.5. Let ℓ be nonnegative, and consider the diagram

$$\begin{array}{ccc} X^{[\ell+n+m],[1],[\ell+m]} & \xleftarrow{p_{123}} & X^{[\ell+n+m],[1],[\ell+m],[1],[\ell]} & \xrightarrow{p_{345}} & X^{[\ell+m],[1],[\ell]} \\ & & \downarrow p_{1245} & & \\ & & X^{[\ell+n+m],[1],[1],[\ell]} & & \end{array}$$

Let

$$\begin{aligned} v &:= p_{123}^*[Q^{[\ell+m+n,\ell+m]}] \cdot p_{345}^*[Q^{[\ell+m,\ell]}] \in A_{2\ell+m+n+2}(V), \\ V &:= p_{123}^{-1}(Q^{[\ell+m+n,\ell+m]}) \cap p_{345}^{-1}(Q^{[\ell+m,\ell]}). \end{aligned}$$

According to Lemma 3.10, the operator $d'_n q_m$ is induced by the class

$$w = p_{1245*}(p_{123}^*c_1(\mathcal{I}_{\ell+m+n,\ell+m}) \cdot v) \in A_{2\ell+m+n+1}(W), \quad W := p_{1245}(V).$$

Let $V' \subset V$ and $W' \subset W$ denote the open subsets of those tuples $(\xi, x, \sigma, y, \zeta)$ and (ξ, x, y, ζ) , respectively, where either $x \neq y$ or $x = y$ but ξ_x is curvilinear. Certainly, $V' = p_{1245}^{-1}(W')$, but in fact we even have that $p_{1245} : V' \rightarrow W'$ is an isomorphism: for the conditions imposed on V' imply that σ is already determined by the remaining data (ξ, x, y, ζ) .

Claim: V' is irreducible of dimension $2\ell + n + m + 2$.

For it follows from Briançon's Theorem that the open part $V' \setminus \{x = y\}$ is irreducible of dimension $2\ell + (n + 1) + (m + 1)$, and tuples of the second kind, i.e. (ξ, x, x, ζ) with ξ_x curvilinear, are easily seen to deform into this open subset.

Claim: $\dim(W \setminus W') < 2\ell + m + n + 1$. In particular, the complement of W' in W cannot support any contribution to w .

Indeed, the set $T = \{(\xi, x, x, \zeta) \mid \xi - \zeta = (n + m)x\}$ has a stratification $T = \coprod_{i \geq 0} T_i$, where the stratum T_i is the locally closed set of all tuples with $\text{length}(\zeta_x) = i$. Let $T'_0 \subset T_0$ be the closed subset that consists of tuples where ξ_x is not curvilinear. Then $W \setminus W' \subset T'_0 \cup T_1 \cup T_2 \dots$. Now T_0 is irreducible of dimension $2\ell + (n + m + 1)$, and T'_0 is a proper closed subset and therefore has strictly smaller dimension. The assertion now follows from Lemma 1.8.

Claim: The intersection of $p_{123}^[Q^{\ell+m+n}]$ and $p_{345}^*[Q^{\ell+m,m}]$ is transversal at general points of V' .*

In fact, the intersection is transversal at all points with $x \neq y$ and ξ curvilinear.

We conclude, that the intersection cycle v equals $[\overline{V'}] + r$, where r is a cycle supported on $p_{1245}^{-1}(W \setminus W')$ and therefore irrelevant for our further computations for dimension reasons. Let us return to the definition of the cycle w .

Identifying V' and W' we see that the variety W' parametrises three families

$$Z \subset \Sigma \subset \Xi \subset W' \times X$$

of subschemes in X . In terms of these we can summarise the discussion above by stating that $q'_n q_m$ is induced by the cycle

$$c_1(p_* \mathcal{I}_{\Sigma/\Xi}) \cdot [W'] \in A_*(W').$$

Having reached this point we pause to reflect what changes in this picture if we exchange the order of the operators q_n and q_m . Up to the usual twist T that flips the factors X in $X^{\ell+m+n, [1], [1], [\ell]}$, not a iota is changed in W' . Indeed, W' parametrises not only three but rather four families of subschemes

$$\begin{array}{ccc} & \Sigma' & \\ & \nearrow & \searrow \\ Z & & \Xi \\ & \searrow & \nearrow \\ & \Sigma'' & \end{array}$$

where Σ' and Σ'' are characterised by the property that at a point $s = (\Xi_s, x, y, Z_s) \in W'$ the subschemes $\Sigma'_s, \Sigma''_s \subset \Xi_s$ are the unique ones with

$$\Sigma'_s - Z_s = mx, \quad \Xi_s - \Sigma'_s = ny$$

and

$$\Sigma''_s - Z_s = ny, \quad \Xi_s - \Sigma''_s = mx.$$

This means: the commutator $[q'_n, q_m]$ is induced by the cycle

$$\left(c_1(p_* \mathcal{I}_{\Sigma'/\Xi}) - c_1(p_* \mathcal{I}_{Z/\Sigma''}) \right) \cdot [W'] \in A_{2\ell+n+m+1}(X^{\ell+n+m, [1], [1], [\ell]}).$$

The ideal sheaves corresponding to the various inclusions between the families Z, Σ', Σ'' and Ξ are related by the following commutative diagram of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{I}_{\Sigma'/\Xi} & \longrightarrow & \mathcal{I}_{Z/\Xi} & \longrightarrow & \mathcal{I}_{Z/\Sigma'} & \longrightarrow & 0 \\ & & \varphi \downarrow & & \parallel & & \uparrow & & \\ 0 & \longleftarrow & \mathcal{I}_{Z/\Sigma''} & \longleftarrow & \mathcal{I}_{Z/\Xi} & \longleftarrow & \mathcal{I}_{\Sigma''/\Xi} & \longleftarrow & 0 \end{array}.$$

The homomorphism

$$p_*\varphi : p_*\mathcal{I}_{\Sigma'/\Xi} \rightarrow p_*\mathcal{I}_{Z/\Sigma''}$$

is an isomorphism off the diagonal $\{x = y\} \subset W'$. On the other hand the closure of $W' \cap \{x = y\}$ equals the image of the ‘diagonal’ embedding $Q^{\ell+m+n,\ell} \rightarrow X^{[\ell+m+n],[1],[1],[\ell]}$. It follows that

$$\left(c_1(p_*\mathcal{I}_{\Sigma'/\Xi}) - c_1(p_*\mathcal{I}_{Z/\Sigma''}) \right) \cdot [W'] = -\mu \cdot [Q^{\ell+m+n,\ell}]$$

where μ is the length of $\text{coker}(p_*\varphi)$ at the generic point of the variety $Q^{\ell+m+n,\ell}$. This proves

$$[q'_n(\alpha), q_m(\beta)] = -\mu \cdot q_{n+m}(\alpha\beta),$$

and it remains to show that

$$\mu = nm.$$

A general point $d = (\xi, x, y, \zeta)$ of $Q^{\ell+m+n,\ell}$ is of the form $(\zeta \cup \eta, x, x, \zeta)$ where $\eta \cap \zeta = \emptyset$ and η is a curvilinear subscheme supported at x . As the computation is local in X we may apply the same reduction process as in the proof of Theorem 3.5: we may assume that $\ell = 0$, that $X = \mathbb{A}^2 = \text{Spec}\mathbb{C}[z, w]$, $x = (0, 0)$ and $I_\zeta = (w, z^n)$. Then there is an open neighbourhood of this point d in W' which is isomorphic to $\mathbb{A}^{n+m+2} = \text{Spec}\mathbb{C}[a_0, \dots, a_{n+m-1}, s, t]$ such that the families Ξ, Σ' and Σ'' are given by the ideals

$$I_\Xi = (w - f(z), (z - t)^n(z - s)^m), \quad I_{\Sigma'} = (w - f(z), (z - s)^m)$$

and

$$I_{\Sigma''} = (w - f(z), (z - t)^n),$$

where $f(z) = a_0 + a_1z + \dots + a_{n+m-1}z^{n+m-1}$. We find

$$p_*\mathcal{O}_{\Sigma''} = \mathbb{C}[\underline{a}, s, t][z]/(z - t)^n$$

and

$$p_*\mathcal{I}_{\Sigma'/\Xi} = (z - s)^m \cdot \mathbb{C}[\underline{a}, s, t][z]/(z - s)^m(z - t)^n.$$

The cokernel of

$$p_*\varphi : (z - s)^m \cdot \mathbb{C}[\underline{a}, s, t][z]/(z - s)^m(z - t)^n \longrightarrow \mathbb{C}[\underline{a}, s, t][z]/(z - t)^n$$

is isomorphic to the $\mathbb{C}[\underline{a}, s, t]$ -module

$$\mathbb{C}[\underline{a}, s, t][z]/((z - s)^m, (z - t)^n) \cong \mathbb{C}[\underline{a}, s + t][z - s, z - t]/((z - s)^m, (z - t)^n).$$

This module is supported along the diagonal $\{s = t\}$ (as we expected), and its stalk at the generic point of the diagonal has length nm (as we had to prove).

Step 2: Assume that m is positive and $-m < n < 0$. First one shows as above that the commutator $[q'_n, q_m]$ is induced by cycles in $A_{2\ell+n+m+1}(X^{[\ell+m+n],[1],[1],[\ell]})$

for each $\ell \geq 0$, which are supported on the diagonally embedded varieties $\mathcal{Q}^{\ell+m+n, \ell}$, so that

$$[\mathfrak{q}'_n(\alpha), \mathfrak{q}_m(\beta)] = -c_{n,m} \cdot \mathfrak{q}_{n+m}(\alpha\beta)$$

for certain constants $c_{n,m}$. In order to determine these constants we apply the commutator $[\cdot, \mathfrak{q}_{-n-m}(1)]$. Then the oscillator relations yield for the right hand side

$$-c_{n,m}(n+m) \int_X \alpha\beta \cdot \text{id}_{\mathbb{H}}.$$

On the other hand

$$[[\mathfrak{q}'_n(\alpha), \mathfrak{q}_m(\beta)], \mathfrak{q}_{-n-m}(1)] = [[\mathfrak{q}'_n(\alpha), \mathfrak{q}_{-n-m}(1)], \mathfrak{q}_m(\beta)]$$

Now

$$\begin{aligned} [\mathfrak{q}'_n(\alpha), \mathfrak{q}_{-n-m}(1)] &= (-1)^m [(\mathfrak{q}_{-n}^\dagger)'(\alpha), \mathfrak{q}_{n+m}^\dagger(1)] \\ &= -(-1)^m [\mathfrak{q}_{n+m}(1), \mathfrak{q}'_{-n}(\alpha)]^\dagger, \end{aligned}$$

which by Step 1 equals $(-1)^m n(n+m) \mathfrak{q}_m(\alpha)^\dagger = n(n+m) \mathfrak{q}_{-m}(\alpha)$. Hence

$$\begin{aligned} [[\mathfrak{q}'_n(\alpha), \mathfrak{q}_m(\beta)], \mathfrak{q}_{-n-m}(1)] &= n(n+m) [\mathfrak{q}_{-m}(\alpha), \mathfrak{q}_m(\beta)] \\ &= n(n+m)(-m) \int_X \alpha\beta \cdot \text{id}_{\mathbb{H}}. \end{aligned}$$

Choose classes α, β with $\int_X \alpha\beta \neq 0$. It follows that $c_{n,m} = nm$.

Step 3: The general case can now be reduced formally to the cases already treated. The assertion is certainly trivial if either $n = 0$ or $m = 0$. If the assertion is known to be true for some pair (n, m) , we may apply the operation \dagger to both sides and find:

$$\begin{aligned} [\mathfrak{q}'_{-n}(\alpha), \mathfrak{q}_{-m}(\beta)] &= (-1)^{n+m} [(\mathfrak{q}_n^\dagger)'(\alpha), \mathfrak{q}_m^\dagger(\beta)] \\ &= -(-1)^{n+m} [(\mathfrak{q}'_n)^\dagger(\alpha), \mathfrak{q}_m^\dagger(\beta)] \\ &= (-1)^{n+m} [\mathfrak{q}'_n(\alpha), \mathfrak{q}_m(\beta)]^\dagger = -nm \cdot (-1)^{n+m} \mathfrak{q}_{n+m}^\dagger(\alpha\beta) \\ &= (-n)(-m) \cdot \mathfrak{q}_{-n-m}(\alpha\beta). \end{aligned}$$

This and the identity

$$[\mathfrak{q}'_n(\alpha), \mathfrak{q}_m(\beta)] = (-1)^{|\alpha| \cdot |\beta|} [\mathfrak{q}'_m(\beta), \mathfrak{q}_n(\alpha)]$$

allow us to reduce anything to cases checked in Step 1 and Step 2. \square

In order to prove part 1 of Theorem 3.11, it remains to treat the case $n + m = 0$. This will be done in two steps. First, we prove a qualitative statement about the structure of the ‘correction term’, and afterwards we determine the precise value of the ‘coefficient’ K_n :

Proposition 3.14 — *There exist rational divisors $K_n \in \text{Pic}(X) \otimes \mathbb{Q}$, $n \in \mathbb{Z}$, with $K_0 = 0$ and $K_{-n} = K_n$ and such that*

$$[\mathfrak{q}'_n(\alpha), \mathfrak{q}_{-n}(\beta)] = n^2 \cdot \int_X K_n \alpha \beta \cdot \text{id}_{\mathbb{H}} \quad (10)$$

for all $\alpha, \beta \in H^*(X)$.

Proof. There is nothing to prove for $n = 0$. Moreover,

$$[\mathfrak{q}'_n(\alpha), \mathfrak{q}_{-n}(\beta)] = (-1)^{|\alpha| \cdot |\beta|} \cdot [\mathfrak{q}'_{-n}(\beta), \mathfrak{q}_n(\alpha)].$$

It follows that if there is a divisor K_n so that (10) holds for n , then (10) also holds for $-n$ with the choice $K_{-n} = K_n$. Hence it suffices to prove the proposition for positive integers n .

Let ℓ be a nonnegative integer and consider the diagram

$$\begin{array}{ccccc} X^{[\ell],[1],[\ell+n]} & \xleftarrow{p_{123}} & X^{[\ell],[1],[\ell+n],[1],[\ell]} & \xrightarrow{p_{345}} & X^{[\ell+n],[1],[\ell]} \\ & & \downarrow p_{1245} & & \\ & & X^{[\ell],[1],[1],[\ell]} & & \end{array}$$

Let

$$\begin{aligned} v &:= p_{123}^* [Q^{[\ell,\ell+n]}] \cdot p_{345}^* [Q^{[\ell+n,\ell]}] \in A_{2\ell+2}(V), \\ V &:= p_{123}^{-1}(Q^{[\ell,\ell+n]}) \cap p_{345}^{-1}(Q^{[\ell+n,\ell]}). \end{aligned}$$

According to Lemma 3.10, the operator $\mathfrak{q}'_{-n} \mathfrak{q}_n$ is induced by the class

$$w = (-1)^n p_{1245*} (p_{123}^* c_1(\mathcal{I}_{\ell,\ell+n}) \cdot v) \in A_{2\ell+1}(W), \quad W := p_{1245}(V).$$

Consider the diagonal part $W \cap \{x = y\}$ first. It is contained in $\bigcup_{i \geq 0} T_i$, where $T_i = \{(\xi, x, x, \zeta) \mid \ell(\xi_x) = \ell(\zeta_x) = i\}$. The closure of T_0 is the diagonal $\Delta \cong X^{[\ell]} \times X \subset X^{[\ell],[1],[1],[\ell]}$ and is therefore irreducible of dimension $2\ell + 2$. Whereas for $i \geq 1$, the set T_i embeds into the irreducible variety $X^{[\ell-i]} \times (X_0^{[i]} \times_X X_0^{[i]})$ of dimension $2(\ell - i) + (i + 1) + (i + 1) - 2 = 2\ell$.

The off-diagonal part $W \cap \{x \neq y\}$ is empty if $\ell < n$. If $\ell \geq n$ it has precisely one irreducible component W' of maximal dimension $2\ell + 2$: it contains as a dense subset the image of the embedding

$$\{(\eta, \xi', \zeta') \in X^{[\ell-n]} \times X_0^{[n]} \times X_0^{[n]} \mid \eta, \xi' \text{ and } \zeta' \text{ are pairwise disjoint}\} \longrightarrow W,$$

$$(\sigma, \xi', \zeta') \mapsto (\sigma \cup \xi, \rho(\xi'), \rho(\zeta'), \sigma \cup \zeta').$$

Since the function $(\xi, x, y, \zeta) \mapsto \ell(\xi_x)$ is semicontinuous and is at least n on W' , it follows that $\overline{W'} \cap \Delta$ is contained in $\bigcup_{\nu \geq n} T_\nu$. In particular, this intersection has

dimension $\leq 2\ell$. As we want to compute a cycle of degree $2\ell + 1$, we may restrict our attention to the open part W' and may disregard the complement of W' in its closure.

$p_{1245}^{-1} : p_{1245}^{-1}(W') \rightarrow W'$ is an isomorphism, which we use to identify W' and the off-diagonal part of V . Now W' parametrises four flat families of subschemes on X : besides the families Ξ and Z of fibrewise length ℓ , these are the families $\Xi \cap Z$ and $\Xi \cup Z$ of fibrewise length $\ell - n$ and $\ell + n$. The contribution of W' to w is the class

$$(-1)^n c_1(p_* \mathcal{I}_{\Xi/\Xi \cup Z}) \cdot [W'] \in A_{2\ell+1}(W').$$

Reversing the order of the operators q'_n and q_n shows that the part of the cycle u inducing the commutator $[q'_n, q_n]$, that is supported on W' , is the class

$$(-1)^n \left(c_1(p_* \mathcal{I}_{\Xi/\Xi \cup Z}) - c_1(p_* \mathcal{I}_{\Xi \cap Z/\Xi}) \right) \cdot [W'].$$

Since the ideal sheaves $\mathcal{I}_{\Xi/\Xi \cup Z}$ and $\mathcal{I}_{\Xi/\Xi \cap Z}$ are isomorphic, this class is zero.

Thus we may fully concentrate on the contribution of the diagonal part Δ . (Also note that for the reversed order $q_n q'_n$ any diagonal parts must be contained in $\bigcup_{\nu \geq n} T_\nu$ and are therefore too small and irrelevant.)

The complement of the open subset $T_0 \cong X^{[\ell]} \times X \setminus \Xi_\ell$ in Δ_0 has codimension ≥ 2 . Locally near $p_{1245}^{-1}(T_0)$ there are isomorphisms between $X^{[\ell+n, \ell]}$ and $X^{[\ell]} \times X^{[n]}$, and similarly between $Q^{[\ell+n, \ell]}$ and $X^{[\ell]} \times X_0^{[n]}$. Hence if $\bar{w} \in A_1(X)$ is the intersection cycle for the special case $\ell = 0$, then the general cycle is simply given by $w = [X^{[\ell]}] \times \bar{w} \in A_{2\ell+1}(X^{[\ell]} \times X)$. But that was all we had to prove: a cycle of this form induces the linear map

$$\alpha \otimes \beta \otimes y \mapsto \int_{\bar{w}} \alpha \beta \cdot y, \quad \alpha, \beta \in H^*(X; \mathbb{Q}), y \in \mathbb{H}$$

□

Corollary 3.15 — *For all positive integers n one has*

$$q'_n(\alpha) = n\mathfrak{L}_n(\alpha) + nq_n(K_n \alpha).$$

Proof. Use the same argument as in the first paragraph of the proof of the main theorem after Corollary 3.12. □

To finish the proof of Theorem 3.11 it remains to show:

Proposition 3.16 — *For all positive integers n the rational divisor defined by Proposition 3.14 is given by*

$$K_n = \frac{n-1}{2} K,$$

where K is the canonical class of the surface X .

This will be done in the next section.

3.4 The vertex operator, completion of the proof

Definition 3.17 — Let $\gamma \in H^*(X)$ be an element which is of even degree though not necessarily homogeneous, and let t be a formal parameter. Define operators $S_m(\gamma)$, $m \geq 0$, by

$$S(\gamma, t) := \sum_{m \geq 0} S_m(\gamma) t^m := \exp \left(\sum_{n > 0} \frac{(-1)^{n-1}}{n} \mathfrak{q}_n(\gamma) \cdot t^n \right).$$

Since γ is of even degree by assumption, any two operators $\mathfrak{q}_n(\gamma)$ and $\mathfrak{q}_{n'}(\gamma)$ commute in the ordinary, i.e. ‘ungraded’ sense. In particular, there is no ambiguity in the meaning of the expression on the right hand side in the definition.

The geometric meaning of the operators S_m is explained by the following theorem: let C be a smooth curve in X . There is an induced closed embedding $\mathcal{S}^n C = C^{[n]} \rightarrow X^{[n]}$. Let $[C] \in H^*(X)$ and $[C^{[n]}] \in H^*(X^{[n]})$ be the corresponding cohomology classes, i.e., the Poincaré dual classes of the fundamental classes of these varieties.

Theorem 3.18 (Nakajima, Grojnowski) — *The following relation holds for all non-negative integers n :*

$$[C^{[n]}] = S_n([C]) \cdot \mathbf{1}.$$

For proofs see [22] and [13]. □

Lemma 3.19 — *Let $\gamma \in H^*(X)$ be an element of even degree. Then*

$$S'(\gamma, t) = S(\gamma, t) \cdot \sum_{n > 0} (-1)^{n-1} t^n \left\{ \mathfrak{L}_n(\gamma) + \mathfrak{q}_n \left(\gamma K_n + \gamma^2 \frac{n-1}{2} \right) \right\}.$$

Proof. Assume first that \mathfrak{a} is an operator of even degree, and that $[\mathfrak{d}, \mathfrak{a}]$ commutes with \mathfrak{a} . Then

$$\begin{aligned} \left(\sum_{n=0}^{\infty} \frac{\mathfrak{a}^n}{n!} \right)' &= \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{i=1}^n \mathfrak{a}^{i-1} \cdot \mathfrak{a}' \cdot \mathfrak{a}^{n-i} \\ &= \sum_{n=1}^{\infty} \frac{1}{n!} \cdot \left\{ n \mathfrak{a}^{n-1} \mathfrak{a}' + \sum_{i=1}^n \mathfrak{a}^{n-2} \cdot (n-i) \cdot [\mathfrak{a}', \mathfrak{a}] \right\} \\ &= \sum_{n=0}^{\infty} \frac{\mathfrak{a}^n}{n!} \cdot \mathfrak{a}' + \sum_{n=1}^{\infty} \frac{\mathfrak{a}^{n-2}}{n!} \binom{n}{2} [\mathfrak{a}', \mathfrak{a}] \\ &= \exp(\mathfrak{a}) \cdot \left\{ \mathfrak{a}' + \frac{1}{2} [\mathfrak{a}', \mathfrak{a}] \right\}. \end{aligned}$$

Next, let \mathfrak{a}_ν be a family of commuting operators of even degree such that any $[\mathfrak{a}'_\nu, \mathfrak{a}_\mu]$ commutes with every \mathfrak{a}_ξ . Then it follows from Step 1 and

$$[\mathfrak{a}'_\mu, \exp(\mathfrak{a}_\nu)] = \exp(\mathfrak{a}_\nu) \cdot [\mathfrak{a}'_\mu, \mathfrak{a}_\nu]$$

that

$$\left(\exp \left(\sum_{\nu} \mathfrak{a}_{\nu} \right) \right)' = \exp \left(\sum_{\nu} \mathfrak{a}_{\nu} \right) \cdot \left\{ \sum_{\nu} \mathfrak{a}'_{\nu} + \frac{1}{2} \sum_{\nu, \mu} [\mathfrak{a}'_{\nu}, \mathfrak{a}_{\mu}] \right\}.$$

Now apply this formula to the family $\mathfrak{a}_{\nu} = \frac{(-1)^{\nu-1}}{\nu} \mathfrak{q}_{\nu}(\gamma) t^{\nu}$ and use our previous results $\mathfrak{a}'_{\nu} = (-1)^{\nu-1} t^{\nu} (\mathfrak{L}_n(\gamma) + \mathfrak{q}_{\nu}(K_{\nu} \gamma))$ and $[\mathfrak{a}'_{\nu}, \mathfrak{a}_{\mu}] = -(-t)^{\nu+\mu} \mathfrak{q}_{\nu+\mu}(\gamma^2)$. One gets $S'(\gamma, t) = S(\gamma, t) \cdot (*)$ with

$$\begin{aligned} (*) &= \sum_{n>0} (-1)^{n-1} t^n (\mathfrak{L}_n(\gamma) + \mathfrak{q}_n(K_n \gamma)) - \frac{1}{2} \sum_{\nu, \mu>0} (-t)^{\nu+\mu} \mathfrak{q}_{\nu+\mu}(\gamma^2) \\ &= \sum_{n>0} (-1)^{n-1} t^n \cdot \left\{ \mathfrak{L}_n(\gamma) + \mathfrak{q}_n(K_n \gamma + \frac{1}{2} N_n \gamma^2) \right\} \end{aligned}$$

where N_n is the number of pairs of positive integers ν and μ that add up to n , i.e., $N_n = n - 1$. \square

Let $C \subset X$ be a smooth projective curve. The boundary $\partial X^{[n]}$ intersects $C^{[n]}$ generically transversely in the boundary $\partial C^{[n]}$ of $C^{[n]}$, i.e. in the set of all tuples with multiple points. The subvarieties $X_0^{[n]}$ and $\partial C^{[n]}$ have complementary dimensions $n + 1$ and $n - 1$ in $X^{[n]}$ and we may compute the intersection number

$$I := \int_{X^{[n]}} [X_0^{[n]}] \cup [\partial C^{[n]}].$$

We will do this first using our algorithmic language, and afterwards using a geometric argument. The comparison of the two results will lead to the identification of the divisors K_n .

Lemma 3.20 — $[X_0^{[n]}] = \mathfrak{q}_n(1_X) \cdot \mathbf{1}$ and $[\partial C^{[n]}] = -2 \cdot S'_n([C]) \cdot \mathbf{1}$.

Proof. The first assertion follows from the definition of the operators \mathfrak{q}_n . By Nakajima's Theorem, $S_n([C]) \cdot \mathbf{1}$ is the class of the submanifold $C^{[n]} \subset X^{[n]}$, and hence according to Lemma 3.8:

$$S'_n([C]) \cdot \mathbf{1} = \mathfrak{d} \cdot S_n([C]) \cdot \mathbf{1} = -\frac{1}{2} [\partial X^{[n]}] \cdot [C^{[n]}] = -\frac{1}{2} [\partial C^{[n]}].$$

\square

Lemma 3.21 —

$$\int_{X^{[n]}} (\mathfrak{q}_n(1_X) \cdot \mathbf{1}) \cdot (S'_n([C]) \cdot \mathbf{1}) = \int_X \left\{ n K_n C + \binom{n}{2} C^2 \right\}.$$

Proof. Indeed,

$$\begin{aligned} \int_{X^{[n]}} (\mathfrak{q}_n(1_X) \cdot \mathbf{1}) \cdot (S'_n([C]) \cdot \mathbf{1}) &= (-1)^n \int_{X^{[0]}} \mathfrak{q}_{-n}(1_X) S'_n([C]) \cdot \mathbf{1} \\ &= (-1)^n \int_{X^{[0]}} [\mathfrak{q}_{-n}(1_X), S'_n([C])] \cdot \mathbf{1}, \end{aligned}$$

since $\mathfrak{q}_{-n}(1_X) \cdot \mathbf{1} = 0$. Now \mathfrak{q}_{-n} commutes with any product $\mathfrak{q}_{i_1} \cdots \mathfrak{q}_{i_s}$ if $s \geq 2$, $i_j > 0$ and $\sum_j i_j = n$. Thus the only summand in S'_n that contributes to the commutator with \mathfrak{q}_{-n} is $(-1)^{n-1} \mathfrak{q}_n(C(K_n + C(n-1)/2))$. Hence

$$[\mathfrak{q}_{-n}(1_X), S'_n([C])] = (-1)^n n \int_X C \left(K_n + \frac{n-1}{2} C \right) \cdot \text{id}_{\mathbb{H}}$$

This proves the lemma. \square

Next, we give the geometric computation of I :

Lemma 3.22 —

$$\int_{X^{[n]}} [X_0^{[n]}] \cdot [\partial C^{[n]}] = -n(n-1) \cdot C(C+K).$$

Proof. We have $[X_0^{[n]}] \cdot [\partial C^{[n]}] = [\partial X^{[n]}] \cdot ([X_0^{[n]}] \cdot [C^{[n]}])$. The intersection of $X^{[n]}$ and $C^{[n]}$ is transversal and is equal to the image of the closed immersion $\Delta : C \rightarrow C^{[n]}$ sending a point c to the unique subscheme of C of length n that is supported in c . Thus

$$I = \deg(\mathcal{O}_{X^{[n]}}(\partial X^{[n]})|_{\Delta(C)}) = \deg(\mathcal{O}_{C^{[n]}}(\partial C^{[n]})|_{\Delta(C)}).$$

The embedding Δ factors through the diagonal embedding $C \rightarrow C^n$ and the quotient map $\pi : C^n \rightarrow C^{[n]}$. Moreover, if $\text{pr}_{ij} : C^n \rightarrow C^2$ denotes the projection to the product of the i -th and j -th factor,

$$\pi^*(\mathcal{O}_{C^{[n]}}(\partial C^{[n]})) \cong \left(\bigotimes_{i < j}^n \text{pr}_{ij}^* \mathcal{O}_{C \times C}(\Delta C) \right)^{\otimes 2}.$$

From this we conclude:

$$\begin{aligned} I = \deg(\Delta^* \mathcal{O}_{C^{[n]}}(\partial C^{[n]})) &= 2 \cdot \binom{n}{2} \deg(\mathcal{O}_{C \times C}(\Delta C)|_{\Delta C}) \\ &= -n(n-1) \cdot C(C+K). \end{aligned}$$

\square

Proof of Proposition 3.16. From Lemma 3.20 and Lemma 3.21 we conclude

$$I = (-2) \cdot C(nK_n + \binom{n}{2} C).$$

Comparison with Lemma 3.22 shows that $K_n = \frac{n-1}{2} K$. \square

This finishes the proof of Theorem 3.11.

4 Towards the ring structure of \mathbb{H}

4.1 Tautological sheaves

There is a natural way to associate to a given vector bundle on X a series of tautological' vector bundles on the Hilbert schemes $X^{[n]}$, $n \geq 0$. The Chern classes of the tautological bundles may be grouped together to form operators on \mathbb{H} .

Consider the standard diagram

$$\begin{array}{ccc} \Xi_n & \subset & X^{[n]} \times X \xrightarrow{q} X \\ & & \downarrow p \\ & & X^{[n]} \end{array}$$

Let F be a locally free sheaf on X . For each $n \geq 0$ the associated *tautological bundle* on $X^{[n]}$ is defined as

$$F^{[n]} := p_*(\mathcal{O}_{\Xi_n} \otimes q^*F).$$

Since p is a flat finite morphism of degree n , $F^{[n]}$ is locally free with

$$\mathrm{rk}(F^{[n]}) = n \cdot \mathrm{rk}(F).$$

Note that $F^{[0]} = 0$ and $F^{[1]} = F$.

Furthermore, if $0 \rightarrow F_1 \rightarrow F \rightarrow F_2 \rightarrow 0$ is a short exact sequence of locally free sheaves on X , the corresponding sequence $0 \rightarrow F_1^{[n]} \rightarrow F^{[n]} \rightarrow F_2^{[n]} \rightarrow 0$ is again exact. Hence sending the class $[F]$ of a locally free sheaf F to $[F^{[n]}]$ gives a group homomorphism

$$-^{[n]} : K(X) \longrightarrow K(X^{[n]}).$$

Definition 4.1 — Let u be a class in $K(X)$. Define operators

$$\mathbf{c}(u) \in \mathrm{End}(\mathbb{H}) \quad \text{and} \quad \mathbf{ch}(u) \in \mathrm{End}(\mathbb{H})$$

as follows: For each $n \geq 0$, the action on $H^*(X^{[n]}; \mathbb{Q})$ is given by multiplication with the total Chern class $c(u^{[n]})$ and the Chern character $ch(u^{[n]})$, respectively.

Let

$$\mathbf{c}(u) = \sum_{k \geq 0} \mathbf{c}_k(u) \quad \text{and} \quad \mathbf{ch}(u) = \sum_{k \geq 0} \mathbf{ch}_k(u)$$

be the decompositions into homogeneous components of bidegree $(0, 2k)$. Since all of these operators are of even degree and only act 'vertically' on \mathbb{H} by multiplication, they commute with each other and in particular with the previously defined boundary operator $\mathfrak{d} = \mathbf{c}_1(\mathcal{O}_X)$.

Moreover, we have

$$\mathbf{c}(u + v) = \mathbf{c}(u) \cdot \mathbf{c}(v) \quad \text{and} \quad \mathbf{ch}(u + v) = \mathbf{ch}(u) + \mathbf{ch}(v)$$

for all $u, v \in K(X)$.

Theorem 4.2 — Let u be a class in $K(X)$ of rank r and let $\alpha \in H^*(X)$. Then

$$[\mathrm{ch}(u), \mathfrak{q}_1(\alpha)] = \exp(\mathrm{ad} \mathfrak{d})(\mathfrak{q}_1(\mathrm{ch}(u)\alpha)),$$

or, more explicitly,

$$[\mathrm{ch}_n(u), \mathfrak{q}_1(\alpha)] = \sum_{\nu=0}^n \frac{1}{\nu!} \mathfrak{q}_1^{(\nu)}(\mathrm{ch}_{n-\nu}(u)\alpha).$$

Similarly,

$$\mathfrak{c}(u) \cdot \mathfrak{q}_1(\alpha) \cdot \mathfrak{c}(u)^{-1} = \sum_{\nu, k \geq 0} \binom{r-k}{\nu} \mathfrak{q}_1^{(\nu)}(\mathfrak{c}_k(u)\alpha).$$

Proof. We may assume that u is the class of a locally free sheaf F . Recall the standard diagram for the incidence variety $X^{[\ell, \ell+1]}$:

$$\begin{array}{ccccc} X & \xleftarrow{\rho} & X^{[\ell, \ell+1]} & \xrightarrow{\psi} & X^{[\ell+1]} \\ & & \downarrow \varphi & & \\ & & X^{[\ell]} & & \end{array}$$

The variety $X^{[\ell, \ell+1]}$ parametrises two families of subschemes of X . Their structure sheaves fit into an exact sequence

$$0 \rightarrow \rho_X^* \mathcal{O}_{\Delta_X} \otimes p^* \mathcal{O}_{X^{[\ell, \ell+1]}}(-E) \rightarrow \psi_X^*(\mathcal{O}_{\Xi_{\ell+1}}) \rightarrow \varphi_X^*(\mathcal{O}_{\Xi_\ell}) \rightarrow 0,$$

where $p : X^{[\ell, \ell+1]} \times X \rightarrow X^{[\ell, \ell+1]}$ is the projection and E is the exceptional divisor. Applying the functor $p_*(\cdot \otimes q^* F)$ to this exact sequence yields

$$0 \rightarrow \rho^* F \otimes \mathcal{O}_{X^{[\ell, \ell+1]}}(-E) \rightarrow \psi^* F^{[\ell+1]} \rightarrow \varphi^* F^{[\ell]} \rightarrow 0. \quad (11)$$

Let $\lambda = c_1(\mathcal{O}_{X^{[\ell, \ell+1]}}(-E))$. Then

$$\psi^* \mathrm{ch}(F^{[\ell+1]}) = \varphi^* \mathrm{ch}(F^{[\ell]}) + \rho^* \mathrm{ch}(F) \cdot \exp(\lambda)$$

and

$$\psi^* \mathfrak{c}(F^{[\ell+1]}) = \varphi^* \mathfrak{c}(F) \cdot \sum_{\nu, k \geq 0} \binom{r-k}{\nu} \lambda^\nu \rho^* \mathfrak{c}_k(F).$$

It follows for any $x \in H^*(X^{[\ell]}; \mathbb{Q})$:

$$\begin{aligned} \mathrm{ch}(F) \mathfrak{q}_1(\alpha)(x) &= \mathrm{ch}(F^{[\ell+1]}) \cdot PD^{-1} \psi_*([X^{[\ell, \ell+1]}] \cap \rho^*(\alpha) \varphi^*(x)) \\ &= PD^{-1} \psi_*([X^{[\ell, \ell+1]}] \cap \psi^*(\mathrm{ch}(F^{[\ell+1]}) \rho^*(\alpha) \varphi^*(x))) \\ &= PD^{-1} \psi_*([X^{[\ell, \ell+1]}] \cap \rho^*(\alpha) \varphi^*(\mathrm{ch}(F^{[\ell]}) x)) \\ &\quad + \sum_{\nu \geq 0} \frac{1}{\nu!} PD^{-1} \psi_*(\lambda^\nu \cdot [X^{[\ell, \ell+1]}] \cap \rho^*(\mathrm{ch}(F)\alpha) \varphi^*(x)) \\ &= \mathfrak{q}_1(\alpha)(\mathrm{ch}(F)x) + \sum_{\nu \geq 0} \frac{1}{\nu!} \mathfrak{q}_1^{(\nu)} \mathfrak{q}_1(\mathrm{ch}(F)\alpha)(x). \end{aligned}$$

Here we used Lemma 3.10 which says that the cycle $\mathcal{X} \cdot [X^{\ell, \ell+1}]$ induces the operator $q_1^{(\nu)}$. This is the equation for the Chern character. The equation for the total Chern class is proved analogously. \square

Remark 4.3 — The sequence (11) was used by Ellingsrud in a recursive method to determine Chern classes and Segre classes of tautological bundles (unpublished, but see [25],[4]). He expresses the classes $(\varphi, \rho)_* c(E)$ in terms of the Segre classes of the universal family $\Xi_{[n]} \subset X \times X^{[n]}$. Thus one needs to control the behaviour of these Segre classes under the induction procedure. This method yields qualitative results on the *structure* of certain classes and integrals, but all attempts to get numbers have ended so far in unsurmountable combinatorial difficulties. \square

Remark 4.4 — The results of the present and the previous section provide an algorithmic description of the multiplicative action of the subalgebra $\mathcal{A} \subset \mathbb{H}$ which is generated by the Chern classes of all tautological bundles: The elements $q_1(\alpha_1) \cdot \dots \cdot q_{i_s}(\alpha_s) \cdot \mathbf{1}$ generate \mathbb{H} as a \mathbb{Q} -vector space. By Corollary 3.12, each such element can be written as a linear combination of expression $w \cdot \mathbf{1}$, where w is a word in an alphabet consisting of \mathfrak{d} and operators $q_i(\alpha)$, $\alpha \in H^*(X; \mathbb{Q})$. By Theorem 4.2 the commutator of $\text{ch}(F)$ with any of these is again a word in this alphabet. And finally, Theorem 3.11 shows how such a word can be expressed in terms of the basic operators q_n . Admittedly, without a further understanding of the algebraic structure this description is useful for computations in $H^*(X^{[\ell]}; \mathbb{Q})$ only for small values of ℓ or if one implements it in some computer algebra system.

4.2 The line bundle case

The results of the previous section suffice to compute the Chern classes of the tautological bundles $L^{[n]}$ associated to a line bundle L in terms of the basic operators.

Theorem 4.5 — *Let L be a line bundle on X . Then*

$$\sum_{n \geq 0} c(L^{[n]}) = \exp \left(\sum_{m \geq 1} \frac{(-1)^{m-1}}{m} q_m(c(L)) \right) \cdot \mathbf{1}.$$

Remark 4.6 — Expanding the term on the right hand side, one realises that the cohomological degree of any summand contained in $H^*(X^{[n]}; \mathbb{Q})$ is $\leq 2n$, and, moreover, the maximal degree $2n$ can only be attained if the arguments of all operators q involved have degree 2. In other words, considering elements of top degree only, the equation of the theorem specialises to

$$\sum_{n \geq 0} c_n(L^{[n]}) = \exp \left(\sum_{m \geq 1} \frac{(-1)^{m-1}}{m} q_m(c_1(L)) \right) \cdot \mathbf{1}. \quad (12)$$

This is Nakajima's result 3.18: for suppose $C \subset X$ is a smooth curve and $L = \mathcal{O}_X(C)$. If $\xi \in X^{[n]}$, the natural homomorphism $\mathcal{O}_X \rightarrow \mathcal{O}_\xi(C)$ vanishes if and only if $\xi \subset C$. Hence the vanishing locus of the global vector bundle homomorphism

$$\mathcal{O}_{X^{[n]}} \longrightarrow (\mathcal{O}_X(C))^{[n]} = L^{[n]}$$

is the subvariety $C^{[n]}$. Therefore $[C^{[n]}] = c_n(L^{[n]})$. Inserting this into (12), we recover Nakajima's formula 3.18

$$\sum_{n \geq 0} [C^{[n]}] = \exp \left(\sum_{m \geq 1} \frac{(-1)^{m-1}}{m} \mathfrak{q}_m([C]) \right) \cdot \mathbf{1}$$

Based on this observation, the theorem was conjectured by L. Göttsche in a letter to G. Ellingsrud and the author.

Proof of the theorem. We shall give two variants of the proof which differ slightly in flavour. Observe that the left hand side in the theorem equals

$$\sum_{n \geq 0} c(L^{[n]}) = \mathfrak{c}(L) \cdot \sum_{n \geq 0} 1_{X^{[n]}} = \mathfrak{c}(L) \cdot \exp(\mathfrak{q}_1(1_X)) \cdot \mathbf{1}.$$

Variant 1. Applying Theorem 4.2 with $F = L$ and $r = 1$ we get

$$\mathfrak{c}(L) \cdot \mathfrak{q}_1(1_X) \cdot \mathfrak{c}(L)^{-1} = \{\mathfrak{q}_1(1_X + c_1(L)) + \mathfrak{q}'_1(1_X)\}.$$

Hence

$$\begin{aligned} \mathfrak{c}(L) \cdot \exp(\mathfrak{q}_1(1_X)) \cdot \mathbf{1} &= \mathfrak{c}(L) \cdot \exp(\mathfrak{q}_1(1_X)) \cdot \mathfrak{c}(L)^{-1} \cdot \mathbf{1} \\ &= \exp(\mathfrak{c}(L) \cdot \mathfrak{q}_1(1_X) \cdot \mathfrak{c}(L)^{-1}) \cdot \mathbf{1} \\ &= \exp(\mathfrak{q}_1(c(L)) + \mathfrak{q}'_1(1_X)) \cdot \mathbf{1} \\ &= \sum_{n \geq 0} \frac{1}{n!} (\mathfrak{q}_1(c(L)) + \mathfrak{q}'_1(1_X))^n \cdot \mathbf{1}. \end{aligned}$$

Expanding the right hand side yields summands which are words in the two symbols $\mathfrak{q}_1(c(L))$ and $\mathfrak{q}'_1(1_X)$. Moving all factors $\mathfrak{q}'_1(1_X)$ within a given word as far to the right as possible using the commutation relations of the main theorem we can write

$$\sum_{n \geq 0} \frac{1}{n!} (\mathfrak{q}_1(c(L)) + \mathfrak{q}'_1(1_X))^n = \mathfrak{A} \cdot \mathbf{1} + \mathfrak{B} \cdot \mathfrak{q}'_1(1_X) \cdot \mathbf{1} = \mathfrak{A} \cdot \mathbf{1},$$

where \mathfrak{A} is a sum of expressions of the form

$$\nu_1! \cdots \nu_s! \cdot \frac{(-1)^{\nu_1-1} \mathfrak{q}_{\nu_1}(c(L))}{\nu_1} \cdots \frac{(-1)^{\nu_s-1} \mathfrak{q}_{\nu_s}(c(L))}{\nu_s}.$$

Let $\alpha = (1^{\alpha_1} 2^{\alpha_2} 3^{\alpha_3} \dots)$ denote a partition and let $|\alpha| := \sum_{i \geq 1} i \alpha_i$, and $\alpha! := \prod_i (i!)^{\alpha_i}$. We get

$$\sum_{n \geq 0} \frac{1}{n!} (\mathfrak{q}_1(c(L)) + \mathfrak{q}'_1(1_X))^n \cdot \mathbf{1} = \sum_{\alpha} N_{\alpha} \frac{\alpha!}{|\alpha|!} \prod_{i \geq 1} \left(\frac{(-1)^{i-1} \mathfrak{q}_i(c(L))}{i} \right)^{\alpha_i} \cdot \mathbf{1}, \quad (13)$$

where the natural number N_{α} counts how often the operator

$$\alpha! \prod_{i \geq 1} \left(\frac{(-1)^{i-1} \mathfrak{q}_i(c(L))}{i} \right)^{\alpha_i}$$

arises from a word in $\mathfrak{q}'_1(1_X)$ and $\mathfrak{q}_1(c(L))$ of length $|\alpha|$. It is not difficult to see that N_{α} equals the number of possibilities to partition a set of $|\alpha|$ elements into subsets in such a way that there are α_i subsets of cardinality i . Hence

$$N_{\alpha} := \frac{1}{\alpha_1! \alpha_2! \dots} \cdot \frac{|\alpha|!}{\alpha!}.$$

Inserting this into equation (13) above one gets

$$\begin{aligned} \sum_{n \geq 0} \frac{1}{n!} (\mathfrak{q}_1(c(L)) + \mathfrak{q}'_1(1_X))^n \cdot \mathbf{1} &= \sum_{\alpha} \prod_{i \geq 1} \frac{1}{\alpha_i!} \left(\frac{(-1)^{i-1} \mathfrak{q}_i(c(L))}{i} \right)^{\alpha_i} \cdot \mathbf{1} \\ &= \prod_{i \geq 1} \sum_{\alpha_i \geq 0} \frac{1}{\alpha_i!} \left(\frac{(-1)^{i-1} \mathfrak{q}_i(c(L))}{i} \right)^{\alpha_i} \cdot \mathbf{1} \\ &= \prod_{i \geq 1} \exp \left(\frac{(-1)^{i-1} \mathfrak{q}_i(c(L))}{i} \right) \cdot \mathbf{1} \\ &= \exp \left(\sum_{i \geq 1} \frac{(-1)^{i-1}}{i} \mathfrak{q}_i(c(L)) \right) \cdot \mathbf{1}. \end{aligned}$$

Variant 2. Starting again from the sequence

$$\mathfrak{c}(L) \cdot \mathfrak{q}_1(1_X) = \{\mathfrak{q}_1(1_X + c_1(L)) + \mathfrak{q}'_1(1_X)\} \cdot \mathfrak{c}(L),$$

we multiply by $\frac{1}{n!} \mathfrak{q}_1(1_X)^n t^n$ from the right and sum up over all $n \geq 0$:

$$\begin{aligned} \frac{d}{dt} \left(\mathfrak{c}(L) \cdot \sum_{n \geq 0} \frac{1}{n!} \mathfrak{q}_1(1_X)^n t^n \right) \cdot \mathbf{1} &= \mathfrak{c}(L) \cdot \sum_{n \geq 0} \frac{1}{n!} \mathfrak{q}_1(1_X)^{n+1} t^n \cdot \mathbf{1} \\ &= \{\mathfrak{q}_1(1_X + c_1(L)) + \mathfrak{q}'_1(1_X)\} \cdot \left(\mathfrak{c}(L) \cdot \sum_{n \geq 0} \frac{1}{n!} \mathfrak{q}_1(1_X)^n t^n \right) \cdot \mathbf{1}. \end{aligned}$$

This means that the series

$$\sum_{n \geq 0} \mathfrak{c}(L^{[n]}) t^n = \mathfrak{c}(L) \cdot \exp(\mathfrak{q}_1(1_X) t) \cdot \mathbf{1}$$

satisfies the linear differential equation

$$\frac{d}{dt}\mathfrak{X} = \{\mathfrak{q}_1(1_X + c_1(L)) + \mathfrak{q}'_1(1_X)\} \cdot \mathfrak{X} \quad (14)$$

with initial condition

$$\mathfrak{X}(0) = \mathbf{1}. \quad (15)$$

On the other hand, consider the operator

$$S(c(L), t) = \exp\left(\sum_{m \geq 1} \frac{(-1)^{m-1}}{m} \mathfrak{q}_m(c(L)) t^m\right).$$

We find

$$\frac{d}{dt}S(c(L), t) = S(c(L), t) \cdot \left(\sum_{m \geq 0} (-1)^m \mathfrak{q}_{m+1} t^m\right),$$

and

$$\begin{aligned} & \left[\{\mathfrak{q}_1(1_X + c_1(L)) + \mathfrak{q}'_1(1_X)\}, S(c(L), t)\right] \\ &= S(c(L), t) \cdot \left(\sum_{m \geq 1} \frac{(-1)^{m-1}}{m} [\mathfrak{q}'_1(1_X), \mathfrak{q}_m(c(L))] t^m\right) \\ &= S(c(L), t) \cdot \left(\sum_{m \geq 1} (-1)^m \mathfrak{q}_{m+1}(c(L)) t^m\right). \end{aligned}$$

This shows

$$\begin{aligned} & \{\mathfrak{q}_1(1_X + c_1(L)) + \mathfrak{q}'_1(1_X)\} \cdot S(c(L), t) \cdot \mathbf{1} \\ &= S(c(L), t) \cdot \left(\sum_{m \geq 1} (-1)^m \mathfrak{q}_{m+1}(c(L)) t^m\right) \cdot \mathbf{1} \\ & \quad + S(c(L), t) \cdot \mathfrak{q}_1(c(L)) \cdot \mathbf{1} \\ &= S(c(L), t) \cdot \left(\sum_{m \geq 0} (-1)^m \mathfrak{q}_{m+1}(c(L)) t^m\right) \cdot \mathbf{1} \end{aligned}$$

Hence $S(c(L), t) \cdot \mathbf{1}$ satisfies the system (14) and (15) as well and therefore equals $c(L) \cdot \exp(\mathfrak{q}_1(1_X)t) \cdot \mathbf{1}$. This proves the theorem. \square

4.3 Top Segre classes

The following problem was posed by Donaldson in connection with the computation of instanton invariants: let n be an integer ≥ 1 , and consider a linear system $|H|$ of dimension $3n - 2$ inducing a map $X \dashrightarrow \mathbb{P}^{3n-2}$. A zero-dimensional subscheme $\xi \in X^{[n]}$ does not impose independent conditions on the linear system $|H|$ if the natural homomorphism

$$H^0(\mathbb{P}^{3n-2}, \mathcal{O}_{\mathbb{P}}(1)) \longrightarrow H^0(\xi, \mathcal{O}_{\xi}(H))$$

fails to be surjective. The subscheme of all such $\xi \in X^{[n]}$ has virtual dimension zero, and its class is given by $c_{2n}(W^\vee)$, where W is the virtual vector bundle

$$H^0(\mathbb{P}^{3n-2}, \mathcal{O}_{\mathbb{P}}(H)) \otimes \mathcal{O}_{X^{[n]}} - \mathcal{O}(H)^{[n]}.$$

Thus the number of those ξ that impose dependent conditions is given by

$$N_n := \int_{X^{[n]}} c_{2n}(-\mathcal{O}(H)^{[n]}) = \int_{X^{[n]}} \mathfrak{c}(-\mathcal{O}(H)) \cdot \frac{\mathfrak{q}_1(1_X)^n}{n!} \cdot \mathbf{1}.$$

More explicitly, N_1 is the degree of the linear system, N_2 is the number of double points, N_3 is the number of trisecants to a surface in \mathbb{P}^3 and N_4 is the number of quadrupels of points on a surface in \mathbb{P}^3 that span a plane.

Problem: Express N_n in terms of intrinsic invariants of X such as the degree $d := H.H$, the intersection $\kappa := H.K$ and $\chi := K.K$ and the topological Euler characteristic χ .

Note that even the fact that such an expression in terms of the given invariants exists is not evident *a priori*. This has been proved by Tikhomirov [25]. It also follows immediately from our approach.

Using our algorithm, we can attack this problem as follows. Theorem 4.2 yields for $F = -\mathcal{O}(H)$ and $r = -1$ the formula:

$$\begin{aligned} \mathfrak{c}(-\mathcal{O}(H)) \cdot \mathfrak{q}_1(1_X) \cdot \mathfrak{c}(-\mathcal{O}(H))^{-1} &= \sum_{\nu, k \geq 0} \binom{-1-k}{\nu} \mathfrak{q}_1^{(\nu)}(c_k(-H)) \\ &= \sum_{\nu \geq 0} (-1)^\nu \mathfrak{q}_1^{(\nu)} \left(\sum_{k=0}^{\infty} \binom{\nu+k}{k} (-H)^k \right) \\ &= \sum_{\nu \geq 0} (-1)^\nu \mathfrak{q}_1^{(\nu)}((1-H+H^2)^{\nu+1}). \end{aligned}$$

Denote the operator sum on the right hand side by \mathfrak{N} . It follows as in the proof of Theorem 4.5 that $\mathfrak{c}(-\mathcal{O}(H)) \cdot \exp(\mathfrak{q}_1(1_X)t) \cdot \mathbf{1}$ satisfies the following differential equation and initial value condition:

$$\frac{d}{dt} \mathfrak{X} = \mathfrak{N} \mathfrak{X} \quad \text{and} \quad \mathfrak{X}(0) = \mathbf{1}.$$

As long as no explicit generating function is available we must be content with the following semi-explicit solution to Donaldson's problem:

$$N_n = \frac{1}{n!} \int_{X^{[n]}} \mathfrak{N}^n \cdot \mathbf{1}.$$

Note that the right hand side is more than a mere reformulation of the definition of N_n : the expression on the right hand side is a linear combination of words in the operators q_1 and \mathfrak{d} and can be explicitly evaluated by applying the rules of Theorem 3.1.1.

Example 4.7 — As a special case, let us compute N_2 . This is the number of secant lines to an embedded surface in \mathbb{P}^5 that pass through a fixed but general point $x \in \mathbb{P}^5$. Hence we should find Severi's double point formula [23] (see also [2]). We have

$$2 \cdot N_2 = \int_{X^{[2]}} \left(\sum_{n \geq 0} (-1)^n q_1^{(n)} (1 - (n+1)H + \binom{n+2}{2} H^2) \right)^2 \cdot \mathbf{1}.$$

Since $q_1^{(n)}(\alpha) \cdot \mathbf{1} = 0$ for all $n > 0$ and for all α , and for degree reasons the integral reduces to

$$2 \cdot N_2 = \int_{X^{[2]}} I$$

with

$$\begin{aligned} I &= \left(q_1^2(H^2 \otimes H^2) + q_1' q_1(2H^2 \otimes H + 3H \otimes H^2) \right. \\ &\quad \left. + q_1'' q_1(6H^2 \otimes 1 + 3H \otimes H + 1 \otimes H^2) \right. \\ &\quad \left. + q_1''' q_1(4H \otimes 1 + 1 \otimes H) + q_1'''' q_1(1 \otimes 1) \right) \cdot \mathbf{1}. \end{aligned}$$

Since $q_1'(\alpha) \cdot \mathbf{1} = 0$, one easily sees that

$$q_1^{(n)} q_1(\alpha \otimes \beta) \cdot \mathbf{1} = -q_2^{(n-1)}(\alpha\beta) \cdot \mathbf{1}.$$

This yields

$$I = (q_1^2(H^2 \otimes H^2) - q_2'(10H^2) - q_2''(5H) - q_2'''(1)) \cdot \mathbf{1}.$$

The term q_2 vanishes for degree reasons. Moreover, for $n \geq 2$ we have

$$\begin{aligned} q_2^{(n)}(\alpha) \cdot \mathbf{1} &= (q_1^2(\delta_*(\alpha)) + q_2(K\alpha))^{(n-1)} \cdot \mathbf{1} \\ &= (-q_2^{(n-2)}(c_2(X)\alpha) + q_2^{(n-1)}(K\alpha)) \cdot \mathbf{1}. \end{aligned}$$

(Note that the composite map $H^*(X) \xrightarrow{\delta} H^*(X) \otimes H^*(X) \xrightarrow{\cup} H^*(X)$ is the multiplication with the self intersection of the diagonal, i.e. the second Chern class $c_2(X)$ of X .) Applying this to I , we find

$$\begin{aligned} I - q_1^2(H^2 \otimes H^2) \cdot \mathbf{1} &= -(10q_2'(H^2) + 5q_2''(H) + q_2'''(1)) \cdot \mathbf{1} \\ &= -(q_2(10H^2 - c_2(X)) + q_2''(5H + K)) \cdot \mathbf{1} \\ &= -q_2'(10H^2 - c_2(X) + 5HK + K^2) \cdot \mathbf{1}. \end{aligned}$$

This yields:

$$I = q_1^2(H^2 \otimes H^2) + \delta_*(-10H^2 + c_2(X) - 5HK - K^2) \cdot \mathbf{1},$$

and therefore

$$2 \cdot N_2 = \int_{X^{[2]}} I = d^2 - 10d - 5\pi - \kappa + \chi.$$

□

Obviously, for higher n , the practical calculation of N_n quickly becomes rather difficult. Already the case of N_3 surpassed my personal calculation skills. Using MAPLE, I computed the following expressions:

$$\begin{aligned} 3! \cdot N_3 &= d^3 - 30d^2 + 224d - 3d(5\pi + \kappa - \chi) \\ &\quad + 192\pi + 56\kappa - 40\chi, \end{aligned}$$

$$\begin{aligned} 4! \cdot N_4 &= d^4 - 60d^3 + d^2(1196 - 30\pi + 6\chi - 6\kappa) \\ &\quad - d(7920 - 1068\pi + 220\chi - 284\kappa) + 3\chi^2 + 1944\chi - 6\chi\kappa \\ &\quad - 30\chi\pi + 75\pi^2 + 3\kappa^2 + 30\kappa\pi - 9042\pi - 3300\kappa, \end{aligned}$$

$$\begin{aligned} 5! \cdot N_5 &= d^5 - 100d^4 + d^3(3740 + 10\chi - 50\pi - 10\kappa) \\ &\quad - d^2(62000 - 3420\pi + 700\chi - 860\kappa) + d(384384 + 15\chi^2 \\ &\quad + 15960\chi - 30\chi\kappa - 150\pi\chi + 15\kappa^2 + 150\kappa\pi - 75610\pi \\ &\quad - 24340\kappa + 375\pi^2) - 400\chi^2 - 117120\chi + 3920\pi\chi + 960\kappa\chi \\ &\quad + 226560\kappa - 4720\kappa\pi - 560\kappa^2 + 530880\pi - 9600\pi^2. \end{aligned}$$

$$\begin{aligned} 6! \cdot N_6 &= d^6 - 150d^5 + d^4(8980 - 15\kappa + 15\chi - 75\pi) \\ &\quad - d^3(268200 - 2020\kappa + 1700\chi - 8340\pi) \\ &\quad + d^2(3996064 + 45\chi^2 + 71100\chi - 90\chi\kappa - 450\chi\pi \\ &\quad + 450\kappa\pi + 1125\pi^2 - 101040\kappa + 45\kappa^2 - 340530\pi) \\ &\quad - d(23761920 + 2850\chi^2 + 1292320\chi - 28020\chi\pi \\ &\quad - 6660\chi\kappa + 3810\kappa^2 + 32820\kappa\pi - 5995740\pi \\ &\quad - 2224040\kappa + 68850\pi^2) + 15\chi^3 + \chi^2(45160 - 45\kappa - 225\pi) \\ &\quad + \chi(8517120 + 1125\pi^2 + 450\kappa\pi - 435030\pi - 123460\kappa + 45\kappa^2) \\ &\quad - 18151200\kappa + 598170\kappa\pi - 1875\pi^3 - 37768560\pi - 1125\kappa\pi^2 \\ &\quad - 15\kappa^3 + 1046790\pi^2 - 225\kappa^2\pi + 80860\kappa^2 \end{aligned}$$

These calculations verify LeBarz' trisecant formula for N_3 [19, Théorème 8] and the computation of N_4 by Tikhomirov and Troshina [26]. The formulae for N_5 and

N_6 seem to be new. I omit the presentation of N_7 : the information is contained in the following analysis of these numerical data.

It is always possible to organise these data into the following form:

$$\sum_{n \geq 0} N_n(-z)^n = \exp \left(- \sum_{m > 0} \frac{z^m}{m} d_m \right).$$

What is surprising is that the polynomials d_m in the variables d , π , κ , and χ should depend *linearly* on the *three* expressions d , $\pi_0 := \pi - 2\kappa$, and $\chi_0 := \chi - 11\kappa$. This holds for $m \leq 7$ according to the computations above, which imply that

$$\begin{aligned} d_1 &= d \\ d_2 &= 10d + 5\pi_0 - \chi_0 \\ d_3 &= 112d + 96\pi_0 - 20\chi_0 \\ d_4 &= 1320d + 1507\pi_0 - 324\chi_0 \\ d_5 &= 16016d + 22120\pi_0 - 4880\chi_0 \\ d_6 &= 198016d + 314738\pi_0 - 70976\chi_0 \\ d_7 &= 2480640d + 4402720\pi_0 - 1012032\chi_0, \end{aligned}$$

and it is only natural to conjecture that this holds in general. Observe also that the sequence of coefficients of χ_0 seems to be the square of the sequence of coefficients of d . More precisely:

Conjecture 4.8 — *Let $f(z) := \sum_{m > 0} 2^{m-2} \binom{3m-1}{m} z^m$. Then there is a power series $g(z) := \sum_{m > 1} (3m-1)\beta_m z^m$ with positive integral coefficients such that*

$$-z \frac{d}{dz} \log \left(\sum_{n \geq 0} N_n(-z)^n \right) = f(z)d + g(z)\pi_0 - f(z)^2 \chi_0. \quad (16)$$

□

The fact that the right hand side in (16) depends linearly on χ_0 can be proved by the methods in the forthcoming paper [4].

We thank Don Zagier for pointing out to us the existence of Sloane's 'Encyclopedia of Integer Sequences' [24]. We had had reasons to believe that the sequence of coefficients of d be divisible by the binomial coefficients $\binom{3m-1}{2}$. After dividing by these, we are left with the sequence 1, 1, 4, 24, 176, 1456. A search for this reduced sequence in the encyclopedia was successful and led to the above given (conjectural) identification of the coefficients of f . Unfortunately, the corresponding 'reduced' sequence of coefficients β_m of π_0 remains mysterious:

$$0, 1, 12, 137, 1580, 18514, 220136 \dots$$

References

- [1] J. Briançon, *Description de $\text{Hilb}^n \mathbb{C}\{x, y\}$* . Inventiones math. 41, 45-89 (1977).
- [2] F. Catanese, *On Severi's proof of the double point formula*. Comm. in Algebra 7 (1979), 763-773.
- [3] J. Cheah, *On the Cohomology of Hilbert Schemes of Points*. J. Alg. Geom. 5 (1996), 479-511.
- [4] G. Ellingsrud, L. Göttsche, M. Lehn, *On the cobordism class of Hilbert schemes of Points on Surfaces*. In preparation.
- [5] G. Ellingsrud, A. Strømme, *On the homology of the Hilbert schemes on points in the plane*. Inv. Math. 87 (1987), 343-352.
- [6] G. Ellingsrud, A. Strømme, *Towards the Chow ring of the Hilbert scheme of \mathbb{P}^2* . J. reine angew. Math. 441 (1993), 33-44.
- [7] G. Ellingsrud, A. Strømme, *An intersection number for the punctual Hilbert scheme of a surface*. Trans. Amer. Math. Soc. to appear.
- [8] B. Fantechi, L. Göttsche, *The cohomology ring of the Hilbert scheme of 3 points on a smooth projective variety*. J. reine angew. Math. 439 (1993), 147-158,
- [9] J. Fogarty, *Algebraic Families on an Algebraic Surface*. Am. J. Math. 10, 511-521 (1968).
- [10] W. Fulton, *Intersection Theory*. Erg. Math. (3. Folge) Band 2, Springer Verlag 1984.
- [11] L. Göttsche, *The Betti numbers of the Hilbert scheme of points on a smooth projective surface*. Math. Ann. 286 (1990), 193-207.
- [12] L. Göttsche, W. Sörgel, *Perverse sheaves and the cohomology of Hilbert schemes of smooth algebraic surfaces*. Math. Ann. 296 (1993), 235-245.
- [13] I. Grojnowski, *Instantons and Affine Algebras I: The Hilbert Scheme and Vertex Operators*. Math. Res. Letters 3 (1996), 275-291.
- [14] A. Grothendieck, *Techniques de construction et théorèmes d'existence en géométrie algébrique IV: Les schémas de Hilbert*. Séminaire Bourbaki, 1960/61, no. 221.
- [15] A. Grothendieck, *Sur quelques points d'algèbre homologique*. Tôhoku Math. J. 9 (1957), 119-221.

- [16] D. Huybrechts, M. Lehn, *The geometry of moduli spaces of coherent sheaves*. Aspects of Mathematics, Vol. E 31. Vieweg Verlag, 1997.
- [17] A. Iarrobino, *Punctual Hilbert Schemes*. Memoirs of the AMS, Volume 10, Number 188, 1977.
- [18] B. Iversen, *Linear determinants with applications to the Picard scheme of a family of algebraic curves*. Lect. Notes Math. 174, Springer Verlag, Berlin (1970).
- [19] P. LeBarz, *Formules pour les trisécantes des surfaces algébriques*. L'Ens. Math. 33 (1987), 1-66.
- [20] I. G. Macdonald, *The Poincaré Polynomial of a Symmetric Product*. Proc. Cambridge Phil. Soc. 58 (1962), 563-568.
- [21] H. Nakajima, *Heisenberg algebra and Hilbert schemes of points on projective surfaces*. Ann. Math. 145 (1997), 379-388.
- [22] H. Nakajima, *Lectures on Hilbert schemes of points on surfaces*. Preprint, University of Tokyo, 1996.
- [23] F. Severi, *Sulle intersezioni delle varietà algebriche e sopra i loro caratteristiche singolarità proiettive*. Mem. Accad. Scienze di Torino, S. II 52 (1902), 61-118. Also in: Memorie Scelte, I, Zuffi (1950), Bologna.
- [24] N. J. A. Sloane, S. Plouffe, *The Encyclopedia of Integer Sequences*. Academic Press, San Diego, 1995. On-line version: http://www.research.att.com/_njas/sequences.
- [25] A. S. Tikhomirov, *Standard bundles on a Hilbert scheme of points on a surface*. In: Algebraic geometry and its applications, Yaroslavl', 1992. Aspects of Mathematics, Vol. E25. Vieweg Verlag, 1994.
- [26] A. S. Tikhomirov, T. L. Troshina, *Top Segre class of a standard vector bundle \mathcal{E}_D^4 on the Hilbert scheme $\text{Hilb}^4(S)$ of a surface S* . In: Algebraic geometry and its applications, Yaroslavl', 1992. Aspects of Mathematics, Vol. E25. Vieweg Verlag, 1994.
- [27] C. Vafa, E. Witten, *A strong coupling test of S-duality*. Nucl. Phys. 431 (1994), 3-77.

Manfred Lehn
 Mathematisches Institut der Georg-August-Universität
 Bunsenstr. 3-5, D-37073 Göttingen, Germany
 e-mail: lehn@uni-math.gwdg.de