

## Moments of Sums

STEVEN FINCH

April 23, 2004

Let  $X_1, X_2, \dots, X_n$  be a sequence of independent random variables. A huge amount of work has been done on estimating the  $L_p$ -norm of the sum of the  $X$ s:

$$\left\| \sum_{k=1}^n X_k \right\|_p = \left\{ \mathbb{E} \left( \left| \sum_{k=1}^n X_k \right|^p \right) \right\}^{1/p}, \quad p > 0.$$

We first discuss Khintchine's inequality [1], which deals with the Rademacher sequence  $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ , where

$$\mathbb{P}(\varepsilon_k = 1) = \mathbb{P}(\varepsilon_k = -1) = 1/2 \quad (\text{symmetric Bernoulli distribution})$$

for each  $k$ . It is known that there exist constants  $A_p, B_p$  such that the bounds

$$A_p \left( \sum_{k=1}^n c_k^2 \right)^{1/2} \leq \left\| \sum_{k=1}^n c_k \varepsilon_k \right\|_p \leq B_p \left( \sum_{k=1}^n c_k^2 \right)^{1/2}$$

hold for arbitrary  $c_1, c_2, \dots, c_n \in \mathbb{R}$  and  $n \geq 1$ . Szarek [2] and Haagerup [3], building on [4, 5, 6, 7, 8, 9], proved that the best such constants are

$$A_p = \begin{cases} \|W\|_p & \text{if } 0 < p \leq p_0 \\ \|Z\|_p & \text{if } p_0 < p < 2 \\ 1 & \text{if } 2 \leq p < \infty \end{cases} = \begin{cases} 2^{1/2-1/p} & \text{if } 0 < p \leq p_0 \\ 2^{1/2} \left( \frac{\Gamma((p+1)/2)}{\sqrt{\pi}} \right)^{1/p} & \text{if } p_0 < p < 2 \\ 1 & \text{if } 2 \leq p < \infty \end{cases},$$
$$B_p = \begin{cases} 1 & \text{if } 0 < p \leq 2 \\ \|Z\|_p & \text{if } 2 < p < \infty \end{cases} = \begin{cases} 1 & \text{if } 0 < p \leq 2 \\ 2^{1/2} \left( \frac{\Gamma((p+1)/2)}{\sqrt{\pi}} \right)^{1/p} & \text{if } 2 < p < \infty \end{cases}$$

where  $W = 2^{-1/2}(\varepsilon_1 + \varepsilon_2)$ ,  $Z$  is Normal(0, 1), and  $p_0 = 1.8474163360\dots$  is the unique solution of the equation

$$\Gamma\left(\frac{p+1}{2}\right) = \frac{\sqrt{\pi}}{2}$$

---

<sup>0</sup>Copyright © 2004 by Steven R. Finch. All rights reserved.

in the interval  $0 < p < 2$ . In words, if  $\sum_{k=1}^n c_k^2 = 1$ , then  $A_1 = 2^{-1/2}$  and  $B_1 = 1$  encompass the average of  $|\pm c_1 \pm c_2 \pm \cdots \pm c_n|$  taken over all  $2^n$  possible choices of signs. See also [10, 11, 12, 13, 14, 15].

A complex analog of Khintchine's inequality deals with the Steinhaus sequence  $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ , where  $\varepsilon_k$  is uniformly distributed on the unit circle  $\{z : |z| = 1\}$  for each  $k$ . We keep notation identical to before, except that we allow  $c_1, c_2, \dots, c_n \in \mathbb{C}$ . The best constants  $A_p, B_p$  in the inequality

$$A_p \left( \sum_{k=1}^n |c_k|^2 \right)^{1/2} \leq \left\| \sum_{k=1}^n c_k \varepsilon_k \right\|_p \leq B_p \left( \sum_{k=1}^n |c_k|^2 \right)^{1/2}$$

were conjectured by Haagerup [16] to be

$$A_p = \begin{cases} \|W\|_p & \text{if } 0 < p \leq p_0 \\ \|Z\|_p & \text{if } p_0 < p < 2 \\ 1 & \text{if } 2 \leq p < \infty \end{cases} = \begin{cases} 2^{1/2} \left( \frac{\Gamma((p+1)/2)}{\sqrt{\pi} \Gamma((p+2)/2)} \right)^{1/p} & \text{if } 0 < p \leq p_0 \\ (\Gamma((p+2)/2))^{1/p} & \text{if } p_0 < p < 2 \\ 1 & \text{if } 2 \leq p < \infty \end{cases},$$

$$B_p = \begin{cases} 1 & \text{if } 0 < p \leq 2 \\ \|Z\|_p & \text{if } 2 < p < \infty \end{cases} = \begin{cases} 1 & \text{if } 0 < p \leq 2 \\ (\Gamma((p+2)/2))^{1/p} & \text{if } 2 < p < \infty \end{cases}$$

where  $W = 2^{-1/2}(\varepsilon_1 + \varepsilon_2)$ ,  $Z = 2^{-1/2}(U + iV)$  with  $U, V$  independent and Normal(0, 1), and  $p_0 = 0.4756170089\dots$  is the unique solution of the equation

$$2^{p/2} \Gamma\left(\frac{p+1}{2}\right) = \sqrt{\pi} \left( \Gamma\left(\frac{p+2}{2}\right) \right)^2$$

in the interval  $0 < p < 2$ . Here, if  $\sum_{k=1}^n |c_k|^2 = 1$ , then  $A_1 = \sqrt{\pi}/2$  and  $B_1 = 1$  encompass an average taken over all "complex signs" rather than only "real signs" as earlier. Sawa [17] announced that he could verify significant portions of Haagerup's conjecture, but only the case  $p \approx 1$  was published. See also [14, 15, 18, 19]. We mention as well the following result [20, 21] for which  $p = 1$  and  $n$  is the parameter of interest:

$$\mathbb{E} \left( \left| \sum_{k=1}^n \varepsilon_k \right| \right) = \begin{cases} \frac{2}{\pi} \int_0^\infty \frac{1 - \cos(t)^n}{t^2} dt & \text{for the real case} \\ \int_0^\infty \frac{1 - J_0(t)^n}{t^2} dt & \text{for the complex case} \end{cases}$$

where  $J_0(t)$  is the zeroth Bessel function of the first kind. On the one hand, we have

$$\frac{2}{\pi} \int_0^\infty \frac{1 - \cos(t)^n}{t^2} dt = \frac{n!}{2^{n-1} m! (n-m-1)!}$$

for the real case, where  $m = \lfloor (n-1)/2 \rfloor$ . On the other hand, the Bessel integral takes on the values 1,  $4/\pi$ , 1.57459723... and 1.79909248... for  $n = 1, 2, 3$  and 4. Keane [22] recently determined that the third value in this list has the following closed-form expression:

$$\frac{1}{8\pi^3} \Gamma\left(\frac{1}{6}\right)^2 \Gamma\left(\frac{1}{3}\right)^2 + 48\pi \Gamma\left(\frac{1}{6}\right)^{-2} \Gamma\left(\frac{1}{3}\right)^{-2} = 1.5745972375\dots$$

but the fourth value still remains open.

We next discuss Rosenthal's inequalities [23]:

$$\left\| \sum_{k=1}^n X_k \right\|_p \leq C_p \cdot \max \left\{ \left( \sum_{k=1}^n \|X_k\|_p^p \right)^{1/p}, \left\| \sum_{k=1}^n X_k \right\|_1 \right\}, \quad p \geq 1$$

for nonnegative random variables and

$$\left\| \sum_{k=1}^n X_k \right\|_p \leq D_p \cdot \max \left\{ \left( \sum_{k=1}^n \|X_k\|_p^p \right)^{1/p}, \left\| \sum_{k=1}^n X_k \right\|_2 \right\}, \quad p \geq 2$$

for symmetric random variables (meaning that the distribution of  $-X$  is the same as the distribution of  $X$ ). A variation of the latter inequality arises if we loosen the restrictive hypothesis "symmetric" to "zero mean"; the constant is then denoted  $E_p$  rather than  $D_p$ . Johnson, Schechtman & Zinn [24] showed that the growth rate of the best constants  $C_p, D_p, E_p$  is  $p/\ln(p)$  as  $p \rightarrow \infty$ ; in contrast, the growth rate for  $B_p$  is only  $\sqrt{p}$ . Subsequent work [25, 26, 27, 28] yielded that

$$C_p = \begin{cases} 1 & \text{if } p = 1 \\ 2^{1/p} & \text{if } 1 < p < 2 \\ \|Q\|_p & \text{if } 2 \leq p < \infty \end{cases}, \quad D_p = \begin{cases} 1 & \text{if } p = 2 \\ (1 + \|Z\|_p^p)^{1/p} & \text{if } 2 < p < 4 \\ \|R - S\|_p & \text{if } 4 \leq p < \infty \end{cases}$$

where  $Q$  is Poisson(1),  $Z$  is Normal(0,1), and  $R, S$  are independent Poisson(1/2) variables. It is known that  $\|Q\|_m^m = \alpha_m$  and  $\|R - S\|_{2m}^{2m} = \beta_m$  for integer  $m$ , where  $\{\alpha_m\}_{m=1}^{\infty} = \{1, 2, 5, 15, 52, 203, \dots\}$  is the sequence of Bell numbers [29, 30]

$$\alpha_m = \frac{1}{e} \sum_{j=0}^{\infty} \frac{j^m}{j!} = \frac{d^m}{dx^m} \exp(\exp(x) - 1) \Big|_{x=0}$$

and  $\{\beta_m\}_{m=1}^{\infty} = \{1, 4, 31, 379, \dots\}$  is the sequence

$$\beta_m = \frac{2}{e} \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \frac{k^{2m}}{j!(j+k)!2^{2j+k}} = \frac{d^{2m}}{dx^{2m}} \exp(\cosh(x) - 1) \Big|_{x=0}.$$

Ibragimov & Sharakhmetov [31] conjectured that

$$E_p = \begin{cases} (1 + \|Z\|_p^p)^{1/p} & \text{if } 2 < p < 4 \\ \|Q - 1\|_p & \text{if } 4 \leq p < \infty \end{cases}$$

and proved that this is true when  $p = 2m$ ; further,  $\|Q - 1\|_{2m}^{2m} = \gamma_m$  and  $\{\gamma_m\}_{m=1}^\infty = \{1, 4, 41, 715, \dots\}$  is the sequence

$$\gamma_m = \frac{1}{e} \sum_{j=0}^{\infty} \frac{(j-1)^{2m}}{j!} = \frac{d^{2m}}{dx^{2m}} \exp(\exp(x) - x - 1) \Big|_{x=0}.$$

Combinatorial interpretations apply for each of the three sequences:  $\alpha_n$  is the number of partitions of an  $n$ -element set into blocks;  $\beta_n$  is the number of partitions of a  $2n$ -element set into blocks, each containing an even number of elements; and  $\gamma_n$  is the number of partitions of a  $2n$ -element set into blocks, each containing more than one element [30].

Define the following Orlicz-type norm:

$$[\Xi]_p = \inf \left\{ \lambda > 0 : \prod_{k=1}^{\infty} \mathbb{E} \left( \left| 1 + \frac{X_k}{\lambda} \right|^p \right) \leq e^p \right\}$$

for an arbitrary sequence  $\Xi = \{X_k\}_{k=1}^\infty$  of independent random variables, for any  $p > 0$ . We mention Latała's inequality [32]:

$$\frac{e-1}{2e^2} \cdot [\Xi]_p \leq \left\| \sum_{k=1}^{\infty} X_k \right\|_p \leq e \cdot [\Xi]_p$$

which holds either if all the  $X$ s are nonnegative and  $p \geq 1$ , or if all the  $X$ s are symmetric and  $p \geq 2$ . Observe here that the bounds do not depend on  $p$ , unlike the earlier inequalities. For the nonnegative case, Hitczenko & Montgomery-Smith [33] improved the left-hand constant  $(e-1)/(2e^2) = 0.116272\dots$  to  $\xi = 0.154906\dots$ , where  $\xi$  is the unique positive solution of the equation

$$\sum_{k=0}^{\infty} \frac{(2k+1)^k}{k!} x^k = e.$$

It is not known if this improvement carries over to the symmetric case, nor whether a calculation of best constants is feasible at present.

## REFERENCES

- [1] A. Khintchine, Über dyadische Brüche, *Math. Z.* 18 (1923) 109–116.
- [2] S. J. Szarek, On the best constants in the Khinchin inequality, *Studia Math.* 58 (1976) 197–208; MR0430667 (55 #3672).
- [3] U. Haagerup, The best constants in the Khintchine inequality, *Studia Math.* 70 (1981) 231–283; MR0654838 (83m:60031).
- [4] J. E. Littlewood, On bounded bilinear forms in an infinite number of variables, *Quart. J. Math.* 1 (1930) 164–174.
- [5] R. R. Hall, On a conjecture of Littlewood, *Math. Proc. Cambridge Philos. Soc.* 78 (1975) 443–445; MR0404151 (53 #7954).
- [6] P. Whittle, Bounds for the moments of linear and quadratic forms in independent random variables (in Russian), *Teor. Veroyatnost. i Primenen.* 5 (1960) 331–335; Engl. transl. in *Theory Probab. Appl.* 5 (1960) 302–305; MR0133849 (24 #A3673).
- [7] S. B. Steckin, On the best lacunary system of functions (in Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* 25 (1961) 357–366; MR0131097 (24 #A951).
- [8] M. L. Eaton, A note on symmetric Bernoulli random variables, *Annals of Math. Statist.* 41 (1970) 1223–1226; MR0268930 (42 #3827).
- [9] R. M. G. Young, On the best possible constants in the Khintchine inequality, *J. London Math. Soc.* 14 (1976) 496–504; MR0438089 (55 #11008).
- [10] B. Tomaszewski, A simple and elementary proof of the Khintchine inequality with the best constant, *Bull. Sci. Math.* 111 (1987) 103–109; MR0886960 (89b:42004).
- [11] R. Komorowski, On the best possible constants in the Khintchine inequality for  $p \geq 3$ , *Bull. London Math. Soc.* 20 (1988) 73–75; MR0916079 (89e:60037).
- [12] F. L. Nazarov and A. N. Podkorytov, Ball, Haagerup, and distribution functions, available online at <http://www.math.msu.edu/~fedja/prepr.html>.
- [13] J. Wissel and J. Turian, On the best constants in the Khintchine inequality, available online at <http://www.ai.mit.edu/people/jude/research/khintchine/paper.html>.

- [14] G. Peshkir and A. N. Shiryaev, Khinchin inequalities and a martingale extension of the sphere of their action (in Russian), *Uspekhi Mat. Nauk*, v. 50 (1995) n. 5, 3–62; Engl. transl. in *Russian Math. Surveys* 50 (1995) 849–904; MR1365047 (96k:60038).
- [15] A. Baernstein and R. C. Culverhouse, Majorization of sequences, sharp vector Khinchin inequalities, and bisubharmonic functions, *Studia Math.* 152 (2002) 231–248; MR1916226 (2003d:60040).
- [16] A. Pełczyński, Norms of classical operators in function spaces, *Astérisque* 131 (1985) 137–162; MR0816744 (87b:47036).
- [17] J. Sawa, The best constant in the Khintchine inequality for complex Steinhaus variables, the case  $p = 1$ , *Studia Math.* 81 (1985) 107–126; MR0818175 (87d:26024).
- [18] G. Peškir, Best constants in Kahane-Khintchine inequalities for complex Steinhaus functions, *Proc. Amer. Math. Soc.* 123 (1995) 3101–3111; MR1283561 (95m:46037).
- [19] H. König and S. Kwapien, Best Khintchine type inequalities for sums of independent, rotationally invariant random vectors, *Positivity* 5 (2001) 115–152; <http://analysis.math.uni-kiel.de/koenig/preprints.html>; MR1825172 (2002a:60023).
- [20] B. Grünbaum, Projection constants, *Trans. Amer. Math. Soc.* 95 (1960) 451–465; MR0114110 (22 #4937).
- [21] H. König, C. Schütt and N. Tomczak-Jaegermann, Projection constants of symmetric spaces and variants of Khintchine’s inequality, *J. Reine Angew. Math.* 511 (1999) 1–42; <http://analysis.math.uni-kiel.de/koenig/preprints.html>; MR1695788 (2000i:46014).
- [22] J. Keane, Exact value of an integral involving Bessel  $J_0$ , unpublished note (2004).
- [23] H. P. Rosenthal, On the subspaces of  $L^p$  ( $p > 2$ ) spanned by sequences of independent random variables, *Israel J. Math.* 8 (1970) 273–303; MR0271721 (42 #6602).
- [24] W. B. Johnson, G. Schechtman and J. Zinn, Best constants in moment inequalities for linear combinations of independent and exchangeable random variables, *Annals of Probab.* 13 (1985) 234–253; MR0770640 (86i:60054).

- [25] I. F. Pinelis and S. A. Utev, Estimates of the moments of sums of independent random variables (in Russian), *Teor. Veroyatnost. i Primenen.* 29 (1984) 554–557; Engl. transl. in *Theory Probab. Appl.* 29 (1984) 574–577; MR0761144 (85m:60034).
- [26] S. A. Utev, Extremal problems in moment inequalities (in Russian), *Limit Theorems of Probability Theory*, ed. A. A. Borovkov, Trudy Inst. Mat. 5, Nauka Sibirsk. Otdel., 1985, pp. 56–75, 175; MR0821753 (87d:60021).
- [27] T. Figiel, P. Hitczenko, W. B. Johnson, G. Schechtman and J. Zinn, Extremal properties of Rademacher functions with applications to the Khintchine and Rosenthal inequalities, *Trans. Amer. Math. Soc.* 349 (1997) 997–1027; MR1390980 (97j:60035).
- [28] R. Ibragimov and S. Sharakhmetov, On an exact constant for the Rosenthal inequality (in Russian), *Teor. Veroyatnost. i Primenen.* 42 (1997) 341–350; Engl. transl. in *Theory Probab. Appl.* 42 (1997) 294–302; MR1474714 (98j:60026).
- [29] S. R. Finch, Lengyel’s constant: Stirling partition numbers, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 316–317.
- [30] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A000110, A000296 and A005046.
- [31] R. Ibragimov and S. Sharakhmetov, The exact constant in the Rosenthal inequality for random variables with mean zero (in Russian), *Teor. Veroyatnost. i Primenen.* 46 (2001) 134–138; Engl. transl. in *Theory Probab. Appl.* 46 (2002) 127–131; MR1968709 (2004b:60054a).
- [32] R. Latała, Estimation of moments of sums of independent real random variables, *Annals of Probab.* 25 (1997) 1502–1513; <http://www.mimuw.edu.pl/~rlatala/publ.html>; MR1457628 (98h:60021).
- [33] P. Hitczenko and S. Montgomery-Smith, A note on sums of independent random variables, *Advances in Stochastic Inequalities*, Proc. 1997 Atlanta conf., ed. T. P. Hill and C. Houdré, Amer. Math. Soc., 1999, pp. 69–73; math.PR/9804068; MR1694763 (2000d:60080).