

Relations between π and the golden ratio ϕ in the form of Bailey-Borwein-Plouffe-type formulas

Jean-Christophe PAIN

CEA, DAM, DIF, F-91297 Arpajon, France
 Université Paris-Saclay, CEA, Laboratoire Matière en Conditions Extrêmes,
 91680 Bruyères-le-Châtel, France

May 19, 2022

Abstract

We provide a family of expressions of π in terms of the golden ratio ϕ in the same spirit of the formula obtained by Bailey, Borwein and Plouffe for π . Connection with cyclotomic polynomials is outlined.

1 Introduction

In 1997, Bailey, Borwein and Plouffe published the following formula [1]:

$$\pi = \sum_{k=0}^{\infty} \frac{1}{16^k} \left(\frac{4}{8k+1} - \frac{2}{8k+4} - \frac{1}{8k+5} - \frac{1}{8k+6} \right), \quad (1)$$

or

$$\pi = \sum_{k=0}^{\infty} \frac{1}{16^k} \frac{(47 + 151k + 120k^2)}{(15 + 194k + 712k^2 + 1024k^3 + 512k^4)}, \quad (2)$$

obtained using the parallel integer relation detection algorithm PSQL [2, 3]. The strength of the latter formula is that it enables one to get the k^{th} base-16 digit of π without computing any other (prior) digit. Many other formulas of this kind, often referred to as “BBP” formulas, were obtained for the first powers of π and other constants (Apéry constant $\zeta(3)$, Catalan constant, $\ln 2$, *etc.*) [4–10]. Finding exact mathematical relations between π and the golden ratio

$$\phi = \frac{1 + \sqrt{5}}{2}. \quad (3)$$

is challenging [11–15]. For instance, Baez obtained the following “Viète-type” formula [16]:

$$\pi = \frac{5}{\phi} \cdot \frac{2}{\sqrt{2 + \sqrt{2 + \phi}}} \cdot \frac{2}{\sqrt{2 + \sqrt{2 + \sqrt{2 + \phi}}}} \cdot \frac{2}{\sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2 + \phi}}}}} \cdots \quad (4)$$

and Chan found Machin-type formulas [17, 18]:

$$\begin{aligned}\pi &= 4 \arctan\left(\frac{1}{\phi}\right) + 4 \arctan\left(\frac{1}{\phi^3}\right) = 8 \arctan\left(\frac{1}{\phi^2}\right) + 4 \arctan\left(\frac{1}{\phi^6}\right) \\ &= 12 \arctan\left(\frac{1}{\phi^3}\right) + 4 \arctan\left(\frac{1}{\phi^5}\right).\end{aligned}\quad (5)$$

Other interesting formulas were also recently published, derived from new forms of Taylor expansions of inverse tangent function [19].

In the present paper, we propose a family of BBP expressions for π in terms of the golden ratio. The number π and the golden ratio ϕ are related by

$$e^{i\frac{\pi}{5}} = \frac{\phi + i\sqrt{3-\phi}}{2}, \quad (6)$$

which gives

$$\cos\left(\frac{\pi}{5}\right) = \frac{\phi}{2} \quad (7)$$

and

$$\sin\left(\frac{\pi}{5}\right) = \frac{\sqrt{3-\phi}}{2} \quad (8)$$

yielding

$$\tan\left(\frac{\pi}{5}\right) = \frac{\sqrt{3-\phi}}{\phi} \quad (9)$$

which can be put in the integral form

$$\pi = 5 \int_0^{\frac{\sqrt{3-\phi}}{\phi}} \frac{dx}{1+x^2}. \quad (10)$$

At this stage, it would be possible to consider the historical Madhava-Gregory-Leibniz, formula

$$\arctan(x) = \int \frac{dx}{1+x^2} = \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} x^{2k+1} \quad (11)$$

to get

$$\pi = \arctan\left(\frac{\sqrt{3-\phi}}{\phi}\right) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} \left(\frac{3-\phi}{\phi^2}\right)^{k+1/2}. \quad (12)$$

However, the approach we follow here, similar to the BBP one, is different. Since

$$x^4 - 1 = (x^2 - 1)(x^2 + 1), \quad (13)$$

it is possible to write

$$\pi = 5 \int_0^{\frac{\sqrt{3-\phi}}{\phi}} \frac{1-x^2}{1-x^4} dx = 5 \sum_{k=0}^{\infty} \int_0^{\frac{\sqrt{3-\phi}}{\phi}} x^{4k} dx - 5 \sum_{k=0}^{\infty} \int_0^{\frac{\sqrt{3-\phi}}{\phi}} x^{4k+2} dx \quad (14)$$

yielding

$$\pi = 5 \sum_{k=0}^{\infty} \left(\frac{\sqrt{3-\phi}}{\phi} \right)^{4k+1} \left[\frac{1}{4k+1} - \left(\frac{3-\phi}{\phi^2} \right) \frac{1}{4k+3} \right] \quad (15)$$

In the same way, using

$$1 - x^8 = (1 + x^2) (1 - x^2 + x^4 - x^6), \quad (16)$$

one gets

$$\begin{aligned} \pi &= 5 \int_0^{\frac{\sqrt{3-\phi}}{\phi}} \frac{1 - x^2 + x^4 - x^6}{1 - x^8} dx \\ &= 5 \sum_{k=0}^{\infty} \int_0^{\frac{\sqrt{3-\phi}}{\phi}} x^{8k} dx - 5 \sum_{k=0}^{\infty} \int_0^{\frac{\sqrt{3-\phi}}{\phi}} x^{8k+2} dx \\ &\quad + 5 \sum_{k=0}^{\infty} \int_0^{\frac{\sqrt{3-\phi}}{\phi}} x^{8k+4} dx - 5 \sum_{k=0}^{\infty} \int_0^{\frac{\sqrt{3-\phi}}{\phi}} x^{8k+6} dx \end{aligned} \quad (17)$$

yielding the final result:

$$\begin{aligned} \pi &= 5 \sum_{k=0}^{\infty} \left(\frac{\sqrt{3-\phi}}{\phi} \right)^{8k+1} \left[\frac{1}{8k+1} - \left(\frac{3-\phi}{\phi^2} \right) \frac{1}{8k+3} \right. \\ &\quad \left. + \left(\frac{3-\phi}{\phi^2} \right)^2 \frac{1}{8k+5} - \left(\frac{3-\phi}{\phi^2} \right)^3 \frac{1}{8k+7} \right]. \end{aligned} \quad (18)$$

2 General case

In the general case, for $p \in \mathbb{N}^*$:

$$x^{4p} - 1 = (x^2 + 1) (-1 + x^2 - x^4 + \dots + x^{2(2p-1)}), \quad (19)$$

i.e.

$$\frac{1}{x^2 + 1} = \frac{1}{(1 - x^{4p})} \sum_{k=0}^{2p-1} (-1)^k x^{2k}, \quad (20)$$

yielding the general formula, $\forall p \in \mathbb{N}^*$:

$$\pi = 5 \sum_{k=0}^{\infty} \left(\frac{\sqrt{3-\phi}}{\phi} \right)^{4pk+1} \sum_{i=0}^{2p-1} \left(\frac{\sqrt{3-\phi}}{\phi} \right)^{2i} \frac{(-1)^i}{4pk + 2i + 1}, \quad (21)$$

which is the main result of the present work. Of course, since

$$\phi^2 = \phi + 1, \quad (22)$$

one can write

$$\pi = 5 \sum_{k=0}^{\infty} \left(\frac{3-\phi}{\phi+1} \right)^{2pk+1/2} \sum_{i=0}^{2p-1} \left(\frac{3-\phi}{\phi+1} \right)^i \frac{(-1)^i}{4pk + 2i + 1}. \quad (23)$$

3 Conclusion

We presented a family of expressions of π in terms of the golden ratio ϕ in the same vein as the BBP formula derived for π . Although a number of relations involving simultaneously π and ϕ exist in the literature, some of same being of the BBP type as well, the relations presented here were, to our knowledge, not published elsewhere. It is hoped that the procedure described in the present paper may stimulate the derivation of further relations.

Appendix A: Connection with cyclotomic polynomials

The cyclotomic polynomials can be defined as

$$\Phi_n(x) = \prod_{\substack{1 \leq k \leq n \\ \gcd(k,n)=1}} \left(x - e^{\frac{2ik\pi}{n}} \right) \quad (24)$$

and satisfy

$$\prod_{d|n} \Phi_d(x) = x^n - 1. \quad (25)$$

One has also

$$\Phi_n(x) = \prod_{d|n} (x^d - 1)^{\mu\left(\frac{n}{d}\right)}, \quad (26)$$

where μ represents Möbius function:

$$\begin{aligned} \mu(n) &= +1 && \text{if } n \text{ is a square-free positive integer with an even number of prime factors,} \\ &= -1 && \text{if } n \text{ is a square-free positive integer with an odd number of prime factors,} \\ &= 0 && \text{if } n \text{ has a squared prime factor.} \end{aligned} \quad (27)$$

If p is an odd prime number, one has

$$\frac{1}{1+x^2} = -\frac{\Phi_1(x)\Phi_2(x)\Phi_p(x)\Phi_{2p}(x)\Phi_{4p}(x)}{(1-x^{4p})} \quad (28)$$

where Φ_n are the cyclotomic polynomials [20]:

$$\Phi_1(x) = x - 1, \quad (29)$$

$$\Phi_2(x) = x + 1, \quad (30)$$

$$\Phi_p(x) = \frac{1-x^p}{1-x} = x^{p-1} + x^{p-2} + \dots + x + 1, \quad (31)$$

$$\Phi_{2p}(x) = \frac{(1-x^{2p})(1-x)}{(1-x^p)(1-x^2)} = x^{p-1} - x^{p-2} + \dots - x + 1, \quad (32)$$

and

$$\Phi_{4p}(x) = \frac{(1-x^{4p})(1-x^2)}{(1-x^{2p})(1-x^4)} = x^{2p-2} - x^{2p-4} + \dots - x^2 + 1. \quad (33)$$

References

- [1] D. H. Bailey, P. Borwein and S. Plouffe, *On the rapid computation of various polylogarithms constants*, Math. Comput. **66**, 903-913 (1997).
- [2] H. R. P. Ferguson, D. H. Bailey and S. Arno, *Analysis of PSLQ, An Integer Relation Finding Algorithm*, Math. Comput. **68**, 351-369 (1999).
- [3] D. H. Bailey and D. J. Broadhurst, *Parallel Integer Relation Detection: Techniques and Applications*, Math. Comput. **70**, 1719-1736 2001.
- [4] D. H. Bailey, *A compendium of BBP-type formulas for mathematical constants*, <https://www.davidhbailey.com/dhbpapers/bbp-formulas.pdf>
- [5] G. Huvent, *Formules BBP* file:///C:/Users/painj/Desktop/BBP-phi/huvent_seminaire.pdf
- [6] K. Adegoke, *A Novel Approach to the Discovery of Binary BBP-type Formulas for Polylogarithm Constants*, Integers **12**, 345-371 (2012).
- [7] V. Adamchik and S. Wagin, *π : A 2000-year search changes direction*, Mathematica in Education and Research **1**, 11-19 (1996).
- [8] V. Adamchik and S. Wagon, *A simple formula for π* , Amer. Math. Monthly **104**, 852-855 (1997).
- [9] H. C. Chan, *π in Terms of ϕ* , The Fibonacci Quarterly **44.2**, 141-144 (2006).
- [10] H. C. Chan, *More formulas for π* , The Fibonacci Quarterly **113**, 452-455 (2006).
- [11] J. M. Borwein and P. B. Borwein, *Pi and the AGM*, John Wiley and Sons, New York, 1987.
- [12] E. Valdebenito, *Pi in Terms of Phi*, <https://vixra.org/pdf/1905.0561v1.pdf>
- [13] E. Valdebenito, *BBP-High-Precision Arithmetic*, <https://vixra.org/pdf/1711.0239v1.pdf>
- [14] S. Anderson and D. Novak, *A connection between ϕ and π* https://hascmathart.weebly.com/uploads/7/6/8/7/7687070/a_connection_between_the_number_pi_and_phi.pdf
- [15] K. Adegoke, *The golden ratio, Fibonacci numbers and BBP-type formulas*, Fibonacci Quarterly **52**, 129-138, (2014).
- [16] J. Baez, *π and the golden ratio*, <https://johncarlosgbaez.wordpress.com/2017/03/07/pi-and-the-golden-ratio/>
- [17] H.-C. Chan, *Machin-type formulas expressing π in terms of ϕ* , The Fibonacci Quarterly **46/47**, 32-37 (2008/2009).
- [18] F. Luca and P. Stanica, *On Machin's formula with powers of the golden section*, Int. J. Number Theory **05**, 973-979 (2009).

- [19] X. Wu, Z. Chen and Y. Zhu, *New BBP-type formulae for π derived from new forms of Taylor expansions of inverse tangent function*, *Mathematics* **10**, 290 (2022).
- [20] H. Riesel, *The Cyclotomic Polynomials*, in Appendix 6. *Prime Numbers and Computer Methods for Factorization*, 2nd ed. Boston, MA: Birkhäuser, pp. 305-308, 1994.